Crystal structure of a human plasma membrane phospholipid flippase

Hanayo Nakanishi[1], Katsumasa Irie[1,2], Katsumori Segawa[3], Kazuya Hasegawa[4], Yoshinori Fujiyoshi[5,6], Shigekazu Nagata[3] & Kazuhiro Abe[1,2,*]

[1]Cellular and Structural Physiology Institute; [2]Graduate School of Pharmaceutical Sciences, Nagoya University, Nagoya 464-8601, Japan; [3]WPI Immunology Frontier Research Center, Osaka University, Osaka 565-0871, Japan; [4]Japan Synchrotron Radiation Research Institute, 1-1-1 Kouto, Sayo 679-5198, Japan; [5]TMDU Advanced Research Institute, Tokyo Medical and Dental University, 1-5-45, Yushima, Bunkyo-ku, Tokyo 113-8510, Japan; [6]CeSPIA Inc., 2-1-1, Otemachi, Chiyoda, Tokyo 100-0004, Japan

Abstract

ATP11C, a member of P4-ATPase flippase, exclusively translocates phosphatidylserine from the outer to the inner leaflets of the plasma membrane, and maintains the asymmetric distribution of phosphatidylserine in the living cell. However, the mechanisms by which ATP11C translocates phosphatidylserine remain elusive. Here we show the crystal structures of a human plasma membrane flippase, ATP11C-CDC50A complex, in an outward-open E2P conformation. Two phosphatidylserine molecules are in a conduit that continues from the cell surface to the occlusion site in the middle of the membrane. Mutations in either of the phosphotidylserine binding sites or along the pathway between significantly impairs specific ATPase and transport activities. We propose a model for phosphatidylserine translocation from the outer to the inner leaflet of the plasma membrane.

Introduction

Phospholipids are asymmetrically distributed between the outer and inner leaflets in the plasma membrane of eukaryotic cells. Aminophospholipids such as phosphatidylserine (PtdSer) and phosphatidylethanolamine (PtdEtn) are confined to the inner leaflet, while phosphatidylcholine and sphingomyelin are enriched in the outer leaflet[1]. The asymmetric distribution of phospholipids is widely conserved in eukaryotes, being tuned for barrier functions and various signal transductions on the plasma membrane[2]. On occasion, this phospholipid asymmetry in the plasma membrane is disrupted, exposing PtdSer on the cell surface. Cells undergoing apoptosis expose PtdSer as an "eat me" signal to phagocytes[3,4]. Activated platelets also display PtdSer as a scaffold for clotting enzyme reactions[5]. The amphipathic nature of phospholipids prevents their spontaneous flip-flop movement across the lipid bilayer in most cases, and the translocation of phospholipids therefore needs membrane proteins to overcome the energetic barrier required for the phospholipid translocation[2].

While scramblases mediate non-specific and bi-directional movement of phospholipids between inner and outer leaflets[6], flippases exhibit ATP-driven, directional and up-hill translocation of phospholipids from the outer to inner leaflets against their concentration gradient across the membrane bilayer[7,8,9]. Different from other members of cation-transporting P-type ATPases[10–13], Type IV P-type ATPase (P4-ATPase) comprises a subfamily of P-type ATPases that transports phospholipid[14]. Among all 14 members of P4-ATPase in humans, ATP11A and ATP11C work as aminophospholipid-specific flippases at the plasma membrane[3,4]. They require an accessary subunit, CDC50A, for their correct localization and the functional expression on the plasma membrane[15–17]. In fact, cells lacking ATP11A and ATP11C, or CDC50A almost completely lose flippase activity for PtdSer and PtdEtn at the plasma membrane, resulting in failure to maintain the asymmetric accumulation of PtdSer in the inner leaflet. Their inactivation causes PtdSer exposure on the cell surface. In apoptotic cells, ATP11A and ATP11C are subjected to a caspase-mediated proteolysis and irreversible inactivation. It is proposed that calcium ions likely also inhibit the flippase activity for calcium-dependent PtdSer exposure in activated platelets or lymphocytes[18]. Regulation of aminophospholipid asymmetry by flippases is physiologically important; ATP11A-deficient mice are embryonic lethal[19], and ATP11C-deficient mice display pleiotropic phenotypes such as B-cell lymphopenia[20,21], cholestasis[22], mild anemia[23], and dystocia. Recently, mutations in the ATP11C gene were identified in patients suffered from anemia[24].

The directional translocation of specific phospholipids mediated by

P4-ATPases is achieved by the cyclic conversion of enzyme conformations, E1, E2 and

their auto-phosphorylated forms E1P and E2P, similar to the Post-Albers type reaction

scheme[14,25] for cation-transporting P-type ATPases. However, despite the huge size of

phospholipids relative to inorganic cations, the mechanism by which P4-flippases

translocate phospholipids across the membrane, the so-called "giant substrate problem"

[26,27], has remained elusive and yet is of considerable interest. Here, we describe a crystal

structure of a *bona fide* human plasma membrane flippase ATP11C-CDC50A complex

in the outward-open E2P conformation, analyzed to 3.9 Å resolution. Our structure has

two PtdSer molecules simultaneously bound in the putative lipid translocating conduit.

The structure, together with functional analyses, enable us to propose a molecular

mechanism for the phospholipid translocation across the plasma membrane by ATP11C.

Results and Discussion

Overall structure of the outward-open conformation

Human ATP11C and CDC50A were expressed using the BacMam system (Fig.

S1, Methods)[28]. Purified and deglycosylated proteins were mixed with

dioleoylphosphatidylcholine (DOPC)[29], and crystallized in the presence of phosphate

analogue[30] beryllium fluoride (BeF$_x$) and dioleoylphosphatidylserine (DOPS). Crystals

were harvested in the presence of excess DOPS, which was key to preservation of

crystal quality. X-ray diffraction data from more than 1,500 individual crystals were

merged, and the structure was determined by molecular replacement using the atomic

model of the E2BeF state of ATP8A1[31] as a search model, at a resolution of 3.9Å with

acceptable statistics of R$_{work}$/R$_{free}$ = 27.9/34.7 (Fig. S1, Table S1). As is seen in most of

the other crystallized P-type ATPases, molecular packing occurs as type I crystals in

which complexes are embedded in the lipid bilayer[32]. The asymmetric unit of the crystal

consists of four protomers of ATP11C-CDC50A. Due to different crystal contacts, the

appearance of the electron density map differs significantly for each protomer (Fig. S2).

Despite limited resolution, however, well-ordered regions, especially the

transmembrane (TM) region of protomer A and B were visible at side chain level (Fig.

S3). The overall molecular conformations of the four protomers are essentially the same,

although structural variations in some of the loop structures exist (Fig. S2). We therefore

focus on the well-ordered protomer A in what follows.

Like other members of the well-characterized, cation-transporting P-type

ATPases[11–13], the up-hill translocation of aminophospholipids by ATP11C is achieved

according to the Post-Albers type reaction scheme[14,25] (Fig. 1A). The outward-open E2P

conformation captures PtdSer or PtdEtn on the outer leaflet and induces

dephosphorylation of E2P, thus PtdSer- or PtdEtn-dependent ATP hydrolysis can be

detected (Fig. 1B), similar to the inward transport of K^+ by Na^+,K^+-ATPase and

H^+,K^+-ATPase. As we included BeF_x and PtdSer for the crystallization, the molecular

conformation[33] was expected to be the outward-open E2P state in which PtdSer is bound

to the conduit facing to the exoplasmic side (Fig. 1A). In fact, the overall structure of

the ATP11C-CDC50A complex (Fig. 1C) is very close to the corresponding E2P

structures of recently reported flippases yeast Drs2p-Cdc50p[34] and human

ATP8A1-CDC50A[31] complexes. The sequence identities of each catalytic subunit are

35.3% and 36.2%, respectively (Fig. S4 for sequence alignment)[35]. The molecular

conformation is also defined by the relative orientations of the cytoplasmic domains[36]

(actuator (A), phosphorylation (P) and nucleotide-binding (N) domains), and the

arrangement of the ten transmembrane (TM) helices of the catalytic subunit ATP11C.

The phosphate analogue BeF_x likely forms a covalent bond to the invariant aspartate in

the [409]DKTG signature sequence, which is covered by the [179]DGES/T motif located on

the surface of the A domain to prevent spontaneous dephosphorylation by the bulk water

(Fig. S5). In this conformation, the N domain is segregated from the P domain, and is

expected to be relatively flexible compared to the other two domains due to the lack of

intimate inter-domain interactions, and this is consistent with its poor electron density

(Fig. S2,3). The relative orientation of the A and P domains in ATP11C is close to those

observed in Drs2p and ATP8A1 E2P states, and clearly different from that in the

ATP8A1 E2-P_i transition state (Fig. S5), indicating that the present ATP11C structure

adopts an E2P state.

The CDC50A subunit consists of two TM helices with a long N-terminal tail

and a short C-terminus on the cytoplasmic side, and a large exoplasmic domain in

which four *N*-linked (Asn107, Asn180, Asn190 and Asn294) and one *O*-linked (Ser292)

glycosylation sites are located. We extensively investigated various combinations of

glycosylation site mutants to improve protein expression and crystal quality, and found

that the Asn190Gln/Ser292Trp double mutant produced the best crystals. Because the

samples were treated with endoglycosidase to remove excess glycans during

purification, an acetylglucosamine (GlcNAc) moiety is retained on each three remaining

N-linked glycosylation sites. Interestingly, PNGase treatment, which truncates all

glycans including core GlcNAc, produced tiny crystals, suggesting that these GlcNAcs

contribute to rather stable crystal packing. We modeled one GlcNAc moiety for each of

the two sites (Asn107, Asn294), and three glycans at a position close to Asn180

(Asn180-GlcNAc-GlcNAc-Man). The three glycans at Asn180 fit into a groove formed

between two lobes (lobe A and B) of CDC50A and Trp323 (in the connecting loop between TM3 and TM4 of ATP11C and conserved in mammalian flippases that requires CDC50A), like a wedge, thereby keeping them together and sterically protecting them from the endoglycosidase during purification (Fig. 1D).

The electron density map of the N domain allowed visualization of its overall folding at C_α level, and made it possible to assign the location of the three caspase recognition sites (Fig. 1E,F). Inactivation of ATP11A or ATP11C by effector caspases requires cleavage of multiple caspase sites located on the N domain, and a single site cleavage is not enough for full inactivation of the flippase[4,18]. There are three caspase-recognition sites in ATP11C (site I – III, Fig. 1, Fig. S4), and these are located on loops at both ends of an α-helix-containing region. Site I and site II are very close to each other at the N-terminal side of the helix, and site III is at the C-terminal side. This region, especially the α-helical part in between caspase recognition sites, is located at the center of the N domain scaffold, and seems like a bolt that holds together the surrounding α-helix and several β-sheets, and thus obviously important for N domain folding. Interaction between this helix and surrounding N domain segments may be sufficient to keep its fold even if one of the caspase sites is cleaved. Cleavage at both ends of the helix must lead to irreversible unfolding of the N domain which is required for the ATP binding. Note, all three sites are exposed on the same side of the N domain with a distance of around 35~40Å in between two regions (site I,II and site III). Such geometry of cleavage sites suggests that instantaneous two-site cleavage may occur by a caspase-3 dimer[37] upon apoptotic signal transduction, rather than single, independent cleavages, given that the distance between the catalytic center of the caspase-3 dimer is around 40Å.

Outward-open conformation

According to the reaction scheme of P4-flippase (Fig. 1A), the outward-open E2P state, mimicked by the BeF_x-binding, is a reaction state responsible for the incorporation of PtdSer from the outer leaflet to the transport occlusion site. A structural requirement for the outward-open conformation is therefore the transient formation of a conduit that physically allows the translocation of the phospholipid head group from the outer leaflet to the occlusion site in the middle of the membrane. We found a longitudinal crevice in the peripheral region of TM in ATP11C (Fig. 2A, Fig. S6). The crevice is composed of TM2, TM4 and TM6 helices and runs along TM4, and is continuous from the exoplasmic surface of the lipid bilayer to the middle of the membrane (Fig. 4B). Thus, the unwound part of TM4 (^{356}PVSM motif), which has been

implicated in the lipid transport[27], is exposed to the hydrophobic bulk lipid. Comparison

with other flippase structures in the corresponding reaction state reveals that the crevice

in ATP11C is wider than that in Drs2p activated form[34] (Fig. S6). In the ATP8A1 E2P

state, a membrane crevice is not formed at all[31], probably due to its longer C-terminal

regulatory domain compared with that in ATP11C (~38 amino acids)[38] (Fig. S4). The

exoplasmic side of the crevice is closed in the PtdSer-occluded E2-P$_i$ transition state of

ATP8A1, and clearly different from that in the ATP11C E2P state (Fig. S6). In ATP11C,

TM2 is kinked at Pro94, an amino acid residue conserved in all mammalian P4-ATPases

(Fig. 2B, Fig. S6), so that the exoplasmic side of TM2 departs from the central axis of

the crevice in this region. This structural feature enables the formation of the wide and

continuous crevice from the exoplasmic side to the PtdSer occlusion site near the

unwound part of TM4. Substitution of Pro94 for alanine significantly reduces apparent

affinities for both PtdSer and PtdEtn (Fig. 2C), indicating that the TM2 kink is key for

developing a crevice structure that is wide enough for passage of a phospholipid head

group. We conclude that the observed longitudinal crevice is in fact the outward facing

conduit that enables the phospholipid translocation.

Well-ordered crystals were generated in the presence of both BeF$_x$ and DOPS.

The thermal stability[39] of the purified ATP11C-CDC50A complex in the presence of

both BeF$_x$ and PtdSer (T_m = 53.5ºC) is markedly higher than those in the absence (T_m =

33.6ºC), or presence of either BeF$_x$ (T_m = 41.3ºC) or PtdSer alone (T_m = 46.4ºC) (Fig.

2D), indicating that the BeF$_x$-bound form binds PtdSer, and the enzyme is likely

accumulats in a distinct PtdSer-bound, but not occluded, E2P form in the crystal. This

conclusion is supported by the fact that an excess of PtdSer is required for preservation

of the crystals when harvested for X-ray diffraction studies (Methods), indicating that

PtdSer dissociates from the binding site at low to zero concentration. In fact, the

electron densities found in the crystal structure (Fig. 1C, Fig. 3) led us to model two

PtdSer molecules in the strucrture.

PtdSer binding at the exoplasmic cavity

There is a cavity at the exoplasmic side of the conduit (Fig. 3), which is formed

by the TM3-4 loop of ATP11C and the exoplasmic domain of CDC50A around the

surface of the membrane outer leaflet. This cavity is connected to the crevice observed

in the TM region (Fig. 3, Fig. S6). Unexpectedly, the omit map showed an extra density

at this position, and one PtdSer can be modeled here (Fig. 3C). The head group of the

PtdSer is accommodated deep in the cavity, the surface of which is highly hydrophilic

and electro-positive because of the many basic amino acids from both ATP11C and

CDC50A arranged here (Fig. 3AB, Fig. S4). A cluster of basic amino acids is also
observed in the cryo-EM structure of ATP8A1[31]. This positively-charged, and
hydrophilic binding cavity would be favorable for attracting the negatively-charged
phosphate group of a phospholipids. The hydrocarbon tails of the bound PtdSer lie
along the membrane plane, projecting towards the membrane lipid phase. Single
replacements of the positively-charged amino acids located at the entrance of the cavity
produced only moderate effects on PtdSer- or PtdEtn-dependent ATPase activities,
probably due to the number of remaining basic amino acids. This interpretation is
supported by the approximately 50% reduction in the V_{max} of the triple mutant
(Arg132Ala^{CDC}/ Arg133Ala^{CDC}/ Lys136Ala^{CDC}) relative to that of wild-type (Fig. 3D).
Beside these positively-charged amino acids, mutations in other hydrophilic amino
acids in this cavity (Asn325Gln, Asp343Ala in ATP11C) produce a large reduction in
apparent affinity for PtdEtn (Fig. 3E). Structural rigidity of the TM3-4 loop may also be
important, because the apparent affinity for PtdEtn is also reduced in the alanine
replacement of Trp323, which tethers the TM3-4 loop to the exoplasmic domain of
CDC50A (Fig. 1D, Fig. 3C). These structural and functional data therefore suggest that
this cavity is a priming site, responsible for the initial incorporation of phospholipids
from the membrane outer leaflet – the beginning of the transport pathway.

The TM3-4 loop in ATP11C is 6 or 5 amino acid longer than that of ATP8A1
and Drs2p, respectively, and less conserved compared with other parts of the protein
(Fig. S4). However, the hydrophilic and electro-positive nature of this cavity is
essentially the same, at least, for the three flippases whose structures have been
determined. Interestingly, some of mutations in the exoplasmic cavity had differing
effects on the apparent affinities for PtdSer and PtdEtn (Fig. 3D,E), suggesting that this
region does indeed associates with the transport ligands. This observation is consistent
with a previously reported chimeric study on yeast flippase Dnf1, in which amino acids
in the TM3-4 loop are shown to contribute to the ligand specificity[26].

PtdSer occlusion site
236

We also identified another PtdSer at the canonical occlusion site in the middle
of the membrane (Fig 4), the position of which is close to that of PtdSer occluded in the
ATP8A1 E2-P$_i$ transition state. We modeled only a part of PtdSer (phospho-L-serine
moiety) at this position according to the observed electron density (Fig. 4A), the other
part including its acyl chains are likely disordered in the bulk lipid phase. Like other
flippases as well as cation-transporting P-type ATPases, the conserved proline (Pro356)
in the PVSM sequence (corresponding to a PISL motif in most of the other P4-ATPases)

gives a characteristic unwinding at the middle of TM4, enabling the accommodation of a PtdSer head group at this position. The conduit ends at Val357 in the PVSM motif (Fig. 4B), which has been implicated as a gating residue for the phospholipids against the cytoplasmic inner leaflet[27], similar to the glutamate residue in the corresponding PEGL motif of P2-type ATPases[40,41]. Hydrophobic residue of Val98 (corresponding to Ile115 in ATP8A2) supports the gating residue Val357 on its cytoplasmic side, and these hydrophobic amino acids form a tight seal as a cytoplasmic gate, which prevents the penetration of the PtdSer head group to the cytoplasmic side in the outward-facing E2P state. In fact, mutation of these amino acids severely impaired PtdSer- or PtdEtn-dependent ATPase activity as well as PtdSer transport activity in the plasma membrane (Fig. 4C-E), consistent with previous predictions for other flippases[26]. Mutation in the hydrophilic amino acids located at the cytoplasmic side of TM1 (Gln63Ala, Arg66Ala, Asn69Ala (corresponds to Asn220 in yeast flippase Dnf1)[42]) also showed reduced V_{max} values for ATPase activity relative to wild-type. These amino acids may contribute to the rigidity of the cytoplasmic gate, or are actually part of the conduit for the inner leaflet when the cytoplasmic gate opens. Val357 in ATP11C corresponds to Ile357 in ATP8A1, Ile364 in ATP8A2 and Ile508 in Drs2p, thus of almost all the P4-ATPases only ATP11A and ATP11C have a valine residue in this position (methionine in ATP8B3). In the case of ATP11C, mutation Val357Ile produces a remarkable reduction in apparent affinity for PtdSer and PtdEtn, while V_{max} is a bit higher than that of wild-type (Fig. 4C). In fact, the transport activity of Val357Ile is comparable to that of wild-type (Fig. 4E). In contrast, the ATPase activities of Val357Ala and Val357Phe, and the transport activity of Val357Phe are significantly reduced compared with those of wild-type. Evidently, correct size is important for gating residue Val357. Mutagenesis studies also reveal the important contribution of the conserved residues around the occlusion site (Phe72, Asn352) and TM5-6 (Lys880, Asn881 and Asn912), all alanine mutants showed significant reductions in either V_{max} or apparent affinity for PtdSer and/or PtdEtn determined through ATPase activity profiles, and transport activities as well, in good agreement with previous studies of ATP8A2[27,43]. Interestingly, Phe343 is not conserved among P4-ATPases, despite its close position to the phospholipid head group. In other PtdSer-transporting P4-ATPases, including ATP8A1, ATP8A2 and Drs2p, Phe343 in ATP11C is replaced with asparagine (Fig. S4), and the hydrophilic residue actually contributes to PtdSer coordination in the ATP8A1 E2-P$_i$ structure[31]. The Phe354Asn mutant in ATP11C showed significantly higher affinity for PtdSer and PtdEtn relative to wild-type while keeping its V_{max} of ATPase activity and PtdSer transport activity comparable to wild-type level. Therefore, a

hydrophilic or smaller side chain in this position is favorable for the accommodation of
the aminophospholipid head group in the occlusion site. It can be speculated that
ATP11C has Val357 and Phe354 instead of the most conserved isoleucine and
asparagine of other PtdSer-dependent flippases, to fine-tune the PtdSer and PtdEtn
transport activity relevant to a plasma membrane flippase. As an important note,
mutations Val98Ala, Phe72Ala, Asn881Ala and Asn912Ala increases the apparent
affinity for PtdSer, but lower that for PtdEtn, relative to the wild-type (Fig. 4D),
therefore these mutants were able to discriminate PtdSer and PtdEtn.
288
289   **A transport model**
290   Two PtdSer molecules bound to both ends of the conduit found in the crystal
structure (Figs. 1,3 and 4) led us to put forward a transport model for the flippase, which
is distinct from other models proposed so far[26,27,44] (Fig. 5). Translocation of the
phospholipid, either PtdSer or PtdEtn is initiated by its binding to the positively-charged
and hydrophilic cavity composed of the CDC50A exoplasmic domain and the ATP11C
TM3-4 loop (Fig. 3, Fig. S7). The electro-positive and hydrophilic nature of the cavity
likely attracts the head group of the phospholipid from the outer leaflet layer of the
membrane. The exoplasmic cavity connects to the longitudinal crevice along TM4 (Fig.
2, Fig. S6). Therefore, once the phospholipid head group is incorporated into the
exoplasmic cavity, it may diffuse along the crevice while keeping its hydrocarbon
chains projecting to the hydrophobic bulk lipid. Val349 in TM4 projects into the conduit
(Fig. 4AB). Its replacement with bulky phenylalanine (Val349Phe) severely impairs
PtdSer- and PtdEtn-dependent ATPase activity as well as PtdSer transport activity
relative to wild-type, in contrast to the moderate effects of alanine substitution (Fig.
4C-E). In addition, phenylalanine mutations of residues close to Val349 (Thr90Phe in
TM2 and Leu350Phe in TM4) also significantly lowers ATPase activity relative to the
wild-type. These amino acid residues are located between, and distant from, the two
substrate binding sites and clearly bulky substitutions impede diffusion of the
phospholipid head group from the exoplasmic cavity to the occlusion site. In the
occlusion site, the phospholipid head group must be coordinated by conserved
hydrophilic amino acids such as Asn352 and Asn881. The gating residue Val357 blocks
further penetration of the phospholipid to the cytoplasmic inner leaflet. PtdSer binding
to the occlusion site may be the signal to induce the conformational change required for
reaching the PtdSer-occluded E2-P_i transition state. Some of the hydrophilic amino
acids in the TM1-2 loop may join in the coordination of the head group, as observed in
the ATP8A1 E2-P_i structure (Fig. S6)[31]. Movement of TM1-2, as a result of the loop

involvement, adjusts the A domain to induce dephosphorylation of E2P. Attainment of
the E2 or E1state, the next step, opens the cytoplasmic gate. A physical pathway from
the occlusion site to the inner leaflet need to be opened. Mutation of amino acid
residues around the cytoplasmic gate largely inhibited phospholipid dependent ATPase
activity (Fig. 4). Movement of the TM1-2 helix bundle outwards may be needed for the
phospholipid to move to the cytoplasmic inner leaflet, although this has not yet been
elucidated.

Translocation of the phospholipid across the two leaflets is energetically
extremely unfavorable due to the amphiphilic nature of phospholipids[45], and the rate of
spontaneous phospholipid flip-flop is order of several hour to several days. In our
envisaged translocation mechanism, the most energy-consuming step may be moving
the hydrophilic head group from the outer membrane surface to the transport conduit. In
this step the phospholipid head group needs to disconnects from the polar interactions
formed with neighboring phospholipids, surrounding membrane proteins as well as
water molecules, and also needs to change its orientation approximately 90º from a
vertical to horizontal orientation in the lipid bilayer. The structure and resulting
transport model answer the fundamental question in the translocation mechanism of
flippase -how phospholipid reaches the transport conduit from the outer leaflet. The
hydrophilic surface of the exoplasmic cavity interior provides an environment similar to
the water-facing membrane surface and lowers the energetic barrier required for
acquisition of the phospholipid head group. Many membrane transport proteins seem to
employ a common strategy for the translocation of their specific substrates; the
environment of the substrate binding site of the protein mimics that of the location from
whence the substrate comes. Flippases are not an exception, and this strategy is applied
in the most energy-consuming step in the sequence of the lipid flipping process by the
ATP11C-CDC50A complex.
342

Acknowledgement

We thank M. Taniguchi for technical assistance, Dr. T. Nishizawa for sharing
unpublished results of the ATP8A1 structure and Dr. D. McIntosh for improving the
manuscript. This work was supported by Grants-in-Aid for the Scientific Research
(17H03653), Basis for Supporting Innovative Drug Discovery and Life Science
Research (BINDS) from Japan Agency for Medical Research and Development
(AMED), Takeda Science Foundation (to K.A.); Core Research for Evolutional Science
and Technology from JST (JPMJCR14M4, to S.N. and K.A.); Grants-in-Aid for
Scientific Research (S), the Japan New Energy and Industrial Technology Development
Organization (NEDO), and the Japan Agency for Medical Research and Development
(AMED) (to Y.F.). This work is a part of a project (2018B2703 and 2019B2707) at
SPring-8. This research is partly supported by the Platform Project for Supporting Drug
Discovery and Life Science Research (BINDS) from AMED under Grant number
JP18am0101070.

Author Contributions Y.F., S.N. and K.A. designed the study. H.N. and K.A. were
responsible for protein expression. H.N. purified and crystallized the protein. H.N. K.S.
and K.A. performed biochemical analysis. H.N., K.H. and K.A. collected X-ray
diffraction data. K.H. merged X-ray diffraction data. K.I. and K.A. analyzed the
structure. K.I. and K.A. interpreted the structure. H.N. K.S. and K.A. wrote the
manuscript with agreement of all authors.

Author Information

Atomic coordination and structure factors for the structures reported in this work have
been deposited in the Protein Data Bank under accession number NXXX.
Correspondence and requests for materials should be addressed to K.A.
(kabe@cespi.nagoya-u.ac.jp).

Materials and Methods

Protein expression and purification

Human *ATP11C* (NCBI: XM_005262405.1)[4] was sub-cloned into a hand-made
vector as described previously[13]. Both of the amino terminal 7 amino acids (ΔN7) and
the carboxyl terminal 38 amino acids (ΔC38) of hATP11C were truncated, and the Flag
epitope tag (DYKDDDDK), hexa-histidine tag, the enhanced green fluorescence protein
(EGFP) followed by a tobacco etch virus (TEV) protease recognition sequence were
attached to the amino terminal of the deletion mutant (ATP11C_cryst). Human *CDC50A*

cDNA (NCBI: NM_018247.3) was sub-cloned into the vector independently. The

Asn190Gln and Ser292Trp double mutation was introduced into the construct to

regulate its glycosylation status (CDC50A_QW). The former mutation is simply aimed

to prevent *N*-linked glycosylation. The latter is to eliminate *O*-linked glycosylation at

Ser292 and at the same time to increase the efficiency of *N*-linked glycosylation at

Asn294[48], because these residues appeared to be glycosylated alternatively. The

heterodimer composed of ATP11C_cryst and CDC50A_QW was successfully expressed

in the plasma membrane using baculovirus-mediated transduction of mammalian

Expi293 cells (Thermo) for 48 h at 31.5 °C as described previously[28,46]. The harvested

cells were directly solubilized with 1.5 % (w/v) n-decyl β-D-maltoside in a lysis buffer

containing 40 mM MES/Tris (pH 6.5), 200 mM NaCl, 2 mM $Mg(CH_3COO)_2$, 1 mM

ATP, 1 mM dithiothreitol, 0.1 mM ethylene

glycol-bis(2-aminoethylether)-*N,N,N',N'*-tetraacetic acid (EGTA) and protease inhibitor

cocktail (Roche) on ice for 20 min. After removing the insoluble material by

ultracentrifugation (200,000×g for 1h), the supernatant was mixed with anti-Flag M2

affinity resin (Sigma-Aldrich) for 1 h at 4 °C. The resin was washed with 20 column

volumes of buffer consisting of 20 mM MES/Tris (pH 6.5), 200 mM NaCl, 5% (v/v)

glycerol, 1 mM $Mg(CH_3COO)_2$, 0.1 mM ATP, 0.1 mM EGTA and 0.03% octaethylene

glycol monododecyl ether ($C_{12}E_8$, Nikko Chemical). Flag-EGFP tagged ATP11C was

eluted with 0.2 mg/ml Flag peptide (Sigma-Aldrich) in the wash buffer. Eluted proteins

were incubated with TEV protease and MBP-fusion endoglycosidase (EndoHf, New

England Biolabs) at 4 °C overnight. Released affinity tags containing Flag-EGFP were

removed from the mixture by a Ni-NTA resin (QIAGEN). The non-absorbed fractions

were concentrated and subjected to a size-exclusion column chromatograph using a

Superose6 Increase column (GE Healthcare), equilibrated in a buffer comprising 20 mM

MES/Tris (pH 6.5), 1%(v/v) glycerol, 50 mM NaCl, 5 mM $MgCl_2$, and 0.03% $C_{12}E_8$.

Peak fractions were collected and concentrated to 10 mg/ml. See Fig. S1 for the purity

of the sample at each step. The concentrated ATP11C samples were mixed with 1 mM

ADP, 0.5 mM $BeSO_4$, 1.5 mM NaF, and 0.1 mg/ml dioleoyl phosphatidylserine (DOPS),

and added to the glass tubes in which a layer of dried dioleoyl phosphatidylcholine

(DOPC) had formed, in a lipid-to-protein ratio of 0.2. $C_{12}E_8$ was added to the glass

tubes in a protein-to-detergent ratio of 0.5 to 2.0, and the mixture was incubated

overnight at 4 °C in a shaker mixer operated at 120 rpm[29]. After removing the insoluble

material by ultracentrifugation, lipidated samples were subjected to crystallisation. Note

that the effect of truncation of the both terminal amino acids of ATP11C, the double

mutation introduced to CDC50A and deglycosylation of CDC50A on the PtdSer- and

PtdEtn-dependent ATPase activity were negligible compared with wild-type without
endoglycosidase treatment.
417

Gene editing for CDC50A
The CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats)-Cas
(CRISPR-associated) system with pX330 vector (Addgene) was used to edit the
CDC50A gene in HEK293S GnT1- cells as described [4]. The sgRNA sequences for
human CDC50A were as follows; 5'-CACCGGGCAACGTGTTTATGTATTA-3' and
5'-AAACTAATACATAAACACGTTGCCC-3'. Target protospacer sequences are
underlined.
425

Crystallization and data collection
Crystals were obtained by vapour diffusion at 20°C. The lipidated 10 mg/ml protein
sample obtained either from Expi293 cell or CDC-KO cells, containing 1 mM ADP, 0.5
mM $BeSO_4$, 1.5 mM NaF, and 0.1 mg/ml DOPS was mixed with reservoir solution
containing 10% (v/v) glycerol, 14-17% PEG4000, 0.4 M $MgSO_4$, and 2 to 5 mM
β-mercaptoethanol. Crystals made using protein samples purified from Expi293 cells
grew in a thin plate-like shape with the dimensions of $800 \times 500 \times 50$ μm in 2 weeks. In
contrast, crystals from CDC-KO cells grew as small crystals usually less than 50 μm
with a polyhedron shape. These crystals were picked up with LithoLoops (Protein Wave
Corporation), and flash frozen in liquid nitrogen. The crystals were harvested in the
presence of 10% (v/v) glycerol, 14-17% PEG4000, 0.4 M $MgSO_4$, 4 mg/ml DOPS, 20
mM MES/Tris (pH 6.5), 50 mM NaCl, 5 mM $MgCl_2$, 5 mM β-mercaptoethanol, 2%
$C_{12}E_8$, 1 mM ADP, 0.5 mM $BeSO_4$ and 1.5 mM NaF. Note, when crystals were
harvested in the absence of DOPS, a few percent approximately of crystals gave X-ray
diffraction better than 4Å. In the presence of 4 mg/ml DOPS, however, the number of
well-diffracting crystals mostly increased to approximately 30% of all harvested crystals.
Despite the different crystal morphologies of the crystals obtained from Expi293 cells
and CDC50A KO cells (Fig. S1), the crystals show the same space group ($P2_12_12_1$) and
unit cell size ($a = 100.5$Å, $b = 232.8$Å, $c = 492.9$ Å, $\alpha = \beta = \gamma = 90°$).
X-ray diffraction data were collected at the SPring-8 beamline BL32XU, BL41XU and
BL45XU. For the large plate-like crystals obtained from Expi293 cells, X-ray
diffraction data were collected by helical scan method[49], or by irradiating micro-focus
beam from the direction perpendicular to the *c*-axis by monitoring crystals on 90° bent
LithoLoop (Fig. S1). Crystals from CDC-KO cells were too small to collect full data set
from each crystal, and well-diffracted crystal could not be determined from its

morphology. Therefore, multiple crystals were mounted on a ϕ1 μm LithoLoop, and the raster scan was performed to identify well-diffracted crystals. After selecting target crystals, 10º small-wedge data were collected from each individual crystal (Fig. S1). Total 1,588 crystals were used for the data collection and some of them were performed automatic manner by using ZOO system[50].

Structural determination and analysis

All the diffraction data from individual 1,588 well-diffracting crystals were processed and merged using automatic data processing system KAMO[47] with XDS[51]. Structure factors were subjected to anisotropy correction using the UCLA MBI Diffraction Anisotropy server[52] (http://services.mbi.ucla.edu/anisoscale/). The structure was determined by molecular replacement with PHASER, using a homology model based on the cryo-EM structure of the E2P state of ATP8A1 (PDB ID: 6K7L) as a search model. Coot[53] was used for cycles of iterative model building and Refmac5[54] and Phenix[55] were used for refinement. The final crystallographic model of BeF$_x$-bound human ATP11C at a 3.8Å resolution, refined to R_{work} and R_{free} of 0.29 and 0.36 was deposited in the PDB with accession code PDB: 6LKN. Figures were prepared using UCSF Chimera[56] and PYMOL (https://pymol.org).

Activity assay using recombinant proteins

To measure the ATPase activity, Flag-EGFP tag connected by the TEV cleavage site to the N-terminal tail of ATP11C_cryst was used to estimate its expression level by fluorescence size-exclusion column chromatography (FSEC)[57]. The original and additional mutant complexes of ATP11C_cryst and the CDC50A_QW were expressed using the BacMam system and purified in a smaller batch format as described above except for TEV protease digestion and endoglycosidase treatment. The purified proteins (the purity of samples used for the ATPase measurement was comparable to lane 4 of SDS-PAGE in Fig. S1) were subjected to an ATPase activity assay as described previously[58]. Briefly, partially purified ATP11C (wild-type or mutants) was suspended in buffer comprising 40 mM HEPES, 2 mM MgCl$_2$, 2 mM ATP, 2% glycerol, 100 mM NaCl, 0.03 mg/ml C$_{12}$E$_8$ (pH 7.0 adjusted by Tris) and indicated concentrations of phospholipids (DOPS or POPE, dissolved as 10 mg/ml stock in 2% C$_{12}$E$_8$), in 96-well plates. Reactions were initiated by incubating the samples at 37 ℃ using a thermal cycler, and maintained for 1h. Reactions were terminated, and the amount of released inorganic phosphate was determined colorimetrically using a microplate reader (TECAN). Samples used for the ATPase measurement were analyzed by FSEC with

monitoring with Trp fluorescence (Ex 280 nm, Em 320 nm) monitoring, and peak

fluorescence values were determined. The peak values of samples were compared to

that of a fully purified sample used for the crystallization whose protein concentration

was accurately determined by UV absorption, and protein concentrations for each

measured sample were estimated. The phospholipid concentration-dependent ATPase

activities were plotted, and data fitted as described previously[43] to estimate apparent

affinities ($K_{0.5}$) and V_{max} using PRISM4 software. For all measurements, data were

duplicated at twelve different phospholipid concentrations for a single measurement,

and at least three independent measurements were conducted for each mutant. The $K_{0.5}$

and V_{max} values plotted in Fig. 3 and 4 are mean and S.D.s estimated from at least three

independent measurements, and representative ones are shown in Fig. S5. Note, the

PtdSer- or PtdEtn-dependent specific ATPase activities of ATP11C_cryst-CDC50A_QW

complex were almost the same as those of wild-type. We therefore refer to

ATP11C_cryst-CDC50A_QW as wild-type in the activity assay for simplicity.

Flippase assay

Flippase activity was determined as described[59]. In brief, *ATP11A-ATP11C*

double-deficient WR19L mutant (*DKO*) cells were transformed with retroviruses

carrying cDNA for human FLAG-tagged CDC50A and EGFP-tagged wild-type ATP11C

or mutants. The stable transformants were then subjected to cell sorting for EGFP with

FACSAriaII (BD Biosciences), cells at the same EGFP-intensity were sorted, and

expanded. Amounts of EGFP-tagged proteins and their localization to the plasma

membrane were verified by Western blotting and a confocal microscope (FV-1000D;

Olympus), respectively. For the flippase assay, *DKO* cells and its transformants

expressing wild-type ATP11C or mutants were incubated with 1 μM NBD-PS

(1-oleoyl-2-{6-[(7-nitro-2-1,3-benzoxadiazol-4-yl)amino]hexanoyl}-*sn*-gylcero-3-phosp

hoserine for 3 min at 20 ℃ in Hanks' balanced salt solution (HBSS) containing 1 mM

$MgCl_2$ and 2 mM $CaCl_2$. The cells were collected by centrifugation, resuspended in

HBSS containing 5 mg/ml fatty acid-free BSA to extract nonincorporated lipids, and

analyzed by FACSCanto II (BD Biosciences).

Thermal stability

Purified samples were incubated at the indicated temperatures for 10 min in the

presence of 40 mM HEPES, 100 mM NaCl, 2 mM $MgCl_2$ (free) with 1 mM $BeSO_4$, 3

mM NaF (BeF) and/or 0.1 mg/ml DOPS (PtdSer). After incubation, samples were

cooled on ice and denatured aggregates removed using a membrane filter (pore size 0.22

µm), and the resulting filtrates were analyzed by size-exclusion column chromatography

using Superose 6 Increase 10/150 GL (GE healthcare). Peak values at the retention time

for the complex were plotted as a function of incubation temperature, and their T_m

values estimated.

References

1. Leventis, P. A. & Grinstein, S. The Distribution and Function of Phosphatidylserine in Cellular Membranes. *Annu. Rev. Biophys.* **39**, 407–427 (2010).

2. Pomorski, T. & Menon, A. K. Lipid flippases and their biological functions. *Cellular and Molecular Life Sciences* **63**, 2908–2921 (2006).

3. Segawa, K. & Nagata, S. An Apoptotic 'Eat Me' Signal: Phosphatidylserine Exposure. *Trends Cell Biol.* **25**, 639–650 (2015).

4. Segawa, K. *et al.* Caspase-mediated cleavage of phospholipid flippase for apoptotic phosphatidylserine exposure. *Science (80-.).* **344**, 1164–1168 (2014).

5. Bevers, E. M. *et al.* The complex of phosphatidylinositol 4,5-bisphosphate and calcium ions is not responsible for Ca2+-induced loss of phospholipid asymmetry in the human erythrocyte: A study in Scott syndrome, a disorder of calcium-induced phospholipid scrambling. *Blood* **86**, 1983–1991 (1995).

6. Williamson, P. Phospholipid Scramblases Supplementary Issue: Cellular Anatomy of Lipid Traffic. 41–44 doi:10.4137/Lpi.s31785

7. Tang, X., Halleck, M. S., Schlegel, R. A. & Williamson, P. A subfamily of P-type ATPases with aminophospholipid transporting activity. *Science (80-.).* **272**, 1495–1497 (1996).

8. Coleman, J. A., Kwok, M. C. M. & Molday, R. S. Localization, purification, and functional reconstitution of the P4-ATPase Atp8a2, a phosphatidylserine Flippase in photoreceptor disc membranes. *J. Biol. Chem.* **284**, 32670–32679 (2009).

9. Shin, H.-W. & Takatsu, H. Substrates of P4-ATPases: beyond aminophospholipids (phosphatidylserine and phosphatidylethanolamine). *FASEB J.* **33**, 3087–3096 (2019).

10. Palmgren, M. G. & Axelsen, K. B. Evolution of P-type ATPases. *Biochim. Biophys. Acta* **1365**, 37–45 (1998).

11. Toyoshima, C., Nakasako, M., Nomura, H. & Ogawa, H. Crystal structure of the calcium pump of sarcoplasmic reticulum Ê resolution. *Nature* **405**, 647–655 (2000).

12. Morth, J. P. *et al.* Crystal structure of the sodium–potassium pump. *Nature* **450**, 1043–1049 (2007).

13. Abe, K., Irie, K., Nakanishi, H., Suzuki, H. & Fujiyoshi, Y. Crystal structures of the gastric proton pump. *Nature* **556**, 214–229 (2018).

14. Andersen, J. P. *et al.* P4-ATPases as phospholipid flippases-structure, function, and enigmas. *Front. Physiol.* **7**, 1–23 (2016).

15.   Saito, K. *et al.* Cdc50p, a protein required for polarized growth, associates with
the Drs2p P-type ATPase implicated in phospholipid translocation in
Saccharomyces cerevisiae. *Mol. Biol. Cell* **15**, 3418–3432 (2004).
16.   Bryde, S. *et al.* CDC50 proteins are critical components of the human class-1 P
4-ATPase transport machinery. *J. Biol. Chem.* **285**, 40562–40572 (2010).
17.   Segawa, K., Kurata, S. & Nagata, S. The CDC50A extracellular domain is
required for forming a functional complex with and chaperoning phospholipid
flippases to the plasma membrane. *J. Biol. Chem.* **293**, 2172–2182 (2018).
18.   Segawa, K., Kurata, S. & Nagata, S. Human type IV P-type ATPases that work
as plasma membrane phospholipid flippases and their regulation by caspase and
calcium. *J. Biol. Chem.* **291**, 762–772 (2016).
19.   Perez-Garcia, V. *et al.* Placentation defects are highly prevalent in embryonic
lethal mouse mutants. (2018). doi:10.1038/nature26002
20.   Yabas, M. *et al.* ATP11C is critical for the internalization of phosphatidylserine
and differentiation of B lymphocytes. *Nat. Immunol. Vol.* **12**, (2011).
21.   Siggs, O. M. *et al.* The P4-type ATPase ATP11C is essential for B lymphopoiesis
in adult bone marrow. *Nat. Immunol.* **12**, (2011).
22.   Siggs, O. M., Schnabl, B., Webb, B. & Beutler, B. X-linked cholestasis in mouse
due to mutations of the P4-ATPase ATP11C. doi:10.1073/pnas.1104631108
23.   Yabas, M. *et al.* Mice Deficient in the Putative Phospholipid Flippase ATP11C
Exhibit Altered Erythrocyte Shape, Anemia, and Reduced Erythrocyte Life Span
*. (2014). doi:10.1074/jbc.C114.570267
24.   Arashiki, N. *et al.* ATP11C is a major flippase in human erythrocytes and its
defect causes congenital hemolytic anemia. *Haematologica* **101**, 559–565 (2016).
25.   Coleman, J. A., Vestergaard, A. L., Molday, R. S., Vilsen, B. & Andersen, J. P.
Critical role of a transmembrane lysine in aminophospholipid transport by
mammalian photoreceptor P4-ATPase ATP8A2. *Proc. Natl. Acad. Sci.* **109**,
1449–1454 (2012).
26.   Baldridge, R. D. & Graham, T. R. Two-gate mechanism for phospholipid
selection and transport by type IV P-type ATPases. *Proc. Natl. Acad. Sci. U. S. A.*
110, E358-67 (2013).
27.   Vestergaard, A. L. *et al.* Critical roles of isoleucine-364 and adjacent residues in
a hydrophobic gate control of phospholipid transport by the mammalian
P4-ATPase ATP8A2. *Proc. Natl. Acad. Sci.* **111**, E1334–E1343 (2014).
28.   Dukkipati, A., Park, H. H., Waghray, D., Fischer, S. & Garcia, K. C. BacMam
system for high-level expression of recombinant soluble and membrane

glycoproteins for structural studies. *Protein Expr. Purif.* **62**, 160–170 (2008).

29.   Gourdon, P. *et al.* HiLiDe-systematic approach to membrane protein

crystallization in lipid and detergent. *Crystal Growth and Design* **11**, 2098–2106

(2011).

30.   Danko, S., Yamasaki, K., Daiho, T., Suzuki, H. & Toyoshima, C. Organization of

cytoplasmic domains of sarcoplasmic reticulum Ca2+-ATPase in E1P and

E1ATP states: A limited proteolysis study. *FEBS Lett.* **505**, 129–135 (2001).

31.   Hiraizumi, M., Yamashita, K., Nishizawa, T. & Nureki, O. Cryo-EM structures

capture the transport cycle of the P4-ATPase flippase. *Science (80-.).* **365**,

1149–1155 (2019).

32.   Norimatsu, Y., Hasegawa, K., Shimizu, N. & Toyoshima, C.

Protein-phospholipid interplay revealed with crystals of a calcium pump. *Nature*

545, 193–198 (2017).

33.   Danko, S. *et al.* ADP-insensitive phosphoenzyme intermediate of sarcoplasmic

reticulum Ca2+-ATPase has a compact conformation resistant to proteinase K,

V8 protease and trypsin. *FEBS Lett.* **489**, 277–282 (2001).

34.   Timcenko, M. *et al.* Structure and autoregulation of a P4-ATPase lipid flippase.

Nature **571**, 366–370 (2019).

35.   Katoh, K., Misawa, K., Kuma, K.-I. & Miyata, T. *MAFFT: a novel method for*

rapid multiple sequence alignment based on fast Fourier transform.

36.   Toyoshima, C., Norimatsu, Y., Iwasawa, S., Tsuda, T. & Ogawa, H. How

processing of aspartylphosphate is coupled to lumenal gating of the ion pathway

in the calcium pump. *Proc. Natl. Acad. Sci.* **104**, 19831–19836 (2007).

37.   MacKenzie, S. H. & Clark, A. C. Death by caspase dimerization. *Adv. Exp. Med.*

Biol. **747**, 55–73 (2012).

38.   Takatsu, H. *et al.* Phospholipid flippase ATP11C is endocytosed and

downregulated following Ca2+mediated protein kinase C activation. *Nat.*

Commun. **8**, (2017).

39.   Hattori, M., Hibbs, R. E. & Gouaux, E. A fluorescence-detection size-exclusion

chromatography-based thermostability assay for membrane protein

precrystallization screening. *Structure* **20**, 1293–1299 (2012).

40.   Inesi, G., Lewis, D., Toyoshima, C., Hirata, A. & De Meis, L. Conformational

fluctuations of the Ca2+-ATPase in the native membrane environment: Effects of

pH, temperature, catalytic substrates, and thapsigargin. *J. Biol. Chem.* **283**,

1189–1196 (2008).

41.   Tsunekawa, N., Ogawa, H., Tsueda, J., Akiba, T. & Toyoshima, C. Mechanism

637   of the E2 to E1 transition in Ca2+ pump revealed by crystal structures of gating

638   residue mutants [Biochemistry]. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 2–7 (2018).

639 42. Roland, B. P. & Graham, T. R. Directed evolution of a sphingomyelin flippase

640   reveals mechanism of substrate backbone discrimination by a P4-ATPase. *Proc.*

641   *Natl. Acad. Sci.* **113**, E4460–E4466 (2016).

642 43. Mikkelsen, S. A. *et al.* Asparagine 905 of the mammalian phospholipid flippase

643   ATP8A2 is essential for lipid substrate–induced activation of ATP8A2

644   dephosphorylation. *J. Biol. Chem.* **294**, 5970–5979 (2019).

645 44. Jensen, M. S. *et al.* Phospholipid flipping involves a central cavity in P4 ATPases

646   OPEN. doi:10.1038/s41598-017-17742-y

647 45. Pomorski, T. G. & Menon, A. K. Lipid somersaults: Uncovering the mechanisms

648   of protein-mediated lipid flipping. *Progress in Lipid Research* (2016).

649   doi:10.1016/j.plipres.2016.08.003

650 46. Goehring, A. *et al.* Screening and large-scale expression of membrane proteins in

651   mammalian cells for structural studies. *Nat. Protoc.* **9**, 2574–2585 (2014).

652 47. Yamashita, K., Hirata, K. & Yamamoto, M. KAMO: towards automated data

653   processing for microcrystals. *Acta Crystallogr. Sect. D Struct. Biol.* **74**, 441–449

654   (2018).

655 48. Murray, A. N. *et al.* Enhanced Aromatic Sequons Increase

656   Oligosaccharyltransferase Glycosylation Efficiency and Glycan Homogeneity.

657   *Chem. Biol.* (2015). doi:10.1016/j.chembiol.2015.06.017

658 49. Flot, D. *et al.* The ID23-2 structural biology microfocus beamline at the ESRF. *J.*

659   *Synchrotron Radiat.* **17**, 107–118 (2010).

660 50. Hirata, K. *et al.* Zoo: An automatic data-collection system for high-throughput

661   structure analysis in protein microcrystallography. *Acta Crystallogr. Sect. D*

662   *Struct. Biol.* **75**, 138–150 (2019).

663 51. Kabsch, W. *et al.* XDS. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 125–132

664   (2010).

665 52. Strong, M. *et al.* Toward the structural genomics of complexes: crystal structure

666   of a PE/PPE protein complex from Mycobacterium tuberculosis. *Proc. Natl.*

667   *Acad. Sci. U. S. A.* **103**, 8060–5 (2006).

668 53. Emsley, P. *et al.* Coot⊡: model-building tools for molecular graphics. *Acta*

669   *Crystallogr. Sect. D Biol. Crystallogr.* **60**, 2126–2132 (2004).

670 54. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal

671   structures. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **67**, 355–367 (2011).

672 55. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for

macromolecular structure solution. *Acta Crystallogr. D. Biol. Crystallogr.* **66**,
213–21 (2010).

56.   Pettersen, E. F. *et al.* UCSF Chimera- A visualization system for exploratory
research and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

57.   Kawate, T. & Gouaux, E. Fluorescence-Detection Size-Exclusion
Chromatography for Precrystallization Screening of Integral Membrane Proteins.
Structure **14**, 673–681 (2006).

58.   Yamamoto, K. *et al.* A single K^+-binding site in the crystal structure of the
gastric proton pump. *eLife* **8**, (2019).

59.   Segawa, K. *et al.* Phospholipid flippases enable precursor B cells to flee
engulfment by macrophages. doi:10.1073/pnas.1814323115

Figures

686
Fig. 1. Crystal structure of ATP11C-CDC50A complex.
(A) Reaction scheme of phospholipid (PL) translocation coupled with ATP-hydrolysis.
Cartoons represent molecular conformations of ATP11C-CDC50A complex (inward- or
outward-, and open or occluded states). (B) PL-dependent ATPase activity by the
purified ATP11C-CDC50A complex. Specific ATPase activities in the presence of
DOPS (red), POPE (blue) or DOPC (black) were plotted as a function of their
concentrations. DOPS gave the highest ATPase activity, POPE showed intermediate,
while there was no detectable ATPase activity in the presence of DOPC. (C) Overall
structure of the outward-open E2P state of ATP11C-CDC50A complex in ribbon
representation. Color of the ATP11C gradually changes from the N-terminus (purple) to
the C-terminus (red). CDC50A subunit is shown in grey ribbon, and attached *N*-linked
glycans were displayed as green sticks. A DOPS molecule (PtdSer) and its hydrophilic
phosphoserine part are shown as spheres in the exoplasmic cavity and the occlusion site
in the middle of the TM domain, respectively. Three cytoplasmic domains are indicated
with different colors, and caspase recognition sequences at the N domain surface are
indicated in red. Gray ribbon in the N domain thus indicates the region that is removed
after the caspase cleavage. (D) Closed view of the exoplasmic region of CDC50A
subunit. Surface represents electron density map (1.5σ). Lobe A (light grey) and lobe B
(dark grey) are shown in different colors, and TM3-4 loop is shown in blue with Trp323
side chain fitted into the map. *N*-linked glycans (green) are highlighted. (E,F) Close-up
of the N domain (light blue) from two different viewpoints. Surface shows electron
density map at 1.0σ contour level. Three caspase recognition sites (I, II and III) and the
region in between them (Gly466-Asn477) are indicated in red and grey, respectively.

Fig. 2. The crevice in the TM region.
(A) Surface representation of the ATP11C-CDC50A complex shows the crevice in the
TM region. Green spheres with CPK coloring represent phosphoserine at the occlusion
site in the crevice, and PtdSer bound to the exoplasmic cavity. Surfaces of the atomic
model of ATP11C and CDC50A are shown in blue and grey, respectively. (B) Close-up
view of the membrane crevice indicated as a dotted box in A. The crevice is mostly
composed of TM2, TM4 and TM6. Pro94 makes a kink at the exoplasmic side of TM2
(yellow dotted lines), which exposes the unwound region of TM4 (PVSM, shown in
spheres) to the lipid bilayer phase. (C) PtdSer- or PtdEtn-dependent ATPase activities of
wild-type (same as in Fig. 1) and Pro94Ala mutant as indicated in the figure. (D)
Thermal stabilities of purified ATP11C-CDC50A complex determined by FSEC (see
Methods). Peak values in the FSEC analysis were plotted as a function of treatment
temperature in the absence (free) or presence of indicated substrates.

728
Fig. 3. Exoplasmic cavity
(A,B) Surface representations of the atomic model of ATP11C-CDC50A according to
their hydrophobicity (A) or Coulombic surface potentials (B). Each model was sliced at
the plane where the phosphate moiety of PtdSer is located, along with the membrane
plane, and opened up to show the surface of the exoplasmic cavity interior. Color codes
as indicated in the figures. (C) PtdSer binding cavity in detail. Some hydrophilic amino
acids surrounding PtdSer are shown as sticks. Color codes as in Fig. 1. Red mesh and
transparent red surface represent the 2Fo-Fc electron density maps around the PtdSer
molecule with contour levels of 1.0σ and 2.0σ, respectively. (D,E) The V_{max} and
apparent affinity for PtdSer or PtdEtn derived from the ATPase activities of indicated
mutants. The ATPase activity per mg of purified protein was determined and analyzed in
the presence of varying concentrations of PtdSer or PtdEtn (Representatives shown in
Fig. S7). The V_{max} is shown as a percentage of the wild-type value (D). The apparent
affinities for phospholipids are expressed as the concentration giving half-maximum
activation ($K_{0.5}$), and plotted as x-fold of wild-type values (E).

Fig. 4. Transmembrane PtdSer occlusion site
(A) PtdSer occlusion site in detail, viewed from perpendicular to the membrane plane.
Figure is displayed as in Fig. 3C. Phospho-L-serine (stick, green with CPK color) is
modeled according to the observed electron density. (B) Phospholipid conduit along
with TM4. Surface of the atomic model (light blue) is shown with superimposed ribbon
model. Only surface model is clipped by the different plane at the position where TM4
is located, so as to show how the conduit runs along TM4. Its clipped surface is seen as
transparent wheat color. Figure is drawn from TM1 and 2 viewpoint, with exoplasmic
side-up. (C,D) V_{max} and $K_{0.5}$ for indicated mutants determined from their PtdSer- or
PtdEtn-dependent ATPase activities, as described in Fig. 3. (E) The PtdSer flippase
activity. *ATP11A-ATP11C* knock-out (*DKO*) T-lymphoma cells expressing wild-type
ATP11C or indicated mutants were incubated with 1 μM NBD-PS for 3 min.
Experiments were performed two or three times, and flippase activity for NBD-PS is
shown as a percentage to that of wild-type ATP11C. Horizontal red bars denote
averages.

762
763
Fig. 5. A phospholipid transport model for ATP11C
(A) Schematic drawing of the transport mechanism by ATP11C-CDC50A complex. In
the outward-open E2P state (present structure), phospholipid enters from the surface of
outer leaflet to the exoplasmic cavity by changing its orientation (upper left). Trapped
phospholipid head group diffuses along the membrane crevice with its hydrophobic tails
extending out to the hydrophobic core of the bilayer (upper right). When the
phospholipid head group reaches Val357 in the middle of the membrane, the phosphate
head group is occluded by closing the crevice with the TM1-2 helix bundle (lower right).
This process is coupled with dephosphorylation of E2P as seen in the ATP8A1 E2-P_i
transition state structure. Complete dephosphorylation may further open the cytoplasmic
gate, and phospholipid is translocated to the cytoplasmic inner leaflet (lower left). (B)
Close-up view of the membrane crevice indicated as dotted box in A. Phospholipid
headgroup traverses from positively-charged exoplasmic cavity to the occlusion site
near Val357 along with TM4.
778
779

Fig. S1. Structural determination of ATP11C-CDC50A complex
(A) Construction of TP11C and CDC50A used in crystallization. See Methods for

details. (B) Purification of ATP11C-CDC50A complex. Lane 1: solubilized cell lysate,
lane 2: pass-through of Flag resin, lane 3: wash fraction, lane 4: elution by Flag peptide
(subjected to ATPase assay), lane 5: TEV proteinase- and endoglycosidase-treated
sample, lane 6: pass-through fraction of Ni-NTA and amylose resin, lane 7: concentrated
peak fractions by size-exclusion chromatography (arrowhead in the left panel).
Arrowheads on the right indicated as follows, a: HSP70, b: EndoHf, c: cleaved eGFP, d:
TEV proteinase. The elution profile of ATP11C-CDC50A complex by size-exclusion
column chromatography is shown on the left. (C) Representative X-ray diffraction
obtained from a plate-like crystal shown in D. Diffraction spots better than 3.6Å were
obtained along the c^*-axis, whereas these are limited to around 4~6Å in directions along
with a^*- and b^*-axes, thus strongly anisotropic. (D,E) Three-dimensional crystals
obtained from the samples purified from Expi293 cells, showing thin, but large
plate-like crystals (D). In contrast, small crystals were obtained from CDC50A-KO cells
(E). (F) Data acquisition strategy. We employed normal type LithoLoops for helical
scan data acquisition from large single crystals. However, because crystals showed
strong anisotropy, we also collected data sets by irradiating X-ray beam from the
direction perpendicular to the c-axis by using 90º bent type LithoLoops. For the small
but well diffracting crystals obtained from CDC50A-KO cells, data from each
individual crystals was collected for 10º. All of these crystals showed identical unit cell
size and symmetry regardless of crystal morphologies and expression cell types, as seen
in the histograms of unit cell dimensions (G). All diffraction data from 1,588 crystals
were finally merged into a single data set (H), and used for the molecular replacement.

806
Fig. S2. Crystal packing.
(A-C) An asymmetric unit contains four protomers (A, B, C and D) shown in blue,
green, yellow and red, respectively. Their symmetry-related molecules are shown in
light colors. Yellow boxes indicate unit cells view from different direction in A-C as
indicated in the figures. (D) Molecules in the asymmetric unit are displayed according
to their temperature factors. Colors gradually change from blue (16) to red (283). (E,F)
Comparison of the molecular conformation of four protomers. Arrowheads indicate loop
structures, the conformations of which are variable among the four protomers. (G)
$2F_o$-F_c electron density maps of the four protomers at the same contour level of 1.5σ.

821
822
823 Fig. S3. Electron density maps
824 Surface represents $2F_o$-F_c electron density maps of the indicated regions with 1.5σ
825 contour level. Color code as in Fig. 1.

Fig. S4. Sequence alignment.
Primary sequences of indicated P4-ATPases isoforms were aligned using software
MAFFT ver.7[35] and large gaps introduced in the non-conserved region were manually

edited. Cytoplasmic domains (A, pink; P, green; N, light blue), secondary structures
(α-helices, β-sheets and TM helices) and mutations introduced for the crystallized
construct (ΔN7, ΔC38) are indicated above the alignment. The degree of conservation
among evaluated sequences is indicated below the alignment. Acidic, basic, hydrophilic
or hydrophobic amino acids are indicated as red, blue, green and black characters,
respectively. Gene and protein ID is as follows; human ATP11C
(NCBI:_XM_005262405.1), human ATP11A (Uniprot: P98196-1), human ATP8A1
(UniProt: Q9Y2Q0-2), bovine ATP8A2 (Genebank: GQ303567.3), and Saccharomyces
cerevisiae Drs2 (UniProt: P39524-1).

Fig. S5. Cytoplasmic domains
(A-D) Comparison of the relative orientation of the cytoplasmic domains viewed along
with the membrane plane. Cytoplasmic domains of ATP11C are shown in worm models
with the same colour code as in Fig. 1C (A). Atomic models of Drs2p in E2P activated
form (B, grey), ATP8A1 E2P form (C, orange) and ATP8A1 E2-P_i form (D, yellow) are
superimposed on the ATP11C structure according to their P domain structures to show
relative orientations of A and N domains. (E-G) Azimuthal position of the A domain is
compared. A domain of Drs2p (E), ATP8A1 E2P state (F) and E2-P_i state (G), and these
models are superimposed on the ATP11C structure (only A domain is highlighted in
magenta, and others are shown in transparent colors) as in A-D. Dotted arrow indicates
the different azimuthal positions between ATP11C E2P state and ATP8A1 E2-P_i state.
Phosphate analogue BeF$_x$ (red spheres) and DGES/T motif (sticks) in ATP11C structure
are highlighted in all figures.

Fig. S6. Comparison of membrane crevices
(A-D) Surface representation of the atomic models of ATP11C E2P state (A), Drs2p
E2P activated form (B), ATP8A1 E2P (C) and E2-P_i state (D). PtdSer and phosphoserine
are indicated as spheres. Brown background indicates approximate location of the lipid
bilayer. (E-H) Comparison of the TM helix arrangement in ribbon representation.
Atomic models of ATP11C (color codes as in Fig. 1), Drs2p (light grey) and ATP8A1
(tan) are aligned according to their TM helices (E,F). ATP8A1 E2-P_i transition state
(light green) is also compared with ATP11C E2P state (G,H) Only catalytic subunits are
shown in the figure, viewed from the exoplasmic side (E,G) or perpendicular to the
membrane plane with exoplasmic side up (F,H). Phosphoserine (sticks) and Pro94
(spheres) in ATP11C, and PtdSer occluded in ATP8A1 E2-P_i state (sticks) are indicated
in the figure.

Fig. S7. Phospholipid-dependence of ATPase activity for mutants.
ATPase activities of indicated mutants are plotted as a function of DOPS (left) or POPE (right) concentration. Mutants are categorized as follows; cytoplasmic gate (1st row), surface of the membrane cleft (2nd row), occlusion site (3rd row), TM3-4 loop at exoplasmic cavity and residues in TM5 and 6 (4th row), and CDC50A exoplasmic domain facing the cavity (5th row). ATPase activity for the wild-type enzyme is shown in all graphs as a control (black lines).

	ATP11C E2P
Data collection	
Resolution (Å) [†]	4.7 × 4.2 × 3.9 (4.0 − 3.9)[‡]
Space group	$P\,2_1 2_1 2_1$
Cell dimensions	
a, b, c (Å)	100.46, 232.82, 498.89
α, β, γ (°)	90, 90, 90
R_{merge}	1.141 (−)[*]
R_{pim}	0.0419 (−)[*]
$I/\sigma I$	15.88 (0.21)
$C/C1/2$	0.92 (0.864)
Completeness (%)	71.32 (11.64)
Redundancy	776.9 (698.5)
Refinement	
Resolution (Å)	50 − 3.9 (4.0 − 3.9)
No. of reflections	82706647 (7342288)
R_{work}/ R_{free} (%)	27.9/34.8 (26.8/32.6)
Wilson B-factor	98.19
No. of atoms	45171
Protein	44727
Ligands	444
Average B-factor	163.19
Protein (Å^2)	163.10
Ligands (Å^2)	172.65
R.m.s deviations	
Bond lengths (Å)	0.007
Bond angles (°)	1.35

Table S1. Data collection and refinement statistics

[†]The diffraction data are anisotropic. The resolution limits given are for the *a**, *b** and *c** axes, respectively.

[‡]Statistics for the highest-resolution shell are shown in parentheses.

[*]Statistics for the highest-resolution shell are not given due to the strong anisotropy.