Unraveling nanotopography of cell surface receptors

Christian Franke[1,6,*], Tomáš Chum[2,6], Zuzana Kvíčalová[2], Daniela Glatzová[2,3], Alvaro Rodriguez[4], Dominic Helmerich[5], Otakar Frank[4], Tomáš Brdička[3], Sebastian van de Linde[6,8,*] and Marek Cebecauer[2,8,*]

[1] Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany

[2] Department of Biophysical Chemistry, J. Heyrovsky Institute of Physical Chemistry, Czech Academy of Sciences, Prague, Czech Republic

[3] Laboratory of Leukocyte Signalling, Department of Institute of Molecular Genetics, Czech Academy of Sciences, Prague, Czech Republic

[4] Department of Electrochemical Materials, J. Heyrovsky Institute of Physical Chemistry, Czech Academy of Sciences, Prague, Czech Republic

[5] Department of Biotechnology and Biophysics, Biocenter, University of Würzburg, D-97074, Würzburg, Germany

[6] Department of Physics, SUPA, University of Strathclyde, Glasgow, United Kingdom

[7] These authors contributed equally: Christian Franke and Tomáš Chum.

[8] These authors contributed equally: Sebastian van de Linde and Marek Cebecauer.

* Corresponding authors: Ch.F. cfranke@mpi-cbg.de, S.v.d.L. s.vandelinde@strath.ac.uk and M.C. marel.cebecauer@jh-inst.cas.cz

ABSTRACT

Cells communicate with the surrounding environment via receptors on their surface. However, it remains unclear how are receptors organized with respect to the complex cell surface morphology. This is mainly due to a lack of accessible, robust and high-resolution methods. Here, we present an approach for mapping the topography of receptors at the cell surface with nanometer precision. The method includes coating of glass coverslips with a glycine hydrogel which preserves the fine membrane morphology while it allows positioning of immobilized cells close to the optical surface. We further developed an advanced and simplified algorithm for the analysis of single molecule localization data acquired in a biplane detection scheme. These advancements enable direct and quantitative mapping of protein distribution on ruffled plasma membrane with near isotropic 3D nanometer resolution. By using CD4 clusters on unstimulated T cells as an example, we demonstrate that nanoclusters observed by 2D restricted methods represent accumulation of this receptor at the tips of membrane protrusions. On the contrary, CD45, which was shown to segregate from T cell nanoclusters, localizes to the membrane basis, thus, avoiding contact with CD4 on the tips. In summary, we provide a simple and straightforward workflow to study molecules and their relations at the complex surface nanomorphology of differentiated metazoan cells.

INTRODUCTION

Supramolecular complexes drive numerous vital processes in cells, such as gene expression, molecular transport or signal transduction. Cellular membranes provide an excellent platform to assemble molecules into complex structures. Indeed, membrane-associated molecules, including surface receptors, were found to form clusters with nanometric dimensions.[1, 2, 3, 4, 5] Lateral interactions and/or actin-cytoskeleton anchorage drive clustering of some receptors (e.g. tetraspanin platforms, focal adhesion complexes). However, the mechanism of cluster assembly for other receptors remains unknown and often controversial (e.g. receptor microclusters on lymphocytes). Indeed, receptor clustering is the subject of intense debate.[6, 7, 8] Several works provided the evidence about the monomeric character of receptors, which were shown to cluster using different experimental setups.[9, 10, 11] Thus, understanding their origins can help to unravel the very existence of receptor clusters. In this work we focus on receptor clustering on the plasma membrane of lymphocytes, but the method and general principles discussed herein apply to the surface of any cell, including those of prokaryotes.

A common feature of supramolecular structures, including membrane receptor clusters, is their small size (< 200 nm), which falls in a range inaccessible by standard light microscopy. Therefore, molecular assemblies in cells are often studied indirectly. For example, total internal reflection fluorescence (TIRF) microscopy enabled to visualize dynamic receptor microclusters in B and T cells.[12, 13] It was the lateral mobility of these entities, which indicated the very existence of clusters in these cells. The size and shape of the observed clusters were irresolvable because the spatial resolution of TIRF microscopy is diffraction limited. Later, super-resolution (SR) microscopy techniques were developed to overpass the diffraction limit and offered a more detailed insight into the architecture of cell receptor clusters (for example refs: [5, 14, 15, 16]).

Currently, single-molecule localization microscopy (SMLM) is the method of choice to study organization of membrane receptors due to its ability to localize emitters (i.e. labelled receptors) with nanometer precision and its potential for quantitative assessments of protein distribution and number.[17, 18, 19, 20] SMLM-based studies confirmed the earlier observations of TIRF microscopy that receptors and associated signaling molecules can cluster in non-stimulated and stimulated immune cells.[21, 22, 23, 24, 25] However, there is an intense discussion about the feasibility of SMLM for the cluster analysis of membrane molecules.[26, 27, 28] Recently, procedures aiming to minimize the impact of methodological artefacts on the SMLM results were developed.[29, 30, 31]

In most of these studies, the SMLM methods were used to generate localization maps by projecting the presumably three-dimensional receptor distribution onto a two-dimensional plane. The precise information about the axial position of emitters (e.g. receptors), and therefore their distribution, is missing. This was an accepted trade-off due to the limited axial resolution of the applied SR methods (including STED and SIM; ref. [32]). Moreover, the plasma membrane is routinely depicted as a rather featureless structure (for example: refs. [33, 34]). Yet differentiated cells of vertebrates are densely covered with membrane protrusions and

invaginations, thus, resulting in a highly three-dimensional nature of their surface.[35] Scanning electron microscopy micrographs demonstrated that finger-like membrane protrusions reminiscent of microvilli dominate the surface of immune cells.[36, 37, 38] Microvilli are dynamic structures with ~100 nm diameter and varying length between 0.5-5 μm.[39] Although less well comprehended on immune cells, tips of microvilli have a potential to accumulate membrane receptors in domains with a diameter ~100 nm in analogy to signaling receptors and channels on epithelial and sensory cells.[40]

Electron microscopy (EM), with its ability to provide information about ultrastructural details, was the key method for characterization of microvilli in the pioneering works.[41] However, EM cannot visualize specific proteins efficiently, since the labelling densities are limited by ligand/antigen accessibility, steric hindrance and electron repulsion.[42, 43, 44] A rapid development of SR light microscopy techniques enabled the visualization of three-dimensional objects with high precision, with the use of highly specific and frequently efficient labelling methods and on living cells (e.g. refs. [45, 46]). However, to characterize receptor distribution on nanometric membrane structures such as microvilli, it is important to develop three-dimensional SR techniques with axial resolution well below 50 nm. Several SMLM based methods were suggested,[47] such as astigmatism,[48] multiplane[46] or biplane imaging,[49, 50] double-helix point spread function[51] or interferometric PALM.[52]

Whereas microscopy techniques with dramatically improved spatial resolution recently became a reality, the (cell) sample preparation for SR imaging still suffers from several caveats. The spatial resolution offered by SMLM comes with the trade of time resolution since thousands of camera frames are needed to render a map, sufficiently representing, for example, a receptor distribution on a cell.[53] Thus, the movement of imaged objects (e.g. proteins) has to be minimized. This is usually accomplished by fixation of cells prior to imaging. Importantly, cells grown or isolated initially in suspension must be attached to the optical surface (e.g. coverslips). Poly-*L*-lysine (PLL; or its isomer poly-*D*-lysine) is used for immobilization of suspension cells in standard protocols.[54] However, attachment to a surface via interaction with PLL leads to the deformation of cells.[55] Similarly, the flattening of cell surfaces is observed when positioning immune cells on adhesion molecules (e.g. ICAMs) directly coated on coverslips or linked to a supported planar lipid bilayer.[56, 57] The use of hydrogel or Matrigel enables to stabilize cells for imaging without observable morphological changes.[58, 59] However, these materials submerge cells into a 3D matrix which leads to their random axial distribution. Single-molecule fluorescence microscopy techniques (including SMLM) depend on an efficient transmission of photons from emitters and, thus, provide the best results for molecules close to the optical surface.[47] Currently, none of the available methods can immobilize cells close to the optical surface without interfering with their complex surface morphology.

CD4 is a surface glycoprotein involved in T cell development and function. It is associated with Lck kinase which drives antigen-specific T cell activation.[60, 61] CD4 also mediates T cell adhesion to the cells which express major histocompatibility complex class II (MHCII) proteins

on their surface. We have recently shown that CD4 accumulates in clusters on the surface of non-stimulated T cells in a palmitoylation-dependent manner.[28] The origin of these CD4 clusters remains unknown. CD45, another glycoprotein that is three times larger than CD4, is expressed at the surface of all lymphocytes.[62] Its intracellular phosphatase domain regulates Lck kinase activity. CD45 was found to segregate from signaling molecules in activated T cells, as predicted by the kinetic segregation model of T cell activation.[63, 64] However, the mechanism responsible for such segregation is incompletely understood due to a lack of information about its precise localization at the surface of immune cells.

Here, we describe a new method for mapping receptors at the nano-topography of the plasma membrane with near isotropic three-dimensional resolution. The workflow involves a newly developed, optimized coating of glass coverslips which preserves the complex morphology of a cell surface while it allows for the positioning of immobilized cells close to the optical surface. Moreover, we adapted the recently reported TRABI method[65] for nanometer precise three-dimensional SMLM imaging to improve data acquisition, processing and drift correction, as well as the integrity of the three-dimensional SMLM data set. This allowed us to employ a straightforward quantitative assessment of the axial distributions of selected cell surface receptors at the nanoscale. We applied this method to reveal the three-dimensional nature of CD4 receptor clusters in non-stimulated T cells. We further uncovered the importance of membrane protrusions for the segregation of CD45 phosphatase from CD4 involved in T cell signaling. The presented method enables molecular studies of cell surface nano-morphology but also of other fluorescent nanoscopic structures.

RESULTS

Preservation of membrane morphology during cell immobilization on coverslips

In order to map receptor distribution at the complex surface of cells and at the nanoscale, the experimental setup must include quantitative labelling of target molecules, an appropriate sample preparation and a nanometer precise three-dimensional imaging method. In this work, labelling of surface receptors was performed by standard immunofluorescence protocols using directly labelled, highly specific antibodies.[66, 67]

Cells grown or isolated in suspension, e.g. T cells, must be immobilized on the optical surface (e.g. coverslips; Fig. 1a-b) in order to facilitate imaging approaches, which require long acquisition times. This is true for most SMLM techniques.[68] Commonly, poly-*L*-lysine (PLL; or poly-*D*-lysine) coating of glass coverslips is used to adhere suspension cells to the optical surface.[55] Negatively charged biomolecules at the cell surface electrostatically interact with the polycationic layer formed by PLL on coverslips.[69, 70] However, such electrostatic interaction can deform the cell surface (Fig. 1a; ref. [55]). Indeed, we have observed a rapid flattening of a T cell surface upon settling to the PLL-coated coverslip, visualized by TIRF microscopy of CD4-GFP fusion protein (Fig. 1c). With small exceptions (red arrowhead), a homogenous distribution of CD4-GFP signal dominated the attached T cell surface already 60

seconds after the first detectable contact with the optical surface (blue arrowheads; and Supplementary Fig. S1a). This is even more pronounced in cells observed 2 and 4 minutes after the first contact. Continuous and homogenous fluorescence signal of CD4-GFP indicates a lack of membrane accumulation in protrusions and invaginations. These data indicate that immobilization on PLL-coated coverslips is responsible for flattening of the T cell surface and, probably, for the variation in the number and size of CD4 clusters detected in our previous work.[28] Moreover, when analyzed by SMLM, diverse distribution patterns of CD4 can be found on T cells immobilized on PLL-coated coverslips. Only few cells have CD4 accumulated in high density regions (Supplementary Fig. S2a), a majority of cells exhibits a random distribution of this receptor (Supplementary Fig. S2b). We hypothesize that the pattern of CD4 distribution in these cells is defined by the time since their first contact with the PLL-coated surface. Of note, during the sample preparation, cells cannot be synchronized with respect to their first contact with the coverslip.

Since, in this work, we were interested in the origin of CD4 clusters on a rough T cell surface, we required an alternative method for coating of coverslips which better preserves their cellular morphology. Coating of an optical surface with glycine was previously used to minimize non-specific background signal in single-molecule fluorescence imaging.[71, 72] However, when applied without prior PLL treatment, glycine forms a hydrogel-like layer on a glass surface (Supplementary Fig. S3a). Our atomic force microscopy (AFM) measurements indicated a continuous surface coating with glycine (Supplementary Fig. S3b). However, the AFM measurements were limited by the softness of the glycine layer and a tendency of this material to adhere to the AFM tip. We were thus unable to determine the exact thickness and stiffness of the glycine layer. The few deformations observed in the tested samples indicated that the average thickness of glycine layer is at least 15 nm (Supplementary Fig. S3c). We further observed crystals on coverslips coated with glycine that were subsequently dried, which supports the notion of a glycine hydrogel. On the contrary, no such precipitate was observed on dried PLL-coated coverslips (Supplementary Fig. S3d). We thus conclude that glycine forms a gel-like structure on the glass surface, which functions as a semi-soft cushion upon cell landing.

To demonstrate the impact of glycine coating on cellular morphology, we imaged T cells, expressing CD4-GFP, during landing on coverslips using TIRF microscopy. In contrast to the PLL-coated samples, highly heterogeneous CD4-GFP signal was detected in cells immobilized on glycine-coated coverslips (Fig. 1d). Intensity line profiles measured across the surface of such T cells exhibit large deviations 1, 2 and 4 minutes after their first detectable contact with a coated coverslip (Supplementary Fig. S1b). These data clearly indicate that glycine-coating allows immobilization of cells close to the optical surface and improves the preservation of cellular morphology when compared to PLL-coating (Fig. 1a).

We further examined the impact of the newly developed coating method on the resting state and viability of immobilized T cells. PLL was reported to induce calcium response in cells attached to the coated surface.[73] Our measurements confirmed that PLL induces calcium

mobilization in unstimulated T cells, albeit much lower compared to antigenic stimulation mimicked by the coverslip coated with specific antibodies (anti-CD3ε) (Fig. 1e and Supplementary Fig. S4). In turn, the immobilization of T cells on glycine-coated coverslips stimulated only a negligible calcium response (Fig. 1e and Supplementary Fig. S4). Similarly, longer cultivation (> 20 min) of T cells on PLL-coated coverslips negatively influenced their viability (Fig. 1f). During the incubation of T cells on the glycine-coated coverslips, we observed much lower effect on the viability of cells compared to the sample on PLL. In summary, our findings indicate that coating of optical surfaces with glycine reduces the stress generated by the charged surface of, for example, PLL-coated coverslips. Moreover, the gel-like structure of the glycine coating preserves the surface morphology of immobilized cells as demonstrated using TIRF (Fig. 1c,d) and allows studying the three-dimensional receptor distribution on the surface of resting T cells with SMLM.

The three-dimensional SMLM method.

In order to capture the entire nanoscopic three-dimensional organization of receptors at the cell surface, we acquired image sequences with a highly inclined and laminated optical sheet (HILO) illumination, which enables single-molecule detection in cells much deeper than in TIRF microscopy (Fig. 2a). At the same time, it restricts the out-of-focus fluorescence from the remaining parts of a cell, thus improving the local signal-to-noise ratio.[74] Prior to the acquisition, we briefly irradiated the whole cell in epifluorescence to further minimize out-of-focus contributions (see **Methods**).

Intensity based biplane SMLM imaging - dTRABI. Previously, we reported the temporal, radial-aperture-based intensity estimation (TRABI) method in combination with a biplane emission detection mode as a powerful three-dimensional imaging tool with nanometer precision.[65, 75] In short, TRABI comprised fitting single-molecule spots by a Gaussian function with an invariant width, a subsequent independent photometric analysis of these spots and a final allocation of the axial coordinate based on a calibration curve. The photometric value was determined from the intensity obtained by the established Gaussian fitting procedure (I_{Fit}) and background-subtracted reference intensity of the spot (I_R), with $I_{Fit} < I_R$.

Here, we used a simplified but superior version of TRABI method where solely the intensity information obtained by the Gaussian fit (I_{Fit}) was used as metric for both channels in a biplane imaging scheme. Synchronized image stacks of the two channels (transmitted and reflected), generated by dividing the emitted fluorescence equally with a 50/50 beam-splitter, were analyzed by the same Gaussian function with an invariant width for every spot. Spots in the reflected channel were mapped to the corresponding spots in the transmitted channel by the nearest neighbor algorithm (Fig. 2b, see **Methods**). By omitting the standard, photometric analysis,[65] the computation time was reduced and the allowed number of localizations per frame was significantly increased. We therefore reckon, that we achieve an improved structural resolution while maintaining the same axial localization precision as the original

TRABI approach (Supplementary Fig. S5). We subsequently called the new method dTRABI (for *direct* TRABI; Fig. 2c).

Fiducial-free drift and tilt correction. In order to ensure optimal axial localization over an extended imaging time as well as accounting for subtle sample drift, we developed a fiducial free approach for its correction (see **Methods**). In short, the axial footprint of the entire structure is tracked over time, resulting in a spatio-temporal drift trace (Fig. 3). This trace is fitted with a high-order polynomial, which serves as the correction term for the raw localizations by linearization. We reckon that in a thin sample layer like the plasma membrane the spatio-temporal distribution of active photo-switches that reside in their on-state is constant over time. The selection of appropriate regions in axially more extended samples can include structures, where the local z-dimension is restricted. Most data sets that were used to conclude the results in the following paragraphs showed axial drift with different orders of magnitude and non-linearity, which could be accounted for by the described correction approach. However, an additional linear axial tilt was observable in some data sets, most likely due to minor imperfections of the sample rest. Since we assume the layer closest to the coverslip to be axially flat on the whole-cell scale, we fitted a three-dimensional plane to the raw image, thereby determining its gradient and thus enabling the linearization of the raw localization data (see **Methods** and Fig. 3).

Quantitative axial localization analysis. For the analysis of the axial receptor distribution, localization files were loaded and processed in Fiji with custom written scripts (see **Methods**).[76, 77] The analysis comprised three steps: generation of quantitative image stacks, segmentation in regions of interest (ROIs) and axial quantification of individual ROIs. (i) A quantitative 3D image stack (z-stack) was generated with 20 nm voxel size in *x*, *y* and *z*. (ii) These stacks were then segmented by automatically generating multiple region of interests (ROIs) with an area of 4 µm² within the interior of the cell (Supplementary Fig. S6). (iii) Then the axial distribution of localizations was analyzed, by counting all localizations per ROI along the z-stack. For each ROI the resulting distribution was fitted to a superposition of two Gaussians (Supplementary Fig. S6). By analyzing properties of these fits, we acquired parameters which provide quantitative information about the axial distribution of surface receptors (see **Methods**).

Imaging CD45 nano-topography using the dTRABI approach

To demonstrate the applicability of dTRABI, we labelled CD45 receptors, which are highly expressed on the surface of T cells (> 100.000 copies/cell; ref. [78]) with Alexa Fluor 647-conjugated antibody (MEM-28) and imaged as described above. Cells were immobilized on glycine-coated coverslips to preserve cell surface morphology. Resulting three-dimensional images with color-coded z-axis represent a footprint of a cell on a coverslip (Fig. 4a). The apparent optical depth, which varies between cells (300-650 nm), depends on the available structure in the individual field of view. CD45-labelled cells exhibited a variety of features that

can extend far away from the coverslip. However, the entire extent of the basal membrane resting on the surface was captured in all displayed and analyzed cells. Fig. 4b shows the three-dimensional image of CD45 which indicates a complex morphology of the cell surface. Magnified images and the corresponding *xz*-projections (Fig. 4c) demonstrate the ability of dTRABI to visualize large membrane protrusions at the edge of the cell (ROIs 1 and 3; blue arrowheads), as well as more subtle membrane extensions towards the exterior (ROIs 1-4; red arrowheads) and interior (ROIs 3 and 5; green arrowheads) of the cell.

The origin of CD4 nanoclusters on the surface of unstimulated T cells

Next, we examined the origin of previously reported receptor (signaling) microclusters.[1, 4, 5] Using two-dimensional SR imaging, we recently showed that CD4 receptors accumulate in high density regions on unstimulated T cells.[28] The three-dimensional organization of these clusters remained unresolved. In this work, we employed our new coverslip coating strategy and dTRABI to study CD4 clusters with near isotropic three-dimensional resolution in immobilized cell with preserved surface nanotopography. To verify the specificity of these structures, we also imaged CD45, which was reported to distribute on the surface of non-stimulated T cells more evenly.[79] To avoid visualization of intracellular molecules, fixed T cells were labelled with specific antibodies in the absence of membrane permeabilization.

First, we confirmed that CD4 accumulates in high density regions in palmitoylation-dependent manner using 2D *d*STORM with high lateral resolution (Supplementary Fig. S7). Next, three-dimensional high-resolution maps of the CD4 surface distribution were acquired by analyzing the same data sets with the dTRABI approach (Fig. 5a). These results provide detailed information about the nanotopography of CD4 at the surface of unstimulated T cells with axial localization precision of 10-20 nm (Supplementary Fig. S5). The nanoscopic three-dimensional view of wild-type CD4 indicates that the receptor is preferentially localized to one topographical level and is clustered (Fig. 5a). Plotting a relative z-position of CD4 molecules in the central area, which avoids cell edges, indicates the accumulation of wild-type CD4 in a narrow space near the coverslip (Fig. 5b and Supplementary Fig. S8). On the contrary, a broad surface distribution is evident from the three-dimensional view of a CD4 variant which cannot be palmitoylated and was previously shown to exhibit random distribution in such unstimulated T cells (CD4 CS1; Fig. 5c; ref. [28]). Our quantitative analysis demonstrates a less constrained distribution of receptor z-positions (Fig. 5d). The plot of averaged z-positions over all tested cells (21 CD4 WT and 18 CD4 CS1 cells) confirms the different axial distribution of the two CD4 variants on T cells (Supplementary Fig. S8). However, the contrast between the two variants is strongly reduced due to a variation in surface complexity of individual cells and distinct regions of the basal membrane. Therefore, the propensity of CD4 molecules to accumulate at the specific topographical levels was first analyzed in square regions of interest (ROIs) of 2×2 µm^2 selected to cover the cell-coverslip contact area over all tested cells (846 CD4 WT ROIs and 1044 CD4 CS1 ROIs in total; Supplementary Fig. S6). As mentioned above, for quantitative analysis, relative z-positions for the localizations in ROIs were fitted with a

sum of two Gaussians (Fig. 5e). From these fits, three parameters were derived, i.e. the spread of the two Gaussians termed z-distribution width (Fig. 5f-g), the peak-to-peak distance (Supplementary Fig. S9) and the width difference of the two Gaussians (Supplementary Fig. S10). These parameters better describe the axial distribution of receptors compared to the plots with the averaged relative z-positions of localizations. Indeed, a very narrow z-distribution width in the range of $90 - 150$ nm in all segments demonstrated strong tendency of wild-type CD4 to accumulate at the specific topographical level in resting T cells (Fig. 4f-g). On the contrary, a lack of palmitoylation in CD4 CS1 variant caused much broader distribution of this receptor on T cell surface, exhibiting a z-distribution width predominantly in the range of $110 - 170$ nm and with a substantial fraction between 150 and 340 nm. The difference between the widths of the two Gaussians is also more pronounced for CD4 CS1 (80 nm instead of 50 nm for CD4 WT) (Supplementary Fig. S9c).

The specificity of CD4 receptor clusters was further emphasized by a less constrained lateral and axial distribution of CD45 in T cells imaged using dTRABI (Fig. 5h,i; 13 cells and 305 ROIs were analyzed). Quantitative analyses performed consistent with the CD4 variants demonstrated a broader axial distribution of CD45 receptor compared to CD4 WT (Fig. 5f-g). We found the range of z-distribution widths to be similar around 90 and 150 nm, but in contrast to CD4 WT, CD45 has an additional large fraction between 150 and 400 nm. This broad CD45 distribution is particularly noticeable when analyzing the peak-to-peak distance of the fitted bimodal distribution to the relative z-position of the receptor (Supplementary Fig. S9b, see **Methods**). The obtained peak-to-peak distances were exponentially distributed and the mean values for CD4 WT and CS1 were 23 and 36 nm, respectively, whereas CD45 was more broadly distributed around 120 nm. More examples of 3D dTRABI images and their quantitative analysis for all three receptors are shown in Supplementary Fig. S10 and Supplementary Fig. S6, respectively.

Segregation of CD45 from CD4 clusters in non-stimulated T cells

When analyzing dTRABI images, we noted areas with a very low number of CD45 localizations (Fig. 6a; yellow arrowheads). Similarly, CD45 exclusion zones were detectable in 2D SOFI images of antibody-labeled T cells (Fig. 6b). The narrow shape of these exclusion zones was reminiscent of CD4 clusters observed on T cells imaged by dTRABI approach (Fig. 5a and Supplementary Fig. S10). We thus wondered whether the CD45 exclusion zones represent areas with CD4 accumulation. T cells expressing CD4 WT fused to photo-switchable protein mEos2 (ref. [28]) were labelled with specific anti-CD45 antibodies and analyzed using two-color SMLM. The data indicates that CD4 and CD45 essentially segregate into two separate zones on the surface of non-stimulated T cells immobilized on glycine-coated coverslips (Fig. 7a,b). Since CD4 clusters to one topographical level near the T cell-coverslip contact sites (Fig. 5a,b,f), we reckon that CD4 preferentially accumulates at the tips of membrane protrusions (e.g. microvilli) and CD45 to the shaft of these structures and their base at the plasma membrane (Fig. 7c). Indeed, we observed the accumulation of CD4 at the tips of large T cell

membrane protrusions in several cells (Fig. 7d,e). In turn, the shaft of protrusions was extensively covered with CD45 signal. Often, CD45 was essentially segregated from CD4 in these nanoscopic membrane structures (Fig. 7e).

Having established a new method which preserves the surface morphology and a resting state of immobilized cells, we analyzed segregation of CD4 and CD45 in living T cells forming contact with the glycine-coated coverslips using confocal microscopy (Fig. 8). Large scale (microscopic) segregation of CD4 and CD45 was observed in membranes forming contact with coverslips (Fig. 8; middle and lower panels). No such segregation was detectable in free-standing areas of the plasma membrane. Nevertheless, our two-color SMLM data indicate that CD4 and CD45 segregate on such membranes, however, at the nanoscopic scale (Fig. 7). In summary, our data suggest that CD4 and CD45 largely segregate on non-stimulated, cultured T cells which is most likely caused by their different preference for specific areas on the complex three-dimensional surface of these cells.

DISCUSSION

In this work, we have introduced a new approach for molecular mapping of the membrane receptor topography with nanometer precision. The method combines the use of glycine-coated optical surfaces which preserve membrane morphology of immobilized cells with an advanced and simplified dTRABI algorithm for quantitative, near isotropic three-dimensional SMLM imaging. Using directly labelled antibodies and dTRABI processing of biplane SMLM data, we achieved 10-20 nm localization precision in all spatial directions. This enabled us to reveal the origin of CD4 clusters and the involvement of cell surface morphology in CD45 segregation from signaling receptor clusters.

Our principal task was to preserve fine structures on the surface of T cells and to analyze the distribution of signaling receptors therein. Even though flat in most common illustrations, the plasma membrane of differentiated metazoan cells is a three-dimensional structure which is densely covered with finger-like protrusions (microvilli, filopodia), ruffles and invaginations (endocytic cavities and podosomes).[40] Its imaging with two-dimensional methods thus leads to simplifications and, potentially, misinterpretations. The availability of SR microscopy capable to resolve nanoscopic three-dimensional structures is currently limited to specialized nanophotonics laboratories. Nevertheless, standard methods of sample preparation for imaging (e.g. on PLL-coated coverslips) systematically damage the fragile surface morphology of studied cells, as indicated also by our data showing that the T cell surface complexity is diminished within few minutes since the first contact with a PLL-coated coverslip (Fig. 1c and Supplementary Fig. S2).

Several materials were previously developed to prevent cell deformation during their immobilization in a three-dimensional space, for example, Matrigel and synthetic hydrogels. Matrigel enables three-dimensional cell cultures but it is derived from the extracellular matrix of mouse tumors and exhibits autofluorescence. On the contrary, fluorescence properties of

synthetic hydrogels (e.g. CyGEL Sustain) can be controlled and the cells prepared in hydrogels display numerous protrusions on their surface, including nanoscopic structures which require SR microscopy for their detection.[58] However, TIRF and HILO sample illumination techniques which can reduce the out-of-focus background signal and, thus, improve the quality of SMLM images, enable fluorescence signal detection solely in the vicinity of the optical surface.[74, 80] Hydrogels do not allow efficient positioning of immobilized cells at the optical glass.

Here, we developed and analyzed glycine coating of coverslips which preserves cell membrane morphology. We observed a complex membrane morphology over 10 mins after the first contact of a cell with the optical surface (Fig. 1d). Moreover, a coating of coverslips with a narrow layer of glycine-hydrogel facilitates positioning of immobilized cells close to the optical surface. Importantly, cells immobilized on glycine exhibit improved viability and reduced non-specific stimulation compared to PLL-immobilized cells. Of note, the preparation of glycine-coated coverslips is straightforward, fast (<30 min) and inexpensive. All these properties qualify glycine-coating of coverslips for the advanced imaging of cells, including high-resolution three-dimensional dTRABI mapping of cell surface molecules.

We have also developed the dTRABI algorithm to allow for faster and computationally cheaper processing of SMLM data when compared to our original TRABI approach. It is built for a quantitative analysis of axial distribution of fluorescent molecules. Importantly, dTRABI can be used with any localization software that supports fixed Gaussian fitting in combination with a biplane detection scheme. In contrast to a classical biplane imaging that compares the width of spots evaluated by free Gaussian PSF fitting, which facilitates an improved localization precision. The ability of dTRABI to visualize fluorescent molecules on complex, irregular, three-dimensional cellular structures can be highlighted on a T cell which was not included in the quantitative analysis of a receptor axial distribution (Supplementary Fig. S11). Complex membrane structures formed by this dying cell were effectively labeled and visualized using dTRABI approach.

Importantly, dTRABI offers a tool for mapping localization of molecules within 1000 nm from the optical surface. Such working-space provides a sufficient axial signal penetration to visualize the eukaryotic plasma membrane with its nanotopography. Under the conditions used in our experiments, the plasma membrane protrusions were positioned 100-400 nm from the optical surface (Fig. 5). Thus, molecules which do not localize to the protrusions can be monitored as well. This was facilitated by folding of membrane protrusions, the length of which often exceeds the axial depth of our method, under the cell body (Fig. 4 and 9a,b). Evidently, a partial loss of molecular localizations in the areas where the basis of the plasma membrane is positioned outside of the optical limit of this method is caused by a diverse positioning of cells and intensive bending of cell-glass contacts (Fig. 9c,d and Supplementary Fig. S10).

A quantitative character of dTRABI data also enabled detailed analysis of receptor axial distribution. We modelled the axial receptor distribution by fitting a sum of two Gaussian functions (double Gaussian) to the data. Although CD4 distribution can be modelled by a

single Gaussian to a high degree of confidence, a double Gaussian is even more accurate and accounts for a slight asymmetric organization of receptors. In contrast, non-palmitoylatable CD4 CS1 mutant does not follow a mono Gaussian distribution and the superposition of two Gaussians is essential. The analysis allowed us to derive quantitative parameters describing axial receptor distribution in detail, i.e. z-distribution width, peak-to-peak distance and width difference. All these analytical approaches emphasized narrow and symmetrical distribution of CD4 (z-distribution width ~110 nm, peak-to-peak distance < 25 nm), while CD4 CS1 and CD45 show broader distributions.

After implementing our new imaging approach, we focused on two important biological questions: i) what is the origin of receptor microclusters in T cells, and ii) how membrane morphology affects segregation of signaling receptors into different areas of the plasma membrane. Both questions cannot be studied by previously reported protocols. Using quantitative SOFI analysis, we have previously shown that CD4 accumulates in high density regions (clusters) in non-stimulated T cells by a process which depends on its post-translational lipid modification, palmitoylation.[28] However, as in many other cases, these data represented two-dimensional projections of receptor localizations on the complex T cell surface. Using dTRABI imaging of non-stimulated T cells immobilized on glycine-coated coverslips, we demonstrate that CD4 accumulates on the tips of membrane protrusions reminiscent of microvilli. Thus, CD4 clusters, and potentially other clusters, represent receptors (or other molecules) trapped at the tips of membrane protrusions, stressing, that intact cell surface morphology is essential for the proper organization of signaling molecules. Importantly, the mutant CD4 CS1 variant which cannot be palmitoylated, distributed more randomly over the complex surface of T cells. This is in agreement with our previous observation that CD4 CS1 does not accumulate in detectable structures on these cells.[28] Targeting to or stabilizing proteins on membrane protrusions thus may be another, previously unreported role of protein palmitoylation.[81]

The best described function of CD4 is to deliver critical Lck kinase to the proximity of T cell receptor for a rapid response to antigens.[60] TCR was recently reported to accumulate in T cell microvilli by projecting a two-dimensional SMLM image onto the morphological map generated by varying-angle TIRF microscopy (VA-TIRFM) of a membrane dye.[82] VA-TIRFM is diffraction-limited and, therefore, does not provide a high-resolution map of cell surface nanotopography. However, the data indicated the accumulation of TCR in domains formed by or dependent on membrane morphology. We thus speculate that CD4 and TCR localize to the protrusions, the tips of which can represent membrane areas with concentrated signaling molecules arranged to rapidly respond to appropriate signals (Fig. 7c). How these molecules accumulate in these structures and whether the tips of protrusions undergo a further compartmentalization remains unclear. In future ventures, we will establish a robust workflow consisting of a reliable sample preparation, labelling and multi-color dTRABI analysis, in order to investigate multi-molecular architecture of these structures. Our current

approach already provides a good basis for such advancement and is adaptable to further questions.

As discussed earlier, CD4 accumulates in clusters at the tips of membrane protrusions. However, several molecules, including CD45, were shown to segregate from signaling microclusters (e.g. refs. [63, 83]). Using dTRABI and glycine-coated coverslips, we demonstrate that CD45 distributes over a broader axial spectrum compared to CD4. Since CD45 segregates from CD4 in non-stimulated T cells under the experimental conditions used in this study, we conclude that CD45 segregates from signaling molecules by localizing to the basis of the plasma membrane and by avoiding highly curved membranes of the protrusions (Fig. 7c). The molecular mechanism of such segregation remains unknown. Even though listed in a high-throughput study of palmitoylated proteins,[84] a direct experimental proof of CD45 palmitoylation is missing. Since non-palmitoylated CD4 CS1 variant exhibits a more random distribution compared to its native variant, we speculate that the lack of CD45 palmitoylation contributes to its preference for the basis of the plasma membrane. We also demonstrate that CD45 microscopically segregates from the contact site of non-stimulated T cells with a glycine-coated surface. No such segregation is observed for CD4. This suggests that CD45 reacts to mechanical forces by segregating into membranes areas unaffected by force. Nevertheless, nanoscopic segregation caused by complex membrane morphology was observed also in membranes unaffected by cell attachment to the optical surface.

In summary, we provide a new and undemanding workflow to study nanotopography of receptors at the cell surface. We highlight the importance of appropriate sample preparation for the imaging of three-dimensional structures. The improved algorithm of dTRABI exemplifies an easy to implement and high-quality method to generate three-dimensional SMLM data. Finally, we demonstrate the applicability of the workflow by answering two important questions related to the localization and function of surface receptors on lymphocytes. Even though used here for human receptors, the method can be implemented for characterization of molecules on the surface of other organisms (e.g. yeast, plants) but also for complex nanomaterials which do not exceed the current axial penetration of dTRABI (~ 1000 nm).

METHODS

Cell culture and transfection

Jurkat T cells (clone E6; ATCC) and their CD4-knock-out variants (Jurkat CD4-KO cells) were grown in RPMI-1640 medium (Gibco) supplemented with 10% fetal bovine serum (Gibco), Non-essential amino-acids (Gibco) and 10mM HEPES under controlled conditions in a humidified incubator at 37°C, and 5 % CO_2 (Eppendorf). Jurkat CD4-KO cells were derived from wild-type Jurkat T cells (clone E6; ATCC) using CRISPR/Cas9 technology as described.[85] Both cell lines were regularly tested for morphology and mycoplasma infection.

For the expression of exogenous proteins, Jurkat T cells and their CD4 KO variant were transiently transfected with plasmid DNA using Neon® transfection system (Thermo Fisher Scientific) according the manufacturer's instructions. Briefly, 1 μg of vector DNA was used per 200 000 cells in 0.5 ml culture medium. The instrument settings were: 3 pulses, each 1350 V for 10 ms. Transfected cells were used within 36 h since the transfection.

Sample preparation for SR microscopy

Cleaning and handling of coverslips and solutions for microscopy

High precision microscopy coverslips (round, 25 mm in diameter; Marienfeld) in Teflon holders (Wash-N-Dry Coverslip Rack; Diversified Biotech Inc.) were cleaned by incubation at 56°C overnight in 2% Hellmanex III (Hellma Analytics) dissolved in ultrapure Milli-Q® water (Millipore) followed by 30 minutes sonication in a heated sonication bath. Several washes in ultrapure water and one additional sonication step were used to remove all traces of Hellmanex III components. Cleaned coverslips were stored in ultrapure water to avoid drying and contamination with particles of a dust from air.

All other glassware was regularly treated with piranha solution (3 parts of 30% hydrogen peroxide and 7 parts of concentrated sulfuric acid) for 45 minutes followed by several washes with ultrapure water. All solutions were made from concentrated stocks stored in glass containers, diluted with ultrapure water and filtered using syringe filters with 0.22 μm pores (TPP) into piranha-treated glassware.

Glycine coating of coverslips

Ultraclean coverslips were coated by applying 0.5 ml of 2M glycine solution and incubation for 20 min at room temperature in the laminar flow box. Afterwards, the liquid phase was aspirated, and the hydrogel was washed with ultrapure water. Coverslips coated with glycine hydrogel were used immediately for immobilization of cells. The hydrogel-coated coverslips can be stored at 4°C covered with a layer of ultrapure water for several days.

PLL coating of coverslips

Ultraclean coverslips were coated by applying 0.5 ml of 0.01% (w/v) PLL solution in ultrapure water and incubation for 20 min at room temperature in the laminar flow box. Unbound PLL was removed by aspiration of the liquid and a single wash with ultrapure water. Dried, PLL-

coated coverslips can be stored for several weeks. For SR imaging, freshly prepared PLL-coated coverslips were used exclusively.

Immobilization and fixation of cells for SR imaging

Jurkat T cells express varying levels of CD4 which are changing when cells are grown in culture. Therefore, we used transiently transfected Jurkat-CD4KO cells with reintroduced CD4-GFP to study CD4 surface distribution with dTRABI. Cells expressing similar levels of CD4-GFP were selected for imaging and data processing.

For SR imaging, cells were resuspended in pre-warmed PBS (made from 10x stock, Gibco) and seeded on glycine-coated coverslips immediately after the aspiration of a coating liquid. Cells were enabled to land on the coated optical surface for 10 minutes at 37°C in the CO_2 incubator. Afterwards, cells were fixed with pre-warmed 4% paraformaldehyde (Electron Microscopy Sciences) containing 2% saccharose in PBS for 45 minutes at room temperature and the process was stopped with 50 mM NH_4Cl (Sigma-Aldrich) in PBS and three rounds of washing with PBS.

Immunofluorescence

For labeling, cells were first incubated with 5% BSA in PBS (Blocking solution) for 1 hour at the ambient temperature to prevent non-specific binding of antibodies. Immunostaining of specific receptors was performed by incubating cells overnight with Alexa® 647 conjugated primary antibodies (human CD4: OKT4, dilution: 1:100, source: Biolegend; human CD45: MEM-28, dilution: 1:2000, source: ExBio) diluted in Blocking solution. The process was performed at the ambient temperature in a humid chamber to avoid drying of the solutions. After removing of redundant antibodies and 3 times washing with PBS for 5 minutes, the cells were post-fixed with 4% PFA for 5 minutes and washed 5 times with PBS.

The identical protocol was applied for the staining of CD45 on Jurkat CD4 KO cells transfected with pXJ41-CD4-mEOS2 plasmid for 2-color 2D SMLM analysis.

Single-Molecule Localization Microscopy

Three dimensional SMLM experiments for dTRABI analysis were performed on a home-built wide-field setup, which is described elsewhere in detail.[65] Raw image stacks were analyzed with rapidSTORM 3.2 (ref. [86]). Herein, the FWHM was set to 300 nm as an invariant parameter. Furthermore, the lower intensity threshold was set to 500 photons and the fit window radius to 1200 nm. All other fit parameters were kept from the default settings in rapidSTORM 3.2. Linear lateral drift correction was applied manually by spatio-temporally aligning distinct structures to themselves. This was facilitated by color-coding of the temporal coordinate with the built-in tool.

Two-color SMLM experiments were performed on a Nikon Eclipse Ti microscope, which is specified elsewhere in detail.[44] Contrary to the case of dTRABI analysis, the FWHM was set as a free fit parameter, but in the limits of 275 – 750 nm, both for Alexa Fluor 647 antibody and mEOS2 fusion protein localizations. All other parameters were kept consistent to the previous

experiments. Prior to imaging, a glass surface with Tetraspeck beads (Thermo Scientific) was imaged with alternating 561 nm and 647 nm excitation to create a nanometer precise map to allow the correction of chromatic shift. The 561 nm excitation channel was then mapped onto the 647 nm after the localization step.

Prior to acquisition, cells were irradiated in epifluorescence illumination mode to turn emitters, which were out-of-focus in the HILO illumination scheme, into the dark state. In all experiments, the length of the acquisition was set to capture the majority of emitters, i.e. imaging was concluded when only a very minor number of active emitters was detectable. Typical acquisition lengths were 60,000-120,000 frames for Alexa Fluor 647 channel and 30,000-60,000 frames for mEOS2 channel, where integration times were set to 20 ms and 16 ms in the single- and dual-color cases, respectively. Hereby, mEOS2 was excited at 561 nm and activated with 405 nm. The activation power density was increased over the time to create an almost constant signal density.

Intensity based biplane imaging – dTRABI

The principle of dTRABI

We previously reported TRABI-based biplane (BP-TRABI; ref. [65]), where, instead of the axially dependent FWHMs of spots in both optical channels, the photometric ratios *P* of the molecules were used. *P* was calculated as the quotient of the fit intensity (I_{Fit}) and reference intensity (I_R), where only the fixed FWHM PSF fit intensity is used. For BP-TRABI, the photometric ratio of both planes ($P_{1,2}$) is calculated according to

$$P_{1,2}(z) = \frac{P_1 - P_2}{P_1 + P_2}$$

(1)

with P_1 and P_2 being the photometric ratio of the individual planes.

The quality of the resulting axial coordinate depended on the precision of the TRABI intensity measurement. In order to exclude any interference from neighboring emitters as well as temporal overlap with other fluorophores, a set of rigorous exclusion criteria for spots was employed. As a consequence, many localizations given by the localization software were rejected by TRABI. To compensate, significantly longer image stacks had to be recorded to ensure structural consistency in the reconstructed images regarding the Nyquist-criterion. Though this was already an inconvenience for common SMLM organic fluorophores, which exhibit high repetition counts, it can be a significant obstacle for imaging approaches utilizing photo-activatable or -convertible fluorescent proteins. Furthermore, in cases of sub-optimal photo-switching rates that lead to high spot densities, the TRABI approach is computationally expensive or could even fail to produce an image due to a high rejection rate of fluorescent spots.

For a reasonable large TRABI radii I_R converges for both spots in both planes, thus Eq. (1) can be effectively simplified to

$$I_{\text{Fit},1,2}(z) = \frac{I_{\text{Fit},1} - I_{\text{Fit},2}}{I_{\text{Fit},1} + I_{\text{Fit},2}}$$

(2)

As equation 2 demonstrates, in order to achieve three-dimensional BP imaging, there is no need for an actual photometric TRABI analysis anymore, since I_R is cancelled out. Therefore, dTRABI allows for higher emitter densities, is quicker and requires less input parameters.

Calibration and allocation

Raw calibration curves were generated by linearly moving the focal plane through the sample plane while imaging a single-molecule surface under dSTORM conditions as previously described [65]. For this, a surface of BSA was doped with BSA molecules, attached to a short DNA sequence, labeled with Cy5. Fitting the raw emission pattern by independent Gaussians with equal FWHM yielded axially dependent single-molecule intensity curves. An axially precise raw calibration function γ is derived according to equation 2. The running median (binning width 25 nm) of the raw data is fitted with a high order polynomial to generate the axial lookup table.

After an initial rough alignment between the channels, experimental localizations from both optical channels were assigned by a framewise, linear nearest neighbor analysis. Here, the distance threshold was set to 500 nm, which seemed a reasonable value for the robust allocation between channels in a semi-sparse single-molecule environment. From these sets of localizations, the axially dependent intensity quotient $I_{\text{Fit},1,2}$ was calculated and roughly allocated to the look-up table (LUT). The final axial coordinate was determined by a linear interpolation of $I_{\text{Fit},1,2}$ between its "left" and "right" nearest neighbor coordinate of the LUT.

Drift and tilt correction

Since the plasma membrane can be seen as flat over the whole-cell scale, we reckoned that it can be used as its own fiducial marker. We traced the spatio-temporal axial footprint of the entire membrane by fitting of the raw localizations by a high-order polynomial in time. This is followed by the straightforward temporal linearization of the localization data, leading to a stable axial mean value over time. In order to instantly assess the quality of three-dimensional SMLM data we suggest the look for white regions in the color-coded image. In our experience, the abundance of these features usually suggests significant axial drift. However, even a subtle axial drift, which cannot be easily recognized by eye, will be detected by the approach described above. Additionally, non-linear drift, commonly occurring due to heating and resting of threads, can be accounted for.

Since we regularly observed a subtle axial tilt in the nanoscopic color-coded images due to the usage of a round coverslip in a magnetic holder, we developed a simple correction workflow (Fig. 3). The axial tilt of the sample is extrapolated by fitting a plane to the raw image data (pixelsize 100 nm). Afterwards, the data is linearized by simply subtracting the precise local plane-value from the raw localization.

Localization precision calculation

Drift and tilt corrected localizations were tracked in time by determining the three-dimensional nearest neighbor distances of localizations in consecutive frames. Localizations

constructing a track with a total inter-localization distance of less than 75 nm were considered to stem from the same fluorophore. For each sample type, we combined the nearest-neighbor tracks from all recorded independent fields of view, calculated the deviation from the mean coordinate of each track for all relevant spatial coordinates and derived a normalized histogram (Supplementary Fig. S5). By fitting these distributions with a Gaussian, we derived the localization precisions as the standard deviation of the mean.

Axial localization distribution analysis

Localization files obtained from rapidSTORM were loaded and processed in Fiji [76] with custom written scripts [77]. First, all localization coordinates in *x, y* and *z* were used to generate a quantitative stack of 2D images with 20 nm pixel size using a separation of 20 nm in *z* (i.e. z-stack). The numeric value in each pixel was equivalent to the number of localizations. Then the z-stack was segmented into defined regions of interest (ROIs). The interior of the cells was manually selected to serve as boundary ROI, in which the edges of the cells were spared. Within this master ROI a set of squared ROIs with 2×2 μm² area was automatically generated. The ROIs were allowed to be confined by the border of the master ROI and only kept when more than 75% of the ROI area, i.e. > 3 μm were preserved (Supplementary Fig. S6).

Afterwards, the localization density of each ROI was analyzed by accumulating the grey values per ROI within the image stack, i.e. stepwise in *z* every 20 nm, and plotted as a function of *z*. The plot was then fitted to a double Gaussian function of the form

$$y = a_1 \exp\left(-\frac{1}{2}\left(\frac{x-m_1}{s_1}\right)^2\right) + a_2 \exp\left(-\frac{1}{2}\left(\frac{x-m_2}{s_2}\right)^2\right),$$

with $a_{1,2}$ as amplitudes, $m_{1,2}$ as mean values and $s_{1,2}$ as standard deviations of each Gaussian.

The three parameters that were derived from this fit were the z-distribution width (Fig. 5e-g) calculated according to

$$\left(a_1 - \frac{FWHM1}{2}\right) + \left(a_2 + \frac{FWHM2}{2}\right) \text{, for } a_2 > a_1 \text{ with } FWHM_{1,2} = s_{1,2} \times 2.355;$$

the peak-to-peak distance (Supplementary Fig. S9) according to $|a_2 - a_1|$;

and the width difference of the two Gaussians (Supplementary Fig. S9) according to $|FWHM_1 - FWHM_2|$. By calculating these fit-derived metrics the axial distribution of localizations of CD4-WT, CD4-CS1 and CD45 was characterized.

Quantifying axial receptor distribution

For quantifying the obtained axial receptor distribution in Fig. 5f, we fitted the z-distribution width of CD4 WT, CD4 CS1 and CD45 with a model of Gaussian functions. CD4 WT was fitted using a mono Gaussian and the range of z-distribution widths was states as $x_c \pm FWHM/2$, with x_c as the center of the Gaussian and as FWHM as full width at half maximum of the distribution. CD4 CS1 and CD45 data were fitted using a double Gaussian and the resulting ranges were stated as $x_{c1} \pm FWHM_1/2$ and $x_{c2} \pm FWHM_2/2$. The peak-to-peak distributions (Supplementary Fig. S9) were modeled with a mono-exponential decay, and the decay

constant of the fit, i.e. the value at which the amplitude of the distribution is reduced to 36.8%, is stated as mean value in this manuscript.

SOFI

Sample preparation, coverslip coating and image acquisition were performed as described above for SMLM imaging. The raw data were analyzed using balanced SOFI algorithm as described before.[28, 87]

Confocal microscopy

Jurkat CD4-KO T cells were transiently transfected with pXJ41-CD4-mEos2 plasmid 24 hours before the imaging. Cells were immobilized, fixed and immunostained using anti-CD45 antibodies (MEM-28) as described above. Three-dimensional confocal imaging was performed on Leica TCS SP8 WLL SMD-FLIM microscope (Imaging Methods Core Facility of BIOCEV, Vestec, Czech Republic) using two-channel acquisition. The images were processed using standard functions of Fiji/ImageJ software (version 1.52p; ref. [76]).

Calcium measurements

Calcium sensor

Plasmid DNA with the ultrafast, genetically encoded calcium sensor $GCaMP6f_u$ was a kind gift from Katalin Török and Silke Kerruth (St George's, University of London). The coding sequence was amplified using primers AATAGATCTGCCACCATGGGCTGCGTGTGCTCCTC and AATGGATCCTCACTTCGCTGTCATCATTTGTACAAA and subcloned into pXJ41 vector using *BglII* and *BamHI* restriction sites as described before.[88]

Sample preparation

For calcium measurements, the coverslips were coated with 0.01% (w/v) poly-*L*-lysine (Sigma-Aldrich) or 2M glycine as described above. For coating with OKT3 (anti-CD3ε) antibody, clean coverslips were incubated with 0.01 μg/μl OKT3 in PBS for 30 minutes at 37°C. After a brief wash with PBS, the coverslips were used within a day. Coated coverslips were mounted into a ChamLide holder (Live Cell Instruments), filled with 500ul of color-free medium, placed on the microscope and focused to the focal plane prior the acquisition. The measurements were performed twenty hours after transfection of Jukrat T cells with the calcium sensor. For image analysis, cells were washed with PBS, resuspended in phenol red-free RPMI-1640 media (Sigma-Aldrich) supplemented with 2mM *L*-glutamine, 10mM HEPES, 1mM CaCl2 and 1mM MgCl2 and dropped onto the prepared coverslip while running the image acquisition.

The microscope setup and image acquisition

Live cell calcium mobilization imaging was performed on a home built TIRF microscope consisting of the IX73 frame (Olympus), UApo N, 100x 1.49 Oil immersion TIRF objective (Olympus) and OptoSplit II image splitter (CAIRN Optics) mounted on the camera port. Samples were illuminated using 200mW 488 nm laser (Sapphire, Coherent) in a TIRF mode and the intensity was regulated by acousto-optic tunable filter (AOTFnC-400.650-TN, AA

Optoelectronics). Fluorescence emission was detected by an EMCCD camera (iXon ULTRA DU-897U, Andor) with EM gain set to 200. Images were taken in 500ms intervals with the exposure time set to 50 ms.

Data processing

Calcium mobilization was quantified by calculating mean fluorescence in cells immobilized on coated coverslips over the period of 10 minutes. Fiji/ImageJ software was used for the analysis.[76] Graphs in Supplementary Fig. S4 represent a mean fluorescence changes in cells immobilized on PLL-, glycine- or OKT3-coated coverslips. The peak intensity (I(max)) was determined by finding a maximum value using Excel software (Microsoft). The kinetics of fluorescence decay was calculated as a ratio I(max)/I(max+5min) and I(max)/I(max+10min). Higher values indicate a more rapid decay of the fluorescence which indicates stronger response. Values of I(max+5min) and I(max+10min) represent the fluorescence intensity 5 and 10 minutes, respectively, after the maximum calcium response (I(max)) was reached. To avoid the impact of the intensity fluctuations, we calculated I(max), I(max+5min) and I(max+10min) as an average of 10 images (frames).

Characterization of T cell surface in contact with a coated coverslip and cell viability measurements

Microscope setup

For imaging of living T cells forming contact with a coated coverslip and for determining the viability of T cells immobilized on a coated coverslip, a home-build inverted microscope system (IX71, Olympus) equipped with 150 mW 488 nm and 150 mW 561 nm lasers (Sapphire, Coherent) was used. Fluorescence emission for cell surface-contact analysis and viability assay was detected by an EMCCD (iXon DU-897, Andor) and a sCMOS (Zyla-4.2-CL10, Andor) cameras, respectively. Two acousto-optic tunable filters (AOTFnC-400.650-TN, AA Optoelectronics) provided fast switching and synchronization of lasers with a camera. For TIRF imaging, the 488 nm laser beam was focused onto the back focal plane of an objective (UApoN 100x, NA = 1.49, Olympus). TIRF illumination was achieved using a manual micrometer-scale tilting mirror mechanism in the excitation pathway (Thorlabs). For cell viability measurements, the laser light was defocused to achieve a homogenous illumination at the objective sample plane (UPlanSApo 10x, NA = 0.4, Olympus). The system was controlled using the µManager software (version 1.4.22; ref. [89]).

Analysis of the T cell surface in contact with a coated coverslip

For T cell-coverslip contact analysis, Jurkat T CD4-KO cells were transfected with pXJ41-CD4-wt-eGFP plasmid 24-hours prior to the measurement. Cells were then harvested and transferred into color-free RMPI medium (Gibco) supplemented with 2 mM L-glutamine (Lonza) pre-warmed to 37°C. In parallel, cleaned coverslips were mounted into a ChamLide holder for imaging equipped with tubing adaptors (Live Cell Instruments) and covered with a coating solution as described above. After washing with ultrapure water, cells were injected into the ChamLide chamber attached to the microscope stage immediately after coating fluid

removal through the attached tubing. Landing cells were selected under transmitted light and the fluorescence was recorded on EMCCD camera using TIRF illumination for 20 minutes. Images were acquired in one second intervals with 50 ms exposure time. The camera EM gain was set to 100. The experiment was performed at 37°C using the environmental chamber (OKO lab) with controlled temperature. Acquired images were processed using standard functions of Fiji/ImageJ software (version 1.52p; ref. [76]).

T cell viability assay

To evaluate the viability of cells in contact with coated coverslips, Jurkat T cells were transfected with mTurquoise-Farnesyl-5 plasmid (#55551, Addgene bank) 24 hours prior to the assay. This was done to mimic the conditions used for all other experiments in this work but to avoid interference of the protein fluorescence with the dyes used for the assay. For the measurement, cells were washed with PBS, transferred into color-free RPMI (Gibco) medium supplemented with 2 mM *L*-glutamine (Lonza), 10% fetal bovine serum (Gibco), 10 mM HEPES and Non-essential amino-acid mixture (Gibco) containing 25 μg/ml (w/v) calcein-AM viability stain and incubated for 10 minutes at 37°C in the CO_2 incubator. Cells were then transferred into a fresh, color-free RPMI medium supplemented with 2 mM *L*-glutamine and 7AAD (1:20 dilution; 7-Amino-Actinomycine D) viability stain (eBioscience) and incubated for another 5 minutes at 37°C in the CO_2 incubator. After loading with the dyes, cells were injected onto the coated coverslips in ChamLide chamber using attached tubing. Imaging was performed for 30 minutes (with 10 minute acquisition intervals) from the time of the fist cell-coverslip contact detection. Two randomly selected ROIs were selected for each time point to avoid the effect of the fluorophore photo-destruction. Calcein and 7AAD dyes were excited separately by 488 nm and 561 nm lasers, respectively, using defocused light to achieve homogenous illumination. Cells were also imaged under transmitted light to control their morphology. To avoid the impact of fluorescence fluctuations, five frames were collected for each channel, ROI and time point. The frames were then summed to generate a collate image for further quantitative analysis. Data analysis was performed in Fiji/ImageJ (version 1.52p, ref. [76]). Cells were counted by determining local maxima for each channel. The final result was calculated as a percentage of 7AAD positive (dying) cells in all detected cells (calcein positive). The experiment was performed at 37°C using the environmental chamber (OKO lab) with controlled temperature.

Atomic force microscopy

Atomic Force Microscopy (AFM) topography images were collected with Dimension Icon AFM (Bruker Instruments). All images were measured in the Peak Force Tapping mode for fluids, using a probe holder for fluid operation and Scanasyst-Fluid probes (Bruker) with a tip radius of 20 nm and a spring constant of 0.7 N/m.

12 mm diameter coverslips were coated with PLL and glycine as described above. A bare coverslip was prepared as a reference. The AFM probe was submerged in the solution in order to land on the coverslip surface. Due to a delicate nature of the samples and to avoid long

measurement times, setpoint and number of lines were set to 400 pN and 256 x 256 lines respectively, with a scan rate of 1 Hz. Images with scan sizes of 1, 2 and 10 μm were collected.

Acknowledgements

We would like to thank Peter Kapusta, Silke Kerruth and Harsha Mavila for technical assistance and professional advice. We are grateful to Markus Sauer (University of Würzburg) for support of the project. C.F. would like to thank Laure Plantard and Jan Peychl from the MPI-CBG LMF for technical support. M.C. acknowledges funding from Czech Science Foundation (19-0704S), S.vdL. acknowledges funding from Academy of Medical Sciences/the British Heart Foundation/the Government Department of Business, Energy and Industrial Strategy/the Wellcome Trust Springboard Award (SBF003\1163). The measurements at the Imaging Methods Core Facility in BIOCEV, Vestec, Czech Republic were supported by MEYS CR grant Z.02.1.01/0.0/0.0/16_013/0001775.

Contributions

C.F., S.vdL. and M.C. conceived the study. C.F., T.C., Z.K., D.G. and D.H. performed the experiments, C.F., T.C., D.G., S.vdL. and M.C. analyzed the data. A.F. measured and analyzed the AFM data. O.F. and T.B. provided research advice. C.F., S.vdL. and M.C. wrote the paper. All authors reviewed and approved the manuscript.

Corresponding authors

Correspondence to Christian Franke, Sebastian van de Linde and Marek Cebecauer

Additional information

Supplementary Information accompanies this paper at …

Competing interests

The authors declare no competing financial interests.

REFERENCES

1. Cebecauer, M., Spitaler, M., Serge, A. & Magee, A.I. Signalling complexes and clusters: functional advantages and methodological hurdles. *J Cell Science* **123**, 309-320 (2010).

2. Pak, A.J. *et al.* Immature HIV-1 lattice assembly dynamics are regulated by scaffolding from nucleic acid and the plasma membrane. *Proc Natl Acad Sci U S A* **114**, E10056-E10065 (2017).

3. Rossier, O. *et al.* Integrins beta1 and beta3 exhibit distinct dynamic nanoscale organizations inside focal adhesions. *Nat Cell Biol* **14**, 1057-1067 (2012).

4. Barreiro, O. *et al.* Endothelial adhesion receptors are recruited to adherent leukocytes by inclusion in preformed tetraspanin nanoplatforms. *J Cell Biol* **183**, 527-542 (2008).

5. Kellermayer, B. *et al.* Differential Nanoscale Topography and Functional Role of GluN2-NMDA Receptor Subtypes at Glutamatergic Synapses. *Neuron* **100**, 106-119 e107 (2018).

6. Sevcsik, E. & Schutz, G.J. With or without rafts? Alternative views on cell membranes. *Bioessays* **38**, 129-139 (2015).

7. Dustin, M.L. & Davis, S.J. TCR signaling: the barrier within. *Nat Immunol* **15**, 136-137 (2014).

8. Bernardino de la Serna, J., Schutz, G.J., Eggeling, C. & Cebecauer, M. There Is No Simple Model of the Plasma Membrane Organization. *Front Cell Dev Biol* **4**, 106 (2016).

9. Brameshuber, M. *et al.* Monomeric TCRs drive T cell antigen recognition. *Nat Immunol* **19**, 487-496 (2018).

10. James, J.R. *et al.* The T Cell Receptor Triggering Apparatus Is Composed of Monovalent or Monomeric Proteins. *J Biol Chem* **286**, 31993-32001 (2011).

11. Gomes de Castro, M.A. *et al.* Differential organization of tonic and chronic B cell antigen receptors in the plasma membrane. *Nat Commun* **10**, 820 (2019).

12. Campi, G., Varma, R. & Dustin, M.L. Actin and agonist MHC-peptide complex-dependent T cell receptor microclusters as scaffolds for signaling. *J Exp Med* **202**, 1031-1036 (2005).

13. Tolar, P., Hanna, J., Krueger, P.D. & Pierce, S.K. The constant region of the membrane immunoglobulin mediates B cell-receptor clustering and signaling in response to membrane antigens. *Immunity* **30**, 44-55 (2009).

14.   Saka, S.K. *et al.* Multi-protein assemblies underlie the mesoscale organization of the plasma membrane. *Nat Commun* **5**, 4509 (2014).

15.   Letschert, S. *et al.* Super-resolution imaging of plasma membrane glycans. *Angew Chem Int Ed Engl* **53**, 10921-10924 (2014).

16.   Mateos-Gil, P., Letschert, S., Doose, S. & Sauer, M. Super-Resolution Imaging of Plasma Membrane Proteins with Click Chemistry. *Front Cell Dev Biol* **4**, 98 (2016).

17.   Puchner, E.M., Walter, J.M., Kasper, R., Huang, B. & Lim, W.A. Counting molecules in single organelles with superresolution microscopy allows tracking of the endosome maturation trajectory. *Proc Natl Acad Sci U S A* **110**, 16015-16020 (2013).

18.   Loschberger, A., Franke, C., Krohne, G., van de Linde, S. & Sauer, M. Correlative super-resolution fluorescence and electron microscopy of the nuclear pore complex with molecular resolution. *J Cell Sci* **127**, 4351-4355 (2014).

19.   Ehmann, N. *et al.* Quantitative super-resolution imaging of Bruchpilot distinguishes active zone states. *Nat Commun* **5**, 4650 (2014).

20.   Jungmann, R. *et al.* Quantitative super-resolution imaging with qPAINT. *Nat Methods* **13**, 439-442 (2016).

21.   Lillemeier, B.F. *et al.* TCR and Lat are expressed on separate protein islands on T cell membranes and concatenate during activation. *Nat Immunol* **11**, 90-96 (2010).

22.   Rossy, J., Owen, D.M., Williamson, D.J., Yang, Z. & Gaus, K. Conformational states of the kinase Lck regulate clustering in early T cell signaling. *Nat Immunol* **14**, 82-89 (2013).

23.   Rossboth, B. *et al.* TCRs are randomly distributed on the plasma membrane of resting antigen-experienced T cells. *Nat Immunol* **19**, 821-827 (2018).

24.   Mattila, P.K. *et al.* The actin and tetraspanin networks organize receptor nanoclusters to regulate B cell receptor-mediated signaling. *Immunity* **38**, 461-474 (2013).

25.   Balint, S., Lopes, F.B. & Davis, D.M. A nanoscale reorganization of the IL-15 receptor is triggered by NKG2D in a ligand-dependent manner. *Sci Signal* **11**, eaal3606 (2018).

26.   Annibale, P., Vanni, S., Scarselli, M., Rothlisberger, U. & Radenovic, A. Identification of clustering artifacts in photoactivated localization microscopy. *Nat Methods* **8**, 527-528 (2011).

27.   Burgert, A., Letschert, S., Doose, S. & Sauer, M. Artifacts in single-molecule localization microscopy. *Histochem Cell Biol* **144**, 123-131 (2015).

28.   Lukes, T. *et al.* Quantifying protein densities on cell membranes using super-resolution optical fluctuation imaging. *Nat Commun* **8**, 1731 (2017).

29.   Levet, F. *et al.* SR-Tesseler: a method to segment and quantify localization-based super-resolution microscopy data. *Nat Methods* **12**, 1065-1071 (2015).

30.   Baumgart, F. *et al.* Varying label density allows artifact-free analysis of membrane-protein nanoclusters. *Nat Methods* **13**, 661-664 (2016).

31.   Spahn, C., Herrmannsdorfer, F., Kuner, T. & Heilemann, M. Temporal accumulation analysis provides simplified artifact-free analysis of membrane-protein nanoclusters. *Nat Methods* **13**, 963-964 (2016).

32.   Vangindertael, J. *et al.* An introduction to optical super-resolution microscopy for the adventurous biologist. *Methods Appl Fluoresc* **6**, 022003 (2018).

33.   Raghunathan, K. & Kenworthy, A.K. Dynamic pattern generation in cell membranes: Current insights into membrane organization. *Biochim Biophys Acta Biomembr* **1860**, 2018-2031 (2018).

34.   Sezgin, E., Levental, I., Mayor, S. & Eggeling, C. The mystery of membrane organization: composition, regulation and roles of lipid rafts. *Nat Rev Mol Cell Biol* **18**, 361-374 (2017).

35.   Jarsch, I.K., Daste, F. & Gallop, J.L. Membrane curvature in cell biology: An integration of molecular mechanisms. *J Cell Biol* **214**, 375-387 (2016).

36.   Fisher, P.J., Bulur, P.A., Vuk-Pavlovic, S., Prendergast, F.G. & Dietz, A.B. Dendritic cell microvilli: a novel membrane structure associated with the multifocal synapse and T-cell clustering. *Blood* **112**, 5037-5045 (2008).

37.   Kim, H.R. *et al.* T cell microvilli constitute immunological synaptosomes that carry messages to antigen-presenting cells. *Nat Commun* **9**, 3630 (2018).

38.   Shalek, A.K. *et al.* Nanowire-mediated delivery enables functional interrogation of primary immune cells: application to the analysis of chronic lymphocytic leukemia. *Nano Lett* **12**, 6498-6504 (2012).

39.   Gorelik, J. *et al.* Dynamic assembly of surface structures in living cells. *Proc Natl Acad Sci U S A* **100**, 5819-5822 (2003).

40.   Lange, K. Fundamental role of microvilli in the main functions of differentiated cells: Outline of an universal regulating and signaling system at the cell periphery. *J Cell Physiol* **226**, 896-927 (2011).

41. Millington, P.F., Critchley, D.R., Tovell, P.W. & Pearson, R. Scanning electron microscopy of intestinal microvilli. *J Microsc* **89**, 339-344 (1969).

42. Griffiths, G. & Lucocq, J.M. Antibodies for immunolabeling by light and electron microscopy: not for the faint hearted. *Histochem Cell Biol* **142**, 347-360 (2014).

43. Schwarz, H. & Humbel, B.M. Correlative light and electron microscopy using immunolabeled sections. *Methods Mol Biol* **1117**, 559-592 (2014).

44. Franke, C. *et al.* Correlative single-molecule localization microscopy and electron tomography reveals endosome nanoscale domains. *Traffic* **20**, 601-617 (2019).

45. Li, D. *et al.* ADVANCED IMAGING. Extended-resolution structured illumination imaging of endocytic and cytoskeletal dynamics. *Science* **349**, aab3500 (2015).

46. Geissbuehler, S. *et al.* Live-cell multiplane three-dimensional super-resolution optical fluctuation imaging. *Nat Commun* **5**, 5830 (2014).

47. von Diezmann, A., Shechtman, Y. & Moerner, W.E. Three-Dimensional Localization of Single Molecules for Super-Resolution Imaging and Single-Particle Tracking. *Chem Rev* **117**, 7244-7275 (2017).

48. Huang, B., Wang, W., Bates, M. & Zhuang, X. Three-dimensional super-resolution imaging by stochastic optical reconstruction microscopy. *Science* **319**, 810-813 (2008).

49. Juette, M.F. *et al.* Three-dimensional sub-100 nm resolution fluorescence microscopy of thick samples. *Nat Methods* **5**, 527-529 (2008).

50. Ram, S., Prabhat, P., Chao, J., Ward, E.S. & Ober, R.J. High accuracy 3D quantum dot tracking with multifocal plane microscopy for the study of fast intracellular dynamics in live cells. *Biophys J* **95**, 6025-6043 (2008).

51. Pavani, S.R. *et al.* Three-dimensional, single-molecule fluorescence imaging beyond the diffraction limit by using a double-helix point spread function. *Proc Natl Acad Sci U S A* **106**, 2995-2999 (2009).

52. Shtengel, G. *et al.* Interferometric fluorescent super-resolution microscopy resolves 3D cellular ultrastructure. *Proc Natl Acad Sci U S A* **106**, 3125-3130 (2009).

53. Dustin, M.L. & Depoil, D. New insights into the T cell synapse from single molecule techniques. *Nat Rev Immunol* **11**, 672-684 (2011).

54. Smith, C.L. Basic Confocal Microscopy. *Curr. Protoc. Neurosci.*, vol. 56. John Wiley & Sons Inc., 2011, pp 2.2.1-2.2.18.

55. Mazia, D., Schatten, G. & Sale, W. Adhesion of cells to surfaces coated with polylysine. Applications to electron microscopy. *J Cell Biol* **66**, 198-200 (1975).

56. Groves, J.T. & Dustin, M.L. Supported planar bilayers in studies on immune cell adhesion and communication. *J Immunol Methods* **278**, 19-32 (2003).

57. Bunnell, S.C. *et al.* T cell receptor ligation induces the formation of dynamically regulated signaling assemblies. *J Cell Biol* **158**, 1263-1275 (2002).

58. Fritzsche, M. *et al.* Cytoskeletal actin dynamics shape a ramifying actin network underpinning immunological synapse formation. *Sci Adv* **3**, e1603032 (2017).

59. Cavo, M. *et al.* A new cell-laden 3D Alginate-Matrigel hydrogel resembles human breast cancer cell malignant morphology, spread and invasion capability observed "in vivo". *Sci Rep* **8**, 5333 (2018).

60. Chakraborty, A.K. & Weiss, A. Insights into the initiation of TCR signaling. *Nat Immunol* **15**, 798-807 (2014).

61. Glatzova, D. & Cebecauer, M. Dual Role of CD4 in Peripheral T Lymphocytes. *Front Immunol* **10**, 618 (2019).

62. Cordoba, S.P. *et al.* The large ectodomains of CD45 and CD148 regulate their segregation from and inhibition of ligated T-cell receptor. *Blood* **121**, 4295-4302 (2013).

63. Varma, R., Campi, G., Yokosuka, T., Saito, T. & Dustin, M.L. T cell receptor-proximal signals are sustained in peripheral microclusters and terminated in the central supramolecular activation cluster. *Immunity* **25**, 117-127 (2006).

64. van der Merwe, P.A. & Dushek, O. Mechanisms for T cell receptor triggering. *Nat Rev Immunol* **11**, 47-55 (2011).

65. Franke, C., Sauer, M. & van de Linde, S. Photometry unlocks 3D information from 2D localization microscopy data. *Nat Methods* **14**, 41-44 (2017).

66. Sattentau, Q.J., Dalgleish, A.G., Weiss, R.A. & Beverley, P.C. Epitopes of the CD4 antigen and HIV infection. *Science* **234**, 1120-1123 (1986).

67. Bazil, V., Hilgert, I., Kristofova, H., Maurer, D. & Horejsi, V. Sialic acid-dependent epitopes of CD45 molecules of restricted cellular expression. *Immunogenetics* **29**, 202-205 (1989).

68.   Schermelleh, L. *et al.* Super-resolution microscopy demystified. *Nat Cell Biol* **21**, 72-84 (2019).

69.   de Kruijff, B. & Cullis, P.R. The influence of poly(L-lysine) on phospholipid polymorphism. Evidence that electrostatic polypeptide-phospholipid interactions can modulate bilayer/non-bilayer transitions. *Biochim Biophys Acta* **601**, 235-240 (1980).

70.   Pachmann, K. & Leibold, W. Insolubilization of protein antigens on polyacrylic plastic beads using poly-L-lysine. *J Immunol Methods* **12**, 81-89 (1976).

71.   van de Linde, S. *et al.* Direct stochastic optical reconstruction microscopy with standard fluorescent probes. *Nat Protoc* **6**, 991-1009 (2011).

72.   Klein, T. *et al.* Live-cell dSTORM with SNAP-tag fusion proteins. *Nat Methods* **8**, 7-9 (2011).

73.   Santos, A.M. *et al.* Capturing resting T cells: the perils of PLL. *Nat Immunol* **19**, 203-205 (2018).

74.   Tokunaga, M., Imamoto, N. & Sakata-Sogawa, K. Highly inclined thin illumination enables clear single-molecule imaging in cells. *Nat Methods* **5**, 159-161 (2008).

75.   Franke, C. & van de Linde, S. Reply to 'Impact of optical aberrations on axial position determination by photometry'. *Nat Methods* **15**, 990-992 (2018).

76.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012).

77.   van de Linde, S. Single-molecule localization microscopy analysis with ImageJ. *J Phys D Appl Phys* **52** (2019).

78.   Hermiston, M.L., Xu, Z. & Weiss, A. CD45: a critical regulator of signaling thresholds in immune cells. *Annu Rev Immunol* **21**, 107-137 (2003).

79.   Fernandes, R.A. *et al.* A cell topography-based mechanism for ligand discrimination by the T cell receptor. *Proc Natl Acad Sci U S A* **116**, 14002-14010 (2019).

80.   Axelrod, D. Cell-substrate contacts illuminated by total internal reflection fluorescence. *J Cell Biol* **89**, 141-145 (1981).

81.   Blaskovic, S., Blanc, M. & van der Goot, F.G. What does S-palmitoylation do to membrane proteins? *FEBS J* **280**, 2766-2774 (2013).

82.   Jung, Y. *et al.* Three-dimensional localization of T-cell receptors in relation to microvilli using a combination of superresolution microscopies. *Proc Natl Acad Sci U S A* **113**, E5916-E5924 (2016).

83.   Hashimoto-Tane, A. *et al.* Micro-adhesion rings surrounding TCR microclusters are essential for T cell activation. *J Exp Med* **213**, 1609-1625 (2016).

84.   Martin, B.R. & Cravatt, B.F. Large-scale profiling of protein palmitoylation in mammalian cells. *Nat Methods* **6**, 135-138 (2009).

85.   Kornete, M., Marone, R. & Jeker, L.T. Highly Efficient and Versatile Plasmid-Based Gene Editing in Primary T Cells. *J Immunol* **200**, 2489-2501 (2018).

86.   Wolter, S. *et al.* rapidSTORM: accurate, fast open-source software for localization microscopy. *Nat Methods* **9**, 1040-1041 (2012).

87.   Geissbuehler, S. *et al.* Mapping molecular statistics with balanced super-resolution optical fluctuation imaging (bSOFI). *Optical Nanoscopy* **1**, 4 (2012).

88.   Chum, T. *et al.* The role of palmitoylation and transmembrane domain in sorting of transmembrane adaptor proteins. *J Cell Science* **129**, 95-107 (2016).

89.   Edelstein, A.D. *et al.* Advanced methods of microscope control using muManager software. *J Biol Methods* **1** (2014).

FIGURE CAPTIONS

Figure 1. Coating of coverslips with glycine preserves receptor distribution and resting state of imaged T cells. a-b) Schematic illustration of T cell landing on PLL- (left panels) and glycine-coated coverslips (right panels). The arrows indicate forces influencing the cell on the coated coverslip. g-force is the sole force affecting T cells on glycine. T cells on PLL are further stretched due to electrostatic interactions of the surface molecules with PLL. The blue strips represent glass coverslip, the light blue strips above the glycine layer (not in scale). **c-d)** Live cell TIRF microscopy of CD4-GFP in T cells landing on PLL-coated (c) or glycine-coated coverslips (d) measured at 37°C. Blue arrowheads indicate areas of rapid flattening of the cell surface and random distribution of CD4 as indicated by line-profiles in Supplementary Fig. S1. Red arrowheads indicate areas with heterogenous distribution of CD4 on the cell surface as indicated by line-profiles in Supplementary Fig. S1. Selected time-points for the representative cells are shown. In total, 14 cells on glycine-coated and 10 cells on PLL-coated coverslips were analyzed. **e)** Calcium response induced by the interaction of Jurkat T cells with coverslips coated with stimulating antibody (anti-CD3ε; OKT3), PLL or glycine as indicated by the changes in GCaMP3-fast fluorescence over 5 min (black bars) and 10 min (grey bars) since the maximal stimulation of cells. Higher values indicate stronger stimulation of cells (see **Methods** and Supplementary Fig. S4 for more details). **f)** Viability of Jurkat T cells interacting with coverslips coated with PLL- (black bars) or glycine-coated (grey bars) coverslips. Dying cells were defined as a fraction of 7-aminoactinomycin-positive cells within the imaged area using a wide-field microscopy (see **Methods** for more details).

Figure 2. Principle of 3D super-resolution bi-plane imaging using dTRABI. a) Overview of the experimental setup applied to perform 3D dTRABI imaging of T cells. HILO illumination of the sample (blue beam; only shown in enlarged box) triggers fluorescence emission (orange), which is split by a 50/50 non-polarizing beamsplitter (50/50 BS) to acquire biplane images on two separate EM-CCD cameras. The respective imaging lenses are shifted along the optical axis to induce a relative defocus of the image detection on synchronized cameras. Spots, apparent in both detection planes are fitted by a Gaussian with identically set FWHM. **b)** Using a piezo stage, the focal plane was linearly moved through the sample plane while imaging a single-molecule surface under dSTORM conditions. Hereby, both cameras were synchronized. Fitting the raw PSFs by independent Gaussians with invariable FWHM yields axially dependent single-molecule intensity curves (upper panel). The relative change of position of the imaging lens in the reflection path is mirrored by the relative shift of the respective intensity curve (indicated by circles). Data points were spline interpolated to guide the eye (solid lines). An axially precise calibration function γ is derived directly from the raw intensities ($I_{Camera1}$, $I_{Camera2}$) of corresponding localizations from both cameras as $γ(z) = (I_{Camera2} − I_{Camera1})(I_{Camera2} + I_{Camera1})^{-1}$. The running median of the raw data (grey squares) is fitted with a high order polynomial (black line) to generate the basis of the axial lookup table (lower panel). **c)** A two-dimensional high-resolution data set is generated from both image stacks (transmission and reflection path) to create a three-dimensional dTRABI data

set according to the calibration. Finally, the transmission localization set is used to render a high-resolution, axially color-coded image of the focused target structure. Scale bars, 5 μm.

Figure 3. Fiducial free drift and tilt correction of dTRABI data. CD4 WT on Jurkat T cells are shown, where the membrane was directly labelled by a primary antibody conjugated to Alexa Fluor 647. **a)** Principle of the fiducial free correction of axial drift. *Left:* Representative, color-coded high-resolution dTRABI image of an axially unstable sample. Due to the temporal change of the axial coordinate, the resulting image does not exhibit axially distinct features. *Middle:* By extracting the axial localization distribution of the thin membrane layer per frame and tracking it over the entire stack, a spatio-temporal drift trace can be plotted (top). Fitting of the raw data (grey squares) by a high-order polynomial (red line) allows the temporal linearization of the localization data, leading to a stable axial mean value over time (bottom). *Right:* Re-rendering of the drift-corrected localization data reveals a color-coded dTRABI image, exhibiting distinct, high-density clusters of CD4 below another, disperse, layer. **b)** Principle of the fiducial free correction of axial tilt. *Left:* Representative, color-coded high-resolution dTRABI image of an axially tilted sample. *Middle:* The axial tilt of the sample is extrapolated by a fitting a plane to the raw image. Afterwards, the data is linearized by subtracting the local plane-value from the raw localization. *Right:* The tilt corrected image exhibits a homogeneous color-code in the lowest data layer, indicating no residual axial tilt. Exemplary drift and tilt were simulated based on real data, where the right column represents experimentally corrected real data on which the drift and tilt was projected. Scale bars, 5 μm.

Figure 4. Cell surface receptor nanotopography visualized by 3D TRABI imaging. a) Schematic illustration of a sample preparation for a receptor nanotopography imaging using dTRABI. A cell in suspension is first immobilized on a glycine-coated coverslip and imaged using dTRABI approach. The resulting 3D dTRABI image represents a footprint of a cell on the optical surface with the color-coded axial position of localizations (right panel). **b)** 3D TRABI image of T cell surface receptor CD45 with selected ROIs exhibiting a broad z-distribution of receptor localizations. **c)** Magnified x-z projections of ROIs 1-5 as in a). Blue arrowheads point to microscopic membrane protrusions at the cell edges, magenta arrowheads to folded nanoscopic protrusions under the cell body and green arrowhead membrane invagination. Scale bars represent 5 μm in b) and 500 nm in c).

Figure 5. CD4 clusters represent the receptor accumulation at the tips of membrane protrusions. **a)** Representative 3D dTRABI image of CD4 WT at the surface of T cell immobilized on a glycine-coated coverslip. **b)** The axial distribution of CD4 WT localizations of the 4 μm² region of interest (ROI) of the cell as in a). **c)** Representative 3D dTRABI image of non-palmitoylatable CD4 CS1 mutant at the surface of T cell immobilized on a glycine-coated coverslip. **d)** The axial distribution of CD4 CS1 localizations of the 4 μm² ROI of the cell in c). **e-g)** Quantitative analysis of the receptor axial (z-axis) distribution on the surface of T cells. Receptor z-distribution values were calculated using a double Gaussian fit to the axial distribution of localizations for each ROI as in b, d and i (see Supplementary Fig. S6 for more examples) and the FWHM range of the two Gaussian functions represent the z-distribution

width as depicted in **e**. Black circles in **e** represent the axial distribution of receptor localizations for a selected ROI, black line the double Gaussian fit, which is the sum of two Gaussians as depicted in green and magenta, dashed lines in grey depict z-distribution width and lines in light grey depict mean values of the Gaussian distributions. The graphs in **f** and **g** represent z-distribution widths obtained from 21 CD4 WT cells with 846 ROIs (black), 18 CD4 CS1 cells with 1044 ROIs (green) and 13 CD45 WT cells with 305 ROIs (magenta). The distributions in f) and g) show relative and absolute occurrence, respectively. Data points in b), d), f) and i) were spline interpolated to guide the eye. **h**) Representative 3D dTRABI image of CD45 at the surface of T cell immobilized on a glycine-coated coverslip. Color-bars in the upper right corner a), c) and h) indicate the axial position of the localizations in the image. **i**) The axial distribution of CD45 localizations of the 4 μm² ROI of the cell in h). Scale bars in a), c) and g): 5 μm.

Figure 6. 3D dTRABI image of CD45 on the surface of resting T cells and its axial distribution analysis. **a**) 3D dTRABI image of CD45 on T cell. **b**) Magnified ROI as in a) with indicated areas lacking CD45 localizations (yellow arrowheads). **c)** 2D SOFI image of CD45 on T cell immobilized on a glycine-coated coverslip. CD45 was labelled using Alexa Fluor-647 conjugated MEM-28 antibody. Yellow arrowheads indicate cell surface areas lacking CD45 signal. Scale bar in: a) 5 μm, b) 1 μm and c) 5 μm.

Figure 7. Nanoscopic segregation of CD4 and CD45 on the surface of resting T cells. a) Representative 2D image of T cell surface sequentially analyzed for CD4 (green) and CD45 (magenta) by SMLM. CD4 was visualized as mEos2 fusion protein (PALM) after transient transfection of cells and the surface CD45 was labelled using Alexa Fluor-647 conjugated MEM-28 antibody (dSTORM). ROIs 1-2 were zoomed to indicate details of proteins' distribution (right hand side). **b**) Intensity line-profiles were measured along the transparent grey regions indicated in ROIs 1-2 (as in a). Green line represents CD4 and magenta CD45 signals. **c**) Schematic illustration of microvillus with indicated structural segments: tip, shaft and the basis. The two panels below indicate the predicted organization of CD4 and CD45 on the protrusions in cells expressing native CD4 (WT; middle panel) or its non-palmitoylatable variant (CS1; lower panel) CD4. **d)** 2D SMLM image of a selected T cell associating with the optical surface which was sequentially analyzed for CD4 and CD45 (as in a). The accumulation of CD4 on the tips of large membrane protrusions was observed in 10 out of 32 imaged T cells. ROIs 3-4 were zoomed to indicate the details of membrane protrusions with accumulated CD4 on their tips (right hand side). **e**) Intensity line-profiles were measured along the transparent grey regions indicated in ROIs 3-4. The arrows indicate the onset of line-profiles. Scale bars, 5 μm. Images from three independent experiments are shown (n=32).

Figure 8. Microscopic segregation of CD4 and CD45 on T cells associating with the optical surface. a) Confocal micrographs (x-z section) of CD4 (green) and CD45 (magenta) in T cells landing on a coverslip coated with glycine. Early (upper panel), intermediate (middle panel) and later stage (lower panel) of T cell landing on a coverslip is presented. **b)** X-Y sections of

3D confocal images of CD4 (green) and CD45 (magenta) selected at, or close to, the optical surface. CD4 and CD45 were labelled as in Fig.7. Scale bars, 5 μm. Representative images from two independent experiments are shown (n=29).

Figure 9. T cell surface morphology on glycine-coated coverslips. a) Schematic illustration of a T cell in suspension. The cell is covered with numerous membrane protrusions of different sizes. The size of Jurkat T cells is on average 10 μm in diameter, the protrusions, even though a majority is small (0.5 – 2 μm length), can extend up to 5 μm from the cell body. **b)** After landing on a glycine-coated coverslip, membrane protrusions of T cells are folding under the cell body but are not rapidly removed as on PLL. **c-d)** With the axial penetration depth of dTRABI being ~ 1 μm, we were unable to detect receptors at the plasma membrane basis of T cells exhibiting displacement of the cell body from the contact site (c) or those with the cavity formed between the cell body and the coated coverslip. The footprints of such cells show discontinuous labeling or a hole in the middle of the image (see the inserts). The blue strips represent glass coverslip, the light blue strips above the glycine layer (not in scale).

Franke et al. 2020, Fig. 1

Franke et al. 2020, Fig. 2

Franke et al. 2020, Fig. 3

Franke et al. 2020, Fig. 4

Franke et al. 2020, Fig. 5

Franke et al. 2020, Fig. 6

Franke et al. 2020, Fig. 7

Franke et al. 2020, Fig. 8