Functional redundancy and crosstalk between flavodiiron proteins and NDH-1 in

Synechocystis **sp. PCC 6803**

Short title: Crosstalk between flavodiiron proteins and NDH-1

Lauri Nikkanen[1], Anita Santana Sánchez[1], Maria Ermakova[1*], Matthias Rögner[2], Laurent Cournac[3],

and Yagut Allahverdiyeva[1#]

Affiliations:

[1]Molecular Plant Biology, Department of Biochemistry, University of Turku, Turku, Finland

[2]Plant Biochemistry, Faculty of Biology & Biotechnology, Ruhr-University Bochum, Bochum,
Germany

[3]Eco&Sols, University of Montpellier, IRD, CIRAD, INRA, Montpellier Supagro, Montpellier,
France

One sentence summary: Flavodiiron proteins and NDH-1 complex ensure survival of
cyanobacterial cells by cooperatively safeguarding the photosynthetic apparatus against excessive
reduction

Author contributions: Y.A., and L.N. conceptualized the research; L.N., A.S.S., M.E., and Y.A. designed the research; M.E. and A.S.S. conducted growth experiments; L.N. performed MIMS and spectrophotometric experiments; A.S.S. performed immunoblotting; all authors analyzed the data; L.N. and Y.A. wrote the article with the input from A.S.S and M.E.

Funding: This work was supported by the Academy of Finland (project #315119 to Y.A. and the Finnish Center of Excellence, project #307335), the Nordforsk Nordic Center of Excellence 'NordAqua' (#82845) and SFB480- Germany

[*]**Current affiliation:** Australian Research Council Centre of Excellence for Translational Photosynthesis, Division of Plant Science, Research School of Biology, The Australian National University, Acton, Australia

#**Author for contact**: Yagut Allahverdiyeva, allahve@utu.fi

ABSTRACT

In oxygenic photosynthetic organisms excluding angiosperms, flavodiiron proteins (FDPs) catalyze light-dependent reduction of O_2 to H_2O. This alleviates electron pressure on the photosynthetic apparatus and protects it from photodamage. In *Synechocystis* sp. PCC 6803, four FDP isoforms function as hetero-oligomers of Flv1 and Flv3 and/or Flv2 and Flv4. An alternative electron transport pathway mediated by the NAD(P)H dehydrogenase-like complex (NDH-1) also contributes to redox hemostasis and the photoprotection of photosynthesis. Four NDH-1 types haven been characterized in cyanobacteria: NDH-1_1 and NDH-1_2, which function in respiration; and NDH-1_3 and NDH-1_4, which function in CO_2 uptake. All four types are involved in cyclic electron transport. Along with single FDP mutants ($\Delta flv1$ and $\Delta flv3$) and the double NDH-1 mutants ($\Delta d1d2$, which is deficient in NDH-$1_{1,2}$ and $\Delta d3d4$, which is deficient in NDH-$1_{3,4}$), we studied triple mutants lacking either one of Flv1 or Flv3, and NDH-$1_{1,2}$ or NDH-$1_{3,4}$. We show that the presence of either Flv1/3 or NDH-$1_{1,2}$, but not NDH-$1_{3,4}$, is indispensable for survival during changes in growth conditions from high CO_2 /moderate light to low CO_2 / high light. Our results suggest functional redundancy and crosstalk between FDPs and NDH-$1_{1,2}$ under the studied conditions, and demonstrate that the functions of FDPs and NDH-$1_{1,2}$ are dynamically coordinated for the efficient oxidation of PSI and for photoprotection under variable CO_2 and light availability.

KEYWORDS: Alternative electron transfer, Flavodiiron proteins, Flv, Mehler-like reaction, NDH-1, photosynthesis, photoprotection, Synechocystis.

INTRODUCTION

Photosynthetic organisms have evolved a variety of different regulatory mechanisms which are important for the protection of the photosynthetic machinery during rapid changes in environmental conditions. Fluctuating light intensities present particular risk to the photosystems due to over-reduction of the photosynthetic electron transport chain (PETC), particularly Photosystem (PS) I (Allahverdiyeva et al., 2015; Tiwari et al., 2016; Shimakawa et al., 2016). To counter this, cyanobacteria, algae, and plants (excluding angiosperms) employ C-type flavodiiron proteins (FDPs) as a strong photoprotective electron sink, directing excess photosynthetic electrons from

downstream of PSI to O_2 (Helman et al. 2003, Allahverdiyeva et al., 2013; Gerotto et al., 2016; Ilik

et al., 2017; Chaux et al., 2017; Jokel et al., 2018; Alboresi et al., 2019). This process is referred to

as the Mehler-like reaction and, in contrast to the Mehler reaction (Mehler 1957), does not produce

harmful reactive oxygen species (ROS) (Vicente et al., 2002; Brown et al., 2019).

The genome of the β-cyanobacterium *Synechocystis* sp. PCC 6803 (hereafter *Synechocystis*)

encodes four isoforms of FDPs, namely Flv1-4, which function in O_2 photoreduction mainly as

hetero-oligomers consisting of either Flv1 and Flv3 or Flv2 and Flv4 (Zhang et al., 2004; Mustila et

al., 2016; Santana-Sanchez et al., 2019). Recently, we showed that under air level [CO_2], Flv1/3 and

Flv2/4 hetero-oligomers function in a coordinated and interdependent manner: Flv1/3 stimulates

strong, but transient O_2 photoreduction upon the onset of illumination or during increases in light

intensity, while Flv2/4 hetero-oligomers catalyze a slower and limited steady state reduction of O_2,

also using electrons downstream of PSI (Santana-Sanchez et al., 2019). In elevated [CO_2] and in air

[CO_2] at alkaline pH 9, the *flv4-flv2* operon encoding Flv2, Flv4, and the Sll0218 proteins is

downregulated (Zhang et al., 2009; Santana-Sanchez et al., 2019). Nevertheless, a low level of

Flv1/3 in elevated [CO_2] stimulates strong steady state O_2 photoreduction, whereas in air [CO_2] at

pH 9, Flv1/3 is solely responsible for strong but transient O_2 photoreduction (Santana-Sanchez et

al., 2019). Whilst *in vitro* assays showed that homo-oligomers of recombinant Flv1, Flv3 and Flv4

can reduce O_2 with NADH and/or NADPH (Vicente et al. 2002, Shimakawa et al. 2015, Brown et

al. 2019), FNR and reduced Fd are yet to be considered as possible donors to FDPs. Moreover, the

use of *Synechocystis* mutants solely overexpressing Flv1 or Flv3 clearly demonstrated that, in

contrast to *in vitro* experiments, homo-oligomers of Flv3 or Flv1 are not involved in O_2

photoreduction *in vivo* (Mustila et al. 2016). Thus, the electron donor to FDPs remains to be

elucidated *in vivo*.

In cyanobacteria, photosynthetic and respiratory electron transfer reactions coexist on thylakoid

membranes and are intricately linked, necessitating strict regulation (Mullineaux, 2014). The

NAD(P)H dehydrogenase-like complex 1 (NDH-1, photosynthetic complex I) is mainly localized in

thylakoid membranes and involved in several bioenergetic reactions including cyclic electron

transfer around PSI (CET), respiration and CO_2 acquisition via the carbon concentrating mechanism

(CCM) (for reviews see Battchikova et al. 2011; Ma and Ogawa 2015; Burnap et al. 2015; Peltier et

al. 2016). The NDH-1 complex is present as several forms with distinct physiological roles: the

NDH-1_1 type features the subunits NdhD1 and NdhF1 and functions as Complex I of the respiratory

electron transfer chain (Zhang et al., 2004; He et al., 2015; Saura and Kaila, 2019); the NDH-1_2

form features the NdhD2 subunit instead of NdhD1, but it remains unclear whether NDH-1$_1$ and

NDH-1$_2$ serve different physiological functions (Peltier et al., 2016). The NDH-1$_3$ and NDH-1$_4$

forms that comprise low-Ci-inducible, high affinity CO_2 acquisition subunits NdhD3, NdhF3,

CupA, and CupS, or constitutive, low-affinity CO_2 acquisition subunits NdhD4, NdhF4, and CupB,

respectively, function in conversion of CO_2 to HCO_3^- (Ohkawa et al., 2000a,b; Shibata et al., 2001;

Zhang et al., 2004; Burnap et al., 2013; Han et al., 2017; Schuller et al., 2019). This process is

possibly enabled by alkaline pockets at the thylakoid membrane created by NDH-1 proton pumping

(Kaplan and Reinhold, 1999). All four NDH-1 types catalyze CET from ferredoxin (Fd) to

plastoquinone (PQ), which is coupled to the pumping of $4H^+/2e^-$ into the thylakoid lumen (He et al.,

2015; Laughlin et al., 2019; Schuller et al., 2019; Saura and Kaila, 2019). However, for NDH-1$_{3,4}$

this is to a lesser extent than for NDH-1$_{1,2}$ (Bernat et al., 2011).

In the current study, we aim to elucidate how the electron transport pathways mediated by FDPs

and NDH-1 cooperate to allow the maintenance of redox poise between the PETC, respiration,

CCM, and CO_2 fixation in the Calvin-Benson-Bassham cycle (CBB) under variable light and C_i

availability. To this end, we employed both biophysical and biochemical methods to characterize

various *Synechocystis* mutant strains with combined deficiencies of both FDP and NDH-1

pathways. Our results provide convincing evidence for the presence of either Flv1/3 or NDH-1$_{1,2}$,

but not NDH-1$_{3,4}$, being indispensable for the survival of *Synechocystis* cells under transitions from

high [CO_2] conditions to the combined stress conditions of air [CO_2] and high light. We show that a

dynamically coordinated and cooperative function of Flv1/3 and NDH-1$_{1,2}$ is required for the

photoprotection of the photosynthetic apparatus of *Synechocystis* cells and discuss the molecular

mechanisms involved.

RESULTS

Simultaneous inactivation of Flv1/3 and NDH-1$_{1,2}$ is lethal upon shift from high to air [CO_2]

and high light

In order to examine whether the simultaneous inactivation of NDH-1 and FDPs has an adverse effect

on cell survival, we monitored the growth of *Synechocystis* wild-type (WT) and various mutant

strains lacking (i) either Flv1 or Flv3 ($\Delta flv1$ and $\Delta flv3$, respectively, Helman et al., 2003), (ii) both

NdhD1 and NdhD2 ($\Delta d1d2$, deficient in NDH-1$_1$ and NDH-1$_2$, Ohkawa et al., 2000), (iii) both NdhD3

108 and NdhD4 ($\Delta d3d4$, deficient in NDH-1$_3$ and NDH-1$_4$, Ohkawa et al., 2000), as well as triple mutants

109 with combined deficiencies of both pathways ($\Delta flv1\ d1d2$, $\Delta flv3\ d1d2$, $\Delta flv1\ d3d4$, $\Delta flv3\ d3d4$) in

110 conditions of differing CO_2 availability and light intensity. The triple mutants exhibited similar or

111 slightly slower growth compared to the WT and other mutant strains under high [CO_2] (3% CO_2) and

112 moderate light (ML, intensity of 50 μmol photons m^{-2}s^{-1} PAR) (Fig. 1A). We also detected no

113 substantial difference in growth when the 3% CO_2/ML- pre-grown cells were diluted (OD$_{750}$=0.1)

114 and shifted to air [CO_2] conditions under constant light intensity (Fig. 1B, Fig. S1A, Fig. S1C). In

115 contrast, when cultures pre-grown in 3% [CO_2]/ML were diluted (OD$_{750}$=0.1) and subjected

116 simultaneously to air [CO_2] and high light stress conditions (HL, intensity of 220 μmol photons m^{-2}s^{-}

117 ^{1}), the growth of both $\Delta flv3\ d1d2$ and $\Delta flv1\ d1d2$ strains was completely inhibited (Fig. 1D).

118 Importantly, $\Delta flv1$, $\Delta flv3$ and $\Delta d1d2$ strains demonstrated growth similar to WT cells under air [CO_2]

119 /HL conditions.

120 We next examined whether pre-adaptation to air [CO_2] would rescue the growth arrest. When cells

121 previously adapted to air [CO_2] /ML were diluted (OD$_{750}$=0.1) and shifted to air [CO_2] /HL, no

122 differences in growth patterns between the strains were detected (Fig. 1C). Importantly, the triple

123 mutants lacking either Flv1 or Flv3 and NDH-1$_{3,4}$ ($\Delta flv1\ d3d4$ and $\Delta flv3\ d3d4$ strains) did not show

124 a lethal phenotype after a shift from 3% [CO_2]/ML to air [CO_2]/HL, although a somewhat reduced

125 growth was observed compared to the control strains (Fig. S1B, Fig. S1D). This suggests that, under

126 the studied conditions, crosstalk between Flv1/3 and NDH-1$_{3,4}$ complexes is not as crucial as with

127 NDH-1$_{1,2}$. Our attempts to create M55/$\Delta flv1$ and M55/$\Delta flv3$ double mutants lacking either Flv1 or

128 Flv3 and the central membrane component of the NDH-1 complex, NdhB, were both unsuccessful.

129 We were unable to obtain any M55/$\Delta flv1$ colonies and the M55/$\Delta flv3$ mutant strain demonstrated

130 poor growth and was not segregated, suggesting that the absence of both Flv1/3 and NDH-1, disrupts

131 essential cell metabolism.

132 We also studied the growth of the mutant strains on agar plates containing BG-11 (Fig. 1E). For these

133 experiments, 3% CO_2/ML-grown cells were diluted and grown on plates either under air [CO_2] / ML

134 or air [CO_2] / HL conditions for 7 days. The $\Delta flv1$ and $\Delta flv3$ and $\Delta d1d2$ mutants did not exhibit any

135 visible differences in growth capacity under these conditions in comparison to the WT. In contrast,

136 the growth of the $\Delta flv1\ d1d2$ and $\Delta flv3\ d1d2$ triple mutants was strongly reduced under air [CO_2] /

137 ML, while no growth was detected under air [CO_2] / HL (Fig. 1E). The $\Delta flv1\ d3d4$ and $\Delta flv3\ d3d4$

triple mutants demonstrated a slow growth phenotype, similarly to that previously reported for $\Delta d3d4$

on agar plates (Ohkawa et al. 2000b).

As the growth phenotypes of $\Delta flv1$ and $\Delta flv3$, $\Delta flv1$ $d1d2$ and $\Delta flv3$ $d1d2$ were similar under all

conditions, we only included $\Delta flv3$ and $\Delta flv3$ $d1d2$ in the majority of subsequent experiments.

Figure 1. Growth of WT and mutant strains under different experimental conditions. All experimental
cultures were started from pre-cultures adjusted to OD_{750}=0.1. **(A)** Cells were pre-cultivated under 3% $[CO_2]$
/ ML (50 μmol photons $m^{-2}s^{-1}$), adjusted to OD_{750}=0.1 and grown in the same conditions. **(B)** Cells were pre-
cultivated under 3% $[CO_2]$/ML and shifted to air $[CO_2]$/ML. **(C)** Cells were pre-cultivated under air
$[CO_2]$/ML and shifted to air $[CO_2]$/HL 220 μmol photons $m^{-2}s^{-1}$). **(D)** Cells were pre-cultivated under 3%
$[CO_2]$/ML and shifted to air $[CO_2]$/HL. The values presented in A–D are means of three independent
biological replicates ±SE. **(E)** Growth of WT and mutant cells on agar plates containing BG-11. Cells grown
under 3% $[CO_2]$ / ML were harvested and spotted on BG-11 agar plates in ten-fold dilution series starting at
OD_{750}=1, incubation was under air $[CO_2]$ /ML and air $[CO_2]$ /HL for 7 days. See also Supplemental Fig. S1.

Either Flv1/3 or NDH-1$_{1,2}$ is required for the oxidation of PSI during sudden increases in light
intensity

The lethal phenotype observed upon the shift to air $[CO_2]$ / HL which was caused by the
simultaneous impairment of the Mehler-like reaction catalyzed by Flv1/3 (Helman et al. 2003,
Allahverdiyeva et al. 2011, 2013) and by NDH-1$_{1,2}$ complex mediated respiration and CET
(Ohkawa et al., 2000b, Bernat et al. 2011) suggests a functional redundancy between these electron
transport pathways. To further investigate this possibility, we used membrane inlet mass
spectrometry (MIMS) to examine the kinetics of O_2 and CO_2 exchange upon HL-illumination of
dark-adapted WT and mutant cells grown for four days in air $[CO_2]$ / ML. In order to distinguish O_2
uptake from photosynthetic gross O_2 evolution under illumination, we enriched cell suspensions
with $^{18}O_2$ prior to measurements. In dark-adapted WT cells, a transient peak in O_2 uptake occurs
during the first minute of illumination (Fig. 2A). We showed recently that this transient peak is
attributed to the Mehler-like reaction catalyzed predominantly by Flv1/3 hetero-oligomers, while
Flv2/4 hetero-oligomers mainly contribute to steady-state light-induced O_2 reduction in an
interdependent manner (Santana-Sanchez et al., 2019). In $\Delta flv3$ cells, light-induced O_2 reduction
was almost abolished and only a slight impairment of the rate of photosynthetic gross O_2 evolution
was observed in comparison to WT cells (Fig. 2B). This result is in agreement with previous studies
(Helman et al., 2003; Allahverdiyeva et al., 2013; Santana-Sanchez et al., 2019).

In contrast to the WT, $\Delta d1d2$ cells lacked a fast decay phase of light-induced O_2 uptake during the
first minute of illumination, demonstrating sustained O_2 photoreduction at high levels (c.a. 75–100
μmol O_2 mg Chl^{-1} hr^{-1}) throughout the illumination period. Meanwhile, gross O_2 evolution was

diminished nearly two-fold compared to the WT (Fig. 2C). Thus, sustained O_2 photoreduction in

$\Delta d1d2$ is not due to increased electron flow from PSII, but most likely due to increased activity of

FDPs. Indeed, $\Delta flv3\ d1d2$ mutant cells showed only minor light-induced O_2 uptake transiently

during the first minute of illumination (Fig. 2D). The $\Delta flv3\ d1d2$ cells demonstrated slower

induction of photosynthetic O_2 evolution following two-component kinetics, and slightly impaired

steady-state gross O_2 evolution compared to WT (Fig. 2D).

The initial peak in CO_2 uptake rate at the onset of illumination likely reflects activation of the CCM

(Liran et al., 2018). Accordingly, this initial peak is absent in the M55 mutant deficient in NDH-1$_{3,4}$

in addition to NDH-1$_{1,2}$, as well as in WT cells grown at pH 6 (Fig. S2). No salient impairment of

CCM was observed in any of the studied strains under air [CO_2] / ML, with CO_2 uptake rates

peaking at c.a. 1.5 mM CO_2 mg Chl^{-1} hr^{-1}. In WT as well as in $\Delta flv3$, CO_2 fixation is then induced

and the respiratory compensation point surpassed after ~1 min of illumination (Fig. 2A-B). In

$\Delta d1d2$, however, induction of CO_2 fixation is severely delayed, and the cells only reached the

compensation point at the end of the 5 min illumination period (Fig. 2C). Intriguingly, although a

slight delay was observed also in $\Delta flv3\ d1d2$ induction of CO_2 fixation was largely recovered in the

triple mutant (Fig. 2D).

Figure 2. O_2 and CO_2 exchange rates in WT and mutant strains. Cells were grown in air [CO_2] / ML for
4 days, after which the cells were harvested and Chl a concentration adjusted to 10 μg ml^{-1} with fresh BG-11.
Cells were dark-adapted for 15 min, and gas exchange was monitored by membrane inlet mass spectrometry
(MIMS) over a 5 min illumination period with 500 μmol photons m^{-2}s^{-1} of white actinic light. Prior to the
measurements, samples were supplemented with $^{18}O_2$ at an equivalent concentration to $^{16}O_2$ to distinguish O_2
uptake from O_2 evolution, and with 1.5 mM NaHCO$_3$. The dashed line in each panel indicates the
compensation point of CO_2 fixation (uptake rate equals respiratory rate). The experiment was repeated with
three independent biological replicates, of which representative measurements are shown.

These MIMS results indicated that combined deficiency of Flv1/3 and NDH-1$_{1,2}$ under standard
growth conditions, air [CO_2] / ML, slightly perturbs photosynthetic electron transfer and the redox
poise between the PETC and cytosolic sink reactions. Next, we examined the ability of the mutant
strains to adjust their photosynthetic activity to variable light conditions by measuring Chl a
fluorescence simultaneously with P700 redox changes in conditions where light intensity
periodically fluctuated between low (25 μmol photons m^{-2}s^{-1}, LL) and high irradiance (530 μmol
photons m^{-2}s^{-1}, HL). Cells lacking Flv3 suffered from transient PSI acceptor side limitation Y(NA)
upon sudden increases in light intensity, resulting in delayed oxidation of PSI (Fig. 3A), which is in
line with previous studies (Helman et al. 2003, Allahverdiyeva et al. 2013). During subsequent 1
min cycles of LL/HL, however, the ability of Δ$flv3$ to oxidize PSI in HL improved, suggesting that
cells were able to acclimate to the fluctuating light condition via a compensatory mechanism that is
distinct from Flv1/3 hetero-oligomers, such as $e.g.$ NDH-1 mediated electron transport.

We detected no acceptor-side limitation during the HL phases in Δ$d1d2$ (Fig. 3A) as PSI was
oxidized similarly to the WT (Fig. 3B). Interestingly, however, Δ$d1d2$ cells exhibited slightly
elevated acceptor-side limitation during the LL phases of the experiment (Fig. 3A), suggesting a
role for NDH-1$_{1,2}$ in maintaining photosynthetic redox poise in light-limited conditions. In the triple
mutant strain Δ$flv3\ d1d2$, transitions from low to high irradiance as well as from dark to light
caused severe limitation on the acceptor side of PSI, resulting in an inability to oxidize PSI during
periodic 1 min HL illumination. Unlike the Δ$flv3$ mutant, Δ$flv3\ d1d2$ did not show improvement in
the PSI acceptor side limitation during subsequent cycles. This implies a compensating activity of
NDH-1 in Δ$flv3$ mutant under the studied conditions. Interestingly, diminished donor side limitation
in Δ$flv3\ d1d2$ resulted in slightly elevated effective yield of PSI during the HL phases (Fig. 3C).
This could indicate the attenuation of pH-dependent limitation of electron transfer at Cyt b_6f, due to
impairment of proton motive force (pmf) generation in CET and the Mehler-like reaction. The

effective yield of PSII was slightly decreased in HL in Δ*flv3 and* Δ*d1d2,* but not in Δ*flv3 d1d2* (Fig. 3D).

Figure 3. Photosynthetic parameters in WT and mutants strains in fluctuating light intensities. (A) Acceptor side limitation of PSI, Y(NA), **(B)** donor side limitation of PSI, Y(ND), as well as **(C)** effective yield of PSI, Y(I) were calculated from absorbance difference changes at 875-830 nm according to (Klughammer and Schreiber, 2008). Chl *a* fluorescence changes were measured simultaneously with PSI redox changes and used to calculate **(D)** effective yield of PSII. Cells were grown in air [CO_2] / ML for 4 days, after which the cells were harvested and Chl *a* concentration adjusted to 10 μg ml^{-1} with fresh BG-11. Cells were dark-adapted for 10 min, and then illuminated with red actinic light intensity alternating between 25 and 530 μmol photons m^{-2}s^{-1} in 1 min periods. Saturating pulses (500 ms of 5000 μmol photons m^{-2}s^{-1}) were provided every 15 seconds. The values shown are averages from 3–4 independent biological replicates ±SE.

The differences in photosynthetic electron transport reported above may also be contingent on altered redox states of the NADP$^+$/ NADPH pool. Therefore, in order to investigate the effect of Flv3 and/or NDH-1$_{1,2}$ deficiency on NADP$^+$/NADPH redox kinetics, we recorded NADPH fluorescence changes from dark-adapted cells simultaneously with Chl *a* fluorescence in conditions where actinic light

intensity fluctuated between 25 (LL) and 530 µmol photons $m^{-2}s^{-1}$ (HL) similarly to Fig. 3. Upon

dark-to-LL transitions, NADPH rapidly accumulated close to a maximal amount in all strains. Re-

oxidation of NADPH then occurred in WT and Δ*flv3d1d2* cells, while in Δ*flv3* a slower oxidation

phase preceded a transient re-reduction phase during the first minute of illumination (Fig. 4A-B).

Strong reduction of the $NADP^+$ pool was also detected in Δ*flv3* during the second HL phase of the

experiment. In Δ*d1d2* cells, very little oxidation of the NADPH pool occurred during illumination,

even at HL-LL transitions (Fig. 4C). Chl *a* fluorescence also remained at an elevated level (Fig. 4C),

suggesting a reduced PQ pool. This was possibly due to delayed activation of CO_2 fixation in the

CBB cycle (Fig. 2C), and impaired CET or dark respiration. In contrast, NADPH was strongly

oxidized in Δ*flv3 d1d2* cells at HL-LL transitions (Fig. 4D), but was predominantly reduced in dark-

adapted cells (Fig. 4D). Upon cessation of illumination, oxidation of the NADPH pool was followed

by transient re-reduction after c.a. 10 s. As observed previously (Schreiber and Klughammer, 2009;

Holland et al., 2015), the re-reduction peak coincided with a secondary post-illumination rise in Chl

a fluorescence (PIFR), while oxidation of NADPH paralleled an initial PIFR. The post-illumination

re-reduction of NADP+ was diminished and both Chl *a* PIFR peaks were missing in Δ*d1d2* (Fig. 4C).

Therefore, the PIFR peaks likely originate from cyclic and respiratory electron transport via NDH-

$1_{1,2}$ (Holland et al., 2015). Both Chl PIFR peaks were however observed in Δ*flv3 d1d2* cells, indicating

involvement of a complementary pathway, possibly via succinate dehydrogenase (Cooley and

Vermaas, 2001), or a CET pathway dependent on PGR5 and PGRL1-LIKE (Yeremenko et al., 2005;

Dann and Leister, 2019). Finally, we monitored NADPH redox kinetics under conditions mimicking

the MIMS experiments, where strong O_2 photoreduction occurs in WT and Δ*d1d2* during the first

minute of dark-to-high light transition (Fig. 2). No substantial differences were observed between the

strains, as close to maximal reduction of $NADP^+$ was obtained within ~0.25 s (Fig. 4F) and

maintained throughout a 40 s illumination period (Fig. 4E). This argues against NADPH being a

primary electron donor to either Flv1/3 or NDH-$1_{1,2}$.

Figure 4. NADPH and Chl *a* fluorescence under fluctuating light in WT and mutant strains. WT (**A**), Δ*flv3* (**B**), Δ*d1d2* (**C**), and Δ*flv3 d1d2* cells (**D**) were grown in air [CO₂] / ML for 4 days, after which the cells were harvested and Chl *a* concentration adjusted to 5 μg ml⁻¹ with fresh BG-11. Cells were dark-adapted for 10 min, and then subjected to illumination with red actinic light alternating between 25 and 530 μmol photons m⁻²s⁻¹ in 1 min periods. NADPH redox changes were monitored with the Dual-PAM 100 spectrophotometer and its 9-AA/NADPH accessory module by measuring fluorescence changes between 420 and 580 nm induced by excitation at 365 nm, as described by (Schreiber and Klughammer, 2009) and

(Kauny and Setif, 2014). Chl *a* fluorescence was recorded simultaneously. **(E)** NADPH redox changes

during illumination of dark-adapted cells with 530 μmol photons m^{-2}s^{-1}. Other experimental details are as in

A-D. **(F)** shows a magnification of the first 2 s of illumination in (E). The values are normalized according to

minimum and maximum fluorescence changes at the onset and cessation of illumination, respectively. The

experiments were repeated with three independent biological replicates, of which representative

measurements are shown. r.u. = relative units.

Redox status of PSI donor and acceptor side electron carriers and buildup of *pmf* during

dark-to-light transitions depend on both Flv1/3 and NDH-1$_{1,2}$

To elucidate the molecular mechanism behind the photosynthetic phenotypes of the studied mutant

strains, we next utilized a DUAL-Klas-NIR 100 spectrophotometer to distinguish between redox

changes of plastocyanin (PC), P700, and ferredoxin (Fd) (Klughammer and Schreiber, 2016;

Schreiber, 2017; Setif et al., 2019) upon exposure of dark-adapted cells to high irradiance (503 μmol

photons m^{-2}s^{-1}). Upon illumination of WT cells, rapid oxidation of P700 and PC occurred, followed

by transient reduction of PC, P700, as well as Fd after ~0.2 s (Fig. 5A). Re-oxidation of all three

electron carriers then ensued after ~0.5 s. In Δ*flv3* cells P700 and PC remained mostly reduced after

the initial oxidation transient until ~3 s, when another transient and partial re-oxidation peak of P700

and PC occurred at ~8 s (Fig. 5B). This was followed by re-reduction of P700, similarly to that

observed recently by Bulychev et al. (2018). After 15 s in light, P700 and PC became gradually

oxidized. In contrast to WT, Fd remained reduced for several seconds in light, and was only slowly

re-oxidized over the 30 s illumination period (Fig. 5B). These observations indicate the importance

of Flv1/3 as an electron sink to O$_2$, accepting electrons presumably from reduced Fd (after ~0.5 s in

light). In Δ*d1d2*, redox changes of PC, P700 and Fd were similar to WT, except that the reduction of

P700$^+$ following initial oxidation as well as reduction of Fd occurred already after ~50 ms (Fig. 5C).

In Δ*flv3 d1d2*, re-reduction of P700$^+$ and reduction of Fd also occurred already after ~50 ms (Fig.

5D), suggesting involvement of NDH-1$_{1,2}$ as an acceptor of electrons from Fd at that stage. Fd was

reduced more quickly in Δ*flv3 d1d2* than in any of the other strains and was slowly re-oxidized over

30 s (Fig. 5D), likely due to shortage of electron acceptors (Fig. 3A). In summary, these NIR-

spectroscopic measurements revealed that Flv1/3 and NDH-1$_{1,2}$ control the redox poise between PSI,

PC, and primary electron acceptor Fd during specific time frames at transitions to high irradiance.

The results are congruent with Fd functioning as the electron donor to both NDH-1$_{1,2}$ and Flv1/3.

To inspect whether the phenotypes observed above depend on differential buildup or regulation of

pmf, we measured *in vivo* changes in the *pmf* by monitoring the absorbance change difference between

500-480 nm, which constitutes the electrochromic shift (ECS) in *Synechocystis* (Viola et al., 2019).

In vivo measurement of light-induced ECS revealed that after 1 s of illumination of dark-adapted WT

cells with 500 μmol photons $m^{-2}s^{-1}$, high *pmf* level was transiently generated, followed by decline

during subsequent seconds (Fig. 5E). After ~10 s of illumination *pmf* again increased towards a

steadier value (Fig. 5E). Congruently with the strong reduction of P700 and Fd (Fig. 5B), the initial

pmf peak after the first second of illumination was heavily dependent on the presence of Flv1/3. In

both Δ*flv3* and Δ*flv3 d1d2 pmf* and thylakoid proton flux (vH+) were lower than in WT after 1 s (Fig.

5G). In Δ*flv3 pmf* remained drastically lower than in WT during the first seconds of illumination, but

differed only slightly from WT thereafter due to diminished conductivity of the thylakoid membrane

(gH+) (Fig. 5F), which is mainly determined by the activity of the ATP synthase (Cruz et al., 2005;

Viola et al., 2019). In Δ*d1d2* thylakoid proton flux was similar to WT (Fig. 5G), likely due to

enhanced FDP activity (Fig. 2C) compensating for impaired CET and respiration. However, *pmf*

remained lower than in WT throughout the 1 min experiment (Fig. 5E) because of elevated

conductivity (Fig. 5F). In contrast, despite drastically lower proton flux (Fig. 5G), Δ*flv3 d1d2* cells

maintained *pmf* close to WT level after the first seconds in light (Fig. 5E) due to diminished

conductivity (Fig. 5F). This indicates that the increased acceptor side limitation of PSI in Δ*flv3 d1d2*

in comparison to Δ*flv3* (Fig. 3A) is not caused by lack of photosynthetic control as a consequence of

lowered *pmf* generation in CET. Based on comparison of ECS kinetics in single, double, and triple

mutant cells studied here, we conclude that both Flv1/3 and NDH-1$_{1,2}$ contribute to proton flux during

transitions from dark to high irradiance. However, deficiency in NDH-1$_{1,2}$ in Δ*d1d2* is mostly

compensated by elevated FDP-activity in terms of proton flux, while ATP synthase activity is

increased, possibly as a response to the delay in activation of carbon fixation (Fig. 2C). In contrast,

impaired thylakoid proton flux in Δ*flv3* during dark-to-light transitions cannot be compensated by

NDH-1. Instead, downregulation of ATP-synthase activity lowers thylakoid conductivity and allows

maintenance of *pmf* (apart from the first seconds of illumination) in Δ*flv3* and, even more

dramatically, in Δ*flv3 d1d2*.

Figure 5. Redox changes of PC, P700, and Fd, and buildup of *pmf* during dark-to-HL transitions. WT
(**A**), Δ*flv3* (**B**), Δ*d1d2* (**C**), and Δ*flv3 d1d2* cells (**D**) were dark-adapted for 10 min, after which absorbance
difference changes at 780–800 nm, 820–870 nm, 840–965 nm and 870–965 nm were measured with the
Klas-NIR 100 spectrometer during 30 s illumination at 503 µmol photons m^{-2}s^{-1} of red actinic light. The PC,
P700, and Fd signals were deconvoluted from the absorbance difference changes using differential model
plots (DMPs) determined for this study (Fig. S3). The upper panels in A–D show magnifications of the first
1.5 seconds of illumination from the lower panels. The experiment was repeated with three independent
biological replicates, of which representative traces are shown. See also Supplemental Fig. S3. (**E**) Buildup

of *pmf* during transitions from dark to 500 μmol photons m^{-2}s^{-1} of green actinic light. Absorbance difference

between 500-480 nm (ECS) was measured with a JTS-10 spectrometer, and light-induced *pmf* was

determined from the dark interval relaxation kinetics (DIRK) of the ECS signal. Means ±SEM from 3–4

independent experiments are shown. The small inset shows *pmf* values after 1s of illumination for a clearer

view. **(F)** Conductivity of the thylakoid membrane (gH+) after one and 30 s of illumination as in (E). gH+

was determined as the inverse of the time constant of first-order post-illumination decay kinetics of the ECS

signal. **(G)** Thylakoid proton flux (vH+), calculated as *pmf* x gH+. **(H)** Representative traces of 500-480 nm

absorbance differences used to determine the values in (E) and post-illumination ECS decay kinetics during a

500 ms dark interval after 30 s in light (lower panels). First-order fits are shown in green. Cells, after growth

in air [CO_2] / 50 μmol photons m^{-2}s^{-1} for 4 days, were harvested and Chl *a* concentration adjusted to 10 μg

ml^{-1} with fresh BG-11 for A–D and to 7.5 μg ml^{-1} for E–J.

PSI content is diminished in triple mutants deficient in Flv1/3 and NDH-1$_{1,2}$ shifted to air [CO_2] and high irradiance

The results from the growth assays, along with the real-time gas exchange, P700 redox change, and

ECS measurements suggested that in Δ*d1d2* cells, an increase in the activity of FDPs compensates

for the absence of a functional NDH-1$_{1,2}$ complex, allowing efficient oxidation of PSI under high

irradiance. Conversely, in Δ*flv1* and Δ*flv3* mutants, over-reduction of the electron transport chain

occurs initially upon exposure to high irradiance, but another mechanism(s) eventually allow

acclimation and survival in high [CO_2] and high light (Helman et al., 2003). Under air [CO_2] Flv1/3

hetero-oligomers catalyse transient O_2 photoreduction, which is why no strong growth phenotype is

observed in Δ*flv1* or Δ*flv3* under constant high irradiance (Allahverdiyeva et al. 2013, Santana-

Sanchez et al. 2019). However, when both Flv1/3 and NDH-1$_{1,2}$ are absent and the electron sink

capacity of the cytosol is not elevated by high [CO_2], cells are unable to oxidize PSI in high light,

possibly resulting in lethal photodamage to PSI. To assess this hypothesis, we employed

immunoblotting to provide estimates of the protein content of PSII (D1) and PSI (PsaB) subunits,

orange carotenoid protein (OCP), large isoform of the ferredoxin-NADP+ oxidoreductase (FNR$_L$),

FDPs, as well as NdhD3 and the bicarbonate transporter SbtA after 24 hours exposure to different

growth conditions.

WT and Δ*flv1* or Δ*flv3* cells grown at 3% [CO_2] / ML and shifted to air [CO_2] / ML at OD=0.1,

accumulated similar levels of the PSII reaction center protein D1 and the PSI subunit PsaB.

371 However, we detected a moderate decrease in the amount of both D1 and PsaB in the $\Delta flv1\ d1d2$

372 and $\Delta flv3\ d1d2$ triple mutants, while an increased amount of both proteins was detected in $\Delta d1d2$

373 (Fig. 6A). After 24 hours exposure to air [CO$_2$] / HL, PsaB amount had decreased even further in

374 $\Delta flv1\ d1d2$ and $\Delta flv3\ d1d2$ (Fig. 6C). Furthermore, 77K fluorescence spectra revealed that while

375 $\Delta flv3\ d1d2$ and WT cells grown under air [CO$_2$] / ML had similar PSI/PSII ratios (Fig. 6D), the 24-

376 hour exposure to air [CO$_2$] / HL caused a dramatic decrease in the relative PSI fluorescence cross

377 section (Fig. 6E). This strongly supports the hypothesis that loss of PSI contributes to the lethality

378 of the shift to air [CO$_2$] / HL. Interestingly, a substantial decrease in D1 content in air [CO$_2$] / HL

379 was also observed in the triple mutants (Fig. 6B-C).

380 Importantly, the $\Delta flv1\ d1d2$ and $\Delta flv3\ d1d2$ strains were unable to induce substantial accumulation

381 of the proteins encoded by the $flv4$-2 operon after a shift from 3% to air [CO$_2$] (Fig. 6A and Fig.

382 6C). Only when pre-grown for 4 days in air [CO$_2$] / ML before being shifted to higher irradiance,

383 the expression of the operon was induced (Fig. 6B). Similarly to $flv4$-2 operon proteins, both

384 NdhD3 and SbtA failed to accumulate after shifts from 3% to air [CO$_2$] / HL in the triple mutants,

385 which likely impairs CCM and contributes to the lethal phenotype of the triple mutants in those

386 conditions. Only small amounts of NdhD3 and SbtA were detected also in triple mutant cells shifted

387 from 3% [CO$_2$] to air [CO$_2$] / ML (Fig. 6A). Closer to WT amount of NdhD3 but only a small

388 amount of SbtA was detected after 24 hours in air [CO$_2$] / HL (shifted from air [CO$_2$] / ML) (Fig.

389 6B). In the $\Delta d1d2$ mutant NdhD3 content was similar to WT, but Flv3 as well as the $flv4$-2 operon

390 proteins (Flv2, Flv4, and Sll0218) were upregulated in all conditions tested (Fig. 6), which likely

391 contributes to the increased rate of O$_2$ photoreduction in that strain (Fig. 2C). The amounts of OCP

392 and of FNR$_L$ were unchanged in the mutant strains in all conditions (Fig. 6).

Figure 6. Immunodetection of proteins related to photosynthesis and CCM and 77K fluorescence emission spectra in WT and mutant strains. Pre-cultures were grown for 3 days at 3% [CO_2] / ML (50 μmol photons m^{-2}s^{-1}) and then pelleted, resuspended in fresh BG-11 at OD$_{750}$=0.15 and **(A)** incubated for 24 hours in air [CO_2] / ML, **(B)** grown for 4 days in air [CO_2] / ML, harvested, resuspended in BG-11 at OD$_{750}$=0.15 and then incubated for 24 hours in air [CO_2] / HL (220 μmol photons m^{-2}s^{-1}), **(C)** incubated for 24 hours in air [CO_2] / HL. Total protein extracts were separated by SDS-PAGE and probed with specific antibodies. The immunoblots shown are representative of 2–3 independent biological replicates. **(D)** 77K fluorescence emission spectra from intact WT and Δ*flv3 d1d2* cells grown for 4 days in air [CO_2] / ML. **(E)** After pre-growth at 3% [CO_2] / ML cells were pelleted, resuspended in fresh BG-11 at OD$_{750}$=0.15 and incubated for 24 hours in air [CO_2] / HL. The cells (D and E) were pelleted, adjusted to [Chl *a*] of 7.5 μg ml^{-1}, and excited with 440 nm light at 77K. The spectra were normalized to the PSII fluorescence peak at 685 nm.

DISCUSSION

Flv1/3 hetero-oligomers have been shown to be essential for photoreduction of O_2 as a rapid response to excessive reduction of PSI, and consequently, for cell survival under fluctuating light conditions (Allahverdiyeva et al., 2013; Santana-Sanchez et al., 2019). Moreover, we recently showed that in air [CO_2], Flv2/4 hetero-oligomers also catalyze low level light-induced steady-state reduction of O_2 on the acceptor side of PSI in a coordinated manner with Flv1/3 (Santana-Sanchez et al., 2019).

In turn, another alternative electron transport component, the NDH-1 complex, functions in CET around PSI, using Fd as electron donor (Saura and Kaila, 2019; Schuller et al., 2019). Besides CET, the NDH-$1_{1,2}$ complexes, incorporating the NdhD1 or NdhD2 subunit, are involved in respiration, whereas NDH-$1_{3,4}$ complexes, incorporating the NdhD3 or NdhD4 subunit, function in CO_2 uptake. In the current study, we have used *Synechocystis* mutant strains with different combinations of deficiencies in Flv1/3 and NDH-1 in order to examine the physiological significance of these alternative electron transport pathways and the possibility of functional interdependence between them. Our results provide compelling evidence for the concomitant physiological functions of Flv1/3 and NDH-$1_{1,2}$ and their dynamic coordination for the efficient oxidation of PSI (thus protecting it from photodamage) under variable light conditions, and when low [CO_2] limits the consumption of reductants in the CBB cycle. The two pathways can compensate for each other's absence to some extent, but absence of both the Flv1/3 hetero-oligomer and NDH-$1_{1,2}$ is lethal when cells are transferred from elevated [CO_2] to the combined condition of air level [CO_2] and high irradiance (Fig. 1D-E). In addition to the absence of these two important pathways, lethality is due to an inability to accumulate low C_i-inducible photoprotective CCM proteins (Fig. 6C). Combined deficiency of Flv1/3 and NDH-$1_{3,4}$ did not result in a lethal phenotype upon similar shifts, indicating that functional redundancy exists specifically between the Mehler-like reaction and NDH-$1_{1,2}$

Coordinated functions of Flv1/3 and NDH-$1_{1,2}$ protect PSI by maintaining redox poise between the PETC and carbon fixation

Cells lacking functional Flv1/3 hetero-oligomers suffer from a transient over-reduction of PSI during sudden increases in light intensity (Fig. 3 and Fig. 5) due to the impairment of the Mehler-

like reaction (Fig. 2, Allahverdiyeva et al., 2013). Whilst NAD(P)H has been proposed as the

electron donor to Flv3 (Vicente et al., 2002; Brown et al., 2019) and Flv1 homo-oligomers (Brown

et al., 2019), *in vivo* experiements have been unsupportive (Mustila et al. 2016), with exact electron

donor(s) to FDP hetero-oligomers yet to be proven. *In vivo* experiments performed in this study

demonstrate that the quick re-oxidation of Fd after 0.5 s (Fig. 5A) in light is absent in the $\Delta flv3$

deletion strains, whereby Fd remains strongly reduced (Fig. 5B). This result, together with a lack of

impairment in $NADP^+$ reduction and oxidation kinetics of $\Delta flv3$ deletion strains (similar conditions,

Fig. 4F) provide strong support for Fd, rather than NAD(P)H, being the primary electron donor to

the Flv1/3 hetero-oligomer *in vivo*. Accordingly, Fd has been shown to interact with Flv1 and Flv3

by Fd-chromatography (Hanke et al., 2011) and with Flv3 by a two-hybrid assay (Cassier-Chauvat

and Chauvat, 2014).

The deficiency of NDH-$1_{1,2}$, in turn, reduced the cells' ability to maintain oxidized P700 and Fd

under high irradiance soon after (~50–200 ms) a dark-to-light transition (Fig. 5C). This brief

timescale may be due to PSI-NDH-1 supercomplexes (Gao et al., 2016) where P700 can be

oxidized rapidly upon the onset of illumination. It has been shown that formation of the NDH-1-PSI

supercomplex is important to keep PSI functional under various stress conditions (Zhao et al. 2017).

NDH-$1_{1,2}$ -deficiency also caused slightly elevated acceptor side limitation of PSI under low

irradiance (Fig. 3A), which was likely due to a lack of NADPH oxidation during the low light

phases of fluctuating light (Fig. 4C). Thus, NDH-$1_{1,2}$ appears to play an important role under low

light, as has been previously suggested for chloroplastic NDH in angiosperms (Yamori et al., 2011;

Yamori et al., 2015) and bryophytes (Ueda et al., 2012). The delayed activation of CBB in $\Delta d1d2$

did not result in an inability to oxidize PSI (Fig. 3B and Fig. 5C), likely due to significant

enhancement of FDP-mediated O_2 photoreduction (Fig. 2C) providing an enlarged electron sink for

the photosynthetic electron transport chain.An opposite order of causation is also possible, whereby

the excessive funneling of photosynthetic electrons to O_2 would cause the delay in induction of CO_2

fixation in $\Delta d1d2$. However, the observation that there is no NADPH shortage at the onset of

illumination in $\Delta d1d2$, and rather the consumption of NADPH during transitions from HL to LL is

impaired (Fig. 4C), suggests that the availability of reductant for the CBB is not the limiting factor.

Nor is it likely CO_2, as CCM is functioning as in WT (Fig. 2C), or ATP, as ATP synthase activity

was even higher than in WT in $\Delta d1d2$ (Fig. 5F). Nevertheless, simultaneous deficiency of FDPs in

addition to NDH-$1_{1,2}$ (in $\Delta flv3\ d1d2$ triple mutant), with the effect of diminishing the flow of

photosynthetic electrons to O_2 photoreduction, mostly rescued the delay in CO_2 fixation (Fig. 2D)

as well as NADPH consumption (Fig. 4D) seen in Δ*d1d2*. Transient O_2 photoreduction at a low rate

was still observed in Δ*flv3 d1d2* during dark-to-HL transitions (Fig. 2D), possibly mediated by the

thylakoid terminal oxidases (Ermakova et al., 2016) or by photorespiration (Allahverdiyeva et al.,

2011). However, the triple mutants suffered from more severe inability to oxidize PSI than Δ*flv3*

during sudden increases in irradiance (Fig. 3). These observations suggest that in addition Flv1/3,

also NDH-1$_{1,2}$ has a role in contributing to oxidation of PSI during changes in light conditions or

carbon availability. Upon deficiency of NDH-1$_{1,2}$ (in Δ*d1d2*), cells prioritize protection of PSI over

efficient CO_2 fixation by upregulating the Mehler-like reaction via an unknown mechanism. The

triple mutants cannot do this, leading to timely induction of CO_2 fixation at the high cost of inability

to oxidize PSI (Fig. 3A and 3) or Fd (Fig. 5D). This results in loss of PSI (Fig. 6), possibly due to

photodamage to its FeS clusters (Tiwari et al., 2016; Shimakawa et al., 2016).

On the mechanism of NDH-1-mediated oxidation of PSI

Contribution of NDH-1 to oxidation of PSI during sudden increases in light intensity is not

unprecedented. In angiosperms, where FDPs have been lost during evolution (Ilik et al., 2017),

Arabidopsis mutants lacking the NDH complex show a delay in oxidation of PSI during increases in

light intensity in comparison to *Arabidopsis* WT (Nikkanen et al., 2018; Shimakawa and Miyake,

2018a). By definition, canonical CET cannot directly increase the relative proportion of oxidized

P700, as electrons from the acceptor side of PSI are shunted back to the intersystem electron

transfer chain. However, NDH-1 may enhance PSI oxidation by at least three alternative, but

mutually non-exclusive mechanisms.

i)   NDH-1-mediated CET is coupled to translocation of protons from cytosol to the thylakoid

489   lumen with a $2H^+/e^-$ stoichiometry (Strand et al., 2017; Saura and Kaila, 2019). It will

490   therefore contribute to buildup of ΔpH, which limits electron transfer to PSI by inhibiting

491   PQH_2 oxidation at Cyt *b6f* (Shimakawa and Miyake, 2018b) and drives ATP synthesis to

492   accommodate the needs of the CBB, increasing its electron sink capacity. As the Mehler-

493   like reaction also contributes to buildup of ΔpH by consuming H^+ on the cytosolic side of

494   the thylakoid membrane and by supporting linear electron flow (Fig. 5), (Allahverdiyeva et

495   al., 2013), enhanced FDP activity in Δ*d1d2* (Fig. 2C) partly compensates for the lack of

496   NDH-1$_{1,2}$ in respect to generation of proton flux (Fig. 5G). However, this fails to explain

497	the exacerbated impairment of P700 and Fd oxidation in Δ*flv3 d1d2* in comparison to Δ*flv3*
498	(Fig. 3 and Fig. 5) because in the triple mutant *pmf* was not lower than in Δ*flv3* (Fig. 5E).
499	Adequate *pmf* is maintained however at the expense of ATP synthase activity (Fig. 5F), and
500	it is likely that diminished ATP production contributes to the increased acceptor side
501	limitation of PSI in Δ*flv3 d1d2* (Fig. 3A). NDH-1$_{1,2}$ could therefore contribute to P700
502	oxidation by enhancing cytosolic sink capacity by providing a more suitable ATP:NADPH
503	ratio for the CBB. Adjustment of the ATP:NADPH ratio closer to the theoretically optimal
504	3:2 has long been considered a fundamental reason for the existence of CET (Kramer et al.,
505	2004; Yamori and Shikanai, 2016).

ii)   NDH-1-mediated respiratory electron transfer, *i.e.* coupling of PQ reduction by NDH-1 to

O_2 reduction by thylakoid terminal oxidases (*i.e.* Cyd and Cox) would also contribute to

oxidation of PSI by relieving electron pressure in the intersystem chain during illumination

(Ermakova et al., 2016), as well as further contribute to ΔpH by consuming H^+ on the

cytosolic side of thylakoids (Cyd) and pumping protons to the lumen (Cox) (Brändén et al.,

2006). Interestingly, Liu and colleagues (2012) have shown that the subcellular localization

of NDH-1 complexes is dependent on the redox state of the PQ pool. An oxidized PQ pool

causes NDH-1 to accumulate at specific clusters in thylakoid membranes where it would be

more likely to transfer electrons (via the PQ pool) to a terminal oxidase. A reduced PQ

pool, in turn, results in a more even distribution of NDH-1 within thylakoids (Liu et al.,

2012). However, it is important to note that terminal oxidases do not have a high electron

sink capacity (Ermakova et al., 2016).

iii)   NDH-1 has been predicted, albeit not yet experimentally shown in photosynthetic

organisms, to be able to function in reverse: to oxidize PQH$_2$ driven by concomitant release

of protons from the thylakoid lumen (Strand et al., 2017). Such reverse activity would

constitute a 'pseudo-linear' electron transfer pathway that would bypass PSI and thereby

prevent its over-reduction. This could occur in conditions where the PQ pool is reduced,

pmf is high, and the Fd pool is oxidized (Strand et al., 2017). Such conditions likely exist

transiently during dark-to-light and LL-to-HL transitions (Fig. 4 and 5) (Strand et al.,

2019). Accordingly, fast re-reduction of P700$^+$ already after 50 ms was observed in Δ*d1d2*

and Δ*flv3 d1d2* during dark-to-HL transitions (Fig. 5C). As supported by the impaired

ability to oxidize Fd in absence of Flv1/3 (Fig. 5), the presence of Flv1/3-catalyzed Mehler-

like reaction would likely be essential in this model in order to maintain the Fd pool in a

sufficiently oxidized state to provide electron acceptors for reverse-functioning NDH-1. It

is noteworthy, however, that NDH-1 reverse activity would also have the effect of lowering

ΔpH, thereby relieving photosynthetic control at Cyt *b6f*. This could counteract the effect of

any reverse NDH-1 activity by increasing electron flow to PSI.

Hypothetical mechanisms for coordination of Flv1/3 and NDH-$1_{1,2}$ activities are shown in Fig. 7.

Inability to induce a strong CCM network contributes to lethality upon shifts to air [CO₂] /

high irradiance in cells deficient in both Flv1/3 and NDH-$1_{1,2}$

Exposure of *Synechocystis* cells to low [CO₂] induces expression of high-affinity CCM-related

genes such as SbtA, a HCO_3^- / Na^+ symporter on the plasma membrane, and NdhD3, which is part

of the NDH-1_3 complex on thylakoids specializing in C_i uptake (Ohkawa et al., 2000; Shibata et al.,

2001, Shibata et al., 2002; Zhang et al., 2004). CCM is energetically expensive, but the large C_i flux

involved in the operation of CCM contributes to dissipation of excess light energy under stress

conditions (Xu et al., 2008; Burnap et al. 2015). NDH-$1_{3,4}$ in particular has a photoprotective role,

using reduced Fd to drive CO_2 conversion to HCO_3^- with concomitant translocation of protons into

the thylakoid lumen.

Despite the low induction level of the CCM proteins NdhD3 and SbtA, the triple mutants (Fig. 6)

survive under standard growth conditions of air [CO₂] / ML (Fig. 1B-C). When a decrease in [CO₂]

is coupled with an increase in irradiance, however, a low amount of CCM, reflected by low

accumulation level of NdhD3 and SbtA in the Δ*flv3d1d2* and Δ*flv3d1d2* triple mutants (Fig. 6C)

fails to dissipate excess energy, thus resulting in photodamage and lethality of those conditions. The

regulatory mechanisms controlling the inability of the triple mutants to accumulate low Ci-inducible

proteins is unclear and remains to be elucidated. Accumulation of $NADP^+$ and α-ketogutarate

inhibits the induction of CCM gene expression via interaction with the transcription factor NdhR

(CcmR) (Daley et al., 2012). NdhR also controls expression of the Flv4-2 operon encoding Flv2,

Flv4, and Sll0218 (Eisenhut et al., 2012), all of which also showed decreased accumulation levels in

the triple mutants (Fig. 6). However, at least in the tested conditions, no substantial increase in the

relative amount of $NADP^+$ was detected in Δ*flv3d1d2* cells (Fig. 4). It is important to note however,

that Flv1/3 and NDH-1$_{3,4}$ cooperation alone is not crucial for cell metabolism, since the Δ*flv1 d3d4*

and Δ*flv3 d3d4* mutant cells do not show clear growth phenotypes (Fig. S1).

CONCLUSIONS

In the current study we have shown that FDPs and NDH-1 function cooperatively to maintain redox

balance between the PETC and cytosolic carbon assimilation upon sudden changes in light intensity

and/or carbon availability. It is likely that both pathways receive electrons primarily from Fd, which

enables coordinated regulation of their activities. Cooperation of FDPs and NDH-1 has an essential

photoprotective role during shifts to air [CO$_2$] /HL, contributing to efficient oxidation of PSI,

buildup of *pmf,* and induction of expression of low C$_i$-specific CCM-related genes. Finally, it is

worth noting that simultaneous removal of Flv1/3 and NDH-1$_{1,2}$ as competitive electron sinks

downstream of PSI may be a useful biotechnological tool for maximizing direction of

photosynthetic electrons to desired pathways under controlled environmental conditions

(McCormick et al., 2013; Thiel et al., 2019; Jokel et al., 2019). Furthermore, our findings about

functional interplay between FDPs and NDH-1 will be highly relevant in projects aiming to enhance

crop productivity via introduction of exogenous FDPs to higher plants (Yamamoto et al., 2016;

Gómez et al., 2018).

Figure 7. Schematic models for the coordination of the activities of FDPs and NDH-1 in preventing

over-reduction of PSI. (A) In conditions where the Fd pool is higly reduced and the PQ pool is not, NDH-

1L transfers electrons from Fd to the PQ pool, coupled to O$_2$ reduction by thylakoid-localised RTOs. NDH-

1L pumps 2H$^+$/e$^-$ into the lumen while FDPs and RTOs consume H$^+$ when reducing O$_2$ to H$_2$O on the

cytosolic side of thylakoids. This increases transthylakoid *pmf*, which in turn drives ATP synthesis as well as

inhibits PQH_2 oxidation at Cyt b_6f, thereby preventing excessive electron supply to PSI. Carbonic anhydrase

activity of NDH1-MS converts HCO_3^- to CO_2, consuming H^+ in the cytosol, likely also coupled to

translocation of H^+ to the lumen, further increasing *pmf*. Increased availability of CO_2 and ATP/NADPH

ratio enhances the electron sink capacity of the CBB on the acceptor side of the PETC. **(B)** If the PQ pool is

highly reduced, Fd pool is oxidized, and *pmf* is high, NDH-1L could hypothetically function in reverse,

transferring electrons from PQH_2 to Fd^+, powered by release of $2H^+/e^-$ from the lumen. A strong Mehler-like

reaction catalyzed by Flv1/3 would be required to maintain a sufficient pool of oxidized Fd, with Flv1/3

likely accepting electron directly from Fd^-. Most components depicted in (A) are omitted from (B) to

enhance clarity, not to suggest their absence. Extent of the colour fill for PQ/PQH_2 and Fd depict level of

reduction.

MATERIALS AND METHODS

Strains and culture conditions

In the current study we used the glucose-tolerant WT strain *Synechocystis* sp. PCC 6803 (Williams,

1988), single mutant strains Δ*flv1* and Δ*flv3* (Helman et al., 2003), the M55 mutant (Δ*ndhB*)

(Ogawa 1991), double mutants Δ*d1d2 and Δd3d4* (Ohkawa et al., 2000a), and triple mutants Δ*flv1*

d1d2, Δflv3 d1d2, Δflv1 d3d4, and Δ*flv3 d3d4* obtained from the CyanoMutants collection

(Nakamura et al., 1999). The triple mutants were constructed by T. Ogawa in Δ*d1d2 and Δd3d4*

backgrounds. All the mutants demonstrated complete segregation. Pre-experimental cultures were

always grown in 30 ml batches of BG-11 medium pH 7.5 (Williams, 1988) under 3% [CO_2] at 30°C

under continuous white light of 50 µmol photons $m^{-2}s^{-1}$ (ML) with agitation. Mutant pre-cultures

were supplemented with the appropriate antibiotics. At the logarithmic growth phase, cells were

harvested and re-suspended in fresh BG-11 without antibiotics at OD_{750} of 0.1–0.2, as described in

the appropriate figure legends. Cells were then shifted to air [CO_2] / 30°C and illuminated

continuously with white light of either 50 or 220 µmol photons $m^{-2}s^{-1}$, as described in the figure

legends.

Gas exchange measurements

Exchange of $^{16}O_2$ (*m/z*=32), $^{18}O_2$ (*m/z*=36), CO_2 (*m/z*=44) was measured *in vivo* with membrane-inlet mass spectrometry (MIMS) as described in Mustila et al. (2016). Harvested cells were resuspended in fresh BG-11 pH 7.5 and adjusted to 10 µg Chlorophyll (Chl) *a* ml^{-1}, and kept for one hour in air $[CO_2]$ / 50 µmol photons m^{-2}s^{-1}. Prior to the measurements, cells were supplemented with $^{18}O_2$ at an equivalent concentration to $^{16}O_2$ and with 1.5 mM NaHCO$_3$. Cells were dark-adapted for 15 min, after which gas exchange was monitored over a 5 min illumination period of 500 µmol photons m^{-2}s^{-1} of white actinic light (AL). The gas exchange rates were calculated according to Beckmann et al., (2009).

Measurement of fluorescence and absorbance changes

Experimental cultures for all spectroscopic experiments were grown for four days in air $[CO_2]$ / 50 µmol photons m^{-2}s^{-1} at 30°C in BG-11 pH 7.5. Chl *a* fluorescence and P700-oxido-reduction (875-830 nm absorbance difference) were simultaneously recorded (Fig. 3) with the Dual-PAM 100 spectrophotometer (Walz, Effeltrich, Germany). Harvested cells were resuspended in fresh BG-11 pH 7.5 and adjusted to 10 µg Chl *a* ml^{-1}, and kept for one hour in air $[CO_2]$ / 50 µmol photons m^{-2}s^{-1} and dark-adapted for 10 min before being subjected to fluctuating light regime alternating between 1 min periods of 25 and 530 µmol photons m^{-2}s^{-1} of red AL. Saturating pulses (500 ms of 5000 µmol photons m^{-2}s^{-1}) were administered at 15 s intervals. Photosynthetic parameters were calculated as follows: Y(NA)=(P$_m$-P$_m$´)/P$_m$; Y(ND)=P/P$_m$; Y(I)=1-Y(ND)-Y(NA) (Klughammer and Schreiber, 2008); Y(II)=(F$_m$´-F)/F$_m$´ (Genty et al., 1989).

A Dual-Klas NIR 100 spectrophotometer (Walz) was used to measure absorbance difference changes at 780–800 nm, 820–870 nm, 840–965 nm and 870–965 nm, from which PC, P700, and Fd redox changes (Fig. 5) were deconvoluted based on differential model plots (DMPs) for PC, P700, and Fd (Klughammer and Schreiber, 2016; Schreiber, 2017; Setif et al., 2019). In WT *Synechocystis*, very fast re-oxidation of Fd upon illumination impedes measurement of model spectra. To circumvent this issue, we measured the DMPs from Δ*flv3 d1d2* cells where P700 oxidation was severely delayed in comparison to WT (Fig. 3B). The DMPs (Fig. S3) were measured using scripts provided with the instrument software. Harvested cells were adjusted to 10 µg Chl *a* ml^{-1} with fresh BG-11 pH 7.5, and kept for one hour in air $[CO_2]$ / 50 µmol photons m^{-2}s^{-1} and

dark-adapted for 30 min before illumination for 30 s with 503 µmol photons m^{-2}s^{-1} of red AL. It is

noteworthy that cytochrome c_6 and (F$_A$F$_B$) redox changes likely contribute to the PC and Fd signals,

respectively (Setif et al., 2019).

NADPH fluorescence changes between 420 and 580 nm, induced by excitation at 365 nm, were

measured simultaneously with Chl *a* fluorescence with the Dual-PAM 100 and its 9-AA/NADPH

accessory module (Walz) (Schreiber and Klughammer, 2009; Kauny and Setif, 2014). Experimental

samples were prepared as above and [Chl *a*] was adjusted to 5 µg ml^{-1}. Cells were dark-adapted for

10 min and subjected to the same fluctuating light regime as in Fig. 3, but without saturating pulses.

Electrochromic shift (ECS) was measured as absorbance difference between 500 and 480 nm (Viola

et al., 2019). Absorbance changes were measured from dark-adapted cell suspensions with a JTS-10

spectrophotometer (BioLogic, Seyssinet-Pariset, France) using 500 and 480 nm CWL 50mm 10nm

FWHM bandpass filters (Edmund Optics, Barrington, NJ, USA) and BG39 filters (Schott, Mainz,

Germany) protecting the light detectors from scattering effects. Absorbance changes induced by

measuring light only were subtracted from changes under ML + AL. ECS values were normalized

to ECS change induced by a single turnover flash provided by a XST-103 xenon lamp (Walz).

Experimental samples were prepared as above and [Chl *a*] was adjusted to 7.5 µg ml^{-1}, and

illuminated with green AL of 500 µmol photons m^{-2}s^{-1} for 1s or 60 s interspersed with 500 ms dark

intervals at 2, 7.5, 18, 30, and 45 s. *Pmf* was calculated as the extent of ECS decrease at the dark

intervals. Thylakoid conductivity (gH+) was calculated as the inverse of the time constant of a first-

order fit to ECS relaxation kinetics during a dark interval, and proton flux (vH+) as *pmf* x gH+

(Cruz. et al., 2005).

77K fluorescence emission spectra were measured with a QE Pro spectrophotometer (Ocean Optics,

Dunedin, FL, USA). Cells were harvested and [Chl *a*] was adjusted to 7.5 µg ml^{-1} with fresh BG-11

pH 7.5. Cells were frozen in liquid N$_2$ and excited at 440 nm. Raw spectra were normalized to PSII-

fluorescence peak at 685 nm.

Protein extraction and immunoblotting

Total protein extracts from cultures were isolated as previously described (Zhang et al., 2009),

separated by SDS-PAGE on 12 % gels with 6M urea, and blotted on PVDF membranes.

Membranes were probed with antibodies raised against D1, PsaB (Agrisera, Vännäs, Sweden, AS10

695), PetH (Kindly shared by H. Matthijs), OCP (kindly shared by D. Kirilovsky), SbtA (kindly

shared by T. Ogawa), NdhD3 (Eurogentec, Liége, Belgium), Flv2, Flv4, Flv3 (Antiprot, Puchheim,

Germany), sll0218 (MedProbe, Oslo, Norway). Horseradish peroxidase –conjugated secondary

antibody (GE Healthcare, Chicago, IL, USA) and Amersham ECL (GE Healthcare) were used for

detection.

Supplemental material

Supplemental Figure S1. Growth of $\Delta flv1$ $d3d4$ and $\Delta flv3$ $d3d4$ mutant strains.

Supplemental Figure S2. O_2 and CO_2 fluxes in M55 and in WT at pH 6.

Supplemental Figure S3. Near-infrared (NIR) differential model blots (DMPs) for deconvolution of

PC, P700, and Fd signals with the Klas-NIR 100 spectrometer.

ACKNOWLEDGEMENTS

We thank Prof. Teruo Ogawa for sharing the FDP and NDH-1 mutants and Dr. Henna Mustila for

construction of the ΔM55 $flv3$ strain. Dr. Duncan Fitzpatrick is thanked for expertise and assistance

with MIMS. We also thank Dr. Pierre Setif and Dr. Gert Schansker for their expert advice on

measuring the DMPs for the Dual-Klas NIR spectrophotometer, and Prof. Jens Appel for critical

reading of the manuscript. We declare no conflicts of interest.

FIGURE LEGENDS

Figure 1. Growth of WT and mutant strains under different experimental conditions. All

experimental cultures were started from pre-cultures adjusted to OD_{750}=0.1. **(A)** Cells were pre-

cultivated under 3% [CO_2] / ML (50 μmol photons $m^{-2}s^{-1}$), adjusted to OD_{750}=0.1 and grown in the

same conditions. **(B)** Cells were pre-cultivated under 3% [CO_2]/ML and shifted to air [CO_2]/ML.

(C) Cells were pre-cultivated under air [CO_2]/ML and shifted to air [CO_2]/HL 220 µmol photons m^{-2}s^{-1}). **(D)** Cells were pre-cultivated under 3% [CO_2]/ML and shifted to air [CO_2]/HL. The values

presented in A–D are means of three independent biological replicates ±SE. **(E)** Growth of WT and

mutant cells on agar plates containing BG-11. Cells grown under 3% [CO_2] / ML were harvested

and spotted on BG-11 agar plates in ten-fold dilution series starting at OD_{750}=1, incubation was

under air [CO_2] /ML and air [CO_2] /HL for 7 days. See also Supplemental Fig. S1.

Figure 2. O_2 and CO_2 exchange rates in WT and mutant strains. Cells were grown in air [CO_2] /

ML for 4 days, after which the cells were harvested and Chl *a* concentration adjusted to 10 µg ml^{-1}

with fresh BG-11. Cells were dark-adapted for 15 min, and gas exchange was monitored by

membrane inlet mass spectrometry (MIMS) over a 5 min illumination period with 500 µmol

photons m^{-2}s^{-1} of white actinic light. Prior to the measurements, samples were supplemented with

$^{18}O_2$ at an equivalent concentration to $^{16}O_2$ to distinguish O_2 uptake from O_2 evolution, and with 1.5

mM $NaHCO_3$. The dashed line in each panel indicates the compensation point of CO_2 fixation

(uptake rate equals respiratory rate). The experiment was repeated with three independent biological

replicates, of which representative measurements are shown.

Figure 3. Photosynthetic parameters in WT and mutants strains in fluctuating light intensities.

(A) Acceptor side limitation of PSI, Y(NA), **(B)** donor side limitation of PSI, Y(ND), as well as **(C)**

effective yield of PSI, Y(I) were calculated from absorbance difference changes at 875-830 nm

according to (Klughammer and Schreiber, 2008). Chl *a* fluorescence changes were measured

simultaneously with PSI redox changes and used to calculate **(D)** effective yield of PSII. Cells were

grown in air [CO_2] / ML for 4 days, after which the cells were harvested and Chl *a* concentration

adjusted to 10 µg ml^{-1} with fresh BG-11. Cells were dark-adapted for 10 min, and then illuminated

with red actinic light intensity alternating between 25 and 530 µmol photons m^{-2}s^{-1} in 1 min periods.

Saturating pulses (500 ms of 5000 µmol photons m^{-2}s^{-1}) were provided every 15 seconds. The values

shown are averages from 3–4 independent biological replicates ±SE.

Figure 4. NADPH and Chl *a* fluorescence under fluctuating light in WT and mutant strains.

WT **(A)**, Δ*flv3* **(B)**, Δ*d1d2* **(C)**, and Δ*flv3 d1d2* cells **(D)** were grown in air [CO_2] / ML for 4 days,

after which the cells were harvested and Chl *a* concentration adjusted to 5 µg ml^{-1} with fresh BG-

11. Cells were dark-adapted for 10 min, and then subjected to illumination with red actinic light

alternating between 25 and 530 µmol photons m^{-2}s^{-1} in 1 min periods. NADPH redox changes were

monitored with the Dual-PAM 100 spectrophotometer and its 9-AA/NADPH accessory module by

measuring fluorescence changes between 420 and 580 nm induced by excitation at 365 nm, as

described by (Schreiber and Klughammer, 2009) and (Kauny and Setif, 2014). Chl *a* fluorescence

was recorded simultaneously. **(E)** NADPH redox changes during illumination of dark-adapted cells

with 530 μmol photons $m^{-2}s^{-1}$. Other experimental details are as in A-D. **(F)** shows a magnification

of the first 2 s of illumination in (E). The values are normalized according to minimum and

maximum fluorescence changes at the onset and cessation of illumination, respectively. The

experiments were repeated with three independent biological replicates, of which representative

measurements are shown. r.u. = relative units.

Figure 5. Redox changes of PC, P700, and Fd, and buildup of *pmf* during dark-to-HL

transitions. WT **(A)**, Δ*flv3* **(B)**, Δ*d1d2* **(C)**, and Δ*flv3 d1d2* cells **(D)** were dark-adapted for 10 min,

after which absorbance difference changes at 780–800 nm, 820–870 nm, 840–965 nm and 870–965

nm were measured with the Klas-NIR 100 spectrometer during 30 s illumination at 503 μmol

photons $m^{-2}s^{-1}$ of red actinic light. The PC, P700, and Fd signals were deconvoluted from the

absorbance difference changes using differential model plots (DMPs) determined for this study

(Fig. S3). The upper panels in A–D show magnifications of the first 1.5 seconds of illumination

from the lower panels. The experiment was repeated with three independent biological replicates, of

which representative traces are shown. See also Supplemental Fig. S3. **(E)** Buildup of *pmf* during

transitions from dark to 500 μmol photons $m^{-2}s^{-1}$ of green actinic light. Absorbance difference

between 500-480 nm (ECS) was measured with a JTS-10 spectrometer, and light-induced *pmf* was

determined from the dark interval relaxation kinetics (DIRK) of the ECS signal. Means ±SEM from

3–4 independent experiments are shown. The small inset shows *pmf* values after 1s of illumination

for a clearer view. **(F)** Conductivity of the thylakoid membrane (gH+) after one and 30 s of

illumination as in (E). gH+ was determined as the inverse of the time constant of first-order post-

illumination decay kinetics of the ECS signal. **(G)** Thylakoid proton flux (vH+), calculated as *pmf* x

gH+. **(H)** Representative traces of 500-480 nm absorbance differences used to determine the values

in (E) and post-illumination ECS decay kinetics during a 500 ms dark interval after 30 s in light

(lower panels). First-order fits are shown in green. Cells, after growth in air [CO_2] / 50 μmol

photons $m^{-2}s^{-1}$ for 4 days, were harvested and Chl *a* concentration adjusted to 10 μg ml^{-1} with fresh

BG-11 for A–D and to 7.5 μg ml^{-1} for E–J.

Figure 6. Immunodetection of proteins related to photosynthesis and CCM and 77K

fluorescence emission spectra in WT and mutant strains. Pre-cultures were grown for 3 days at

3% [CO_2] / ML (50 μmol photons $m^{-2}s^{-1}$) and then pelleted, resuspended in fresh BG-11 at

OD_{750}=0.15 and **(A)** incubated for 24 hours in air [CO_2] / ML, **(B)** grown for 4 days in air [CO_2] /

ML, harvested, resuspended in BG-11 at OD_{750}=0.15 and then incubated for 24 hours in air [CO_2] /

HL (220 μmol photons $m^{-2}s^{-1}$), **(C)** incubated for 24 hours in air [CO_2] / HL. Total protein extracts

were separated by SDS-PAGE and probed with specific antibodies. The immunoblots shown are

representative of 2–3 independent biological replicates. **(D)** 77K fluorescence emission spectra from

intact WT and Δ*flv3 d1d2* cells grown for 4 days in air [CO_2] / ML. **(E)** After pre-growth at 3%

[CO_2] / ML cells were pelleted, resuspended in fresh BG-11 at OD_{750}=0.15 and incubated for 24

hours in air [CO_2] / HL. The cells (D and E) were pelleted, adjusted to [Chl *a*] of 7.5 μg ml^{-1}, and

excited with 440 nm light at 77K. The spectra were normalized to the PSII fluorescence peak at 685

nm.

Figure 7. Schematic models for the coordination of the activities of FDPs and NDH-1 in

preventing over-reduction of PSI. (A) In conditions where the Fd pool is higly reduced and the

PQ pool is not, NDH-1L transfers electrons from Fd to the PQ pool, coupled to O_2 reduction by

thylakoid-localised RTOs. NDH-1L pumps $2H^+/e^-$ into the lumen while FDPs and RTOs consume

H^+ when reducing O_2 to H_2O on the cytosolic side of thylakoids. This increases transthylakoid *pmf*,

which in turn drives ATP synthesis as well as inhibits PQH_2 oxidation at Cyt *b₆f*, thereby preventing

excessive electron supply to PSI. Carbonic anhydrase activity of NDH1-MS converts HCO_3^- to

CO_2, consuming H^+ in the cytosol, likely also coupled to translocation of H^+ to the lumen, further

increasing *pmf*. Increased availability of CO_2 and ATP/NADPH ratio enhances the electron sink

capacity of the CBB on the acceptor side of the PETC. **(B)** If the PQ pool is highly reduced, Fd pool

is oxidized, and *pmf* is high, NDH-1L could hypothetically function in reverse, transferring

electrons from PQH_2 to Fd^+, powered by release of $2H^+/e^-$ from the lumen. A strong Mehler-like

reaction catalyzed by Flv1/3 would be required to maintain a sufficient pool of oxidized Fd, with

Flv1/3 likely accepting electron directly from Fd^-. Most components depicted in (A) are omitted

from (B) to enhance clarity, not to suggest their absence. Extent of the colour fill for PQ/PQH_2 and

Fd depict level of reduction.

LITERATURE CITED

Alboresi A, Storti M, Cendron L, Morosinotto T (2019) Role and regulation of class-C flavodiiron

proteins in photosynthetic organisms. Biochem J 476: 2487-2498, doi/10.1042/BCJ20180648.

Allahverdiyeva Y, Ermakova M, Eisenhut M, Zhang P, Richaud P, Hagemann M, Cournac L, Aro

EM (2011) Interplay between Flavodiiron Proteins and Photorespiration in Synechocystis sp. PCC

6803. J Biol Chem 286: 24007-24014. doi/10.1074/jbc.M111.223289.

Allahverdiyeva Y, Mustila H, Ermakova M, Bersanini L, Richaud P, Ajlani G, Battchikova N,

Cournac L, Aro EM (2013) Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and

photosynthesis under fluctuating light. Proc Natl Acad Sci USA 110: 4111-4116,

doi/10.1073/pnas.1221194110.

Allahverdiyeva Y, Suorsa M, Tikkanen M, Aro EM (2015) Photoprotection of photosystems in

fluctuating light intensities. J Exp Bot 66: 2427-2436. doi/10.1093/jxb/eru463.

Allakhverdiev SI, Murata N (2004) Environmental stress inhibits the synthesis de novo of proteins

involved in the photodamage-repair cycle of Photosystem II in Synechocystis sp PCC 6803.

Biochim Biophys Acta Bioenerg 1657: 23-32. doi/10.1016/j.bbabio.2004.03.003.

Beckmann K, Messinger J, Badger MR, Wydrzynski T, Hillier W (2009) On-line mass

spectrometry: membrane inlet sampling. Photosynth Res 102: 511-522. doi/10.1007/s11120-009-

9474-7.

Bernat G, Appel J, Ogawa T, Rögner M (2011) Distinct Roles of Multiple NDH-1 Complexes in the

Cyanobacterial Electron Transport Network as Revealed by Kinetic Analysis of P700(+) Reduction

in Various ndh-Deficient Mutants of Synechocystis sp. Strain PCC6803. J Bacteriol 193: 292-295,

doi/10.1128/JB.00984-10.

Brändén G, Gennis RB, Brzezinski P (2006) Transmembrane proton translocation by cytochrome c

oxidase. Biochim Biophys Acta 1757: 1052-1063, doi/10.1016/j.bbabio.2006.05.020.

Brown KA, Guo Z, Tokmina-Lukaszewska M, Scott LW, Lubner CE, Smolinski S, Mulder DW,

Bothner B, King PW (2019) The oxygen reduction reaction catalyzed by Synechocystis sp. PCC

6803 flavodiiron proteins. Sustain Energy Fuels 3: 3191-3200, doi/10.1039/c9se00523d.

Bulychev AA, Cherkashin AA, Muronets EM, Elanskaya IV (2018) Photoinduction of electron

transport on the acceptor side of PSI in Synechocystis PCC 6803 mutant deficient in flavodiiron

proteins Flv1 and Flv3. Biochim Biophys Acta Bioenerg 1859, 1086-1095,

doi/10.1016/j.bbabio.2018.06.012.

Burnap RL, Hagemann M, Kaplan A (2015) Regulation of CO_2 concentrating mechanism in

cyanobacteria. Life (Basel) 5: 348-371.

Burnap RL, Nambudiri R, Holland S (2013) Regulation of the carbon-concentrating mechanism in

the cyanobacterium *Synechocystis* sp PCC6803 in response to changing light intensity and inorganic

carbon availability. Photosynth Res. 118, 115-124, doi/10.1007/s11120-013-9912-4.

Cassier-Chauvat C, Chauvat F (2014) Function and regulation of ferredoxins in the cyanobacterium,

Synechocystis PCC6803: recent advances. Life 4: 666–680.

Chaux F, Burlacot A, Mekhalfi M, Auroy P, Blangy S, Richaud P, Peltier G (2017). Flavodiiron

Proteins Promote Fast and Transient O-2 Photoreduction in Chlamydomonas. Plant Physiol 174:

1825-1836. doi/10.1104/pp.17.00421.

Cooley JW, Vermaas W (2001) Succinate dehydrogenase and other respiratory pathways in

thylakoid membranes of Synechocystis sp strain PCC 6803: Capacity comparisons and

physiological function. J Bacteriol 183: 4251-4258, doi/10.1128/JB.183.14.4251-4258.2001.

Cruz JA, Avenson TJ, Kanazawa A, Takizawa K, Edwards GE, Kramer DM (2005) Plasticity in

light reactions of photosynthesis for energy production and photoprotection. J Exp Bot 56: 395-406.

Daley SM, Kappell AD, Carrick MJ, Burnap RL (2012) Regulation of the cyanobacterial CO2-

concentrating mechanism involves internal sensing of NADP+ and α-ketoglutarate levels by

transcription factor CcmR. PLoS One 7: e41286. doi/10.1371/journal.pone.0041286.

Dann M, Leister D (2019) Evidence that cyanobacterial Sll1217 functions analogously to PGRL1 in

enhancing PGR5-dependent cyclic electron flow. Nat Commun 10: 5299. doi/10.1038/s41467-019-

13223-0.

Eisenhut M, Georg J, Klähn S, Sakurai I, Mustila H, Zhang P, Hess WR, Aro EM (2012) The

antisense RNA As1_flv4 in the Cyanobacterium Synechocystis sp. PCC 6803 prevents premature

expression of the flv4-2 operon upon shift in inorganic carbon supply. J Biol Chem 287: 33153-

33162.

Ermakova M, Huokko T, Richaud P, Bersanini L, Howe CJ, Lea-Smith DJ, Peltier G,

Allahverdiyeva Y (2016) Distinguishing the Roles of Thylakoid Respiratory Terminal Oxidases in

the Cyanobacterium Synechocystis sp PCC 6803. Plant Physiol 171: 1307-1319.

doi/10.1104/pp.16.00479.

Gao F, Zhao J, Chen L, Battchikova N, Ran Z, Aro EM, Ogawa T, Ma W (2016) The NDH-1L-PSI

Supercomplex Is Important for Efficient Cyclic Electron Transport in cyanobacteria. Plant Physiol

172: 1451-1464. doi/10.1104/pp.16.00585.

Genty B, Briantais JM, Baker NR (1989) The Relationship between the Quantum Yield of

Photosynthetic Electron-Transport and Quenching of Chlorophyll Fluorescence. Biochim Biophys

Acta 990: 87-92. doi/10.1016/S0304-4165(89)80016-9.

852 Gerotto C, Alboresi A, Meneghesso A, Jokel M, Suorsa M, Aro EM, Morosinotto T (2016)

853 Flavodiiron proteins act as safety valve for electrons in Physcomitrella patens. Proc Natl Acad Sci

854 USA 113: 12322-12327, doi/10.1073/pnas.1606685113.

855 Gómez R, Carrillo N, Morelli MP, Tula S, Shahinnia F, Hajirezaei MR, Lodeyro AF (2018) Faster

856 photosynthetic induction in tobacco by expressing cyanobacterial flavodiiron proteins in

857 chloroplasts. Photosynth Res 146: 129-136.

858 Hanke GT, Satomi Y, Shinmura K, Takao T, Hase T (2011) A screen for potential ferredoxin

859 electron transfer partners uncovers new, redox dependent interactions. Biochim Biophys Acta

860 Proteins Proteomics 1814: 366-374.

861 Han X, Sun N, Xu M, Mi H (2017) Co-ordination of NDH and Cup proteins in CO2 uptake in

862 cyanobacterium Synechocystis sp PCC 6803. J Exp Bot 68: 3869-3877. doi/10.1093/jxb/erx129.

863 He Z, Zheng F, Wu Y, Li Q, Lv J, Fu P, Mi H (2015) NDH-1L interacts with ferredoxin via the

864 subunit NdhS in Thermosynechococcus elongatus. Photosynth Res 126: 341-349,

865 doi/10.1007/s11120-015-0090-4.

866 Helman Y, Tchernov D, Reinhold L, Shibata M, Ogawa T, Schwarz R, Ohad I, Kaplan A (2003)

867 Genes encoding a-type flavoproteins are essential for photoreduction of O-2 in cyanobacteria. Curr

868 Biol 13: 230-235, doi/10.1016/S0960-9822(03)00046-0.

869 Holland SC, Kappell AD, and Burnap RL (2015) Redox changes accompanying inorganic carbon

870 limitation in Synechocystis sp. PCC 6803. Biochim Biophys Acta 1847: 355-363.

871 doi/10.1016/j.bbabio.2014.12.001.

Ilik P, Pavlovic A, Kouril R, Alboresi A, Morosinotto T, Allahverdiyeva Y, Aro EM, Yamamoto H,

Shikanai T (2017) Alternative electron transport mediated by flavodiiron proteins is operational in

organisms from cyanobacteria up to gymnosperms. New Phytol 214: 967-972.

doi/10.1111/nph.14536.

Jokel M, Johnson X, Peltier G, Aro EM, Allahverdiyeva Y (2018). Hunting the main player

enabling Chlamydomonas reinhardtii growth under fluctuating light. Plant J 94: 822-835,

doi/10.1111/tpj.13897.

Jokel M, Nagy V, Tóth SZ, Kosourov S, Allahverdiyeva Y (2019) Elimination of the flavodiiron

electron sink facilitates long-term H2 photoproduction in green algae. Biotechnol Biofuels 12: 280.

doi/10.1186/s13068-019-1618-1.

Kaplan A, Reinhold L (1999) CO2 concentrating mechanisms in photosynthetic microorganisms.

Annu Rev Plant Physiol Plant Mol Biol 50: 539-570. doi/10.1146/annurev.arplant.50.1.539.

Kauny J, Setif P (2014) NADPH fluorescence in the cyanobacterium Synechocystis sp PCC 6803:

A versatile probe for in vivo measurements of rates, yields and pools. Biochim Biophys Acta

Bioenerg 1837: 792-801. doi/10.1016/j.bbabio.2014.01.009.

Klughammer C. Schreiber U (2008) Saturation Pulse method for assessment of energy conversion

in PSI. PAM Application Notes 1: 11-14.

Klughammer C, Schreiber U (2016) Deconvolution of ferredoxin, plastocyanin, and P700

transmittance changes in intact leaves with a new type of kinetic LED array spectrophotometer.

Photosynth Res 128: 195-214.

Kramer DM, Avenson TJ, Edwards GE (2004) Dynamic flexibility in the light reactions of

photosynthesis governed by both electron and proton reactions. Trends Plant Sci 9: 349-357.

Laughlin TG, Bayne AN, Trempe JF, Savage DF, Davies KM (2019) Structure of the complex I-

like molecule NDH of oxygenic photosynthesis. Nature 566: 411–414.

Liran O, Shemesh E, Tchernov D (2018) Investigation into the CO_2 concentrating step rates within

the carbon concentrating mechanism of Synechocystis sp PCC6803 at various pH and light

intensities reveal novel mechanistic properties. Algal Res 33: 419-429.

doi/10.1016/j.algal.2018.06.020.

Liu L, Bryan SJ, Huang F, Yu J, Nixon PJ, Rich PR, Mullineaux CW (2012) Control of electron

transport routes through redox-regulated redistribution of respiratory complexes. Proc Natl Acad

Sci USA 109: 11431-11436. doi/10.1073/pnas.1120960109.

Mehler AH (1957) Studies on reactions of illuminated chloroplasts. I. Mechanism of the reduction

of oxygen and other Hill reagents. Arch Biochem Biophys 33: 65–77.

Ma W, Ogawa T (2015) Oxygenic Photosynthesis-specific Subunits of Cyanobacterial NADPH

Dehydrogenases. IUBMB Life 67: 3-8. doi/10.1002/iub.1341.

McCormick AJ, Bombelli P, Lea-Smith DJ, Bradley RW, Scott AM, Fisher AC, Smith AG, Howe

CJ (2013) Hydrogen production through oxygenic photosynthesis using the cyanobacterium

Synechocystis sp PCC 6803 in a bio-photoelectrolysis cell (BPE) system. Energy Environ Sci 6:

2682-2690. doi/10.1039/c3ee40491a.

Mullineaux CW (2014) Co-existence of photosynthetic and respiratory activities in cyanobacterial

thylakoid membranes. Biochim Biophys Acta Bioenerg 1837: 503-511.

doi/10.1016/j.bbabio.2013.11.017.

Mustila H, Paananen P, Battchikova N, Santana-Sanchez A, Muth-Pawlak D, Hagemann M, Aro

EM, Allahverdiyeva Y (2016) The Flavodiiron Protein Flv3 Functions as a Homo-Oligomer During

Stress Acclimation and is Distinct from the Flv1/Flv3 Hetero-Oligomer Specific to the O-2

Photoreduction Pathway. Plant Cell Physiol 57: 1468-1483. doi/10.1093/pcp/pcw047.

Nakamura Y, Kaneko T, Miyajima N, Tabata S (1999) Extension of CyanoBase. CyanoMutants:

repository of mutant information on *Synechocystis* sp strain PCC6803. Nucleic Acids Res 27: 66-

68. doi/10.1093/nar/27.1.66.

Nikkanen L, Toivola J, Trotta A, Guinea Diaz M, Tikkanen M, Aro EM, Rintamäki E (2018)

Regulation of cyclic electron flow by chloroplast NADPH-dependent thioredoxin system. Plant

Direct 2: 1-24.

Ogawa T (1991) A gene homologous to the subunit-2 gene of NADH dehydrogenase is essential to

inorganic carbon transport of Synechocystis PCC6803. Proc Natl Acad Sci USA 88: 4275-4279.

Ohkawa H, Price D, Badger M, Ogawa T (2000a) Mutation of *ndh* genes leads to inhibition of CO2

uptake rather than HCO3- uptake in *Synechocystis* sp. strain PCC 6803 J Bacteriol 182: 2591-2596.

Ohkawa H, Pakrasi HB, Ogawa T (2000b) Two types of functionally distinct NAD(P)H

dehydrogenases in *Synechocystis* sp strain PCC6803. J Biol Chem 275: 31630-31634.

doi/10.1074/jbc.M003706200.

Peltier G, Aro EM, Shikanai T (2016) NDH-1 and NDH-2 Plastoquinone Reductases in Oxygenic

Photosynthesis. Annu Rev Plant Biol 67: 55-80.

Santana-Sanchez A, Solymosi D, Mustila H, Bersanini L, Aro EM, Allahverdiyeva Y (2019)

Flavodiiron proteins 1-to-4 function in versatile combinations in O-2 photoreduction in

cyanobacteria. eLife 8: e45766. doi/10.7554/eLife.45766.

Saura P, Kaila VRI (2019) Molecular dynamics and structural models of the cyanobacterial NDH-1

complex. Biochim Biophys Acta Bioenerg 1860: 201-208. doi/10.1016/j.bbabio.2018.11.010.

Schreiber U, Klughammer C (2009) New NADPH/9-AA module for the DUAL-PAM-100:

Description, operation and examples of application. PAM Application Notes 2: 1-13.

Schreiber U (2017) Redox changes of ferredoxin, P700, and plastocyanin measured simultaneously

in intact leaves. Photosynth Res 134: 343-360.

Schuller JM, Birrell JA, Tanaka H, Konuma T, Wulflhorst H, Cox N, Schuller SK, Thiemann J,

Lubitz W, Setif P, Ikegamis T, Engel BD, Kurisu G, Nowaczyk MM (2019) Structural adaptations

of photosynthetic complex I enable ferredoxin-dependent electron transfer. Science 363: 257-260.

doi/10.1126/science.aau3613.

Setif P, Boussac A, Krieger-Liszkay A (2019) Near-infrared in vitro measurements of photosystem

I cofactors and electron-transfer partners with a recently developed spectrophotometer. Photosynth

Res 142: 307-319. doi/10.1007/s11120-019-00665-2.

Shibata M, Ohkawa H, Kaneko T, Fukuzawa H, Tabata S, Kaplan A, Ogawa T (2001) Distinct

constitutive and low-CO2-induced CO2 uptake systems in cyanobacteria: Genes involved and their

951 phylogenetic relationship with homologous genes in other organisms. Proc Natl Acad Sci USA 98:

952 11789-11794. doi/10.1073/pnas.191258298.

953 Shibata M, Katoh H, Sonoda M, Ohkawa H, Shimoyama M, Fukuzawa H, Kaplan A, Ogawa T

954 (2002) Genes essential to sodium-dependent bicarbonate transport in cyanobacteria - Function and

955 phylogenetic analysis. J Biol Chem 277: 18658-18664, doi/10.1074/jbc.M112468200.

956 Shimakawa G, Shaku K, Miyake C (2016) Oxidation of P700 in Photosystem I Is Essential for the

957 Growth of Cyanobacteria. Plant Physiol 172: 1443-1450. doi/10.1104/pp.16.01227.

958 Shimakawa G, Miyake C (2018a) Changing frequency of fluctuating light reveals the molecular

959 mechanism for P700 oxidation in plant leaves. Plant Direct 2: e00073.

960 Shimakawa G, Miyake C (2018b) Oxidation of P700 Ensures Robust Photosynthesis. Front Plant

961 Sci 9: 1617. doi/10.3389/fpls.2018.01617.

962 Strand DD, D'Andrea L, Bock R (2019) The plastid NAD(P)H dehydrogenase-like complex:

963 structure, function and evolutionary dynamics. Biochem J 476: 2743-2756.

964 doi/10.1042/BCJ20190365.

965 Strand DD, Fisher N, Kramer DM (2017) The higher plant plastid NAD(P)H dehydrogenase-like

966 complex (NDH) is a high efficiency proton pump that increases ATP production by cyclic electron

967 flow. J Biol Chem 292: 11850-11860.

968 Thiel K, Patrikainen P, Nagy C, Fitzpatrick D, Pope N, Aro EM, Kallio P (2019) Redirecting

969 photosynthetic electron flux in the cyanobacterium Synechocystis sp. PCC 6803 by the deletion of

970 flavodiiron protein Flv3. Microb Cell Fact 18: 189-2. doi/10.1186/s12934-019-1238-2

Tikkanen M, Mekala NR, Aro EM (2014) Photosystem II photoinhibition-repair cycle protects

Photosystem I from irreversible damage. Biochim Biophys Acta 1837: 210-215.

doi/10.1016/j.bbabio.2013.10.001.

Tiwari A, Mamedov F, Grieco M, Suorsa M, Jajoo A, Styring S, Tikkanen M, Aro EM (2016)

Photodamage of iron-sulphur clusters in photosystem I induces non-photochemical energy

dissipation. Nat Plants 2: 16035.

Ueda M, Kuniyoshi T, Yamamoto H, Sugimoto K, Ishizaki K, Kohchi T, Nishimura Y, Shikanai T

(2012) Composition and physiological function of the chloroplast NADH dehydrogenase-like

complex in Marchantia polymorpha. Plant J 72: 683-693, doi/10.1111/j.1365-313X.2012.05115.x.

Vicente JB, Gomes CM, Wasserfallen A, Teixeira M (2002) Module fusion in an A-type

flavoprotein from the cyanobacterium Synechocystis condenses a multiple-component pathway in a

single polypeptide chain. Biochem Biophys Res Commun 294: 82-87. doi/10.1016/S0006-

291X(02)00434-5.

Viola S, Bailleul B, Yu J, Nixon P, Selles J, Joliot P, Wollman F (2019) Probing the electric field

across thylakoid membranes in cyanobacteria. Proc Natl Acad Sci USA 116: 21900-21906.

doi/10.1073/pnas.1913099116.

Williams JGK (1988) Construction of Specific Mutations in Photosystem-Ii Photosynthetic

Reaction Center by Genetic-Engineering Methods in Synechocystis-6803. Meth Enzymol: 167.

766-778.

Xu M, Bernát G, Singh A, Mi H, Rögner M, Pakrasi HB, Ogawa T (2008) Properties of mutants of

Synechocystis sp. strain 6803 lacking inorganic carbon sequestration systems. Plant Cell Physiol

49: 1672-1677. doi/10.1093/pcp/pcn139.

Yamamoto H, Takahashi S, Badger MR, Shikanai T (2016) Artificial remodelling of alternative

electron flow by flavodiiron proteins in Arabidopsis. Nat Plants 2: 16012. doi/

10.1038/nplants.2016.12.

Yamori W, Sakata N, Suzuki Y, Shikanai T, Makino A (2011) Cyclic electron flow around

photosystem I via chloroplast NAD(P)H dehydrogenase (NDH) complex performs a significant

physiological role during photosynthesis and plant growth at low temperature in rice. Plant J 68:

966-976.

Yamori W, Shikanai T, Makino A (2015) Photosystem I cyclic electron flow via chloroplast NADH

dehydrogenase-like complex performs a physiological role for photosynthesis at low light. Sci Rep

5: 15593.

Yamori W, Shikanai T (2016) Physiological Functions of Cyclic Electron Transport Around

Photosystem I in Sustaining Photosynthesis and Plant Growth. Annu Rev Plant Biol 67: 81-106.

Yeremenko N, Jeanjean R, Prommeenate P, Krasikov V, Nixon PJ, Vermaas W, Havaux M,

Matthijs H (2005) Open reading frame ssr2016 is required for antimycin A-sensitive photosystem I-

driven cyclic electron flow in the cyanobacterium Synechocystis sp PCC 6803. Plant Cell Physiol

46: 1433-1436, doi/10.1093/pcp/pci147.

Zhang P, Battchikova N, Jansen T, Appel J, Ogawa T, Aro EM (2004) Expression and functional

roles of the two distinct NDH-1 complexes and the carbon acquisition complex

NdhD3/NdhF3/CupA/Sll1735 in Synechocystis sp PCC 6803. Plant Cell 16: 3326-3340.

doi/10.1105/tpc.104.026526.

Zhang P, Allahverdiyeva Y, Eisenhut M, Aro EM (2009) Flavodiiron Proteins in Oxygenic

Photosynthetic Organisms: Photoprotection of Photosystem II by Flv2 and Flv4 in Synechocystis sp

PCC 6803. PLoS One 4: e5331. doi/10.1371/journal.pone.0005331.

Zhao J, Gao F, Fan DY, Chow WS, Ma W (2017) NDH-1 Is Important for Photosystem I function

of *Synechocystis* sp. strain PCC 6803 under environmental stress conditions. Front Plant Sci 8:

2183. doi/10.3389/fpls.2017.02183.

1019

1020