Human antimicrobial peptide, LL-37, induces non-inheritable

reduced susceptibility to vancomycin in *Staphylococcus aureus*

Cathrine Friberg[1#], Jakob Haaber[1$], Martin Vestergaard[1], Anaëlle Fait[1], Veronique Perrot[2], Bruce Levin[2], Hanne

Ingmer[1]*

1) Department of Veterinary and Animal Sciences, University of Copenhagen, Stigbøjlen 4, DK-1870

Frederiksberg C, Denmark

Current address: Novo Nordisk, Hagedornsvej 1, DK-2820 Gentofte, Denmark

$ Current address: SNIPRbiome, Lerso Parkallé 44, DK-2100 Copenhagen, Denmark

2) Department of Biology, Emory University, Atlanta, GA, USA

Running title: LL-37 reduces vancomycin susceptibility

Keywords: *Staphylococcus aureus* USA300, VISA, antimicrobial peptides, LL-37, vancomycin

*Corresponding author (hi@sund.ku.dk)

ABSTRACT

Antimicrobial peptides (AMPs) are central components of the innate immune system providing protection

against pathogens. Yet, serum and tissue concentrations vary between individuals and with disease conditions.

We demonstrate that the human AMP LL-37 lowers the susceptibility to vancomycin in the community-

associated methicillin-resistant *S. aureus* (CA-MRSA) strain FPR3757 (USA300). Vancomycin is used to treat

serious MRSA infections, but treatment failures occur despite MRSA strains being tested susceptible according

to standard susceptibility methods. Exposure to physiologically relevant concentrations of LL-37 increased the

minimum inhibitory concentration (MIC) of *S. aureus* towards vancomycin by 75%, and resulted in shortened

lag-phase and increased colony formation at sub-inhibitory concentrations of vancomycin. Computer

simulations using a mathematical antibiotic treatment model indicated that a small increase in MIC might

decrease the efficacy of vancomycin in clearing a *S. aureus* infection. This prediction was supported in a

Galleria mellonella infection model, where exposure of *S. aureus* to LL-37 abolished the antimicrobial effect of

vancomycin. Thus, physiological relevant concentrations of LL-37 reduce susceptibility to vancomycin,

indicating that tissue and host specific variations in LL-37 concentrations may influence vancomycin

susceptibility *in vivo*.

Introduction

Bacterial resistance to antibiotics is generally associated with genetic changes, such as point mutations in the core genome or acquisition of resistance genes present on mobile genetic elements[1] or intrinsic resistance to antimicrobials, such as in-permeability of the outer membrane of Gram-negative bacteria or the active efflux of antimicrobials provided by chromosomally encoded efflux pumps[2]. Previously, we showed that exposure to the synthetic antimicrobial peptide (AMP) colistin reduces susceptibility to vancomycin in the clinically important, methicillin resistant *Staphylococcus aureus* (MRSA) strain USA300 via induction of the GraRS cell wall regulon[3]. *S. aureus* is a human pathogen that causes a variety of diseases and particularly the MRSA clones are becoming widely distributed both in the hospital environment and in the community[4]. MRSA infections are often treated with the glycopeptide antibiotic vancomycin. The CLSI susceptibility breakpoint for vancomycin is 2 µg/ml[5]. However, in the last decade vancomycin-intermediate *S. aureus* strains (VISA) have emerged with reduced vancomycin susceptibility (MIC = 4-8 µg/ml)[6]. These strains are characterized by chromosomal mutations that reduce negative cell wall charge[7], increase cell wall thickness[8] or reduce Triton-X mediated autolysis[9]. Even for non-VISA strains vancomycin treatment failures have been linked to strains displaying MIC in the higher end of the susceptibility range (≥1.5 µg/ml)[5,10-13]. Our previous work revealed that upon colistin exposure, *S. aureus* cells transiently and reversibly induce cellular changes that result in a phenotypic VISA-like appearance[3]. This reduced vancomycin susceptibility state will remain undetectable by routine clinical testing and thus the MIC of vancomycin upon colistin exposure *in vivo* may be higher than expected from such testing.

Colistin is a cationic AMP used in treatment of Gram-negative infections, however AMPs are also major components of the innate immune system and a critical defence against a large number of pathogens[14]. Human AMPs are small, often cationic peptides with broad-spectrum antimicrobial activity caused by membrane-damaging pore-formation[14]. An example is the cathelicidin, LL-37[15], which is expressed in a variety of immune cells, including polymorphonuclear leukocytes (PMN), the first line of defence against invading bacterial

56   pathogens. In the present study, we demonstrate that the human AMP LL-37 transiently reduces susceptibility

57   to vancomycin.

Materials and methods

Bacterial strains and growth conditions

S. aureus USA300 FPR3757[16] was obtained from the American Type Culture Collection (ATCC). *S. aureus* strains were routinely grown in Mueller-Hinton (MH) medium (Sigma) at 37°C in Erlenmeyer flasks with aeration.

Minimum inhibitory concentration

To assess vancomycin susceptibility of *S. aureus* exposed to LL-37 (Sigma), *S. aureus* FPR3757 was diluted to 10^5 CFU/ml in Mueller-Hinton (MH) media and incubated for 1 h (200 rpm, 37 °C). Pre-treatment with either 10 or 20 µg/ml LL-37 for 1 h was followed by supplementation of vancomycin (Sigma) and the minimum inhibitory concentration (MIC) was determined in a 96-well microtiter plates (200 rpm, 37 °C, 24 h using technical duplicates and biological triplicates).

Vancomycin susceptibility testing by plating

Exponential phase *S. aureus* FPR3757 cultures were adjusted to $OD_{600} = 0.25$ in pre-heated MH broth and was then treated with sub-inhibitory concentrations of LL-37 (0, 5, 10, 20 and 40 µg/ml) for 1 h. The cultures were adjusted to $OD_{600} = 0.1$ before spotting 10 µl aliquots of serial four-fold dilutions on freshly prepared MH agar plates containing increasing concentrations of vancomycin (0, 1, 1.25 and 1.5 µg/ml).

In one experiment, cultures of *S. aureus* FPR3757 were grown with 10 µg/ml LL-37 for 60 min and LL-37 was removed from the medium by spinning (5,000g, 5 min, room temperature) and re-suspended in pre-heated MH broth in a subset of the culture, while exposure to LL-37 was maintained in the other subset. After 0, 60 and 90 min, the cultures where plated on agar plates containing vancomycin (1.5 µg/ml).

The agar plates were inspected for growth following 24 h incubation.

Growth experiments

Overnight cultures were diluted to approx. 10^5 CFU/ml in fresh MH broth, and incubated for 1 h at 37°C and

200 rpm. The culture was then split in appropriate aliquots and treated with LL-37 or left untreated for 1 h at

37°C and 200 rpm. The untreated and pre-treated cultures were then split in appropriate aliquots and treated

with vancomycin (Sigma) or left untreated. These were then further aliquoted into the wells of a 100-well

honeycomb plate and incubated in a Bioscreen C Analyzer (Oy Growth Curves Ab Ltd, Helsinki, Finland) at 37°C

and medium continuous shaking.

Determination of Triton X-100 induced autolysis

Strains were grown in MH broth at 37°C, 180 rpm, to mid-exponential phase. The cultures were diluted to an

OD_{600} of 0.15 in warm MH, grown to an OD_{600} of 0.25, and 10 µg/ml LL-37 was added to one subset of cultures

while another subset of cultures served as the unexposed control. After 1 h, the cultured cells were washed

twice in ice-cold sterile distilled water and resuspended in the same volume of 0.05 M Tris-HCl (pH 7.2)

containing 0.05% Triton X-100. Cells were incubated at 30°C, and the OD_{600} was measured every 30 min. Data

are expressed as the percent loss of OD_{600} at the indicated times compared to findings at time zero. Each data

point represents the mean and standard deviation from three independent experiments.

A model of antibiotic treatment

Using previous models[17], the following model was derived to test the effect of a higher MIC on clearance when

treated with the same antibiotic concentration. In a continuous flow at rate *w* a bacterial population (P) is

limited by the concentration of a resource (R; µg/ml). An antibiotic of concentration (A; µg/ml) is dosed every 8

hours and the concentration declines exponentially at rate *d* per day, due to flow from the site of infection. The

population replicates at maximum rates V_{MAX} and the rates of growth are proportional to the concentration of

the resource and antibiotic. For the antibiotic-mediated killing, we assume hill functions so that at an antibiotic

concentration of A and resource concentration of R, the net growth rate of the P population is,

$$\psi_P(A,R) = \left(V_{MAX} - \left(\frac{(V_{MAX} - V_{MIN})\left(\frac{A}{MIC}\right)^k}{\left(\frac{A}{MIC}\right)^k - \left(\frac{V_{MIN}}{V_{MAX}}\right)} \right) \right) \frac{R}{(R+k)}$$

where V_{MIN} is the minimum growth (maximum kill) rates and MIC the corresponding minimum inhibitory

concentrations of the antibiotic. The parameter κ controls the shape of this function[17]. To account for the

resource limitation, we use the function R/(R+k) where k, the Monod constant, is the concentration of the

limiting resource where in the absence of the antibiotic the rate of is half its maximum value. The bacteria take

up resource at rates proportional to their maximum growth rates and a conversion efficiency parameter, e

(μg/cell). The population is maintained in continuous culture (chemostat-like) habitat into which resources

from a reservoir at a concentration of C μg/ml enter at a constant rate, w per hour, which is the same rate at

which excess resources, bacteria and antibiotics are removed. With these definitions and assumption, the rate

of change in the densities of the different populations of bacteria and concentration of antibiotics

$$\frac{dR}{dt} = w(C - R) - e\frac{R}{R+k}V_{MAX}P$$

$$\frac{dP}{dt} = \psi_P(A,R)P - wP$$

$$\frac{dA}{dt} = -A(w+d)$$

where d is the rate of decay in the concentration of the antibiotic.

The simulation was programmed in Berkeley Madonna[TM] using Euler.

Galleria mellonella infection model

Twenty healthy 5th instar larvae each weighing 250 mg were injected with 2.5 x 10^6 CFU of *S. aureus* FPR3757

followed by 10 µg/ml LL-37 or 0.9 % NaCl for controls. One hour post infection, the larvae were treated with 15

mg/kg bodyweight of vancomycin, corresponding to a relevant clinical dose. The LL-37 pre-treatment and

antimicrobial therapy was repeated every 24 h for a total of 72 h and survival was monitored. Controls for

toxicity of LL-37 combined with the antimicrobials as well as for non-treated *S. aureus* infection were included

with all larvae receiving an equal number of injections of either active compound or 0.9 % NaCl.

Statistics

The data were analyzed in GraphPad Prism 7 (GraphPad Software Inc.) using one-way analysis of variance

(ANOVA) with a *post hoc* analysis of Dunnett's multiple comparison tests, where *P < 0.05* (*), *P < 0.01* (**) and

P < 0.001 (***). The survival data was analyzed by using the Kaplan-Meier method and statistical difference

determined, using log-rank test.

Results

Pre-exposure to LL-37 reduces S. aureus susceptibility to vancomycin

We have previously shown that pre-exposure of *S. aureus* to the antimicrobial peptide, colistin, reduces

vancomycin susceptibility[3]. To address if exposure to the human antimicrobial peptide LL-37 also affects

vancomycin susceptibility we initially measured the minimum inhibitory concentration (MIC) towards

vancomycin in the presence or absence of LL-37. While untreated *S. aureus* FPR3757 displayed vancomycin MIC

of 0.8 µg/ml, pre-treatment with LL-37 (MIC_{LL-37} = >128 µg/ml) for one hour at sub-inhibitory concentrations, 10

or 20 µg/ml, displayed vancomycin MIC of 1.2 and 1.4 µg/ml, respectively. To further examine the effect of LL-

37 on vancomycin susceptibility, we spotted serial dilutions of cultures pre-treated with various sub-inhibitory

concentrations of LL-37 on agar plates containing increasing concentrations of vancomycin. Cultures exposed

to 40 µg/ml LL-37 were able to form colonies in the presence of 1.5 µg/ml vancomycin in contrast to untreated

cells, where no growth was observed (Figure 1).

When observing the growth dynamics of treated and untreated cells in liquid cultures, pre-treatment with LL-

37 notably reduced the vancomycin induced lag-phase measured as the time from inoculation until density

reached OD_{600} = 0.1 (Figure 2a and b). In the presence of vancomycin (at 0.75 µg/ml) the lag-phase was 24.5

(±1.9) h for untreated cells, whereas the lag-phase was 8.1 (±2.5) h following pre-exposure to 20 µg/ml LL-37

(p<0.001). Shortening of the lag phase was proportional to the concentration of LL-37 used for pre-treatment

(Figure 2a and b). In the absence of vancomycin, LL-37 treatment did not affect growth kinetics (Figure 2a).

When determining maximum growth rate, no effect of LL-37 was observed in the vancomycin-treated or

control cultures (Figure S1), indicating that shortening of lag-phase rather than growth rate is important for the

LL-37 induced reduction in vancomycin susceptibility.

LL-37 induces reversible vancomycin tolerance

To examine if continuous LL-37 exposure is needed for the reduced vancomycin susceptibility we monitored

the LL-37 exposure time required before being able to observe reduced susceptibility to vancomycin and how

long the cells retained the reduced susceptibility after removal of LL-37. We found that the phenotype was

observed already after 10 minutes of exposure to LL-37 and that it lasted approximately 60 minutes after LL-37

was removed from the growth medium (Figure 3). These results demonstrate that the reduced vancomycin

susceptibility induced by LL-37 exposure is transient and non-inherited.

LL-37 reduces autolysis

Reduced susceptibility to vancomycin in genetically stable VISA strains are characterized phenotypically by a

decrease in Triton X-100 induced autolysis[9]. We examined the effect of LL-37 exposure (10 µg/ml) on autolysis

as described previously[3,7]. Compared to an un-treated control, the pre-treated culture showed less Triton X-100

induced autolysis (Figure 4).

In silico model

Clinical isolates of vancomycin susceptible MRSA strains with MIC in the higher end of the susceptibility range

(≥1.5 µg/ml) are linked to vancomycin treatment failure[5]. Such treatment failures may be related to

unidentified host factors, which are causing an increase in *S. aureus* vancomycin MIC[18]. Since we observed an

increase in vancomycin MIC after exposure to relevant concentrations of human LL-37 in the laboratory, we

explored the potential clinical implications *in silico* using a mathematical antibiotic treatment model to

simulate situations of constant concentrations of antibiotic and variable MIC. As illustrated by this model, if the

concentration of antibiotic is significantly higher (≈5x) than the MIC, small increments in MIC do not markedly

affect the rate of clearance (Figure 5). However, the lower the antibiotic concentration is at the site of

infection, the more impact the increased MIC will have on the rate of clearance. Maximum allowable

vancomycin concentration during treatment is limited by the nephrotoxic properties of the drug[19] and thus

clinical data[5] and our computer simulations indicate that the relatively small increase in MIC observed when *S.*

aureus is exposed to clinically relevant LL-37 concentrations may have a negative impact on the efficacy of

vancomycin in clearing an infection.

LL-37 abolishes antimicrobial effect of vancomycin in Galleria mellonella

The observed *in vitro* reduction in vancomycin susceptibility after LL-37 exposure combined with the

predictions of treatment failure associated with modest increments in MIC from our computer model

prompted us to test the efficacy of vancomycin in treating *Galleria mellonella* wax moth larvae[20] infected with

S. aureus FPR3757 in the presence or absence of LL-37. In the absence of LL-37, vancomycin reduced the

mortality inflicted by the MRSA strain. However, in the presence of LL-37, the antimicrobial potential of

vancomycin was completely abolished (P<0.001), rendering the larvae as susceptible to *S. aureus* killing as the

non-treated *S. aureus* infection (Figure 6).

Discussion

Antimicrobial chemotherapy is currently based on standardized *in vitro* antimicrobial susceptibility testing, however the assay can face limitations in predicting *in vivo* treatment efficacy[21]. The lack of predictive value is partly attributed to the fact that the growth media in which the susceptibility tests are conducted, do not sufficiently represent the *in vivo* environment. Several factors, such as nutrient availability, oxygen level, pH and presence of innate immune components, have been identified to affect antibiotic susceptibility in both Gram-positive and –negative species[22-25]. Here we show that the presence of the antimicrobial peptide LL-37 of the innate immune response antagonizes the activity of vancomycin against *S. aureus in vitro*. This effect is transient and is also evident in the *G. mellonella* infection model, where the presence of LL-37 with vancomycin significantly reduces survival of *G. mellonella* larvae compared with vancomycin treatment alone. Our data suggest that the MIC values reported from *in vitro* assays may underestimate the vancomycin MIC *in vivo* in the presence of LL-37.

Vancomycin is one of the most important antibiotics for treatment of MRSA infections[26]. The CLSI susceptibility breakpoint for vancomycin is 2 µg/ml[5]. Treatment failures reported for infections with MRSA strains being susceptible to vancomycin has led to discussions whether the MIC breakpoint at 2 µg/ml should be further decreased[5]. A meta-analysis associated high-MIC (≥ 1.5 µg/ml) MRSA isolates with increased risk of mortality compared with low-MIC (< 1.5 µg/ml) MRSA isolates[5], thus indicating that even minor increases in vancomycin MIC can negatively affect treatment outcome and survival of patients with MRSA infections.

The impact of LL-37 on vancomycin susceptibility under *in vivo* conditions may vary from person to person. Plasma levels of LL-37 in healthy individuals have been measured to be approximately 1 µg/ml, but with variations between individuals from 0.5 to 3 µg/ml[27]. In areas of bacterial infection the local concentration will reach even higher levels due to de-granulation of PMNs, e.g. in bronchoalveolar lavage fluid, where LL-37 concentrations from 5 to 25 µg/ml have been measured in infected individuals[28,29]. Furthermore, LL-37 levels

are also influenced by disease conditions, where for example psoriasis patients have increased LL-37 levels[30],

while atopic dermatitis patients have reduced levels[31]. Thus, it is conceivable that person-to-person variation in

LL-37 levels could influence susceptibility to vancomycin pointing to the importance of host factors in

antimicrobial chemotherapy.

Collectively, our results indicate that a better understanding of the host's role in the development of reduced

vancomycin susceptibility could be important to design treatment regimens and improve clinical outcome.

Acknowledgement

This research was funded by grants from the U.S. National Institutes of General Medical Science GM098175-17 to B.R. L.; from the Danish Council for Independent Research, Technology and Production to J.H. (12-126289) and H.I. (7017-00079B) and the European Union Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No. 765147 to A. F. and H.I.

The funding sources had no influence on study design, data analysis or decision to submit the article for publication.

C.F., J.H., B.L. and H.I. conceived and designed the study. Experiments were performed by C.F., A.F. and J.H. Analysis of data and drafting of the manuscript was performed by C.F., J.H., M.V., V.P., B.L. and H.I. All authors read and approved the final manuscript.

Competing interests

None to declare.

Ethical approval

Not required.

References

References

1   Blair, J. M., Webber, M. A., Baylay, A. J., Ogbolu, D. O. & Piddock, L. J. Molecular mechanisms of
antibiotic resistance. *Nature Reviews Microbiology* **13**, 42-51 (2015).
2   Cox, G. & Wright, G. D. Intrinsic antibiotic resistance: mechanisms, origins, challenges and solutions.
International Journal of Medical Microbiology **303**, 287-292 (2013).
3   Haaber, J. *et al.* Reversible antibiotic tolerance induced in *Staphylococcus aureus* by concurrent drug
exposure. *Mbio* **6**, e02268-02214 (2015).
4   Chambers, H. F. & DeLeo, F. R. Waves of resistance: *Staphylococcus aureus* in the antibiotic era. *Nature*
Reviews Microbiology **7**, 629-641 (2009).
5   Van Hal, S., Lodise, T. P. & Paterson, D. L. The clinical significance of vancomycin minimum inhibitory
concentration in *Staphylococcus aureus* infections: a systematic review and meta-analysis. *Clinical*
Infectious Diseases **54**, 755-771 (2012).
6   Howden, B. P., Davies, J. K., Johnson, P. D., Stinear, T. P. & Grayson, M. L. Reduced vancomycin
susceptibility in *Staphylococcus aureus*, including vancomycin-intermediate and heterogeneous
vancomycin-intermediate strains: resistance mechanisms, laboratory detection, and clinical
implications. *Clinical microbiology reviews* **23**, 99-139 (2010).
7   Meehl, M., Herbert, S., Götz, F. & Cheung, A. Interaction of the GraRS two-component system with the
VraFG ABC transporter to support vancomycin-intermediate resistance in *Staphylococcus aureus*.
Antimicrobial agents and chemotherapy **51**, 2679-2689 (2007).
8   Sieradzki, K. & Tomasz, A. Alterations of cell wall structure and metabolism accompany reduced
susceptibility to vancomycin in an isogenic series of clinical isolates of *Staphylococcus aureus*. *Journal*
of bacteriology **185**, 7103-7110 (2003).
9   Koehl, J. L. *et al.* Cell wall composition and decreased autolytic activity and lysostaphin susceptibility of
glycopeptide-intermediate *Staphylococcus aureus*. *Antimicrobial agents and chemotherapy* **48**, 3749-
3757 (2004).
10   Howden, B. P. *et al.* Treatment outcomes for serious infections caused by methicillin-resistant
Staphylococcus aureus with reduced vancomycin susceptibility. *Clinical Infectious Diseases* **38**, 521-528
(2004).
11   Lodise, T. *et al.* Relationship between vancomycin MIC and failure among patients with methicillin-
resistant *Staphylococcus aureus* bacteremia treated with vancomycin. *Antimicrobial agents and*
chemotherapy **52**, 3315-3320 (2008).
12   Dombrowski, J. C. & Winston, L. G. Clinical failures of appropriately-treated methicillin-resistant
Staphylococcus aureus infections. *Journal of Infection* **57**, 110-115 (2008).
13   Moore, M. R., Perdreau-Remington, F. & Chambers, H. F. Vancomycin treatment failure associated with
heterogeneous vancomycin-intermediate *Staphylococcus aureus* in a patient with endocarditis and in
the rabbit model of endocarditis. *Antimicrobial agents and chemotherapy* **47**, 1262-1266 (2003).
14   De Smet, K. & Contreras, R. Human antimicrobial peptides: defensins, cathelicidins and histatins.
Biotechnology letters **27**, 1337-1347 (2005).
15   Gudmundsson, G. H. *et al.* The human gene *FALL39* and processing of the cathelin precursor to the
antibacterial peptide LL-37 in granulocytes. *European Journal of Biochemistry* **238**, 325-332 (1996).
16   Diep, B. A. *et al.* Complete genome sequence of USA300, an epidemic clone of community-acquired
meticillin-resistant *Staphylococcus aureus*. *The Lancet* **367**, 731-739 (2006).
17   Regoes, R. R. *et al.* Pharmacodynamic functions: a multiparameter approach to the design of antibiotic
treatment regimens. *Antimicrobial agents and chemotherapy* **48**, 3670-3676 (2004).
18   Holmes, N. E. *et al.* Antibiotic choice may not explain poorer outcomes in patients with *Staphylococcus*
aureus bacteremia and high vancomycin minimum inhibitory concentrations. *Journal of Infectious*
Diseases **204**, 340-347 (2011).

19   Horey, A., Mergenhagen, K. A. & Mattappallil, A. The Relationship of Nephrotoxicity to Vancomycin
Trough Serum Concentrations in a Veteran's Population: A Retrospective Analysis (November).
Ann.Pharmacother. (2012).
20   Ramarao, N., Nielsen-Leroux, C. & Lereclus, D. The insect *Galleria mellonella* as a powerful infection
model to investigate bacterial pathogenesis. *Journal of visualized experiments: JoVE* (2012).
21   Doern, G. V. & Brecher, S. M. The clinical predictive value (or lack thereof) of the results of *in vitro*
antimicrobial susceptibility tests. *Journal of Clinical Microbiology* **49**, S11-S14 (2011).
22   Ersoy, S. C. *et al.* Correcting a fundamental flaw in the paradigm for antimicrobial susceptibility testing.
EBioMedicine **20**, 173-181 (2017).
23   Hughes, D. & Andersson, D. I. Environmental and genetic modulation of the phenotypic expression of
antibiotic resistance. *FEMS microbiology reviews* **41**, 374-391 (2017).
24   Kubicek-Sutherland, J. Z. *et al.* Host-dependent induction of transient antibiotic resistance: a prelude to
treatment failure. *EBioMedicine* **2**, 1169-1178 (2015).
25   Sakoulas, G., Kumaraswamy, M., Kousha, A. & Nizet, V. Interaction of antibiotics with innate host
defense factors against *Salmonella enterica* Serotype Newport. *mSphere* **2**, e00410-00417 (2017).
26   Liu, C. *et al.* Clinical practice guidelines by the Infectious Diseases Society of America for the treatment
of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clinical infectious*
diseases, e18–e55 (2011).
27   Sørensen, O., Cowland, J. B., Askaa, J. & Borregaard, N. An ELISA for hCAP-18, the cathelicidin present
in human neutrophils and plasma. *Journal of immunological methods* **206**, 53-59 (1997).
28   Chen, C. I.-U., Schaller-Bals, S., Paul, K. P., Wahn, U. & Bals, R. β-defensins and LL-37 in bronchoalveolar
lavage fluid of patients with cystic fibrosis. *Journal of Cystic Fibrosis* **3**, 45-50 (2004).
29   Schaller-Bals, S., Schulze, A. & Bals, R. Increased levels of antimicrobial peptides in tracheal aspirates of
newborn infants during infection. *American Journal of Respiratory and Critical Care Medicine* **165**, 992-
995 (2002).
30   Batycka-Baran, A., Maj, J., Wolf, R. & Szepietowski, J. The new insight into the role of antimicrobial
proteins-alarmins in the immunopathogenesis of psoriasis. *Journal of immunology research* **2014**
(2014).
31   Ong, P. Y. *et al.* Endogenous antimicrobial peptides and skin infections in atopic dermatitis. *New*
England Journal of Medicine **347**, 1151-1160 (2002).

Figure legends

Figure 1. LL-37 influences vancomycin susceptibility of *S. aureus*.

Exponential phase *S. aureus* FPR3757 cultures were pretreated with 0, 5, 10, 20 and 40 µg/ml LL-37 for 1 h

before spotting serial four-fold dilutions on MH agar containing increasing concentrations of vancomycin.

Figure 2. LL-37 influences vancomycin susceptibility of *S. aureus* in liquid culture.

(a) The growth of *S. aureus* FPR3757 with (black lines) or without (grey lines) exposure to 0.75 µg/ml

vancomycin after treatment for 1 h with 0, 5, 10 or 20 µg/ml LL-37. The curves are an average of five replicates.

(b) The lag-phase of the same cultures depicted in (a) (n=5). Error bars display 95% confidence intervals.

**P<0.01; *P<0.05.

Figure 3. Transient induction of reduced vancomycin susceptibility by LL-37.

(a) Cultures of *S. aureus* FPR3757 were grown without (left) or with (right) 10 µg/ml LL-37 for 10 minutes and

spotted on vancomycin plates (1.5 µg/ml). (b) Cultures of *S. aureus* FPR3757 were grown with 10 µg/ml LL-37

for 60 min and LL-37 was removed from the medium in a subset of the culture (left), while exposure to LL-37

was maintained in the other subset (right). After 0, 60 and 90 min, the cultures where plated on agar plates

containing vancomycin (1.5 µg/ml).

Figure 4. LL-37 exposure reduces Triton X-100 mediated autolysis.

LL-37 induced autolysis measured as reduction of bacterial density over time in a 0.05% Triton X-100 solution

containing treated (10 µg/ml LL-37) or untreated cultures. Error bars display 95% confidence intervals, (n=3).

Figure 5. Computer simulation of the effects of increased vancomycin MIC on the clearance of an infection

treated with the same antibiotic concentration.

The full, dashed or double lines represent populations with MICs of 0.8, 1.2 or 1.4 µg/ml, respectively. The

black, green and orange lines represent exposure to 10, 5 or 2.5 µg/ml antibiotic, dosed every 8 hours.

Parameter values: $V_{MAX}=1$, $V_{MIN}=-3$, k=1, $e=5 \times 10^{-7}$, w=0.5 and C=1000.

Figure 6. Influence of LL-37 on vancomycin treatment of *S. aureus* infecting *G. mellonella*.

Survival of *G. mellonella* wax moth larvae infected with *S. aureus* FPR3757 and with or without LL-37 (10 µg/ml)

exposure followed by treatment with vancomycin (15 mg/kg bodyweight). Controls for non-treated *S. aureus*

infection and non-infected larvae exposed to combinations of the drugs were included. All larvae received

same number of injections. n=20 larvae for each condition.

Figures

Figure 1

Figure 2

(a)

(b)

Figure 3

(a)

(b)

Figure 4

Figure 5

Figure 6

Supplementary figure

Figure S1. LL-37 effect on maximum growth rate.

From the growth experiments as described in Materials and Methods, the maximum growth rates were
determined for *S. aureus* cultures pre-exposed to LL-37 (0 to 20 µg/ml) for 1 h before being exposed to
0 or 0.75 µg/ml vancomycin. Error bars represent s.d. of 3 biological replicates each containing 3
technical replicates.