1

Earliest experience of a relatively rare sound but not a frequent sound causes long term changes in the adult auditory cortex

Authors: Muneshwar Mehra[1], Adarsh Mukesh[1] and Sharba Bandyopadhyay[1,2*]

[1] Advanced Technology Development Centre, IIT Kharagpur, [2] Department of Electronics and Electrical Communication Engineering, IIT Kharagpur

[*]Corresponding author

Pages: 33 (+ 24 Supplementary Material), Figures: 7 (+ 6 Supplementary), Tables: (4 Supplementary), Videos: (2 Supplementary)

Author contributions: MM performed all experiments, AM assisted. MM, AM and SB analyzed data. AM did modeling and theoretical derivation. SB supervised the work. SB, MM and AM wrote the manuscript.

Acknowledgements: MM thanks CSIR for Fellowship, AM thanks MHRD for PMRF, SB thanks India Alliance, IIT Kharagpur, MHRD and SRIC Cell, IIT KGP for funding. This work was supported by the DBT/Wellcome Trust India Alliance Fellowship/Grant IA/I/11/2500270 awarded to SB.

Abstract

Sensory experience during a critical period alters sensory cortical responses and organization. We find that the earliest sound driven activity in the mouse auditory cortex (ACX) starts before ear-canal opening (ECO). Effects of auditory experience before ECO on ACX development are unknown. We find that mouse ACX subplate neurons (SPNs), crucial in thalamocortical maturation, respond to sounds before ECO showing oddball selectivity. Before ECO, SPNs are more selective to oddball sounds in auditory streams than thalamo-recipient layer 4 (L4) neurons and not after ECO. We hypothesize that SPNs' oddball selectivity can direct development of L4 responses before ECO. Exposing mice before ECO with a rarely occurring tone in a stream of another tone occurring frequently leads to the strengthening of the adult cortical representation of the rare tone, but not that of the frequent tone. Results of control exposure experiments at multiple developmental windows and also with only a single tone corroborate the observations. A computational network model of known thalamic inputs to SPNs and L4 explains the observed developmental plasticity. Information-theoretic analysis with sparse coding assumptions also predicts the observations. Thus, salient low probability sounds in the earliest auditory environment cause long term changes in the ACX.

Introduction

Sensory cortical circuitry development is activity-dependent and is based on the sensory environment to which an animal is exposed[1,2]. Even the earliest experiences, like in utero exposure to a mother's voice, may lead to maternal speech preference in newborns[3] and near term fetal responses to the mother's spoken words[4]. It is well known that in the auditory system, altering the natural auditory environment during specific periods (critical period[5-7]) of development alters cortical circuitry and organization remarkably[8,9]. However, it is not known if the auditory system has the capability to adapt to environmental changes occurring before the known critical period. SPNs, the firstborn cortical neurons[10,11], present in large numbers during and before the critical period[5,12], play a crucial role in the development of functional cortical circuitry and organization[13-17]. Further, SPNs in the ACX are known to be the earliest to respond to sounds at least in ferrets[18]. However, the properties of early SPN sound responses, which are likely to play a crucial role in ACX development, are unknown, especially in the mice. SPNs receive inputs from the thalamic axons projecting to L4[10] in the ACX[11,19] and drive L4 activity over a time window of development[10,20]. Thus, it is important to know what kinds of natural stimuli specifically activate SPNs as their activity coincident with thalamic inputs can sculpt the earliest feedforward thalamocortical circuitry[13,14] in the ACX. SP and L4 neurons play a crucial role in the maturation of thalamocortical circuitry, so it becomes important to know how the

response properties of SPNs and L4 neurons change, starting from the onset of ACX sound
driven activity.

Although ACX responses have been shown to exist in ferrets[18] at ages equivalent to P6
(postnatal day 6) in mice[21], no study reports the presence of sound driven activity in the ACX of
mice at ages before ECO (P11/12). Low threshold hearing onset coincides with ECO[5,22] in the
mouse, with outer hair cell (OHC) based sensitivity allowing detection of near 0 dB SPL sounds.
The developing auditory pathway is functional before ECO[23] as known from ACX sound driven
responses obtained by surgically opening the ear canal[24] in mice. However, moderate-intensity
sounds, for example, at conversational levels (~70 dB SPL) may drive spiking activity in the ACX
before ECO without surgically opening the ear canal, through bone conduction[25]. The presence
of responses before ECO suggests that activity-driven plasticity in the ACX starts earlier than
previously thought. However, previous studies with a single tone exposure at ages before ECO,
show no changes in tonotopy[5,7] or ACX organization in the adult. Thus, even if responses in the
ACX exist before ECO, it is important to understand the response properties in more naturalistic
stimulus paradigms to test the possibility of activity driven ACX plasticity starting before ECO.

Thus, we first investigate the possibility of ACX responses of the L4 cortical plate and SPNs in
mice with ear canals closed as would be required for the earliest natural sensory experience to
drive cortical activity. With multiple techniques, we show that ACX sound-driven activity with
ear canals closed start as early as postnatal day 7 (P7). We further show that, before ECO SPNs
have unique response properties. SPNs, unlike cortical plate neurons, show selectivity to
oddball sounds in a stream of repeating sounds before ECO. While after ECO, cortical plate L4
neurons show stronger deviant detection than remaining SPNs. The observed deviant detection
shown by SPNs in the earliest phase can direct the thalamus to L4 synapse[13,14]. The above
opens up the possibility that specific kinds of most initial sensory experience can modify cortical
development. For example, exposure to music or the structured sound environment during the
fetal period leads to long term plastic effects with enhanced responsiveness to the sounds
experienced in the prenatal period[26]. We use a new auditory exposure paradigm with a low
probability, salient/oddball tone (deviant, D) in a stream of a repeatedly occurring tone
(standard, S). We show that with continuous exposure to the above S-D acoustic environment
at ages before the established auditory critical period[5–7,27] (before ECO), can remarkably alter
the functional auditory cortical (ACX) responses, which persist into adulthood. The alterations
are specific to the deviant or rare exposure tone (D). Performing control exposures with only a
single tone during the same developmental window do not alter responses in the adult ACX.
The observed plasticity is an outcome of the unique deviant detection properties we find of
SPNs, before ECO, with strong responses to only the rare stimulus (D). With a computational
network model derived from our experimental results and known thalamus to SP to L4
connectivity, we show how such plasticity occurs. Further, using mutual information

maximization and sparse coding principles, we show that the outcome of early exposure with
our new paradigm can be predicted theoretically. Hence our results revise the established
timelines and concept of the auditory critical period and may generalize to other sensory
systems.

Results

Mouse ACX and IC have auditory responses before ECO

We performed widefield fluorescence imaging of the ACX in Thy1-GCaMP6f (Jax Labs) at early
ages (Fig. 1Ai) before (n=10; P7-P8=3, P09=4, P10=4) and after ECO (P13, Fig. 1Aii), to
investigate the presence of sound driven Ca^{2+} responses in the ACX. After ECO, by P13, mice
show strong Ca^{2+} responses and defined rostro-caudal gradient of high to low frequencies[28]
marking out the primary ACX (A1) and other auditory responsive regions (Fig. 1Aii). Since the
auditory pathway is immature yet functional before ECO, we next investigate the presence of
auditory responses to tones in the ACX of mice before ECO[5,22] without surgically opening the
ear canals. We found that sound-evoked Ca^{2+} responses are present in the mouse ACX at high
(90 dB SPL) to moderate intensities (70 dB SPL, in awake state) at ages before ECO (Fig. 1Aiii-vi
and Fig. 1B, Supplementary Video 1). With decreasing age, before ECO, progressively less
percentage of areas of the cortical surface showed significant responses to multiple tone
frequencies (Fig. 1Aiii-vi, Table S1), from 80% in P13 to 3% in P7. The percentage of frequencies
of the tone stimuli tested, that showed significant responses in widefield imaging also
progressively reduced (Table S1) from 71% in P13 to 20% in P7. Both the above reflect gradual
maturation of the ACX starting from the onset of sound responsive activity at P7 to achieving
tonotopy in A1 by P13. For further confirmation of sound-evoked responses in the auditory
pathway before ECO widefield fluorescence imaging in the inferior colliculus (IC) of Thy1-
GCaMP6f (Jax Labs) mice at P9-10 was performed. As in the ACX, we found significant sound-
evoked responses in widefield fluorescence on the IC surface (Fig. 1C).

L2/3 and L4 ACX single neurons have responses before ECO

129 The presence of significant auditory responses in Ca^{2+} dependent fluorescence in widefield
130 imaging (Fig. 1) could potentially indicate either the activity of input axons or spiking responses
131 of, neurons in the ACX, or both. To rule out only input axon activity in the ACX widefield
132 imaging, we performed 2-photon Ca^{2+} imaging of single neurons in L2/3 and L4 of the ACX of
133 mice (14 mice) before ECO and after ECO (2 mice; P12-P13) (Fig. 2A) with high to moderate

Figure 1. Widefield fluorescence imaging-based responses to tones in the ACX and IC of Gcamp6f mouse before ECO. A) i) Experimental timeline. ii) Widefield fluorescence imaging performed at P13 Gcamp6f mouse shows defined rostro-caudal gradient of high to low frequencies. Smoothed df/f traces with error bars representing SEMs of selected regions (square) responding to selected frequencies (color) are shown to the right of each image. iii-vi) Widefield fluorescence imaging-based significant response regions (white contours) in the ACX to 90 dB SPL tone in 4 different ages (P07-P10) before ECO. The black horizontal bar represents the stimulus duration. B) As in B, widefield fluorescence imaging in the ACX based significant response 80 and 70 dB SPL tone at P09 and P10 awake mouse. Respective df/f traces with SEMs of the selected region are shown at the bottom of each figure.   C) As in A, widefield fluorescence responses before ECO in the IC at P9 and P10 to 90 dB SPL tone. The "."(Purple) represents the first significant window, while the "*"(Purple) represents the most significant window. Sound-evoked Ca^{2+} responses are present in the mouse ACX at high (90 dB SPL) to moderate intensities (70 dB SPL) at ages before ECO (one-sided paired t-test, *p<0.05,**p<0.01 and ***p<0.001, Black).

intensities of tones. Each 50 ms long tone stimulus was repeated 40 times and consisted of 3-5
frequencies from 6-48 kHz, ½ octave apart, presented at 60-90 dB SPL, with a 5-7s gap between
repetitions. Single neurons in L2/3 in awake mice showed Ca^{2+} transients in response to tones
at ages before ECO (P7-P9, Fig. 2Bi, Supplementary Video 2). The population mean df/f in
response to different frequency tones (Fig. 2Bii) was investigated, considering all the neurons
that responded with at least 25% reliability (fraction of repetitions showing a significant
response, Methods). We found an increase in the mean population response strength before
ECO (14 mice; P07-P08=3, P09=5, P10-P11=6, ANOVA with Tukey post hoc analysis, $F_{(2,576)}$ =
33.5, p <0.0001, Fig. 2Ci), and a decreasing trend in peak latency with age (Fig. 2Bii). The above
indicated a gradual maturation of responses up to ECO (P12-13) in L2/3 ACX neurons. Along
with an increase in tone response strength over age, response reliability increased significantly
with age (Fig. 2Cii, 14 mice; P07-P08=3, P09=5, P10-P11=6, ANOVA with Tukey post hoc
analysis, $F_{(2,6953)}$=375.84, p<0.0001) before ECO, also showing gradual response
development. The percentage of overall cases (imaged neuron and tone stimulus pairs) that
showed significant responses also showed a gradual increase (Pearson correlation, r=0.76, n=6,
p=0.078) with age from 6.84% at P7 to 12.5% at P12/13 (Table S2). All the above observations
indicate the presence of sound driven Ca^{2+} activity before ECO in single neurons and that such
activity varies with age in a manner expected from increasing maturity of the neural pathway.

The widefield Ca^{2+} responses at moderate intensities (Fig. 1B) were also ruled out to reflect only
input axon activity as single neurons of L2/3 showed clear Ca^{2+} transients in response to tones
down to 60 dB SPL (Fig. 2Di). The population mean df/f traces of all neurons (4 mice; 90 dB:
359, 80 dB: 348, 70 dB: 556, 60dB: 144 neurons) that responded significantly (with 25%
reliability, Methods) to tones at the 4 intensities used, showed an increase in response strength
with increasing intensity, except 80 dB SPL (Fig. 2Dii). The percentage of cases of single-neuron
tone stimulus pairs that showed significant responses also showed a similar trend, increasing
from 7% at 60 dB SPL to 40.5% at 90 dB SPL (Table S3).

The previous set of data were collected with usually a little over 1s of baseline preceding the
stimulus. In order to rule out a contribution from any coincident spontaneous oscillating
activity[29] to the response, we obtained extended 2-photon imaging data before ECO (3 mice, P9
and P10, 6 ROIs) with >2s of baseline (Figs. 2E) and also with and without sound stimulation
(Fig. 2F). The population mean df/f traces of all neurons in L2/3, showing significant activity
above baseline in response to tones at P9 (666 neurons) and P10 (138 neurons) portray the
same nature of responses as in Fig. 2Bii. The long duration mean baseline activity from 40
repetitions fluctuated around 0, as expected. We confirmed the presence of sound-evoked
responses before ECO, considering any coincident ongoing spontaneous oscillations by
comparing mean df/f traces of all neurons imaged in with and without sound cases (3 mice, 461
neurons, P9-P10, Fig. 2F). No significant activity was observed under no sound condition.

Figure 2. Sound evoked activity of ACX L2/3 and L4 neurons in the Gcamp6f mouse before ECO. A) Experimental timeline. B) i) Example traces of df/f of single L2/3 neurons (arrows in 2-P mean images of acquired frames with cells expressing GCamp6f at age P07 and P08, right) in response to 0 dB attenuation tones, respectively at three different ages (P07-P09). The black vertical line on the bottom of each trace represents stimulus onset. The corresponding green vertical shading bar in each example trace represents a significant moving window (3 frames) which shows the earliest significant p-value among all the frames (corresponding p-value shown above, one-sided paired t-test, *p<0.05). The error bars show the SEMs of 40 repetitions. B) ii) Population mean df/f of the age groups (color) of L2/3 neurons shown in Fig. Bi. C) i) Box plot showing response peak strength of all L2/3 neurons imaged at different ages (14 mice; P07-P08=3, P09=5, P10-P11=6, ANOVA with Tukey post hoc analysis, corresponding significance-value shown to the right). C) ii) Box plot showing response reliability of all L2/3 neurons imaged at different ages, as shown above (ANOVA with Tukey post hoc analysis, ***p<0.001, corresponding significance-value shown to the right). D) Similarly, as shown in B) i), are example traces of df/f of single L2/3 neurons of ACX in response to 90dB-60dB SPL tones at age group P9-P10 of an awake mouse, with corresponding significance values of the earliest significant window (3 frames) obtained. ii) The population mean df/f of L2/3 neurons across different sound intensity level (90dB-60dB SPL, 4 mice; 90 dB: 359, 80 dB: 348, 70 dB: 556, 60dB: 144 neurons). E) Extended 2-photon Ca^{2+} imaging data, showing population mean df/f (below) at two ages (P09, 2mice, 666 neurons and P10, 138 neurons) before ECO, indicating long (>2s) spontaneous window. F) Single neuron population mean df/f of the same age groups in response to sound (Left) and the same neurons without sound stimulation (Right), with 40 repetitions in each case (P09-P10, 3mice, 461 neurons). G) i) Three example traces of df/f of single L4 neurons (Cyan) in response to 0 dB attenuation tones at two different frequencies (Above row, 10 Khz and Below row, 17 kHz). The green patch represents the earliest significant moving window obtained with respective significance-values. ii) The population mean df/f of L4 neurons which showed significant responses (P9-10, 2 mice, 64 neurons). L2/3 and L4 ACX single neurons respond to sound before ECO (one-sided paired t-test, *p<0.05,**p<0.01 and ***p<0.001, Black asterisk).

As in L2/3, putative L4 single neurons based on the depth of 2-photon imaging (300-350
microns from pia) also showed Ca^{2+} responses. Mean df/f based responses of 3 representative
L4 neurons to 2 frequencies are shown in Fig. 2Gi. As in L2/3, the population mean of all
significantly responding cases in L4 (2 mice, 64 neurons) showed responses to sound with >2s of
baseline activity showing no spontaneous oscillatory activity (Fig. 2Gii). Thus based on 2-photon
Ca^{2+} imaging of single neurons, we find both L2/3 and L4 to be tone sound responsive before
ECO, showing responses as early as P7 and down to moderate intensities of 60 dB SPL.
Together, the above data indicate that widefield fluorescence imaging-based sound responses
in ACX are not due to only input axon related activity.

SPNs in mice are driven by sounds before ECO

Two-photon Ca^{2+} imaging of single neurons, simultaneously from L4 and SPNs was beyond our
scope[30]. Although SPNs are known to be the first to respond to sounds in ferrets, equivalent
properties of SPNs in the mouse at the corresponding age of P6[18] are unknown. Further, our
Ca^{2+} responses do not show the nature of spiking activity in L4, thus requiring extracellular
electrophysiology in mice before ECO. To reliably target SPNs with electrodes at different
developmental ages, we first characterized the depth of SPNs based on SPN specific
markers[31,32] CTGF and Complexin 3 (Fig. 3B, green and red respectively) using IHC and based on
cell morphology from Nissil stains[33] (Fig. 3C). As the thickness of the cortex also varied with age,
we quantified the SPN (Fig. S1E) and L4 (Fig. 3C, Fig. S1D) depth profile normalized to cortical
thickness. The most consistent data was obtained with Nissil stains, and thus absolute depth as
a function of age (P6 to P18) was used for electrode placements. Representative histology data
are shown in Fig. S1A-E. The thickness of the SPN layer reduced with age as known[12,34].

We next performed extracellular recordings in the mouse before ECO (P7-P12) from the ACX
SPNs and L4 simultaneously with custom made staggered electrodes arrays (Fig. 3A, 4*4 arrays
x-y spacing 125 microns, 8 in SPN and 8 in L4, 400 or 600 microns apart in-depth, Microprobe,
USA). Sound-evoked local field potentials (LFPs, Methods) simultaneously obtained from SP
(>=800 microns; P8-P9, 3 mice, 72/536 and P10-P11, 4mice, 138/1248) and L4 (300-350
microns, P8-P9, 3 mice, 65/536 and P10-P11, 4 mice, 140/1248 locations) before ECO showed
significant sound-driven activity (Figs. 3Di-ii, Figs. 3Ei-ii). Longer lasting oscillatory LFPs were
observed at ages before ECO than after ECO (P12-P13, Figs. 3Fi-ii). The above results are in
accordance with the pattern of LFP activity observed in ferrets at ages before ECO[18]. The
population mean LFPs (Figs. 3D-Fii) were quantified with the root mean square (RMS) values in
50 ms bins normalized by the mean RMS values of the baseline (200 ms preceding the stimulus
onset). Significant responses in both L4 (P8-P9; one-sided, paired t-test, t(64)=8.18, p<0.0001,

Figure 3. L4 and SP Neurons in the mouse ACX respond to tones before ECO. A) Schematic of types of 4x4 electrode arrays used – either all electrodes were of same lengths (i), or they had alternate short and long electrodes (ii) either 600 μm apart or 400 μm apart. B) SP depth determined with immunohistochemistry (IHC) by labelling SPNs with antibodies of SPN markers Complexin-3 (Red) and CTGF (Green). C) Depth of L4 (Cyan) and SP (Magenta) over age determined from cell morphology with Nissl stains. D-E) i) LFP in a single channel in L4 (Cyan, Above) and SP (Magenta, below), in response to tone at age, P09 and P10. The green patch represents the window across which the significance was calculated (L4 and SP ACX single neurons respond to sound before ECO (one-sided paired t-test, *p<0.05, **p<0.01 and ***p<0.001)., see methods). The horizontal red bar represents the stimulus onset. D-E) ii) Population average of all the significant recording sites and SEM of RMS values of LFP with corresponding p-values (one-sided paired t-test, ***p<0.001). The Red vertical line represents stimulus onset. F) i-ii) Similar plots of single-site and mean population LFP activity at ages after ECO (P12-P13), as shown in D-E) i-ii), respectively. G-H) i) Dot raster and z-score of an L4 (Cyan, Above two plots) and SP (Magenta, Bottom two plots) neuron, in response to tone at age, P9 and P10. The Red vertical line represents stimulus onset. Inset to the right represents spike shape. G-H) ii) Population mean PSTHs of all the recorded SP (P8-9, 4 mice; P10-P11, 7mice) and L4 (P8-9, 4 mice; P10-P11, 9 mice) units across different age groups which had a z-score>1 in at least 1-time bin within 500ms after stimulus onset, with respective significance-values (one-sided paired t-test). The Red vertical line represents the stimulus onset. I) i-ii) Similar plots of single-unit and mean population spike activity at ages after ECO (P12-P13) as shown in, G-H) i-ii), respectively. J) i) Latencies of L4 and SP neurons recorded simultaneously in P8-P11 (before ECO), ii) P12-13 and P14-15 age groups. iii) The population mean latencies revealed lowering of latencies with age and SPNs having lower latency than L4 neurons at ages before ECO and P12-P13 (one-sided unpaired t-test, *p<0.05 and **p<0.01).

P10-P11; t(139)=11.42 p<0.0001) and SP (P8-P9; t(71)=9.377, p<0.0001, P10-P11; t(137)=10.43
p<0.0001) were observed before ECO that last for > 1s (Figs. 3Di-ii and Figs. 3Ei-ii, Top and
Bottom) unlike after ECO (Figs. 3Fi-ii, Top and Bottom, L4 and SP, respectively).

Single units in L4 (P8-P9, 4 mice; P10-P11, 9 mice, cyan) and SPN (P8-P9, 4 mice; P10-P11,
7mice, magenta) layer showed the presence of spiking responses to tones before ECO (Figs.
3Gi-ii and Figs. 3Hi-ii, Top and Bottom rows respectively) and after ECO (Figs 3Ii-ii). Sample dot
rasters (Top) and z-scores (Bottom) of spiking responses in 50ms time bins shown in Figs. 3Gi-ii
and 3Hi-ii indicate the presence of significant sound-evoked spiking before ECO. Population
PSTHs of all significantly responding units in SP (P8-P9; one-sided paired t-test, t(59)=1.733,
p=0.043; P10-11, t(259)=10.07, p<0.0001) and L4 (P8-P9; SP, t(51)=5.738, p<0.0001; P10-P11,
t(451)=12.678, p<0.0001) show significant sound driven activity before ECO, which get stronger
with age (Figs. 3G-I ii). The response peak latencies (Methods) of L4 and SP single-units were
obtained at ages before (P09 example trace, Fig. 3Ji) and after ECO (P12 example trace, Fig.
3Jii),. Before ECO and at P12-P13 the mean peak latencies of SPNs were found to be
significantly lower than L4 neurons (one-sided t-test, before ECO, t(439)=-2.003, p=0.022; P12-
P13, t(252)=-2.935, p=0.002, Fig. 3Jiii). The above results suggest that SPNs respond at shorter
latency relative to L4 neurons before ECO, and the response properties mature with age. In
subsets of experiments, at multiple ages, confirmation of electrode location in ACX was done
with DiI crystal insertion in the recording site and observing labelling of MGBv (Fig. S1A). Also,
in higher age groups, A1 recording was confirmed with the presence of general tonotopy (Fig.
S1B) based on the spatial location of best frequencies (BFs) of units from multiple electrode
penetrations. Electrode tracks were also routinely confirmed to be in the SPN layer (Fig. S1Di).
In a subset of experiments in SCNN1-tdtomato mice (Jax 9613, P14, Fig. S1Dii) with staggered
electrodes, confirmation of electrodes being simultaneously in L4 (L4 neurons expressing
tdtomato) and SPN (from IHC, Fig. S1Dii) was done (3 mice, P12, P14 and P17). Thus based on
the data from multiple techniques, we conclude that neurons in the different cortical layers and
SP layer respond to sounds before ECO, without surgically opening the ear canal. Auditory
responses are present as early as P7 in mice and respond down to 60 dB SPL tones.

SPNs, unlike L4 neurons, show oddball selectivity before ECO

Having established the presence of auditory responses before ECO, we investigate the response
properties of SPNs and L4 neurons in the context of stimulus-specific adaptation[35–37] (SSA). We
use oddball stimuli with a stream of standard tokens (S) with an embedded deviant token (D),
to mimic the natural situation of a continuous auditory environment with important sounds
occurring with low probability[35,37] (Fig. 4A). The stimuli consisted of 50 ms long sound tokens
presented at 3.3 or 4 Hz, of the pattern SSS...SDS...S, (D at 8[th] or 7[th] position of 15 or 10 total

tokens) for the oddball response characterization. One of the S and D tokens was a tone (f,
range 6-34 kHz) and the other a broadband noise (N, bandwidth 6-48 kHz with equivalent
sound level). Responses were collected in pairs with the S and D swapped to normalize for a
neuron's inherent selectivity for a particular S or D, (Fig. 4Ai). Each SD set of a pair was
presented with 10-40 repetitions and 5-7s silence duration between the end of the previous
and start of the next stream. A common selectivity index (CSI) was used to quantify a neuron's
deviant detection strength:

$\text{CSI}=[R_D(f)+R_D(N)-R_S(f)-R_S(N)] / [R_D(f)+R_D(N)+R_S(f)+R_S(N)]$[35];

where $R_S(.)$ and $R_D(.)$ denote response to S and D, respectively. Two cases for $R_S(.)$ were
considered: mean response to the first S or onset and mean response for all the S's; the deviant
detection strength values are referred to as CSI(F) and CSI respectively. To define oddball
selectivity, if a neuron evoked higher responses to D compared to S in an oddball sequence with
corresponding CSI value positive, then it was considered to exhibit oddball selectivity. We refer
firing rate based oddball selectivity as deviant detection.

ACX L4 (300-350 microns, 11 mice, 84 neurons) and SP (>800 microns, 11 mice, 34 neurons)
single units responses were obtained before ECO in response to the presentation of the SD
stimulus for f-N and its swap. Sample PSTHs of SPN before ECO shows the preference of the
oddball or to the deviant tone (f) token (Fig. 4Aii). In general, response to the SD stimulus
showed peaks in the response following the D-token, mainly in SPN and the first S-token or
onset, primarily in L4 (Fig. 4B, L4 top row, SP bottom row). The earliest response peaks of L4
and SPN at these ages with the SD stimuli occur in a 250 ms window with a latency of 50-100
ms from the first token and in a similar window following the deviant (Fig. 4B). Comparing the
mean response to the S-token and that of the D-token (Fig. 4B, S: green, D: red) shows the
above. In the overall population mean response before ECO (P8-P12) for all L4 neurons, and all
SPNs recorded indicate the presence of deviant detection in SP and not L4 (Fig. 4Ci-ii). Further,
based on the spontaneous activity (100 ms preceding the stimulus), z-scores of the example
neurons and population of neurons (Fig. S2) demonstrate a significant response to D and S and
that in the population SP has significant activity at the deviant and not L4. To quantify the
selectivity of the oddball, we obtaining spike rates in 250 ms windows because of the width of
the peaks and calculated CSI(F) and CSI for all SPNs and L4 neurons. We observed higher
deviant detection in SP than in L4 for both CSI (one-sided unpaired t-test, $t(116)=3.666$,
$p<0.001$, Fig. 4Di) and CSI(F) ($t(116)=3.954$, $p<0.001$, Fig. 4Dii).

To rule out spurious deviant detection due to coincident spontaneous oscillating activity during
the stimulus, we also considered CSIs expected by chance. 1000 random window pair sets each
were considered in random-CSI calculations. Two non-overlapping windows 250 ms long were
taken randomly in the standard region before the deviant or after the deviant not considering
the deviant response. The mean CSIs were not significantly different between L4 and SP in
>90% of the random window sets. More importantly, the distributions of mean random-CSIs for

Figure 4. SPNs, unlike L4 neurons, show higher oddball selectivity before ECO. A) i) Schematic showing the Tone-Noise (f-N) oddball stimulation protocol with the swap on the right. A) ii) PSTHs of a single SP neuron, showing higher response at deviant tone (f) token (Swap, 250 ms response window gray patch, Right). B) Example traces of spike rate of 3 L4 neurons (upper 2 rows) and 3 SP neurons (lower 2 rows) in the first 3 columns, showing response to tone (f, Standard in Green and Deviant in Red, Above) and noise (N, Standard in Green and Deviant in Red, Below). A light gray patch is the mean rate window (250 ms) across which CSI(F) was computed. C) i) Mean population spike rate of all L4 and SP neurons (upper and lower two plots, respectively) in response to tone (f, Standard in Green and Deviant in Red, Above) and noise (Standard in Green and Deviant in Red, below) presented at 3.33 Hz. C) ii) Same as C i), mean population spike rate of L4 and SP neurons in responses to tone and noise, delivered at 4Hz. D) i-ii) Population (3.3 and 4 Hz combined) Cumulative distribution of CSI(F) and CSI for SP and L4 (Magenta and Cyan trace, respectively) show significantly higher oddball selectivity in SP (one-sided unpaired t-test, ***p<0.001). Similar Cumulative distributions with mean CSI calculated for each of 1000 random pair of SD (Standard-Deviant) response windows selections for S response and D response either before deviant (Light dashed lines, Di) or after deviant (Light Dashed lines, Dii). No significant difference between L4 and SP in >90% of the random window sets.

SPN and L4 showed that the means were not significantly different from 0 (dotted lines, Fig 4Di-
ii).

Further to ensure that our results were not sensitive to the choice of latency and window size,
we also varied the response window from 50 ms to 300 ms (in steps of 50 ms) and latency from
0 to 350 ms (in steps of 50 ms). We obtained the CSIs for all neurons, and we found for what
combinations SPNs had significantly higher oddball selectivity than L4 neurons (Fig. S3). For a
substantial range of values around the chosen latency and window size, the same results were
obtained, indicating deviant detection in SPN and not in L4 before ECO.

Unlike before ECO, L4 shows higher oddball selectivity than SP following ECO

Oddball selectivity in adults is a common phenomenon in all cortical layers of ACX[38]. Given the
observed absence of oddball selectivity in populations of L4 neurons before ECO, we
hypothesize that such selectivity in L4 starts after ECO. Thus, we next investigated how the
oddball selectivity changes in L4 and SPNs after ECO. Single-unit responses to the same SD
paradigm with *f-N* sound tokens (Fig. 4A) were collected simultaneously from L4 (P12-P13, 10
mice, 45 units, 57 cases; P14-P15, 8 mice, 66 units, 115 cases; P16-21, 4 mice, 20 units, 31
cases; P22-P28, n=6 mice, 38 units, 97 cases ; >P28, 7 mice, 44 units, 63 cases) and SPNs after
ECO from P12 to the juvenile/adult at >P28 (P12-P13, n=10 mice, 54 SPN units, 101 SPN cases;
P14-P15, n=8 mice, 59 SPN units, 67 SPN cases). We again quantify the selectivity based on
CSI(F) and CSI. Comparison of population PSTH segments in response to tone (Fig. 5Ai) or the
noise (Fig. 5Aii) as the first S token (green), all S tokens (blue) and the deviant (red) show how
selectivity to the oddball changes in L4 and SP over age after ECO starting from P12-13. Since
direct comparisons are confounded by the inherent selectivity to the *f* or *N*, we compare CSIs
over age (Fig. 5Aiii and iv) including the CSIs before ECO. L4 neurons start showing oddball
selectivity immediately after ECO, which gradually declines to stable values with age till
adulthood (CSI(F): ANOVA with Tukey post hoc analysis, $F(4,358)=37.68$, $p<0.0001$; CSI:
$F(4,358)=12.74$, $p<0.0001$). On the other hand, SPNs with a significantly lower number at P12-
P13 compared to peak at P8[34], show significantly lower CSIs than L4 after ECO (CSI(F): one-sided
unpaired t-test, P12-13$_{L4}$ vs P12-13$_{SP}$, $t(156)=-1.944$, $p=0.027$; P14-15$_{L4}$ vs P14-15$_{SP}$, $t(180)=-$
1.10, $p=0.13$; CSI: P12-13$_{L4}$ vs P12-13$_{SP}$, $t(156)=-1.621$, $p=0.050$; P14-15$_{L4}$ vs P14-15$_{SP}$, $t(180)=-$
1.41, $p=0.07$). We considered both CSI(F) and CSI since at early ages due to high adaptation
(decrease in time constant with age: $F_{L4}(4,455)=11.92$, $p<0.0001$) in neurons with robust
responses at the first S token (responding tokens increases: $F(4,455)=6.34$, $p<0.0001$) and not
any others (Fig. S4). The same trend was observed in both CSIs, showing that before ECO SPNs
are stronger deviant detectors than L4 while after ECO the relative deviant detection strength
switches.

Above, we used *N-f* SD stimuli to mimic a repeating broadband auditory environment with the
tone as a low probability stimulus (like a pup or adult mouse vocalizations[39] in a normal
auditory environment). Since deviant selectivity has been studied primarily with two tones[35,37]
(f_1 and f_2), we also used an f_1-f_2 SD stimulus set, to investigate if our results also hold in such a
case. Further, using tones as both tokens allows more direct comparisons between responses.
f_1 and f_2 were spaced either 1/4, 1/2 or 1 octave apart. Data were obtained from multiple L4
(P12-P13, 3 mice, 28 units, 47 cases; P14-P15, 3 mice, 41 units, 63 cases; P16-P21, 3 mice, 11
units, 15 cases; P22-P28, 4 mice, 24 units, 53 cases; >P28, 3 mice, 26 units, 74 cases) and SP
(P12-P13, 3 mice, 9 SPN units, 14 SPN cases; P14-P15, 3 mice, 34 SPN units, 51 SPN cases) single
units across different developmental windows. Data were collected from single units
simultaneously, and hence f_1 and f_2 were at different distances from the BF of each of the units.
One of the frequencies (f_1 or f_2) was chosen to be at or near the BF of the majority of units. The
responses to the SD paradigm were used to compare mean PSTH segments in response to the
first S token, all other S tokens and the D token (Fig. 5B) as previously. A similar trend of
decrease in CSIs in L4 over age was observed, as with *N-f* stimuli (CSI(F): ANOVA with Tukey
post hoc analysis, $F(4,247)=30.42$, $p<0.0001$; CSI: $F(4,247)=13.83$, $p<0.0001$)(Fig. 5B). From
CSI(F) and CSI, we found that L4 neurons are more selective (CSI(F): one-sided unpaired t-test,
$P12-13_{L4}$ vs $P12-13_{SP}$, $t(59)=1.8731$, $p=0.033$; $P14-15_{L4}$ vs $P14-15_{SP}$, $t(112)=2.866$, $p=0.002$; CSI:
$P12-13_{L4}$ vs $P12-13_{SP}$, $t(59)=3.343$, $p<0.001$; $P14-15_{L4}$ vs $P14-15_{SP}$, $t(112)=2.163$, $p=0.016$) to
oddballs than SPNs after ECO.

The observed high oddball selectivity in SPN or L4 at different developmental ages could be a
reflection of subcortical deviant selectivity[40,41]. The MGBv is the main thalamic nucleus from
which A1, L4 and SPN get inputs[11,19] (Fig. 5Ci-ii). Thus we quantified CSIs from responses of
MGBv single units to the *N-f* SD stimuli (Fig. 5Ciii-iv). In both age groups considered (P14-15 and
P24-26) we find that CSI(F) and CSI were lower than that of L4 (CSI(F): $P14-15_{L4}$ vs $P14-P15_{MGBv}$,
$t(180)=-4.373$, $p<0.0001$; $P22-28_{L4}$ vs $P24-P26_{MGBv}$, $t(163)=-0.031$, $p=0.488$; CSI: $P14-15_{L4}$ vs $P14-$
$P15_{MGBv}$, $t(180)=-7.370$, $p<0.0001$; $P22-28_{L4}$ vs $P24-P26_{MGBv}$, $t(163)=-6.209$, $p<0.0001$) and SPN
(CSI(F): $P14-15_{SP}$ vs $P14-P15_{MGBv}$: $t(132)=-2.881$, $p=0.002$; CSI: $P14-15_{SP}$ vs $P14-P15_{MGBv}$: $t(132)=-$
4.006, $p<0.0001$) in developing mice and also lower than L4 in the adult. Thus, the observed
oddball selectivity even at the early ages is emergent in the cortex and expectedly a network
phenomenon[42].

Figure 5. Sound evoked activity of ACX and MGB neurons after ECO. A) i) Schematic showing the Tone-Noise (f-N) oddball stimulation protocol with the swap. ii) Mean population PSTH of all significant responding neurons (L4 neurons above, SP neurons below), indicating the deviant response (Red), the first standard response (Green), and the mean of responses at all the standards (blue) to tone (Above 2 rows) and noise (Below, 2 rows) for different age groups after ECO. A) iii-iv) Overall mean oddball selectivity for noise and tone tokens measured by CSI(F) (above) and CSI (below), respectively as a function of age. B) i) Similar mean population PSTH for tone-tone (f1-f2) oddball stimulation protocol. B) ii-iii) As in A) iii-iv), Overall mean oddball selectivity for tone stimuli measured by CSI(F) and CSI respectively as a function of age. C) i) Example of electrode tracks (white arrowheads) shown in Nissl stains of brain slices showing recording locations in MGB of the thalamus. ii) Shows a magnified view of the relevant part of a coronal section with MGB, in which single-unit recordings from MGBv was performed (P14). iii) Mean and standard errors of CSI(F) (above) and CSI (below) for the thalamic neurons, by age groups, are shown. *p<0.05,**p<0.01 and ***p<0.001.

Exposure in a standard deviant acoustic environment before and after ECO show differential developmental plasticity for the standard and the deviant frequencies

Having found higher deviant detection in ACX SP than in L4 before ECO (P7-P11) and the opposite after ECO (P12-P15) during the known critical period, we next investigated the functional implications of our observations. Previous studies[5,7] have shown that there is no change in the ACX organization in adult rodents following exposure to a continuous stream of a single frequency tone before ECO. Our results indicate that while L4 and SPNs are responsive before ECO, the above observation is expected given the strong SSA observed in SPNs before ECO (Fig. 4D), responding to the D stimulus and not the S. We hypothesize that because of higher deviant detection by SPNs before ECO (Fig. 4D), and because SPNs drive L4 neurons[19] at early ages, exposure to an SD auditory environment before ECO could influence activity-driven plasticity of the ACX.

Adult ACX frequency representation of normally reared mice (EX0) was compared with mice exposed to an SD sound environment (S occurring with probability 0.9 and D with probability 0.1), at 4 different developmental stages between P6 and P21 (Fig. 6A, EX1-EX4). We exposed each of the groups to an f1-f2 combination of S=12 kHz and D=17 kHz. Additionally, to rule out frequency combination specific effects, for the most relevant age groups, namely P6-11 (EX2) and P11-16 (EX3), we also did additional experiments with another f1-f2 combination (P6-11: S=20 kHz and D=14 kHz, P11-16: S=14 kHz and D=10 kHz). In all the above cases stimuli at 90 dB SPL were used to ensure sound driven activity in ACX before ECO. Since we found that with ear closed, responses in the ACX existed for as low as 60 dB SPL, for our exposure study to be functionally relevant we did further experiments by exposing the P6-11 group at 70 dB SPL also (EX5). Finally, as stimulus control, we performed exposures of P6-11 (EX6) and P11-P16 (EX7) groups with an SS paradigm in which cases only one frequency (12 kHz) was used (EX6 and EX7 respectively).

Single-unit recordings in L4 (depth 350-450 m) in response to pure tones (13 frequencies, 6-48 kHz, 1/8th octaves apart, 50-90 dB SPL, 10 dB steps) from A1 and AAF were performed in the exposed (EX1-EX7) and normally reared (EX0) mice at P28-42 (Fig. 6A). All significant response firing rates from the tuning curves (cartoon tuning curves, Fig. 6B) for each of EX0 to EX7 were obtained. EX1, EX4 and control (EX0) had the same median firing rate (rank-sum test, 12 sp/s for every case, p_{EX1}=0.8475, p_{EX4}=0.1875). All other groups, EX2-3 and EX5-7 show, increased median firing rates (rank-sum test, 16-24 sp/s, $p_{EX2,3,5,6,7}$ <0.0001 compared to EX0. Thus with auditory environment manipulations in the most relevant groups P6-P11 and P11-P16 (the known critical period), the overall median firing rates changed. With our limited resolution BF mapping, we did not find any apparent over-representation of any particular frequency in BF maps showing broad tonotopy (A1 and AAF) on the cortical surface (Fig. S5A) in any of the

exposure cases, unlike other studies with the equivalent of the EX3 group (Fig. S5B). Other than
the lower spatial resolution of single unit sampling, the above may also be due to parameters of
exposure (sound frequency, duration and repetition rate). We considered only mice where
recordings based on BF gradients were in A1/AAF.

Absence of over-representation of any particular frequency and increased firing rates in groups
exposed during the known critical period and groups exposed during the period immediately
preceding critical period, suggested plasticity in the strength of responses. Thus, we next
investigated if the effect of exposure was on firing rate responses to tones and if changes were
frequency specific. From the tuning curves, the population mean response rates to a reference
frequency (RF) were plotted as a function of the unit's BF relative to the RF (Fig. 6B, Methods).
The RF could be the frequency of S or D or that of SS. For S/D exposures (EX1-5) we obtained
two such plots, with S as RF and D as RF (green and red, respectively), for SS exposures (EX6-7),
S was the RF (blue). For the control group C (gray) since there was no actual RF, each of the 13
frequencies presented was considered as RFs separately. Based on the 13 mean rate profiles as
a function of BF re each of the 13 RFs, the maximum rate at each of the BFs (re RF) is
considered as the upper boundary of observed rates at RF in control (EX0, Fig. 6C, gray), the
lower bound being '0'.

The population mean rate as a function of BF relative to RF (re RF) for EX2&6, EX3&7 and EX5
(Fig. 6Ci, ii and iii respectively) showed differential effects on response strength at BF to the S
and D for SD exposure before ECO and after ECO. Exposure after ECO in the known critical
period of P11-15, the S frequency responses at S as BF are strengthened, in comparison to that
of D (EX3; Fig. S6, one-sided unpaired t-test, t(107)=2.65, p=0.0045). However, before ECO, the
opposite effect is observed (EX2; Fig. S6, one-sided unpaired t-test, t(120)=3.08, p=0.0013). No
such changes are observed in exposure before P6 (EX1; Fig. S6, one-sided unpaired t-test,
t(81)=0.15, p=0.55) and after P15(EX4; Fig.S6, one-sided unpaired t-test, t(58)=0.88, p=0.81).
Similarly at 70 dB SPL exposure (EX5; Fig. S6, t(57)=1.507, p=0.06). Only in the case of SS for
P11-15 exposure (EX7), there is an increase of S responses of neurons with S as BF beyond the
boundary of control (gray dashed line, Fig. S6, one-sample t-test, t(17)=2.68, p=0.015). A higher
response than the boundary of control for D was found before ECO SD exposure at both 90
(EX2, one-sample t-test, t(42)=2.80, p=0.007) and 70 dB SPL (EX5, one-sample t-test, t(38)=5.06,
p<0.0001). Comparisons (significant or not) with a population of 13 control rate profiles are
marked on respective bars (gray '*' or NS). When considering each set of exposure frequencies,
for the merged groups of SD exposure with different sets S and D frequencies, the response to
D is greater than that of S before ECO (one-sided unpaired t-test, t(90)=3.07, p=0.0014, 20/14
kHz and t(28)=0.95, p=0.17, 12/17 kHz had comparatively less tuning curves, both 90 dB SPL
and t(57)=1.5, p=0.06 12/17, 70 dB SPL). After ECO, the response for D was found to be less
than that of S (t(51)=1.53 p=0.06 14/10 kHz and t(54)=2.11 p=0.019 12/17 kHz).

Figure 6. Exposure to an SD (Standard-Deviant) stimulus protocol, before ECO, induces long term plastic changes. A) Different exposure
paradigms by age and type of exposure stimulus. B) Schematic explaining the construction of Reference frequency (RF) curves (see methods). C)
i) RF curves of P6-P11 (before ECO) for SD and SS (only standard) exposure protocol at 90 dB SPL. ii) P11-P15 (after ECO), SD and SS protocol at
90 dB SPL, and iii) P6-P11 SD protocol at 70 dB SPL. The grey line represents the control group (EX0), showing the maximum boundary for all the
13 frequencies (see methods). iv) Bar plots of all the exposure groups showing RF rates at BF normalized by the rates at all the other
frequencies 0.5 octaves away on either side (see methods). The stripped bars in the P6-P11 group represent the SD exposure protocol at 70 dB
SPL (EX5). The grey bar represents the control. *p<0.05,**p<0.01.

Absolute rates at BF for all frequencies could, in general, be higher after exposure. Although we
did not get any over-representation, we did get a higher median rate for the relevant exposed
groups. We investigated the effect of the exposure paradigm on the rates of all the neurons
with respect to their BFs. To do that we made reference frequency plots (Fig. 6B, see methods)
and normalized the BF rates of the neurons with the BF as RF (center peak values of the plots)
by the mean BF rates of the neurons with BF >1/2 octave (two bins on either side) away from
the exposed frequencies. A value more than 1 will signify a higher rate enhancement at the BF
for the neurons with the BF as RF, while a lower value will indicate a higher rate enhancement
at the BF for neurons with BF away from the RF. We found that in the SD exposure the
contribution of the D frequency in the rate of neurons with D as BF is significantly higher before
ECO (EX2) as compared to the contribution of S frequency in the rate of neurons with S as BF
(Fig. 6Civ, bar plots, one-sided unpaired t-test, t(120)=2.15, p=0.016 at 90 dB and t(57)=1.76,
p=0.041 at 70 dB SPL) and the opposite is true after ECO (EX3, one-sided unpaired t-test,
t(107)=2.71, p=0.004). The contribution of S in the rate of neurons with S as BF in the SD
exposure before ECO (EX2) is lower than 1 (one-sample t-test, t(78)=-2.75, p=0.0078) but does
not show any effect after ECO (EX3, one sample t-test t(75)=-0.83 p=0.39). Contribution of S at S
as BF is not significantly different than 1 in the SS exposure both before (one-sample t-test
t(6)=-0.37 p=0.72) and after ECO (t(17)=-1.23 p=0.23). As in the case of absolute rates, we
looked into the frequency pairs individually and got similar results for before ECO (EX2, one-
sided unpaired t-test, t(90)=2.16, p=0.016 20/14 kHz and t(28)=0.95, p=0.17 12/17 kHz had
comparatively less tuning curves, both 90 dB SPL and t(57)=1.71, p=0.041 12/17, 70 dB SPL) and
that of D is less than that of S after ECO (EX3, t(51)=1.61 p=0.056 14/10 kHz and t(54)=2.19
p=0.016 12/17 kHz). Thus there is a differential long term plasticity of responses to the low
probability sound frequency depending on the period of development during which animals are
exposed in the SD environment. It coincides with the differential oddball selectivity before and
after ECO observed in the ACX L4 and SPNs involved early in development, in setting up the ACX
circuitry.

Model binary network with observed oddball selectivity of SPNs and L4 shows a stronger
representation of deviant frequency before ECO and standard frequency after ECO

Having found long term plastic changes specific to the deviant in the ACX with exposure to an
SD environment before ECO (P6-P11), we next explain our observations based on known
circuitry of SPN, L4 and thalamic inputs[11,19]. SPNs receive stronger thalamic inputs than L4 early
on and SPNs project to L4 neurons[19] (Fig. 7A). In our simultaneous SP-L4 recordings with
staggered electrodes before ECO, 42 pairs of neurons (between L4 and SP) were found to be
connected (Fig. 7B-C and Methods) out of 136 simultaneously recorded L4-SP unit pairs. After

ECO (P12-15), we found 176/244 L4-SP unit pairs to be connected. The direction of connections
was obtained based on significant cross-correlations at positive or negative time lags and
confirmed via granger-causality test. (SP->L4, magenta or L4->SP, cyan, corresponding examples
in Fig. 7B and population data in Fig. 7C). Assuming columnar sizes of ~100µm, we grouped the
pairs as columnar ($\leq$ 125 µm) or not (>125 µm), based on the lateral distance between the
electrodes from which connected pairs were obtained (Fig 7C). Observed directionality of the
connected pairs before ECO suggests the existence of connections from SPN->L4 (12/16 pairs
showed connections from SPN->L4 within a spatial distance of $\leq$ 125 µm, while 13/26 pairs
were found to be connected at a distance of >125 µm, Fig 7C, first column, top). However, the
L4->SP connection probability within $\leq$ 125 µm (4/16 connected pairs) was found to be
significantly lower than SPN->L4 (one-tailed z test, z= -2.828, p=0.002, Fig 7C, first column, top).
Moreover, L4->SP connections were found to be significantly higher than SP->L4 at ages after
ECO (one-tailed z test, z=4.746, p<0.0001, Fig 7C, first column, bottom). The summary of
connectivity between L4 neurons and SPNs (Fig. 7C) shows that SP->L4 connections are
significantly reduced (one-tailed z-test, z= -2.600, p=0.004) from before ECO to after ECO.
Granger causality (GC) analysis[43,44] also revealed similar connectivity profiles between SP and L4
at ages before and after ECO, as observed using cross-correlation (Fig. 7C, second column, top
& bottom). It is also evident from firing rates of L4 neurons that the thalamic drive increases
after ECO as also observed in slice recordings[19]. We also found that recurrent connections,
obtained with cross-correlation as well as GC, within SP (SP->SP) and within L4 (L4->L4) were
differentially altered from before ECO to after ECO (Fig. 7D). Significantly more recurrent
connections were present within SP than within L4 before ECO(z=5.437, p<0.0001, Fig. 7D,
Above) and which switched after ECO (z=9.655, p<0.0001, Fig. 7D, Below). SP to SP recurrent
connections decreased with ECO while that in L4 increased, very much like the observed CSI
values in SPN and L4 before and after ECO, expected from the role played by recurrent
connections in deviant detection[42]. Our observations on deviant detection strength measured
by CSIs are thus further corroborated by the lateral connection changes observed in the two
stages in the two layers.

Thus at the earliest stages stimuli through thalamic inputs that drive SPNs would drive the L4
neurons. Such activation of L4 neurons with coincident activity in the thalamus to L4 synapse
would strengthen the synapse following Hebbian plasticity. Thus the observed deviant
detection of the SPNs responding to D and not the S would strengthen the synapse on L4
conveying the D information. On the other hand, after ECO, the thalamus to L4 synapses are
already stronger than before ECO, and the L4 neuron responds to both the S and D stimulus,
preferring S more. The SP to L4 synapse is weaker than before ECO and is not as strong in
evoking spikes in L4 by itself. Thus on exposure to SD after ECO, synapses conveying D
information from the thalamus weaken, while that conveying S information strengthens.

Figure 7. The binary network model of a developing ACX with subplate (SP) and thalamocortical (TC) inputs both before ECO, and after ECO, the
size of the blobs represent the synaptic weight which changes across time. A) A network schematic showing standard (S, as Green in SD and as
Blue in SS paradigm) and deviant (D, as Red in SD Paradigm) TC inputs to both subplate and layer IV and a subplate input to layer IV (top),
Standard-Deviant (SD) and Only Standard (SS) stimulation protocol (bottom). B) Example pair of correlograms for SP to L4 (left) and L4 to SP
(right) connections. The grey plot represents the bootstrapped shuffled data (control). The "*" represents significant time windows (95% CI) for
both before ECO and after ECO. An example EPSP profile, showing synaptic depression. C) Population bar plots showing connection probability
between SP and L4 for different distances according to cross-correlation(left) and Granger-causality (right). D) Connection probability for SP-SP
and L4-L4 connections. E) i) An example EPSP profile, showing synaptic depression. ii) The asymmetric STDP learning followed by all the plastic
synapses, showing a larger LTD window. The synaptic weight modifications occurred over -80 to +40ms of post-pre spike times. F) Example
PSTHs showing higher oddball selectivity in subplate (Magenta) than layer IV (Cyan) before ECO and the opposite after ECO. G) Temporal
evolution of the relevant synaptic strengths (S) in the two exposure periods (i: Before ECO, SD: EX2; SS: EX6 and ii: After ECO, SD: EX3; SS: EX7)
for simulations of exposure with SD or SS protocol (the inset schematic shows the final point of the simulation with the blob sizes representing
relative synaptic weights). H) i) Plots showing the mean spike time difference (post-pre) across time and ii) mean relative change in synaptic
weight (Δw/w). *p<0.05,**p<0.01 and ***p<0.001.

We test the above explanation for the observed exposure dependent developmental plasticity
based on L4 and SPN deviant detection through a binary network model (Supplementary
Material, Fig. 7A) of early developing TC circuitry of the ACX[19] similar to that considered in the
visual cortex[14]. To have a minimal model that could potentially explain the results we did not
consider lateral connections, although the presence of recurrence will, in principle, only
improve the results. Thalamic inputs selective for either of two inputs, S or D (green and red,
Fig. 7A) are considered to make synapses on an SPN, sending collateral inputs to L4 neuron[10].
The SP and L4 neurons are modelled as integrate and fire neurons with absolute and relative
refractory[14]. The SPN neuron makes a synapse onto the L4 neuron as known[10]. All synapses
were considered depressing in nature[42,45] such that L4 and SPN responses show similar
adaptation observed early in development (Fig. 7Ei and Fig. S4). Also, all synapses except the
thalamic inputs to SPN are plastic following an asymmetric Hebbian spike-time-dependent
plasticity (STDP[14]) rule (Fig. 7Eii). The abscissa in Fig. 7Eii represents the difference in pre-and
post-synaptic spike times ($\delta\tau$, post − pre), with positive $\delta\tau$ leading to LTP and negative $\delta\tau$
leading to LTD. The ordinate represents the percentage relative change in synaptic weight
($\Delta w/w$) for each pre and postsynaptic spike pair.

We consider the model with two different starting points, one representing the period before
ECO (Fig. 7A, left) and another after ECO (Fig. 7A, right). In the earlier developmental period
(before ECO), the SP CSI was 0.41, and that of L4 was -0.02, and in the later developmental
period SP CSI was -0.17, and that of L4 was 0.4 (Fig. 7F), as in our observations (Figs. 5Aiii-iv).
The differences in the initial point of the network (Supplementary Material) in the two cases
(Fig. 7A left and right) are supported by other data[19] and our observations of larger time
constants later in development (Fig. S4), increased thalamic synaptic weights on L4 due to
normal development, lower SPN to L4 connectivity (Fig. 7B-C) and reduced SPN rates through
higher thresholds. We use the same exposure protocols SD and SS for the two periods to show
the nature of how the thalamic inputs to L4 develop (Fig. 7G). Before ECO with higher oddball
selectivity in SP than in L4, with an SD exposure, the D thalamic input to L4 strengthens faster
than the S thalamic input (Fig. 7Gi). On the other hand, after ECO, with the oddball selectivity
reversed in SP and L4, D to L4 synapse weakens and S to L4 strengthens, as L4 spikes more due
to thalamic input spikes in S and not for that of D (Fig. 7Gii). The above is evident when
considering the relative pre and postsynaptic spike times ($\delta\tau$, post − pre) driving LTP or LTD of
the S->L4 and D->L4 synapses in the two cases – before and after ECO (Fig. 7Hi, Supplementary
Methods). The mean strength of weight change occurring in the synapses based on all $\Delta w/w$ for
each of the pre and postsynaptic spike pairs separated by $\delta\tau$ also show the same effect on
synaptic strength change (Fig. 7Hii). Mean $\delta\tau$(D->L4) is positive before ECO and negative after
ECO with a significant difference. Thus the effect of SD exposure before ECO is overall long term
potentiation of the D->L4 synapse while that after ECO the same synapse weakens as observed

in the model results (Fig. 7G). The opposite is true for the S->L4 synapse. Before ECO with an SS
exposure the S thalamic input does not show strengthening at a fast rate (as D, above) leading
to the normal representation of the S input (Fig. 6Civ, bar plot P6-11) equivalent to S in the SD
case. After ECO, with the only S stimulus also the S inputs, expectedly strengthens leading to
normal responses to S as observed (Fig. 6Civ, barplot).

Sparse coding and information maximization principles imply a strengthening of response to
low probability stimuli

While a network model mechanistically addresses the observations, we theoretically address
the observations of early exposure to a low probability stimulus leading to the strengthening of
its responses and its weakening in later exposure case. Our observations and model results are
contrary to the general ideas of long term plasticity - a repeated presentation of a stimulus in
the critical period leads to the strengthening of the representation of that particular stimulus in
the long term[1,5,7,9]. Auditory cortical spiking activity is known to be sparse and sparse coding
principles theoretically explain several known receptive field types in the auditory pathway[46,47].
We assumed that a minimal activity-based coding in single neurons underlying sparse
representation in populations and that such representation is one of the goals of the
developing auditory system while maximizing information about the stimulus in the responses.
With these principles, we use an objective function (Eqn. 1) as the desired optimization
performed by activity-driven plasticity.

$CI_k = I(R_k, T) - \rho(r_0 + x a_k)$ ... (1a)

$CI_{total} = \sum_{k=1}^{N} CI_k = \sum_{k=1}^{N} I(R_k, T) - \rho\left(N r_0 + x \sum_{k=1}^{N} a_k\right)$ ... (1b)

In Eq. 1, $I(R_k, T)$ is the mutual information (*MI*) between the stimulus (a random variable *T* which
is either S or D, with probabilities (1-*x*) and *x*, where *x* is small and is considered as the
probability of the deviant stimulus) and the response rate in the k^{th} token is R_k, ρ is a constant,
N is the total number of stimulus tokens. CI_k is considered as the constrained information
function at the k^{th} token. The mean ensemble response rates are assumed to be r_0 and
$(r_0 + a_k)$ for S and D at the k^{th} stimulus respectively, and assumed to have a Gaussian
distribution both having variance σ^2. The mathematical proof of the optimization is provided in
the supplementary material. However, the basis is as follows. The mutual information would
increase as the separation between the two rates (a_k) increases. However, since the sparse
activity requires minimal activity by single neurons to code the two stimuli, to minimize the
summed activity, *a* must be positive and small. So the response to the deviant needs to be
larger than that of the standard. However, the relative weightage of *MI* and the summed
activity (ρ) can allow *a* to be increasing or decreasing and still cause CI_k to be increasing.

We show that the condition for which CI_{total} keeps increasing by considering $\Delta CI = CI_n - CI_m$

($n>m$) with the assumption that the sequence $\{a_k\}$ is monotonic is

$$\Delta CI = CI_n - CI_m = x\left[\frac{a_n{}^2 - a_m{}^2}{2\sigma^2} + \Lambda(a_n) - \Lambda(a_m) - \rho\{a_n - a_m\}\right]$$

669
$$= x(a_n - a_m)\left[\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} - \rho\right] \ldots (2)$$

From the last expression in Eq. 2, under 2 different ranges of ρ, ΔCI is increasing with *an*

increasing or *a* decreasing respectively. The theoretical analysis suggests that before and after

ECO, the relative importance of coding or discrimination and minimal activity is changed. Based

on the above, sparseness in later stages (after ECO), when spike rates are higher, needs to be

weighted more than earlier stages (before ECO) to explain the differential effect observed

before and after ECO in response to the low probability stimulus.

Discussion

Earliest sound driven activity in the mouse auditory pathway

We show that the mouse auditory pathway, ACX and IC, is driven by sounds with ear canals

closed as early as P7 and down to intensities of 60 dB SPL. The above result is perhaps not

surprising as auditory brainstem responses (ABR) in mice have been shown before ECO[23] and

that there are long term changes in prestin expression in mice when exposed to loud sounds at

these early ages[23] (P6-P11). Studies based on sound-evoked[23] and spontaneous activity[48] show

that the central auditory pathway is functional before ECO. However, the peripheral auditory

system may be immature at those ages. The external auditory meatus in mice reopens at P7,

and the outer ear is fully mature by that time[49]. Further, lever arm lengths of middle ear bones

are at 70% of mature values but have normal lever ratios by P7[50]. The oval window is also of

normal size by P7[50]. Most importantly, mechano-transduction currents from outer hair cells

(OHC) and inner HCs (IHC) become maximal in the cochlear base by P2 and at the apex by

P5[51,52], and thus IHCs are fully mature by P5[56]. However, OHCs are immature at P5 with

nonlinear capacitance (NC) of OHCs starting at P7 and maturing fully at P18[53]. The development

of OHC NC from P7 is correlated with the expression of prestin, the motor protein responsible

for the OHC NC[53] and hence hearing sensitivity. Thus, barring OHC NC, the key to low threshold

(0 dB SPL) hearing and high auditory nerve fibre sensitivity, all evidence suggests that the

peripheral auditory system is mature. Thus moderate to high-intensity sound-evoked activity in

the auditory pathway is possible at ages before ECO in mice. However, no study to the best of

our knowledge has shown auditory responses in mice before ECO. So far, all studies considering

sound-evoked activity and early critical period in the auditory system have considered hearing
onset to coincide with ECO. Thus our results open up an earlier time window of activity-driven
plasticity to occur in the auditory pathway. Previous work on ferrets[18] also shows such a
possibility, where the authors show that SPNs are the first to respond to sounds at an
equivalent developmental stage of ferrets as we find in mice.

Auditory development and deviant detection

Post-ECO development of basic response properties like tuning, tuning width, thresholds to
tonal sounds or broadband noise and other basic stimuli have been studied[5,54]. Our study not
only shows that the starting point of auditory activity driven development is before ECO but
also provides data on how deviant detection changes over development into adulthood. We
show that the well-known deviant detection property of the auditory pathway[35,42] is present in
the early stages and is stronger in cortical neurons than the thalamus. We also show that
deviant detection is strongest at early ages and reduces to the values observed in the adult.
More importantly, we find that SPNs are deviant selective at the earliest stages of sound driven
activity, before ECO, while cortical L4 neuron populations are not deviant selective at those
ages. Since SSA underlying deviant selectivity is primarily a network phenomenon[42], it may be
surprising that SPNs are deviant selective at ages when thalamocortical and cortico-cortical
connections are forming. However, it is well known that SPNs form synapses with each
other[11,55,56]. Correlated with our observations of higher oddball selectivity in SPNs, SPNs are
found to have more recurrent connections before ECO than after ECO and more than that of L4
neurons before ECO. After ECO, L4 neurons have more recurrent connections than before. The
differential nature of recurrent connectivity in L4 and SPN before and after ECO also
corroborates with our observations of differential oddball selectivity in the two populations,
before and after ECO. Thus the nature of recurrent connectivity change we observe during early
development supports the changes in the strength of deviant detection as measured by CSIs.

Early auditory experience-based plasticity, SPNs and critical period

Broad tonotopic organization in the primary auditory cortex (A1)[57,58] is known to be altered
with over-representation of a particular frequency in the adult rats/mice which were exposed
to that frequency tone during the critical period (P12 to P15, after ECO)[5,7]. In light of our
present results with auditory responses present before the above period, we revisit exposure to
a single frequency tone before the critical period. The previous studies[5,7] did not find any
evidence of adult cortical over-representation of the exposure frequency when exposing mice
to that frequency before ECO. Thus, the window of P12-15 in mice and a similar window in rats

was defined as the critical period[5,7]. Our data explain the lack of over-representation with
exposure before ECO despite the presence of auditory responses during that period. The SPNs
driving thalamocortical synapses to mature are found to have deviant detection at these early
stages and thus SPNs, and consequently, L4 neurons do not respond to the repeated tone
presentations. Therefore in our SS exposure results before ECO (P6-11, EX6) also, we do not
find any plastic strengthening of responses to the S frequency as expected from above.
However, the unique deviant detection properties of the SPNs before ECO, as hypothesized
show strengthening of response to the deviant tone with an SD exposure before ECO. It is
expected due to the lack of or weak responses to the repeating S frequency and responses to
the D frequency. However, SD exposure following ECO shows the opposite since the SPNs are
less deviant selective and thalamic inputs are now stronger, driving L4 responses directly.
Similarly, the SS exposure after ECO also shows expected results. The mechanism underlying
our observations is summarized in our model results. Of course, adaptation time constants play
a role in the observed deviant detection and hence our exposure results.

Since SSA is primarily a network phenomenon with recurrent connections playing a key role,
the observed changes in recurrent connections cannot be ruled out in understanding our
results. In our current model, we did not consider the recurrent connections. Extension of our
minimal binary network model to a recurrent network is expected to also produce the same
results as the effects of deviant detection and SSA, underlying our observations would remain
unchanged. Our model is only a minimal neural structure capturing the key mechanisms
explaining the observations. Further, the model currently does not include inhibition, an
element crucial in development, as synaptic plasticity rules of inputs/outputs of inhibitory
neurons at such ages are not available. It likely plays a pivotal role as the maturation of
inhibition is controlled by SPNs[14]. Also, the effect of the inhibitory switch, occurring before ECO,
on the activity driven plasticity observed before ECO, needs to be investigated. The closure of
critical period, around P16 also coincides with the observed near absence of SPNs in mice from
P15. Thus mechanisms implicated in critical period closure like inhibition, adenosine, cholinergic
inputs[6,59–61], could also be tied to SPNs, since SPNs guide development of subcortical
projections. Finally, exposures with selective inhibition of SPN activity during relevant periods
of development are required to establish the SPN driven cortical development further.

We thus suggest that to understand experience-dependent developmental plasticity better, the
period before ECO needs to be considered and possibly revise timelines of the critical period.
The concept of the auditory critical period has been modified by showing that it can be
stimulus-specific[62]. Similarly, we suggest the presence of a critical period that is deviant
selective before ECO.

Implications of the SPN deviant detection critical period

We show in mice that relatively low probability salient stimuli before ECO at ages equivalent to
human gestational week 25[21] cause long term plasticity. The above potentially explains
observations in humans of maternal voice recognition in infants[4,63]. The results also suggest
that prenatal sensory environment can influence cortical development and may also generalize
to other sensory systems given the unique response properties of the SPNs. The generalization
to other systems or other stimuli, in this case, is supported by our theoretical predictions.
Although our results are derived based on tones as stimuli, the possibility of coding of auditory
objects by the neurons in the ACX can also be addressed through similar experiments. For
example, specific response selectivity to species-specific vocalizations in mice and other species
could also be explained based on the low probability salient occurrence of such sounds during
early development. Corresponding experiments for other sensory systems can lead to a better
understanding of cortical development.

Methods

All animal experiments were approved by the Institutional Animal Ethics Committee
(IAEC) of Indian Institute of Technology Kharagpur. Detailed methods are provided in the
Supplementary Materials. Thy1-Gcamp6f (Jax Labs) or C57BL6/J strain mice (NIN, Hyderabad,
India or Jax Labs) across different developmental time windows were deeply anaesthetized for
surgery (2-2.5% isoflurane), and prepared for in-vivo recordings after making a craniotomy (3-4
mm diameter, based on stereotaxic coordinates and landmarks). To investigate the presence of
response before ECO, widefield and 2-Photon Ca^{+2} imaging was performed on Thy1-Gcamp6f
expressing animal across different developmental window starting at ages before the ear canals
were opened (P7 onwards). Few awake recordings were performed to probe responses at
modest sound levels (70dB SPL). For in-vivo electrophysiological recordings from A1/AAF or
MGB, 16 channel electrodes (Microprobes) were lowered in SP and L4 of ACX using a
micromanipulator inside a sound chamber (IAC Acoustics). LFP and spiking responses to sounds
(played through TDT free-field speakers ES1, attenuated by PA5 and generated with TDT RX6)
were obtained with a unity gain headstage and preamps (Plexon Inc) and stored using National
Instruments DAQ card. Sound generation, data acquisition and online/offline analysis were
done with custom-written MATLAB routines. Anatomical labelling and confirmation of
recordings from A1/AAF and from L4 or SP were done using multiple approaches (lipophilic
tracer injections in recording site, Nissil sections and IHC). Exposure in specific auditory
environments of mouse litters for the different time windows was done in another sound
chamber (IAC Acoustics) maintaining an 8hr-16hr light-dark cycle. A computational model based
on synaptic plasticity and fee-forward synaptic depression was added to demonstrate the high

oddball selectivity before ECO. A mathematical commentary based on mutual information

maximization was further done to provide theoretical support for our findings.

References

1. Schreiner, C. E. & Polley, D. B. Auditory map plasticity: Diversity in causes and consequences. *Current Opinion in Neurobiology* vol. 24 143–156 (2014).

2. Hensch, T. K. Critical period plasticity in local cortical circuits. *Nature Reviews Neuroscience* (2005) doi:10.1038/nrn1787.

3. Decasper, A. J. & Fifer, W. P. Of human bonding: Newborns prefer their mothers' voices. *Science (80-.).* (1980) doi:10.1126/science.7375928.

4. Voegtline, K. M., Costigan, K. A., Pater, H. A. & DiPietro, J. A. Near-term fetal response to maternal spoken voice. *Infant Behav. Dev.* (2013) doi:10.1016/j.infbeh.2013.05.002.

5. de Villers-Sidani, E., Chang, E. F., Bao, S. & Merzenich, M. M. Critical Period Window for Spectral Tuning Defined in the Primary Auditory Cortex (A1) in the Rat. *J. Neurosci.* (2007) doi:10.1523/JNEUROSCI.3227-06.2007.

6. Takesian, A. E., Bogart, L. J., Lichtman, J. W. & Hensch, T. K. Inhibitory circuit gating of auditory critical-period plasticity. *Nat. Neurosci.* (2018) doi:10.1038/s41593-017-0064-2.

7. Barkat, T. R., Polley, D. B. & Hensch, T. K. A critical period for auditory thalamocortical connectivity. *Nat. Neurosci.* (2011) doi:10.1038/nn.2882.

8. Zhou, X. & Merzenich, M. M. Enduring effects of early structured noise exposure on temporal modulation in the primary auditory cortex. *Proc. Natl. Acad. Sci.* **105**, 4423–4428 (2008).

9. Zhang, L. I., Bao, S. & Merzenich, M. M. Persistent and specific influences of early acoustic environments on primary auditory cortex. *Nat. Neurosci.* **4**, 1123–1130 (2001).

10. Luhmann, H. J., Kirischuk, S. & Kilb, W. The Superior Function of the Subplate in Early Neocortical Development. *Front. Neuroanat.* **12**, (2018).

11. Kanold, P. O. & Luhmann, H. J. The Subplate and Early Cortical Circuits. *Annu. Rev. Neurosci.* (2010) doi:10.1146/annurev-neuro-060909-153244.

12. Hoerder-Suabedissen, A. & Molnár, Z. Molecular diversity of early-born subplate neurons. *Cereb. Cortex* **23**, 1473–1483 (2013).

13. Kanold, P. O., Kara, P., Reid, R. C. & Shatz, C. J. Role of subplate neurons in functional maturation of visual cortical columns. *Science (80-.).* (2003) doi:10.1126/science.1084152.

14. Kanold, P. O. & Shatz, C. J. Subplate Neurons Regulate Maturation of Cortical Inhibition and Outcome of Ocular Dominance Plasticity. *Neuron* (2006) doi:10.1016/j.neuron.2006.07.008.

15. Ghosh, A. & Shatz, C. J. Involvement of subplate neurons in the formation of ocular
dominance columns. *Science (80-.).* (1992) doi:10.1126/science.1542795.

16. Ghosh, A., Antonini, A., McConnell, S. K. & Shatz, C. J. Requirement for subplate neurons
in the formation of thalamocortical connections. *Nature* (1990) doi:10.1038/347179a0.

17. McConnell, S. K., Ghosh, A. & Shatz, C. J. Subplate neurons pioneer the first axon
pathway from the cerebral cortex. *Science (80-.).* (1989) doi:10.1126/science.2475909.

18. Wess, J. M., Isaiah, A., Watkins, P. V & Kanold, P. O. Subplate neurons are the first
cortical neurons to respond to sensory stimuli. *Proc. Natl. Acad. Sci.* (2017)
doi:10.1073/pnas.1710793114.

19. Zhao, C., Kao, J. P. Y. & Kanold, P. O. Functional Excitatory Microcircuits in Neonatal
Cortex Connect Thalamus and Layer 4. *J. Neurosci.* (2009) doi:10.1037/pha0000116.

20. Hanganu, I. L., Kilb, W. & Luhmann, H. J. Functional synaptic projections onto subplate
neurons in neonatal rat somatosensory cortex. *J. Neurosci.* (2002) doi:20026716.

21. Clancy, B., Darlington, R. B. & Finlay, B. L. Translating developmental time across
mammalian species. *Neuroscience* (2001) doi:10.1016/S0306-4522(01)00171-3.

22. Polley, D. B., Thompson, J. H. & Guo, W. Brief hearing loss disrupts binaural integration
during two early critical periods of auditory cortex development. *Nat. Commun.* (2013)
doi:10.1038/ncomms3547.

23. Chang, A. *et al.* Specific Influences of Early Acoustic Environments on Cochlear Hair Cells
in Postnatal Mice. *Neural Plast.* (2018) doi:10.1155/2018/5616930.

24. Meng, X. *et al.* Transient Subgranular Hyperconnectivity to L2/3 and Enhanced Pairwise
Correlations during the Critical Period in the Mouse Auditory Cortex. *Cereb. Cortex* (2020)
doi:10.1093/cercor/bhz213.

25. Sohmer, H., Perez, R., Sichel, J. Y., Priner, R. & Freeman, S. The pathway enabling external
sounds to reach and excite the fetal inner ear. *Audiol. Neuro-Otology* (2001)
doi:10.1159/000046817.

26. Partanen, E., Kujala, T., Tervaniemi, M. & Huotilainen, M. Prenatal music exposure
induces long-term neural effects. *PLoS One* (2013) doi:10.1371/journal.pone.0078946.

27. Hensch, T. K. Critical period plasticity in local cortical circuits. *Nature Reviews
Neuroscience* (2005) doi:10.1038/nrn1787.

28. Stiebler, I., Neulist, R., Fichtel, I. & Ehret, G. The auditory cortex of the house mouse:
Left-right differences, tonotopic organization and quantitative analysis of frequency
representation. *J. Comp. Physiol. - A Sensory, Neural, Behav. Physiol.* (1997)
doi:10.1007/s003590050140.

29. Blankenship, A. G. & Feller, M. B. Mechanisms underlying spontaneous patterned activity
in developing neural circuits. *Nature Reviews Neuroscience* (2010) doi:10.1038/nrn2759.

30. Yildirim, M., Sugihara, H., So, P. T. C. & Sur, M. Functional imaging of visual cortical layers
and subplate in awake mice with optimized three-photon microscopy. *Nat. Commun.*
(2019) doi:10.1038/s41467-018-08179-6.

31. Osheroff, H. & Hatten, M. E. Gene expression profiling of preplate neurons destined for
the subplate: Genes involved in transcription, axon extension, neurotransmitter
regulation, steroid hormone signaling, and neuronal survival. *Cereb. Cortex* (2009)
doi:10.1093/cercor/bhp034.

32. Hoerder-Suabedissen, A. *et al.* Novel markers reveal subpopulations of subplate neurons
in the murine cerebral cortex. *Cereb. Cortex* (2009) doi:10.1093/cercor/bhn195.

33. Viswanathan, S., Bandyopadhyay, S., Kao, J. P. Y. & Kanold, P. O. Changing Microcircuits
in the Subplate of the Developing Cortex. *J. Neurosci.* **32**, 1589–1601 (2012).

34. Wang, W. Z. *et al.* Subplate in the developing cortex of mouse and human. *J. Anat.* (2010)
doi:10.1111/j.1469-7580.2010.01274.x.

35. Ulanovsky, N., Las, L. & Nelken, I. Processing of low-probability sounds by cortical
neurons. *Nat. Neurosci.* (2003) doi:10.1038/nn1032.

36. Malmierca, M. Neuronal adaptation, novelty detection and regularity encoding in
audition. *Front. Syst. Neurosci.* (2014) doi:10.3389/fnsys.2014.00111.

37. Pérez-González, D. & Malmierca, M. S. Adaptation in the auditory system: an overview.
Front. Integr. Neurosci. (2014) doi:10.3389/fnint.2014.00019.

38. Szymanski, F. D., Garcia-Lazaro, J. A. & Schnupp, J. W. H. Current source density profiles
of stimulus-specific adaptation in rat auditory cortex. *J. Neurophysiol.* (2009)
doi:10.1152/jn.00240.2009.

39. Holy, T. E. & Guo, Z. Ultrasonic songs of male mice. *PLoS Biol.* **3**, 1–10 (2005).

40. Parras, G. G. *et al.* Neurons along the auditory pathway exhibit a hierarchical
organization of prediction error. *Nat. Commun.* (2017) doi:10.1038/s41467-017-02038-6.

41. Antunes, F. M., Nelken, I., Covey, E. & Malmierca, M. S. Stimulus-specific adaptation in
the auditory thalamus of the anesthetized rat. *PLoS One* (2010)
doi:10.1371/journal.pone.0014071.

42. Yarden, T. S. & Nelken, I. Stimulus-specific adaptation in a recurrent network model of
primary auditory cortex. *PLoS Comput. Biol.* (2017) doi:10.1371/journal.pcbi.1005437.

43. Granger, C. W. J. Investigating Causal Relations by Econometric Models and Cross-
spectral Methods. *Econometrica* (1969) doi:10.2307/1912791.

44. Francis, N. A. *et al.* Small Networks Encode Decision-Making in Primary Auditory Cortex.
Neuron (2018) doi:10.1016/j.neuron.2018.01.019.

45. Mill, R., Coath, M., Wennekers, T. & Denham, S. L. A neurocomputational model of
stimulus-specific adaptation to oddball and markov sequences. *PLoS Comput. Biol.* (2011)
doi:10.1371/journal.pcbi.1002117.

46. Hromádka, T., DeWeese, M. R. & Zador, A. M. Sparse representation of sounds in the
unanesthetized auditory cortex. *PLoS Biol.* (2008) doi:10.1371/journal.pbio.0060016.

47. Smith, E. C. & Lewicki, M. S. Efficient auditory coding. *Nature* (2006)
doi:10.1038/nature04485.

48. Babola, T. A. *et al.* Homeostatic Control of Spontaneous Activity in the Developing
Auditory System. *Neuron* (2018) doi:10.1016/j.neuron.2018.07.004.

49. Anthwal, N. & Thompson, H. The development of the mammalian outer and middle ear.
J. Anat. (2016) doi:10.1111/joa.12344.

50. Huangfu, M. & Saunders, J. C. Auditory development in the mouse: Structural maturation
of the middle ear. *J. Morphol.* (1983) doi:10.1002/jmor.1051760302.

51. Kim, K. X. & Fettiplace, R. Developmental changes in the cochlear hair cell
mechanotransducer channel and their regulation by transmembrane channel-like
proteins. *J. Gen. Physiol.* (2013) doi:10.1085/jgp.201210913.

52. Lelli, A., Asai, Y., Forge, A., Holt, J. R. & Géléoc, G. S. G. Tonotopic gradient in the
developmental acquisition of sensory transduction in outer hair cells of the mouse
cochlea. *J. Neurophysiol.* (2009) doi:10.1152/jn.00136.2009.

53. Abe, T. *et al.* Developmental expression of the outer hair cell motor prestin in the mouse.
J. Membr. Biol. (2007) doi:10.1007/s00232-007-9004-5.

54. Chang, E. F., Bao, S., Imaizumi, K., Schreiner, C. E. & Merzenich, M. M. Development of
spectral and temporal response selectivity in the auditory cortex. *Proc. Natl. Acad. Sci. U.*
S. A. (2005) doi:10.1073/pnas.0508239102.

55. Hirsch, S. & Luhmann, H. J. Pathway-specificity in N-methyl-d-aspartate receptor-
mediated synaptic inputs onto subplate neurons. *Neuroscience* (2008)
doi:10.1016/j.neuroscience.2008.01.068.

56. Kanold, P. Subplate neurons: crucial regulators of cortical development and plasticity.
Front. Neuroanat. (2009) doi:10.3389/neuro.05.016.2009.

57. Bandyopadhyay, S., Shamma, S. A. & Kanold, P. O. Dichotomy of functional organization
in the mouse auditory cortex. *Nat. Neurosci.* (2010) doi:10.1038/nn.2490.

58. Issa, J. B. *et al.* Multiscale Optical Ca2+Imaging of Tonal Organization in Mouse Auditory

Cortex. *Neuron* (2014) doi:10.1016/j.neuron.2014.07.009.

59.   Courtney E. Yaeger, Dario Ringach, J. T. T. Neuromodulatory control of localized dendritic
spiking in critical period cortex. *Nature* **567**, 100–104 (2019).

60.   Blundon, J. A. *et al.* Restoring auditory cortex plasticity in adult mice by restricting
thalamic adenosine signaling. *Science (80-.).* (2017) doi:10.1126/science.aaf4612.

61.   Vickers, E. D. *et al.* Parvalbumin-Interneuron Output Synapses Show Spike-Timing-
Dependent Plasticity that Contributes to Auditory Map Remodeling. *Neuron* (2018)
doi:10.1016/j.neuron.2018.07.018.

62.   Bhumika, S. *et al.* A Late Critical Period for Frequency Modulated Sweeps in the Mouse
Auditory System. *Cereb. Cortex* (2020) doi:10.1093/cercor/bhz262.

63.   Mehler, J. *et al.* A precursor of language acquisition in young infants. *Cognition* (1988)
doi:10.1016/0010-0277(88)90035-2.

Supplementary Material

Detailed Methods

Subjects

All animal experiments were approved by the Institutional Animal Ethics Committee
(IAEC) of Indian Institute of Technology Kharagpur.

Acute craniotomy for wide-field imaging and 2-photon Ca^{2+} imaging

A standard surgical procedure was performed as mentioned above in in-vivo electrophysiology
recordings. Animals were initially anaesthetized under isoflurane (induction at 5% isoflurane,
followed by 2% isoflurane). After removing the skin and tissue above the skull, a metallic head
plate was implanted above the estimated auditory cortical area. Craniotomy of around 3mm
diameter was performed over the left auditory area. Smaller size craniotomy was preferred, as
it gave us the advantage to minimize brain motion. A craniotomy was filled with 1.5% low
melting-point Agarose. Immediately after that, a 5mm coverslip was imbedded over the
craniotomy to dampen the pulsation. Animals were then transferred to the 2-photon imaging
setup, and the level of anaesthesia was reduced down to 0.1-0.2% isoflurane during the
recordings. For recording responses at lower sound levels, awake recordings were performed at
age P9-P10. For awake recordings, Immediately after completing the surgery, pups were
transferred to the sound chamber and allow them to recover from anaesthesia. This time
window also allowed pups to habituate before the recordings were performed. Throughout the
experiment, the animal temperature was maintained at 37°C by a heating pad.

Wide-Field Imaging

For wide-field calcium imaging, imaging signals were acquired using a 14-bit CoolSNAP
HQ2 CCD camera (Photometrics). Images of surface vasculature were acquired using blue LED
illumination (470 nm), and wide-field GCaMP6f signals were recorded at ~30 Hz using blue
illumination (470 nm)[1]. Fluorescence signals were acquired at 4x binning of 1040x1392 pixels,
using a 4X objective (Olympus). Based on the auditory landmarks (Medial cerebral artery (MCA)
and rhinal vein position) and the responses to different frequencies, the region of interest and
the focal plane of the image (150-300 µm) were moved to desired coordinates for performing
2-photon imaging.

1022

1023

1024

Two-Photon Calcium Imaging

Two-photon calcium imaging was performed using a commercially available 2-photon microscopy system (Prairie View Technologies). The microscopy was controlled by Prairie View software (Prairie View Technologies). For imaging, the excitation beam of wavelength 860nm generated from Insight laser (Spectra-Physics, USA) was directed over the exposed brain volume. The laser was delivered through 20X/0.8 NA water immersion objective (2mm WD, Olympus). The laser power was adjusted from 50mW to 80mW depending upon the condition of the specimen. Frames in the region of interests (~150 μm x 200 μm with 1.16 μm pixel size) were imaged at ~4-6 Hz (160-250ms frame period, 1.2-4 μs dwell time) with the stimulus presented at usually the 6th (to the 10th) frame in a sequence of 20-30 frames per stimulus.

Auditory stimulation for imaging

The stimulus was presented through the electrostatic speaker (ES1) placed 2cm away from the right ear of the mouse pups Stimulus was generated through custom-written software in MATLAB (Mathworks), passed through TDT RZ6 multifunctional processor. Acoustic calibrations were performed using microphone 4939 (Brüel & Kjær, Denmark) and showed a typical flat (+/- 7 dB) calibration curve from 4-60 kHz. Auditory stimuli consisted of broadband noise (6-48 kHz, 40 repetitions), tones (40 repetitions of respective frequencies), and series of a sound token representing SD paradigm (30 repetitions). Only for one of the animals (P12, after ear canal opening), 5 repetitions of each pure frequency tone was delivered. Each of the stimuli was interleaved with a gap of 5s. The loudest intensities of sounds at 80-90 dB SPL were presented to ear, to allow mechanotransduction at younger ages (before ear canal opening). However, in awake recordings, 50-90 dB of sounds were presented at age P9-P10.

Single units spike times were obtained from the acquired spike channel data using threshold crossing and spike sorting with custom-written software in MATLAB.

Anatomical labelling

Nissl staining was performed at different age groups to identify the SP layer and to guide in-vivo electrophysiology experiments. Mice of ages P6 (n=2 mice), P13 (n=2 mice), P16 (n=2 mice) and P18 (n=2 mice) were initially perfused, and the brains were harvested. Since ear canal opening is after P11 and as the cortical size stabilizes by P21, the above ages were chosen initially for the characterization. Nissl stains were performed after obtaining coronal sections of the auditory regions based on mouse atlas[2]. Immunohistochemistry was also performed on brain slices across different developmental time points (P06-P07, n=2 mice; P08-P11, n=3 mice, P12-13, n=5mice; P14-P16, n=2mice; P17-P19, n=2 mice; P20-28, N=3 mice; >P28, n=2 mice) with known markers of SP which are Complexin 3 (Cplx3); Synaptic Systems (122302) and Connective Tissue-Derived Growth Factor (CTGF); Santa Cruz (sc-14939). Paraformaldehyde

(PFA) fixed brain tissue was embedded in a 3% agarose block. Brain embedded block was fixed
on a vibratome (Leica) sectioning platform. 50 micron thin sections of auditory slices were cut,
blocked and permeabilized in a buffer containing, 1% BSA, 0.3% Triton X and 10% Normal
Donkey Serum (for CTGF) or 10% Normal Goat Serum (for Cplx3). Sections were incubated in
primary antibodies (Cplx3: 1:500 and CTGF: 1:500) overnight and followed by washing with 1X
PBS five times. After that, sections were incubated for two hours in Secondary antibodies
(Cplx3: anti-rabbit 594, Invitrogen and CTGF: anti-goat 488, Invitrogen) diluted in the buffer, 1%
BSA, 3% serum and 0.3% Triton X at room temperature. Sections were rinsed, mounted and
imaged under epi-fluorescence microscopy. Depth of SP in different regions and at different
developmental time scales was determined from these initial studies and was used as a guide
for performing electrophysiology recordings. Further confirmation about recordings from input
layer (Layer IV) was also obtained from SCNN1-td-tomato (cross of SCNN1-cre and floxed-td-
tomato: JAX mice) mice, which have SCNN1 expressing neurons (specific to Layer IV) labelled,
allowing confirmation of recordings in Layer IV. Sections were imaged using epi-fluorescence
microscopy (Xcite illumination, Bruker), and were analyzed for cytoarchitecture by using
custom-written codes in MATLAB.

In-vivo extracellular single-unit recordings

Initially, animals were anaesthetized under deep anaesthesia (5% Isoflurane) inside the
induction chamber. Initial Induction step helps in maintaining anaesthesia level in animals even
after bringing down the anaesthesia levels (1.5-2%) during surgery. Core body temperature was
continuously maintained at 37^{o}C with a heating pad throughout the experiment. A central
incision was made, and the left temporal portion of the skull was exposed. Exposed left
temporalis was cleaned with the application of alcohol. After that, tissue clearance was done by
rubbing the skull with 3% of Hydrogen peroxide. A 5mm diameter region was marked on the
left hemisphere. Skull was firmly fixed after keeping the mark at the centre of a head plate.
Craniotomy was performed over the estimated auditory cortical area. After completing a
craniotomy, the experimental animal was shifted immediately to a soundproof chamber (IAC
Acoustics). The depth of anaesthesia was continuously monitored throughout the experiment
with paw pinches. During recording sessions, anaesthesia was brought down to a concentration
of 0.5-1%, to get robust responses from primary auditory cortex.

Extracellular recordings were performed from the left auditory cortex using tungsten
Micro Electrodes Array (MEA) of impedance 3-5 Mega ohms (MicroProbes, USA). A 4X4
custom-designed metal MEA with an inter-electrode spacing of 125 microns was lowered into
the auditory cortex using a micromanipulator (MP-285, Sutter Instrument Company, Novato,
CA). We also performed recording using arrays separated by 400 or 600 microns in length to
record across laminae simultaneously (for example, Layer IV and SP, Fig. 3A). Signals were

acquired after passing through a unity gain headstage (Plexon, HST16o25) and followed by
PBX3 (Plexon) preamp with a gain of X1000. Wideband signal (used to extract LFP, 0.7 Hz to 6
kHz) and spike signals (150 Hz to 8 kHz) were obtained in parallel and acquired through National
Instruments Data Acquisition Card (NI-PCI-6259) at 20 kHz sampling rate, controlled through
custom-written MATLAB (Mathworks, Natick, USA) routines. Further, all online/offline analysis
was performed using custom-written MATLAB routines. A laminar profile of responses was
obtained from the primary auditory cortex (A1) by advancing the electrodes in depth at 100-
150 micron steps.

Since the size and location of A1 over-development varies, it is essential to confirm the
location of recording to be A1. For A1 recordings additional physiological information based on
tuning – the existence of broad tonotopy, across electrodes in the recording array, latency of
responses, selectivity to tone stimuli was used to confirm recordings in primary auditory areas
(A1 and Anterior Auditory Field (AAF)). Based on the location of electrode tracks and
comparison with mouse brain atlas[70] and landmarks (location relative to the rhinal vein, size of
the hippocampus in coronal slices) also provides additional confirmation of recording location.
In many experiments after recording a DiI crystal[3], a lipophilic tracer was inserted into the
cortex at the recording location. After allowing passive travel of the tracer for 20-30 days at
35°C, brain slices were cut to look for labelled fibres and cell bodies in the ventral division of
the MGB (lemniscal structure projecting to A1) allowing confirmation of recordings in A1 Fig.
S1A). We further confirmed recordings from ACX by electrode location (using electrode tracks)
in primary regions and also from broad tonotopy observed (Fig. S1B). The recording depths
targeted for L4 and SPN were based on known depths of L4 and SPNs in ACX[34]; also, in a
separate group of mice (P8-P11, n=3, P12-P13, n=7; P14-P16, n=5) depth of SPNs was
determined to be above 800 μm based on two methods. First by cell morphology in Nissl stains
of coronal slices (Fig. S1C) and second by immunostaining coronal slices containing A1 with
antibodies of SPN markers[32,33], *Complexin3* and *CTGF*, at different ages (Fig. S1E) and depth
range of L4 from the expression of *SCNN1* (B6.Cg-Tg(Scnn1a-cre)3Aibs/J mouse, Jax Labs, 09613
crossed with CAG-tdTomato mouse Jax Labs, 07909), an L4 specific marker[34]. In a subset of
mice, we confirmed electrode recording site to be in SP layer from electrode tracks along with
immunostaining as above (Fig. S1E) and electrodes to be in L4 by performing recordings in the
SCNN1-cre-td-Tomato (n=3, P12, P14 and P18) mice (Fig. S1D). Depths of other layers (L2/3 and
L5/6) were determined based on the above results.

Auditory stimulation for in-vivo electrophysiology

The stimulus was presented inside the soundproof chamber, 10 cm away from the right ear
(contralateral) of the mouse, with TDT electrostatic speakers (ES1) driven by TDT drivers ED1
after attenuation by TDT attenuators (PA5) generated through TDT RX6 using custom made

software written in MATLAB. The acoustic calibrations, performed with microphone 4939 (Brüel
& Kjær, Denmark), of the ES1 speakers (TDT) in the sound chamber, showed a typical flat (+/- 7
dB) calibration curve from 4-60 kHz.

1135

Exposure protocol

For another set of experiment, animal litters were exposed to sound sequences inside a
soundproof chamber with maintained 8-hr light and 16hr dark cycle, continuously for 5 days in
4 age groups (P0-P5/6, P6-P10/11, P11-P15/16 and P16-P21).

The litters were kept along with the breeders in a cage. Each exposure stimulus was synthesized
online and presented through an electrostatic speaker, kept 5cm above the cage, using custom-
written codes on MATLAB. All the age groups were presented with a sequence of stimuli having
50 ms sound tokens with a 250 ms inter-token gap at 70 and 90 dB SPL. The sequence (SD) had
a token of two frequencies, a standard (S) having an arrival probability of 90% and a deviant (D)
arriving with 10% probability. For only standard exposure (SS), there was no D frequency. After
sound exposure, animals were returned immediately to home cages in the animal house and
reared up in a normal environment until experiments were performed. During recording
sessions, mapping of the primary auditory cortex was performed. Tone pips (50 ms, 5 ms up
down ramps, 6-48 kHz, 0.25 octaves difference at 10-50 dB above threshold, 10 dB increments,
5 repetitions of each frequency-Intensity combination) was presented to the right ear.
Extracellular responses were recorded from thalamo-recipient layer (Deep Layer III and layer IV)
at depth 350-500 microns. Frequency responses area (FRA) was constructed of each unit for a
minimum of three sound levels played. Single units from A1 having response peak latency
within 9 ms to 31 ms were included in the analysis. This window was selected based on
population mean and variance of response peak latencies obtained across all sound levels.

1156

1157 **Data Analysis**
1158

Wide Field imaging analysis (Fig. 1)

Wide Field imaging analysis was performed using custom codes written in MATLAB
(Mathworks). For analysis of the wide-field data, raw images collected were binned at 4x and
then smoothed with a moving window of 4x4. Further analysis was done on the smoothed
images. To find the significant pixels, a one-sided t-test across all the iterations was performed
between the 5 frames immediately preceding the stimulus onset and a moving window of 5

frames after the stimulus onset in all 130 µm by 130 µm square areas. The pixels which gave a p-value of less than 0.05 on at least one window after the stimulus onset, were considered for further analysis. The mean change in df/f with time traces was constructed for pixels showing significant positive responses. These traces were further smoothed with a moving average window of 5 frames, and a one-sided t-test was performed again to find the first significant post-stimulus frame and the frame with the most significant response.

Table S1

Sl.no.	Age	Tonal stimuli played (~90dB SPL, in kHz)	Significant responsive frequencies and respective % responding areas	Fraction of frequencies played showed responses (%)
1.	P-7	8.5,12,17,24,48	8.5kHz: 2.86%	20.00%
2.	P-8	6,8.5,12,17,24,34,48	6kHz: 0.56%; 8.5kHz: 3.67%	28.57%
3.	P-9	6,8.5,12,17,24,34,48	6kHz: 26.9%; 17kHz: 3.44%; 24kHz: 0.56%; 48kHz: 1.31%	57.14%
4.	P-10	6,8.5,12,17,24,34,48	8.5kHz: 37.12%; 12kHz: 4.38%; 17kHz: 0.5%; 48kHz: 19.93%	57.14%
5.	P-13	6,8.5,12,17,24,34,48	6kHz: 43.8%; 8.5kHz: 79.8%; 12kHz: 13.54%; 17kHz: 70.8% 24kHz: 22.9%	71.42%

2-photon imaging analysis (Fig. 2)

2-photon imaging analysis was performed using custom codes written in MATLAB (Mathworks).
Imaging sequences were aligned by performing X-Y drift correction. Cells were selected
manually by selecting the centre point of the cell on the motion-corrected mean image. ROI (5
µm radius) were drawn based on the cell centres. Raw fluorescence signal over time (F) of the
selected ROIs across all frames were extracted. For each trial, relative fluorescence was
computed by using $\Delta F/F_0 = (F - F_0)/ F_0$), where F_0 corresponds to baseline fluorescence. Baseline
fluorescence amplitude was estimated by calculating the mean of fluorescence values over all
the frames preceding stimulus, except the first 2 frames. Neurons on stimulation to tones and
noise were considered to be significantly responsive as follows. One-sided paired t-tests were
performed between mean df/f over 3 of the successive frames before the stimulus and each
one of 6 windows, of 3 consecutive frames after stimulus onset (up to the 8[th] frame, ~1.5-1.8s).
If any of the 6 moving windows showed a significant increase in mean df/f (40 repetitions) over
baseline, the neuron was considered responsive.

Table S2.

Sl.no.	Age	Fraction of cases (all neurons in response to different frequency tone at 0 dB attenuation) that showed significant responses.
1.	P-7	6.84%
2.	P-8	5.6%
3.	P-9	10.6%
4.	P-10	15.4%
5.	P-11	12.3%
6.	P12-P13	12.4%

1192

1193

1194

1195   Table S3.

Sl.no.	Tone Intensity	Fraction of cases (all neurons in response to different intensities of tones) that showed significant responses.
1.	90 dB	40.50%
2.	80 dB	12.77%
3.	70 dB	33.04%
4	60 dB	7.02%

1196

For significant responses in single trials to compute reliability, a trial was deemed responsive if
the mean df/f of 4 frames (~1s) after the stimulus after stimulus onset was higher than 1*STD
(STD obtained from 40 repetitions) of the mean of the 4 baseline frames preceding the
stimulus. Population analysis was performed on only those neurons which evoked significant
responses to >25% of the trials (>25% Reliability). Reliability was computed as the proportion of
trials that showed significant responses out of 40 trials.

1203

Identification of Subplate Layer (Fig. 3, Fig. S1)
The identification of the SP Layer was based on separation drawn by sparse cell zone above,
and the horizontally oriented cytoarchitecture of the SP neurons. Based on the above features,
manual points (perpendicular to lamina) using MATLAB on slice images were selected to define
the depth and width of the SP layer. A similar analysis was performed too on the
immunolabelled brain slice images.

1210

LFP analysis (Fig. 3D-F)
For LFP analysis, baseline shifting was performed to have a mean of baseline 200 ms preceding
the stimulus at 0. To obtain LFP, acquired wideband signals were notch filtered (50Hz,
Butterworth 8th order, to remove AC supply line noise) and then band-pass filtered (between 5
Hz and 300 Hz, Butterworth 2nd order). LFPs were smoothed further using a Gaussian window,
SD 150 ms. LFP responses crossing the threshold of +/- 200 mV throughout the recorded time
window were excluded from population analysis. A spontaneous window of 400ms before the

onset and a response window of 500ms after the onset was considered for testing the significance of LFP response. The entire stretch of 900ms (spontaneous + response) was divided into 9 bins of 100ms each, making 4 bins in the spontaneous window and 5 bins in the response window. One-sided paired t-test was performed between the mean of the four spontaneous bins and each of the five response bins and only those traces with a p-value less than 0.05 in at least one of the five response bins were taken for further analysis. The normalized population RMS plots were plotted for these traces, where each of the post-onset RMS values was normalized with the mean of the 4 spontaneous RMS values.

Rate Calculation

Neuronal firing rate was computed by calculating the mean response within stimulus duration or token. Only single units that responded (compared to spontaneous activity 400ms preceding stimulus, 2-tailed t-test, $p<0.05$) to at least one of the stimuli (sound token (*f* or *N*) as standard or deviant) used were considered for analysis.

Latency Analysis (Fig. 3J)

For each neuron, tuning parameters (Best Frequency (BF) and Latency) were defined using custom-written codes in MATLAB. The best frequency for a neuron after ECO is the frequency which elicits a maximum response within stimulus duration (50 ms) to a given frequency at a given sound level. At ages after ECO, Latency to stimulus onset was defined as the time bin (5 ms resolution) at which peak response was observed for a BF. For calculating latency at ages before ECO, we first considered all Psths of neurons binned at 5ms with z score >1 within 500ms after stimulus onset. After that, a peak response within 300ms after stimulus onset was selected as the latency to the presented stimulus.

Adaptation Time Constants (Fig. S4)

A time-course analysis was performed to fit the response rates for SSS…..SDS….SS stimuli. Across all age groups and cortical layers, we employed exponential fitting to only those responsive units which have significant responses for tone as standards. To compute exponential decay time constants, the average spike rate in each token before the deviant was fitted with a first-degree exponential curve, based on least square error method. Fitting was performed using the 'fit' function in MATLAB.

1256 ***Construction of Reference frequency plots (Fig. 6)***

1257

1258 To investigate if the effect of exposure on firing rate were frequency-specific, the reference
1259 frequency plots (RF plots) were constructed to show the rate for the chosen reference
1260 frequency (S or D) as a function of its distance from the best frequency for each tuning curve.

1261

1262

1263 Table S4.

Age of exposure	Protocol (all 90 dB SPL, unless specified)	Litter	Animals	Units	Tuning curves
Control	No exposure	2	3+3	400	815
P6-P11	SD (12kHz-17kHz) SD (20kHz-14kHz)	2 1	2+3 3	256 197	504 525
P6-P11	SD (12kHz-17kHz) at 70 dB SPL	2	2+3	158	804
P11-P16	SD (12kHz-17kHz) SD (14kHz-10kHz)	1 1	5 3	367 233	700 553
P16-P21	SD (12kHz-17kHz)	2	2+2	130	255
P0-P5	SD (12kHz-17kHz)	2	2+2	183	370
P11-P16	SS (only 12kHz)	1	3	327	642
P6-P11	SS (only 12kHz)	1	3	194	487

1264

1265

1266

For example, if the reference frequency (RF) is chosen to be 12kHz (S), and for a given tuning
curve with the best frequency at 24kHz, the rate at RF is 12kHz is 10sp/s, then the RF plot will
have a value of 10sp/s 4 bins (1 octave) to the left of centre for that tuning curve. This is done
for all the tuning curves, and the population RF curve was plotted. The grey bar for the control
was constructed by repeating the same process 13 times considering each frequency as the
reference frequency and finding the maximum value of the rates at each bin, thus signifying the
maximum boundary. The bar plots were constructed by normalizing the RF rates at the best

frequency (centre of the RF curve). The normalization was done by dividing it with the mean of
all the BF rates except the ones within a distance of a half octave on both the sides of the RF
curve.

1277

1278

Connectivity between pairs (Fig. 7)

Cross-correlation between pairs of PSTHs (nrep= 20-30 repetitions of SD stimulus and swap, bin
size = 5ms) was used to detect connections between simultaneously recorded pairs of neurons
(SP-L4 pairs). Bootstrap analysis was done to get variability in the cross-correlograms (MATLAB
'xcorr' function, up to lag of 20 ms) by randomly resampling (same number of repeats from the
existing 20-30 repeats) with replacement, 100 times. Thus a mean and SD (dark line with
errorbar (Fig. 7B) of the correlograms were obtained. To compare with spurious cross-
correlations, all repeats from both neurons were mixed to give 2*nrep (number of repetitions)
spike trains, and randomly a number of reps of them were assigned to each of the neurons, and
from these PSTHs, cross-correlograms were obtained with mean and SD by repeating the
process 100 times (gray line with errorbar, Fig. 7B). Non-overlap of errorbars were considered
to be significant connections between the pair of neurons. We used 1.65-SD for data in P12 and
above groups and 1-SD in before ECO group due to the large difference in firing rates of single
units in the two cases.
The cross-correlation at best provides a linear time-shifted dependence between the activities
of L4 and SP, although convincing enough for our result, it is unable to capture the changing
temporal co-variability between the two signals. To address this issue, we also verified the
connectivity proportion results using granger causality, which provides more stringent criteria
to find the direction of the information flow (causation). We performed this between the mean
PSTH in 5ms binning of the SP and L4 responses.

1299

1300 **Details of Binary Network Model of L4 and SP with TC Inputs**

1301

1302 We built a binary neural network consisting of two separate thalamic input units for standard
1303 and deviant. The thalamic units project to a sub-plate neuron and further to a cortical neuron.
1304 The sub-plate neuron also projects to the cortical neuron. A similar model was used to
1305 demonstrate ocular dominance in visual cortex[4,5]. The spike rate of the two thalamic inputs are
1306 given by,

1307

1308
$$f_{std}(r) = \begin{cases} 8 & when\ a\ standard\ token\ comes \\ 0.15 & when\ a\ deviant\ token\ comes \\ 0.1 & spontaneous\ rate \end{cases}$$

$$f_{dev}(r) = \begin{cases} 0.15 & when\ a\ standard\ token\ comes \\ 8 & when\ a\ deviant\ token\ comes \\ 0.1 & spontaneous\ rate \end{cases}$$

1309

1310 The synaptic weight of the thalamo-subplate projections was kept constant at a value of 0.2.
1311 The projections received by the cortical neuron was kept plastic and learning was implemented
1312 following the below plasticity rule.

1313

1314
$$\frac{\Delta w}{w} = \begin{cases} Amp_{strength} * e^{-\frac{\Delta t}{\tau_{strength}}} & potentiation \\ -Amp_{weak} * e^{-\frac{\Delta t}{\tau_{weak}}} & depression \end{cases}$$

1315

1316 Where $\Delta t = |t_{post\ synaptic} - t_{pre\ synptic}|$
1317 $Amp_{strength} = 0.015 \qquad \tau_{strength} = 13\ ms$
1318 $Amp_{weak} = 0.021 \qquad \tau_{weak} = 20\ ms$
1319

We performed simulations for two regimes, one for the early development phase and one for
the later. For the early phase, we kept the synaptic weight of the sub-plate projections at a
value of 0.22 and thalamo-cortical projections at a smaller value of 0.02. It was done to mimic
the early stages of cortical development where the thalamic projections are unable to drive the
cortical activity alone and are aided by the sub-plate for driving the cortical activity[1,2,6].
In the later phase, both these weights were kept at an intermediate value of 0.04 and 0.12
respectively. This reflects the intermediate stage of development where the thalamo-cotical
projections are strengthening, and the sub-plate-cortical connections are weakening[2,6].
All the synapses were kept depressing throughout the simulation with the following equations
governing their dynamics[7],

1330

1331
$$\frac{dx_r}{dt} = -M.\frac{x_r}{\tau_{re}} + \frac{x_i}{\tau_{ir}} \qquad \frac{dx_e}{dt} = +M.\frac{x_r}{\tau_{re}} - \frac{x_e}{\tau_{ei}} \qquad \frac{dx_e}{dt} = \frac{x_e}{\tau_{ei}} - \frac{x_i}{\tau_{ir}}$$

1332

1333 Where M=1 whenever a spike occurs.

1334

1335 For thalamus, $\quad \tau_{re} = 0.9\ ms \qquad \tau_{ei} = 10\ ms \qquad \tau_{ir} = 4200\ ms\ (early)\ \&\ 100ms\ (later)$

1336

1337 For sub-plate, $\quad \tau_{re} = 0.9\ ms \qquad \tau_{ei} = 27\ ms \qquad \tau_{ir} = 4200\ ms\ (early)\ \&\ 100ms\ (later)$

1338

1339 The value of τ_{ir} was chosen in accordance with our electrophysiology data which shows a
1340 decrease in adaptation in later ages.

1341

1342 The sub-plate and the cortical neuron were modelled on the principles of integrate and fire
1343 neuron[1]. The membrane potential can be written as follows,

1344
$$V_{sp}(t) = g_{std}(t-1) * w_{s1}(t-1) * x_{r_{std}}(t-1) + g_{dev}(t-1) * w_{s2}(t-1) * x_{r_{dev}}(t-1)$$
$$V_{l4}(t) = g_{std}(t-1) * w_{l1}(t-1) * x_{r_{std}}(t-1) + g_{dev}(t-1) * w_{l2}(t-1) * x_{r_{dev}}(t-1)$$
$$+ g_{sp}(t-1) * w_{sp}(t-1) * x_{r_{sp}}(t-1)$$

1345

1346 Where g(t) is the conductance and can be written as,

1347

$$g(t) = spk\_trn * k(t)$$

1348 $k(t)$ is the convolution kernel
$$k(t) = e^{-(t-t_{spk})/\tau_{syn}}$$
$$\tau_{syn} = 10\ ms$$

1349

1350   The threshold of Layer IV neuron was kept constant at 0.1 for both the regimes, while that of
1351   the subplate neuron was kept at 0.05 for the early regime and 0.15 for the late regime.

1352

1353   The refractory period dynamics after each spike was modelled with a separate exponential
1354   given by[1],

1355

$$V_{\Delta}(t) = \beta * e^{-(t-t_{spk})/\tau_{refr}}$$
$$\tau_{refr} = 2\ ms$$
$$\beta = 5$$

1356

1357   The voltage drops by the above value after every spike. The refractory period lasts for 20ms
1358   after the spike. A leak of 10% was also incorporated[1].

1359

1360   The model gets a drive from the thalamic activity, modelled by uncorrelated Poisson spikes in
1361   the two thalamic units. The stimulus train consisted of standard and deviant tokens occurring
1362   with a probability of 0.9 and 0.1, respectively in the standard-deviant protocol and consisted of
1363   only standard tokens in case of only standard protocol. Each token is 50ms long separated by a
1364   250ms inter-token-interval with a sampling rate of 1000/sec.

1365

1366

1367   **Theoretical Analysis for Optimization based on Mutual Information Maximization and**
1368   **Sparseness**

1369

1370   We consider the same stimulus train that we used in the exposure experiments.
1371   We assume the probability of deviant to be x and that of a standard to be 1-x

1372

1373

$$p(t_{std}) = 1 - x \qquad p(t_{dev}) = x$$

1374

1375   The responses for each of the two tokens are normally distributed

1376

1377 $$p(r|t_{std}) = \frac{1}{\sigma\sqrt{2\pi}}\exp\left(-\left(\frac{(r-r_o)^2}{2\sigma^2}\right)\right) or\ g_s \quad p(r|t_{dev}) = \frac{1}{\sigma\sqrt{2\pi}}\exp\left(-\left(\frac{(r-(r_o+a_k))^2}{2\sigma^2}\right)\right) or\ g_d$$

1378

1379   We consider the variance σ^2 and mean response for the standard token r_o to be constant. The
1380   mean response for the deviant token varies with each successive token, and its difference from
1381   mean standard response takes values a_k for every kth token. We assume all the tokes to be
1382   independent and without any history dependence (no adaptation). Thus,

1383

$$p(r) = xg_d + (1 - x)g_s$$

1384

1385 We wish to know the trend of $f(k)$ with successive tokens that will enhance the discriminability
1386 between the standard and the deviant token. To do that, we use mutual information as a
1387 measure of discriminability paired with a sparse coding constraint and try to maximize it.
1388 Our constrained information for the k^{th} token is,
1389

$$CI_k = I(R_k, T) - \rho(r_o + x a_k)$$

1390

Where $I(R_k, T)$ denotes the mutual information between the response at the kth token and
the stimulus. ρ is the langrage multiplier which is constraining the mean response at the kth
token. The lagrangian has been constructed to minimize the mean response at the kth token
(sparse representation).
We need to find the value of ρ, which will maximize the mutual information between the entire
stimulus train and the total response along with maintaining a sparse representation
throughout the stimulus period. We can write the total constrained information as the sum of
the constrained information of independent individual tokens,
1399

$$CI_{total} = \sum_{k=1}^{N} CI_k = \sum_{k=1}^{N} I(R_k, T) - \rho\left(N r_o + x \sum_{k=1}^{N} a_k\right)$$

1400

1401 Where N is the total number of tokens.
1402 For any token, $I(R_k, T)$ can be written as,
1403

$$I(R,T) = \int_0^{r_{max}} \left[\sum_T \left\{ p(r,t) \ln\left(\frac{p(r,t)}{p(r)p(t)}\right) \right\} \right] dr$$

$$= \int_0^{r_{max}} \left[\sum_T \left\{ p(r|t)\, p(t) \ln\left(\frac{p(r|t)p(t)}{p(r)p(t)}\right) \right\} \right] dr$$

$$= \int_0^{r_{max}} \left[\sum_T \left\{ p(r|t)\, p(t) \ln\left(\frac{p(r|t)}{p(r)}\right) \right\} \right] dr$$

$$= \int_0^{r_{max}} \left[p(r|t_{dev})\, p(t_{dev}) \ln\left(\frac{p(r|t_{dev})}{p(r)}\right) + p(r|t_{std})\, p(t_{std}) \ln\left(\frac{p(r|t_{std})}{p(r)}\right) \right] dr$$

$$= \int_0^{r_{max}} \left[x g_d \ln\left(\frac{g_d}{x g_d + (1-x)g_s}\right) + (1-x)g_s \ln\left(\frac{g_s}{x g_d + (1-x)g_s}\right) \right] dr$$

$$= \int_0^{r_{max}} \left[x g_d \ln g_d + (1-x)g_s \ln g_s - (x g_d + (1-x)g_s) \ln(x g_d + (1-x)g_s) \right] dr$$

1404 $$= -\frac{1}{2}\ln(2\pi e\sigma^2) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x g_d + (1-x)g_s) \right] dr$$

1405 $$= -\frac{1}{2}\ln(2\pi e\sigma^2) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\{g_s(x A(r) + (1-x))\} \right] dr \qquad where\ A(r) = \frac{g_d}{g_s}$$

$$= -\frac{1}{2}\ln(2\pi e\sigma^2) - x\int_0^{r_{max}} [g_d \ln g_s]\, dr - (1-x)\int_0^{r_{max}} [g_s \ln g_s]\, dr - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x A(r) + (1-x)) \right] dr$$

$$= -\frac{1}{2}\ln(2\pi e\sigma^2) + \frac{x}{2}\ln(2\pi e\sigma^2) + \frac{x a^2}{2\sigma^2} + \frac{1-x}{2}\ln(2\pi e\sigma^2) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x A(r) + (1-x)) \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x A(r) + (1-x)) \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\left\{ (1-x)\left(\frac{x A(r)}{1-x} + 1\right) \right\} \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \ln(1-x)\int_0^{r_{max}} [x g_d + (1-x)g_s]\, dr - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\left(\frac{x A(r)}{1-x} + 1\right) \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \ln(1-x) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\left(\frac{x A(r)}{1-x} + 1\right) \right] dr$$

1406 We wish to use the Taylor expansion of ln in the above expression. To do that we need,
1407

$$\frac{x A(r)}{1-x} < 1$$

$$x < \frac{1}{1 + A(r)}$$

1408

1409 We see that $A(r)$ is the ratio of two normal distributions and it can be reduced to,

$$A(r) = exp\left(\frac{a(2r - 2r_o - a)}{2\sigma^2}\right)$$

1410

1411 Since x is a constant and is only dependent on the stimulus, we set a limit for x that will make
1412 this expression valid for all the possible rates,

1413

$$x < \frac{1}{1 + exp\left(\frac{a(2r - 2r_o - a)}{2\sigma^2}\right)}$$

1414 If we assume that on the 1st token (when the discrimination has not yet started), the mean rate
1415 for standard and deviant is the same (a=0), we see that,

1416

$$x < \frac{1}{2}$$

1417

1418 Thus we proceed further with an assumption that the probability of the arrival of the deviant is
1419 less than 50% and the results that we discuss later will hold true only on this assumption.
1420 We now perform the Taylor series expansion,

1421

$$\frac{xa^2}{2\sigma^2} - \ln(1-x) - \int_0^{r_{max}} \left[(xg_d + (1-x)g_s)\sum_{i=1}^{\infty}\left\{\frac{-1^{i+1}}{n}\left(\frac{xA(r)}{1-x}\right)^i\right\}\right] dr$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\int_0^{r_{max}}\{g_d\{A(r)\}^i\} dr\right] - (1-x)\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\int_0^{r_{max}}\{g_s\{A(r)\}^i\} dr\right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}\int_0^{r_{max}}\left\{exp\left(-\frac{(r-(r_o+a))^2 + i(a^2 - 2a(r-r_o))}{2\sigma^2}\right)\right\} dr\right]$$

$$- (1-x)\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}\int_0^{r_{max}}\left\{exp\left(-\frac{(r-r_o)^2 + i(a^2 - 2a(r-r_o))}{2\sigma^2}\right)\right\} dr\right]$$

1427 =

1428 $\frac{xa^2}{2\sigma^2} - \ln(1-x) - x\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}\int_0^{r_{max}}\left\{exp\left(-\frac{((r-r_o)^2+a^2-2(r-r_o)a+ia^2-2ia(r-r_o))}{2\sigma^2}\right)\right\} dr\right] -$

1429 $(1-x)\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}\int_0^{r_{max}}\left\{exp\left(-\frac{(r-r_o)^2+ia^2-2ia(r-r_o)}{2\sigma^2}\right)\right\} dr\right]$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}exp\left(-\frac{(i+1)a^2}{2\sigma^2}\right)\int_0^{r_{max}}\left\{exp\left(-\frac{((r-r_o)^2 - 2(r-r_o)a - 2ia(r-r_o))}{2\sigma^2}\right)\right\} dr\right]$$

$$- (1-x)\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}exp\left(-\frac{ia^2}{2\sigma^2}\right)\int_0^{r_{max}}\left\{exp\left(-\frac{(r-r_o)^2 - 2ia(r-r_o)}{2\sigma^2}\right)\right\} dr\right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}exp\left(-\frac{(i+1)a^2}{2\sigma^2}\right)\int_0^{r_{max}}\left\{exp\left(-\frac{(r^2 + r_o^2 - 2rr_o - 2ra + 2r_o a - 2iar + 2iar_o)}{2\sigma^2}\right)\right\} dr\right]$$

$$- (1-x)\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}exp\left(-\frac{ia^2}{2\sigma^2}\right)\int_0^{r_{max}}\left\{exp\left(-\frac{r^2 + r_o^2 - 2rr_o - 2iar + 2iar_o}{2\sigma^2}\right)\right\} dr\right]$$

1430

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}exp\left(-\frac{(i+1)a^2 + r_o^2 + 2r_o a + 2iar_o}{2\sigma^2}\right)\int_0^{r_{max}}\left\{exp\left(-\frac{(r^2 - 2(r_o + a + ia)r)}{2\sigma^2}\right)\right\} dr\right]$$

$$- (1-x)\sum_{i=1}^{\infty}\left[\frac{-1^{i+1}x^i}{i(1-x)^i}\frac{1}{\sigma\sqrt{2\pi}}exp\left(-\frac{ia^2 + r_o^2 + 2iar_o}{2\sigma^2}\right)\int_0^{r_{max}}\left\{exp\left(-\frac{r^2 - 2(r_o + ia)r}{2\sigma^2}\right)\right\} dr\right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x)$$

$$- x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{(i+1)a^2 + r_o^2 + 2r_o a + 2iar_o - (r_o + a + ia)^2}{2\sigma^2}\right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-(r_o+a+ia))^2}{2\sigma^2}\right) \right\} dr \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{ia^2 + r_o^2 + 2iar_o - (r_o + ia)^2}{2\sigma^2}\right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-(r_o+ia))^2}{2\sigma^2}\right) \right\} dr \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(-\frac{(i+1)a^2 + r_o^2 + 2r_o a + 2iar_o - (r_o + a + ia)^2}{2\sigma^2}\right) \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(-\frac{ia^2 + r_o^2 + 2iar_o - (r_o + ia)^2}{2\sigma^2}\right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(\frac{ia^2(i+1)}{2\sigma^2}\right) \right] - (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(\frac{ia^2(i-1)}{2\sigma^2}\right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^{i+1}}{i(1-x)^i} exp\left(\frac{ia^2(i+1)}{2\sigma^2}\right) + \frac{-1^{i+1}x^i}{i(1-x)^{i-1}} exp\left(\frac{ia^2(i-1)}{2\sigma^2}\right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x + x \sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} exp\left(\frac{i(i+1)a^2}{2\sigma^2}\right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x + x\Lambda(a)$$

1431 Where $\Lambda(a) = \sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} exp\left(\frac{i(i+1)a^2}{2\sigma^2}\right) \right]$

1432

1433 On a careful examination, we see that the above expression will always be positive.

1434 We find the change in constrained information between two tokens m&n (m<n)

1435

$$\Delta CI = CI_n - CI_m = \{MI_n - MI_m\} - \rho x\{a_n - a_m\}$$
$$= x \left[\frac{a_n^2 - a_m^2}{2\sigma^2} + \Lambda(a_n) - \Lambda(a_m) - \rho\{a_n - a_m\} \right]$$
$$= x(a_n - a_m) \left[\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} - \rho \right]$$

1436 We want the above expression to be greater than or equal to zero. Thus, we get two cases,

1437

1438 Case 1: $a_n > a_m$

1439

1440 We need,

1441

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} > \rho$$

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} \left\{ exp\left(\frac{i(i+1)a_n^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_m^2}{2\sigma^2}\right) \right\} \right]}{a_n - a_m} \geq \rho$$

1442

1443 We see that as a increases, the 2nd term in the above expression increases with the exponential
1444 of the square of the increase in a. The 2nd term overall will be much larger than 1. If we assume
1445 a monotonic behavior in a, the least value of $\Lambda(a_n) - \Lambda(a_m)$ will be between the first two
1446 tokens. Thus we can say that,

1447

$$\rho \leq \frac{a_2 + a_1}{2\sigma^2} + \frac{\sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} \left\{ exp\left(\frac{i(i+1)a_2^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_1^2}{2\sigma^2}\right) \right\} \right]}{a_2 - a_1}$$

1448

1449 The above expression sets a limit on the value of ρ or in other words, the weightage of
1450 sparseness which will keep the mutual information increasing.
1451 This case corresponds to the early age results, where the responses for deviant was greater
1452 than that of standard (a>0).
1453
1454 Case 2: $a_n < a_m$
1455 We need,

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} \leq \rho$$

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} \left\{ exp\left(\frac{i(i+1)a_n^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_m^2}{2\sigma^2}\right) \right\} \right]}{a_n - a_m} \leq \rho$$

1456
1457 We see that as a decreases, the 2nd term in the above expression decreases with the
1458 exponential of the square of the increase in a. The 2nd term overall will be much less than -1. If
1459 we assume a monotonic behavior in a, the largest value of $\Lambda(a_n) - \Lambda(a_m)$, will be between
1460 the first two tokens. Thus we can say that,
1461

$$\rho \geq \frac{a_2 + a_1}{2\sigma^2} + \frac{\sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} \left\{ exp\left(\frac{i(i+1)a_2^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_1^2}{2\sigma^2}\right) \right\} \right]}{a_2 - a_1}$$

1462
1463 The above expression sets a limit on the value of ρ or in other words, the weightage of
1464 sparseness which will keep the mutual information increasing.
1465 This case corresponds to the later age results, where the responses for the standard was
1466 greater than that of deviant (a<0).

1478

1479

1480

1481

1482 **Supplementary Figures**

1483

1484

1485

Figure S1. A) Representative confirmation of recording in A1 in lower age groups P12, P13, P18
with DiI injections in the recording site with labelling in MGBv (inset). B) Sample locations of
recordings of single units mapped onto the cortical surface showing broad clearer tonotopy in
P29 than in P15. C) Thickness of SP as a percentage of cortical thickness over age determined
from cell morphology with Nissl stains (Above). D) Slices were cut from brains (fixed) in which
single-unit recordings were performed in the ACX (representative examples at 3 ages, P12-P14,
Above) and IHC was performed. Electrode tracks show recordings performed in SPN and in case
of dual recordings (second electrode type, Fig. 3A) in SCNN1-td-tomato mouse (P14) show
electrodes in L4 and SPN simultaneously.

Figure S2. A) i) Example traces of zscores of 3 L4 neurons (upper 2 rows) and 3 SP neurons (lower 2 rows) in the first 3 columns, showing response to tone (f, Standard in Green and Deviant in Red, Above) and noise (N, Standard in Green and Deviant in Red, Below). A light gray patch is the mean rate window (250 ms) across which CSI(F) was computed. B) i) Mean population zscore of all L4 and SP neurons (upper and lower two plots respectively) in response to tone (f, Standard in Green and Deviant in Red, Above) and noise (Standard in Green and Deviant in Red, below) presented at 3.33 Hz. B) ii) Same as B) i), mean population zscore of L4 and SP neurons in responses to tone and noise, delivered at 4Hz.

Figure S3. Top and middle row shows the mean CSI values of the population of L4 and SP neurons respectively for different choice of response latencies and response window sizes. Asterisks mark the response latency and response window size combination used in Fig. 4 CSI calculations. The two columns are for the 2 different CSIs used CSI and CSI(F). Bottom row shows which combinations of response attribute have significantly higher CSI in SP than in L4 (brown, blue indicates NS).

1530

1531

1532

Figure S4. Ai) Population CDFs of adaptation time constants (as -1/time constant) for L4 neurons (Cyan) and SP neurons (Magenta) across different age groups (SP only for P12-15). Adaptation time constants, determined from exponential fits to mean responses to each successive token of Standards, increase with age. Aii) Shows population CDFs (for the same set of neurons and age groups as in the top row) of the number tokens in which there were significant responses in the first set of 7 (or 6) tokens before the deviant for the same population of neurons. B) Mean -1/Adaptation time constants for the data in Ai. Adaptation time constant increases with age (ANOVA, p<0.0001). C) Boxplots showing range with median, IQR and outliers for data in Aii. Significant responding tokens increases with age (ANOVA, p<0.0001).

1543
1544

Figure S5. A) Four example images of location (along the cortical surface) of BFs on the cortical surface in the craniotomy of single units recorded in L4 at ages P28 and above (Fig. 6A single-unit recording) identified in color (colorbar, bottom right) for four exposure groups EX2 and EX1 in the top row (SD exposure) and EX5 and EX6 in the bottom row (SS exposure). Broad tonotopy showing A1 and AAF are clear in the images. B) Bar plot shows the % change in an area of each of the preferred frequencies in A1 and AAF between control (EX0 group of mice) and the only standard P10-P15 group (EX6).

1552

1556 Figure S6. Bar plot summarizes population mean absolute response to RF for neurons with BF at
1557 RF of all 6 EX cases (2 bars each, red and green for SD exposure cases and 1 blue for SS). Mean
1558 of the 13 control cases shown in gray bar in the centre. The stripped bar here represents the
1559 p6-p10 70 dB SPL exposure protocol (EX5).

1560

1561

1562

1563

Supplementary Videos

Supplementary Video 1: This video demonstrates the widefield imaging-based sound-evoked responses to tone (17kHz, 20dB attenuation, ~ 70 dB SPL) in ACX of an awake Thy1-Gcamp6f mouse at P9. Video acquired at 10 fps.

Supplementary Video 2: This video demonstrates the 2-Photon imaging-based sound-evoked responses to tone (24kHz, 0dB attenuation, ~90 dB SPL) in ACX of a Thy1-Gcamp6f mouse at P7. Video acquired at 5 fps.

Videos are available at

https://drive.google.com/drive/folders/1UvD6NXDpHhnKrEB1N2zaoAW4S-gl7jDi

References

1.   Gee, J. M. *et al.* Imaging Activity in Neurons and Glia with a Polr2a-Based and Cre-Dependent GCaMP5G-IRES-tdTomato Reporter Mouse. *Neuron* (2014) doi:10.1016/j.neuron.2014.07.024.

2.   Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates*. *São Paulo, Academic Press* (2012). doi:10.1364/OE.20.020998.

3.   Torii, M. & Levitt, P. Dissociation of corticothalamic and thalamocortical axon targeting by an EphA7-mediated mechanism. *Neuron* (2005) doi:10.1016/j.neuron.2005.09.021.

4.   Kanold, P. O. & Luhmann, H. J. The Subplate and Early Cortical Circuits. *Annu. Rev. Neurosci.* (2010) doi:10.1146/annurev-neuro-060909-153244.

5.   Kanold, P. O. & Shatz, C. J. Subplate Neurons Regulate Maturation of Cortical Inhibition and Outcome of Ocular Dominance Plasticity. *Neuron* (2006) doi:10.1016/j.neuron.2006.07.008.

6.   Wess, J. M., Isaiah, A., Watkins, P. V & Kanold, P. O. Subplate neurons are the first cortical neurons to respond to sensory stimuli. *Proc. Natl. Acad. Sci.* (2017) doi:10.1073/pnas.1710793114.

7.   Mill, R., Coath, M., Wennekers, T. & Denham, S. L. A neurocomputational model of stimulus-specific adaptation to oddball and markov sequences. *PLoS Comput. Biol.* (2011) doi:10.1371/journal.pcbi.1002117.