1

Earliest experience of rare but not frequent sounds cause long term changes in the adult auditory cortex

Authors: Muneshwar Mehra[1], Adarsh Mukesh[1] and Sharba Bandyopadhyay[1,2*]

[1] Advanced Technology Development Centre, IIT Kharagpur, [2] Department of Electronics and Electrical Communication Engineering, IIT Kharagpur

[*]Corresponding author

Pages: 27 (+ 44 Supplementary Material), No of Figures: 6 (+ 20 Supplementary +5 Tables)

Author contributions: MM performed all experiments, AM assisted. MM, AM and SB analyzed data. AM did modelling and theoretical derivation. SB supervised the work. SB, MM and AM wrote the manuscript.

Acknowledgements: MM thanks CSIR for Fellowship, AM thanks MHRD for PMRF, SB thanks India Alliance, IIT Kharagpur, MHRD and SRIC for funding. SB gives special thanks to PPC and SDG for support and POK for imbibing interest in the subplate.

Abstract

Early sensory experience during the critical period of development is known to alter sensory
cortical response properties and organization. We find that the earliest sound driven activity
in the mouse auditory cortex (ACX) can be detected before earcanal opening (ECO). How
auditory experience at these ages may influence auditory cortical development is unknown.
Particularly, subplate neurons (SPNs), involved in sensory thalamocortical maturation, in the
auditory cortex, are also found to be driven by sounds at ages before ECO in mice. We find
that SPNs are selective to low probability deviant sounds in auditory streams before ECO,
more so than thalamo-recipient, layer 4 (L4) neurons and not after ECO. We hypothesize
that SPNs with their deviant selectivity can direct development of L4 responses before ECO.
Exposing mice before ECO with a rarely occurring tone in a stream of another tone occurring
frequently leads to strengthening of cortical representation of the rare tone, but not the
frequent tone in the adult. Control exposure experiments with only a frequently occurring
tone and also in other developmental age windows corroborate the importance of low
probability sounds in auditory development. With a computational network model of known
thalamic inputs to SPNs and L4 we explain the observed developmental plasticity. An
information theoretic analysis with sparse coding assumptions also predicts the
observations. Thus, salient low probability sounds in the earliest auditory environment
causes long term changes in the auditory cortex.

Introduction

Important and salient sensory stimuli occur with low probability, while high probability
stimuli, occurring repeatedly, are usually of less importance and are ignored by an organism.
Sensory cortical circuitry development is activity dependent and is based on the sensory
environment to which an animal is exposed[1,2]. Particularly in the auditory system, altering
the natural auditory environment during specific periods (critical period[3–5]) of early
development remarkably alters cortical circuitry and organization[6,7]. SPNs, the first born
cortical neurons[8,9], play a crucial role in the development of functional cortical circuitry and
organization[10–14]. Here, with a new auditory exposure paradigm before the established
auditory critical period[2–5] (before ECO) using low probability salient/deviant stimuli in the
early auditory environment, we show that the functional auditory cortical (ACX) responses
are remarkably altered into adulthood. The alterations are specific to the deviant or rare
exposure stimulus. The observed plasticity is an outcome of the unique deviant detection
properties we find of SPNs, before ECO, with strong responses to only the rare stimulus. A
computational network model derived from our experimental results show how such
plasticity occurs. Further, using mutual information maximization and sparse coding
principles, we show that the outcome of early exposure with our new paradigm can be
predicted theoretically. Hence our results revise the established timelines and concept of
auditory critical period and may generalize to other sensory systems.

The development of basic response properties of neurons, in the ACX of the rodent ECO
(P12) onwards, like tuning, tuning width, thresholds to tonal sounds or broadband noise and
other basic stimuli have been studied[3,15]. In spite of the importance of low probability
stimuli[16], developmental profile of response properties to streams or sequences of sounds
which are more naturalistic are not known. The changes in response properties of neurons
in the ACX, in a lamina specific manner, over the developmental time course, during the
critical period have also not been studied. The early transient SP layer is an integral part of
the thalamocortical (TC) projection pathway[9–11,17–19], in the auditory or other sensory areas.
Although in-vivo sound driven responses in ACX SPNs of ferrets exist[20], they are not fully
characterized and not known to exist in mice. SPNs receive collateral inputs from the
thalamic axons projecting to layer IV (L4)[8] in the ACX[9,18], but their functional role, if any, in
the development of the ACX have not been studied.

Results

Given the importance of low probability stimuli[16] and stimulus specific adaptation in the
adult[21,22], single unit responses in L4 and SPNs in A1 and AAF of the developing mice to an
oddball stimulus paradigm were first investigated. Single unit responses of the oddball
stimulus were obtained from lightly anaesthetized mice (0.5-0.75% isoflurane in oxygen)
from postnatal day 12, P12 (after ECO) up to P48 (N=51, male and female). A1 and/or AAF
was targeted with custom designed microelectrode arrays (4x4 MEA, 125 μm apart, all 16 of
the same length or 8 each staggered by 400 μm or 600 μm, for simultaneous L4 and SPN
recording, Fig. S1A), based on vasculature (Fig. S1B)[23–25]. Recording areas and depths were
confirmed to be in the primary ACX and from L4 and SP (Figs. S1-3). First, basic responses to
broadband noise and tones, at multiple sound levels were obtained (50-100 ms long, gap 2-
7s, tones 6-48 kHz in 1/4[th] or 1/2 oct steps). Next, a series of sound tokens (n = 15 or 10),
each 50 ms long (at 4 Hz), of the pattern SSS…SDS…S, with S as Standard and D as Deviant
(at 8[th]/7[th] position of 15/10 total tokens) were used for the oddball response
characterization. S and D could be two distinct tones (6-34 kHz) or a tone and a broadband
noise (6-48 kHz with equivalent sound level). To normalize for a neuron's inherent
selectivity for a particular S or D, responses were always collected in pairs with the S and D
swapped (Fig. S4A). Each SD set of a pair was presented with 10-40 repetitions with a 1-7s
silence duration between the end of the previous and start of next stream. To quantify a
neuron's selectivity to the deviant we used the common selectivity index

CSI = $[R_D(f_1)+R_D(f_2)-R_S(f_1)-R_S(f_2)]/[R_D(f_1)+R_D(f_2)+R_S(f_1)+R_S(f_2)]$ or

$[R_D(f)+R_D(N)-R_S(f)-R_S(N)] / [R_D(f)+R_D(N)+R_S(f)+R_S(N)]$[16];

where $R_S(.)$ and $R_D(.)$ denote response to S and D respectively. Three cases for $R_S(.)$ were
considered: mean response to first S or onset, mean response to S immediately preceding

the deviant or mean response for all the S's; the selectivity values are referred to as CSI(F),
CSI(P) and CSI respectively.

Early responses in the auditory cortex show higher deviant selectivity than in the adult

Single unit responses to the SD paradigm with 2 different tones, f_1 and f_2, spaced either 1/4,
1/2 or 1 octave apart were recorded in mice at different ages. Data were simultaneously
collected from multiple single units and hence f_1 and f_2 were at various differences from the
BF of each of the units. Based on online observations one of the frequencies (f_1 or f_2) was
chosen to be at or near the BF of the majority. Figure 1 shows representative responses of 3
single units in L4 and SP during the early developing period after ECO (P12-P15, Fig. 1A-C).
Plots of spike dot raster, in response to tones, are shown in Fig. 1A-Ci (top) along with the
peri-stimulus time histograms (PSTHs, bottom) for the response to chosen f_1 and f_2 (dark
and light grey rectangles in raster). The responses to the SD paradigm (raster plots and
PSTHS) for the same units are shown in Fig. 1A-Cii (time scale expanded in Fig. 1A-Ciii).
Reliable responses were observed to tones and SD stimuli in the P12-15 group of animals
with wider tuning (Methods) in L4 than in SP (0.34 and 0.25 octaves, one sided unpaired t-
test, df=101, p<0.010). Simultaneous recordings in L4 and SP (Fig. 1D) showed lower peak
response latencies (~5 ms more than first spike latency) for BF tones in SP than in L4 in P12-
P13 group (mean 27.38 and 32.27 ms, *p*<0.0001, df=246 two-sided unpaired *ttest*, Fig. 1E)
and in P14-P15 group (mean 27.08 and 30.52 ms, *p* <0.05, df=78 Fig. 1E).

CSI from the pairs of SD responses (Fig. 1A-Cii) were calculated and the mean CSIs observed
as a function of separation of frequencies used (CSI(F), 0.5 octave: p<0.0001, df=48, two-
sided unpaired *ttest*; 1 octave: p=0.0031, df=61 Fig. 1F, and CSI, 0.5 octave: p<0.0001, df=48;
1 octave: p<0.0001, df=61 Fig. 1G) for L4 (6 mice, 69 units, 110 cases) were higher than that
of SPN (6 mice, 43 units, 65 cases), except at ¼ octave separation for the population of
neurons in P12-15 (p<0.099, df=48, Fig. 1F & p<0.2535, df=48 Fig. 1G). CSI(F) is biased to be
low due to the large onset response, evident from the lower values of CSI(F) than CSI (Fig.
1F-I). On average, independent of f_1-f_2, SP neurons had lower deviant selectivity than L4
neurons and they reduced from P12 to P15. Further, as age progressed from ECO (P12) to
the adult the CSI values and hence deviant selectivity reduced (P12-13$_{L4}$ vs P16-21$_{L4}$: df=60,
p=0.0019, two sided unpaired t-test; P12-13$_{L4}$ vs P22-28$_{L4}$: df=98, p<0.0001; P12-13$_{L4}$ vs
>P28$_{L4}$: df=119, p<0.0001; P14-15$_{L4}$ vs P14-15$_{SP}$: df=112, p=0.005 Fig. 1H & P12-13$_{L4}$ vs P16-
21$_{L4}$: p<0.0001; P12-13$_{L4}$ vs P22-28$_{L4}$: p<0.0001; P12-13$_{L4}$ vs >P28$_{L4}$: p<0.0001; P12-13$_{L4}$ vs
P12-13$_{SP}$: df=59, p=0.0011; P14-15$_{L4}$ vs P14-15$_{SP}$: p=0.033 Fig.1I), showing that adults had
much weaker selectivity to the deviant than in early ages. The reduction in selectivity to the
deviant with age was not due to the distribution of choices of f_1 and f_2 in the different age
groups (Fig. S4).

Since deviant selectivity is dependent on f_1-f_2 (Fig. 1F&G) and hence can be confounding we
also used an *N-f* pair (a noise token and a single frequency, Fig. 2 and Fig. S5&S6). Since the

noise token is broadband, in this case neurons would show deviant selectivity if there are parallel channels followed by integration based on spectral contrast of sounds[26]. Robust responses were present for the broadband noise token, both, when presented by itself (Fig. S5Bi-Gi, sometimes showing inhibition, Fig. S5Ci) and in the SD paradigm (Fig. 2). Single units responded to noise, both in L4 (Fig. 2A and Fig. S5) and in SP (Fig. 2B and Fig. S6) of A1 not only in the age group P12-P15 but also, expectedly, in L4 neurons for ages above P15. Minimal SPNs were observed beyond P15 (Fig. S1D) and hence SPNs were not targeted beyond P15. In the case of noise and tone as well, the same age dependence of deviant selectivity was observed as with tones. The difference in L4 and SP were similar to that observed with tones for both CSI(F) and CSI. CSI(P) in which $R_S(.)$ is the response rate for the standard token preceding the deviant also showed the same effects (two sided unpaired t-test, P12-13$_{L4}$ vs P22-28$_{L4}$: df=98, p<0.0001; P12-13$_{L4}$ vs >P28$_{L4}$: df=119, p=0.0464 Fig. S7).

In both the SD responses the age dependence of deviant selectivity (P12-13$_{L4}$ vs P16-21$_{L4}$: df=86, p=0.025, two sided unpaired t-test; P12-13$_{L4}$ vs P22-28$_{L4}$: df=152, p<0.0001; P12-13$_{L4}$ vs >P28$_{L4}$: df=118, p<0.0001 Fig. 2C & P12-13$_{L4}$ vs P16-21$_{L4}$: p=0.022; P12-13$_{L4}$ vs P22-28$_{L4}$: p<0.0001; P12-13$_{L4}$ vs >P28$_{L4}$: p<0.0001 Fig. 2D) is found to be correlated with changes in adaptation time constant[27] with age. An exponential decay fit to responses at each successive S token from the onset for every unit, shows that the mean of the time constants increases with age (Fig. S8Ai,B; P12-13$_{L4}$ vs P16-21$_{L4}$: df=74, p=0.027, two sided unpaired t-test; P12-13$_{L4}$ vs P22-28$_{L4}$: df=209, p<0.0001; P12-13$_{L4}$ vs >P28$_{L4}$: df=148, p=0.0026). Similarly the number of S tokens, from the onset, to which neurons respond significantly also increase with age (Wilcoxon ranksum test, P12-P13/P21-P27, P<0.001, (Fig. S8Aii,C), expected from increasing decay time constants. These results also indicate that A1 neurons even early in development do not necessarily only integrate information in parallel frequency channels but also possibly along parallel spectral contrast channels; as responses to deviant tone (high spectral contrast) is observed contrary to the idea that a broadband noise token as S, should cause all frequency channels to adapt, leading to lack of deviant tone response. Response to tones and noise can be passed on to the cortex early in development as parallel channels of information. The above is further corroborated by lower CSI and CSI(F) observed in MGBv computed from *N-f* SD responses (Fig. S9B; A1 P14-15$_{L4}$ vs MGBv P14-15: df=180, p<0.0001, two sided unpaired t-test; A1, P14-15$_{SP}$ vs MGBv P14-15: df=132, p=0.0046; A1 P22-28$_{L4}$ vs MGBv P22-28: df=165, p=0.9755; Fig. S9C A1 P14-15$_{L4}$ vs MGBv P14-15: df=180, p<0.0001, two sided unpaired t-test; A1, P14-15$_{SP}$ vs MGBv P14-15: df=132, p<0.0001; A1 P22-28$_{L4}$ vs MGBv P22-28: df=165, p<0.0001) in two age groups P14-15 (n=2, 27 units, 71 cases) and P24-26 (n=3, 42 units, 79 cases). The higher deviant selectivity in L4 and SP especially in the developing mice (P12-P15) is a cortical phenomenon and is not a reflection of thalamic inputs. The CSI values obtained in the MGBv in P14-15 age group were drastically lower than that observed in L4 (CSI, A1 P14-15$_{L4}$ vs MGBv P14-15: two sided unpaired t-test df=180, p<0.0001) and SPN and lower than L4 at higher ages (P24-26). In both cases, two tones and noise-tone, the CSI values are larger than

182 that obtained with a long stimulus protocol[21] as we use a stream with a single deviant that is
183 repeated.

Mouse ACX neurons have auditory responses before ECO

Low threshold hearing onset coincides with ECO, however high intensity sounds may drive
spiking activity in the ACX before ECO. Auditory cortical responses have been shown to exist
in ferrets[20] at ages equivalent to P6 in mice[28], however no study reports presence of sound
driven activity in the ACX of mice at ages before ECO (P11/12). However, auditory brainstem
responses in mice have been shown before ECO[29] and also long term changes in prestin
expression have been reported in mice exposed to loud sounds at these early ages[29] (P6-
P11). Although the central auditory pathway is functional[29,30] showing spontaneous activity
the peripheral auditory system maybe immature. In mice, the external auditory meatus
reopens at P7 and the outer ear or pinna is fully mature by P7[31] and middle ear bones lever
arm lengths are at 70% of mature values with normal lever ratios by P7[32]. The oval window
is also of normal size by P7[32]. Most importantly, mechano-transduction currents from outer
hair cells (OHC) and inner HCs (IHC) become maximal in the cochlear base by P2 and at the
apex by P5[33,34]. IHCs are fully mature by P5[35] while OHCs are immature with nonlinear
capacitance (NC) of OHCs, key to low threshold hearing and high sensitivity of auditory
nerve fibers, only starts at P7 and is fully mature at P18. The development of OHC NC is
correlated with the expression of the motor protein, prestin, responsible for the OHC NC[35].
Thus high intensity sound evoked activity is possible at ages before ECO in mice.

To investigate the possibility of auditory responses to high intensity noise and tones in the
ACX of mice before ECO[3,36] we performed additional experiments using single unit
extracellular recordings, wide-field fluorescence imaging and 2-photon Ca^{2+} imaging in Thy1-
GCaMP6f (Jax Labs) at ages P7-11 before ECO (Fig. 3 and Fig. S10-14). Sound evoked long
lasting LFP's were present as early as P08 (before ECO, Fig. S10A-B). Single unit activity
collected from SP and L4, with 5 repetitions of tones or noise did not show significant
responses at P08-10 (Fig. S10C). Sound driven spiking activity in the ACX before ECO, was
confirmed first with widefield fluorescence imaging in Thy-1-GCamp6f mice (Fig. 3A and
Fig.S11A). Over age, from P07 to P13, progressively larger areas of the cortical surface
showed significant responses to multiple tone frequencies (~90 dB SPL, 40 repetitions), with
the presence of clear tonotopy in A1 by P13 (Fig. 3A). The presence of significant responses
in Ca^{2+} dependent fluorescence could potentially indicate activity in input axons to the ACX
and/or spiking responses in ACX in response to sounds as early as P7. We confirm presence
of sound driven activity lower in the auditory pathway by performing widefield fluorescence
imaging in the inferior colliculus (IC) of Thy1-GCaMP6f (Jax Labs) mice at P9-10 (Fig. 3B).
Since 5 repetitions were not sufficient to show significant activity in ACX single units while
40 repetitions showed significant sound evoked responses in widefield fluorescence, in both
the ACX and the IC, we hypothesize that responses at these early ages have low reliability (%

of responding trials, Methods). Moreover, the percentage of responsive cortical surface area
and fraction of frequencies that showed significant responses increase with age (Fig 3A, Fig.
S11A, Table S3). Both the above indicate gradual maturation of the ACX.

To rule out, only input axon activity in the ACX widefield imaging, 2-photon Ca^{2+} imaging of
single neurons in L2/3 of the ACX of mice before ECO was also performed with 40
repetitions each of noise bursts and different tone frequencies (Methods, Fig. S11B-C). A
small fraction of significantly responsive neurons to different tones and broadband noise
was found in L2/3 from P07 to P11, before ECO (Table S4-S5). Considering all the neurons
that responded with at least 25% reliability (Fig. 3Ci), the population mean df/f in each age
was investigated (Fig. 3Cii). Increase in the population mean response strength and
decreasing trend in peak latency with age was found (Fig. 3Cii, any frequency tone, Fig. S12,
each frequency tone and Fig. S13Ai-ii, noise bursts). The above indicated a gradual
maturation of responses up to ECO (P12-13). Also, significant positive correlation is found in
response reliability with age (Fig. 3Ciii, Fig. S13Aiii) before ECO, showing gradual response
development.

Although with 40 repetitions of tone/noise, detecting spurious responses due to high
spontaneous activity observed before ECO are unlikely, we used two approaches to rule out
such a possibility. In additional 2-photon Ca^{2+} imaging experiments in P9 and P10 mice
(before ECO), first we considered >2s baseline before tone presentation. Considering only
neurons with >25% reliability of responses (Fig. 3Di) and their population mean df/f (Fig.
3Dii), we found that the baseline did not show any such activity as observed following the
stimulus onset. Secondly, we also repeated the same imaging experiments with and without
sound stimulation. The df/f of the neurons that responded with >25% reliability with sound
presentation (Fig. 3Ei, left) was also obtained for the same durations without sound
presentation (Fig. 3Ei, right). The population means in the 2 cases (Fig. 3Eii) clearly show the
presence of sound driven activity in the ACX L2/3 single neurons before ECO.

Two-photon Ca^{2+} imaging of single neurons, simultaneously from L4 and SPNs was beyond
our scope. To investigate the nature of spiking activity in L4 and the SP, with larger number
of repetitions of tone and noise stimuli (due to the observed low reliability), single unit
recordings were done in mice with ear canals closed (n=6, P8-11) with staggered electrodes
(Fig. S1A). Unlike with 5 repetitions of stimuli (Fig. S10C), z-scores of spiking responses in
50ms time bins (Fig. 3Fi, and Fig S13Bi) show presence of significant sound evoked spiking
in SP and L4 neurons before ECO. Population PSTHs of all significantly responding units in SP
and L4 (Fig. 3Fii and Fig. S13Bii) show significant sound driven activity before ECO, which get
stronger with age.

 SPNs unlike L4 and L2/3 neurons, show deviant selectivity before ECO

As responses to tones and noise exist in the mouse ACX before ECO, we investigated deviant
selectivity properties in ACX L2/3, L4 and SPNs. 2-photon Ca^{2+} imaging of the ACX L2/3 single
neurons as early as P07 show significant responses to S and D in the *N-f* SD protocol (Fig.
S14Ai-ii). Mean response profiles of all significantly responding L2/3 neurons show
responses to both the S and D (Fig. S14Aiii-iv). However, distribution of CSI (mean -0.037)
values of single neurons based on mean df/f (Fig. S14Av) indicates no preference for the D,
in L2/3 single neurons.

Activity of single units during presentation of the SD protocol for f_1-f_2 and *f-N* (30 and 20
reps, 3.3 or 4 Hz repetition rate, Fig. 4Ai) showed peaks in response following the deviant
(mainly in SPN) and onset (mainly in L4) as shown in the first 3 columns of Fig. 4A (dot
rasters and PSTHs of 3 L4 neurons, dark gray, upper four rows, SD and swap and 3 SPNs,
light gray, lower four rows). Population mean PSTHs for all L4 neurons and all SPNs recorded
(P8-P12, before ECO) indicate the presence of deviant detection in SP and not L4 (Fig. 4Aii-
iii, 3.3 Hz and 4 Hz). Further, based on the spontaneous activity (100 ms preceding the
stimulus), z-scores of the example neurons and population of neurons (Fig. S15AB)
demonstrate responses (95% CI, one-sided), particularly following the onset and the
deviant. In the population mean z-score, the earliest peaks occur in L4 in a 250 ms window
with a latency of 50-100 ms from the first token and in SP in a similar window following the
deviant. Considering a latency of 100 ms, because of observations in LFPs and by obtaining
spike rates in 250 ms windows because of width of significant LFP rms values (50 ms sound
and 200 ms gap), CSI(F), CSI and CSI(P) were calculated for L4 and SPN (Fig. 4Bi-iii). There is
higher deviant selectivity in SP than in L4 (two sided unpaired t-test, p<0.01 for all CSIs).
Since in single trials for single neurons the responses were not reliable, we also varied the
response window from 50 ms to 300 ms (in steps of 50 ms) and latency from 0 to 350 ms (in
steps of 50 ms) and obtained the CSIs for all neurons and we found for what combinations
SPNs had significantly higher deviant selectivity than L4 neurons (Fig. S16). Thus in spite of
low reliability of responses in the population L4 did not show higher deviant selectivity than
SP. Thus neurons in the mouse ACX respond to sounds before ECO with SPNs showing
responses to the deviant. Possibly, depending on the strength of SPN to L4 connections, the
L4 may or may not have a response. Ten pairs of neurons (between L4 and SP) were found
to be connected out of 27 simultaneously recorded L4-SP pairs (Methods) before ECO (Fig.
S17). Observed directionality of the connected pairs suggests existence of connections from
SPN to L4 (6/12 pairs showed connections from SPN to L4 within spatial distance of 125
microns while 4/15 pairs were found to be connected at distance of >125 microns, p<0.11).
Directional connectivity as a function of spatial distance clearly suggests columnar
organisation of SPN to L4 connections in ACX[19] before ECO (Fig. S17B). Due to the low
reliability of responses (Fig. 3Diii-Eiii), to rule out spurious deviant selectivity, we considered
1000 random pairs of windows, in CSI calculations considering two non-overlapping

windows 250 ms randomly taken in the standard region before the deviant (Fig. 4Biv) or after the deviant (Fig. 4Bv) where the mean CSIs were not significantly different between L4 and SP in >95% of the random pairs. The distributions of mean random-CSIs for SPN and L4 also show that the means were not significantly different from 0. Thus SPNs show significantly stronger responses to the deviant than L4 before ECO and lower than L4 after ECO. Such early context dependence of responses in SPNs allow for the possibility of SPNs in sound driven sculpting of TC inputs earlier in development than previously studied[11].

Exposure in a standard deviant acoustic environment before and after ECO show differential developmental plasticity for the standard and the deviant frequencies

Broad tonotopic organization in the primary auditory cortex (A1)[23,24] is known to be altered with over-representation of a particular frequency in the adult rats/mice which were exposed to that frequency tone during the critical period (P12 to P15, after ECO)[3,5]. However, as observed (Fig. 3-4), spiking and LFP (Fig. 3-4 and Fig. S10-S14) responses in the L2/3, SP and L4 in ACX are present before ECO. Further it is known that the auditory pathway is active before P12 (in rodents), with the TC circuitry maturing[37–39], functional hair cells by P6[40] and SPNs in A1 responds to sounds[20] (in ferrets, at ages developmentally equivalent to mouse P7)[28]. In spite of such activity, exposure of rodents to a single frequency tone from P9-P12, no changes were found in broad tonotopic organization with single unit BF mapping of A1, when the exposed pups were adults[3,5]. SPNs and L4 neurons in the mouse A1 respond to sounds as early as P08-P12 (before ECO, Fig. 3-4 and Fig. S10-12) and SPNs act as deviant detectors with high context dependence (Fig 4 and Fig S13). Such response properties and SPN to L4 connections (Fig. S17)[18] coupled with activity dependent plasticity[1,6] in the early auditory system leads to the hypothesis that exposure to a continuous train of a fixed frequency tone (S) with an embedded low probability (10%) tone (D), before ECO would cause enhanced responses to the D frequency in the adult. Spiking in the SPNs drives L4 spiking[18,19] in a correlated manner, with thalamic spikes in neurons projecting to SPN and L4 would drive strengthening of thalamus to L4 synapse[11]. The above suggests that exclusive responses to the deviant in SPNs would lead to strengthening of the thalamic inputs tuned to the deviant frequency. Mice were exposed (Fig. 5A) to such an SD sound protocol (Methods) from P6-10/11 (S=12 kHz, D=17 kHz or S=20 kHz, D=14 kHz, EX2, Fig. 5A) and single unit recordings in L4 (depth 350-450 μm) in response to pure tones (13 frequencies, 6-48 kHz, 1/8[th] oct apart, 50-90 dB SPL, 10 dB steps) from A1 and AAF were performed in the mice at P28-42. As age controls we did exposures of the SD (S=12 kHz, D=17 kHz) protocol at multiple age groups – P0-5/6 (EX4), P11-15/16 (also, S=14 kHz, D=10 kHz, EX1) and P16-21 (EX3). For stimulus control, exposure of mice in age groups P6-10/11 (EX6) and P11-15/16 (EX5) were done with only a single frequency (SS protocol, S=12 kHz). Further, a group with normal rearing was considered as control (EX0).

All significant response firing rates from the tuning curves (cartoon tuning curves, Fig. 5B) for each of EX0 to EX6 show that EX3, EX4 and control (EX0) have same median firing rate (ranksum test, 12 sp/s for every case, p_{EX3}=0.8475, p_{EX4}=0.1875 Fig. 5C). All other groups, EX1-2, EX5-6 show increased median firing rates (*ranksum* test, 16-24 sp/s, $p_{EX1,2,5,6} < 10^{-6}$, Fig. 5C) compared to EX0. The cumulative distributions (cdfs) of mean firing rates at the respective reference frequencies considered in the different exposure groups (green: S, red: D and blue: SS, Fig. 5D) are within the boundaries (thick gray lines) of EX0 cdfs (except at very low rates in EX3-4). Since any particular frequency out of 13 frequencies could be a reference we considered each of the frequencies in EX0 as a reference frequency (mean with STD, thin gray lines, Fig. 5D) to find the limits of the cdfs in the control. The minimum and the maximum from the 13 cdfs of control are considered as the boundaries. With our limited resolution mapping, we did not find any apparent over representation of any particular frequency in BF maps showing broad tonotopy (A1 and AAF) on the cortical surface (Fig. S18A) in any of the exposure cases (EX1-2 and EX5-6), unlike other studies with EX5 group[3,5] (Fig. S18B). Other than lower spatial resolution of single unit sampling, it may also be due to parameters of exposure (sound frequency, duration and repetition rate).

From the tuning curves, population mean response rates to the relevant reference frequency (RF) were plotted as a function of BF (re RF in octaves, Fig. 5B). For S/D exposures (EX1-4) we obtained two plots, with S as RF and D as RF (green and red respectively), for SS exposures (EX5-6), S was the RF (blue) and for the control C (gray) each of the 13 frequencies were considered as RFs separately. The maximum rate in the profile (as a function of BF re RF) of the 13 reference frequencies (EX0 case) is considered as the upper boundary of observed rates at RF in control (Fig. S19 and Fig. 5E), minimum being '0'. When considering normalized firing rates (with rate at BF as 100%) the population mean responses to reference frequencies (for EX1-2 and EX5-6) were largely within the maximum boundary suggesting no frequency specific change in normalized rates compared to control (Fig. S19). The population mean rate as a function of BF re RF for EX2&5 and EX1&6 (Fig. 5Ei and ii) show differential effects on response strength at BF to the S and D for SD exposure before ECO and after ECO. After ECO in the critical period of P11-15, the S frequency responses (with S as BF) are strengthened in comparison to that of D, while before ECO the opposite effect is observed. No such changes are observed in exposure before P6 and after P15 (Fig. S20 bar plots, two sided unpaired t-test, P11-15 standard vs deviant: df=107, p=0.009; P6-10 standard vs deviant: df=120, p=0.002). In comparison when SS exposure is done after ECO there is a strengthening of responses to S and not after ECO. Only in the case of SS for P11-15 exposure there is increase of S responses of neurons with S as BF beyond the boundary of control (gray dashed line, Fig. S20) and that for D before ECO SD exposure. Comparisons (significant or not) with the population of 13 control rate profiles are marked on respective bars (gray '*' or NS, Fig. S20). When considering each set of exposure frequencies, for the merged groups of SD exposure with different sets S and D frequencies, the response to D is greater than that of S before ECO (p<0.01 20/14 kHz and p<0.07 12/17 kHz) and that of D is

less than that of S after ECO (p<0.01 14/10 kHz and p<0.01 12/17 kHz). Further, when
normalizing the response rates by the mean response at BF of other frequencies (>1/2
octave away from the exposed frequencies) we find that in the SD exposure the response to
D of neurons with D as BF is significantly higher before ECO and is significantly lower after
ECO (Fig. 5E, barplots, unpaired t-test, P11-15 standard vs deviant: p<0.05, df=107; P6-10
standard vs deviant: p<0.05, df=120). Response to S in SD exposure before ECO is lower than
1 with p<0.051 and no effect is seen in S after ECO. Response to S at S as BF is not
significantly different from 1 in the SS exposure before or after ECO. Thus there is a
differential long term plasticity of responses to the low probability sound frequency
depending on the period of development during which animals are exposed in the SD
environment. It coincides with the differential deviant selectivity before and after ECO
observed in the ACX L4 and SPNs involved early in development, in setting up the ACX
circuitry.

Model binary network with observed deviant selectivity of SPNs and L4 shows stronger
representation of deviant frequency before ECO and standard frequency after ECO

To explain the observed exposure dependent developmental plasticity based on L4 and SPN
deviant selectivity we use a binary network model (Supplementary Material, Fig. 6A) of early
developing TC circuitry of the ACX[18] similar to that considered in the visual cortex[11].
Thalamic inputs selective for either of two inputs, S or D (green and red, Fig. 6A) are
considered to make synapses on an SPN, sending collateral inputs to a L4 neuron[8]. The SP
and L4 neurons are modelled as integrate and fire neurons with absolute and relative
refractory[11]. The SPN neuron makes a synapse onto the L4 neuron as known[9] (Fig. S17). All
synapses are considered depressing in nature[41,42](Fig. 6B) such that L4 and SPN responses to
oddball paradigm show similar adaptation observed early in development (Fig. 1-2, Fig. S8).
Also, all synapses except the thalamic inputs to SPN are plastic changing their strengths with
an asymmetric Hebbian spike timing dependent plasticity (STDP) rule (Fig. 6C). We consider
the model with two different starting points, one representing the period before ECO and
another after ECO (Fig. 6D & Ei-ii). In the earlier developmental period (before ECO), the SP
CSI was 0.41 and that of L4 was -0.02 and in the later developmental time period SP CSI was
-0.17 and that of L4 was 0.4 (Fig. 6D), mimicking our observations (Figs. 1,2&4). The
differences in the initial point of the network (Supplementary Material) in the two cases are
supported by other data[18] and our observations of larger time constants later in
development (Fig. S8) and increased thalamic synaptic weights on L4 due to normal
development, lower SPN to L4 connectivity (Fig. S17B) and reduced SPN rates through
higher thresholds. We use the same exposure protocols SD and SS for the two periods to
show the nature of how the thalamic inputs to L4 develop (Fig. 6E).

During early development with higher deviant selectivity in SP than in L4, with an SD
exposure, the D thalamic input to L4 strengthens faster than that of the S, which will lead to

strengthening of responses to the D stimulus in the adult (Fig. 5E). Whereas in the same
period with an SS exposure the S thalamic input does not show strengthening at a fast rate
(as D, above) leading to normal representation of the S input (Fig. 5E, bar plot P6-11)
equivalent to S in the SD case. On the other hand later in development with L4 showing
higher deviant selectivity than SP, but higher time constants of response decay with more
responses to the S frequency now, shows that in the SD protocol the S thalamic input
synapse strengthens faster with the D input weakening as the SP has lower drive for D than
before. Further, with the only S stimulus also the S inputs, expectedly strengthens leading to
normal responses to S as observed (Fig. 5E, barplot). Thus higher correlated activity
between the L4 and thalamus introduced by SP during the D stimulus because of the high SP
deviant selectivity in the SD exposure, causes increased responses to the low probability D
stimulus in the adult. The lack of such activity and lowering of deviant selectivity in SP and
more responses to S in the later part of development the SD exposure now shows the
opposite effect.

Sparse coding and information maximization principles imply strengthening of response to
low probability stimuli

While a network model mechanistically addresses the observations, we theoretically
address the observations of early exposure to a low probability stimulus leading to
strengthening of its responses and its weakening in later exposure case. Our observations
and model results are contrary to the usual ideas of long term plasticity - a repeated
presentation of a stimulus in the critical period leads to strengthening of the representation
of that particular stimulus in the long term[1,3,5,7]. Auditory cortical spiking activity is known to
be sparse and sparse coding principles theoretically explain a number of known receptive
field types in the auditory pathway[43–45]. Assuming minimal activity based coding in single
neurons underlying sparse representation in populations and that such representation is
one of the goals of the developing auditory system while maximizing information about the
stimulus in the responses we use an objective function (Eqn. 1) as the desired optimization
performed by activity driven plasticity.

$CI_k = I(R_k, T) - \rho(r_o + x a_k)$ … (1a)
$CI_{total} = \sum_{k=1}^{N} CI_k = \sum_{k=1}^{N} I(R_k, T) - \rho\left(N r_o + x \sum_{k=1}^{N} a_k\right)$ … (1b)
441

In Eq. 1, $I(R_k, T)$ is the mutual information (*MI*) between the stimulus (a random variable *T*
which is either S or D, with probabilities (1-*x*) and *x*, where *x* is small and is considered as the
probability of the deviant stimulus) and the response rate in the k^{th} token is R_k, ρ is a
constant, *N* is the total number of stimulus tokens. CI_k is considered as the constrained
information function at the k^{th} token. The mean ensemble response rates are assumed to be
r_0 and $(r_0 + a_k)$ for S and D at the k^{th} stimulus respectively, and assumed to have a
Gaussian distribution both having variance σ^2. The mathematical proof of the optimization
is provided in the supplementary material. However, the basis is as follows. The mutual

information would increase as the separation between the two rates (a_k) increases.
However since sparse activity requires minimal activity by single neurons to code the two
stimuli, to minimize the summed activity, a must be positive and small. So the response to
the deviant needs to be larger than that of the standard. However, the relative weightage of
MI and the summed activity (ρ) can allow a to be increasing or decreasing and still cause CI_k
to be increasing.

We show that the condition for which CI_{total} keeps increasing by considering $\Delta CI = CI_n -$
CI_m ($n>m$) with the assumption that the sequence $\{a_k\}$ is monotonic is

$$\Delta CI = CI_n - CI_m = x\left[\frac{a_n{}^2 - a_m{}^2}{2\sigma^2} + \Lambda(a_n) - \Lambda(a_m) - \rho\{a_n - a_m\}\right]$$

458
$$= x(a_n - a_m)\left[\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} - \rho\right] \dots (2)$$

From the last expression in Eq. 2, under 2 different ranges of ρ, ΔCI is increasing with a
increasing or a decreasing respectively. The theoretical analysis suggests that before and
after ECO the relative importance of coding or discrimination and minimal activity is
changed. Based on the above, sparseness in later stages (after ECO), when spike rates are
higher, needs to be weighted more than earlier stages (before ECO) to explain the
differential effect observed before and after ECO in the response to the low probability
stimulus.

Discussion

Auditory critical period has long been studied using a single tone presentation throughout
the critical period of P12-P15, starting with low threshold hearing coinciding with ECO[15,46].
The frequency over-representation observed with such exposures have been shown to be
absent in exposure periods other than the 'critical period'. Multiple studies have also shown
possible mechanisms of closure of such a critical period[4,47–49], however, the starting of
critical period in spite of those studies, remain unclear. In our results we show that
responses in the mouse ACX exist before ECO as in another study[20] in equivalent ages in the
ferret. Further, we find that a key element in the development of thalamocortical circuitry,
the SPNs, have unique response properties, showing high selectivity to deviant stimuli. Since
spiking in SPNs drive spiking in the thalamo-recipient L4 neurons, developmentally, the
stimulus statistics itself is able to guide thalamocortical potentiation, through a pre-
attentive mechanism. The results of exposure experiments show that relative to a
repeatedly occurring standard frequency, before ECO, representation of a low probability
deviant stimulus is strengthened more. Thus low probability salient stimuli before ECO at
ages equivalent to human gestational week 25[28] cause long term plasticity. It potentially
explains observations in humans of maternal voice recognition in infants[50,51]. The results

also suggest that prenatal sensory environment, can influence cortical development and
may also generalize to other sensory systems given the unique response properties of the
SPNs. Although our results are derived based on tones as stimuli, the possibility of coding of
auditory objects by the neurons in the ACX can also be addressed through similar
experiments. For example, specific response selectivity to species specific vocalizations in
mice and other species[52,53] could also be explained based on the low probability salient
occurrence of such sounds during early development. Corresponding experiments for other
sensory systems can lead to better understanding of cortical development.

Our model results, with two inputs explain the observations and an extension to a whole
network can also produce similar results. The model currently does not include inhibition,
an element crucial in development, as synaptic plasticity rules of inputs/outputs of
inhibitory neurons at such ages are not available. It is likely that it plays a crucial role as the
maturation of inhibition is controlled by SPNs[11]. Also effect of inhibitory switch, occurring
before ECO, on the activity driven plasticity observed before ECO, needs to be investigated.
The closure of critical period, around P16 also coincides with the observed near absence of
SPNs in mice from P15. Thus mechanisms implicated in critical period closure like inhibition,
adenosine, cholinergic inputs[4,47–49], could also be tied to SPNs since SPNs guide
development of subcortical projections. Finally, exposures with selective inhibition of SPN
activity during relevant periods of development are required to establish the SPN driven
cortical development further.

Methods

All animal experiments were approved by Institutional Animal Ethics Committee
(IAEC) of Indian Institute of Technology Kharagpur. Detailed methods are provided in the
Supplementary Materials. C57BL6/J strain mice (National Institute of Nutrition, Hyderabad,
India or Jax Labs) across different developmental time windows or in young adults (P29-P48)
were deeply anesthetized for surgery (2-2.5% isoflurane) and prepared by making a
craniotomy (3-4 mm diameter, based on stereotaxic co-ordinates and landmarks) for in-vivo
electrophysiological recordings from A1/AAF or MGB, with 16 channel electrodes
(Microprobes for Lifesciences) under isoflurane (0.5-1%) in a sound chamber (IAC Acoustics).
LFP and spiking responses to sounds (played through TDT free-field speakers ES1,
attenuated by PA5 and generated with TDT RX6) were obtained with a unity gain headstage
and preamps (Plexon Inc) and stored using National Instruments DAQ card. Sound
generation, data acquisition and online/offline analysis were done with custom written
MATLAB routines. Anatomical labelling and confirmation of recordings from A1/AAF and
from L4 or SP was done using multiple approaches (lipophilic tracer injections in recording
site, nissil sections and IHC). Exposure in specific auditory environments of mouse litters for
the different time windows were done in another sound chamber (IAC Acoustics)
maintaining a 8hr-16hr light dark cycle.

     Figure Legends

Figure 1. Deviant selectivity of SP and L4 reduces with age in A1 with tone tokens. A-C) Spike
dot rasters and PSTHs of 3 neurons (2 in L4 and 1 in SP identified in title) in the developing
A1; i: pure tone rasters followed by PSTHs of chosen SD and frequencies, dark and light gray
bars in rasters; ii: pair of dot rasters and PSTHs to oddball presentations (SD and swap); iii:
time expanded PSTH at onset and deviant in ii. D) Dot rasters of response to tones of
example SP and L4 recorded simultaneously. E) Latencies of L4 and SP neurons recorded
simultaneously in P12-13 and P14-15 age groups showing lowering of latencies with age and
SPNs having lower latency than L4 neurons. F-G) Bar plots of mean deviant selectivity
measured by CSI(F) and CSI respectively for different f_1-f_2 values for L4 (dark) and SP (light).
H-I) Overall mean deviant selectivity for tone stimuli measured by CSI(F) and CSI respectively
as a function of age.

Figure 2. Deviant selectivity of SP and L4 reduces with age in A1 with tone and noise tokens.
A-B) Pairs of PSTHs to oddball presentations (SD and swap) for 6 representative L4 neurons
(A) of different ages (identified in the titles) and that of 3 SPNs (B). C-D) Overall mean
deviant selectivity for noise and tone tokens measured by CSI(F) and CSI respectively as a
function of age.

Figure 3. Neurons in the mouse ACX and IC respond to sounds before ECO. A) Wide field
fluorescence imaging based significant response (p<0.05, paired ttest, 40 repetitions)
regions (white contours) in 3 different ages before ECO and P13, after ECO. Smoothed df/f
traces with SEMs of selected regions (square) responding to selected frequencies (color) are
shown to the right of each image. The black bar represents the stimulus duration. B) As in A,
widefield fluorescence responses before ECO in the IC at P9 and P10. C) i) Mean df/f (40
repetitions) of L2/3 neurons (each row) in response to a single frequency tone (before and
after, P12-13, ECO). White line shows stimulus onset. ii) Population mean df/f of the age
groups (color) of neurons shown in Fig. Ci. iii) Box plot showing response reliability of all
L2/3 neurons imaged at the different ages. Straight line shows linear fit between %
reliability and age with the p value of correlation between them. D) i-ii) Extended 2-photon
Ca^{2+} imaging data showing mean df/f (40 repetitions) of single neurons (each row) and
population mean df/f (below) at two ages before ECO, showing long (>2s) spontaneous
window. E) Single neuron mean df/f and population mean df/f of the same age groups in
response to sound (i) and the same neurons without sound stimulation (ii), with 40
repetitions in each case. F) i) z-scores of single unit responses to tones at different age
groups before ECO in L4 and SP. z-score was computed in each 50 ms bin for each case (each
row is 1 unit in response to 1 frequency tone, 50 ms long, 10-30 repetitions). Top set are
units from L4 bottom set are for SPNs. The white vertical line represents stimulus onset. ii)
Population mean PSTHs of all the recorded units across different age groups which had a z-

score>1 in at least 1 time bin, with respective p-values (paired t-test). The black lines represent the stimulus onset.

Figure 4. SPNs unlike L4 neurons show deviant selectivity before ECO. A) i) Dot rasters and PSTHs of 3 L4 neurons (upper 4 rows, SD and Swap) and 3 SP neurons (lower 4 rows, SD and Swap) are shown in the first 3 columns (Insets: Raster for deviant responses and peaks). ii) Mean population PSTHs of all L4 and SP neurons (upper and lower two plots respectively) to SD (30 reps, 3.3 Hz) and its swap. iii) Same as Aii, for responses to SD (20 reps, 4Hz). B) i-iii) Population (3.3 and 4 Hz combined) histograms of CSI(F), CSI and CSI(P) for SP and L4 (light and dark gray bars respectively) show significantly higher deviant selectivity in SP. iv-v) Similar histograms with mean CSI calculated for each of 1000 random pair of SD response windows selections for S response and D response either before deviant (iv) or after deviant (v).

Figure 5. Exposure to an SD stimulus environment before ECO causes long term plasticity of responses to the D sound and not after ECO. A) Different exposure paradigms (EX0-6) by age and exposure stimulus are represented. B) Cartoon of different tuning curves and construction of profile of response to RF as a function of BF re RF. C) Boxplots for all significant rate responses in all tuning curves collected in the different EX groups. D) CDF plots of response rate distributions of different RFs considered in the different cases. Thin gray line shows mean an STD of CDFs of 13 such CDFs of control data with each of the 13 frequencies considered as RF. Thick gray lines show boundaries of the control CDFs. E) Profile of population mean response rates for 4 exposure cases (EX2&5, S,D and SS in i and EX1&6, S,D and SS in ii), and maximum boundary of 13 response profiles of control data shown in thick gray. Inset bar plot summarizes normalized (by mean response at BF to other frequencies >1/2 octave away from RF) population mean response to RF for neurons with BF at RF of all 6 EX cases (2 bars each, red and green for SD exposure cases and 1 blue for SS).

Figure 6. A binary network model of developing ACX with SP and TC inputs can explain observations of early exposure driven plasticity. A) Cartoon of network of SP and L4 neurons with 2 TC inputs to both SPN and L4, one selective for S and one for D. SPN projects to L4 (gray synapse). B) All synapses were depressing in nature with the example EPSP profile shown. C) The asymmetric STDP learning rule followed by all synapses on the L4 neuron is shown. LTD window was longer and modifications occurred over -80 to 40 ms of post–pre spike times. D) Sample PSTHs in earliest part of development (representing before ECO) shows high deviant selectivity in SP and low in L4 in the model neurons. E) Temporal evolution of the relevant synaptic strengths in the two exposure periods (i: Before ECO and ii: After ECO) for simulations of exposure with SD or SS protocol.

References

1. Schreiner, C. E. & Polley, D. B. Auditory map plasticity: Diversity in causes and consequences. *Current Opinion in Neurobiology* vol. 24 143–156 (2014).

2. Hensch, T. K. Critical period plasticity in local cortical circuits. *Nature Reviews Neuroscience* (2005) doi:10.1038/nrn1787.

3. de Villers-Sidani, E., Chang, E. F., Bao, S. & Merzenich, M. M. Critical Period Window for Spectral Tuning Defined in the Primary Auditory Cortex (A1) in the Rat. *J. Neurosci.* (2007) doi:10.1523/JNEUROSCI.3227-06.2007.

4. Takesian, A. E., Bogart, L. J., Lichtman, J. W. & Hensch, T. K. Inhibitory circuit gating of auditory critical-period plasticity. *Nat. Neurosci.* (2018) doi:10.1038/s41593-017-0064-2.

5. Barkat, T. R., Polley, D. B. & Hensch, T. K. A critical period for auditory thalamocortical connectivity. *Nat. Neurosci.* (2011) doi:10.1038/nn.2882.

6. Zhou, X. & Merzenich, M. M. Enduring effects of early structured noise exposure on temporal modulation in the primary auditory cortex. *Proc. Natl. Acad. Sci. U. S. A.* (2008) doi:10.1073/pnas.0800009105.

7. Zhang, L. I., Bao, S. & Merzenich, M. M. Persistent and specific influences of early acoustic environments on primary auditory cortex. *Nat. Neurosci.* **4**, 1123–1130 (2001).

8. Luhmann, H. J., Kirischuk, S. & Kilb, W. The Superior Function of the Subplate in Early Neocortical Development. *Front. Neuroanat.* **12**, (2018).

9. Kanold, P. O. & Luhmann, H. J. The Subplate and Early Cortical Circuits. *Annu. Rev. Neurosci.* (2010) doi:10.1146/annurev-neuro-060909-153244.

10. Kanold, P. O., Kara, P., Reid, R. C. & Shatz, C. J. Role of subplate neurons in functional maturation of visual cortical columns. *Science (80-.).* (2003) doi:10.1126/science.1084152.

11. Kanold, P. O. & Shatz, C. J. Subplate Neurons Regulate Maturation of Cortical Inhibition and Outcome of Ocular Dominance Plasticity. *Neuron* (2006) doi:10.1016/j.neuron.2006.07.008.

12. Ghosh, A. & Shatz, C. J. Involvement of subplate neurons in the formation of ocular dominance columns. *Science (80-.).* (1992) doi:10.1126/science.1542795.

13. Ghosh, A., Antonini, A., McConnell, S. K. & Shatz, C. J. Requirement for subplate neurons in the formation of thalamocortical connections. *Nature* (1990) doi:10.1038/347179a0.

14. McConnell, S. K., Ghosh, A. & Shatz, C. J. Subplate neurons pioneer the first axon pathway from the cerebral cortex. *Science (80-.).* (1989)

doi:10.1126/science.2475909.

15.   Chang, E. F., Bao, S., Imaizumi, K., Schreiner, C. E. & Merzenich, M. M. Development
of spectral and temporal response selectivity in the auditory cortex. *Proc. Natl. Acad.*
Sci. U. S. A. (2005) doi:10.1073/pnas.0508239102.

16.   Ulanovsky, N., Las, L. & Nelken, I. Processing of low-probability sounds by cortical
neurons. *Nat. Neurosci.* (2003) doi:10.1038/nn1032.

17.   Ghosh, A. & Shatz, C. J. A role for subplate neurons in the patterning of connections
from thalamus to neocortex. *Development* (1993).

18.   Zhao, C., Kao, J. P. Y. & Kanold, P. O. Functional Excitatory Microcircuits in Neonatal
Cortex Connect Thalamus and Layer 4. *J. Neurosci.* (2009) doi:10.1037/pha0000116.

19.   Viswanathan, S., Bandyopadhyay, S., Kao, J. P. Y. & Kanold, P. O. Changing
Microcircuits in the Subplate of the Developing Cortex. *J. Neurosci.* **32**, 1589–1601
(2012).

20.   Wess, J. M., Isaiah, A., Watkins, P. V & Kanold, P. O. Subplate neurons are the first
cortical neurons to respond to sensory stimuli. *Proc. Natl. Acad. Sci.* (2017)
doi:10.1073/pnas.1710793114.

21.   Pérez-González, D. & Malmierca, M. S. Adaptation in the auditory system: an
overview. *Front. Integr. Neurosci.* (2014) doi:10.3389/fnint.2014.00019.

22.   Nieto-Diego, J. & Malmierca, M. S. Topographic Distribution of Stimulus-Specific
Adaptation across Auditory Cortical Fields in the Anesthetized Rat. *PLoS Biol.* (2016)
doi:10.1371/journal.pbio.1002397.

23.   Bandyopadhyay, S., Shamma, S. A. & Kanold, P. O. Dichotomy of functional
organization   in   the   mouse   auditory   cortex.   *Nat.   Neurosci.*   (2010)
doi:10.1038/nn.2490.

24.   Issa, J. B. *et al.* Multiscale Optical Ca2+Imaging of Tonal Organization in Mouse
Auditory Cortex. *Neuron* (2014) doi:10.1016/j.neuron.2014.07.009.

25.   Sawatari, H. *et al.* Identification and characterization of an insular auditory field in
mice. *Eur. J. Neurosci.* (2011) doi:10.1111/j.1460-9568.2011.07926.x.

26.   Barbour, D. L. & Wang, X. Contrast tuning in auditory cortex. *Science (80-.).* (2003)
doi:10.1126/science.1080425.

27.   Mill, R., Coath, M., Wennekers, T. & Denham, S. L. Abstract stimulus-specific
adaptation models. *Neural Computation* (2011) doi:10.1162/NECO_a_00077.

28.   Clancy, B., Darlington, R. B. & Finlay, B. L. Translating developmental time across
mammalian species. *Neuroscience* (2001) doi:10.1016/S0306-4522(01)00171-3.

29.   Chang, A. *et al.* Specific Influences of Early Acoustic Environments on Cochlear Hair
Cells in Postnatal Mice. *Neural Plast.* (2018) doi:10.1155/2018/5616930.

30. Babola, T. A. *et al.* Homeostatic Control of Spontaneous Activity in the Developing
Auditory System. *Neuron* (2018) doi:10.1016/j.neuron.2018.07.004.

31. Anthwal, N. & Thompson, H. The development of the mammalian outer and middle
ear. *J. Anat.* (2016) doi:10.1111/joa.12344.

32. Huangfu, M. & Saunders, J. C. Auditory development in the mouse: Structural
maturation of the middle ear. *J. Morphol.* (1983) doi:10.1002/jmor.1051760302.

33. Kim, K. X. & Fettiplace, R. Developmental changes in the cochlear hair cell
mechanotransducer channel and their regulation by transmembrane channel-like
proteins. *J. Gen. Physiol.* (2013) doi:10.1085/jgp.201210913.

34. Lelli, A., Asai, Y., Forge, A., Holt, J. R. & Géléoc, G. S. G. Tonotopic gradient in the
developmental acquisition of sensory transduction in outer hair cells of the mouse
cochlea. *J. Neurophysiol.* (2009) doi:10.1152/jn.00136.2009.

35. Abe, T. *et al.* Developmental expression of the outer hair cell motor prestin in the
mouse. *J. Membr. Biol.* (2007) doi:10.1007/s00232-007-9004-5.

36. Polley, D. B., Thompson, J. H. & Guo, W. Brief hearing loss disrupts binaural
integration during two early critical periods of auditory cortex development. *Nat.*
Commun. (2013) doi:10.1038/ncomms3547.

37. Ruebsamen, R. & Lippe, W. The development of cochlear function. in *Development of*
the auditory system (eds. Rubel, E., Poppe, A. & Fay, R.) 193–270 (Springer, 1997).

38. Shnerson, A. & Pujol, A. Development: anatomy, electrophysiology, and behavior. in
The auditory psychobiology of the mouse (ed. Willot, J.) 395–425 (Charles C. Thomas,
1983).

39. Pujol, R., Lavigne-Rebillard, M. & Lenoir, M. Development of sensory and neural
structures in the mammalian cochlea. in *Development of the auditory system* (eds.
Rubel, E., Popper, A. & Fay, R.) 146–192 (Spinger, 1997).

40. Tritsch, N. X. *et al.* Calcium action potentials in hair cells pattern auditory neuron
activity before hearing onset. *Nat. Neurosci.* (2010) doi:10.1038/nn.2604.

41. Mill, R., Coath, M., Wennekers, T. & Denham, S. L. A neurocomputational model of
stimulus-specific adaptation to oddball and markov sequences. *PLoS Comput. Biol.*
(2011) doi:10.1371/journal.pcbi.1002117.

42. Yarden, T. S. & Nelken, I. Stimulus-specific adaptation in a recurrent network model
of   primary   auditory   cortex.   *PLoS   Comput.   Biol.*   (2017)
doi:10.1371/journal.pcbi.1005437.

43. Hromádka, T., DeWeese, M. R. & Zador, A. M. Sparse representation of sounds in the
unanesthetized auditory cortex. *PLoS Biol.* (2008) doi:10.1371/journal.pbio.0060016.

44. Lewicki, M. S. Efficient coding of natural sounds. *Nat. Neurosci.* (2002)
doi:10.1038/nn831.

45. Smith, E. C. & Lewicki, M. S. Efficient auditory coding. *Nature* (2006)
doi:10.1038/nature04485.

46. Sun, Y. J. *et al.* Fine-tuning of pre-balanced excitation and inhibition during auditory
cortical development. *Nature* **465**, 927–931 (2010).

47. Blundon, J. A. *et al.* Restoring auditory cortex plasticity in adult mice by restricting
thalamic adenosine signaling. *Science (80-.).* (2017) doi:10.1126/science.aaf4612.

48. Vickers, E. D. *et al.* Parvalbumin-Interneuron Output Synapses Show Spike-Timing-
Dependent Plasticity that Contributes to Auditory Map Remodeling. *Neuron* (2018)
doi:10.1016/j.neuron.2018.07.018.

49. Courtney E. Yaeger, Dario Ringach, J. T. T. Neuromodulatory control of localized
dendritic spiking in critical period cortex. *Nature* **567**, 100–104 (2019).

50. Mehler, J. *et al.* A precursor of language acquisition in young infants. *Cognition* (1988)
doi:10.1016/0010-0277(88)90035-2.

51. Voegtline, K. M., Costigan, K. A., Pater, H. A. & DiPietro, J. A. Near-term fetal response
to   maternal   spoken   voice.   *Infant   Behav.   Dev.*   (2013)
doi:10.1016/j.infbeh.2013.05.002.

52. Portfors, C. V., Roberts, P. D. & Jonson, K. Over-representation of species-specific
vocalizations   in   the   awake   mouse   inferior   colliculus.   *Neuroscience*   (2009)
doi:10.1016/j.neuroscience.2009.04.056.

53. Wang, X. & Kadia, S. C. Differential Representation of Species-Specific Primate
Vocalizations in the Auditory Cortices of Marmoset and Cat. *J. Neurophysiol.* (2001)
doi:10.1152/jn.2001.86.5.2616.

54. Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic
Coordinates*. *São Paulo, Academic Press* (2012). doi:10.1364/OE.20.020998.

55. Torii, M. & Levitt, P. Dissociation of corticothalamic and thalamocortical axon
targeting   by   an   EphA7-mediated   mechanism.   *Neuron*   (2005)
doi:10.1016/j.neuron.2005.09.021.

56. Osheroff, H. & Hatten, M. E. Gene expression profiling of preplate neurons destined
for the subplate: Genes involved in transcription, axon extension, neurotransmitter
regulation, steroid hormone signaling, and neuronal survival. *Cereb. Cortex* (2009)
doi:10.1093/cercor/bhp034.

57. Hoerder-Suabedissen, A. *et al.* Novel markers reveal subpopulations of subplate
neurons   in   the   murine   cerebral   cortex.   *Cereb.   Cortex*   (2009)
doi:10.1093/cercor/bhn195.

58. Gee, J. M. *et al.* Imaging Activity in Neurons and Glia with a Polr2a-Based and Cre-
Dependent   GCaMP5G-IRES-tdTomato   Reporter   Mouse.   *Neuron*   (2014)
doi:10.1016/j.neuron.2014.07.024.

  Figures

Figure 1. Deviant selectivity of SP and L4 reduces with age in A1 with tone tokens. A-C) Spike dot rasters and PSTHs of 3 neurons (2 in L4 and 1 in SP identified in title) in the developing A1; i: pure tone rasters followed by PSTHs of chosen SD and frequencies, dark and light gray bars in rasters; ii: pair of dot rasters and PSTHs to oddball presentations (SD and swap); iii: time expanded PSTH at onset and deviant in ii. D) Dot rasters of response to tones of example SP and L4 recorded simultaneously. E) Latencies of L4 and SP neurons recorded simultaneously in P12-13 and P14-15 age groups showing lowering of latencies with age and SPNs having lower latency than L4 neurons. F-G) Bar plots of mean deviant selectivity measured by CSI(F) and CSI respectively for different f_1-f_2 values for L4 (dark) and SP (light). H-I) Overall mean deviant selectivity for tone stimuli measured by CSI(F) and CSI respectively as a function of age.

757 **Figure 2.** Deviant selectivity of SP and L4 reduces with age in A1 with tone and noise tokens.
758 A-B) Pairs of PSTHs to oddball presentations (SD and swap) for 6 representative L4 neurons
759 (A) of different ages (identified in the titles) and that of 3 SPNs (B). C-D) Overall mean
760 deviant selectivity for noise and tone tokens measured by CSI(F) and CSI respectively as a
761 function of age.

Figure 3. Neurons in the mouse ACX and IC respond to sounds before ECO. A) Wide field fluorescence imaging based significant response (p<0.05, paired ttest, 40 repetitions) regions (white contours) in 3 different ages before ECO and P13, after ECO. Smoothed df/f

traces with SEMs of selected regions (square) responding to selected frequencies (color) are shown to the right of each image. The black bar represents the stimulus duration. B) As in A, widefield fluorescence responses before ECO in the IC at P9 and P10. C) i) Mean df/f (40 repetitions) of L2/3 neurons (each row) in response to a single frequency tone (before and after, P12-13, ECO). White line shows stimulus onset. ii) Population mean df/f of the age groups (color) of neurons shown in Fig. Ci. iii) Box plot showing response reliability of all L2/3 neurons imaged at the different ages. Straight line shows linear fit between % reliability and age with the p value of correlation between them. D) i-ii) Extended 2-photon Ca^{2+} imaging data showing mean df/f (40 repetitions) of single neurons (each row) and population mean df/f (below) at two ages before ECO, showing long (>2s) spontaneous window. E) Single neuron mean df/f and population mean df/f of the same age groups in response to sound (i) and the same neurons without sound stimulation (ii), with 40 repetitions in each case. F) i) z-scores of single unit responses to tones at different age groups before ECO in L4 and SP. z-score was computed in each 50 ms bin for each case (each row is 1 unit in response to 1 frequency tone, 50 ms long, 10-30 repetitions). Top set are units from L4 bottom set are for SPNs. The white vertical line represents stimulus onset. ii) Population mean PSTHs of all the recorded units across different age groups which had a z-score>1 in at least 1 time bin, with respective p-values (paired t-test). The black lines represent the stimulus onset.

Figure 4. SPNs unlike L4 neurons show deviant selectivity before ECO. A) i) Dot rasters and PSTHs of 3 L4 neurons (upper 4 rows, SD and Swap) and 3 SP neurons (lower 4 rows, SD and Swap) are shown in the first 3 columns (Insets: Raster for deviant responses and peaks). ii) Mean population PSTHs of all L4 and SP neurons (upper and lower two plots respectively) to SD (30 reps, 3.3 Hz) and its swap. iii) Same as Aii, for responses to SD (20 reps, 4Hz). B) i-iii) Population (3.3 and 4 Hz combined) histograms of CSI(F), CSI and CSI(P) for SP and L4 (light and dark gray bars respectively) show significantly higher deviant selectivity in SP. iv-v) Similar histograms with mean CSI calculated for each of 1000 random pair of SD response windows selections for S response and D response either before deviant (iv) or after deviant (v).

Figure 5. Exposure to an SD stimulus environment before ECO causes long term plasticity of responses to the D sound and not after ECO. A) Different exposure paradigms (EX0-6) by age and exposure stimulus are represented. B) Cartoon of different tuning curves and construction of profile of response to RF as a function of BF re RF. C) Boxplots for all significant rate responses in all tuning curves collected in the different EX groups. D) CDF plots of response rate distributions of different RFs considered in the different cases. Thin gray line shows mean an STD of CDFs of 13 such CDFs of control data with each of the 13 frequencies considered as RF. Thick gray lines show boundaries of the control CDFs. E) Profile of population mean response rates for 4 exposure cases (EX2&5, S,D and SS in i and EX1&6, S,D and SS in ii), and maximum boundary of 13 response profiles of control data shown in thick gray. Inset bar plot summarizes normalized (by mean response at BF to other frequencies >1/2 octave away from RF) population mean response to RF for neurons with BF at RF of all 6 EX cases (2 bars each, red and green for SD exposure cases and 1 blue for SS).

Figure 6. A binary network model of developing ACX with SP and TC inputs can explain observations of early exposure driven plasticity. A) Cartoon of network of SP and L4 neurons with 2 TC inputs to both SPN and L4, one selective for S and one for D. SPN projects to L4 (gray synapse). B) All synapses were depressing in nature with the example EPSP profile shown. C) The asymmetric STDP learning rule followed by all synapses on the L4 neuron is shown. LTD window was longer and modifications occurred over -80 to 40 ms of post–pre spike times. D) Sample PSTHs in earliest part of development (representing before ECO) shows high deviant selectivity in SP and low in L4 in the model neurons. E) Temporal evolution of the relevant synaptic strengths in the two exposure periods (i: Before ECO and ii: After ECO) for simulations of exposure with SD or SS protocol.

 Supplementary Material

Detailed Materials and Methods

Subjects

All animal experiments were approved by Institutional Animal Ethics Committee
(IAEC) of Indian Institute of Technology Kharagpur.

In-vivo electrophysiological recordings were performed on C57BL6/J strain mice
(National Institute of Nutrition, Hyderabad, India or Jax Labs) across different
developmental time windows in two different contexts, tone-tone (f_1-f_2) pairs (P12-P13, n=3
mice, 28 L4 units, 9 SPN units, 47 L4 cases, 14 SPN cases, ; P14-P15, n=3 mice , 41 L4 units,
34 SPN units, 63 L4 cases, 51 SPN cases; P16-P21, n=3 mice, 11 units, 15 cases; P22-P28,
n=4 mice, 24 units, 53 cases; >P28, n=3 mice, 26 units, 74 cases) and Noise-tone (N-f) pairs
(P12-P13, n=10 mice, 45 L4 units, 54 SPN units, 57 L4 cases, 101 SPN cases ; P14-P15, n=8
mice, 66 L4 units, 59 SPN units, 115 L4 cases, 67 SPN cases; P16-21, n=4 mice, 20 units, 31
cases; P22-P28, n=6 mice, 38 units, 97 cases ; >P28, 7 mice, 44 units, 63 cases). For
experiments in mice before ECO single unit or LFP responses with 5 stimulus repetitions for
tone and noise we recorded in 9 mice (38 units: P08=1; P09, n=2; P10, n=2; P11, n=3; P12,
n=1). In an additional 8mice, single unit responses to tones, noise (10-30 repetitions) and SD
paradigm with 30 repetitions were obtained. Over all before ECO n=15animals were
included to have responses (P08, n=2; P09, n=4; P10, n=4; P11, n=4; P12, n=1)) out of 38
animals tried. For recording L2/3 responses, 2-photon Ca^{+2} imaging was performed on Thy1-
Gcamp6f (n=12, P07=2; P08=1; P09=3; P10=3; P11=1; P12=1; P13=1) mice. In a subset of the
above animals (n=9, P07=1; P08=1; P09=3; P10=3; P13=1) wide field fluorescence imaging
was performed.

Anatomical labelling

In order to guide in-vivo electrophysiology and identify SP layer based on
morphology and cytoarchitecture of neurons, Nissl staining was performed at different age
groups. Mice of ages P6 (n=2 mice), P13 (n=2 mice), P16 (n=2 mice) and P18 (n=2 mice) were
initially perfused and the brains were harvested. Since ear canal opening is after P11 and as
the cortical size stabilizes by P21, the above ages were chosen initially for the
characterization. Nissl stains were performed after obtaining coronal sections of the
auditory regions based on mouse atlas[54]. For further confirmation, immunohistochemistry
was performed on brain slices across different developmental time points (P06-P07, n=2
mice; P08-P11, n=3 mice, P12-13, n=5mice; P14-P16, n=2mice; P17-P19, n=2 mice; P20-28,
N=3 mice; >P28, n=2 mice) with known markers of SP which are Complexin 3 (Cplx3);
Synaptic Systems (122302) and Connective Tissue Derived Growth Factor (CTGF); Santa Cruz
(sc-14939). Paraformaldehyde (PFA) fixed brain tissue was embedded in a 3% agarose
block. Brain embedded block was fixed on a vibrotome (Leica) sectioning platform. 50

micron thin sections of auditory slices were cut, blocked and permeabilized in a buffer
containing, 1% BSA, 0.3% Triton X and 10% Normal Donkey Serum (for CTGF) or 10% Normal
Goat Serum (for Cplx3). Sections were incubated in primary antibodies (Cplx3: 1:500 and
CTGF: 1:500) overnight and followed by washing with 1X PBS five times. Thereafter, sections
were incubated for two hours in Secondary antibodies (Cplx3: anti rabbit 594, Invitrogen
and CTGF: anti goat 488, Invitrogen) diluted in buffer, 1% BSA, 3% serum and 0.3% triton X
at room temperature. Sections were rinsed, mounted and imaged under epifluorescence
micreoscopy. Depth of SP in different regions and at different developmental time scales
was determined from these initial studies and was used as a guide for performing
electrophysiology recordings. Further confirmation about recordings from input layer (Layer
IV) was also obtained from SCNN1-td-tomato (cross of SCNN1-cre and floxed-td-tomato: JAX
mice) mice, which have SCNN1 expressing neurons (specific to Layer IV) labelled, allowing
confirmation of recordings in Layer IV. Sections were imaged using epi-fluorescence
microscopy (Xcite illumination, Bruker) and analysed for cytoarchitecture using custom
written codes in MATLAB.

In-vivo extracellular single unit recordings

Initially animals were anaesthetised with deep anaesthesia (5% Isoflurane) inside the
induction chamber. Initial Induction aids in maintaining anaesthesia level in animals even
after bringing down the anaesthesia levels (1.5-2%) during surgery. Core body temperature
was maintained constantly at $37^{o}C$ with a heating pad, throughout the experiment. A central
incision was made and the left temporal portion of the skull was exposed. Exposed left
temporalis was cleaned with application of alcohol. Thereafter, tissue clearance was done
by rubbing the skull with 3% of Hydrogen peroxide. A 5mm diameter region was marked on
the left hemisphere. Skull was firmly fixed after keeping the mark at the centre of a head
plate. Craniotomy was performed over the estimated auditory cortical area. After
performing craniotomy, experimental animal was shifted immediately to a sound proof
chamber (IAC Acoustics). The depth of anaesthesia was monitored constantly throughout
the experiment with paw pinches. During recording sessions anaesthesia was brought down
to a concentration of 0.5-1%, in order to get robust responses from primary auditory cortex.

Extracellular recordings were performed from the left auditory cortex using tungsten
Micro Electrodes Array (MEA) of impedance 3-5 Mega ohms (MicroProbes, USA). A 4X4
custom designed metal MEA with inter electrode spacing of 125 microns was lowered into
the auditory cortex using a micromanipulator (MP-285, Sutter Instrument Company,
Novato, CA). We also performed recording using arrays separated by 400 or 600 microns in
length to record across laminae simultaneously (for example, Layer IV and SP, Fig. S1).
Signals were acquired after passing through a unity gain headstage (Plexon, HST16o25)
followed by PBX3 (Plexon) preamp with gain of X1000, to obtain wideband signal (used to
extract LFP, 0.7 Hz to 6 kHz) and spike signals (150 Hz to 8 kHz) in parallel and acquired

through National Instruments Data Acquisition Card (NI-PCI-6259) at 20 KHz sampling rate, controlled through custom written MATLAB (Mathworks, Natick, USA) routines. Further, all online/offline analysis was performed using custom written MATLAB routines. To obtain laminar profile of responses from primary auditory cortex (A1), multiple depths were covered by advancing the electrodes in depth at 100-150 micron steps.

Since the size and location of A1 over development varies, it is important to confirm location of recording to be A1. For A1 recordings additional physiological information based on tuning – existence of broad tonotopy, across electrodes in the recording array, latency of responses, selectivity to tone stimuli was used to confirm recordings in primary auditory areas (A1 and Anterior Auditory Field (AAF)). Based on location of electrode tracks and comparison with mouse brain atlas[54] and landmarks (location relative to rhinal vein, size of hippocampus in coronal slices) also provides additional confirmation of recording location. In many experiments after recording a DiI crystal[55], a lipophilic tracer, was inserted into the cortex at the recording location. After allowing passive travel of the tracer for 20-30 days at 35^{o}C , brain slices are cut to look for labelled fibres and cell bodies in the ventral division of the MGB (lemniscal structure projecting to A1) allowing confirmation of recordings in A1 Fig. S1C). We further confirmed recordings from electrode location in primary regions subsequently, from broad tonotopy observed (Fig. S1B). The recording depths targeted for L4 and SPN were based on known depths of L4 and SPNs in ACX[19]; also, in a separate groups of mice (P12-P13, n=5; P14-P16, n=3) depth of SPNs was determined to be above 900 μm based on two methods. First by cell morphology in Nissl stains of coronal slices (Fig. S1D) and second by immunostaining coronal slices containing A1 with antibodies of SPN markers[56,57], *Complexin3* and *CTGF* (Fig. S2), at different ages (Fig. S2C) and depth range of L4 from expression of *SCNN1* (B6.Cg-Tg(Scnn1a-cre)3Aibs/J mouse, Jax Labs, 09613 crossed with CAG-tdTomato mouse Jax Labs, 07909), a L4 specific marker[19]. In a subset of mice in which electrophysiological recordings were performed we confirmed electrode recording site to be in SP layer from electrode tracks along with immunostaining as above (Fig. S3A) and electrodes to be in L4 by performing recordings in the *SCNN1-cre*-td-Tomato (P14 and P18) mice (Fig. S3). Depths of other layers (L2/3 and L5/6) were determined based on the above results.

Acute craniotomy for widefield imaging and 2-photon Ca^{2+} imaging

A standard surgical procedure was performed as mentioned above in in-vivo electrophysiology recordings.   Animals were initially anaesthetised under isoflurane (induction at 5% isoflurane followed by 2% isoflurane). After removing the skin and tissue above the skull, a metallic head plate was implanted above the estimated auditory cortical area. Craniotomy of around 3mm diameter was performed over left auditory area. Smaller size craniotomy was preferred, as it gave us the advantage to minimize brain motion. Thereafter, craniotomy was filled with 1.5% low melting-point Agarose and immediately a

5mm coverslip was imbedded over the craniotomy. Animals were then transferred to the 2-photon imaging setup and the level of anaesthesia was reduced down to 0.1-0.2% isoflurane during the recordings.

Auditory stimulation for in-vivo electrophysiology

Stimulus was presented inside sound proof chamber, 10 cm away from the right ear (contralateral) of the mouse, with TDT electrostatic speakers (ES1) driven by TDT drivers ED1 after attenuation by TDT attenuators (PA5) generated through TDT RX6 using custom made software written in MATLAB. The acoustic calibrations, performed with microphone 4939 (Brüel & Kjær, Denmark), of the ES1 speakers (TDT) in the sound chamber, showed a typical flat (+/- 7 dB) calibration curve from 4-60 kHz.

Auditory stimulation for imaging

Stimulus was presented through electrostatic speaker (ES1) placed 2cm away from the right ear of the mouse pups Stimulus was generated through custom written software in MATLAB (Mathworks), passed through TDT RZ6 multifunctional processor. Acoustic calibrations were performed using microphone 4939 (Brüel & Kjær, Denmark) and showed a typical flat (+/- 7 dB) calibration curve from 4-60 kHz. Auditory stimuli consisted of broadband noise (6-48 kHz, 40 repetitions), tones (40 repetitions of respective frequencies), and series of sound token representing SD paradigm (30 repetitions). Only for one of the animals (P12, after ear canal opening) 5 repetitions of each tonal frequency were delivered. Each of the stimuli was interleaved with a gap of 5s. The loudest intensities of sounds at 80-90 dB SPL were presented to ear, in order to allow mechanotransduction at younger ages (before ear canal opening).

Single units spike times were obtained from the acquired spike channel data using threshold crossing and spike sorting with custom written software in MATLAB.

Wide Field Imaging

For wide field calcium imaging, imaging signals were acquired using a 14-bit CoolSNAP HQ2 CCD camera (Photometrics). Images of surface vasculature were acquired using blue LED illumination (470 nm), and wide field GCaMP6f signals were recorded at ~30 Hz using blue illumination (470 nm)[58]. Fluorescence signals were acquired at 4x binning of 1040x1392 pixels, using a 4X objective (Olympus). Based on the auditory landmarks (Medial cerebral artery (MCA) and rhinal vein position) and the responses to different frequencies, the region of interest and the focal plane of the image (150-300 µm) were moved to desired coordinates for performing 2-photon imaging.

Two-Photon Calcium Imaging

Two photon calcium imaging was performed using a commercially available 2-photon microscopy system (Prairie View Technologies). The microscopy was controlled by Prairie View software (Prairie View Technologies). For imaging, excitation beam of wavelength 860nm generated from Insight laser (Spectra Physics, USA) was directed over the exposed brain volume. The laser was delivered through 20X/0.8 NA water immersion objective (2mm WD, Olympus). The laser power was adjusted from 50mW to 80mW depending upon the condition of the specimen. Frames in region of interests (~150 μm x 200 μm with 1.16 μm pixel size) were imaged at ~4-6 Hz (160-250ms frame period, 1.2-4 μs dwell time) with the stimulus presented at usually the 6th (to the 10th) frame in a sequence of 20-30 frames per stimulus.

Exposure protocol

For another set of experiment, animal litters were exposed to sound sequences inside a soundproof chamber with maintained 8-hr light and 16hr dark cycle, continuously for 5 days in 4 age groups (P0-P5/6, P6-P10/11, P11-P15/16 and P16-P21).

Table S1

Age of exposure	Protocol	Litter	Animals	Units	Tuning curves
Control	No exposure	2	3+3	400	815
P6-P11	SD (12kHz-17kHz) SD (20kHz-14kHz)	1+1	5+3	256 197	504 525
P11-P16	SD (12kHz-17kHz) SD (14kHz-10kHz)	1+1	2+3	367 233	700 553
P16-P21	SD (12kHz-17kHz)	1+1	2+2	130	255
P0-P5	SD (12kHz-17kHz)	1+1	2+2	183	370
P11-P16	SS (only 12kHz)	1	3	327	642
P6-P11	SS (only 12kHz)	1	3	194	487

The litters were kept along with the breeders in a cage. Each exposure stimulus was synthesised online and presented through electrostatic speaker, kept 5cm above the cage, using custom written codes on MATLAB. All the age groups were presented with sequence of stimuli having 50 ms sound tokens with a 250 ms inter-token gap at 90 dB SPL. The sequence (SD) had token of two frequencies, a standard (S) having an arrival probability of 90% and a deviant (D) arriving with 10% probability. For only standard exposure (SS), there

was no D frequency. After sound exposure animals were returned immediately to home
cages in animal house and reared up in normal environment until experiments were
performed. During recording sessions, mapping of primary auditory cortex was performed.
Tone pips (50 ms, 5 ms up down ramps, 6-48 kHz, 0.25 octaves difference at 10-50 dB above
threshold, 10 dB increments, 5 repetitions of each frequency-Intensity combination) was
presented to the right ear. Extracellular responses were recorded from thalamo-recipient
layer (Deep Layer III and layer IV) at depth 350-500 microns. Frequency responses area
(FRA) was constructed of each unit for minimum of three sound levels played. Single units
from A1 having response peak latency within 9 ms to 31 ms were included in the analysis.
This window was selected on the basis of population mean and variance of response peak
latencies obtained across all sound levels.

1015

1016 **Data Analysis**

1017

1018 *Identification of Subplate Layer*

1019

1020 SP Layer from nissl staining was based on compartmentalization drawn by cell sparse zone
1021 above and the horizontally oriented cytoarchitecture of the SP layer. Based on above
1022 features manual points (perpendicular to lamina) using MATLAB on slice images were
1023 selected to define depth and width of the SP layer. Similar analysis was performed too on
1024 the immunolabelled brain slice images.

1025

1026 *Rate Calculation*
1027 Neuronal firing rate was computed by calculating mean response within stimulus duration
1028 or token. Only single units that responded (compared to spontaneous activity 400ms
1029 preceding stimulus, 2-tailed t-test, $p<0.05$) to at least one of the stimuli (sound token (*f* or
1030 *N*) as standard or deviant) used were considered for analysis.

1031

Latency Analysis (Fig. 1D)
For each neuron, tuning parameters (Best Frequency (BF), Latency and Tuning width) were
defined using custom written codes in MATLAB. Best frequency for a neuron is the
frequency which elicits maximum response within stimulus duration (50 ms) to a given
frequency at given sound level. Latency to stimulus onset was defined as the time bin (5 ms
resolution) at which peak response was observed for a BF.
Tuning width was computed using the Full width Half Maximum (FWHM). Only single peak
tuning curves were considered for tuning width analysis. A continuous stretch of
frequencies crossing the threshold (half of the maximum amplitude) was considered as the
tuning width of a neuron.

1042

Adaptation Time Constants (Fig. S8)

A time course analysis was performed to fit the response rates for a SSS…..SDS….SS stimuli. Across all age groups and cortical layers, we employed exponential fitting to only those responsive units which have significant responses for tone as standards. To compute exponential decay time constants, the average spike rate in each token before the deviant was fitted with a first degree exponential curve, based on least square error method. Fitting was performed using the 'fit' function in MATLAB.

LFP Analysis (Fig. 3A, Fig. S10)

For LFP analysis, baseline shifting was performed to have mean of baseline 200 ms preceding the stimulus at 0. To obtain LFP, acquired wideband signals were notch filtered (50Hz, Butterworth 8^{th} order, to remove AC supply line noise) and then bandpass filtered (between 1 Hz and 300 Hz, Butterworth 2^{nd} order). LFPs were smoothed further using a Gaussian window, SD 150 ms. Due to the unreliability of responses as observed in the spiking data before ECO, for our population analysis we considered all the LFP recorded for all presented frequencies in order to quantify the response magnitude in terms of population mean amplitude and mean RMS value (across 50ms bin). LFP responses crossing the threshold of 200 mV throughout the recorded time window were excluded from population analysis.

Connectivity between pairs (Fig. S17)

Cross-correlation between pairs of PSTHs (nrep= 20-30 repetitions of SD stimulus and swap, bin size = 2ms) was used to detect connections between simultaneously recorded pairs of neurons (SP-L4 pairs). Bootstrap analysis was done to get variability in the cross-correlograms (MATLAB xcorr function, up to lag of 20 ms) by randomly resampling (same number of repeats from the existing 20-30 repeats) with replacement, 100 times. Thus a mean and SD (dark line with errorbar (Fig. S17A) of the correlograms were obtained. To compare with spurious cross correlations, all repeats from both neurons were mixed to give (2*nrep spike trains) and randomly nrep of them were assigned to each of the neurons and from these PSTHs cross-correlograms were obtained with mean and SD by repeating the process 100 times (gray lie with errorbar, Fig. S17A). Non-overlap of errobars were considered to be significant connections between the pair of neurons. We used 1.65-SD for data in P12 and above groups and 1-SD in before ECO group due to large difference in firing rates of single units in the two cases.

1083 Table S2

Interelectrode distance		Connected pairs /Total pairs SP ——→ L4	Connected pairs /Total pairs L4 ——→ SP
125	Before ECO	6/12	4/12
	P12-P13	27/102	22/102
200-300	Before ECO	3/12	3/12
	P12-P13	16/78	16/78
350-400	Before ECO	0/2	0/2
	P12-P13	5/10	4/10
>400	Before ECO	1/1	1/1
	P12-P13	2/10	4/10

1084

1085 Imaging Analysis

Wide Field imaging analysis was performed using custom codes written in MATLAB
(Mathworks). For analysis of the wide field data raw images collected were binned at 4x and
then smoothed with a moving window of 4x4. Further analysis was done on the smoothed
images. In order to find the significant pixels, a one-sided t-test across all the iterations was
performed between the 5 frames immediately preceding the stimulus onset and a moving
window of 5 frames after the stimulus onset in all 130 μm by 130 μm square areas. Only
those pixels which gave a p-value less than 0.05 on at least one window after the stimulus
onset were considered for further analysis. The mean change in df/f with time traces was
constructed for pixels showing significant positive responses. These traces were further
smoothed with a moving average window of 5 frames and a one-sided t-test was performed
again to find the first significant post-stimulus frame and the frame with the most significant
response. Some of these traces which showed the maximum response are shown in the
main text (Fig.3A) and supplementary figure (Fig. S11C).

1099

1100

1101

1102

1103

1104

1105

1106

1107 Table S3

Sl.no.	Age	Tonal stimuli played (~90db SPL, in kHz)	Significant responsive frequencies and respective % responding areas	Fraction of frequencies played showed responses (%)
1.	P-7	8.5,12,17,24,48	8.5kHz: 2.86%	20.00%
2.	P-8	6,8.5,12,17,24,34, 48	6kHz: 0.56%; 8.5kHz: 3.67%	28.57%
3.	P-9	6,8.5,12,17,24,34, 48	6kHz: 26.9%; 17kHz: 3.44%; 24kHz: 0.56%; 48kHz: 1.31%	57.14%
4.	P-10	6,8.5,12,17,24,34, 48	8.5kHz: 37.12%; 12kHz: 4.38%; 17kHz: 0.5%; 48kHz: 19.93%	57.14%
5.	P-13	6,8.5,12,17,24,34, 48	6kHz: 43.8%; 8.5kHz: 79.8%; 12kHz: 13.54%; 17kHz: 70.8% 24kHz: 22.9%	71.42%

1108

2-photon imaging analysis was performed using custom codes written in MATLAB
(Mathworks). Imaging sequences were aligned by performing X-Y drift correction. Cells
were selected manually by selecting the centre point of the cell on the motion corrected
mean image. ROI (5 μm radius) were drawn based on the cell centres. Raw Fluorescence
signal over time (F) of the selected ROIs across all frames were extracted. For each trial,
relative fluorescence was computed by using $\Delta F/F_0 = (F - F_0)/ F_0$), where F_0 corresponds to
baseline fluorescence. Baseline fluorescence amplitude was estimated by computing mean
of fluorescence values over 5 frames before the stimulus onset. Neurons on stimulation to
tones and noise were considered to be significantly responsive as follows. One-sided paired
ttest's were performed between mean df/f over 3 of the successive frames before the
stimulus and each one of 4 windows, of 3 successive frames after stimulus onset (up to the
6^{th} frame, ~1.2-1.5s). If any of the 4 moving windows showed a significant increase in mean
df/f (40 repetitions) over baseline, the neuron was considered responsive.

Table S4

Sl.no.	Age	Fraction of cases (all neurons in response to different frequency tone) that showed significant responses.
1.	P-7	5.78%
2.	P-8	4.13%
3.	P-9	5.91%
4.	P-10	3.08%
5.	P-11	2.31%
6.	P12-P13	8.23%

1123

1124   Table S5

Sl.no.	Age	Fraction of cases (all neurons in response to broadband noise) that showed significant responses.
1.	P-7	5.35%
2.	P-8	10.25%
3.	P-9	8.11%

1125

For significant responses in single trials to compute reliability, a trial was deemed responsive
if the mean df/f of 4 frames (~1s) after the stimulus after stimulus onset was greater than
1*STD (STD obtained from 40 repetitions) of the mean of the 4 baseline frames preceding
the stimulus. Given the unreliability of the responses before ear canal opening, for the
population traces (Fig. 3B), only those neurons were considered in the pool which evoked
significant responses to >25% of the trials (>25% reliability). Reliability was computed as the
proportion of trials that showed significant responses (above).

In case of the SD paradigm, each SD set of a pair (SD and swap) was collected with 30
repetitions. Cells in the pair of datasets (SD and swap) were matched for further analysis, as
the imaging for each SD dataset lasted long (> 5 mins), there could be drifts between two
datasets. Thus the matching was based on correlations (MATLAB function corr2) between
different relative x-y translations (+/- 5.8 μm or 5 pixels) between the two x-y drift corrected
(see above) mean images acquired in each SD set of a pair. The relative x-y translation that
gave the maximum correlation was used to then consider same cells in both datasets if the

cell centers were within 5.8 μm. Only cells that matched in the 2 datasets were considered
further. Many neurons did not show significant responses (d-prime<1, between pre stimulus
frames df/f values and post stimulus, S onset or post D) in the average df/f of all 30
repetitions. However, as responses had low reliability, significant responses (d-prime>1)
were observed in stretches of 10 repetitions of the stimuli (moving average was performed
of 10 repetitions to check for significant responses). Response window of 6 frames after
standard and deviant token onset and the baseline window of 3 frames before stimulus
onset were considered for drawing significance. Pairs of neurons which intermittently
responded to standards or deviants in at least 3 moving repetition windows (of 10
repetitions) were considered for population analysis. Mean df/f traces could become
negative and hence the mean df/f of each trial was shifted by adding the minimum df/f over
all frames before calculating CSI from 2-photon Ca^{2+} imaging data. CSI was calculated from
mean translated (as above) df/f of 6 frames following S onset and eight frames after D for
each set in a pair. CSI is calculated as done with mean rates, replacing rate by df/f. Since
other than the onset token and deviant token there were no responses in single units at
early ages these mean values likely do not include effects of other S's.

Details of Binary Network Model of L4 and SP with TC Inputs
1157

1158 We built a binary neural network consisting to two separate thalamic input units for
1159 standard and deviant. The thalamic units project to a sub-plate neuron and further to a
1160 cortical neuron. The sub-plate neuron also projects to the cortical neuron. A similar model
1161 was used to demonstrate ocular dominance in the visual cortex[11,9]. The spike rate of the two
1162 thalamic inputs are given by,
1163

1164
$$f_{std}(r) = \begin{cases} 8 & when\ a\ standard\ token\ comes \\ 0.15 & when\ a\ deviant\ token\ comes \\ 0.1 & spontaneous\ rate \end{cases}$$
$$f_{dev}(r) = \begin{cases} 0.15 & when\ a\ standard\ token\ comes \\ 8 & when\ a\ deviant\ token\ comes \\ 0.1 & spontaneous\ rate \end{cases}$$

1165

1166 The synaptic weight of the thalamo-subplate projections were kept constant at a value of
1167 0.2. The projections received by the cortical neuron was kept plastic and learning was
1168 implemented following the below plasticity rule.
1169

1170
$$\frac{\Delta w}{w} = \begin{cases} Amp_{strength} * e^{-\frac{\Delta t}{\tau_{strength}}} & potentiation \\ -Amp_{weak} * e^{-\frac{\Delta t}{\tau_{weak}}} & depression \end{cases}$$

1171
1172 Where $\Delta t = |t_{post\ synaptic} - t_{pre\ synptic}|$
1173 $Amp_{strength} = 0.015$ $\tau_{strength} = 13\ ms$
1174 $Amp_{weak} = 0.021$ $\tau_{weak} = 20\ ms$
1175

We performed simulations for two regimes, one for the early development phase and one
for the later. For the early phase, we kept the synaptic weight of the sub-plate projections at
a value of 0.22 and that of the thalamo-cortical projections at a smaller value of 0.02. This
was done to mimic the early stages of cortical development where the thalamic projections
are unable to drive the cortical activity alone and are aided by the sub-plate for driving the
cortical activity[1,2,20].

In the later phase, both these weights were kept at an intermediate value of 0.04 and 0.12
respectively. This reflects the intermediate stage of development where the thalamo-cotical
projections are strengthening and the sub-plate-cortical connections are weakening[2,20].

All the synapses were kept depressing throughout the simulation with the following
equations governing their dynamics[41],

1187

1188
$$\frac{dx_r}{dt} = -M.\frac{x_r}{\tau_{re}} + \frac{x_i}{\tau_{ir}} \qquad \frac{dx_e}{dt} = +M.\frac{x_r}{\tau_{re}} - \frac{x_e}{\tau_{ei}} \qquad \frac{dx_e}{dt} = \frac{x_e}{\tau_{ei}} - \frac{x_i}{\tau_{ir}}$$

1189

1190   Where M=1 whenever a spike occurs.

1191

1192   For thalamus, $\tau_{re} = 0.9\ ms$ $\tau_{ei} = 10\ ms$ $\tau_{ir} = 4200\ ms\ (early)\ \&\ 100ms\ (later)$

1193

1194   For sub-plate, $\tau_{re} = 0.9\ ms$ $\tau_{ei} = 27\ ms$ $\tau_{ir} = 4200\ ms\ (early)\ \&\ 100ms\ (later)$

1195

1196   The value of τ_{ir} was chosen in accordance with our electrophysiology data which shows a
1197   decrease in adaptation in later ages.

1198

1199   The sub-plate and the cortical neuron were modelled on the principles of integrate and fire
1200   neuron[1]. The membrane potential can be written as follows,

1201

$$V_{sp}(t) = g_{std}(t-1) * w_{s1}(t-1) * x_{r_{std}}(t-1) + g_{dev}(t-1) * w_{s2}(t-1) * x_{r_{dev}}(t-1)$$
$$V_{l4}(t) = g_{std}(t-1) * w_{l1}(t-1) * x_{r_{std}}(t-1) + g_{dev}(t-1) * w_{l2}(t-1) * x_{r_{dev}}(t-1)$$
$$+ g_{sp}(t-1) * w_{sp}(t-1) * x_{r_{sp}}(t-1)$$

1202

1203   Where g(t) is the conductance and can be written as,

1204

1205
$$g(t) = spk\_trn * k(t)$$
$k(t)$ is the convolution kernel
$$k(t) = e^{-(t-t_{spk})/\tau_{syn}}$$
$$\tau_{syn} = 10\ ms$$

1206

1207   The threshold of Layer IV neuron was kept constant at 0.1 for both the regimes, while that
1208   of the subplate neuron was kept at 0.05 for the early regime and 0.15 for the late regime.

1209

1210   The refractory period dynamics after each spike was modelled with a separate exponential
1211   given by[1],

1212

$$V_\Delta(t) = \beta * e^{-(t-t_{spk})/\tau_{refr}}$$
$$\tau_{refr} = 2\ ms$$
$$\beta = 5$$

1213

1214   The voltage drops by the above value after every spike. The refractory period lasts for 20ms
1215   after the spike. A leak of 10% was also incorporated[1].
1216

1217   The model gets a drive from the thalamic activity, modeled by uncorrelated Poisson spikes
1218   in the two thalamic units. The stimulus train consists of standard and deviant tokens
1219   occurring with a probability of 0.9 and 0.1 respectively in the standard-deviant protocol and
1220   consisted of only standard tokens in case of only standard protocol. Each token is 50ms long
1221   separated by a 250ms inter-token-interval with a sampling rate of 1000/sec.
1222

1223

1224   **Theoretical Analysis for Optimization based on Mutual Information Maximization and**
1225   **Sparseness**
1226

1227   We consider the same stimulus train that we used in the exposure experiments.
1228   We assume the probability of deviant to be x and that of standard to be 1-x
1229

1230

$$p(t_{std}) = 1 - x \qquad p(t_{dev}) = x$$

1231

1232   The responses for each of the two tokens are normally distributed
1233

1234 $p(r|t_{std}) = \frac{1}{\sigma\sqrt{2\pi}}\exp\left(-\left(\frac{(r-r_o)^2}{2\sigma^2}\right)\right)$ or g_s $p(r|t_{dev}) = \frac{1}{\sigma\sqrt{2\pi}}\exp\left(-\left(\frac{(r-(r_o+a_k))^2}{2\sigma^2}\right)\right)$ or g_d

1235

1236   We consider the variance σ^2 and mean response for the standard token r_o to be constant.
1237   The mean response for the deviant token varies with each successive token and its
1238   difference from the mean standard response takes values a_k for every kth token. We
1239   assume all the tokes to be independent and without any history dependence (no
1240   adaptation). Thus,
1241

$$p(r) = xg_d + (1 - x)g_s$$

1242

1243   We wish to know the trend of $f(k)$ with successive tokens that will enhance the
1244   discriminability between the standard and the deviant token. To do that, we use mutual
1245   information as a measure of discriminability paired with a sparse coding constraint and try
1246   to maximize it.
1247   Our constrained information for the k[th] token is,
1248

$$CI_k = I(R_k, T) - \rho(r_o + xa_k)$$

1249

Where $I(R_k, T)$ denotes the mutual information between the response at the kth token and
the stimulus. ρ is the langrage multiplier which is constraining the mean response at the kth
token. The langragian has been constructed to minimize the mean response at the kth token
(sparse representation).
We need to find the value of ρ which will maximize the mutual information between the
entire stimulus train and the total response along with maintaining a sparse representation

throughout the stimulus period. We can write the total constrained information as the sum
of the constrained information of independent individual tokens,
1258

$$CI_{total} = \sum_{k=1}^{N} CI_k = \sum_{k=1}^{N} I(R_k, T) - \rho \left(N r_o + x \sum_{k=1}^{N} a_k \right)$$

1259

1260   Where N is the total number of tokens.
1261   For any token, $I(R_k, T)$ can be written as,
1262

$$I(R,T) = \int_0^{r_{max}} \left[\sum_T \left\{ p(r,t) \ln \left(\frac{p(r,t)}{p(r)p(t)} \right) \right\} \right] dr$$

$$= \int_0^{r_{max}} \left[\sum_T \left\{ p(r|t) \, p(t) \ln \left(\frac{p(r|t)p(t)}{p(r)p(t)} \right) \right\} \right] dr$$

$$= \int_0^{r_{max}} \left[\sum_T \left\{ p(r|t) \, p(t) \ln \left(\frac{p(r|t)}{p(r)} \right) \right\} \right] dr$$

$$= \int_0^{r_{max}} \left[p(r|t_{dev}) \, p(t_{dev}) \ln \left(\frac{p(r|t_{dev})}{p(r)} \right) + p(r|t_{std}) \, p(t_{std}) \ln \left(\frac{p(r|t_{std})}{p(r)} \right) \right] dr$$

$$= \int_0^{r_{max}} \left[x g_d \ln \left(\frac{g_d}{x g_d + (1-x)g_s} \right) + (1-x)g_s \ln \left(\frac{g_s}{x g_d + (1-x)g_s} \right) \right] dr$$

$$= \int_0^{r_{max}} \left[x g_d \ln g_d + (1-x)g_s \ln g_s - (x g_d + (1-x)g_s) \ln(x g_d + (1-x)g_s) \right] dr$$

1263 $$= -\frac{1}{2} \ln(2\pi e \sigma^2) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x g_d + (1-x)g_s) \right] dr$$

1264 $$= -\frac{1}{2} \ln(2\pi e \sigma^2) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\{ g_s(x A(r) + (1-x)) \} \right] dr \qquad \text{where } A(r) = \frac{g_d}{g_s}$$

$$= -\frac{1}{2} \ln(2\pi e \sigma^2) - x \int_0^{r_{max}} [g_d \ln g_s] \, dr - (1-x) \int_0^{r_{max}} [g_s \ln g_s] \, dr - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x A(r) + (1-x)) \right] dr$$

$$= -\frac{1}{2} \ln(2\pi e \sigma^2) + \frac{x}{2} \ln(2\pi e \sigma^2) + \frac{x a^2}{2\sigma^2} + \frac{1-x}{2} \ln(2\pi e \sigma^2) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x A(r) + (1-x)) \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln(x A(r) + (1-x)) \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\left\{ (1-x) \left(\frac{x A(r)}{1-x} + 1 \right) \right\} \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \ln(1-x) \int_0^{r_{max}} [x g_d + (1-x)g_s] \, dr - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\left(\frac{x A(r)}{1-x} + 1 \right) \right] dr$$

$$= \frac{x a^2}{2\sigma^2} - \ln(1-x) - \int_0^{r_{max}} \left[(x g_d + (1-x)g_s) \ln\left(\frac{x A(r)}{1-x} + 1 \right) \right] dr$$

1265   We wish to use the Taylor expansion of ln in the above expression. To do that we need,
1266

$$\frac{x A(r)}{1-x} < 1$$

$$x < \frac{1}{1 + A(r)}$$

1267

1268   We see that $A(r)$ is the ratio of two normal distributions and it can be reduced to,

$$A(r) = exp \left(\frac{a(2r - 2r_o - a)}{2\sigma^2} \right)$$

1269

1270   Since x is a constant and is only dependent on the stimulus, we set a limit for x that will
1271   make this expression valid for all the possible rates,
1272

$$x < \frac{1}{1 + exp \left(\frac{a(2r - 2r_o - a)}{2\sigma^2} \right)}$$

1273   If we assume that on the 1st token (when the discrimination has not yet started), the mean
1274   rate for standard and deviant is the same (a=0), we see that,

1275

$$x < \frac{1}{2}$$

1276

1277 Thus we proceed further with an assumption that the probability of the arrival of the
1278 deviant is less than 50% and the results that we discuss later will hold true only on this
1279 assmption.
1280 We now perform the Taylor series expansion,
1281

$$\frac{xa^2}{2\sigma^2} - \ln(1-x) - \int_0^{r_{max}} \left[(xg_d + (1-x)g_s) \sum_{i=1}^{\infty} \left\{ \frac{-1^{i+1}}{n} \left(\frac{xA(r)}{1-x} \right)^i \right\} \right] dr$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \int_0^{r_{max}} \{g_d\{A(r)\}^i\} \, dr \right] - (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \int_0^{r_{max}} \{g_s\{A(r)\}^i\} \, dr \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-(r_o+a))^2 + i(a^2 - 2a(r-r_o))}{2\sigma^2} \right) \right\} dr \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-r_o)^2 + i(a^2 - 2a(r-r_o))}{2\sigma^2} \right) \right\} dr \right]$$

1287 =

1288 $$\frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} \int_0^{r_{max}} \left\{ exp\left(-\frac{((r-r_o)^2 + a^2 - 2(r-r_o)a + ia^2 - 2ia(r-r_o))}{2\sigma^2} \right) \right\} dr \right] -$$

1289 $$(1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-r_o)^2 + ia^2 - 2ia(r-r_o)}{2\sigma^2} \right) \right\} dr \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{(i+1)a^2}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{((r-r_o)^2 - 2(r-r_o)a - 2ia(r-r_o))}{2\sigma^2} \right) \right\} dr \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{ia^2}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-r_o)^2 - 2ia(r-r_o)}{2\sigma^2} \right) \right\} dr \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{(i+1)a^2}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r^2 + r_o^2 - 2rr_o - 2ra + 2r_oa - 2iar + 2iar_o)}{2\sigma^2} \right) \right\} dr \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{ia^2}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{r^2 + r_o^2 - 2rr_o - 2iar + 2iar_o}{2\sigma^2} \right) \right\} dr \right]$$

1290

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{(i+1)a^2 + r_o^2 + 2r_oa + 2iar_o}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r^2 - 2(r_o + a + ia)r)}{2\sigma^2} \right) \right\} dr \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{ia^2 + r_o^2 + 2iar_o}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{r^2 - 2(r_o + ia)r}{2\sigma^2} \right) \right\} dr \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x)$$

$$- x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{(i+1)a^2 + r_o^2 + 2r_oa + 2iar_o - (r_o + a + ia)^2}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-(r_o + a + ia))^2}{2\sigma^2} \right) \right\} dr \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} \frac{1}{\sigma\sqrt{2\pi}} exp\left(-\frac{ia^2 + r_o^2 + 2iar_o - (r_o + ia)^2}{2\sigma^2} \right) \int_0^{r_{max}} \left\{ exp\left(-\frac{(r-(r_o + ia))^2}{2\sigma^2} \right) \right\} dr \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(-\frac{(i+1)a^2 + r_o^2 + 2r_oa + 2iar_o - (r_o + a + ia)^2}{2\sigma^2} \right) \right]$$

$$- (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(-\frac{ia^2 + r_o^2 + 2iar_o - (r_o + ia)^2}{2\sigma^2} \right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(\frac{ia^2(i+1)}{2\sigma^2} \right) \right] - (1-x) \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^i}{i(1-x)^i} exp\left(\frac{ia^2(i-1)}{2\sigma^2} \right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - \sum_{i=1}^{\infty} \left[\frac{-1^{i+1}x^{i+1}}{i(1-x)^i} exp\left(\frac{ia^2(i+1)}{2\sigma^2} \right) + \frac{-1^{i+1}x^i}{i(1-x)^{i-1}} exp\left(\frac{ia^2(i-1)}{2\sigma^2} \right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x + x \sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} exp\left(\frac{i(i+1)a^2}{2\sigma^2} \right) \right]$$

$$= \frac{xa^2}{2\sigma^2} - \ln(1-x) - x + x\Lambda(a)$$

1291 Where $\Lambda(a) = \sum_{i=1}^{\infty} \left[\frac{-1^i}{i(i+1)} \frac{x^i}{(1-x)^i} exp\left(\frac{i(i+1)a^2}{2\sigma^2} \right) \right]$

1292

1293 On a careful examination, we see that the above expression will always be positive.

1294 We find the change in constrained information between two tokens m&n (m<n)

1295

$$\Delta CI = CI_n - CI_m = \{MI_n - MI_m\} - \rho x\{a_n - a_m\}$$

$$= x\left[\frac{a_n^2 - a_m^2}{2\sigma^2} + \Lambda(a_n) - \Lambda(a_m) - \rho\{a_n - a_m\}\right]$$

$$= x(a_n - a_m)\left[\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} - \rho\right]$$

1296 We want the above expression to be greater than or equal to zero. Thus, we get two cases,

1297

1298 Case 1: $a_n > a_m$

1299

1300 We need,

1301

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} > \rho$$

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\sum_{i=1}^{\infty}\left[\frac{-1^i}{i(i+1)}\frac{x^i}{(1-x)^i}\left\{exp\left(\frac{i(i+1)a_n^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_m^2}{2\sigma^2}\right)\right\}\right]}{a_n - a_m} \geq \rho$$

1302

1303 We see that as a increases, the 2nd term in the above expression increases with the
1304 exponential of the square of the increase in a. The 2nd term over all will be much larger than
1305 1. If we assume a monotonic behavior in a, the least value of $\Lambda(a_n) - \Lambda(a_m)$ will be
1306 between the first two tokens. Thus we can say that,

1307

$$\rho \leq \frac{a_2 + a_1}{2\sigma^2} + \frac{\sum_{i=1}^{\infty}\left[\frac{-1^i}{i(i+1)}\frac{x^i}{(1-x)^i}\left\{exp\left(\frac{i(i+1)a_2^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_1^2}{2\sigma^2}\right)\right\}\right]}{a_2 - a_1}$$

1308

1309 The above expression sets a limit on the value of ρ or in other words the weightage of
1310 sparseness which will keep the mutual information increasing.
1311 This case corresponds to the early age results, where the responses for deviant was greater
1312 than that of standard (a>0).

1313

1314 Case 2: $a_n < a_m$

1315 We need,

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\Lambda(a_n) - \Lambda(a_m)}{a_n - a_m} \leq \rho$$

$$\frac{a_n + a_m}{2\sigma^2} + \frac{\sum_{i=1}^{\infty}\left[\frac{-1^i}{i(i+1)}\frac{x^i}{(1-x)^i}\left\{exp\left(\frac{i(i+1)a_n^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_m^2}{2\sigma^2}\right)\right\}\right]}{a_n - a_m} \leq \rho$$

1316

1317 We see that as a decreases, the 2nd term in the above expression decreases with the
1318 exponential of the square of the increase in a. The 2nd term over all will be much less than -
1319 1. If we assume a monotonic behavior in a, the largest value of $\Lambda(a_n) - \Lambda(a_m)$ will be
1320 between the first two tokens. Thus we can say that,

1321

$$\rho \geq \frac{a_2 + a_1}{2\sigma^2} + \frac{\sum_{i=1}^{\infty}\left[\frac{-1^i}{i(i+1)}\frac{x^i}{(1-x)^i}\left\{exp\left(\frac{i(i+1)a_2{}^2}{2\sigma^2}\right) - exp\left(\frac{i(i+1)a_1{}^2}{2\sigma^2}\right)\right\}\right]}{a_2 - a_1}$$

The above expression sets a limit on the value of ρ or in other words the weightage of sparseness which will keep the mutual information increasing.

This case corresponds to the later age results, where the responses for standard was less than that of deviant (a<0).

Supplementary Figure Legends

Figure S1. A) Schematic of types of 4x4 electrode arrays used – either all electrodes were of same lengths (i) or they had alternate short and long electrodes (ii) either 600 μm apart or 400 μm apart. B) Sample locations of recordings of single units mapped onto the cortical surface showing broad clearer tonotopy in P29 than in P15. C) Representative confirmation of recording in A1 in lower age groups P12, P13, P18 with DiI injections in the recording site with labelling in MGBv (inset). D) Thickness of SP as a percentage of cortical thickness and depth of L4 and SP over age determined from cell morphology with Nissl stains.

Figure S2. SP depth and thickness determined with immunohistochemistry (IHC) by labelling SPNs with antibodies of SPN markers Complexin-3 (A) and CTGF (B).

Figure S3. Slices were cut from brains (fixed) in which single unit recordings were performed in the ACX (representative examples at 3 ages, in A) and IHC was performed. Electrode tracks show recordings performed in SPN and in case of dual recordings (second electrode type, Fig. S1A) in SCNN1-td-tomato mouse show electrodes in L4 and SPN simultaneously (B).

Figure S4. Top and bottom row show bar plots of mean CSI(F) and mean CSI for neurons in L4 at different f_1 and f_2 separations.

Figure S5. A) Schematic of SD tokens and its swap, which is the stimulus presentation protocol. B-G) Data from six representative L4 units at different ages (P12-P38, same as in Fig. 2), given in titles with respective CSI(F), are shown as dot rasters and PSTHs in response to the 4 stimulus sets (i-iv) – noise at multiple sound levels, pure tones at different frequencies, f-N SD stimulus and its swap. The chosen frequency and sound level of noise used in the SD stimuli are marked on the respective dot rasters in i and ii with gray light and dark rectangles, along with PSTHs of those responses in the bottom plot of i and ii. PSTHS in iii and iv also shown in Fig. 2A. CSI(F) values are given in parentheses for the 6 neurons in B-G.

Figure S6. Same set of plots as in Fig. S5B (for L4 neurons of Fig. 2A), for the 3 representative SPN responses shown in Fig. 2B. CSI(F) values are given in parentheses for the 3 neurons in A-C.

Figure S7. A) Histograms of deviant selectivity (*N-f* case) measured with CSI values - CSI(F) in i, CSI in ii and CSI(P) in iii, for the entire population of neurons of L4 (top row in each plot i-

iii) and SP (bottom row) as a function of age (5 age groups considered). B) Same plot as in Fig. 1H-I, for CSI(P) for f_1-f_2 SD stimulus.

Figure S8. Ai) Shows the population CDFs of adaptation time constants (as -1/time constant) for L4 neurons (dark gray) and SP neurons (light gray) over different age groups (SP only for P12-15). Adaptation time constants, determined from exponential fits to mean responses to each successive token of Standards, increase with age. Aii) Shows population CDFs (for the same set of neurons and age groups as in the top row) of the number tokens in which there were significant responses in the first set of 7 (or 6) tokens before the deviant for the same population of neurons. B) Mean -1/Adaptation time constants for the data in Ai. C) Boxplots showing range with median, IQR and outliers for data in Aii.

Figure S9. A) Shows a sample image of the relevant part of a coronal section with MGB, in which single unit recordings from MGBv was performed (P14). B) Histograms of CSI(F) in the top row and CSI in the bottom row are shown for all single units recorded in MGBv from two age groups (left P14-15, right P24-26). C) Mean and standard errors of CSI(F) and CSI for the data in B, by age groups, are shown.

Figure S10. A) i) LFP in a single channel in SP, in response to noise obtained from 5 repetitions at P08. ii) Population average with SEM of the same in above. iii) Population average and SEM of rms values of LFP. B) i) Population average of LFP in SP and L4 to tones from 5 repetitions ii) Population average and SEM of rms values of LFP in SP and L4 to the same as above. iii) Representative, sample traces of LFP (average of 5 repetitions) obtained in P9 and P10 from SP and L4 (1 case each) showing responses to tones (each row, 6-48 kHz, ½ octave apart ~90 dB SPL, 50 ms tones). C) Dot rasters and PSTHs of spiking activity of 4 neurons (2 in L4 and 2 in SP, identified in each column) before ECO with pure tone presentations show unreliable spiking with 5 repetitions (thick black line – stimulus onset).

Figure S11 A) Left: Image of craniotomy in which wide field imaging was performed in a P7 mouse. Blue shaded regions showed positive maximum df/f post stimulus, with widefield imaging. Right: A sample trace shows the smoothed mean df/f across the frames of the region marked by the blue square on the craniotomy. Black horizontal bar represents the stimulus ('*' corresponds to minimum p-value among all frames, '.' represents the first significant window). B-C) Example traces of df/f of single neurons (arrows in 2-photon mean images of frame with cells expressing GCamp6f, bottom) in response to broadband noise (C) and tones (B) respectively at three different ages. The corresponding gray and black vertical lines represent stimulus onset. The error bars show the standards error of 40 repetitions. The red and black bars represent the earliest window over which the mentioned p-values were achieved.

Figure S12. Mean df/f traces of all the cells having more than 25% reliability in responses (significant response in >10 repetitions out of 40 repetitions) to tonal stimuli (frequencies in Hz mentioned) at the different ages. Respective p-values and degree of freedom for one-sided paired ttest is also shown. Imaging was performed with 40 repetitions of each frequency in P07-P11 (before ECO) and 5 repetitions in P12-13 after ECO. All imaged cells are included in traces for P12-13 irrespective of reliability.

1412

1413   Figure S13. A) i-ii) Similar plot as in Fig. 3B i-iii, for in-vivo 2-photon Ca^{2+} imaging based

1414   responses to broadband noise. . Bi-ii) Same as in Fig.3Ei-ii, for responses to broadband noise

1415   before ECO.

1416

Figure S14. A) i-ii) Representative df/f traces of responding single neurons in L2/3 to the SD

stimulus, obtained with 2-photon Ca^{2+} imaging (i: S=tone, D=noise, TN and ii: S=noise,

D=tone, NT) at three different ages before ECO. The grey/black and black/grey vertical lines

represent S onset and D respectively. Black ($p<0.2$) and red ($p<0.05$) bars (above traces)

mark timing of significant responses (mean of 3 frames) to the standard and the deviant. iii-

iv) Population mean df/f (311 neurons, n=3 (P7-P9)) in response to the two stimuli TN (iii)

and NT (iv). v) Distribution of CSI values for the population of L2/3 single neurons.

1424

Figure S15. A) i) Figure S15. A) i) the z-scores of responses in SD stimulus (20 reps, 4Hz) (left

column) and Swap (right column) for 3 single units in L4 (top 3 rows) and 3 single units in SP

(bottom 3 rows) are shown (same units as in Fig. 4B, first 3 columns). A) ii) Mean z-scores for

the population of single units in L4 (top row) and that in SP (bottom row) are shown for the

SD stimulus (left) and the corresponding Swap (right). B) i-ii) Same as Ai-ii, for responses to

SD stimulus with 30 reps and at rate of 3.33 Hz.

1431

1432

Figure S16. Top and middle row show the mean CSI values of the population of L4 and SP

neurons respectively for different choice of response latencies and response window sizes.

Asterisks marks the response latency and response window size combination used in Fig. 4

CSI calculations. The three columns are for the 3 different CSIs used CSI, CSI(F) and CSI(P).

Bottom row shows which combinations of response attribute have significantly higher CSI in

SP than in L4 (brown, blue indicates NS).

1439

Figure S17. A) 2 examples of cross-correlograms each for before ECO (top row) and after

ECO (bottom wor) showing connections detected from SP to L4 (left) and L4 to SP (right).

Blue and red colored lines indicate mean of cross-correlograms with errorbars and gray lines

with errorbars show the same for shuffled data (see detailed methods). B) Bar plots

summarize the data in Table S2 showing probability of connections from SP to L4 (left) and

L4 to SP (right) at within 125 microns or greater distances at the 2 age groups before ECO

and P12-13, after ECO).

1447

Figure S18. A) Four example images of location (along cortical surface) of BFs on the cortical

surface in the craniotomy of single units recorded in L4 at ages P28 and above (Fig. 5A single

unit recording) identified in color (colorbar, bottom right) for four exposure groups EX2 and

EX1 in the top row (SD exposure) and EX5 and EX6 in the bottom row (SS exposure). Broad

tonotopy showing A1 and AAF are clear in the images. B) Bar plot shows % change in area of

each of the preferred frequencies in A1 and AAF between control (EX0 group of mice) and

exposed animals (EX5 group of mice).

1455

Figure S19. Similar plots as in Fig. 5E, for the profile of population mean normalized rates to the relevant RF (color legend) for the 4 exposure groups (EX5 and EX6 left, EX2 and EX1 right) as a function of BF (re RF). Normalization was done by considering the maximum rate in each tuning curve (rate at BF) to be 100%. Gray profile shows the same for the maximum of 13 similar plots derived from the control data (C) considering each of the 13 frequencies used as the RF.

Figure S20. Bar plot summarizes population mean response to RF for neurons with BF at RF of all 6 EX cases (2 bars each, red and green for SD exposure cases and 1 blue for SS). Mean of the 13 control cases shown in gray bar in the centre.

Supplementary Figures

 Figure S1. A) Schematic of types of 4x4 electrode arrays used – either all electrodes were of
 same lengths (i) or they had alternate short and long electrodes (ii) either 600 μm apart or
 400 μm apart. B) Sample locations of recordings of single units mapped onto the cortical
 surface showing broad clearer tonotopy in P29 than in P15. C) Representative confirmation
 of recording in A1 in lower age groups P12, P13, P18 with DiI injections in the recording site
 with labelling in MGBv (inset). D) Thickness of SP as a percentage of cortical thickness and
 depth of L4 and SP over age determined from cell morphology with Nissl stains.

Figure S2. SP depth and thickness determined with immunohistochemistry (IHC) by labelling
SPNs with antibodies of SPN markers Complexin-3 (A) and CTGF (B).

1553

1554
1555
1556

Figure S3. Slices were cut from brains (fixed) in which single unit recordings were performed in the ACX (representative examples at 3 ages, in A) and IHC was performed. Electrode tracks show recordings performed in SPN and in case of dual recordings (second electrode type, Fig. S1A) in SCNN1-td-tomato mouse show electrodes in L4 and SPN simultaneously (B).

1597 Figure S4. Top and bottom row show bar plots of mean CSI(F) and mean CSI for neurons in
1598 L4 at different f_1 and f_2 separations.

Figure S5. A) Schematic of SD tokens and its swap, which is the stimulus presentation protocol. B-G) Data from six representative L4 units at different ages (P12-P38, same as in Fig. 2), given in titles with respective CSI(F), are shown as dot rasters and PSTHs in response to the 4 stimulus sets (i-iv) – noise at multiple sound levels, pure tones at different frequencies, f-N SD stimulus and its swap. The chosen frequency and sound level of noise used in the SD stimuli are marked on the respective dot rasters in i and ii with gray light and dark rectangles, along with PSTHs of those responses in the bottom plot of i and ii. PSTHS in iii and iv also shown in Fig. 2A. CSI(F) values are given in parentheses for the 6 neurons in B-G.

1640 Figure S6. Same set of plots as in Fig. S5B (for L4 neurons of Fig. 2A), for the 3 representative
1641 SPN responses shown in Fig. 2B. CSI(F) values are given in parentheses for the 3 neurons in
1642 A-C.

1644
1645

Figure S7. A) Histograms of deviant selectivity (*N-f* case) measured with CSI values - CSI(F) in i, CSI in ii and CSI(P) in iii, for the entire population of neurons of L4 (top row in each plot i-iii) and SP (bottom row) as a function of age (5 age groups considered). B) Same plot as in Fig. 1H-I, for CSI(P) for f_1-f_2 SD stimulus.

Figure S8. Ai) Shows the population CDFs of adaptation time constants (as -1/time constant) for L4 neurons (dark gray) and SP neurons (light gray) over different age groups (SP only for P12-15). Adaptation time constants, determined from exponential fits to mean responses to each successive token of Standards, increase with age. Aii) Shows population CDFs (for the same set of neurons and age groups as in the top row) of the number tokens in which there were significant responses in the first set of 7 (or 6) tokens before the deviant for the same population of neurons. B) Mean -1/Adaptation time constants for the data in Ai. C) Boxplots showing range with median, IQR and outliers for data in Aii.

1673

1674
1675
1676
1677 Figure S9. A) Shows a sample image of the relevant part of a coronal section with MGB, in
1678 which single unit recordings from MGBv was performed (P14). B) Histograms of CSI(F) in the
1679 top row and CSI in the bottom row are shown for all single units recorded in MGBv from two
1680 age groups (left P14-15, right P24-26). C) Mean and standard errors of CSI(F) and CSI for the
1681 data in B, by age groups, are shown.

Figure S10. A) i) LFP in a single channel in SP, in response to noise obtained from 5
repetitions at P08. ii) Population average with SEM of the same in above. iii) Population
average and SEM of rms values of LFP. B) i) Population average of LFP in SP and L4 to tones
from 5 repetitions ii) Population average and SEM of rms values of LFP in SP and L4 to the
same as above. iii) Representative, sample traces of LFP (average of 5 repetitions) obtained
in P9 and P10 from SP and L4 (1 case each) showing responses to tones (each row, 6-48 kHz,
½ octave apart ~90 dB SPL, 50 ms tones). C) Dot rasters and PSTHs of spiking activity of 4
neurons (2 in L4 and 2 in SP, identified in each column) before ECO with pure tone
presentations show unreliable spiking with 5 repetitions (thick black line – stimulus onset).

Figure S11 A) Left: Image of craniotomy in which wide field imaging was performed in a P7 mouse. Blue shaded regions showed positive maximum df/f post stimulus, with widefield imaging. Right: A sample trace shows the smoothed mean df/f across the frames of the region marked by the blue square on the craniotomy. Black horizontal bar represents the stimulus ('*' corresponds to minimum p-value among all frames, '.' represents the first significant window). B-C) Example traces of df/f of single neurons (arrows in 2-photon mean images of frame with cells expressing GCamp6f, bottom) in response to broadband noise (C) and tones (B) respectively at three different ages. The corresponding gray and black vertical lines represent stimulus onset. The error bars show the standards error of 40 repetitions. The red and black bars represent the earliest window over which the mentioned p-values were achieved.

1729
Figure S12. Mean df/f traces of all the cells having more than 25% reliability in responses (significant response in >10 repetitions out of 40 repetitions) to tonal stimuli (frequencies in Hz mentioned) at the different ages. Respective p-values and degree of freedom for one-sided paired ttest is also shown. Imaging was performed with 40 repetitions of each frequency in P07-P11 (before ECO) and 5 repetitions in P12-13 after ECO. All imaged cells are included in traces for P12-13 irrespective of reliability.

1736
1737
1738

1739
1740
1741
1742

Figure S13. A) i-ii) Similar plot as in Fig. 3B i-iii, for in-vivo 2-photon Ca^{2+} imaging based
responses to broadband noise. . Bi-ii) Same as in Fig.3Ei-ii, for responses to broadband noise
before ECO.

Figure S14. A) i-ii) Representative df/f traces of responding single neurons in L2/3 to the SD stimulus, obtained with 2-photon Ca^{2+} imaging (i: S=tone, D=noise, TN and ii: S=noise, D=tone, NT) at three different ages before ECO. The grey/black and black/grey vertical lines represent S onset and D respectively. Black ($p<0.2$) and red ($p<0.05$) bars (above traces) mark timing of significant responses (mean of 3 frames) to the standard and the deviant. iii-iv) Population mean df/f (311 neurons, n=3 (P7-P9)) in response to the two stimuli TN (iii) and NT (iv). v) Distribution of CSI values for the population of L2/3 single neurons.

1756
Figure S15. A) i) the z-scores of responses in SD stimulus (20 reps, 4Hz) (left column) and
Swap (right column) for 3 single units in L4 (top 3 rows) and 3 single units in SP (bottom 3
rows) are shown (same units as in Fig. 4B, first 3 columns). A) ii) Mean z-scores for the
population of single units in L4 (top row) and that in SP (bottom row) are shown for the SD
stimulus (left) and the corresponding Swap (right). B) i-ii) Same as Ai-ii, for responses to SD
stimulus with 30 reps and at rate of 3.33 Hz.
1763
1764
1765
1766

Figure S16. Top and middle row show the mean CSI values of the population of L4 and SP neurons respectively for different choice of response latencies and response window sizes. Asterisks marks the response latency and response window size combination used in Fig. 4 CSI calculations. The three columns are for the 3 different CSIs used CSI, CSI(F) and CSI(P). Bottom row shows which combinations of response attribute have significantly higher CSI in SP than in L4 (brown, blue indicates NS).

Figure S17. A) 2 examples of cross-correlograms each for before ECO (top row) and after ECO (bottom wor) showing connections detected from SP to L4 (left) and L4 to SP (right). Blue and red colored lines indicate mean of cross-correlograms with errorbars and gray lines with errorbars show the same for shuffled data (see detailed methods). B) Bar plots summarize the data in Table S2 showing probability of connections from SP to L4 (left) and L4 to SP (right) at within 125 microns or greater distances at the 2 age groups before ECO and P12-13, after ECO).

Figure S18. A) Four example images of location (along cortical surface) of BFs on the cortical surface in the craniotomy of single units recorded in L4 at ages P28 and above (Fig. 5A single unit recording) identified in color (colorbar, bottom right) for four exposure groups EX2 and EX1 in the top row (SD exposure) and EX5 and EX6 in the bottom row (SS exposure). Broad tonotopy showing A1 and AAF are clear in the images. B) Bar plot shows % change in area of each of the preferred frequencies in A1 and AAF between control (EX0 group of mice) and exposed animals (EX5 group of mice).

1806
1807
1808
Figure S19. Similar plots as in Fig. 5E, for the profile of population mean normalized rates to
the relevant RF (color legend) for the 4 exposure groups (EX5 and EX6 left, EX2 and EX1
right) as a function of BF (re RF). Normalization was done by considering the maximum rate
in each tuning curve (rate at BF) to be 100%. Gray profile shows the same for the maximum
of 13 similar plots derived from the control data (C) considering each of the 13 frequencies
used as the RF.

1840
1841
1842

1843 Figure S20. Bar plot summarizes population mean response to RF for neurons with BF at RF
1844 of all 6 EX cases (2 bars each, red and green for SD exposure cases and 1 blue for SS). Mean
1845 of the 13 control cases shown in gray bar in the centre.

1846

1847

1848

1849

1850

1851

1852

1853

1854

References

1. Schreiner, C. E. & Polley, D. B. Auditory map plasticity: Diversity in causes and consequences. *Current Opinion in Neurobiology* vol. 24 143–156 (2014).
2. Hensch, T. K. Critical period plasticity in local cortical circuits. *Nature Reviews Neuroscience* (2005) doi:10.1038/nrn1787.
3. de Villers-Sidani, E., Chang, E. F., Bao, S. & Merzenich, M. M. Critical Period Window for Spectral Tuning Defined in the Primary Auditory Cortex (A1) in the Rat. *J. Neurosci.* (2007) doi:10.1523/JNEUROSCI.3227-06.2007.
4. Takesian, A. E., Bogart, L. J., Lichtman, J. W. & Hensch, T. K. Inhibitory circuit gating of auditory critical-period plasticity. *Nat. Neurosci.* (2018) doi:10.1038/s41593-017-0064-2.
5. Barkat, T. R., Polley, D. B. & Hensch, T. K. A critical period for auditory thalamocortical connectivity. *Nat. Neurosci.* (2011) doi:10.1038/nn.2882.
6. Zhou, X. & Merzenich, M. M. Enduring effects of early structured noise exposure on temporal modulation in the primary auditory cortex. *Proc. Natl. Acad. Sci. U. S. A.* (2008) doi:10.1073/pnas.0800009105.
7. Zhang, L. I., Bao, S. & Merzenich, M. M. Persistent and specific influences of early acoustic environments on primary auditory cortex. *Nat. Neurosci.* **4**, 1123–1130 (2001).
8. Luhmann, H. J., Kirischuk, S. & Kilb, W. The Superior Function of the Subplate in Early Neocortical Development. *Front. Neuroanat.* **12**, (2018).
9. Kanold, P. O. & Luhmann, H. J. The Subplate and Early Cortical Circuits. *Annu. Rev. Neurosci.* (2010) doi:10.1146/annurev-neuro-060909-153244.
10. Kanold, P. O., Kara, P., Reid, R. C. & Shatz, C. J. Role of subplate neurons in functional maturation of visual cortical columns. *Science (80-.).* (2003) doi:10.1126/science.1084152.
11. Kanold, P. O. & Shatz, C. J. Subplate Neurons Regulate Maturation of Cortical Inhibition and Outcome of Ocular Dominance Plasticity. *Neuron* (2006) doi:10.1016/j.neuron.2006.07.008.
12. Ghosh, A. & Shatz, C. J. Involvement of subplate neurons in the formation of ocular dominance columns. *Science (80-.).* (1992) doi:10.1126/science.1542795.
13. Ghosh, A., Antonini, A., McConnell, S. K. & Shatz, C. J. Requirement for subplate neurons in the formation of thalamocortical connections. *Nature* (1990) doi:10.1038/347179a0.
14. McConnell, S. K., Ghosh, A. & Shatz, C. J. Subplate neurons pioneer the first axon pathway from the cerebral cortex. *Science (80-.).* (1989) doi:10.1126/science.2475909.
15. Chang, E. F., Bao, S., Imaizumi, K., Schreiner, C. E. & Merzenich, M. M. Development of spectral and temporal response selectivity in the auditory cortex. *Proc. Natl. Acad. Sci. U. S. A.* (2005) doi:10.1073/pnas.0508239102.
16. Ulanovsky, N., Las, L. & Nelken, I. Processing of low-probability sounds by cortical neurons. *Nat. Neurosci.* (2003) doi:10.1038/nn1032.
17. Ghosh, A. & Shatz, C. J. A role for subplate neurons in the patterning of connections from thalamus to neocortex. *Development* (1993).
18. Zhao, C., Kao, J. P. Y. & Kanold, P. O. Functional Excitatory Microcircuits in Neonatal

Cortex Connect Thalamus and Layer 4. *J. Neurosci.* (2009) doi:10.1037/pha0000116.

19. Viswanathan, S., Bandyopadhyay, S., Kao, J. P. Y. & Kanold, P. O. Changing Microcircuits in the Subplate of the Developing Cortex. *J. Neurosci.* **32**, 1589–1601 (2012).

20. Wess, J. M., Isaiah, A., Watkins, P. V & Kanold, P. O. Subplate neurons are the first cortical neurons to respond to sensory stimuli. *Proc. Natl. Acad. Sci.* (2017) doi:10.1073/pnas.1710793114.

21. Pérez-González, D. & Malmierca, M. S. Adaptation in the auditory system: an overview. *Front. Integr. Neurosci.* (2014) doi:10.3389/fnint.2014.00019.

22. Nieto-Diego, J. & Malmierca, M. S. Topographic Distribution of Stimulus-Specific Adaptation across Auditory Cortical Fields in the Anesthetized Rat. *PLoS Biol.* (2016) doi:10.1371/journal.pbio.1002397.

23. Bandyopadhyay, S., Shamma, S. A. & Kanold, P. O. Dichotomy of functional organization in the mouse auditory cortex. *Nat. Neurosci.* (2010) doi:10.1038/nn.2490.

24. Issa, J. B. *et al.* Multiscale Optical Ca2+Imaging of Tonal Organization in Mouse Auditory Cortex. *Neuron* (2014) doi:10.1016/j.neuron.2014.07.009.

25. Sawatari, H. *et al.* Identification and characterization of an insular auditory field in mice. *Eur. J. Neurosci.* (2011) doi:10.1111/j.1460-9568.2011.07926.x.

26. Barbour, D. L. & Wang, X. Contrast tuning in auditory cortex. *Science (80-.).* (2003) doi:10.1126/science.1080425.

27. Mill, R., Coath, M., Wennekers, T. & Denham, S. L. Abstract stimulus-specific adaptation models. *Neural Computation* (2011) doi:10.1162/NECO_a_00077.

28. Clancy, B., Darlington, R. B. & Finlay, B. L. Translating developmental time across mammalian species. *Neuroscience* (2001) doi:10.1016/S0306-4522(01)00171-3.

29. Chang, A. *et al.* Specific Influences of Early Acoustic Environments on Cochlear Hair Cells in Postnatal Mice. *Neural Plast.* (2018) doi:10.1155/2018/5616930.

30. Babola, T. A. *et al.* Homeostatic Control of Spontaneous Activity in the Developing Auditory System. *Neuron* (2018) doi:10.1016/j.neuron.2018.07.004.

31. Anthwal, N. & Thompson, H. The development of the mammalian outer and middle ear. *J. Anat.* (2016) doi:10.1111/joa.12344.

32. Huangfu, M. & Saunders, J. C. Auditory development in the mouse: Structural maturation of the middle ear. *J. Morphol.* (1983) doi:10.1002/jmor.1051760302.

33. Kim, K. X. & Fettiplace, R. Developmental changes in the cochlear hair cell mechanotransducer channel and their regulation by transmembrane channel-like proteins. *J. Gen. Physiol.* (2013) doi:10.1085/jgp.201210913.

34. Lelli, A., Asai, Y., Forge, A., Holt, J. R. & Géléoc, G. S. G. Tonotopic gradient in the developmental acquisition of sensory transduction in outer hair cells of the mouse cochlea. *J. Neurophysiol.* (2009) doi:10.1152/jn.00136.2009.

35. Abe, T. *et al.* Developmental expression of the outer hair cell motor prestin in the mouse. *J. Membr. Biol.* (2007) doi:10.1007/s00232-007-9004-5.

36. Polley, D. B., Thompson, J. H. & Guo, W. Brief hearing loss disrupts binaural integration during two early critical periods of auditory cortex development. *Nat. Commun.* (2013) doi:10.1038/ncomms3547.

37. Ruebsamen, R. & Lippe, W. The development of cochlear function. in *Development of the auditory system* (eds. Rubel, E., Poppe, A. & Fay, R.) 193–270 (Springer, 1997).

38. Shnerson, A. & Pujol, A. Development: anatomy, electrophysiology, and behavior. in
The auditory psychobiology of the mouse (ed. Willot, J.) 395–425 (Charles C. Thomas,
1983).

39. Pujol, R., Lavigne-Rebillard, M. & Lenoir, M. Development of sensory and neural
structures in the mammalian cochlea. in *Development of the auditory system* (eds.
Rubel, E., Popper, A. & Fay, R.) 146–192 (Spinger, 1997).

40. Tritsch, N. X. *et al.* Calcium action potentials in hair cells pattern auditory neuron
activity before hearing onset. *Nat. Neurosci.* (2010) doi:10.1038/nn.2604.

41. Mill, R., Coath, M., Wennekers, T. & Denham, S. L. A neurocomputational model of
stimulus-specific adaptation to oddball and markov sequences. *PLoS Comput. Biol.*
(2011) doi:10.1371/journal.pcbi.1002117.

42. Yarden, T. S. & Nelken, I. Stimulus-specific adaptation in a recurrent network model
of primary auditory cortex. *PLoS Comput. Biol.* (2017)
doi:10.1371/journal.pcbi.1005437.

43. Hromádka, T., DeWeese, M. R. & Zador, A. M. Sparse representation of sounds in the
unanesthetized auditory cortex. *PLoS Biol.* (2008) doi:10.1371/journal.pbio.0060016.

44. Lewicki, M. S. Efficient coding of natural sounds. *Nat. Neurosci.* (2002)
doi:10.1038/nn831.

45. Smith, E. C. & Lewicki, M. S. Efficient auditory coding. *Nature* (2006)
doi:10.1038/nature04485.

46. Sun, Y. J. *et al.* Fine-tuning of pre-balanced excitation and inhibition during auditory
cortical development. *Nature* **465**, 927–931 (2010).

47. Blundon, J. A. *et al.* Restoring auditory cortex plasticity in adult mice by restricting
thalamic adenosine signaling. *Science (80-.).* (2017) doi:10.1126/science.aaf4612.

48. Vickers, E. D. *et al.* Parvalbumin-Interneuron Output Synapses Show Spike-Timing-
Dependent Plasticity that Contributes to Auditory Map Remodeling. *Neuron* (2018)
doi:10.1016/j.neuron.2018.07.018.

49. Courtney E. Yaeger, Dario Ringach, J. T. T. Neuromodulatory control of localized
dendritic spiking in critical period cortex. *Nature* **567**, 100–104 (2019).

50. Mehler, J. *et al.* A precursor of language acquisition in young infants. *Cognition* (1988)
doi:10.1016/0010-0277(88)90035-2.

51. Voegtline, K. M., Costigan, K. A., Pater, H. A. & DiPietro, J. A. Near-term fetal response
to maternal spoken voice. *Infant Behav. Dev.* (2013)
doi:10.1016/j.infbeh.2013.05.002.

52. Portfors, C. V., Roberts, P. D. & Jonson, K. Over-representation of species-specific
vocalizations in the awake mouse inferior colliculus. *Neuroscience* (2009)
doi:10.1016/j.neuroscience.2009.04.056.

53. Wang, X. & Kadia, S. C. Differential Representation of Species-Specific Primate
Vocalizations in the Auditory Cortices of Marmoset and Cat. *J. Neurophysiol.* (2001)
doi:10.1152/jn.2001.86.5.2616.

54. Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic*
Coordinates. São Paulo, Academic Press (2012). doi:10.1364/OE.20.020998.

55. Torii, M. & Levitt, P. Dissociation of corticothalamic and thalamocortical axon
targeting by an EphA7-mediated mechanism. *Neuron* (2005)
doi:10.1016/j.neuron.2005.09.021.

56. Osheroff, H. & Hatten, M. E. Gene expression profiling of preplate neurons destined

1993 for the subplate: Genes involved in transcription, axon extension, neurotransmitter
1994 regulation, steroid hormone signaling, and neuronal survival. *Cereb. Cortex* (2009)
1995 doi:10.1093/cercor/bhp034.

1996 57. Hoerder-Suabedissen, A. *et al.* Novel markers reveal subpopulations of subplate
1997 neurons in the murine cerebral cortex. *Cereb. Cortex* (2009)
1998 doi:10.1093/cercor/bhn195.

1999 58. Gee, J. M. *et al.* Imaging Activity in Neurons and Glia with a Polr2a-Based and Cre-
2000 Dependent GCaMP5G-IRES-tdTomato Reporter Mouse. *Neuron* (2014)
2001 doi:10.1016/j.neuron.2014.07.024.

2002