Visual feature tuning of superior colliculus neural reafferent responses after fixational microsaccades

Fatemeh Khademi[1,2], Chih-Yang Chen[1], and Ziad M. Hafed[1,2]*

[1]Werner Reichardt Centre for Integrative Neuroscience, Tuebingen University, Tuebingen, Germany 72076
[2]Hertie Institute for Clinical Brain Research, Tuebingen University, Tuebingen, Germany 72076

*Correspondence to: ziad.m.hafed@cin.uni-tuebingen.de

Abbreviated title:

Visual coding by microsaccades in midbrain

Corresponding author address:

Ziad M. Hafed
Werner Reichardt Centre for Integrative Neuroscience
and
Hertie Institute for Clinical Brain Research
Otfried-Mueller Str. 25
Tuebingen, 72076
Germany
Phone: +49 7071 29 88819

Number of words:
Abstract: 250
Introduction: 650
Discussion: 1361

Number of figures:
10

Number of pages:
49

Acknowledgements:
We were funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation; projektnummer 276693517—SFB 1233). We were also funded by the Werner Reichardt Centre for Integrative Neuroscience (DFG Excellence Cluster: EXC 307).

Conflicts of Interests:
None

Abstract

The primate superior colliculus (SC) is causally involved in microsaccade generation. Moreover, visually-responsive SC neurons across this structure's topographic map, even at peripheral eccentricities much larger than the tiny microsaccade amplitudes, exhibit significant modulations of evoked response sensitivity when stimuli appear peri-microsaccadically. However, during natural viewing, visual stimuli are normally stably present in the environment and are only shifted on the retina by eye movements. Here we investigated this scenario for the case of microsaccades, asking whether and how SC neurons respond to microsaccade-induced image jitter. We recorded neural activity from two male rhesus macaque monkeys. Within the response field (RF) of a neuron, there was a stable stimulus consisting of a grating of one of three possible spatial frequencies. The grating was stable on the display, but microsaccades periodically jittered the retinotopic RF location over it. We observed clear short-latency visual reafferent responses after microsaccades. These responses were weaker, but earlier (relative to new fixation onset after microsaccade end), than responses to sudden stimulus onsets without microsaccades. The reafferent responses clearly depended on microsaccade amplitude, as well as microsaccade direction relative to grating orientation. Our results indicate that one way for microsaccades to influence vision is through modulating how the spatio-temporal landscape of SC visual neural activity represents stable stimuli in the environment. Such representation strongly depends on the specific pattern of temporal luminance modulations expected from the relative relationship between eye movement vector (size and direction), on the one hand, and spatial visual pattern layout, on the other.

Significance statement

Despite their small sizes and slow speeds, microsaccades during fixation still jitter retinal images of stable visual objects in the environment. This means that retinotopically organized neurons' response fields experience temporal modulations of luminance as a visual pattern is shifted by microsaccades. We investigated the impacts of such modulations on superior colliculus (SC) activity. We found that SC neurons exhibit visual reafferent activity bursts with very short latency after microsaccade end. Moreover, these bursts reflect not only the spatial luminance profiles of the patterns, but also how such luminance profiles are temporally modulated by eye movement size and direction. These results indicate that the SC continuously represents visual patterns, even as they are jittered retinotopically by the smallest possible saccades.

Introduction

Microsaccades (reviewed in Martinez-Conde et al., 2004; Engbert, 2006; Rolfs, 2009; Hafed, 2011; Hafed et al., 2015; Ahissar et al., 2016; Poletti and Rucci, 2016; Krauzlis et al., 2017) are small saccades that occur periodically during prolonged gaze fixation. In comparison to larger saccades, with peak velocities as high as ~800 deg/s, microsaccades (having peak velocities well below 100 deg/s) decidedly cause milder disruptions of retinal images. Nonetheless, these fixational eye movements still jitter retinal images, and they are therefore expected to influence visual neural activity due to the sequence of motion blur followed by stable visual stimulus that they cause. Indeed, microsaccades are associated with post-movement visual reafferent responses in a variety of brain areas (Leopold and Logothetis, 1998; Martinez-Conde et al., 2000; Snodderly et al., 2001; Martinez-Conde et al., 2002; Kagan et al., 2008; Bosman et al., 2009; Herrington et al., 2009). However, the bulk of past work has focused primarily on the mere presence of reafferent responses, but not necessarily on their properties. While such presence may be relevant for issues like the refreshing of retinal images by microsaccades, perhaps to help in counteracting visual fading (Martinez-Conde et al., 2004), understanding the properties of post-microsaccadic visual reafference is additionally important for clarifying the mechanisms of visual neural coding when the sensor for vision (the eye) is continually mobile (Snodderly et al., 2001; Rucci et al., 2007; Kagan et al., 2008; Kuang et al., 2012).

In this study, we explored the properties of visual reafference after microsaccades in superior colliculus (SC) visual activity. The SC has historically been studied in primate animal models primarily from a saccade motor control perspective (Robinson, 1972; Gandhi and Katnani, 2011; Basso and May, 2017; Krauzlis et al.,

2017). Indeed, the SC is causally involved in microsaccade generation (Hafed et al.,

2009; Hafed and Krauzlis, 2012; Hafed et al., 2013; Willeke et al., 2019). However,

the SC, even in primates, also exhibits significant visual responses (Goldberg and

Wurtz, 1972), and recent results have revealed visual pattern analysis capabilities of

SC neurons (Hafed and Chen, 2016; Hall and Colby, 2016; Herman and Krauzlis,

2017; Chen and Hafed, 2018; Chen et al., 2018; Hafed, 2018; Chen et al., 2019).

This leaves the question of how the SC represents continuously presented stimuli (as

opposed to stimulus onsets) open, and how such representation is modulated by

microsaccades.

In all of our previous investigations of SC neural modulations by microsaccades, we

presented stimuli suddenly around the time of these eye movements (Hafed and

Krauzlis, 2010; Chen et al., 2015; Hafed et al., 2015; Bellet et al., 2017; Chen and

Hafed, 2017; Chen et al., 2019). Whether and how SC neurons would react to stable

visual stimuli being jittered on the retina by microsaccades are questions that have

not yet been explored. Investigating these questions would be important to support

the notion that the SC may be viewed as much as an early visual area as it is viewed

as a late motor control area, and it would also be in line with a potential role for the

SC in signaling visual salience and priority in the environment (Veale et al., 2017;

White et al., 2017b; White et al., 2017a; White et al., 2019).

In our experiments, we exploited our recent observations of visual feature tuning of

SC neurons for different spatial frequencies (Chen et al., 2018), and we compared

such tuning to the case when the stimuli were stably present in the environment and

only jittered retinotopically by microsaccades. We analyzed the relative relationships

between microsaccade sizes and directions, on the one hand, and grating spatial

frequencies, on the other; we found that the specific pattern of temporal luminance

modulations caused within a given RF by a given microsaccade dictates how the

neuron "updates" its post-microsaccadic representation of the otherwise stable

stimulus. This suggests that the SC can contribute to visual coding of scenes during

active visual fixation.

Materials and Methods

Ethics approvals

All monkey experiments were approved by ethics committees at the Regierungspräsidium Tübingen. The experiments were in line with the European Union directives, and the German laws, governing animal research.

Laboratory setups

Monkey neurophysiology experiments were performed in the same laboratory environment as that described in (Chen et al., 2015; Chen et al., 2018; Chen et al., 2019).

Animal preparation

We collected behavioral and neural data from 2 adult, male rhesus macaques (macaca mulatta). Monkeys N and P (aged 7 years and weighing 8 and 7 kg, respectively) were implanted with scleral search coils to allow measuring eye movements using the electromagnetic induction technique (Fuchs and Robinson, 1966; Judge et al., 1980). The monkeys were also implanted with a head holder to stabilize their head during the experiments, as well as recording chambers to access the right and left SC's. Details on all implant surgeries were provided earlier (Chen and Hafed, 2013; Chen et al., 2015).

Monkey behavioral tasks

Each monkey performed a simple fixation task. Prior to running this main task (which we describe below), we first assessed the visual response field (RF) of a recorded SC neuron using standard fixation and saccade behavioral paradigms described earlier (Chen et al., 2015; Chen and Hafed, 2018; Chen et al., 2018; Chen et al.,

2019). Briefly, the monkey sat in a dark room in front of a gray display with luminance

21 cd/m^2. A white (72 cd/m^2) spot (square of 8.5 x 8.5 min arc) was then presented at

different display locations while the monkey was maintaining gaze fixation on a

similar spot. Visual responses to the appearing spot were used to assess the visual

RF location and boundaries. In some variants, the monkey also made a later

saccade to the stimulus.

We then ran the main task of this study. A similar fixation spot to that described

above was first presented at the center of the display for the monkey to fixate. After

the monkey fixated the spot for ~490 ms (+/- 80 ms s.d.), a high-contrast (100%)

Gabor grating appeared at a location centered on the recorded neuron's RF. The

radius of the grating was tailored to the size of the RF, and the spatial frequency

could be 0.56, 2.22, or 4.44 cycles per degree (cpd). These spatial frequencies are

known to be effective in driving visual responses in the SC (e.g. Chen et al., 2018).

The grating persisted for ~1300-3000 ms until trial end. If the monkey maintained

fixation until trial end, the monkey was rewarded with a juice/water drop.

In all experiments, we used a vertical Gabor grating. While the rhesus macaque SC

does exhibit orientation tuning (Chen and Hafed, 2018), our experience has been

that tuning curves are broad enough to include notable responses for vertical

gratings as well. We thus used vertical gratings in a variety of our recent studies

(Chen et al., 2015; Chen and Hafed, 2017; Chen et al., 2018), and there were

measurable visual responses. Because microsaccades vary in direction when they

occur, this still allowed us to explore the relative relationships between movement

vector direction and stimulus pattern orientation (see Results).

Behavioral analyses

We detected saccades and microsaccades using established methods in our laboratory (Chen and Hafed, 2013; Bellet et al., 2019). We manually inspected each trial to correct for false alarms or misses by the automatic algorithms, which were rare. We also marked blinks or noise artifacts for later removal. We characterized microsaccade times in order to align neural activity to them (for analyzing visual reafferent responses). We also characterized microsaccade amplitudes (i.e. sizes) and directions in order to explore interactions between movement vector and stimulus spatial frequency.

Neural firing rate analyses

We analyzed data from 55 well-isolated single neurons (23 from monkey N and 32 from monkey P). Our criteria for analyzing neurons were the presence of a visual evoked response for at least one of the spatial frequencies that we presented, as well as the availability of >10 trials per presented spatial frequency in a given recording. Our numbers of neurons are typical of recent SC studies exploring visual responses (White et al., 2017a; White et al., 2019), and the response profiles that we observed were sufficiently homogeneous, and consistent with our other recorded tasks in the SC (Hafed and Chen, 2016; Chen and Hafed, 2017; Chen et al., 2018), to make our sample of neurons representative of SC neurons in general.

The neurons described in this study included both purely visual and visual-motor neurons, but we did not notice substantial differences in the present analyses (focused on visual responses only) between the two types. We therefore did not separate analyses for these two functional types of SC neurons. We did, however, ensure that none of our neurons exhibited movement-related discharge for

microsaccade generation (Hafed et al., 2009; Hafed and Krauzlis, 2012; Willeke et

al., 2019). Such discharge would come at the time of potential visual reafferent

responses and mask such responses. Given that our recorded neurons (see Results)

were not extremely foveal (Chen et al., 2019), exclusion of microsaccade-related

discharge was trivially easy.

We used firing rates from the RF mapping tasks alluded to above in order to confirm

that our placement of gratings during the experiments was valid. We then analyzed

the neural data in the main task. We converted spike times to firing rates as we have

done in our prior studies (Chen et al., 2015). We then performed subsequent

analyses on firing rates, as we describe next.

To measure stimulus-evoked visual responses, we aligned firing rates to grating

onset. We grouped trials based on condition (e.g. 0.56 cpd gratings), and we ensured

that there were no microsaccades occurring between -100 and 150 ms relative to

stimulus onset. This was done in order to avoid other influences of microsaccades on

SC evoked visual responses, which we have characterized elsewhere (Hafed and

Krauzlis, 2010; Chen et al., 2015; Chen and Hafed, 2017). Specifically, when a

stimulus onset occurs in the vicinity of a microsaccade onset, the subsequent

stimulus-evoked response in SC neurons is modulated; our goal here was to

characterize the unmodulated stimulus evoked-response and relate it to visual

reafferent responses when a microsaccade jittered the image of a stably presented

grating over the recorded RF.

When summarizing measurements across neurons, we measured peak firing rate in

a given condition as the peak of the average firing rate of the condition across trials.

That is, we grouped all trials of the same type, and we then averaged firing rates

aligned on stimulus onset. We then defined a measurement interval 0-150 ms after

stimulus onset during which we searched for a peak in the average firing rate curve.

The peak value and peak time provided our estimate of peak firing rate of the neuron

and visual response latency of the neuron.

When we visualized population firing rate curves across neurons in a given condition,

we first normalized each neuron's firing rate to a maximum of 1 before averaging the

curves of the pooled neurons. Specifically, in a given condition (e.g. onset of a

grating of 0.56 cpd spatial frequency), we pooled all trials and obtained an average

firing rate curve. If the response to this condition was to be designated as the

normalization reference, we then divided the firing rate curves of the same neuron in

all individual trials (and all individual conditions) by the peak of this average firing rate

curve. This way, all firing rates of the neuron across all trials and conditions were

normalized to the firing rate of a given condition (e.g. 0.56 cpd). In different analyses,

we either normalized by the peak visual evoked response to 0.56 cpd gratings (e.g.

Fig. 4) or by the peak visual evoked response to the preferred spatial frequency of

the neuron (e.g. Fig. 5). In this case, neuron "preference" was defined based on

response strength (i.e. a preferred stimulus caused the strongest evoked response);

in a lot of neurons, the preferred spatial frequency was indeed 0.56 cpd (Chen et al.,

2018). It should also be noted here that in all analyses in which we presented

normalized average firing rates, we also made sure to report raw activity

measurements on a per individual neuron basis (e.g. Figs. 4, 5, 7, 8).

To statistically determine whether a neuron had a significant visual response for a

given spatial frequency compared to baseline activity, we defined two measurement

intervals. The first was 0-150 ms before stimulus onset and served as the baseline

firing rate of the neuron. The second was 0-150 ms after stimulus onset and served

as the visual evoked epoch. Across repetitions of a given condition, if the average

firing rate in the second interval was statistically higher (at a p-value of <0.01) than

the average firing rate in the first interval (Wilcoxon rank sum test if numbers of

repetitions were <16; t-test otherwise), then the neuron possessed a statistically

significant visual response.

For some analyses, we were also interested in whether individual neurons

possessed statistically significant tuning to an individual spatial frequency (from

among the three that we sampled in our experiments). To do that, we defined an

interval of 0-150 ms after stimulus onset. For each neuron and spatial frequency, we

estimated the peak firing rate per trial. We then performed a one-way ANOVA with

the main factor being spatial frequency. If the ANOVA showed a significant main

effect of spatial frequency (at a p-value of <0.05), we also performed a post-hoc t-test

(at a p-value of <0.01). This allowed us to group the neurons into four groups:

preferring the lowest spatial frequency (0.56 cpd); preferring the intermediate spatial

frequency (2.22 cpd); preferring the highest spatial frequency (2.22 cpd); or not

preferring any of the three presented spatial frequencies (no significant main effect in

the one-way ANOVA).

To measure visual reafferent responses, we used the same procedures above, but

we aligned firing rates to microsaccade end (that is, new fixation onset). When we

normalized firing rate curves (see above), we always normalized to visual evoked,

not visual reafferent, responses. This allowed us to quantitatively compare the

strength of visual reafferent SC responses relative to stimulus-driven visual evoked

responses. Also, when statistically assessing whether there was a significant visual

reafferent response, we used the same concept of two measurement intervals as

described above. However, in this case, the "baseline" interval was 50-150 ms before

microsaccade onset, and the "response" interval was 0-100 ms after microsaccade

end. The latter interval was early (compared to visual evoked measurement intervals

above) because the visual reafferent responses had shorter latencies from

microsaccade end (see Results).

In all analyses, we only analyzed visual reafferent responses for microsaccades that

occurred when the grating was already stably present on the display (that is, already

present inside the neuron's RF; see Results). Specifically, we only considered

microsaccades occurring from 300 ms after grating onset until 100 ms before trial

end. The former time value was to ensure that the visual evoked response after

sudden stimulus onset had subsided; the latter time value was to ensure that we had

enough trial repetitions across all collected data.

To further understand and support our observations from spiking activity as assessed

with firing rate measurements, we also analyzed local field potentials (LFP's) as a

highly sensitive population measure of SC visual tuning properties around the

recording electrodes (Ikeda et al., 2015; Hafed and Chen, 2016; Chen and Hafed,

2017, 2018; Chen et al., 2018); LFP's were additionally shown to be useful in

analyses from other brain areas beyond the SC (e.g. Neupane et al., 2016). We used

similar procedures to those described earlier (Hafed and Chen, 2016; Chen and

Hafed, 2017, 2018; Chen et al., 2018). Briefly, we started with the wide-band signals

digitized at 40 KHz by our data acquisition system, which also initially applied a

hardware filter passing the range of 0.7 Hz to 6 KHz frequencies. We then removed

electrical power line noise artifacts using IIR notch filters at 50, 100, and 150 Hz.

Finally, we applied a zero-phase-lag low pass filter with cutoff frequency of 300 Hz,

and we down sampled the resulting data to 1 KHz. This gave us continuous signals

of LFP modulations analogous to continuous firing rate curves obtained from

individual spike times. For each recording site, we averaged the individual-trial LFP

curves and aligned them to either stimulus onset or microsaccade onset/end.

Experimental design and statistical analysis

Our experiments underwent within-neuron analyses, and we report summaries from

each neuron individually. We also enforced minimum numbers of trials for analyses

as described above. When separating neurons according to their preferred spatial

frequencies (e.g. Fig. 5), we only showed spatial frequencies for which there were

enough neurons preferring them to allow reasonable interpretations of the results (i.e.

with enough sampled neurons).

Given the strong congruence of the visual evoked responses (to stimulus onsets)

with our recent observations on SC spatial frequency tuning properties (Chen et al.,

2018), we are confident that our numbers of neurons are sufficient to represent the

overall SC population and summarize our observations on visual reafferent

responses. Moreover, the use of two monkeys is viewed as a minimal compromise

between using animals in research, on the one hand, and ensuring replicability of

results across larger populations of research subjects, on the other.

We report measures of variance for all analyses and figures, and we also show

individual neuron results for all analyses. Statistical tests are reported at relevant

portions of Results, and in separate analyses (not shown), we also further

investigated all of our statistical tests using alternative methods involving

randomization tests.

When reporting p-values, we used the following conventions for purposes of clarity: if

a p-value was less than 0.0001, we reported it simply as $p<0.0001$; if a p-value was

higher, we reported its exact value.

Data availability

All data presented in this paper are stored in institute computers and are available

upon reasonable request.

Results

We explored the visual consequences of fixational eye movements on neural activity

in the rhesus macaque SC. We did so by presenting a Gabor grating of a given

orientation and spatial frequency at a visual location corresponding to the visual RF

location of a recorded neuron (Fig. 1a). The grating was stationary and persisted on

the display for up to approximately 3000 ms, but it was continuously moved relative

to the recorded neuron's retinotopic RF by incessant fixational eye movements.

Across sessions, we recorded from primarily parafoveal neurons in both the right and

left SC (average preferred eccentricity: 3.5 deg +/- 0.39 deg s.e.m.; n=55 neurons),

and the grating radii were tailored in each experiment according to the recorded RF

size and location. We explored the visual reafferent response after a microsaccade

jittered the grating over the recorded neuron's RF. In all cases, we analyzed the

activity of visually-responsive SC neurons, whether they also had a saccade-related

motor burst or not, but we ensured that none of our neurons had a motor burst for

microsaccades in any direction (Hafed et al., 2009; Willeke et al., 2019). Otherwise, it

would have been difficult to differentiate between whether neural activity after

microsaccades was related to visual reafference or to the microsaccade-related

discharge itself.

Microsaccades are associated with a visual reafferent response in SC

We first compared visual reafferent responses after microsaccades to responses

when the grating first appeared inside a neuron's RF. As shown in Fig. 1a, each

monkey fixated a central fixation spot (Materials and Methods), and we presented a

vertical grating filling the RF; RF size was initially assessed with small spot stimuli in

initial RF mapping tasks (Materials and Methods). The example trial shown in Fig. 1b

demonstrates how stimulus onset elicited a short-latency visual response, and how

the neuron continued to fire occasionally afterwards, and particularly after

microsaccades. We characterized the post-stimulus and post-microsaccadic neural

responses. The great majority of microsaccades that we analyzed had a radial

amplitude of <0.2 deg (<12 min arc; Fig. 1c), and they were associated with peak

velocities significantly less than approximately 70 deg/s (medians: 10 deg/s and 23

deg/s for monkeys P and N, respectively).

393
Figure 1 Task design. (a) The monkeys fixated, and we presented an RF stimulus comprising of Gabor
gratings having different spatial frequencies (schematized below the top panel). **(b)** Example eye
position trace (blue) and neural spiking activity (each black tick mark represents an action potential)
from a sample recorded SC neuron. Shortly after stimulus onset, there was a burst of action potentials
(the visual evoked response). Subsequently, the neuron occasionally emitted activity, with bursts after
microsaccades (visual reafferent responses); the black upward triangles indicate microsaccade onsets.
We analyzed such bursts during a sustained fixation interval with the stimulus stably present over the
RF (Materials and Methods). **(c)** In both monkeys, microsaccade amplitudes that we analyzed were
primarily smaller than 0.2 deg (12 min arc), as shown by the histograms of amplitude probabilities.

For each presented spatial frequency, we plotted the visual evoked response of the

same example neuron of Fig. 1 (Fig. 2a), as well as the visual reafferent response

after microsaccades (Fig. 2b); Figure 2c shows the same visual reafferent response

as in Fig. 2b, but now aligned to microsaccade onset rather than microsaccade end.

For the visual reafferent response (Fig. 2b, c), we only analyzed microsaccades for

which the grating was already displayed to the monkey. That is, we picked a

sustained time interval starting 300 ms after grating onset and ending 100 ms before

trial end, and we only analyzed microsaccades during this interval. This interval

started after the initial stimulus-driven visual evoked response of the neuron had

subsided, such that the visual reafferent response was analyzed when there was

already a stable stimulus inside the RF (the microsaccades were small enough to not

move the presented stimulus outside the RF when they occurred).

420
Figure 2 Activity of an example SC neuron after stimulus onset and after microsaccades. (a)
Spiking activity as a function of time after stimulus onset. Each tick mark in the top panel represents an
action potential, and each row of action potentials represents a trial. Trials were sorted by the spatial
frequency of the RF stimulus. The bottom panel shows the same data but now plotted as firing rate
curves. Error bars denote s.e.m. across trials. The neuron had a stronger and earlier visual burst for the
lowest spatial frequency. The numbers of trials per spatial frequency are shown in the top panel. **(b)** For
each microsaccade occurring while the grating was visible (that is, in a sustained interval after grating
onset; Materials and Methods), we aligned the same neuron's activity to microsaccade end (labeled
"offset" in the figure). The neuron emitted a visual reafferent burst, which also depended on spatial
frequency. Note how the visual reafferent response (relative to new fixation onset after a microsaccade)
occurred earlier than the stimulus evoked response in **a**. Error bars in this case denote s.e.m. across
the number of microsaccades analyzed. **(c)** Same as in **b** but now aligned to microsaccade onset.

As can be seen from the visual evoked response (Fig. 2a), this example neuron

preferred the lowest presented spatial frequency (0.56 cpd), emitting the strongest

peak firing rate. Moreover, the neuron's visual response latency also increased with

increasing spatial frequency, consistent with recent observations in the SC (Chen et

al., 2018). In the visual reafferent response (Fig. 2b, c), the neuron's activity was

significantly weaker, but it maintained the preference for the lowest spatial frequency.

Specifically, the peak firing rate after microsaccades for the lowest spatial frequency

(0.56 cpd) was 104 spikes/s when aligned to microsaccade end; this was 38% of the

peak firing rate for the same spatial frequency after stimulus onset. For the

intermediate spatial frequency (2.22 cpd), the peak firing rate of the neuron after

microsaccades was 62 spikes/s (31% of the stimulus evoked response), and the

peak firing rate for the highest spatial frequency (4.44 cpd) was 12 spikes/s (18% of

the stimulus evoked response). Note that due to variability of microsaccade

durations, aligning on microsaccade end (new fixation onset) rather than

microsaccade onset resulted in marginally weaker visual reafferent responses in the

former alignment relative to the latter (compare panels in Fig. 2b, c). For example, for

the lowest spatial frequency (0.56 cpd), the peak visual reafferent response was 104

spikes/s when aligned on microsaccade end and 112 spikes/s when aligned on

microsaccade onset.

Across the population, we measured the visual evoked response after stimulus onset

as the peak firing rate emitted by a neuron during an interval 0-150 ms after stimulus

onset (Materials and Methods). We also measured the visual reafferent response as

the peak firing rate during an interval 0-100 ms after microsaccade end; the intervals

were chosen based on the timings of bursts after stimulus onsets or microsaccades

observed across our population (Materials and Methods). Almost every single neuron

that we recorded from showed a weaker visual reafferent response than a visual

evoked response (Fig. 3a-c). In fact, the visual reafferent response was often

seemingly absent (especially for the grating with the highest spatial frequency), even

when a neuron responded after stimulus onset.

To statistically test whether the visual reafferent response was present or not in a

given neuron, we compared, for each neuron, average firing rate in the interval 0-100

ms after microsaccade end (the visual reafference epoch) to average firing rate in the

interval 50-150 ms before microsaccade onset (the baseline epoch). If the post-

microsaccade firing rate across repetitions was higher than the pre-microsaccade

firing rate at a p-value of <0.01 (assessed using a t-test or ranksum test; see

Materials and Methods), we deemed the neuron to possess a significant visual

reafferent response. Otherwise, the neuron lacked a significant visual reafferent

response. The neurons colored in gray in Fig. 3a-c were the neurons without any

significant visual reafferent response. For the lowest spatial frequency (0.56 cpd; Fig.

3a), these non-responding neurons comprised 36% of our recorded population

(20/55); for the intermediate spatial frequency (2.22 cpd; Fig. 3b), this fraction was

42% (23/55); and, finally, for the highest spatial frequency (4.44 cpd; Fig. 3c), 71%

(39/55) of our recorded neurons showed no visual reafferent response after

microsaccades. Note that for the visual evoked response after stimulus onset, all

neurons that we analyzed showed a response for all spatial frequencies. Thus, the

reafferent response was sometimes weak enough to simply disappear completely.

However, this might be expected: the highest spatial frequency (which showed the

least likelihood of significant visual reafferent response) was anyway associated with

the weakest visual evoked response overall (compare the x-axis ranges of Fig. 3a-c)

(Chen et al., 2018).

490
Figure 3 Properties of visual evoked and visual reafferent SC responses across the population.
(a) For the lowest spatial frequency (0.56 cpd), we plotted peak firing rate after microsaccade end (the visual reafferent response) as a function of peak firing rate after stimulus onset (the visual evoked response). The gray symbols denote the neurons for which the visual reafferent response was not present (that is, statistically indistinguishable from baseline activity without microsaccades; Materials and Methods). **(b, c)** Same as **a** for the intermediate (**b**) and highest (**c**) spatial frequencies (2.22 cpd and 4.44 cpd, respectively). Visual reafferent responses were always weaker than visual evoked responses (p<0.0001, <0.0001, and <0.0001 across all neurons; t-test comparing reafferent to evoked responses for low, intermediate, and high spatial frequencies, respectively). **(d-f)** Same as **a-c** but now plotting the time to peak firing rate instead of the value of peak firing rate. For the visual reafferent responses, time to peak firing rate was measured from microsaccade end (p<0.0001, <0.0001, and <0.0001 across all neurons for low, intermediate, and high spatial frequencies, respectively). **(g-i)** Same as **d-f**, but measuring time to peak firing rate in the visual reafferent responses from microsaccade onset instead of microsaccade end (p=0.006, <0.0001, and 0.3 for low, intermediate, and high spatial frequencies, respectively).

The results above suggest, so far, that the SC exhibits visual reafferent responses

after microsaccades, similar to other early visual areas (Leopold and Logothetis,

1998; Martinez-Conde et al., 2000; Snodderly et al., 2001; Martinez-Conde et al.,

2002; Kagan et al., 2008; Bosman et al., 2009; Herrington et al., 2009). The strength

of these responses can be variable, and even absent, in some cases. We next

describe the latencies of these responses before returning to the potential reasons

for such variability, and complete absence in some cases.

Visual reafferent responses after microsaccades are earlier than visual evoked

responses after stimulus onset

Since a new fixation begins after the end of the prior eye movement, visual reafferent

responses are frequently aligned in analyses on movement end (that is, new fixation

onset; e.g. Kagan et al., 2008). With such a convention, visual reafferent responses

after microsaccades occurred in our data set significantly earlier than visual evoked

responses after stimulus onset. This is already evident in the example neuron of Fig.

2. As can be seen, the time to peak firing rate after stimulus onset in this neuron was

44 ms, 52 ms, and 66 ms for the low, intermediate, and high spatial frequencies,

respectively. On the other hand, after microsaccade end, these times were 27 ms, 28

ms, and 40 ms instead. Across the population, the visual reafferent response (when it

did occur) had a latency (to peak firing rate) of 33 ms +/- 2.5 ms s.e.m. from

microsaccade end for the lowest spatial frequency, 40 ms +/- 3 ms s.e.m. for the

intermediate spatial frequency, and 46 ms +/- 6 ms s.e.m. for the highest spatial

frequency (versus 61 ms +/- 2 ms s.e.m., 62 ms +/- 3 ms s.e.m., and 68 ms +/- 6 ms

s.e.m. for the stimulus driven visual evoked response, respectively). Thus, the post-

microsaccadic visual reafferent respons was ~24 ms earlier than the visual evoked

response (Fig. 3d-f). These results are consistent with other studies for large

saccades in other areas (e.g. Price et al., 2005; Ibbotson et al., 2008; Kagan et al.,

2008; Rajkai et al., 2008; White et al., 2017a).

Note that the latencies in Fig. 2 and Fig. 3d-f after microsaccades might appear too

fast given visual conduction delays in the early visual system. However, the stimulus

jittering that happens on the retina because of a microsaccade (or saccade) actually

starts with the movement onset itself. That is, neurons do not merely respond to the

stimulus at the start of the new fixation, but they respond to the entire spatiotemporal

pattern of stimulus movement caused by the eye movement. Therefore, it is expected

that the visual reafferent response is affected by the entire event of a microsaccade,

including the initial motion component, and not just reacting to the new stable image

of the grating after the end of the eye movement. Consistent with this, when we

computed the timing of the visual reafferent response now relative to microsaccade

onset rather than microsaccade end, we found that it had similar or longer latency

than the original stimulus-driven visual evoked response (Fig. 3g-i). This delaying

(Fig. 3g-i) and weakening (Fig. 3a-c) of the visual reafferent response (when

compared to the stimulus-driven visual evoked response) is expected due to the

motion blur associated with the microsaccadic movements themselves.

Visual reafferent responses after microsaccades do not always reflect the visual

tuning of SC neurons assessed with the evoked response

The visual consequence of any given eye movement should depend on the size and

direction of the movement relative to the stimulus pattern (that is, how the spatial

luminance modulation in the stimulus gets translated over a given RF). This means

that the visual reafferent response might not always be a replica (albeit weakened;

Fig. 3) of the visual evoked response. We investigated this issue, which is under-

represented in the neurophysiology literature of post-saccadic reafference, in our

recorded SC neurons.

We first asked whether response strength and response latency obeyed known

observations about SC visual responses to Gabor gratings (Chen et al., 2018). In the

visual evoked response, we found that most of our neurons (47%) preferred the

lowest spatial frequency in terms of firing rate magnitude, consistent with (Chen et

al., 2018). To demonstrate this, we normalized each neuron's visual evoked

response to the peak firing rate observed after the onset of the grating with the

lowest spatial frequency (0.56 cpd). We then divided all firing rate measurements

across trials, and across spatial frequencies, by this measurement. We also used the

same normalization factor for the visual reafferent response. We then pooled all

neurons' normalized firing rates in Fig. 4a for the visual evoked response and in Fig.

4d for the visual reafferent response (for simplicity, we only show the reafferent

response aligned on microsaccade end, rather than also aligned on microsaccade

onset, in this figure). In terms of response magnitude, tuning was generally

maintained in the visual reafferent response. That is, the visual reafferent response

was strongest for the lowest spatial frequency and weakest for the highest spatial

frequency. This idea is also shown in Fig. 4b, e, in which individual neuron

distributions are presented under the different stimulus presentation conditions of

low, intermediate, and high spatial frequencies; there was a significant main effect of

spatial frequency (one-way ANOVA) in the visual evoked response ($F_{(2,162)}=10.89$,

$p<0.0001$) and also the visual reafferent response ($F_{(2,162)}=7.89$, $p=0.0005$).

Figure 4 Comparison of response tuning to different spatial frequencies in either the visual evoked response or the visual reafferent response. (a) Firing rates of all neurons after stimulus onset. Each neuron's firing rate curve was first normalized by the firing rate curve obtained for the lowest spatial frequency. Then, all neurons were combined (n=55 neurons). Error bars denote s.e.m. across neurons. As can be seen, the population preferred the lowest spatial frequency in terms of response magnitude and also response latency (shorter for lower spatial frequencies). **(b)** Individual neuron measurements of peak firing rate (unnormalized) after stimulus onset. The measurements (from the same neurons as in **a**) are separated based on stimulus spatial frequency. **(c)** For the same neurons, we now calculated time to peak firing rate, as a proxy for visual response latency. The visual evoked response was earliest for the lowest spatial frequency (Chen et al., 2018). **(d)** For the same neurons and the same normalization factor as in **a**, we plotted the visual reafferent response. Error bars denote s.e.m. across neurons. The visual reafferent response was weak, but still preferred the lowest spatial frequency. **(e, f)** Same as **b**, **c** but now for the visual reafferent response.

Interestingly, the rank ordering of response latencies with increasing spatial

frequency, which was observed earlier (Chen et al., 2018), was also present in the

visual reafferent response. Specifically, in Fig. 4c, we plotted the time to peak firing

rate in the visual evoked response of neurons, and confirmed the earlier observations

(Chen et al., 2018). In Fig. 4f, we plotted the time to peak firing rate of the visual

reafferent response after microsaccade end. The range of measurements was

naturally broader and more variable than for the visual evoked response (Fig. 4c), but

this is expected given the variable durations and speeds of individual microsaccades

(introducing variability in both the amount and duration of the motion blur caused by

eye movements before a new "fixation" was eventually established). Nonetheless,

there was a clear trend for increasing response latency with increasing spatial

frequency, as in the visual evoked response (also seen in the example neuron of Fig.

2): there was a significant main effect of spatial frequency (one-way ANOVA) in the

visual evoked response ($F_{(2,162)}=30.27$, $p<0.0001$) and also the visual reafferent

response ($F_{(2,162)}=4.54$, $p=0.0121$).

These results indicate that even though microsaccades are very small (Fig. 1c) and

relatively slow eye movement events (significantly less than 70 deg/s peak velocity),

they still refresh SC activity to represent a stimulus that was moved over the neuronal

RF's by them. Such refreshing is mild (Fig. 3), but still present (Fig. 4), and it is

generally faithful to the visual evoked tuning of neurons.

Having said the above, at the individual neuron level, we did observe cases of the

tuning of a given neuron apparently changing in the visual reafferent response when

compared to the visual evoked response. For example, we grouped the neurons that

preferred the lowest (26 neurons) or intermediate (16 neurons) spatial frequency. We

then repeated the same analyses of Fig. 4a, d but this time normalizing firing rates to

the peak firing rate of the preferred spatial frequency (rather than to the lowest spatial

frequency as in Fig. 4a, d). Neurons preferring the lowest spatial frequency in the

visual evoked response still primarily preferred the lowest spatial frequency in the

visual reafferent response (Fig. 5a, b). Specifically, the left panel of Fig. 5a shows an

analysis similar to that of Fig. 4a. In the right panel of the same figure, we plotted the

tuning curves of individual neurons for the three different spatial frequencies. All

neurons in this group preferred the lowest spatial frequency in the stimulus-driven

visual evoked response (as per the design of the analysis). Now, for the visual

reafferent response after microsaccades (Fig. 5b), the tuning curves of the same

neurons largely still preferred the lowest spatial frequency stimulus (with a milder

reafferent response strength; Fig. 3). However, some neurons actually responded in

a stronger fashion for the intermediate spatial frequency (2.22 cpd) rather than the

lowest spatial frequency (5/26; 19%). This effect was even more striking when we

repeated the same tuning curve analysis for neurons preferring the intermediate

spatial frequency in the visual evoked response; these neurons often preferred the

lowest spatial frequency in the visual reafferent response (Fig. 5c, d); that is, the

visual reafferent response was strongest for the lowest rather than the intermediate

spatial frequency (11/16 neurons; 69%). A similar effect occurred for the neurons

preferring the highest spatial frequency, although there were too few neurons in this

condition (especially with a significant visual reafferent response); we therefore do

not show the results for the neurons preferring the highest spatial frequency.

Figure 5 Differences in tuning between the visual evoked and visual reafferent responses of SC neurons. (a) The left panel shows normalized firing rate (similar to Fig. 4a) for all neurons preferring the lowest spatial frequency. Firing rates were normalized by the visual evoked response to the lowest spatial frequency (the preferred stimulus). Error bars denote s.e.m. across neurons. The right panel shows the individual neuron tuning curves. **(b)** For the same neurons, we plotted the visual reafferent response aligned on microsaccade end. The same normalization factor as in **a** was used. The right panel shows the tuning curves of the same neurons during the visual reafferent response. Most (21/26) neurons still preferred the lowest spatial frequency, but some preferred the intermediate spatial frequency (5/26; the differently colored tuning curves). **(c, d)** Same as **a**, **b** but now for the neurons preferring the intermediate spatial frequency. Firing rate curves were now normalized by the visual evoked response after presentation of the intermediate spatial frequency (the preferred stimulus). The tuning curves in **d** were different from those in **c**. Only 3/16 neurons still preferred the intermediate spatial frequency in the visual reafferent response; most (11/16) preferred the lowest spatial frequency instead.

These results suggest that when the visual reafferent response is weak (that is, for the intermediate and highest spatial frequencies), then it may be most sensitive to interactions between the microsaccadic eye movement's vector and the stimulus properties, resulting in an apparent change in visual feature tuning of the neuron during the visual reafferent response. We directly demonstrate the evidence supporting this hypothesis in the next section; in other words, there is visual coding of neural patterns caused by the specific microsaccadic eye movement vectors.

Visual reafferent responses after microsaccades depend on the temporal image

luminance modulations over RF's caused by the shifting eye positions

To clarify the potential importance of luminance modulations caused by a given eye

movement for the results of Fig. 5 above, we further investigated the relationships

between microsaccade properties and stimulus properties. Specifically, in the

experiments described so far, the grating was vertical (Materials and Methods). This

means that a purely horizontal microsaccade would cause luminance modulation

over the RF as the eye movement displaces the retinotopic image of the grating

horizontally (orthogonal to the grating), whereas a purely vertical microsaccade

would not cause such luminance modulation (Fig. 6a). Moreover, depending on the

spatial frequency of the grating, the size of the shift caused by a purely horizontal

microsaccade should influence the absolute amplitude luminance modulation

experienced at a given retinotopic RF location (Fig. 6b). For example, for a horizontal

microsaccade and a vertical grating of 0.56 cpd, the largest luminance modulation

over an RF expected by the eye movement would occur for a large microsaccade of

approximately 0.9 deg amplitude (54 min arc) (Fig. 6b). Therefore, both

microsaccade size and microsaccade direction should influence the strength of the

visual reafferent response in SC neurons, resulting in an apparent change in tuning

in some cases, as we saw in Fig. 5. We tested for this idea by isolating

microsaccades of specific sizes and directions.

703
Figure 6 Potential impacts of microsaccade sizes and directions on properties of the visual
reafferent response of SC neurons. (a) A movement orthogonal to grating orientation (horizontal in
this case) is expected to cause luminance modulation in time as a result of eye-movement induced shift
in retinotpic stimulus position; a parallel eye movement is expected to cause no modulation. Therefore,
movement direction relative to stimulus orientation should modulate the visual reafferent response. **(b)**
In addition, the amplitude of a movement in relation to the spatial frequency of the stimulus should cause
predictable changes in luminance modulation. Here, assuming a purely orthogonal eye movement, the
absolute value of amplitude modulation expected for a given spatial frequency is plotted as a function
of eye movement amplitude. For example, for the lowest spatial frequency, a very large microsaccade
is needed to cause a strong luminance modulation over a neuron's RF (the dashed vertical line denotes
9 min arc). **(c, d)** Microsaccade amplitude probabilities like in Fig. 1c, but now plotted separately for the
horizontal and vertical components of microsaccades. **(e, f)** Amplitude probabilities as in **c**, **d** but now
for predominantly orthogonal (horizontal) microsaccades. These were movements with directions within
+/- 22.5 deg from horizontal. The parallel component of these movements was, expectedly, much
reduced. Predominantly parallel microsaccades had similar histograms (not shown), except that the
small-amplitude distributions were now in the orthogonal, rather than parallel, component of the
movements.

To do so, we first inspected the constraints on expected luminance modulations that

were imposed on us by our observed microsaccade amplitudes (Fig. 1c). Both

monkeys had microsaccades of different directions, having different sizes of

horizontal or vertical microsaccade components (Fig. 6c, d). This allowed us to

develop specific expectations (Fig. 6b) on the impacts of, say, microsaccade size or

microsaccade direction on the visual reafferent response to a given spatial frequency

(e.g. 0.56 cpd). We first investigated the effects of microsaccade size, and we then

moved on to next investigate the impacts of microsaccade orientation relative to

grating orientation. For the latter, we picked microsaccades with angles that were

within +/- 22.5 deg from either horizontal or vertical, to isolate movements that were

either primarily orthogonal or primarily parallel to the vertical grating, respectively. For

example, in Fig. 6e, f, we measured the distributions of predominantly orthogonal

microsaccades (very small "parallel" amplitude component in the figure) in each

monkey, which we could use in conjunction with Fig. 6b to predict potential effect

sizes for different stimulus spatial frequencies.

In terms of microsaccade size, we picked all horizontal microsaccades (within +/-

22.5 deg from horizontal in direction), and we binned their amplitudes as being either

smaller or larger than 0.15 deg (9 min arc; this binning was, once again, constrained

by our observed microsaccade amplitude distributions in Figs. 1c, 6c-f). Given the

predicted luminance modulation caused by a given movement size (Fig. 6b), one

should expect that the larger microsaccades would be associated with larger visual

reafferent responses, but only for the lowest and intermediate spatial frequencies.

Specifically, between the amplitudes of 9 min arc and ~20-22 min arc (the upper limit

of the great majority of our microsaccades; Fig. 1c), the expected luminance

modulations caused by the eye movements are higher for both low and intermediate

spatial frequencies than the expected luminance modulations caused by smaller

microsaccades with amplitudes <9 min arc (Fig. 6b). For the highest spatial

frequency, the same ranges of microsaccade amplitudes that were available to us

(Figs. 1c, 6c-f) would not be expected to reveal any systematic effect. This was

indeed the case. In Fig. 7a, d, g, we plotted the visual reafferent response for "small"

(<9 min arc) and "large" (>9 min arc) horizontal microsaccades, across the different

stimulus spatial frequencies. The violin plots (Fig. 7b, e, h) show raw measurements

of the visual reafferent response (peak firing rate of a neuron in the interval 0-100 ms

from microsaccade end), and the firing rate curves (Fig. 7a, d, g) show mean curves

across the population of neurons (after each neuron's firing rate curve was first

normalized by the peak visual response after stimulus onset for the lowest spatial

frequency; as in Fig. 4 above). Error bars denote s.e.m. across neurons. As can be

seen, the visual reafferent response was stronger for larger microsaccades (within

the range of microsaccade amplitudes that we were able to analyze), but not for the

highest spatial frequency. We confirmed this statistically in Fig. 7c, f, i by plotting the

measurements in Fig. 7b, e, h as paired observations (reafferent response after large

microsaccades versus reafferent response after small microsaccades). The clearest

effects were present for the low and intermediate spatial frequencies but not for the

highest spatial frequency ($p < 0.0001$, $p < 0.0001$, and $p = 0.61$, respectively; t-test).

Therefore, consistent with the predictions of Fig. 6b, microsaccade size had a clear

influence on post-microsaccadic visual reafferent responses in the SC, and in a

manner that demonstrated dependence on temporal luminance modulations over the

RF caused by the eye movements.

Figure 7 Dependence of the visual reafferent response on microsaccade size in relation to spatial pattern layout. (a) We plotted the average normalized visual reafferent response of neurons (normalized as in Fig. 4) when the RF stimulus had a spatial frequency of 0.56 cpd. The two curves indicate the reafferent response for microsaccades >9 min arc in amplitude or <9 min arc in amplitude, respectively. Error bars denote s.e.m. (across averaged neurons). **(b)** Individual neuron measurements for the same data as in **a**. The distribution of activity for larger microsaccades was higher than that for smaller microsaccades (p<0.0001; t-test). **(c)** The same data but now plotted as paired measurements, showing enhanced an visual reafferent reafferent response for the larger microsaccades. **(d-f)** Similar results for the intermediate spatial frequency (p<0.0001; t-test). **(g-i)** For the highest spatial frequency, there was no clear difference in the strength of the visual reafferent response as a function of larger or smaller microsaccades (p=0.61; t-test).

In terms of microsaccade directions, we expected that movements orthogonal to the

gratings (that is, predominantly horizontal movements) should have the strongest

visual reafferent response since they cause the strongest luminance modulations of

the gratings over the recorded RF (Fig. 6a). We therefore separated microsaccades

according to whether they were predominantly horizontal or predominantly vertical

(Fig. 8). In each case, and as stated above, we defined predominantly horizontal or

predominantly vertical as all movements with directions within +/- 22.5 deg from the

designated direction. The strongest visual reafferent response occurred for the

horizontal microsaccades (labeled orthogonal in the figure since the gratings used

were vertical), but (unlike with microsaccade size effects in Fig. 7) this effect was

now strongest for the intermediate and high spatial frequencies and not for the lowest

spatial frequency. Specifically, in both the average firing rate curves (Fig. 8a, d, f) as

well as the individual neuron distributions (Fig. 8b, e, h), we could easily observe

stronger visual reafferent responses for orthogonal compared to parallel

microsaccades in the intermediate and high spatial frequencies; also see the paired

measurements of the individual neuron distributions shown in Fig. 8c, f, i (p=0.0114,

0.0002 for the intermediate and high spatial frequencies, respectively; t-test).

However, there was no difference in the visual reafferent response when the

microsaccades jittered the image of a low spatial frequency grating either horizontally

or vertically (orthogonally or in a parallel fashion; p=0.63; t-test). Given the overall

sizes of our microsaccades (Figs. 1c, 6c-f), this was expected because the lowest

spatial frequency would have required substantially larger eye movements to cause

strong luminance modulations (Fig. 6b).

816
Figure 8 Dependence of the visual reafferent response on microsaccade direction relative to the
orientation of luminance modulation patterns in a visual stimulus. This figure is formatted
identically to Fig. 7, except that the comparison is now between microsaccades that were predominantly
orthogonal to the grating in the RF or microsaccades that were predominantly parallel. Orthogonal
microsaccades were associated with a stronger visual reafferent response than parallel microsaccades,
except for the lowest spatial frequency (for which our observed microsaccade sizes were too small to
result in appreciable differences in luminance modulations; Fig. 6b). P-values for statistical comparisons
for each spatial frequency are reported in the text.

Therefore, to summarize our results so far, we found that a significant component of

the effects of microsaccades on vision is to modulate how SC visual neural activity

represents stimuli that are already stably present in the environment (Figs. 1-4). The

interaction between the visual transient caused by the eye movement (that is, the

retinotopic motion caused by the movement) and the movement vector itself relative

to the underlying stimulus pattern will vary how SC neural activity is modulated (Figs.

6-8), and therefore how image representations are ultimately decoded downstream

depending on the needed behavioral readout of SC activity. This can give the

potential appearance of changes in visual tuning of SC neurons during the visual

reafferent response interval (e.g. Fig. 5).

We finally aimed to further support the above interpretations by analyzing LFP

signals around our recording electrodes. We previously saw that LFP measurements

in the SC provide a highly sensitive measure of population activity (Hafed and Chen,

2016; Chen and Hafed, 2017, 2018; Chen et al., 2018). We therefore analyzed visual

evoked LFP deflections across our population, as well as microsaccade-aligned LFP

deflections (Materials and Methods). We first replicated our recent observations that

the visual evoked LFP response preferred low spatial frequencies (Chen et al.,

2018). This replication is shown in Fig. 9a (in a similar format to that used in Chen et

al., 2018); stronger LFP deflections (more negative) occurred for the lowest spatial

frequencies. We then aligned the LFP signals on microsaccade end (Fig. 9b) or

microsaccade onset (Fig. 9c), similar to how we aligned firing rates earlier (e.g. Fig.

2). We found that when a microsaccade happened with a stable visual stimulus near

the aggregate population RF location around the recorded sites, there was a clear

visual reafferent LFP response. Interestingly, even though the reafferent response

was weaker than the visual evoked response (compare the y-axis ranges in Fig. 9a

and Fig. 9b, c), there was clear feature tuning of the reafferent response: the visual

reafferent LFP response was strongest (most negative) for the lowest spatial

frequency, and progressively weakened with increasing spatial frequency (Fig. 9b, c),

just like the visual evoked LFP response (Fig. 9a). Moreover, we found that the post-

microsaccadic LFP response was a genuine reafferent response because the LFP

signal aligned on microsaccades when there was no visual stimulus on the display at

all had a much weaker deflection (inset in Fig. 9c; also see Chen and Hafed, 2017).

Therefore, our LFP analyses supported the notion of visual feature tuning of

reafferent neural responses in the SC after microsaccades.

864
Figure 9 A visual reafferent response in LFP signals in the primate SC. (a) We plotted the average
stimulus-driven LFP deflection across all of our recorded SC sites (error bars denote s.e.m. across
sites). Each colored curve shows the LFP response after the onset of a given spatial frequency (color
coded as in the legend in **b**) (n=55 sessions). The LFP signal reflected a general population preference
for low spatial frequencies in the visual evoked response (Chen et al., 2018). **(b)** For the same recording
sites, we aligned the LFP signal to microsaccade end when a grating of a given spatial frequency
(different colors) was in the aggregate RF location represented by the recorded SC site for a given
session. Consistent with the firing rate data (e.g. Figs. 2-4), there was a clear visual reafferent LFP
response, which also reflected the feature tuning of the visual evoked LFP response in **a**. That is, there
was a stronger response (stronger negativity) after microsaccades for low spatial frequencies. Note that
the visual reafferent LFP response, despite showing similar feature tuning to **a**, was weaker in strength
(compare y-axes in **a**, **b**). **(c)** Same data as in **b**, but now aligned on microsaccade onset. Similar
observations were made. The inset shows that when microsaccades occurred in the absence of a visual
stimulus on the display, the microsaccade-aligned LFP response was much weaker (less negative) than
when a stimulus was present. This confirms that the results in **b**, **c** were genuine reflections of SC
reafferent responses. The different colors in the inset denote that stimulus-absent microsaccades were
from trials in which a stimulus of a given spatial frequency (color coded) would later appear.
882

More importantly, when considering the impacts of microsaccade size as a function

of spatial frequency, exactly like we did in our firing rate analyses (Fig. 7), we found

consistent results in the visual reafferent LFP response (Fig. 10a-c): there was a

stronger reafferent LFP response for larger rather than smaller microsaccades (that

were orthogonal to the gratings), but only when low and intermediate spatial

frequencies were being jittered on the retina by the eye movements. Microsaccade

size (for our observed range: Figs. 1c, 6c-f) did not matter when high spatial

frequencies were stably present in the environment. As stated above, these results

are consistent with expectations of temporal luminance modulations caused by

microsaccades, for the given range of movement amplitudes that was present in our

data (Fig. 6a, b).

899
Figure 10 Dependence of the visual reafferent LFP response on microsaccade size and direction
relative to spatial pattern layout. (a-c) Similar analyses to those in Fig. 7, but this time on
microsaccade-aligned LFP signals. For each spatial frequency, we compared the visual reafferent LFP
response when the microsaccade that jittered the displayed stimulus was >9 min arc in amplitude or <9
min arc in amplitude. There was a stronger visual reafferent LFP response for the larger microsaccades,
except for the highest spatial frequency (consistent with Figs. 6b, 7). Error bars denote s.e.m. across
sessions. **(d-f)** Similar analyses to those in Fig. 8, but this time on microsaccade-aligned LFP signals.
The comparison is now between microsaccades orthogonal or parallel to the displayed stimulus, as
opposed to larger or smaller as in **a-c**. Consistent with Figs. 6b, 8, there was a clearly stronger visual
reafferent LFP response for the intermediate and high spatial frequencies, but not for the low spatial
frequency.

Similarly, when we separated microsaccades based on their directions (orthogonal

and parallel) relative to the gratings, the same predictions (Fig. 6a, b) suggested

stronger reafferent responses for orthogonal rather than parallel gratings, but only for

intermediate and high spatial frequencies. This was, again, clearly the case in our

visual reafferent LFP responses (Fig. 10d-f), just like in Fig. 8 for spiking activity.

Therefore, both our firing rate and LFP analyses demonstrated that the SC

continuously updates its representation of stably present stimuli in the visual

environment, and particularly with periodic post-microsaccadic reafferent responses

that critically depend on interactions between the movement vector (dictating

temporal modulations of luminance over an RF), on the one hand, and the spatial

layout of the stimuli, on the other.

Discussion

We investigated the properties of the visual reafferent response in SC neurons after microsaccades jittered stable stimuli inside the neurons' RF's. We found that the SC exhibits a visual reafferent response that is clearly visible in firing rates and LFP signals. The visual reafferent response is significantly weaker than the stimulus-driven visual evoked response, and sometimes also absent. Moreover, the visual reafferent response exhibits feature tuning properties generally similar to those of the visual evoked response. However, at the individual neuron level, some neurons may appear to alter their feature tuning preferences in the visual reafferent response when compared to the visual evoked response. Investigating the detailed interactions between microsaccade sizes/directions and image patterns led us to conclude that such alteration of feature tuning likely reflects the impacts of the temporal modulations of luminance caused over an RF by a microsaccade. Such modulations depend not only on the spatial frequency of the stimulus itself, but also on the eye movement vector (size/direction).

Our results indicate that the SC provides a continuous representation of a visual scene, with periodic "updating" through visual reafferent responses even with tiny microsaccades. In recent work, we investigated foveal visual neurons in the SC, and we found that these neurons reflect how slow fixational drift eye movements can bring the fixated spot in and out of the neurons' RF's (Chen et al., 2019). This causes variability in the neurons' firing activity during fixation. The current results go beyond such previous work by showing that even parafoveal and peripheral SC neurons continue to represent visual scenes as they are jittered on the retina by microsaccades. This is interesting because it demonstrates that the visual properties of the SC that seem to be very relevant for orienting behavior, such as RF size and

visual sensitivity (Hafed and Chen, 2016; Chen et al., 2018; Hafed, 2018), also

extend to visual representations when stimuli are stable in the environment and

moved retinotopically by eye movements. It would be interesting in future studies to

explore how the SC's representation of stable environments can influence

subsequent visuo-motor behavior, exactly through its visual reafferent responses. For

example, do enhanced reafferent responses after microsaccades facilitate

subsequent saccades in the same direction as the extra-foveal bursting neurons?

One interesting consequence of results like those shown in Fig. 5, with apparent

changes in feature tuning of some neurons after microsaccades, is that there may be

inherent ambiguities in individual neurons' response strengths as a consequence of

eye movements. While it may be conceivable that individual neurons can change

their feature tuning curves across eye movements, similar in principle to how spatial

RF's can be significantly altered around the time of saccades (Duhamel et al., 1992;

Walker et al., 1995; Zirnsak et al., 2014), we think that a more likely source of the

alterations that we observed in our experiments (Fig. 5) was the expected temporal

luminance modulation over the recorded RF caused by eye movements (Figs. 6-8,

10). Specifically, we found that there were predictable dependencies of the visual

reafferent response strength on both the stimulus spatial profiles as well as the eye

movement vectors. This functionally means that eye movements contribute to visual

coding by enhancing some feature properties (Rucci et al., 2007; Rucci and Victor,

2015; Rucci et al., 2018), and in our case, this happened even for microsaccades.

For example, edges in an image orthogonal to the microsaccades would be

enhanced significantly more than edges parallel to the microsaccades (Fig. 8, 10d-f).

More interestingly, such visual coding also happens within the SC, a structure that

has been primarily thought of for many years as being more of a motor control

structure rather than a visual structure.

Related to the above, another interesting consequence of our results is that the post-

microsaccadic SC visual reafferent response may impose on peripheral SC neurons

a temporal tag of updated visual representations that is somewhat independent of

position coding (because of the small sizes of the eye movements). Specifically, in

natural viewing, very small saccades may be frequently required to scan features of

far objects (e.g. features of a mountain peak viewed with the naked eye on a hiking

adventure, or features of the far road ahead when driving at high speed on a

highway). Because the RF's of peripheral SC neurons, particularly in the lower visual

field, are relatively big (Hafed and Chen, 2016), such tiny saccades (Fig. 1c) are too

small to move the images being jittered on the retina by the eye movements outside

or away from the neurons' RF's. Nonetheless, according to our results, the temporal

modulations caused by the small retinal jitter would still cause a reafferent response

in the peripheral neurons. Therefore, peripheral edges or other visual features may

be enhanced periodically by small saccades in natural viewing, even when these

small saccades may be used for high acuity foveal tasks (like inspecting far objects).

This periodic "updating" maintains the peripheral image representations that may be

needed to re-orient gaze with larger eye movements to specific peripheral features of

interest. It also means that the SC visual reafferent response that we studied after

microsaccades here is not restricted to being a highly constrained laboratory

phenomenon with enforced fixation like in our tests. Rather, it can still happen in

many natural scenarios in which very small saccades would be frequently employed

anyway.

1002

Combined with results from other visual brain areas (Leopold and Logothetis, 1998;

Martinez-Conde et al., 2000; Snodderly et al., 2001; Martinez-Conde et al., 2002;

Kagan et al., 2008; Bosman et al., 2009; Herrington et al., 2009), the above temporal

tagging idea that we alluded to (i.e. the periodic updating of image processed

representations by microsaccadic or larger saccadic eye movements) may in reality

be an almost global phenomenon across the whole brain. Indeed, several brain areas

that may be concerned with other sensory (e.g. auditory) or motor (e.g. arm

movement) modalities are also sensitive to visual stimulation. As a result, and

because vision comes through a mobile eye, these same brain areas may be

sensitive to eye-movement-induced visual stimulation due to retinal image

movements (i.e. potential visual reafferent responses). This means that visual

processing is almost never immune from the impacts of individual saccadic or

microsaccadic eye movements. In fact, eye movements cause phase-resetting of

visual and oculomotor systems (Gaarder et al., 1966; Hafed and Ignashchenkova,

2013; Bellet et al., 2017), and there are very long-lasting impacts of such resetting on

perception (Bellet et al., 2017). All of this leads to an urgent conclusion that vision, by

virtue of its continually mobile sensor, cannot and should not be studied in isolation of

orienting behaviors, in general, and eye movements, in particular.

1021

Finally, a component of eye movements that still occurs in the absence of saccades

and microsaccades is that of slow ocular drifts. Such fixational drifts are related to

microsaccades and saccades, because drift properties are clearly related to, and

potentially affected by, the occurrence of the latter more rapid eye movements (e.g.

(Chen and Hafed, 2013). Moreover, drifts continue to occur in between

microsaccades, even when strict gaze fixation is required. This means that the same

idea of temporal luminance modulations over an RF that we described in this study

would still be valid for SC neurons with slow ocular drifts. However, given the much

smaller sizes of drift eye movements (~ 1 min arc scale) and their much lower speeds

(< 0.5 deg/s or less) when compared to even microsaccades, the amount of

displacements and potential motion blur associated with drift eye movements would

be expected to be much smaller than even those associated with the very small

movements that we encountered in our experiments (Figs. 1c, 6). It remains to be

seen whether SC visual responses would be sensitive to such minute, almost

indiscernible, image modulations associated with slow ocular drift eye movements. In

our future experiments, this will be explicitly tested, and particularly with causal

manipulations of real-time retinal image stabilization (and careful eye movement

recording) in order to allow us to establish even more strong evidence that SC

neurons provide a continuous, real-time representation of our visual environment that

is reformatted by active oculomotor behavior during fixation.

1042

1043

1044

1045

References

Ahissar E, Arieli A, Fried M, Bonneh Y (2016) On the possible roles of
microsaccades and drifts in visual perception. Vision Res 118:25-30.

Basso MA, May PJ (2017) Circuits for Action and Cognition: A View from the
Superior Colliculus. Annu Rev Vis Sci.

Bellet J, Chen CY, Hafed ZM (2017) Sequential hemifield gating of alpha- and beta-
behavioral performance oscillations after microsaccades. J Neurophysiol
118:2789-2805.

Bellet ME, Bellet J, Nienborg H, Hafed ZM, Berens P (2019) Human-level saccade
detection performance using deep neural networks. J Neurophysiol 121:646-
661.

Bosman CA, Womelsdorf T, Desimone R, Fries P (2009) A microsaccadic rhythm
modulates gamma-band synchronization and behavior. J Neurosci 29:9471-
9480.

Chen CY, Hafed ZM (2013) Postmicrosaccadic enhancement of slow eye
movements. The Journal of neuroscience : the official journal of the Society for
Neuroscience 33:5375-5386.

Chen CY, Hafed ZM (2017) A neural locus for spatial-frequency specific saccadic
suppression in visual-motor neurons of the primate superior colliculus. J
Neurophysiol 117:1657-1673.

Chen CY, Hafed ZM (2018) Orientation and Contrast Tuning Properties and
Temporal Flicker Fusion Characteristics of Primate Superior Colliculus
Neurons. Front Neural Circuits 12:58.

Chen CY, Ignashchenkova A, Thier P, Hafed ZM (2015) Neuronal Response Gain
Enhancement prior to Microsaccades. Curr Biol 25:2065-2074.

Chen CY, Hoffmann KP, Distler C, Hafed ZM (2019) The Foveal Visual
Representation of the Primate Superior Colliculus. Curr Biol 29:2109-2119
e2107.

Chen CY, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency
sensitivity in macaque midbrain. Nat Commun 9:2852.

Duhamel JR, Colby CL, Goldberg ME (1992) The updating of the representation of
visual space in parietal cortex by intended eye movements. Science 255:90-
92.

Engbert R (2006) Microsaccades: A microcosm for research on oculomotor control,
attention, and visual perception. Prog Brain Res 154:177-192.

Fuchs AF, Robinson DA (1966) A method for measuring horizontal and vertical eye
movement chronically in the monkey. J Appl Physiol 21:1068-1070.

Gaarder K, Koresko R, Kropfl W (1966) The phasic relation of a component of alpha
rhythm to fixation saccadic eye movements. Electroencephalography and
clinical neurophysiology 21:544-551.

Gandhi NJ, Katnani HA (2011) Motor functions of the superior colliculus. Annual
review of neuroscience 34:205-231.

Goldberg ME, Wurtz RH (1972) Activity of superior colliculus in behaving monkey. I.
Visual receptive fields of single neurons. J Neurophysiol 35:542-559.

Hafed ZM (2011) Mechanisms for generating and compensating for the smallest
possible saccades. Eur J Neurosci 33:2101-2113.

Hafed ZM (2018) Superior Colliculus: A Vision for Orienting. Curr Biol 28:R1111-
R1113.

Hafed ZM, Krauzlis RJ (2010) Microsaccadic suppression of visual bursts in the
primate superior colliculus. J Neurosci 30:9542-9547.

Hafed ZM, Krauzlis RJ (2012) Similarity of superior colliculus involvement in
microsaccade and saccade generation. J Neurophysiol 107:1904-1916.

Hafed ZM, Ignashchenkova A (2013) On the dissociation between microsaccade rate
and direction after peripheral cues: microsaccadic inhibition revisited. J
Neurosci 33:16220-16235.

Hafed ZM, Chen CY (2016) Sharper, Stronger, Faster Upper Visual Field
Representation in Primate Superior Colliculus. Curr Biol 26:1647-1658.

Hafed ZM, Goffart L, Krauzlis RJ (2009) A neural mechanism for microsaccade
generation in the primate superior colliculus. Science 323:940-943.

Hafed ZM, Lovejoy LP, Krauzlis RJ (2013) Superior colliculus inactivation alters the
relationship between covert visual attention and microsaccades. Eur J
Neurosci 37:1169-1181.

Hafed ZM, Chen C-Y, Tian X (2015) Vision, perception, and attention through the
lens of microsaccades: mechanisms and implications. Frontiers in systems
neuroscience 9:167.

Hall NJ, Colby CL (2016) Express saccades and superior colliculus responses are
sensitive to short-wavelength cone contrast. Proc Natl Acad Sci U S A
113:6743-6748.

Herman JP, Krauzlis RJ (2017) Color-Change Detection Activity in the Primate
Superior Colliculus. eNeuro 4.

Herrington TM, Masse NY, Hachmeh KJ, Smith JE, Assad JA, Cook EP (2009) The
effect of microsaccades on the correlation between neural activity and
behavior in middle temporal, ventral intraparietal, and lateral intraparietal
areas. J Neurosci 29:5793-5805.

Ibbotson MR, Crowder NA, Cloherty SL, Price NS, Mustari MJ (2008) Saccadic
modulation of neural responses: possible roles in saccadic suppression,
enhancement, and time compression. J Neurosci 28:10952-10960.

Ikeda T, Boehnke SE, Marino RA, White BJ, Wang CA, Levy R, Munoz DP (2015)
Spatio-temporal response properties of local field potentials in the primate
superior colliculus. Eur J Neurosci 41:856-865.

Judge SJ, Richmond BJ, Chu FC (1980) Implantation of magnetic search coils for
measurement of eye position: an improved method. Vision Res 20:535-538.

Kagan I, Gur M, Snodderly DM (2008) Saccades and drifts differentially modulate
neuronal activity in V1: effects of retinal image motion, position, and
extraretinal influences. J Vis 8:19 11-25.

Krauzlis RJ, Goffart L, Hafed ZM (2017) Neuronal control of fixation and fixational
eye movements. Philos Trans R Soc Lond B Biol Sci 372.

Kuang X, Poletti M, Victor JD, Rucci M (2012) Temporal encoding of spatial
information during active visual fixation. Current biology : CB 22:510-514.

Leopold DA, Logothetis NK (1998) Microsaccades differentially modulate neural
activity in the striate and extrastriate visual cortex. Exp Brain Res 123:341-
345.

Martinez-Conde S, Macknik SL, Hubel DH (2000) Microsaccadic eye movements and
firing of single cells in the striate cortex of macaque monkeys. Nat Neurosci
3:251-258.

Martinez-Conde S, Macknik SL, Hubel DH (2002) The function of bursts of spikes
during visual fixation in the awake primate lateral geniculate nucleus and
primary visual cortex. Proc Natl Acad Sci U S A 99:13920-13925.

Martinez-Conde S, Macknik SL, Hubel DH (2004) The role of fixational eye
movements in visual perception. Nat Rev Neurosci 5:229-240.

Neupane S, Guitton D, Pack CC (2016) Two distinct types of remapping in primate
cortical area V4. Nat Commun 7:10402.

Poletti M, Rucci M (2016) A compact field guide to the study of microsaccades:
Challenges and functions. Vision Res 118:83-97.

Price NS, Ibbotson MR, Ono S, Mustari MJ (2005) Rapid processing of retinal slip
during saccades in macaque area MT. J Neurophysiol 94:235-246.

Rajkai C, Lakatos P, Chen CM, Pincze Z, Karmos G, Schroeder CE (2008) Transient
cortical excitation at the onset of visual fixation. Cerebral cortex 18:200-209.

Robinson DA (1972) Eye movements evoked by collicular stimulation in the alert
monkey. Vision Res 12:1795-1808.

Rolfs M (2009) Microsaccades: small steps on a long way. Vision Res 49:2415-2441.

Rucci M, Victor JD (2015) The unsteady eye: an information-processing stage, not a
bug. Trends Neurosci.

Rucci M, Ahissar E, Burr D (2018) Temporal Coding of Visual Space. Trends Cogn
Sci 22:883-895.

Rucci M, Iovin R, Poletti M, Santini F (2007) Miniature eye movements enhance fine
spatial detail. Nature 447:851-854.

Snodderly DM, Kagan I, Gur M (2001) Selective activation of visual cortex neurons
by fixational eye movements: implications for neural coding. Vis Neurosci
18:259-277.

Veale R, Hafed ZM, Yoshida M (2017) How is visual salience computed in the brain?
Insights from behaviour, neurobiology and modelling. Philos Trans R Soc
Lond B Biol Sci 372.

Walker MF, Fitzgibbon EJ, Goldberg ME (1995) Neurons in the monkey superior
colliculus predict the visual result of impending saccadic eye movements.
Journal of neurophysiology 73:1988-2003.

White BJ, Itti L, Munoz DP (2019) Superior colliculus encodes visual saliency during
smooth pursuit eye movements. Eur J Neurosci.

White BJ, Kan JY, Levy R, Itti L, Munoz DP (2017a) Superior colliculus encodes
visual saliency before the primary visual cortex. Proc Natl Acad Sci U S A
114:9451-9456.

White BJ, Berg DJ, Kan JY, Marino RA, Itti L, Munoz DP (2017b) Superior colliculus
neurons encode a visual saliency map during free viewing of natural dynamic
video. Nat Commun 8:14263.

Willeke KF, Tian X, Buonocore A, Bellet J, Ramirez-Cardenas A, Hafed ZM (2019)
Memory-guided microsaccades. Nat Commun 10:3710.

Zirnsak M, Steinmetz NA, Noudoost B, Xu KZ, Moore T (2014) Visual space is
compressed in prefrontal cortex before eye movements. Nature 507:504-507.
1184

1185

1186