The spatial structure of feedforward information in mouse

primary visual cortex

Jun Zhuang*[1], Rylan S. Larsen[1], Kevin T. Takasaki[1], Naveen D. Ouellette[1], Tanya L. Daigle[1],
Bosiljka Tasic[1], Jack Waters[1], Hongkui Zeng[1], R. Clay Reid[1]

1 Allen Institute for Brain Science, Seattle, WA, USA, 98109

* Correspondence: junz@alleninstitute.org

Abstract

Location-sensitive and motion-sensitive units are the two major functional types of feedforward projections from lateral genicular nucleus (LGN) to primary visual cortex (V1) in mouse. The distribution of these inputs in cortical depth remains under debate. By measuring the calcium activities of LGN axons in V1 of awake mice, we systematically mapped their functional and structural properties. Although both types distributed evenly across cortical depth, we found that they differ significantly across multiple modalities. Compared to the location-sensitive axons, which possessed confined spatial receptive fields, the motion-sensitive axons lacked spatial receptive fields, preferred lower temporal, higher spatial frequencies and had wider horizontal bouton spread. Furthermore, the motion-sensitive axons showed a strong depth-dependent motion direction bias while the location-sensitive axons showed a depth-independent OFF dominance. Overall, our results suggest a new model of receptive biases and laminar structure of thalamic inputs to V1.

Introduction

Parallel pathways along hierarchical processing stages are a prominent feature of early visual systems in carnivores (Stone et al., 1979; Lennie, 1980) and primates (Nassi and Callaway, 2009), where structured thalamocortical projections from dorsal lateral geniculate nucleus (LGN) to primary visual cortex (V1) serve as major inputs to cortical computations and shape cortical single-cell response properties (e.g., orientation selectivity, Hubel and Wiesel, 1962; Reid and Alonso, 1995) as well as the functional architecture at larger scale (Jin et al., 2011; Kremkow et al., 2016).

In recent years, the mouse visual system has become a popular model system to study visual physiology. When compared with higher mammals, the mouse LGN shows more diverse response properties. Particularly, besides cells with canonical ON and/or OFF concentric receptive fields (Grubb & Thompson, 2003; Denman et al., 2016, Suresh et al., 2016), a significant portion of the cells in mouse LGN are sensitive to motion (Marshel et al., 2012; Piscopo et al, 2013; Zhao et al., 2013; Durand et al., 2016) which may be inherited from its retinal inputs (Huberman et al, 2009; Cruz-Martin et al, 2014; Rompani et al., 2017; Dhande et al., 2015; Román Rosón et al., 2019). How this motion information is transmitted at the thalamocortical stage is less clear. It has been proposed that the motion information is conveyed by the direction-selective (DS) cells from LGN shell to the superficial layer of V1 while the location information is conveyed by cells in LGN core with concentric receptive fields to deeper layers (Cruz-Martin et al., 2014, Seabrook et al., 2017), forming parallel pathways similar to those in carnivores and primates. But the evidence for this model is scarce and

controversial, while one study found mouse layer 4 in V1 receives slightly less DS inputs than

layer 1 (Kondo et al., 2016), another study showed layer 4 receives the same amount of DS

inputs as superficial layers (Sun et al., 2016).

To investigate the spatial distributions of LGN inputs in V1, we used genetic and viral tools to

label mouse LGN neurons with calcium indicator and systematically measured the response

properties of their axon terminals in V1 at different depths. We show that the motion-sensitive

axons and location-sensitive axons represent two major functional types that differ significantly

in functional properties, spatial distribution, and axonal morphology, demonstrating that there

are at least two parallel thalamocortical pathways in mouse visual system.

Results

Selective labeling and functional imaging of LGN axons in V1

To image LGN axons in V1, we used the Vipr2-IRES2-Cre-neo transgenic mice. Crossing Vipr2-IRES2-Cre-neo mice with reporter line Ai14 drove tdTomato expression in the cells from LGN, ventral posterolateral nucleus, ventral posteromedial nucleus of thalamus, suprachiasmatic nucleus and central amygdala nucleus but not in the lateral posterior nucleus (LP) of thalamus and cortex (Figure 1A left, Allen Brain Atlas Transgenic Characterization Experiments: 576523754, 576524006, 577720920, 577721229). In V1, densely labeled axons were visible in layer 1 through layer 4 (Figure 1A right). Among the tdTomato positive nuclei, only LGN projects to V1, suggesting that the labeled axons in V1 were from LGN. To deliver calcium indicator for *in vivo* imaging, we injected a mixture of AAV viruses containing Cre-dependent GCaMP6s and Cre-independent mRuby3 directly into the LGN of Vipr2-IRES2-Cre-neo mice (Figure 1B). Three weeks after injection, the somatic expression of GCaMP was restricted in LGN while mRuby was expressed along the injection track through cortex and hippocampus (Figure 1C left). Importantly, the axons from these LGN cells were also labeled with GCaMP throughout superficial and middle layers in V1 (Figure 1C right).

We then performed two-photon calcium imaging of LGN axons 50-400 μm below the pia in awake mouse V1 (Figure 1D). The imaging was done in a columnar fashion: for a given cortical location, calcium signals from boutons were collected from 3 to 8 planes at different depths under pia. In total, 26 volumes (158 image planes from 6 mice) were imaged and calcium activity traces were extracted from 50171 regions of interest (ROIs) representing putative

boutons (Table S1). During each imaging session, we measured spatial receptive fields (RFs),

tuning for orientation/direction, spatial/temporal frequency (SF/TF), and spontaneous activity.

Many ROIs showed strong calcium activities (Figure 1E and Video S1), significant spatial RFs

(Figure 1F and S2A), and responses to gratings (Figure 1G-H). In some experiments, aberrations

were reduced with adaptive optics, thereby increasing fluorescence intensity and signal-to-

noise ratio. The response properties measured with and without adaptive optics were

substantially similar (Figure S1), so they were pooled together. In summary, our labeling

strategy enabled us to selectively label LGN neurons and characterize the functional properties

of their axons in V1.

LGN boutons can be grouped into axons based on highly correlated activities.

To map individual sources of LGN inputs to V1, it is necessary to distinguish boutons that are

from single axons from boutons that are from different axons. In each imaging plane, there

were subgroups of ROIs showing highly correlated activities indicating that they were likely

from same axons (Figure 1I-J, and Video S1). To identify these subgroups, we calculated an

event-based pairwise correlation between all pairs of boutons, followed by a hierarchical

clustering procedure (Methods, modified from Liang et al., 2018). Overall, the clustering

faithfully preserved the pairwise correlation distance (Figure 1M) and the correlation

coefficients from bouton pairs within and between clusters showed non-overlapping

distribution (Figure 1N). The boutons from single clusters showed highly correlated calcium

activities (Figure 1J) and near-identical response properties (Figure 1K-L), further confirming

their high likelihood of being from single axons. We found the clusters generated with full

calcium traces also represented the correlation structure of spontaneous activities (Figure S3),

thus all imaging planes were processed using the full calcium traces. We use "bouton" when

referring to individual ROIs, "axon" to indicate bouton cluster with high temporal correlations,

and "unit" to refer to either entity. For each axon, we used spatially summed calcium traces

from all its boutons to calculate its response properties (Method).

Figure 1. Labeling strategy, calcium imaging and bouton clustering for LGN axons in V1.

(A) tdTomato positive cells in thalamus (left) and axons in V1 (right) from a Vipr2-IRES-Cre-neo;Ai14 mouse.

(B) Labeling strategy for calcium imaging.

(C) GCaMP (green) and mRuby (red) expressions in a mouse prepared as in (B).

(D) Sketch of in vivo imaging setup.

(E) Mean projection of calcium activities from an example imaging plane at 400 μm below pia.

(F) The ON and OFF zscore receptive fields of the ROIs marked in (E).

(G) The averaged calcium responses of ROIs marked in (E) to different grating conditions. Only the peak TFs for each ROI are shown.

(H) The direction tuning curves of the ROIs marked in (E).

(I) Another example imaging plane at 350 μm below pia showing all active ROIs (grey) and 3 example clusters after axon grouping (colored).

(J) Calcium traces from the three marked axons. Colored traces: Traces from individual boutons. Gray traces: merged traces for the cluster. Vertical gray boxes: detected events. Numbers: mean pair-wise correlation coefficients.

(K) Superimposed orientation/direction tuning curves of individual boutons from each cluster.

(L) Superimposed receptive fields of individual boutons from each cluster.

(M) Clustered correlation coefficient matrix for the example imaging plane in (I). Colored boxes: clusters marked in (I).

(N) Normalized distribution of correlation coefficients for bouton pairs within clusters and between clusters. Solid lines: data from the imaging plane in (I). Dashed lines: data from all imaging planes.

Two populations of LGN axons with distinct functional and structural properties.

We found no depth-dependent differences in population average of direction/orientation

selectivity, preferred TF, and RF strength, while preferred SF was slightly higher at greater

depths (Figure S4). These results were consistent with a previous report (Sun et al, 2016), but

did not support the prediction of depth-dependent projection of motion-sensitive inputs

(Seabrook et al., 2017). We then focused our analysis on motion sensitivity and location

sensitivity, which are most likely to represent parallel pathways suggested by previous studies

(Piscopo et al., 2013, Suresh et al., 2016). We found that location-sensitive boutons (boutons

with significant RFs, RF boutons) were more likely to have low direction selectivity than the

boutons without RFs (Figure 2A) while the motion-selective boutons (boutons with gDSI > 0.5,

DS boutons) were more likely to have low RF strength than non-DS boutons (Figure 2B). Only a

small portion of boutons were both location-selectivite and motion-selectivite (RFDS boutons

5.32% of all boutons, Figure 2C-D), indicating DS boutons and RF boutons are largely separate

functional types. Consistent with Figure S4, we found no depth dependence in the number of

DS and RF boutons up to 400 μm from the pia surface (Figure 2E). Similar functional dichotomy

was also found in axons (Figure S5A-D).

Do DS and RF groups differ in other aspects? We first compared the SF and TF tuning between

DS and RF boutons. We found that while the distribution of preferred TF and SF from RF

boutons spread widely across the spectrum tested, DS boutons preferred only high SFs and low

TFs (Figure 2F-G, preferred SF: DS vs. RF, 0.102 ± 0.036 vs. 0.004 ± 0.063 cpd, U=188907, p<10^{-}

^{100}; preferred TF: DS vs. RF, 1.45 ± 0.59 vs. 2.56 ± 2.80 Hz, U=9421173, p<10^{-100}, Mann Whitney

U test). Similar results were found in axons (Figure S5E-F). Secondly, we compared the

normalized population axon masks between these two groups and found that the DS axons had

a greater bouton spread than the RF axons (Figure 2H-I, mean distance from bouton to axon

center, DS axons: 42.2 ± 27.3 µm, n=4093; RF axons: 39.7 ± 27.9 µm, n=9423; U=18121755,

p=1.22×10^{-8}, Mann Whitney U test). Consistently, the DS axons also showed higher bouton

counts, greater max bouton distances, and larger axon coverage area than RF axons (Figure S6).

These results indicate that the DS axons might have more extensive axon arbors than RF axons.

In summary, our data showed there are at least two separate groups of LGN inputs with one

carrying motion information (DS units), preferring high SF and low TF, and having wider axon

arbors, and the other carrying location information (RF units), preferring a wider wide range of

SFs and TFs, and having narrower axon arbors.

Figure 2. Motion-sensitive and location-sensitive axons are two major functional types of LGN inputs with distinct response properties and axon morphologies.

(A) Distribution of gDSI of boutons with (RF) or without (nRF) significant spatial receptive fields. Dashed line: DS threshold.

(B) Distribution of receptive field strength of DS and non-DS (nDS) boutons. Dashed line: RF threshold

(C) Counts of each functional type. RF group was broken into ON, OFF and ON-OFF groups. Numbers: the portion of DS units in each group.

(D) Venn diagram describing relations among the four non-overlapping groups.

(E) The ratio of each of these four groups across depths.

(F) and (G) Preferred SF and TF distributions, respectively.

(H) Stacked axon masks for both DS axons and RF axons.

(I) Normalized bouton distribution by distance to axon center.

Depth-dependent motion direction bias in DS group

We next examined the preferred direction distribution of DS boutons, which strongly biased

towards nasal and temporal directions (Figure S7) consistent with a previous report (Marshel et

al., 2012). Interestingly, in a given imaging plane, the DS population was often biased towards

either nasal or temporal directions (e.g., the temporal direction in the example plane, Figure

S2B). When broken into different depths, the direction bias showed a marked trend across

cortical depth: it gradually switched from nasal direction at 50 μm to temporal direction at 400

μm (Figure 3A). To quantify this depth-dependent bias, we defined a metric: nasal-temporal

index (NTI, Figure 3B inset). A negative NTI means more nasal-preferring units than temporal-

preferring units in a given population and a positive NTI vice versa. Consistent with the

observations above, the superficial (< 225 μm) boutons had a nasal bias (NTI = 0.245, Figure 3B

magenta), and the deep (> 225 μm) boutons had a temporal bias (NTI = -0.259, Figure 3B,

green). This trend was significant across all mice (Figure 3C, NTI: n=6, superficial vs. deep, 0.22 ±

0.09 vs. -0.11 ± 0.11, W=0, p=0.028) and all volumes (Figure 3D, NTI: n=17, superficial vs. deep,

0.25 ± 0.10 vs. -0.03 ± 0.12, W=17, p=0.005), and was insensitive to the DS threshold (Figure

3E). A similar depth-dependent bias was also found in DS axons (Figure S8).

Figure 3. The motion-sensitive LGN boutons showed depth-dependent bias of motion direction preference.

(A) Preferred direction distribution of DS boutons at each imaging depth. Numbers: NTI explained by the inset in (B). Histograms: distributions of gDSI at the given depth. Vertical scale: 100 boutons. Vertical dashed line: DS threshold.

(B) Preferred direction distribution of superficial half vs. deep half of all DS boutons.

(C) and (D), Superficial vs. deep NTI comparisons for each mouse and each volume, respectively.

(E) NTI across depth with different DS thresholds.

 OFF dominance in RF group

 Like previous reports in other species (Jin et al., 2008; Yeh et al., 2009; Jimenez et al., 2018), our

 RF boutons showed a strong OFF dominance. There were more OFF subfields than ON subfields

 per imaging plane (Figure 4A, all depths: n=121, ON vs. OFF, 54.5 ± 31.7 vs. 73.3 ± 52.1,

 W=1471, p=4.3x10^{-8}) and this difference was consistent across cortical depth (ON vs. OFF,

 superficial: n=73, 53.6 ± 31.5 vs. 72.8 ± 52.7, T=536, p=1.3x10^{-5}; deep: n=48, 55.8 ± 31.9 vs. 74.2

 ± 51.1, W=234, p=8.1x10^{-5}). Additionally, the OFF subfields were larger than ON subfields

 (Figure 4B, all depths, ON vs. OFF: n=121, 83.3 ± 2.0 vs. 102.1 ± 2.7 deg^2, W=1480, p=1.1x10^{-8})

 also consistent across depth (ON vs. OFF, superficial: n=73, 81.5 ± 2.4 vs. 102.1 ± 3.8 deg^2,

 T=532, p=6.8x10^{-6}; deep: n=48, 86.0 ± 3.4 vs. 102.0 ± 3.5 deg^2, W=240, p=3.6x10^{-4}). A similar

 OFF dominance was also found in RF axons (Figure S9).

 Another form of "OFF dominance" that was prominent in other mammalian species (Kremkow

 et al., 2016; Lee et al., 2016), reduced retinotopic scatter for OFF cells, also presented in our

 data. We generated population retinotopic maps for both ON and OFF RF units by pooling maps

 from all imaging planes after alignment and normalization (Method). The pooled maps showed

 a clear retinotopic gradient (Figure 4C) consistent with the established retinotopic map of

 mouse V1 (Garrett et al., 2014; Zhuang et al, 2017), with altitude increasing along the anterior-

 posterior axis and azimuth increasing along the lateral-medial axis. More importantly, the

 retinotopic gradient appeared less noisy for OFF subfields than ON subfields. To quantify this

 difference, we performed linear regression of altitude against anterior-posterior locations and

 azimuth against lateral-medial locations. As expected, the regression fits better for the OFF

retinotopy than for the ON retinotopy (Figure 4D). When the same analysis was performed on

individual volumes, the differences between ON and OFF retinotopy were significant (except

the R^2 for altitude regression) and consistent with the entire population (Figure 4E-F, n=19, ON

vs. OFF, standard error for altitude: 0.008 ± 0.001 vs. 0.006 ± 0.002 deg, W=37, p=0.02;

standard error for azimuth: 0.011 ± 0.002 vs. 0.009 ± 0.004 deg, W=31, p=0.01; R^2 for altitude:

0.032 ± 0.008 vs. 0.054 ± 0.029, W=51, p=0.076; R^2 for azimuth: 0.071 ± 0.012 vs. 0.105 ± 0.049,

W=35, p=0.016).

In summary, the OFF inputs in the RF group are more numerous, possess larger subfield, and

have less retinotopic scatter than the ON inputs.

Figure 4. Location-sensitive LGN boutons were OFF dominant.

(A) and (B) ON vs. OFF receptive subfield counts and areas of RF boutons, respectively.

(C) Normalized retinotopy along cortical surface for RF boutons.

(D) The linear regression of retinotopy location against the cortical distance.

(E) and (F) ON vs. OFF regression standard error of estimate and R^2 for each imaged volume, respectively.

Axon counts underlie functional specific bouton distribution

Finally, we asked what structure principle can give rise to the motion direction bias in DS

boutons and the OFF dominance in RF boutons. One possibility is that axons with different

functions might have different number of boutons ("bouton model", Figure S10A and D, left).

Alternatively, there could be different numbers of axons between different functional groups

while having similar bouton number per axon ("axon model", Figure S10A and D, right). The

bouton model predicts different degrees of direction bias and OFF dominance for boutons and

axons, while the axon model predicts similar degrees for boutons and axons. In favor of the

axon model, our data showed the axons have similar NTIs (Figure S8 and S10B) and OFF

dominance (Figure S9A and S10E) as the boutons. Our data also showed no difference in axon

morphology between subtypes within DS and RF groups (Figure S10C and F), further supporting

the axon model.

Discussion

This study systematically investigated the functional properties and the spatial distributions of

feedforward LGN axons in V1 with unprecedented labeling specificity. We showed the motion-

sensitive (DS) axons and location-sensitive (RF) axons represent two major functional types that

were largely distinct, with only ~5% of units falling between groups (Figure 2C and S5C).

Compared to the RF axons, the DS axons preferred low temporal and high spatial frequencies

and had a wider bouton spread. Furthermore, although neither type had depth-dependent

distribution in V1, the DS group showed a depth-dependent bias in motion direction preference

while the RF group showed a depth-independent OFF dominance. These findings support the

notion that, in the thalamocortical projections of the mouse visual system, there are

functionally and structurally distinct parallel pathways.

LGN specific labeling

We found that Vipr2-IRES-Cre-neo mice allow superior specificity in LGN labeling than wildtype

or Scnn1a-Cre mice used in previous studies (Kondo et al, 2016; Sun et al., 2016; Jaepel et al.,

2017), because they effectively prevent off-target labeling in cortex and LP (Figure 1A and 1C), a

region adjacent to LGN that projects substantially to V1 (Roth et al. 2016). This new labeling

strategy may account for the different results between this study and previous studies, for

instance if the Vipr2-IRES2-Cre-neo mice only label a subset of all LGN relay neurons. While we

cannot disprove this from our data, we note that the proportions of the two main functional

types in our dataset (RS axon: 53%, DS axon: 22%) are similar to what has been reported by

previous electrophysiological studies (RF cells: 52%, DS/OS cells: 11% Piscopo et al., 2013; DS

cells ~25%, Zhao et al. 2013), which argues against significant functional bias in our labeling

strategy.

Parallel pathways

Based on recent understanding of parallel visual pathways in the mouse (reviewed in Seabrook

et al., 2017), our RF units most likely correspond to X and Y units (Cleland et al., 1971;

Hoffmann et al., 1972), and DS units correspond to W units in cat LGN (Wilson & Stone, 1975;

Wilson et al., 1976). This is supported by our finding that DS axons had more extensive arbors

than RF axons (Figure 2H-I and S6), consistent with the morphological differences between W

and X/Y LGN axons in cats (Gilbert & Wiesel, 1979; Humphrey et al., 1985; Anderson et al.,

2009). However, we found the DS units preferred low TF and high SF (Figure 2F-G), notably

different from cat's W LGN cells which prefer lower SF than X- and Y-cells (Sur & Sherman,

1982). Furthermore, we did not find that DS LGN cells project predominantly to superficial

layers in V1 as predicted by the model (Seabrook et al., 2017), or mainly to layer 4 and 6 as has

been seen for DS cells in other species (Hei et al., 2014; Bereshpolova et al., 2019). Instead,

both RF and DS axons in our dataset distributed uniformly throughout all superficial depths

measured (Figure 2E and S4), as had been reported before (Sun et al., 2016, cf. Kondo et al.,

2016). Thus, the relationship between mouse LGN DS cells, W cells in the cat, and DS cells in

other species needs further investigation.

Our statement that DS units tend to not have significant spatial RF should be taken with

caution. The sparse-noise stimulus (Zhuang et al., 2013; Durand et la., 2016; de Vries et al.,

2020) we used to map the spatial RF is effective if the unit responds to brief small flashes, but

will perform poorly if the unit responds best to larger or more spatially structured stimuli, or

requires a longer temporal duration. Nonetheless, our conclusions about parallel pathways are

unlikely to be undermined because (1) a previous study with more structured mapping stimuli

showed similar function dichotomy (Piscopo et al, 2013); and (2) we found other significant

functional and structural differences between DS axons and RF axons (Figure 2F-I).

Motion direction bias of DS axons

Consistent with previous studies (Marshel et al., 2012; Piscopo et al., 2013; Zhao et al., 2013;

Durand et al., 2016; Sun et al., 2016; Kondo et al., 2016), we found that a significant population

of LGN units are direction-selective (22% for axons, 23% for boutons, Figure 2C, Table S1).

When pooling all our DS units together, the preferred directions strongly bias towards the nasal

and temporal directions (Figure S7) consistent with Marshel et al. 2012 but unlike other studies

(Sun et al., 2016: towards nasal direction; Kondo et al., 2016: towards four cardinal axes). These

differences may be due to the different labeling strategies, visual stimuli, and analysis methods.

The nasal/temporal direction biases have ethological relevance since they can be used to detect

self-generated motion during active behavior, as has been demonstrated at both retina (Sabbah

et al, 2017) and visual thalamus (Roth et al., 2016). Interestingly, our data show the direction

bias changes systematically along with cortical depth (Figure 3 and S8). The functional relevance

of this surprising organization is unknown and will be a topic for further investigation.

OFF dominance

It is well established in the early visual system of carnivores and primates that the OFF

responses are faster and stronger than ON responses (Jin et al, 2008, 2011; Yeh et al., 2009;

Mazade et la., 2019; Kremkow et al., 2018). The OFF responses are also stronger in mouse V1

cells (Jimenez et al., 2018). We extend these findings to mouse LGN axons (Figure 4A-B and S9).

Our data further show that the OFF subfields have less retinotopic scatter than ON subfields

(Figure 4C-F), an organization underlying the structure of orientation maps in V1 of other

species (Kremkow et al., 2016; Lee et al., 2016). Considering that mouse V1 lacks a systematic

orientation map (Ohki et al., 2007), our results suggest such polarity-dependent retinotopy is

conserved across various visual cortical architectures through evolution.

Incomplete axon morphology.

The axon morphology estimates of this study were performed on single two-photon imaging

planes (~ 190 x 190 μm in the XY plane and a few microns in Z). Given that the arbor of an LGN

axon in V1 can be more than 500 μm in all three dimensions (Antonini et al., 1999), our

measurements were substantially incomplete. We reasoned that, with such crude

measurements, any subtle differences in axon morphology would fail to reach significance.

However, to our surprise, the DS axons and RF axons showed significant differences

consistently across all morphology measurements (Figure 2H-I and S6). This indicates that there

should be dramatic differences between DS and RF axons in their full morphology, but further

studies are needed to fully characterize the anatomical differences between the two

physiological types.

Methods

Surgery and animal preparation

In total, six Vipr2-IRES2-Cre-neo mice (3 male, 3 female) were used in this study. The surgery

included a stereotaxic viral injection and a cranial window/head-plate implantation. During the

injection, a glass pipette back-loaded with a 1:3 mixture of AAV1-CAG-mRuby (custom made

from plasmid addgene 107744, titer: 1.6×10^{12} vg/ml) and AAV1-Syn (or CAG)-FLEX-GCaMP6s

(addgene: 100845-AAV1, titer 4.3×10^{13} vg/ml or 100842-AAV1, titer 1.8×10^{13} vg/ml,

respectively) was slowly lowered into left LGN (2.3 mm posterior 2.3 mm lateral from bregma,

2.6 mm below pia) through a burr hole. 5 minutes after reaching the targeted location, 200 nL

of virus mixture was injected into the brain over 10 minutes by a hydraulic nanoliter injection

system (Nanoject III, Drummond). The pipette then stayed for an additional 10 minutes before

it was slowly retracted out of the brain. Immediately after injection, a titanium head-plate and

a 5 mm glass cranial window were implanted over left V1 (de Vries et al., 2020; for detailed

protocol see Goldey et al., 2014) allowing *in vivo* two-photon imaging during head fixation.

After surgery, the animals were allowed to recover for at least 5 days before retinotopic

mapping with intrinsic signal during anesthesia (for detailed retinotopic protocol, see Juavinett

et al., 2017). After retinotopic mapping, animals were handled and habituated to the imaging

rig for two additional weeks (de Vries et al., 2020) before *in vivo* two-photon imaging.

All experiments and procedures were approved by the Allen Institute Animal Care and Use

Committee.

Histology

To characterize the Cre expression pattern, Vipr2-IRES2-Cre-neo mice were crossed with the

Ai14 reporter line (Madisen et al., 2010) to generate Vipr2-IRES2-Cre-neo/wt; Ai14/wt animals.

Mice heterozygous for both Cre and the reporter were perfused and brains collected. Briefly,

mice were anesthetized with 5% isoflurane and 10 ml of saline (0.9% NaCl) followed by 50 ml of

freshly prepared 4% paraformaldehyde (PFA) was pumped intracardially at a flow rate of 9

ml/min. Brains were immediately dissected and post-fixed in 4% PFA at room temperature for

3-6 hours and then overnight at 4 °C. After fixation, brains were incubated in 10% and then 30%

sucrose in PBS for 12-24 hours at 4 °C before being cut into 50 μm sections by a freezing-sliding

microtome (Leica SM 2101R). Sections with LGN and V1 were then mounted on gelatin-coated

slides and cover-slipped with mounting media (Prolong Diamond Antifade Mounting Media,

P36965, ThermoFisher).

To verify the injection location and GCaMP expression, the brain tissues from all experimental

mice were collected and sectioned with a similar procedure after all imaging sessions with

additional steps with antibody staining to enhance the GCaMP signal before mounting and

coverslipping. During antibody staining, sections containing LGN and V1 were blocked with 5%

normal donkey serum and 0.2% Triton X-100 in PBS for one hour, incubated in an anti-GFP

primary antibody (1:5000 diluted in the blocking solution, Abcam, Ab13970) for 48-72 hours at

4 °C, washed the following day in 0.2% Triton X-100 in PBS and incubated in a Alexa-488

conjugated secondary antibody (1:500, 703-545-155, Jackson ImmunoResearch) and DAPI.

The sections were then imaged with Zeiss AxioImager M2 widefield microscope with a 10x/0.3

NA objective. Fluorescence from antibody enhanced GCaMP and mRuby3 were extracted from

filter sets Semrock GFP-1828A (excitation 482/18 nm, emission 520/28 nm, dichroic cutoff 495

nm) and Zeiss # 20 (excitation 546/12 nm, emission 608/32 nm, dichroic cutoff 560 nm),

respectively.

In vivo **two-photon imaging**

Following the completion of surgical recovery, retinotopic mapping, and habituation (usually

more than three weeks after initial surgery), thalamocortical axons from LGN labeled with

GCaMP6s were evident in superficial- and mid-layers (0 – 500 µm) in V1 through the cranial

window (Figure 1E). We choose 8 depths for in vivo two-photon imaging (50, 100, 150, 200,

250, 300, 350 and 400 µm below pia). Imaging was done in a columnar fashion: at each cortical

location calcium activities at 3-8 depths were imaged plane-by-plane over multiple sessions

with a conventional 2-photon microscope or in a single session with a multi-plane 2-photon

microscope (described below). In both microscopes, a 16x/0.8 NA water immersion objective

(Nikon 16XLWD-PF) was rotated to 24 degrees from horizontal to image visual cortex using a

commercial rotating head (Sutter MOM). mitted light was first split by a 735 nm dichroic mirror

(FF735-DiO1, Semrock). The short-wavelength light was filtered by a 750 nm short-pass filter

(FESH0750, Thorlabs) and a 470-588 nm bandpass emission filter (FF01-514/44-25, Semrock)

before collected as GCaMP signal, while the long-wavelength light was filtered by a 538-722 nm

band-pass emission filter (FF01-630/92-30, Semrock) before collected as mRuby signal. Image

acquisition was controlled using Vidrio ScanImage software for both scopes (Pologruto et al.,

2003, Vidrio LLC). To maintain constant immersion of the objective, we used gel immersion

(Genteal Gel, Alcon).

Single-plane two-photon imaging. Four mice were imaged with two-photon excitation

generated by laser illumination from a Ti:sapphire laser (Coherent Chameleon Ultra II) tuned to

920 nm. A single z-plane (179.2 x 179.2 μm with 512 x 512 pixels resolution) was imaged for

each session at a frame rate of about 30 Hz with an 8 KHz resonate scanner (Cambridge

Technology, CRS 8K).To maintain constant imaging depth automatic z-drift correction functions

were implemented for experiments using the MOM motors. Briefly, a correction z-stack (± 50

μm from targeted depth, 2 μm step depth) was recorded before each imaging session and,

during the session, the current imaging plane was continuously compared to each plane in the

correction z-stack. If a drift in depth was detected, the stage was automatically adjusted to

compensate for the drift, thus maintaining constant imaging depth. We found this procedure

crucial to our experiments since the boutons are small objects and a few-micron-drift in depth

would result in imaging a different set of boutons.

Multi-plane two-photon imaging. The other two mice were imaged by a custom-built, multi-

plane two-photon microscope (DeepScope, Liu et al., 2018) in which a liquid crystal spatial light

modulator (SLM; HSP-512, Meadowlark Optics) shapes the pupil wavefront to implement fast-

focusing and adaptive optics. The objective pupil was slightly underfilled (~0.65 effective NA)

and correction of systemic aberrations was performed with fluorescent beads to maintain near

diffraction-limited focusing over a 200 um range. Two-photon excitation was produced by laser

light from a commercial solid-state laser (Spectra-Physic Insight X3 laser) tuned to 940 nm. With

this microscope, we simultaneously recorded calcium activity from planes at 5 different depths

(50, 100, 150, 200, 250 μm) in single imaging sessions. Individual frames (125 x 125 μm with

512 x 512 pixels resolution) were acquired at an overall framerate of ~37 Hz with a volume rate

of 7.4 Hz. The DeepScope showed nearly zero z-drift for a prolonged duration (< 2 μm over 24

hours), so we did not implement z-correction in sessions using DeepScope.

In another set of experiments, we used DeepScope to assess the effect of adaptive optics

adjusted on individual animals. The correction procedure was similar to the method described

in Sun et al., 2016. For two mice, 1 μm beads (Thermo Fisher, F8821) were deposited on top of

the brain surface under the coverglass during the initial surgery. Prior to the imaging session,

modal optimization over 12 Zernike modes (up to j = 15 Noll ordering, excluding piston and tilt)

was run to identify the SLM pattern that maximized the beads' fluorescent signal. Then, during

the imaging session, this SLM pattern was put on and off alternatively for consecutive two-

photon imaging frames (one frame on, one frame off, repeated) and drifting gratings were

displayed. After imaging, the interleaving movie was separated into two movies: one with

adaptive optics and the other without. Bouton's tuning properties were then extracted from

each movie and compared against each other.

All imaging sessions were performed during head fixation with the standard Allen Institue Brain

Observatory *in vivo* imaging stage (de Vries et al., 2020).

Visual stimulation

All visual stimuli were generated and displayed by Retinotopic_Mapping python package

(https://github.com/zhuangjun1981/retinotopic_mapping, Zhuang et al., 2017) over PsychoPy

software (https://www.psychopy.org, Peirce et al., 2019) on a 24-inch LCD monitor (ASUS

PA248Q, frame rate 60 Hz, 1920 x 1200 pixels, mean luminance 45.3 cd/m^2) placed 15 cm from

the mouse's right eye (covering 120° x 95° of monocular visual space). We displayed locally

sparse noise and/or full-field drifting grating in each imaging session to measure receptive fields

and orientation/direction/spatial and temporal frequency tuning properties, respectively. In

most sessions, we also displayed a five-minute full-field mid-luminance gray to measure

spontaneous activity. For locally sparse noise, bright and dark squares (5° x 5°) were displayed

in a random sequence on a grid tiling the entire monitor. At any given time, multiple squares

could be displayed but the minimum distance between those squares should be no less than

50°. Each square lasted 100 ms and in total was displayed 50 times. For drifting gratings, the

combinations of 12 directions (every 30°), 3 spatial frequencies (SF, 0.01, 0.04 and 0.16 cpd)

and 3 temporal frequencies (TF, 1, 4, 15 Hz) were displayed. Each display lasted 1 second and

was spaced by 1-second mean luminance grey period. In total, 3 x 3 x 12 + 1 (blank) = 109

conditions were randomly displayed in each iteration and the whole sequence contained 13

iterations. All stimuli were spherically corrected so that they were presented with accurate

visual angles on the flat screen (Zhuang et al., 2017).

2-photon Image preprocessing

The recorded two-photon movies for each imaging plane were first temporally averaged across

5 frames (Sutter scope) or 2 frames (DeepScope), then were motion-corrected using rigid body

transform based on phase correlation by a custom-written python package

(https://github.com/zhuangjun1981/stia/tree/master/stia, Zhuang et al., 2017). We performed

motion correction on the red channel (mRuby) and applied the offsets to the green channel

(GCaMP). To generate regions of interest (ROIs) for boutons, the motion-corrected movies were

further temporally downsampled by a factor of 3 and then processed with constrained non-

negative matrix factorization (CNMF, Pnevmatikakis et al., 2016) implemented in the CaImAn

python library (https://github.com/flatironinstitute/CaImAn, Giovannucci et al., 2018). The

resulting ROI masks were normalized by subtracting the mean and dividing the standard

deviation. Then pixels below 3 were set to 0. These ROIs were further filtered by their size (ROIs

smaller than 1.225 μm^2 or larger than 12.25 μm^2 were excluded), position (ROIs within the

motion artifacts were excluded) and overlap (for ROIs with more than 20% overlap, the smaller

ones were excluded). For each retained ROI, a neuropil ROI was created as the region between

two contours by dilating the ROI's outer border by 1 and 8 pixels excluding the pixels within the

union of all ROIs. The calcium trace for each ROI was calculated by the mean of pixel-wise

product between the ROI and each frame of the movie, and its neuropil trace was calculated in

the same way using its neuropil ROI. To remove the neuropil contamination, the neuropil

contribution of each ROI's calcium trace was estimated by a linear model and optimized by

gradient descendent regression with a smoothness regularization (Zhuang et al., 2017; de Vries

et al., 2020). Nonetheless, the estimates of neuropil contamination in our datasets were

generally low (percentage of neuropil signal that contributed to the ROIs signal: 11.5 ± 0.5%)

likely due to the small volume and sparse distribution of afferent axons in V1. As reported

previously, a high skewness is an indication of active calcium activity (Mukamel et al., 2009,

Dipoppa et al, 2018). The ROIs with skewness greater than 0.6 were defined as "active" boutons

and were included in the following study, while others were excluded (Table S1). For each

imaging session, a comprehensive file in Neurodata Without Borders (nwb) 1.0 format was

generated to store and share metadata, visual stimuli, all preprocessing results, and final

calcium traces using the "ainwb" package ([https://github.com/AllenInstitute/nwb-](https://github.com/AllenInstitute/nwb-api/tree/master/ainwb)

[api/tree/master/ainwb](https://github.com/AllenInstitute/nwb-api/tree/master/ainwb)).

For the neuropil subtracted calcium traces with a minimum value below 1, an offset was added

to the whole trace so that the trace minimum was 1. This procedure ensured a positive baseline

for all units in the df/f calculation (below).

Bouton clustering

The correlation-based bouton clustering procedure was based on previously reported

algorithms (Liang et al., 2018) and the same procedure was performed on each imaging plane.

First, we detected calcium events for each active bouton in the imaging plane. The calcium

trace was smoothed by a gaussian filter with sigma of 0.1 seconds. Then event onsets were

detected as up-crosses over a threshold of 3 standard deviations above the mean. A period 3

seconds before and after each onset was defined as an event window and the union of all event

windows for a particular ROI was saved. Second, for any given pair of boutons, the union of

event windows from both boutons was calculated and the calcium trace within the union

window was extracted and concatenated for each bouton. Then a Pearson correlation

coefficient of the two concatenated traces was calculated for this pair. By performing this

procedure on every pair of active ROIs, we generated a correlation coefficient matrix for each

imaging plane. We found the event detection important because it confined the correlation to

the period only at least one bouton in the pair was active thus avoided correlating the noise

during the inactive period. Third, the correlation coefficient matrices were further thresholded

to reduce noise: the correlation coefficients for a given bouton were maintained if the

coefficients were larger than 0.5 or if they exceeded 3 standard deviations above the mean

value of all the coefficients between this bouton and all others. Otherwise, they will be set to 0.

Third, a hierarchy clustering was performed to a given imaging plane using "1 – thresholded

correlation coefficient matrix" as the distance matrix using Scipy.cluster.hierarchy library with a

"weighted" method

(https://docs.scipy.org/doc/scipy/reference/generated/scipy.cluster.hierarchy.linkage.html).

For each imaging plane, we calculated a cophenetic correlation distance to evaluate the

clustering performance. Fourth, we use a threshold of 1.3 to separate clusters since ~ 1.5 shows

up as a relatively natural cut-off in the dendrograms. This threshold appeared somewhat

conservative on the clustered correlation coefficient matrix (Figure 1M). Fifth, after clustering,

each cluster was identified as a separate axon and its calcium trace was calculated as mean

calcium trace of all boutons belonging to this cluster, weighted by the sum of their ROI weights.

For each detected axon, three values were extracted to estimate the morphology: (1) bouton

number of this axon; (2) maximum distance among all bouton pairs belonging to this axon; (3)

area of the convex polygon encapsulated by all the boutons belonging to this axon. For axons

with only one bouton, the metrics 2 and 3 were set to be "nan" and were excluded from

statistical analysis.

To generate stacked axon mask in Figure 2H, we first selected axons with at least two boutons

for each group (DS axons or RF axons). Then the binary mask of each individual axon was

rotated and scaled into the same standard orientation (up: anterior, left: lateral) and centered

to its own center of mass. All centered axon masks were summed together to generate a

summed population axon mask for each group. Each summed population axon mask was then

divided by its own maximum value, generating a normalized population axon mask (Figure 2H).

Axons with both significant RF and direction selectivity were excluded from this analysis to

avoid double counting.

Receptive field analysis

We calculated the units' spatial receptive fields from their responses to the locally sparse noise

stimulus. For a given square flash at a particular location with a particular sign, the calcium

traces within a temporal window around the square onsets were extracted and aligned by the

square onsets. An event-triggered average trace was then calculated as the mean trace across

all aligned traces. The mean value of the averaged trace in the window [-0.5, 0] seconds from

the onset was calculated as the baseline and event-triggered df/f trace was calculated as

(average trace - baseline) / baseline. The response amplitude to this particular square was then

calculated as the mean df/f in the window [0, 0.5] second from the stimulus onset. By repeating

this procedure for all squares presented in locally sparse noise stimuli, a 2d amplitude map was

generated tiling the entire visual space covered by the monitor for each sign ("ON" from bright

squares and "OFF" from dark squares). From the amplitude map, a z-score map was calculated

by subtracting the mean and dividing the standard deviation of the entire map. The z-score map

was then smoothed by a Gaussian filter with a sigma of 1 pixel and upsampled by a ratio of 10

with cubic interpolation. The resulting map was defined as the receptive field map with a

resolution of 0.5 degrees. The peak value of ON and OFF receptive field map combined was

defined as RF strength plotted in Figure 2B and S4E. We defined a receptive field map with a

peak value exceeding 1.6 as having a significant spatial RF. For each significant spatial RF, an RF

mask was generated by thresholding either with a value of 1.6 (for maps with a peak less than

4) or with a value of 40% of its peak (for maps with a peak greater than 4, Zhuang et al., 2013).

The RF area (plotted in Figure 4B and S9B) was calculated as the area of the binary mask after

thresholding. The RF center was calculated as the location of the center of mass from the

weighted mask after thresholding.

For the retinotopy analysis in Figure 4C-D, we rotated and scaled each imaging plane so that

they had the same orientation of cortical surface (up: anterior, down: posterior, left: lateral,

right: medial) with same pixel size. For each of these planes, we obtained all significant RFs

from boutons in this plane and calculated their centers. The population RF was then calculated

by summing all significant RFs and thresholding with the same steps described above. Then the

relative RF location of each individual RF was calculated as the vector pointing from the

population RF center to the individual RF center. The relative RF locations were then

normalized by subtracting mean and dividing by standard deviation (z-score relative RF

location). To generate the normalized retinotopy map for each imaging plane, the individual z-

score relative RF locations were color-coded and plotted to the center of their boutons within

the rotated and scaled imaging plane. Since the retinotopy gradient is relatively uniform across

mouse V1 (Zhuang et al., 2017), these procedures allowed us to superimpose the normalized

retinotopy maps from all imaging planes together. We performed this for ON and OFF subfields

and altitude and azimuth separately generating four retinotopic maps with respect to each sign

and visual space dimension (Figure 4C). To quantify retinotopic gradient and scatter, we

performed linear regression between cortical locations and z-score relative RF locations from

these four superimposed retinotopic maps with relevant axis (anterior-posterior axis against

altitude and medial-lateral axis against azimuth, Figure 4D).

Grating response analysis

The units' responses to drifting gratings were analyzed by the event-triggered average

procedure similar to the RF response analysis. The difference is the temporal window to

calculate response was [0, 1] second from the grating onsets since the duration of each grating

was 1 second. We first define a unit having significant responses to the drifting gratings if it

reached two criteria: 1) it responded to at least one grating condition with a z-score greater

than 3 and 2) the Oneway ANOVA analysis of df/f responses across all grating conditions

(including blank trials) had a p-value less than 0.01. For a unit that had a significant response to

gratings, its peak condition was defined as the conditions with the highest df/f response. Its

direction tuning curve was extracted as its df/f responses to different directions with the TF and

SF the same as the peak condition and, if the minimum response from this curve was below

zero, an offset was added to the whole curve so that the minimum was zero. Its SF and TF

tuning curves were extracted by a similar procedure. From direction tuning curve, we calculated

the global orientation index (gOSI) as

546
$$gOSI = \frac{\left|\sum R_j e^{i2\theta_j}\right|}{\sum R_j}$$

547   Where j represents different direction conditions, R represents df/f response in a given

548   direction and θ represents the direction. The preferred orientation as the angle of $\sum R_j e^{i2\theta_j}$.

549   We calculated global direction index (gDSI) as

550
$$gDSI = \frac{\left|\sum R_j e^{i\theta_j}\right|}{\sum R_j}$$

551   And the preferred direction as the angle of $\sum R_j e^{i\theta_j}$. From the SF tuning curve, we calculated

552   the preferred SF as

553
$$peak\ SF = 2^{\left(\frac{\sum R_j log_2 SF_j}{\sum R_j}\right)}$$

554   Where j represents different SF conditions. From TF tuning curve, we calculated the preferred

555   TF as

556
$$peak\ TF = 2^{\left(\frac{\sum R_j log_2 TF_j}{\sum R_j}\right)}$$

Where j represents different TF conditions. For the units that did not have significant grating

response, all these measurements were set to be "nan". We defined a unit to be direction-

selective if it had a gDSI greater than 0.5 and orientation-selective if it had a gOSI greater than

0.3.

For each imaging plane, we calculated a nasal-to-temporal index (NTI, see main text):

$$NTI = \frac{num_{nasal} - num_{temporal}}{num_{nasal} + num_{temporal}}$$

Where num_{nasal} is the count of DS units that had preferred direction in the range of [-45°, 45°]

and $num_{temporal}$ is the count of DS units that had preferred direction in the range of [135°, 225°].

Only the planes with num_{nasal} + $num_{temporal}$ greater than 10 were included in the analysis. We

also calculated an ON-OFF index:

$$ON - OFF\ Index = \frac{num_{ON} - num_{OFF}}{num_{ON} + num_{OFF}}$$

Where num_{ON} is the count of RF units that had only ON subfield and num_{OFF} is the count of RF

units that had only OFF subfield. Only the planes with num_{ON} + num_{OFF} greater than 10 were

included in the analysis.

The imaging preprocessing, nwb packaging, bouton clustering, receptive field analysis, grating

response analysis, and functional type classification were performed by a custom-written

python package "corticalmapping"

(https://github.com/zhuangjun1981/corticalmapping/tree/master/corticalmapping).

Statistics

To control the variability across imaging planes (different axon/bouton density, vasculature

pattern, expression level), almost all statistics were extracted from each plane and separated

for different functional types if necessary. For statistics that only one number can be drawn

from each imaging plane (bouton/axon count, NTI, etc), they were presented as mean ±

standard deviation. For statistics that a population distribution can be drawn from each imaging

plane (gDSI, gOSI, peak SF/TF, RF strength, RF area, bouton per axon, max bouton distance,

axon coverage area), mean for each plane was calculated first and mean ± standard error was

reported across imaging planes. The comparisons between different functional types within

imaging plane were performed by Wilcoxon rank-sum test, and comparisons between imaging

planes were performed by Mann-Whitney U test since the distribution of the statistics were

highly skewed and non-gaussian. We have verified our results of these two tests by paired t-

test and independent t-test respectively, and they all agreed with the non-parametric tests (not

shown). In the main text, the significant tests were Wilcoxon rank-sum test if not specified. All

significance tests were two-tailed.

Acknowledgments

Funding was provided by the Allen Institute for Brain Science and award number NS104949

from the National Institute of Neurological Disorders and Stroke. The contents of this

manuscript are solely the responsibility of the authors and do not necessarily represent the

official views of the National Institutes of Health and National Institute of Neurological

Disorders and Stroke. We thank Dr. Roger Tsien for providing tdTomato construct. We thank

many staff members of the Allen Institute, especially the In Vivo Sciences team for surgeries

and the Manufacturing and Processing Engineering team for hardware support. We wish to

thank the Allen Institute for Brain Science founder, Paul G. Allen, for his vision, encouragement,

and support.

References

Anderson JC, da Costa NM, Martin KA (2009) The W cell pathway to cat primary visual cortex. J Comp Neurol 516:20-35.

Antonini A, Fagiolini M, Stryker MP (1999) Anatomical correlates of functional plasticity in mouse visual cortex. J Neurosci 19:4388-4406.

Bereshpolova Y, Stoelzel CR, Su C, Alonso JM, Swadlow HA (2019) Activation of a Visual Cortical Column by a Directionally Selective Thalamocortical Neuron. Cell Rep 27:3733-3740 e3733.

Cleland BG, Dubin MW, Levick WR (1971) Sustained and transient neurones in the cat's retina and lateral geniculate nucleus. J Physiol 217:473-496.

Cruz-Martin A, El-Danaf RN, Osakada F, Sriram B, Dhande OS, Nguyen PL, Callaway EM, Ghosh A, Huberman AD (2014) A dedicated circuit links direction-selective retinal ganglion cells to the primary visual cortex. Nature 507:358-361.

de Vries, S.E.J., Lecoq, J.A., Buice, M.A. et al. (2020) A large-scale standardized physiological survey reveals functional organization of the mouse visual cortex. Nat Neurosci 23:138-151

Denman, D.J., and Contreras, D. (2016). On Parallel Streams through the Mouse Dorsal Lateral Geniculate Nucleus. Front. Neural Circuits 10, 20.

Dhande OS, Stafford BK, Lim JA, Huberman AD (2015) Contributions of Retinal Ganglion Cells to Subcortical Visual Processing and Behaviors. Annu Rev Vis Sci 1:291-328.

Durand S, Iyer R, Mizuseki K, de Vries S, Mihalas S, Reid RC (2016) A Comparison of Visual Response Properties in the Lateral Geniculate Nucleus and Primary Visual Cortex of Awake and Anesthetized Mice. J Neurosci 36:12144-12156.

Garrett ME, Nauhaus I, Marshel JH, Callaway EM (2014) Topography and areal organization of mouse visual cortex. J Neurosci 34:12587-12600.

Gilbert CD, Wiesel TN (1979) Morphology and intracortical projections of functionally characterised neurones in the cat visual cortex. Nature 280:120-125.

Grubb MS, Thompson ID (2003) Quantitative characterization of visual response properties in the mouse dorsal lateral geniculate nucleus. J Neurophysiol 90:3594-3607.

Hei X, Stoelzel CR, Zhuang J, Bereshpolova Y, Huff JM, Alonso JM, Swadlow HA (2014) Directional selective neurons in the awake LGN: response properties and modulation by brain state. J Neurophysiol 112:362-373.

Hoffmann KP, Stone J, Sherman SM (1972) Relay of receptive-field properties in dorsal lateral geniculate nucleus of the cat. J Neurophysiol 35:518-531.

Hubel DH, Wiesel TN (1962) Receptive fields, binocular interaction and functional architecture
in the cat's visual cortex. J Physiol 160:106-154.

Huberman AD, Wei W, Elstrott J, Stafford BK, Feller MB, Barres BA (2009) Genetic identification
of an On-Off direction-selective retinal ganglion cell subtype reveals a layer-specific subcortical
map of posterior motion. Neuron 62:327-334.

Humphrey AL, Sur M, Uhlrich DJ, Sherman SM (1985) Termination patterns of individual X- and
Y-cell axons in the visual cortex of the cat: projections to area 18, to the 17/18 border region,
and to both areas 17 and 18. J Comp Neurol 233:190-212.

Jaepel, J., Hubener, M., Bonhoeffer, T., and Rose, T. (2017). Lateral geniculate neurons
projecting to primary visual cortex show ocular dominance plasticity in adult mice. Nat Neurosci
20, 1708–1714.

Jimenez LO, Tring E, Trachtenberg JT, Ringach DL (2018) Local tuning biases in mouse primary
visual cortex. J Neurophysiol 120:274-280.

Jin JZ, Weng C, Yeh CI, Gordon JA, Ruthazer ES, Stryker MP, Swadlow HA, Alonso JM (2008) On
and off domains of geniculate afferents in cat primary visual cortex. Nat Neurosci 11:88-94.

Jin J, Wang Y, Lashgari R, Swadlow HA, Alonso JM (2011) Faster thalamocortical processing for
dark than light visual targets. J Neurosci 31:17471-17479.

Kondo S, Ohki K (2016) Laminar differences in the orientation selectivity of geniculate afferents
in mouse primary visual cortex. Nat Neurosci 19:316-319.

Kremkow J, Jin J, Wang Y, Alonso JM (2016) Principles underlying sensory map topography in
primary visual cortex. Nature 533:52-57.

Kremkow J, Alonso JM (2018) Thalamocortical Circuits and Functional Architecture. Annu Rev
Vis Sci 4:263-285.

Lee KS, Huang X, Fitzpatrick D (2016) Topology of ON and OFF inputs in visual cortex enables an
invariant columnar architecture. Nature 533:90-94.

Lennie P (1980) Parallel visual pathways: a review. Vision Res 20:561-594.

Liang L, Fratzl A, Goldey G, Ramesh RN, Sugden AU, Morgan JL, Chen C, Andermann ML (2018) A
Fine-Scale Functional Logic to Convergence from Retina to Thalamus. Cell 173:1343-1355.

Marshel JH, Kaye AP, Nauhaus I, Callaway EM (2012) Anterior-posterior direction opponency in
the superficial mouse lateral geniculate nucleus. Neuron 76:713-720.

Mazade R, Jin J, Pons C, Alonso JM (2019) Functional Specialization of ON and OFF Cortical
Pathways for Global-Slow and Local-Fast Vision. Cell Rep 27:2881-2894 e2885.

Nassi JJ, Callaway EM (2009) Parallel processing strategies of the primate visual system. Nat Rev
Neurosci 10:360-372.

Ohki K, Reid RC (2007) Specificity and randomness in the visual cortex. Curr Opin Neurobiol
17:401-407.

Piscopo DM, El-Danaf RN, Huberman AD, Niell CM (2013) Diverse visual features encoded in
mouse lateral geniculate nucleus. J Neurosci 33:4642-4656.

Reid RC, Alonso JM (1995) Specificity of monosynaptic connections from thalamus to visual
cortex. Nature 378:281-284.

Román Rosón, M., Bauer, Y., Kotkat, A.H., Berens, P., Euler, T., and Busse, L. (2019). Mouse
dLGN Receives Functional Input from a Diverse Population of Retinal Ganglion Cells with
Limited Convergence. Neuron 102, 462-476.e8.

Rompani SB, Mullner FE, Wanner A, Zhang C, Roth CN, Yonehara K, Roska B (2017) Different
Modes of Visual Integration in the Lateral Geniculate Nucleus Revealed by Single-Cell-Initiated
Transsynaptic Tracing. Neuron 93:767-776 e766.

Roth MM, Dahmen JC, Muir DR, Imhof F, Martini FJ, Hofer SB (2016) Thalamic nuclei convey
diverse contextual information to layer 1 of visual cortex. Nat Neurosci 19:299-307.

Sabbah S, Gemmer JA, Bhatia-Lin A, Manoff G, Castro G, Siegel JK, Jeffery N, Berson DM (2017)
A retinal code for motion along the gravitational and body axes. Nature 546:492-497.

Seabrook TA, Burbridge TJ, Crair MC, Huberman AD (2017) Architecture, Function, and
Assembly of the Mouse Visual System. Annu Rev Neurosci 40:499-538.

Siegle JH et al. (2019) A survey of spiking activity reveals a functional hierarchy of mouse
corticothalamic visual areas. bioRxiv:805010.

Stone J, Dreher B, Leventhal A (1979) Hierarchical and parallel mechanisms in the organization
of visual cortex. Brain Res 180:345-394.

Sun W, Tan Z, Mensh BD, Ji N (2016) Thalamus provides layer 4 of primary visual cortex with
orientation- and direction-tuned inputs. Nat Neurosci 19:308-315.

Sur M, Sherman SM (1982) Linear and nonlinear W-cells in C-laminae of the cat's lateral
geniculate nucleus. J Neurophysiol 47:869-884.

Suresh, V., Ciftcioglu, U.M., Wang, X., Lala, B.M., Ding, K.R., Smith, W.A., Sommer, F.T., and
Hirsch, J.A. (2016). Synaptic Contributions to Receptive Field Structure and Response Properties
in the Rodent Lateral Geniculate Nucleus of the Thalamus. J Neurosci 36, 10949–10963.

Wilson PD, Stone J (1975) Evidence of W-cell input to the cat's visual cortex via the C laminae of
the lateral geniculate nucleus. Brain Res 92:472-478.

Wilson PD, Rowe MH, Stone J (1976) Properties of relay cells in cat's lateral geniculate nucleus:
a comparison of W-cells with X- and Y-cells. J Neurophysiol 39:1193-1209.

Yeh CI, Xing D, Shapley RM (2009) "Black" responses dominate macaque primary visual cortex
v1. J Neurosci 29:11753-11760.

Zhao X, Chen H, Liu X, Cang J (2013) Orientation-selective responses in the mouse lateral
geniculate nucleus. J Neurosci 33:12751-12763.

Zhuang J, Stoelzel CR, Bereshpolova Y, Huff JM, Hei X, Alonso JM, Swadlow HA (2013) Layer 4 in
primary visual cortex of the awake rabbit: contrasting properties of simple cells and putative
feedforward inhibitory interneurons. J Neurosci 33:11372-11389.

Zhuang J, Ng L, Williams D, Valley M, Li Y, Garrett M, Waters J (2017) An extended retinotopic
map of mouse cortex. Elife 6:e18372.

Method References

Method References

Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD (2018) Vision and
Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron
98:602-615 e608.

Giovannucci A, Friedrich J, Gunn P, Kalfon J, Brown BL, Koay SA, Taxidis J, Najafi F, Gauthier JL,
Zhou P, Khakh BS, Tank DW, Chklovskii DB, Pnevmatikakis EA (2019) CaImAn an open source
tool for scalable calcium imaging data analysis. Elife 8:e38173.

Goldey GJ, Roumis DK, Glickfeld LL, Kerlin AM, Reid RC, Bonin V, Schafer DP, Andermann ML
(2014) Removable cranial windows for long-term imaging in awake mice. Nat Protoc 9:2515-
2538.

Juavinett AL, Nauhaus I, Garrett ME, Zhuang J, Callaway EM (2017) Automated identification of
mouse visual areas with intrinsic signal imaging. Nat Protoc 12:32-43.

Liu R, Ball N, Brockill J, Kuan L, Millman D, White C, Leon A, Williams D, Nishiwaki S, de Vries S,
Larkin J, Sullivan D, Slaughterbeck C, Farrell C, Saggau P (2018) Multi-plane Imaging of Neural
Activity From the Mammalian Brain Using a Fast-switching Liquid Crystal Spatial Light
Modulator. bioRxiv:506618.

Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD, Hawrylycz
MJ, Jones AR, Lein ES, Zeng H (2010) A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. Nat Neurosci 13:133-140.

Mukamel EA, Nimmerjahn A, Schnitzer MJ (2009) Automated analysis of cellular signals from
large-scale calcium imaging data. Neuron 63:747-760.

Peirce J, Gray JR, Simpson S, MacAskill M, Hochenberger R, Sogo H, Kastman E, Lindelov JK
(2019) PsychoPy2: Experiments in behavior made easy. Behav Res Methods 51:195-203.

Pnevmatikakis EA, Soudry D, Gao Y, Machado TA, Merel J, Pfau D, Reardon T, Mu Y, Lacefield C,
Yang W, Ahrens M, Bruno R, Jessell TM, Peterka DS, Yuste R, Paninski L (2016) Simultaneous
Denoising, Deconvolution, and Demixing of Calcium Imaging Data. Neuron 89:285-299.

Pol6gruto TA, Sabatini BL, Svoboda K (2003) ScanImage: flexible software for operating laser
scanning microscopes. Biomed Eng Online 2:13.