Eukaryotic-like ribosomal RNA in Asgard archaea

Authors: Petar I. Penev[1,2], Sara Fakhretaha-Aval[1,3], Vaishnavi J. Patel[4], Jamie J. Cannone[4], Robin R. Gutell[4], Anton S. Petrov[1,3*], Loren Dean Williams[1,2,3*], Jennifer B. Glass[1,2,5*]

Affiliations:

[1]NASA Center for the Origin of Life, Georgia Institute of Technology, Atlanta, GA 30332-0400, USA

[2]School of Biological Sciences, Georgia Institute of Technology, North Avenue, Atlanta, GA 30332, USA

[3]School of Chemistry and Biochemistry, Georgia Institute of Technology, 901 Atlantic Dr, Atlanta, GA 30332, USA

[4]Department of Integrative Biology, The University of Texas at Austin, 2415 Speedway #C0930, Austin, TX 78712, USA

[5]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, 311 Ferst Dr, Atlanta, GA 30332, USA

*Correspondence to: anton.petrov@biology.gatech.edu; loren.williams@chemistry.gatech.edu; jennifer.glass@eas.gatech.edu

Abstract

The ribosome's common core connects all life back to a common ancestor and serves as a window to relationships among organisms. In eukaryotes, the common core is decorated by expansion segments (ES's) that vastly increase ribosomal RNA (rRNA) size. Supersized ES's have not been observed previously in Archaea, and the origin of eukaryotic ES's remains enigmatic. We discovered that the large subunit (LSU) rRNA of two Asgard phyla, *Lokiarchaeota* and *Heimdallarchaeota,* considered to be the closest modern archaeal cell lineages to Eukarya, bridge the gap in size between prokaryotic and eukaryotic LSU rRNA. The elongated LSU rRNAs in *Lokiarchaeota* and *Heimdallarchaeota* stem from the presence of two eukaryotic-sized ES's, ES9 and ES39. We applied chemical footprinting experiments to study the structure of *Lokiarchaeota* ES39. Furthermore, we used covariation and sequence analysis to study the evolution of Asgard ES39's and ES9's. By defining the common eukaryotic ES39 signature old, we found that Asgard ES39's have more and longer helices than eukaryotic ES39's. While Asgard ES39's have sequences and structures distinct from eukaryotic ES39's, we found an overall conservation of a three-way junction across the Asgard species that matches eukaryotic ES39 topology. The similarities between Asgard and eukaryotic ES39 rRNA sizes and structural topologies are consistent with the accretion model of ribosomal evolution. The common ancestor of eukaryotes and Asgard archaea likely had already had a three-way junction topology in the ES39 region, complexity previously only known in eukaryotes. Eukaryotic-like expansions within Asgard rRNA are consistent with the two-domain tree of life.

Introduction

The ribosome connects all life on Earth back to the Last Universal Common Ancestor (LUCA) (Woese and Fox 1977). The small ribosomal subunit (SSU) decodes mRNA and the large ribosomal subunit (LSU) links amino acids together to produce coded protein. Both subunits are made of ribosomal RNA (rRNA) and ribosomal protein (rProtein). All cytoplasmic ribosomes contain a structurally conserved universal common core, comprised of 2800 nucleotides and 28 rProteins, and including the peptidyl transferase center (PTC) in the LSU and the decoding center (DCC) in the SSU (Melnikov, et al. 2012; Bernier, et al. 2018). The rRNA of the common core is a reasonable approximation of the rRNA in LUCA and is most similar to rRNA of extant bacteria (Melnikov, et al. 2012; Petrov, et al. 2014b; Bernier, et al. 2018).

In Eukarya, the rRNA of the common core is elaborated by expansion segments (ES's, **Fig. 1**) (Veldman, et al. 1981; Clark, et al. 1984; Hassouna, et al. 1984; Gonzalez, et al. 1985; Michot and Bachellerie 1987; Bachellerie and Michot 1989; Gutell 1992; Lapeyre, et al. 1993; Gerbi 1996; Schnare, et al. 1996). ES's emerge from a small number of conserved sites on the common core and are excluded from regions of essential ribosomal function such as the DCC, the PTC, and the subunit interface (Ben-Shem, et al. 2010; Anger, et al. 2013). Expansion segments are larger and more numerous on the LSU than on the SSU; across phylogeny, size variation of the SSU rRNA is around 10% of that of LSU rRNA (Gutell 1992; Gerbi 1996; Bernier, et al. 2018).

ES's mutate quickly, at rates much greater than those of common core rRNA (Gonzalez, et al. 1985; Gonzalez, et al. 1988; Ajuh, et al. 1991; Gerbi 1996). For example, ES7 of *D. melanogaster* (fruit fly) is AU-rich and lacks sequence homology with the GC-rich ES7 of *A. albopictus* (Asian tiger mosquito). The high rate of mutation is indicated by the substantial sequence differences in species of the order Diptera that diverged only 200-250 million years ago (Aris-Brosou and Yang 2002; Kumar, et al. 2017). Although sequence conservation of ES's is low, secondary and tertiary structures are excellent sources of information on ES homology and evolution. For this reason, early studies used rRNA secondary structures to establish phylogenetic relationships (Woese, et al. 1990a).

The recent discovery and characterization of the Asgard archaeal superphylum suggests that the last Asgard and eukaryotic common ancestor (LAsECA) contained key components of eukaryotic cellular systems (Spang, et al. 2015; Klinger, et al. 2016; Eme, et al. 2017; Zaremba-

Niedzwiedzka, et al. 2017; Narrowe, et al. 2018; Spang, et al. 2019). Eukaryotic signature proteins (ESPs) found in Asgard archaea are involved in cytoskeleton, trafficking, ubiquitination, and translation (Spang, et al. 2015; Zaremba-Niedzwiedzka, et al. 2017 ; Akıl and Robinson 2018; Melnikov, et al. 2019). Asgard archaea also contain several homologs of eukaryotic ribosomal proteins (Hartman and Fedorov 2002; Spang, et al. 2015; Zaremba-Niedzwiedzka, et al. 2017). Metazoan rRNAs contain supersized ES's of hundreds of nucleotides (nts); hereafter, we define "supersized" ES's as those with >100 nts. Before our work here, it was not known if Asgard rRNAs could contain eukaryotic-like features such as supersized ES's. Eukaryotic-like, supersized ES's have not been observed previously in Bacteria or Archaea and were considered unique to eukaryotes (Ware, et al. 1983; Clark, et al. 1984; Hassouna, et al. 1984; Gerbi 1996; Melnikov, et al. 2012).

Here, we use computational and experimental approaches to characterize structure and evolution of Asgard rRNAs. We find that rRNAs of *Lokiarchaeota* and *Heimdallarchaeota*, both from the Asgard phyla, are elaborated on the LSU by supersized eukaryotic-like ES's. In size and complexity, *Lokiarchaeota* and *Heimdallarchaeota* LSU ES's exceed those of protists rRNAs and rival those of metazoan rRNAs. No es's were found in SSU rRNA of *Lokiarchaeota*. Our data suggest that the ES's of *Lokiarchaeota*, *Heimdallarchaeota*, and Eukarya can be traced back to a common ancestor (LAsECA). These results are consistent with Eukarya branching from within the Asgard superphylum.

Results

Comparative analysis reveals broad patterns of LSU rRNA size relationships. Previously, we developed the SEREB MSA (Sparse and Efficient Representation of Extant Biology, Multiple Sequence Alignment) for comparative analysis of sequences from the translation system (Bernier, et al. 2018). The SEREB MSA is alignments of a sparse and unbiased representation of all major phyla and is designed specifically to understand phenomena that are universal to all of biology. The MSA was manually curated and extensively cross-validated. The SEREB MSA is useful as a seed to study a variety of evolutionary phenomena. Recently, we augmented the SEREB MSA to include additional metazoan sequences, allowing us to characterize ES's and their evolution in metazoans (Mestre-Fos, et al. 2019a; Mestre-Fos,

et al. 2019b). Here, we augmented the SEREB MSA to include 21 sequences from the Asgard
superphylum and 12 sequences from deeply branching protists (**supplementary datasets S1,2**).

The SEREB MSA indicates that LSU rRNA size relationships are governed by the
general pattern: Bacteria (2725-2960 nts, n=61 [n is number of species]) < Archaea excluding
Lokiarchaeota (2886 to 3094 nts, n=48) < Eukarya excluding parasites (3300-5200 nts, n=41;
supplementary figure S1). The SEREB MSA demonstrates that archaeal rRNAs frequently
contain micro-expansion segments (μ-ES's) at positions of attachment of eukaryotic ES's. For
example, in Archaea, μ-ES39 is connected at the site of attachment of the large ES39 in
eukaryotes.

Lokiarchaeota are intermediate between Eukarya and Archaea in LSU rRNA size.
The Asgard augmentation of the SEREB MSA revealed unexpectedly large *Lokiarchaeota* LSU
rRNAs. *Lokiarchaeota* LSU rRNAs range from 3100 to 3300 nts (n=7). *Lokiarchaeota* rRNAs
are close to or within the observed size range of eukaryotic LSU rRNAs (**supplementary figure
S1**). Other Asgard species have LSU sizes in the upper archaeal range (from 3038 to 3142 nts,
n=6). Supersized archaeal ES's, which rival eukaryotic ESs, are observed in *Lokiarchaeota* and
Heimdallarchaeota spp. These ES's connect to the universal common core rRNA at the sites of
attachment of eukaryotic ES9 and ES39 and archaeal μ-ES9 and μ-ES39 (**Fig. 1**). Here we
explored the Asgard augmentation of the SEREB MSA to investigate the structure, distribution,
and evolution of rRNA expansions of Asgard archaea.

Figure 1: Secondary structures of the LSU rRNA from Bacteria, Archaea, and Eukarya. (A) *Escherichia coli* (Bacteria), (B) *Saccharomyces cerevisiae* (fungus, Eukarya); (C) *Pyrococcus furiosus* (Archaea); (D) *Drosophila melanogaster* (insect, Eukarya); (E) *Lokiarchaeota* F3 (Archaea); (F) *Homo sapiens* (primate, Eukarya). Secondary structures in panels A, B, C, D, and F are taken from *Petrov, et al. (2014a)*. Secondary structure in panel E is from this study. Universal rRNA common core is shown in blue lines (not shaded). ES9 is shown with a green line. ES39 is shown with a magenta line. Helices in panel E are abbreviated as Helix 1 (H1), helix 98 (H98), helix *a* (H*a*), helix *a1* (H*a1*), helix *a2* (H*a2*) and helix *b* (H*b*). ES's and helices not present in the common core are shaded in gray. Sizes of secondary structures are to scale. The numbering scheme of Noller, et al. (1981) and Leffers, et al. (1987) were used to label the helices and ES's.

ES9 and ES39 in some Asgard archaea are larger than μ-ES's of other archaea and ES's of protists. The MSA shows that ES39 in *Lokiarchaeota* ranges in size from 95 to 200 nts and in *Heimdallarchaeota* is from 115 to 132 nts. Other Asgard species have shorter ES39's: *Thorarchaeota*, *Odinarchaeota*, and *Freyarchaeota* ES39 sequences are between 50-60 nts. These Asgard ES39 sizes are well within the range of eukaryotic ES39, which is around 80 nts in a variety of protists, 138 nts in *Saccharomyces cerevisiae,* 178 nts in *Drosophila melanogaster,* and 231 nts in *Homo sapiens* (**Fig. 2, supplementary data S3**). For *Candidatus* Lokiarchaeota archaeon 1244-F3-H4-B5 (*Lokiarchaeota* B5), the primary focus of our work here, ES39 is 191 nts (**Figs. 2, 3**). Two of the *Lokiarchaeota* species have 95 nt ES39's (**Fig. 2, supplementary data S2**), slightly below our definition of a supersized ES, but still more than twice as long as the largest archaeal μ-ES39, *Pyrococcus furiosus* (45 nts, **Fig. 1C, supplementary dataset S3**). All Asgard have larger ES39 sequences than other archaea. Archaea from the *Euryarchaeota* phylum, except for the classes *Methanococci* and *Thermococci*, lack ES39 entirely (less than 5 nts) and μ-ES39 of species from other non-Asgard archaea varies between 0 and 45 nts (**supplementary dataset S3**).

Amongst Asgard archaea, only *Lokiarchaeota* spp. exhibit large ES9. *Lokiarchaeota* ES9 ranges from 29 to 103 nts (103 nts in *Lokiarchaeota* B5), compared to 29 nts in *S. cerevisiae,* 44 nts in *D. melanogaster* and *H. sapiens*, and 111 nts in *G. theta* (**supplementary figure S2**). In other non-Asgard archaea, ES9 varies between 20-30 nts. ES9 and ES39 are the primary contributors to the large size of *Lokiarchaeota* LSU rRNAs compared to the LSU rRNAs of other archaea. Outside of the Asgard superphylum*,* archaea lack supersized ES's.

Supersized ES's of *Lokiarchaeota* are transcribed *in situ*. To assess whether *Lokiarchaeota* ES's are transcribed, we assembled metatranscriptomic reads from sediment from the Gulf of Mexico, known to contain *Lokiarchaeota* sequences (Yergeau, et al. 2015; Cai, et al. 2018). We did not find metatranscriptomes that contain sequences from *Heimdallarchaeota*. Multiple transcripts from *Lokiarchaeota*-like LSU ribosomes contain ES9 and ES39 sequences, confirming that *Lokiarchaeota* ES's are indeed transcribed *in situ* (**Fig. 4D**; **supplementary dataset S3**).

Figure 2: Distribution of ES39 lengths from the three domains of life, including Asgard archaea. The number of nucleotides calculated between alignment positions 4891 and 5123 (*H. sapiens* numbering) of the LSU alignment for each species (**supplementary dataset S1**). The box shows the quartiles of the dataset. Whiskers extend to show the rest of the distribution, except for points that are determined to be outliers using a function of the inter-quartile range. Bacteria sequences are gray, *Lokiarchaeota* sequences are green, other Asgard sequences are blue, other archaeal sequences are purple, eukaryotic sequences are red, and sequences from metatranscriptomic contigs (**supplementary dataset S2**) for which there is no species determination are black. Abbreviations: *Ec: Escherichia coli*; *Pf*: *Pyrococcus furiosus*; *La B5: Lokiarchaeota B5*; *La GC: Lokiarchaeota GC14_75*; *Ha B18: Heimdallarchaeota B18G1*; *Ha RS: Heimdallarchaeota RS678*; *Ps: Prometheoarchaeum syntrophicum*; *Ta: Thorarchaeota*; *Oa*: *Odinarchaeota LCB4*; *Fa*: *Freyaarchaeota*; *Hs*: *Homo sapiens; Dm*: *Drosophila melanogaster*; *Sc*: *Saccharomyces cerevisiae*; *Tb*: *Trypanosoma brucei*; *Hi*: *Hexamita inflata*. *: parasitic representatives of the deeply branching eukaryotic group Excavata. †: Excluding the Asgard superphylum.

Asgard LSU rRNA contain the common core. We determined the extent of structural similarity of *Lokiarchaeota* LSU rRNAs with those of eukaryotes. We combined computation and experiment to model the secondary structure of *Lokiarchaeota* B5 LSU rRNA (**Fig. 1E; supplementary figure S3**). Secondary and three-dimensional structures of several eukaryotic and archaeal ribosomes provided a basis for modeling by homology. Like all other LSU rRNAs, *Lokiarchaeota* LSU rRNA contains the rRNA common core, which is trivial to model because the backbone atoms of the common core are highly conserved in all cytosolic ribosomes.

Figure 3: **Secondary structure model of** *Lokiarchaeota* **B5 ES39.** ES39 spans nucleotide positions 3006-3196 of the *Lokiarchaeota* B5 LSU rRNA. Canonical base-pairing positions are indicated with black lines. Helices are annotated with colored labels: blue – Helix 98, purple – Helix *a*, pink – Helix *a1*, orange – Helix *a2*, gold – Helix *b*. Figure was generated with RiboVision (Bernier, et al. 2014).

ES39 has a well-defined structural core in eukaryotes. Before comparing eukaryotic and archaeal ES39's, we characterized the conserved core of eukaryotic ES39, which we call the ES39 signature fold. We compared experimental three-dimensional structures of rRNAs in a wide range of ribosomes, from protists to primates (Ben-Shem, et al. 2010; Klinge, et al. 2011; Khatter, et al. 2015; Li, et al. 2017). The ES39 signature fold, which is common to all eukaryotic ES39s, consists of helix 98 (H98; 20-30 nts), helix *b* (H*b*; 40-50 nts), and their linkage by three unpaired 15 nts long segments of rRNA (**supplementary figure S4)**. These

unpaired segments are tightly associated with the ribosomal surface. In all eukaryotes, two of the unpaired segments interact with eukaryotic-specific α-helical extensions on rProteins uL13 and eL14 **(supplementary figure S6).** The third unpaired segment interacts with ES7 and rProtein aL33 **(supplementary figures S4, S7, S8)** (Khatter, et al. 2015). H*b* is terminated by a conserved GNRA tetraloop (**supplementary figures S4 and S5**). The ES39 signature fold is conserved in structure but not in sequence.

The ES39 signature fold can be decorated by an additional helix. Many eukaryotes possess a third helix (H*a*) that projects from the ES39 signature fold **(supplementary figure S4)**.

H*a* increases in size from simple unicellular eukaryotes, such as *Tetrahymena thermophila* (no

helix) and *Toxoplasma gondii* (10 nts), to more complex eukaryotes such as *S. cerevisiae* (18 nts)

and *Drosophila melanogaster* (20 nts). H*a* is largest in *Chordata* (106 nts for *H. sapiens*;

supplementary figure S4).

Initial *Lokiarchaeota* B5 ES39 secondary models were predicted by two methods. A

preliminary secondary structural model of ES39 of *Lokiarchaeota* B5 was generated using mfold

(Zuker 2003) (**Fig. 3**). The program mfold predicts a minimum free energy secondary structures

using experimental nearest-neighbor parameters. We selected the mfold model with lowest free

energy for further investigation. A preliminary secondary structural model assumed

Lokiarchaeota ES39 was structurally homologous with ES39 of *H. sapiens*. The mfold model

was confirmed to be correct, and the *H. sapiens* homology model was determined to be incorrect

by covariation analysis and Selective 2' Hydroxyl Acylation analyzed by Primer Extension

(SHAPE) reactivity data as described below.

Covariation supports the mfold model for the secondary structure of ES39 of

***Lokiarchaeota* B5.** Covariation, or coupled changes in paired nucleotides across phylogeny, can

help reveal RNA secondary structure (Levitt 1969; Ninio, et al. 1969; Woese, et al. 1980; Noller,

et al. 1981; Gutell, et al. 1993; Gutell, et al. 1994). Base-pairing relationships can be detected

through covariation. We compared co-variation predictions of each secondary model with

observed co-variation in the MSAs. The secondary structure predicted by mfold is consistent

with observed covariation **(Fig. 4)** while secondary structure predicted by *H. sapiens* homology

model is not. The observation of covarying nucleotides supports the model determined by mfold.

Figure 4: **Secondary structure of *Lokiarchaeota* B5 ES39 from experiment and computation.** (A) 1-D topology map of base pairs. The primary sequence of ES39 is on the horizontal line, Arcs indicate base pairs. Each helix is a distinct color. (B) SHAPE reactivity for ES39 mapped onto the secondary structure. Darker color indicates less flexible (paired) rRNA. (C) Base pairing covariation within the Asgard superphylum mapped on the secondary structure. Darker color indicates covarying (paired) rRNA across Asgard. Unpaired rRNA, for which no covariation data can be calculated, is gold. (D) SHAPE reactivity and base-pairing conservation mapped onto the ES39 MSA of Asgard sequences. *Lokiarchaeota* B5

numbering was used. The secondary structure is indicated with colored arrows bellow the alignment and
as colored background. SHAPE reactivity is indicated with a bar graph above the secondary structure
annotation, colors of the bars are consistent with panel B. Base-pairing conservation is indicated with a
bar graph bellow the secondary structure annotation; colors of the bars are consistent with panel C. Panel
D was generated with Jalview (Waterhouse, et al. 2009). Helices are labeled with colored text in each
panel; blue, helix 98; violet, helix *a*; pink, helix *a1*; orange, helix *a2*; yellow, helix *b*. Full sequence
names and sequencing project identifiers are available in **supplementary dataset S2**. Both SHAPE
reactivity and covariation are normalized. *: positions of conserved GNRA tetraloops.

Chemical footprinting confirms the mfold model for secondary structure of ES39 of

Lokiarchaeota **B5.** We further tested the secondary structural model of ES39 from

Lokiarchaeota B5 using SHAPE. This experimental method provides data on RNA flexibility at

nucleotide resolution (Merino, et al. 2005; Wilkinson, et al. 2006). SHAPE reactivity is generally

high for unpaired nucleotides, which are more flexible, and low for paired nucleotides, which are

less flexible. SHAPE has been widely used to probe the structure of rRNA (Leshin, et al. 2011;

Lavender, et al. 2015; Gomez Ramos, et al. 2017; Lenz, et al. 2017) and other RNAs (Wilkinson,

et al. 2005; Gilbert, et al. 2008; Stoddard, et al. 2008; Watts, et al. 2009; Novikova, et al. 2012;

Spitale, et al. 2013; Huang, et al. 2014). The SHAPE results from *Lokiarchaeota* B5 ES39 rRNA

(**supplementary figure S9**) are in agreement with the secondary structure based on mfold,

which, as shown above, is consistent with observed co-variation. Reactivity is low for paired

nucleotides in the mfold model and is high in loops and bulges (**Fig. 4B**). The accuracy of the

SHAPE data is supported by the observation of relatively high reactivity at the vast majority of

unpaired nucleotides and low reactivity for most paired nucleotides of the mfold model. The

Lokiarchaeota SHAPE data are not consistent with models that force *Lokiarchaeota* ES39 to

conform to the *H. sapiens* secondary structure.

Asgard ES39 deviates from the eukaryotic ES39 signature old. In eukaryotic ES39,

the junction of helices H98, H*a*, and H*b* contains three unpaired segments, each 15 nts. In

Lokiarchaeota B5, ES39 lacks unpaired segments greater than 8 nts and contains fewer unpaired

nucleotides (**Fig. 3**). *Lokiarchaeota* B5 ES39 is composed of four short helices, each up to 38 nts

(H98; Helix *a1*:H*a1*; Helix *a2*:H*a2*; and H*b*), and one long helix (H*a*: 72 nts). H98 and H*b*

connect in a three-way junction with H*a* at the base of ES39. H*a1* and H*a2* split H*a* at the top of

ES39 in a three-way junction (**Fig. 3**).

We modeled and visualized secondary structures of ES39 sequences from additional

Asgard species (**Fig. 5, supplementary figure S10**). None of these secondary structures

exhibited unpaired regions longer than 10 nts. ES39 of all Asgard archaea contain a three-way

junction that connects H98, H*a,* and H*b*, as observed in ES39 of *Lokiarchaeota* B5 (**Fig. 3**). Such three-way junctions can indicate locations at which accretions in the ribosomal structure took place (Petrov, et al. 2014b). Furthermore, ES39 in *Heimdallarchaeota* B18G1 had an additional branching of H*a* into H*a1* and H*a2,* mirroring the morphology of ES39 in *Lokiarchaeota* B5 (**Fig. 5, supplementary figure S10**). Despite the common branching morphology, the length of the individual helices varies substantially between different species (**Figs. 4D, 5**).

Figure 5: **ES39 secondary structures mapped on Asgard phylogeny**. Secondary structures of *Pyrococcus furiosus* (Archaea), *Lokiarchaeota* B5 (Asgard archaea), *Promethearchaeum syntropicum*

*(*Asgard archaea), *Thorarchaeota* (Asgard archaea), *Odinarchaeota LCB4* (Asgard archaea),
Heimdallarchaeota B18 G1 and *RS678* (Asgard archaea), *Saccharomyces cerevisiae* (Eukarya), and
Homo sapiens (Eukarya). Helix abbreviations are the same as in **Fig. 1E**. Helices 98 and *b* are highlighted
in orange. The phylogenetic tree topology is from Williams, et al. (2020). Ancestral clades on the
phylogenetic tree are labeled as LAECA: Last Archaeal and Eukaryotic Common Ancestor; LAsECA:
Last Asgardian and Eukaryotic Common Ancestor; LLaCA: Last Lokiarchaeal Common Ancestor;
LOTCA: Last Odin- and Thor-archaeota Common Ancestor; LHaCA: Last Heimdallarchaeota Common
Ancestor; LECA: Last Eukaryotic Common Ancestor. Possible ancestral structures are indicated next to
ancestral the clades. Minimal common structure from related extant species is shown with bold dashes.
The uncertainty in ancestral sizes of helix *a* are shown with light dashes. *: Occurrence of strand
dissociation in ES39 at LECA due to loss of helix 1 is indicated with a red gradient.

Asgard ES39 is located within an archaeal structural environment in the ribosome.
ES39 in Eukarya protrudes from helices 94 and 99 of the ribosomal common core **(Fig. 6)**. In
three dimensions, ES39 is close to ES7 and rProteins uL13, eL14, and aL33 (**supplementary**
figures S4, S6, S7). These elements in all Asgard species are more similar to Archaea than to
Eukarya. Additionally, ribosomes of both *Lokiarchaeota* and *Heimdallarchaeota*, like those of
all Archaea, contain helix 1 (H1), which is in direct contact with H98 at the base of ES39,
whereas eukaryotes lack H1. Combined with the eukaryotic-like size of *Lokiarchaeota* and
Heimdallarchaeota ES39, these characteristics predict that ribosomes from these two Asgard
groups have a unique structure in this region.

Figure 6: **Three dimensional structures of ES39 and its neighborhood from species representatives of the major life domains and Asgard archaea.** (A) *Escherichia coli* (Bacteria), (B) *Pyrococcus furiosus* (Archaea), (C) Asgard archaea; (D) *Homo sapiens* (Eukarya). Helices are abbreviated as helix 1 (H1), helix 94 (H94), helix 98 (H98), helix 99 (H99) and helix *b* (H*b*). Helix 98 is orange, helix 1 is green, helices 94 and 99 are blue. The *P. furiosus* structure is used as model for the Asgard structures. Likely structure of Asgard helices 98 and *b* are shown with transparency. Likely position and direction of the Asgard ES39 continuation is indicated with a black dashed line. Direction of eukaryotic ES39 continuation is indicated with a black dashed line.

The pathway of ES39 evolution appears to be unique. In general, ES's have increased in size over evolution via accretion. Growth by insertion of one rRNA helix into another leaves a structural mark. These growth processes are indicated by the joining of a rRNA helical trunk to an rRNA helical branch at a highly localized three- or four-way junction (an insertion fingerprint) (Petrov, et al. 2014b; Petrov, et al. 2015). Basal structure is preserved when new rRNA is acquired. For instance, ES7 shows continuous growth over phylogeny, expanding from LUCA to Archaea to protists to metazoans to mammals (Petrov, et al. 2014b; Bernier, et al. 2018). The accretion model predicts that H98, at the base of ES39, would superimpose in

Bacteria, Archaea, and Eukarya, but in fact H98 does not overlap in superimposed 3D structures (**Fig. 6**). The archaeon *P. furiosus* has a slightly extended and bent H98 compared to the bacterium *E. coli* (**Fig. 6**). This spatial divergence is likely due to the difference in how *E. coli* H98 and *P. furiosus* H98 interact with H1 of the LSU. *E. coli* H98 interacts within the H1 minor groove through an A-minor interaction, while in *P. furiosus* H98 is positioned on top of H1 (**Fig. 6**). H1 is absent in eukaryotes (**Fig. 1**), allowing H98 to occupy the position of H1 (**Fig. 6**) and causing a strand dissociation in the eukaryotic ES39 structure.

Structurally, *Lokiarchaeota* and *Heimdallarchaeota* ES39 may extend in a different direction than eukaryotic ES39. *Lokiarchaeota* and *Heimdallarchaeota* spp. have larger ES39 than other archaea (**Fig. 2**) and possess H1, unlike Eukarya (**Fig. 6**). We predict that *Lokiarchaeota* and *Heimdallarchaeota* ES39 have an archaeal-like interaction with H1 through H98 and H*b* (**Fig. 6**). In Asgard species, the highly variable H*a* (**Figs. 4D, 5**) likely grows out from the three-way junction between H98 and H*b*, perpendicular to the eukaryotic H*a* (**Fig. 6**). While H*a* of eukaryotic ES39 is pointed in the direction of the sarcin-ricin loop, the large H*a* of *Lokiarchaeota* and *Heimdallarchaeota* is likely pointed in the direction of the central protuberance or the exit tunnel. The difference in directionality between Asgard and eukaryotic ES39 arises from the loss of H1 in Eukarya, where H98 and H*b* partially occupy the vacated space.

Weak sequence homology between ES39 sequences. We searched for ES39 homology between Asgards and deeply branching protists. MSAs between Asgard (n=16) and eukaryotic (n=15) ES39 sequences were compared with eukaryotic-only and Asgard-only MSAs. The percent identities in the eukaryotic-only ES39 MSA varied from 6 to 90% (**supplementary figure S11**). As expected, more closely related eukaryotes had greater identities (40-90% within *Chordata*; **supplementary figure S11**). Common core rRNA, with up to 80% identity even between distantly related species, was much more conserved than ES rRNA, (**supplementary figure S12**). Within the *Lokiarchaeota*, ES39 sequences shared 40-60% identity (**supplementary figure S13**). The extent of similarity between Asgard and eukaryotic ES39 sequences (10-30% identities; **supplementary figure S14**) was similar to the extent of similarity between distantly related eukaryotes (**supplementary figure S11**).

Conservation of a GNRA-tetraloop capping ES39 H*b* between Eukarya and Asgard. The MSA of H*b* from ES39 shows a GNRA tetraloop that is conserved in both eukaryotes and

Asgard archaea. Three sequences from protists exhibit sequence similarity with H*b* sequences

from Asgard archaea (**supplementary figure S5**).

Discussion

The recent discovery of the archaeal Asgard superphylum, which contain multitudes of ESPs,

has redefined our understanding of eukaryotic evolution (Spang, et al. 2015; Zaremba-

Niedzwiedzka, et al. 2017). Incorporation of Asgard species into phylogenies has changed our

perspective on the relationship of Archaea and Eukarya in the tree of life (Hug, et al. 2016;

Fournier and Poole 2018; Williams, et al. 2020). Here, we incorporate rRNA secondary structure

and three-dimensional interactions into comparative analyses. Our work extends structure-based

methods of comparative analysis to *Lokiarchaeota* and *Heimdallarchaeota* rRNAs, and

mechanistic models for the evolution of common rRNA features in Eukarya and some Asgard

archaea. We identified profound structural similarities of rRNAs, indicating the possibility of

functional parallels between eukaryotes and Asgard archaea.

Lokiarchaeota **and** *Heimdallarchaeota* **rRNAs have unique eukaryotic-like features.**

The ribosome is a window to relationships among organisms (Woese and Fox 1977; Hillis and

Dixon 1991; Olsen and Woese 1993; Fournier and Gogarten 2010; Hug, et al. 2016) and was a

driver of ancient evolutionary processes (Agmon, et al. 2005; Smith, et al. 2008; Bokov and

Steinberg 2009; Fox 2010; Petrov, et al. 2015; Melnikov, et al. 2018). Previous work has

revealed rules of ribosomal variation over phylogeny (Hassouna, et al. 1984; Gerbi 1996;

Melnikov, et al. 2012; Bernier, et al. 2018), as well as mechanisms of ribosomal change over

evolutionary history (Petrov, et al. 2014b; Petrov, et al. 2015; Kovacs, et al. 2017; Melnikov, et

al. 2018). Building off those studies, we assessed the extent to which *Lokiarchaeota* and

Heimdallarchaeota ribosomes follow or deviate from previously established rules of ribosomal

sequence and structure variation (Bowman, et al. 2020). We found that *Lokiarchaeota* and

Heimdallarchaeota ribosomes:

• contain the universal common core of rRNA and rProteins (this work) (Spang, et al.

2015; Bernier, et al. 2018),

• confine rRNA variability of structure and size to ES's/μ-ES's (this work),

- restrict ES's to universally conserved sites on the common core (Ware, et al. 1983; Clark, et al. 1984; Hassouna, et al. 1984; Michot and Bachellerie 1987; Bachellerie and Michot 1989; Lapeyre, et al. 1993; Gerbi 1996),
- avoid ES attachment from the ribosomal interior or near functional centers (Ben-Shem, et al. 2010; Anger, et al. 2013), and
- concentrate variability in structure and size on LSU rRNA, not SSU rRNA (Gerbi 1996; Bernier, et al. 2018).

The ribosomes of *Lokiarchaeota*, but not of *Heimdallarchaeota,* violate the established pattern of increasing rRNA length in the order: Bacteria < Archaea ≪ protists ≪ metazoans (Melnikov, et al. 2012; Petrov, et al. 2014b; Bernier, et al. 2018). *Lokiarchaeota* LSU rRNAs are much longer than predicted for archaea; in fact, *Lokiarchaeota* rRNA eclipses the length of rRNA in many protists. *Lokiarchaeota* ES39 is larger than ES39 in protists and some metazoans. ES9 and ES39 in *Lokiarchaeota* are larger than in any other archaea known to date.

Two scenarios for the evolution of ES39. Two exclusive scenarios explain the similarities of ES39 in Eukarya and Asgard archaea. One scenario is based on a three-domain tree of life and predicts parallel evolution of ES39, while the other assumes close common ancestry of Asgard archaea and Eukarya. Our results do not exclude either one completely; therefore, we will discuss both.

Parallel evolution of ES39 between Asgard archaea and Eukarya. The difference in directionality of H*a* in Asgard and Eukarya combined with the high evolutionary rates in ES's, and the lack of structural or sequence similarity, implies parallel evolution of ES39 with differential functions. The parallel growth of LSU ES's in Asgard and Eukarya can be the result of similar selection pressures. One explanation of the similar ES sizes between *Lokiarchaeota*, *Heimdallarchaeota,* and higher eukaryotes (e.g. *Chordata*) may be slow growth rates and small population sizes, which are associated with greater number of benign mutations (Lynch 2007; Rajon and Masel 2011). Both *Lokiarchaeota* and *Chordata* have low growth rates and small populations (Imachi, et al. 2020), that might lead to benign increase of rRNA in ES ribosomal regions.

Common ancestry of ES39 in Asgard and Eukarya. Supersized ES's were thought to be unique to eukaryotic ribosomes. We now know that to be incorrect: two Asgard phyla exhibit supersized ES's. ES sequences are highly variable, even among closely related species

(**supplementary figure S11**), making ES's impractical for sequence-based determination of

ancient ancestry. Secondary and three dimensional structures are more conserved than

sequences, allowing a more accurate detection of distant ancestry between species (Woese, et al.

1990a). By integrating the accretion model of ribosomal evolution (Petrov, et al. 2014b; Petrov,

et al. 2015) and structure-informed phylogeny, we reveal ancestries of Eukarya and Archaea.

The accretion model suggests ES39 in LAsECA contains three helices. An important

structural difference between Asgard and other archaea is the three-helical topology of ES39. All

Asgard have three helices in ES39, while other Archaea either lack ES39 altogether or contain a

short-bent helix consisting of H98 and H*b* (**Figs. 2,5,6**). The accretion model predicts that ES39

of the Last Archaeal and Eukaryotic Common Ancestor (LAECA) contained H98 and H*b* (**Fig.**

5); they comprise the minimal form of ES39 shared between Asgard and other archaea (**Figs. 4**

and **5**). The insertion of H*a* between H98 and H*b* at LAsECA is consistent with the minimal

ES39 (**Fig. 5**). Subsequently, the three-helical topology would be inherited by all Asgard species,

consistent with variable H*a* sizes (**Figs. 4** and **5**). Similarly, eukaryotes have highly conserved

H98 and H*b* while H*a* is variable (**supplementary figures S4 and S5**). Eukarya differ from

Asgard archaea in the unpaired rRNA segments that connect H98, H*b*, and H*a* (**supplementary**

figure S4). The absence of insertion fingerprints means that we lack a detailed model for the

relative ages of eukaryotic H98, H*a*, and H*b*. It is reasonable to assume that LECA had a three-

helical ES39 inherited from one of its Asgardian sister clades. This ancient three-helical ES39

underwent restructuring and strand dissociation upon the loss of H1 (**Figs. 5, 6**). Subsequently,

the unpaired structure of ES39 was inherited by all eukaryotes (**Fig. 5**).

Implications for eukaryogenesis. The mechanisms of transition from prokaryotes to

eukaryotes hold some of the most exciting and fundamental questions about evolution of cellular

complexity. With the discovery of Asgard archaea, the study of eukaryogenesis transitioned into

finding eukaryotic features in archaeal species. This study contributes to a growing body of

eukaryotic specific features in Asgard archaea, specifically supersized ES39 and ES9 in

Lokiarchaeota and *Heimdallarchaeota*. We suggest that ES's are archaeal features that are

elaborated in Eukarya, and Asgard. The precise pathway of evolution for ES's is hard to

ascertain because ES sequences undergo rapid evolution. We can, however, make predictions

about ES functions, given their general location on the ribosomal surface and the known

functions of eukaryotic ES's. ES39 is located on the ribosomal surface between the sarcin-ricin

loop, the end of the exit tunnel and the central protuberance, and has been shown in eukaryotes
to bind factors related to N-terminal acetylation of nascent peptides (Knorr, et al. 2018). Among
a plethora of functions, N-terminal acetylation takes part in protein protection of degradation
from N-terminal ubiquitination (Hershko, et al. 1984). *Lokiarchaeota*, *Heimdallarchaeota*, and
Odinarchaeota contain the necessary enzymes for protein ubiquitination, an eukaryotic hallmark
(Zaremba-Niedzwiedzka, et al. 2017). Except for *Odinarchaeota* there is a correlation between
Asgard phyla that contain large ES39 and Asgard phyla that contain proteins necessary for
ubiquitination. It is possible that Asgard and eukaryotic ES39 grew in response to the increased
rates of ubiquitination. By binding and guiding the N-terminal acetylation system to the exit
tunnel, ES39 can protect nascent peptides from ubiquitination. Increased importance of protein
quality control is logical for organisms with low rates of growth and small population sizes such
as *Chordata* and *Lokiarchaeota* (Imachi, et al. 2020).

The specific roles of μ-ES's and ES's over phylogeny are unknown but are likely
complex, polymorphic, and pleotropic. The observation of μ-ES's in Archaea, ES's in Eukarya,
and supersized ES's in Asgard species suggest that (i) the roots of ribosomal complexity are
inextricably intertwined with the roots of Eukarya, and (ii) the roots of Eukarya penetrate more
deeply into the archaeal domain than previously recognized, conclusions that are consistent with
recent phylogenetic results that root Eukarya within the archaeal branch (Williams, et al. 2020).

Conclusions

Lokiarchaeota and *Heimdallarchaeota* ribosomes contain supersized ES39's with structures that
are distinct from eukaryotic ES39's. *Lokiarchaeota* ES9's are larger than eukaryotic ES9's. To
date, *Lokiarchaeota* and *Heimdallarchaeota* are the only prokaryotic phyla with supersized ES's,
bringing the size of their LSUs close to those of Eukarya. *Lokiarchaeota* and *Heimdallarchaeota*
ES39 likely grow outward from the ribosomal surface in a different direction than eukaryotic
ES39's. Our findings raise the possibility that eukaryotic-sized ES's existed on the ribosomal
surface before LECA, suggesting that ribosomal complexity is more deeply rooted than
previously thought. Our data, based on RNA structure, provide support for primary sequence-
based studies that revealed a two-domain tree with Eukarya rooted within the Asgard
superphylum.

Materials and Methods

Genome sequencing, assembly, and binning

      Sample collection. Sediments were cored from deep seafloor sediment at ODP site 1244 (44°35.1784´N; 125°7.1902´W; 895 m water depth) on the eastern flank of Hydrate Ridge ~3 km northeast of the southern summit on ODP Leg 204 in 2002 (Tréhu, et al. 2003) and stored at -80°C at the IODP Gulf Coast Repository.

      DNA extraction. DNA was extracted from sediment from F3-H4 (18.10 meters below the seafloor) using a MO-BIO PowerSoil total RNA Isolation Kit with the DNA Elution Accessory Kit, following the manufacturer protocol without beads. Approximately 2 grams of sediments were used for the extraction from six extractions (12 g total) and DNA pellets from the two replicates from each depth were pooled together. DNA concentrations were measured using a Qubit 2.0 fluorometer with dsDNA High Sensitivity reagents (Invitrogen, Grand Island, NY, USA). DNA yield was 7.5 ng per gram of sediment.

      Multiple displacement amplification, library preparation, and sequencing. Genomic DNA was amplified using a REPLI-g Single Cell Kit (Qiagen, Germantown, MD, USA) using UV-treated sterile plasticware and reverse transcription-PCR grade water (Ambion, Grand Island, NY, USA). Quantitative PCR showed that the negative control began amplifying after 5 hr of incubation at 30°C, and therefore, the 30°C incubation step was shortened to 5 hr using a Bio-Rad C1000 Touch thermal cycler (Bio-Rad, Hercules, CA, USA). DNA concentrations were measured by Qubit. Two micrograms of MDA-amplified DNA were used to generate genome libraries using a TruSeq DNA PCR-Free Kit following the manufacturer's protocol (Illumina, San Diego, CA, USA). The resulting libraries were sequenced using a Rapid-Run on an Illumina HiSeq 2500 to obtain 100 bp paired-end reads. Metagenomic sequences were deposited into NCBI as accession numbers SAMN07256342-07256348 (BioProject PRJNA390944).

      Metagenome assembly, binning, and annotation. Demultiplexed Illumina reads were mapped to known adapters using Bowtie2 in local mode to remove any reads with adapter contamination. Demultiplexed Illumina read pairs were quality trimmed with Trim Galore (Martin 2011) using a base Phred33 score threshold of Q25 and a minimum length cutoff of 80 bp. Paired-end reads were then assembled into contigs using SPAdes assembler (Bankevich, et al. 2012) with --meta option for assembling metagenomes, iterating over a range of k-mer values

(21,27,33,37,43,47,51,55,61,65,71,75,81,85,91,95). Assemblies were assessed with reports

generated with QUAST (Gurevich, et al. 2013). Features on contigs were predicted through the

Prokka pipeline with RNAmmer for rRNA, Aragorn for tRNA, Infernal and Rfam for other non-

coding RNA and Prodigal for protein coding genes (Laslett and Canback 2004; Lagesen, et al.

2007; Hyatt, et al. 2010; Nawrocki and Eddy 2013; Seemann 2014; Kalvari, et al. 2017). Each

genomic bin was searched manually for 23S rRNA sequences. The *Lokiarchaeota* F3-H4-B5 bin

(estimated 2.8% completeness and 0% contamination) was found to contain a 3300 nt 23S rRNA

sequence. The *Lokiarchaeota* F3-H4-B5 bin was deposited into NCBI as BioSample

SAMN13223206 and GenBank genome accession number WNEK00000000.

Environmental 23S rRNA transcript assembly and bioinformatic analysis

Species selection for analysis. To efficiently sample the tree of life, we used species

from the SEREB database (Bernier, et al. 2018) as a starting set. These species were selected to

sparsely sample all major phyla of the tree of life where allowed by available LSU rRNA

sequences. We further added LSU sequences from species of recently discovered phyla in the

archaeal and eukaryotic domains (**supplementary dataset S2**). These additional species were

again selected as representatives from major phyla. To gain higher level of evolutionary

resolution in the Asgard superphylum we added complete LSU sequences from several available

Asgard species and constructed assemblies from available meta-transcriptomes. All species used

in our analysis with their respective phylogenetic groups are listed in **supplementary dataset**

S3.

Assembly. Publicly available environmental meta-transcriptomic reads were downloaded

from NCBI BioProject PRJNA288120 (Yergeau, et al. 2015). Quality evaluation of the reads

was performed with FastQC (Andrews 2012) and trimming was done with TrimGalore (Martin

2011). Assembly of SRR5992925 was done using the SPADES (Bankevich, et al. 2012)

assembler with --meta and --rna options, to evaluate which performs better. Basic statistic

measures such as Nx, contig/transcript coverage and length were compared (**supplementary**

datasets S4, S5) yielding better results for the rnaspades assembler. All subsequent meta-

transcriptomic datasets were assembled with rnaspades.

Identifying ribosomal RNA sequences. BLAST databases were constructed (Altschul,

et al. 1990) from the resulting contig files and they were queried for ribosomal regions

characteristic of the Asgardian clade (ES39/ES9 sequences from GC14_75). Additionally, the program quast (Gurevich, et al. 2013) with --rna-finding option was used.

SEREB Multiple Sequence Alignment (MSA) augmentation. High scoring transcripts, as well as genomic sequences from added species, were included in the SEREB MSA (Bernier, et al. 2018) using the program mafft (Katoh and Standley 2013) with the --add option. Known intronic regions (Cannone, et al. 2002) were removed from new sequences. The highly variable region of ES39 was manually aligned using structural predictions from mfold (Zuker 2003).

Defining ES regions within Asgard archaea. To compare eukaryotic, archaeal, and Asgard ES regions we defined the basal helices of ES growth for ES9 and ES39. We calculated ES sizes by including all nucleotides within the start and end of the basal helices. In eukaryotes ES9 is a region of a cumulative expansion between helices 29, 30, and 31 (see region D3 in Hassouna, et al. (1984)). For ES9 we used helix 30, since the growth in Lokiarchaeota was contained within that helix. For ES39 we used H98, consistent with previous observations of the variability within that region (Gerbi 1996).

Sequence homology between Asgard and eukaryotic ES39 sequences. To uncover any sequence conservation within the ES39 region of the LSU between Asgard archaea and eukaryotes, we used the augmented SEREB MSA. The ES39 sequences for all eukaryotes and Asgard archaea were extracted from the global LSU alignment. Chordate eukaryotic sequences were removed as they were too GC rich, parasitic eukaryotic sequences were also removed. Multiple different automated alignment methods were used (Do, et al. 2005; Sievers, et al. 2011; Katoh and Standley 2013). Following the automated methods, we applied manual alignment guided by secondary structure predictions. Pairwise percent identities for the alignments where calculated with Ident and Sim from the Sequence Manipulation Suite (Stothard 2000).

LSU size comparison. The LSU size comparison was based on the transcribed gene for the LSU, which is comprised of a single uninterrupted rRNA sequence for bacteria and archaea (**Fig. 1A,C,E**), and is comprised of multiple concatenated rRNA sequences for the fragmented eukaryotic rRNA gene (**Fig. 1B,D,F**). The 5S rRNA, which is essentially constant, is excluded from the size calculation. The comparison takes into account whether the rRNAs are from endosymbionts and pathogens, which tend to contain reduced genomes, metabolisms, and translation systems (Peyretaillade, et al. 1998; Moran 2002; McCutcheon and Moran 2011).

Secondary structure models. To model the secondary structure of *Candidatus*
Lokiarchaeota archaeon 1244-F3-H4-B5 LSU rRNA, we used the secondary structure of *P.*
furiosus (Petrov, et al. 2014a) and a multiple sequence alignment (MSA) of archaeal LSU rRNAs
broadly sampled over the phylogeny (**supplementary dataset S1**). Locations of expansion
segments were unambiguously identified from the MSA. Due to the low percent identity (<50%)
(Bernhart and Hofacker 2009) we applied *ab initio* modelling for ES regions. GNRA tetraloops
are highly conserved structural motifs in rRNA structure (Woese, et al. 1990b; Heus and Pardi
1991; Nissen, et al. 2001). Conserved GNRA tetraloop sequences from the Asgard sequences
were used to determine the limits of variable helices in ES regions. The secondary structures of
the ES's were predicted by mfold (Zuker 2003).

Covariation. To verify the secondary structures of the highly variable ES regions base-
pairing conservation was calculated with the program Jalview (Waterhouse, et al. 2009). Gaps
from the MSA were ignored in the calculation to produce comparable results about available
regions. The base-pairing model of secondary structures of ES9 (**supplementary figure S15**)
and ES39 (**Figs. 4C,D**) was generated in the Jalview annotation format and used for the base-
pairing conservation calculation.

Defining the eukaryotic ES39 signature old. To identify the structurally invariant part
of ES39 in eukaryotes, we used superimposition based on the common core within domain VI of
the ribosomal structures from 4 eukaryotes (*Tetrahymena thermophila*, *Toxoplasma gondii*,
Saccharomyces cerevisiae, *Homo sapiens*; **supplementary figure S4**). Initially the *Drosophila*
melanogaster ribosomal structure (PDB ID: 4V6W) was used in identifying the core. However,
as it has additional loops elongating the unpaired regions, we excluded it from our analysis.
Drosophila melanogaster is known to have AU-enriched ES's; therefore, it is not surprising that
it has perturbations in its ES39.

ES39 rRNA SHAPE analysis

Synthesis of *Lokiarchaeota* ES39 rRNA. pUC57 constructs containing T7 promoter and
the gene encoding *Lokiarchaeota* ES39 rRNA was linearized using HindIII restriction enzyme.
Lokiarchaeota ES39 rRNA was synthesized by *in vitro* transcription using HiScribe™ T7 High
Yield RNA Synthesis Kit; New England Biolabs. RNA was then precipitated in
ethanol/ammonium acetate and purified by G25 size exclusion chromatography

(illustraTMNAPTM-10, GE Healthcare). RNA purity was assayed by denaturing gel
electrophoresis.

SHAPE reaction. Selective 2'-hydroxyl acylation analyzed by primer extension
(SHAPE; Wilkinson, et al. 2006) was performed to chemically probe local nucleotide flexibility
in ES39 rRNA. *In vitro*-transcribed ES39 rRNA was added to folding buffer (180 mM NaOAc,
50mM Na-HEPES (pH 8.0) and 1 mM 1,2- diaminocyclohexanetetraacetic acid (DCTA)) to
obtain 400 nM RNA in total volume of 80 μL. RNA was annealed by cooling from 75 °C to
25 °C at 1 °C/min. RNA modification reaction was performed with final concentration of 100
mM benzoyl cyanide (Sigma) prepared in dimethyl sulfoxide (DMSO). Non-modified RNA
samples were incubated with DMSO. Reactions were carried out for 2 min at room temperature.
Modified RNAs and control sample were purified by precipitation in ethanol and ammonium
acetate at 20 °C for 2 hr. RNA samples were centrifuged at 4°C for 10 min. The RNA pellets
were washed with 100 μL of 80% ethanol for two times and dried out using a SpeedVac vacuum
concentrator. TE buffer [22 μL of 1 mM EDTA and 10 mM Tris-Cl (pH 8.0)] was added to each
sample to resuspend the pellet.

Reverse transcription. Reverse transcription was conducted on 20 μL of modified RNAs
and unmodified RNA sample as a control, in presence of 8 pmol 5'[6-FAM] labeled primer (5'-
GAACCGGACCGAAGCCCG-3'), 2 mM DTT, 625 μM of each deoxynucleotidetriphosphate
(dNTP), and 5 μL of reverse transcription (RT) 5X first-strand buffer [250 mM Tris-HCl (pH
8.3), 375 mM KCl, 15 mM $MgCl_2$]. To anneal the primer, samples were heated at 95°C for 30
secs, held at 65°C for 3 min, and then 4°C for 10 min. RT mixtures were incubated at 52°C for
2 min before addition of 1 μL (200 U) of Superscript III Reverse transcriptase (Invitrogen) and
reactions were incubated at 55°C for 2 hr. later, RT terminated by heating samples at 70°C for 15
min. Chain termination sequencing reaction was performed on 10 pmol unmodified RNA
prepared in TE buffer, 8 pmol 5'[6-FAM] labeled primer, with a ratio of 1:1 dideoxynucleotide
(ddNTP) to dNTP. A sequencing reaction was performed with the same condition without
ddNTPs.

Capillary electrophoresis of RT reaction products and data analysis. Capillary
electrophoresis of RT reactions was performed as described previously (Hsiao, et al. 2013). For
each reaction 0.6 μL DNA size standard (Geneflo™ 625), 17.4 μl Hi-Di Formamide (Applied
Biosystems), and 2 μL of RT reaction mixture were loaded in a 96-well plate. Samples were

heated at 95°C for 5 min before electrophoresis and the RT products were resolved using applied

biosystems. SHAPE data were processed using a Matlab scripts as described previously

(Athavale, et al. 2012). SHAPE profile was mapped onto ES39 rRNA secondary structure with

the RiboVision program (Bernier, et al. 2014).

Acknowledgments: We thank Cecilia Kretz and Piyush Ranjan for sample preparation and

genome binning. We thank Brett Baker for providing an unpublished *Heimdallarchaeota*

sequence. We thank Jessica Bowman, Santi Mestre-Fos, Aaron Engelhart, Anthony Poole, and

Betül Kaçar for helpful conversations.

Funding: This research was supported by National Aeronautics and Space Administration grants

NNX16AJ29G, the Center for the Origin of Life grant 80NSSC18K1139, and a Center for Dark

Energy Biosphere Investigations (C-DEBI) Small Research Grant. This is C-DEBI contribution

[will be provided upon paper acceptance]. This research was supported in part through research

cyberinfrastructure resources and services provided by the Partnership for an Advanced

Computing Environment (PACE) at the Georgia Institute of Technology, Atlanta, Georgia, USA.

Author contributions: ASP, LDW, and JBG conceptualized the research. PIP curated the data

and performed computational analysis. ASP, LDW, and JBG acquired the funding and

administered the project. SF-A performed the SHAPE experiments. JJC and VJP performed

analyses. ASP, LDW, RRG, and JBG supervised the research. RRG validated the research. PIP

and VJP prepared the figures. PIP, JBG, and LDW wrote the manuscript with input from all

authors.

Competing interests: Authors declare no competing interests.

Data and materials availability: Metagenomic sequences were deposited into NCBI as

accession numbers SAMN07256342-07256348 (BioProject PRJNA390944). The *Lokiarchaeota*

F3-H4-B5 bin was deposited into NCBI as BioSample SAMN13223206 and GenBank genome

accession number WNEK00000000. The *Lokiarchaeota* F3-H4-B5 23S rRNA gene is in the

reverse complement of contig WNEK01000002.1, nucleotide positions 251-3579. Scripts used

for generation of data figures and tables, as well as MSAs of the ES39 region are available at

https://github.gatech.edu/LDWLab/Asgard_LSU_project.

References:

References:

Agmon I, Bashan A, Zarivach R, Yonath A. 2005. Symmetry at the active site of the ribosome:
structural and functional implications. In. Biological Chemistry. p. 833.

Ajuh PM, Heeney PA, Maden BE. 1991. Xenopus borealis and Xenopus laevis 28S ribosomal
DNA and the complete 40S ribosomal precursor RNA coding units of both species. Proc Biol Sci
245:65-71.

Akıl C, Robinson RC. 2018. Genomes of Asgard archaea encode profilins that regulate actin.
Nature 562:439-443.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.
J Mol Biol. 215:403-410.

Andrews S. 2012. FastQC: a quality control tool for high throughput sequence data. Available at
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Anger AM, Armache JP, Berninghausen O, Habeck M, Subklewe M, Wilson DN, Beckmann R.
2013. Structures of the human and Drosophila 80S ribosome. Nature 497:80-85.

Aris-Brosou S, Yang Z. 2002. Effects of Models of Rate Evolution on Estimation of Divergence
Dates with Special Reference to the Metazoan 18S Ribosomal RNA Phylogeny. Syst Biol
51:703-714.

Athavale SS, Gossett JJ, Hsiao C, Bowman JC, O'Neill E, Hershkovitz E, Preeprem T, Hud NV,
Wartell RM, Harvey SC, et al. 2012. Domain III of the T. thermophilus 23S rRNA folds
independently to a near-native state. RNA 18:752-758.

Bachellerie JP, Michot B. 1989. Evolution of large subunit rRNA structure. The 3' terminal
domain contains elements of secondary structure specific to major phylogenetic groups.
Biochimie 71:701-709.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko
SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing. Journal of computational biology : a journal of
computational molecular cell biology 19:455-477.

Ben-Shem A, Jenner L, Yusupova G, Yusupov M. 2010. Crystal structure of the eukaryotic
ribosome. Science 330:1203-1209.

Bernhart SH, Hofacker IL. 2009. From consensus structure prediction to RNA gene finding.
Briefings in Functional Genomics 8:461-471.

Bernier CR, Petrov AS, Kovacs NA, Penev PI, Williams LD. 2018. Translation: The Universal
Structural Core of Life. Mol Biol Evol. 35:2065-2076.

Bernier CR, Petrov AS, Waterbury CC, Jett J, Li F, Freil LE, Xiong X, Wang L, Migliozzi BL,
Hershkovits E, et al. 2014. RiboVision suite for visualization and analysis of ribosomes. Faraday
Discuss 169:195-207.

Bokov K, Steinberg SV. 2009. A hierarchical model for evolution of 23S ribosomal RNA.
Nature 457:977-980.

Bowman JC, Petrov AI, Frenkel-Pinter M, Penev PI, Williams LD. 2020. Root of the Tree: The
Significance, Evolution, and Origins of the Ribosome. Chemical Reviews.

Cai M, Liu Y, Zhou Z, Yang Y, Pan J, Gu J-D, Li M. 2018. Asgard archaea are diverse,
ubiquitous, and transcriptionally active microbes. bioRxiv:374165.

Cannone JJ, Subramanian S, Schnare MN, Collett JR, D'Souza LM, Du Y, Feng B, Lin N,
Madabusi LV, Muller KM, et al. 2002. The comparative RNA web (CRW) site: an online
database of comparative sequence and structure information for ribosomal, intron, and other
RNAs. BMC Bioinformatics 3:2.

Clark CG, Tague BW, Ware VC, Gerbi SA. 1984. Xenopus laevis 28S ribosomal RNA: a
secondary structure model and its evolutionary and functional implications. Nucleic Acids Res.
12:6197-6220.

Do CB, Mahabhashyam MS, Brudno M, Batzoglou S. 2005. ProbCons: Probabilistic
consistency-based multiple sequence alignment. Genome Res 15:330-340.

Eme L, Spang A, Lombard J, Stairs CW, Ettema TJG. 2017. Archaea and the origin of
eukaryotes. Nature Rev Microbiol. 15:711-723.

Fournier GP, Gogarten JP. 2010. Rooting the Ribosomal Tree of Life. Mol Biol Evol. 27:1792-
1801.

Fournier GP, Poole AM. 2018. A Briefly Argued Case That Asgard Archaea Are Part of the
Eukaryote Tree. Frontiers in Microbiology 9.

Fox GE. 2010. Origin and evolution of the ribosome. Cold Spring Harb Perspect Biol.
2:a003483.

Gerbi S. 1996. Expansion segments: regions of variable size that interrupt the universal core
secondary structure of ribosomal RNA. In: Ribosomal RNA: Structure, Evolution, Processing
and Function in Protein Synthesis. eds: Zimmermann RA, Dahlberg AE. Boca Raton, FL Telford
- CRC Press. 71. p. 87.

Gilbert SD, Rambo RP, Van Tyne D, Batey RT. 2008. Structure of the SAM-II riboswitch bound
to S-adenosylmethionine. Nat Struct Mol Biol. 15:177-182.

Gomez Ramos LM, Degtyareva NN, Kovacs NA, Holguin SY, Jiang L, Petrov AS, Biesiada M,
Hu MY, Purzycka KJ, Arya DP, et al. 2017. Eukaryotic Ribosomal Expansion Segments as
Antimicrobial Targets. Biochemistry 56:5288-5299.

Gonzalez IL, Gorski JL, Campen TJ, Dorney DJ, Erickson JM, Sylvester JE, Schmickel RD.
1985. Variation among human 28S ribosomal RNA genes. Proc Natl Acad Sci USA. 82:7666-
7670.

Gonzalez IL, Sylvester JE, Schmickel RD. 1988. Human 28S ribosomal RNA sequence
heterogeneity. Nucleic Acids Res. 16:10213-10224.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 29:1072-1075.

Gutell RR. 1992. Evolutionary Characteristics of 16S and 23S rRNA Structure. In: The Origin
and Evolution of the Cell. eds: H. Hartman KM. p. 243-309.

Gutell RR, Gray MW, Schnare MN. 1993. A compilation of large subunit (23S and 23S-like)
ribosomal RNA structures: 1993. Nucleic Acids Res. 21:3055-3074.

Gutell RR, Larsen N, Woese CR. 1994. Lessons from an evolving rRNA: 16S and 23S rRNA
structures from a comparative perspective. Microbiological reviews 58:10-26.

Hartman H, Fedorov A. 2002. The origin of the eukaryotic cell: a genomic investigation. Proc
Natl Acad Sci USA. 99:1420-1425.

Hassouna N, Mithot B, Bachellerie J-P. 1984. The complete nucleotide sequence of mouse 28S
rRNA gene. Implications for the process of size increase of the large subunit rRNA In higher
eukaryotes. Nucleic Acids Res. 12:3563-3583.

Hershko A, Heller H, Eytan E, Kaklij G, Rose IA. 1984. Role of the alpha-amino group of
protein in ubiquitin-mediated protein breakdown. Proc Natl Acad Sci USA. 81:7021-7025.

Heus HA, Pardi A. 1991. Structural features that give rise to the unusual stability of RNA
hairpins containing GNRA loops. Science 253:191-194.

Hillis DM, Dixon MT. 1991. Ribosomal DNA: Molecular Evolution and Phylogenetic Inference.
The Quarterly Review of Biology 66:411-453.

Hsiao C, Lenz TK, Peters JK, Fang P-Y, Schneider DM, Anderson EJ, Preeprem T, Bowman JC,
O'Neill EB, Lie L, et al. 2013. Molecular paleontology: a biochemical model of the ancestral
ribosome. Nucleic Acids Res. 41:3373-3385.

Huang H, Suslov NB, Li NS, Shelke SA, Evans ME, Koldobskaya Y, Rice PA, Piccirilli JA.
2014. A G-quadruplex-containing RNA activates fluorescence in a GFP-like fluorophore. Nat
Chem Biol. 10:686-691.

Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,
Hernsdorf AW, Amano Y, Ise K, et al. 2016. A new view of the tree of life. Nature Microbiology
1:16048.

Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic
gene recognition and translation initiation site identification. BMC Bioinformatics 11:119.

Imachi H, Nobu MK, Nakahara N, Morono Y, Ogawara M, Takaki Y, Takano Y, Uematsu K,
Ikuta T, Ito M, et al. 2020. Isolation of an archaeon at the prokaryote–eukaryote interface. Nature
577:519-525.

Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, Bateman A, Finn
RD, Petrov AI. 2017. Rfam 13.0: shifting to a genome-centric resource for non-coding RNA
families. Nucleic Acids Res. 46:D335-D342.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol. 30:772-780.

Khatter H, Myasnikov AG, Natchiar SK, Klaholz BP. 2015. Structure of the human 80S
ribosome. Nature 520:640.

Klinge S, Voigts-Hoffmann F, Leibundgut M, Arpagaus S, Ban N. 2011. Crystal Structure of the
Eukaryotic 60S Ribosomal Subunit in Complex with Initiation Factor 6. Science 334:941-948.

Klinger CM, Spang A, Dacks JB, Ettema TJG. 2016. Tracing the Archaeal Origins of Eukaryotic
Membrane-Trafficking System Building Blocks. Mol Biol Evol. 33:1528-1541.

Knorr AG, Schmidt C, Tesina P, Berninghausen O, Becker T, Beatrix B, Beckmann R. 2018.
Ribosome–NatA architecture reveals that rRNA expansion segments coordinate N-terminal
acetylation. Nat Struct Mol Biol. 26:35-39.

Kovacs NA, Petrov AS, Lanier KA, Williams LD. 2017. Frozen in Time: The History of
Proteins. Mol Biol Evol. 34:1252-1260.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. Mol Biol Evol. 34:1812-1819.

Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer:
consistent and rapid annotation of ribosomal RNA genes. Nucleic Acids Res. 35:3100-3108.

Lapeyre B, Michot B, Feliu J, Bachellerie J-P. 1993. Nucleotide sequence of the
Schizosaccharomyces pombe 25S ribosomal RNA and its phylogenetic implications. Nucleic
Acids Res. 21:3322-3322.

Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in
nucleotide sequences. Nucleic Acids Res. 32:11-16.

Lavender CA, Lorenz R, Zhang G, Tamayo R, Hofacker IL, Weeks KM. 2015. Model-Free RNA
Sequence and Structure Alignment Informed by SHAPE Probing Reveals a Conserved Alternate
Secondary Structure for 16S rRNA. PLoS Comp Biol. 11:e1004126-e1004126.

Leffers H, Kjems J, Østergaard L, Larsen N, Garrett RA. 1987. Evolutionary relationships
amongst archaebacteria: A comparative study of 23 S ribosomal RNAs of a sulphur-dependent
extreme thermophile, an extreme halophile and a thermophilic methanogen. J Mol Biol. 195:43-
61.

Lenz TK, Norris AM, Hud NV, Williams LD. 2017. Protein-free ribosomal RNA folds to a near-
native state in the presence of Mg2+. RSC Advances 7:54674-54681.

Leshin JA, Heselpoth R, Belew AT, Dinman J. 2011. High throughput structural analysis of
yeast ribosomes using hSHAPE. RNA biology 8:478-487.

Levitt M. 1969. Detailed molecular model for transfer ribonucleic acid. Nature 224:759-763.

Li Z, Guo Q, Zheng L, Ji Y, Xie Y-T, Lai D-H, Lun Z-R, Suo X, Gao N. 2017. Cryo-EM
structures of the 80S ribosomes from human parasites Trichomonas vaginalis and Toxoplasma
gondii. Cell Res. 27:1275-1288.

Lynch M. 2007. The origins of genome architecture. Sunderland, Mass. Sinauer Associates.
ISBN 978-0-87893-484-3.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 17:10-12.

McCutcheon JP, Moran NA. 2011. Extreme genome reduction in symbiotic bacteria. Nature Rev
Microbiol. 10:13-26.

Melnikov S, Ben-Shem A, Garreau de Loubresse N, Jenner L, Yusupova G, Yusupov M. 2012.
One core, two shells: bacterial and eukaryotic ribosomes. Nat Struct Mol Biol. 19:560.

Melnikov S, Manakongtreecheep K, Söll D. 2018. Revising the Structural Diversity of
Ribosomal Proteins Across the Three Domains of Life. Mol Biol Evol. 35:1588-1598.

Melnikov SV, Kwok HS, Manakongtreecheep K, van den Elzen A, Thoreen CC, Soll D. 2019.
Archaeal ribosomal proteins possess nuclear localization signal-type motifs: implications for the
origin of the cell nucleus. Mol Biol Evol:671966.

Merino EJ, Wilkinson KA, Coughlan JL, Weeks KM. 2005. RNA structure analysis at single
nucleotide resolution by selective 2'-hydroxyl acylation and primer extension (SHAPE). J Am
Chem Soc. 127:4223-4231.

Mestre-Fos S, Penev PI, Richards JC, Dean WL, Gray RD, Chaires JB, Williams LD. 2019a.
Profusion of G-quadruplexes on both subunits of metazoan ribosomes. PLoS One 14:e0226177.

Mestre-Fos S, Penev PI, Suttapitugsakul S, Hu M, Ito C, Petrov AS, Wartell RM, Wu R,
Williams LD. 2019b. G-Quadruplexes in Human Ribosomal RNA. J Mol Biol. 431:1940-1955.

Michot B, Bachellerie JP. 1987. Comparisons of large subunit rRNAs reveal some eukaryote-
specific elements of secondary structure. Biochimie 69:11-23.

Moran NA. 2002. Microbial minimalism: genome reduction in bacterial pathogens. Cell
108:583-586.

Narrowe AB, Spang A, Stairs CW, Caceres EF, Baker BJ, Miller CS, Ettema TJG. 2018.
Complex Evolutionary History of Translation Elongation Factor 2 and Diphthamide Biosynthesis
in Archaea and Parabasalids. Genome biology and evolution 10:2380-2393.

Nawrocki EP, Eddy SR. 2013. Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics 29:2933-2935.

Ninio J, Favre A, Yaniv M. 1969. Molecular model for transfer RNA. Nature 223:1333-1335.

Nissen P, Ippolito JA, Ban N, Moore PB, Steitz TA. 2001. RNA tertiary interactions in the large
ribosomal subunit: the A-minor motif. Proc Natl Acad Sci U S A 98:4899-4903.

Noller HF, Kop J, Wheaton V, Brosius J, Gutell RR, Kopylov AM, Dohme F, Herr W, Stahl DA,
Gupta R, et al. 1981. Secondary structure model for 23S ribosomal RNA. Nucleic Acids Res.
9:6167-6189.

Novikova IV, Hennelly SP, Sanbonmatsu KY. 2012. Structural architecture of the human long
non-coding RNA, steroid receptor RNA activator. Nucleic Acids Res. 40:5034-5051.

Olsen GJ, Woese CR. 1993. Ribosomal RNA: a key to phylogeny. The FASEB Journal 7:113-
123.

Petrov AS, Bernier CR, Gulen B, Waterbury CC, Hershkovits E, Hsiao C, Harvey SC, Hud NV,
Fox GE, Wartell RM, et al. 2014a. Secondary Structures of rRNAs from All Three Domains of
Life. PLoS One 9:e88222.

Petrov AS, Bernier CR, Hsiao C, Norris AM, Kovacs NA, Waterbury CC, Stepanov VG, Harvey
SC, Fox GE, Wartell RM, et al. 2014b. Evolution of the ribosome at atomic resolution. Proc Natl
Acad Sci USA. 111:10251-10256.

Petrov AS, Gulen B, Norris AM, Kovacs NA, Bernier CR, Lanier KA, Fox GE, Harvey SC,
Wartell RM, Hud NV, et al. 2015. History of the ribosome and the origin of translation. Proc
Natl Acad Sci USA. 112:15396-15401.

Peyretaillade E, Biderre C, Peyret P, Duffieux F, Méténier G, Gouy M, Michot B, Vivarès CP.
1998. Microsporidian Encephalitozoon cuniculi , a unicellular eukaryote with an unusual
chromosomal dispersion of ribosomal genes and a LSU rRNA reduced to the universal core.
Nucleic Acids Res. 26:3513-3520.

Rajon E, Masel J. 2011. Evolution of molecular error rates and the consequences for
evolvability. Proc Natl Acad Sci USA. 108:1082-1087.

Schnare MN, Damberger SH, Gray MW, Gutell RR. 1996. Comprehensive Comparison of
Structural Characteristics in Eukaryotic Cytoplasmic Large Subunit (23 S-like) Ribosomal RNA.
J Mol Biol. 256:701-719.

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068-2069.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert
M, Söding J, et al. 2011. Fast, scalable generation of high-quality protein multiple sequence
alignments using Clustal Omega. Mol Syst Biol 7:539.

Smith TF, Lee JC, Gutell RR, Hartman H. 2008. The origin and evolution of the ribosome.
Biology Direct 3:16.

Spang A, Saw JH, Jørgensen SL, Zaremba-Niedzwiedzka K, Martijn J, Lind AE, van Eijk R,
Schleper C, Guy L, Ettema TJ. 2015. Complex archaea that bridge the gap between prokaryotes
and eukaryotes. Nature 521:173-179.

Spang A, Stairs CW, Dombrowski N, Eme L, Lombard J, Caceres EF, Greening C, Baker BJ,
Ettema TJG. 2019. Proposal of the reverse flow model for the origin of the eukaryotic cell based
on comparative analyses of Asgard archaeal metabolism. Nature Microbiology 4:1138-1148.

Spitale RC, Crisalli P, Flynn RA, Torre EA, Kool ET, Chang HY. 2013. RNA SHAPE analysis
in living cells. Nat Chem Biol. 9:18-20.

Stoddard CD, Gilbert SD, Batey RT. 2008. Ligand-dependent folding of the three-way junction
in the purine riboswitch. RNA 14:675-684.

Stothard P. 2000. The sequence manipulation suite: JavaScript programs for analyzing and
formatting protein and DNA sequences. BioTechniques 28:1102, 1104.

Tréhu AM, Bohrmann G, Rack FR, Torres ME, Richter C, Bangs NL, Barr SR, Borowski WS,
Claypool GE, Collett TS, et al. 2003. ODP Leg 204; gas hydrate distribution and dynamics
beneath southern Hydrate Ridge. JOIDES Journal 29:5.

Veldman GM, Klootwijk J, de Regt VC, Planta RJ, Branlant C, Krol A, Ebel JP. 1981. The
primary and secondary structure of yeast 26S rRNA. Nucleic Acids Res. 9:6935-6952.

Ware VC, Tague BW, Graham Clark C, Gourse RL, Brand RC, Gerbi SA. 1983. Sequence
analysis of 28S ribosomal DNA from the amphibian Xenopus laevis. Nucleic Acids Res.
11:7795-7817.

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2—a
multiple sequence alignment editor and analysis workbench. Bioinformatics 25:1189-1191.

Watts JM, Dang KK, Gorelick RJ, Leonard CW, Bess JW, Jr., Swanstrom R, Burch CL, Weeks
KM. 2009. Architecture and secondary structure of an entire HIV-1 RNA genome. Nature
460:711-716.

Wilkinson KA, Merino EJ, Weeks KM. 2005. RNA SHAPE Chemistry Reveals Nonhierarchical
Interactions Dominate Equilibrium Structural Transitions in tRNAAsp Transcripts. J Am Chem
Soc. 127:4659-4667.

Wilkinson KA, Merino EJ, Weeks KM. 2006. Selective 2'-hydroxyl acylation analyzed by
primer extension (SHAPE): quantitative RNA structure analysis at single nucleotide resolution.
Nat Protoc. 1:1610-1616.

Williams TA, Cox CJ, Foster PG, Szöllősi GJ, Embley TM. 2020. Phylogenomics provides
robust support for a two-domains tree of life. Nat Ecol Evol. 4:138-147.

Woese CR, Fox GE. 1977. Phylogenetic structure of the prokaryotic domain: the primary
kingdoms. Proc Natl Acad Sci USA. 74:5088-5090.

Woese CR, Kandler O, Wheelis ML. 1990a. Towards a natural system of organisms: proposal
for the domains Archaea, Bacteria, and Eucarya. Proc Natl Acad Sci USA. 87:4576-4579.

Woese CR, Magrum LJ, Gupta R, Siegel RB, Stahl DA, Kop J, Crawford N, Brosius J, Gutell R,
Hogan JJ, et al. 1980. Secondary structure model for bacterial 16S ribosomal RNA:
phylogenetic, enzymatic and chemical evidence. Nucleic Acids Res. 8:2275-2293.

Woese CR, Winker S, Gutell RR. 1990b. Architecture of ribosomal RNA: constraints on the
sequence of "tetra-loops". Proc Natl Acad Sci U S A 87:8467-8471.

Yergeau E, Maynard C, Sanschagrin S, Champagne J, Juck D, Lee K, Greer CW. 2015.
Microbial community composition, functions and activities in the Gulf of Mexico, one year after
the Deepwater Horizon accident. Appl Environ Microbiol. 81:5855-5866.

Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester E, Seitz
KW, Anantharaman K, Starnawski P, Kjeldsen KU, et al. 2017. Asgard archaea illuminate the
origin of eukaryotic cellular complexity. Nature 541:353.

Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. Nucleic
Acids Res. 31:3406-3415.