Title:

Chemical map–based prediction of nucleosome positioning using the

Bioconductor package nuCpos

Authors:

Hiroaki Kato[1], Mitsuhiro Shimizu[2] and Takeshi Urano[1]

Institutional addresses:

1. Department of Biochemistry, Shimane University School of Medicine, Izumo,

Shimane 693-8501, Japan

2. Department of Chemistry, Graduate School of Science and Engineering,

Program in Chemistry and Life Science, School of Science and Engineering,

Meisei University, Hino, Tokyo 191-8506, Japan

Email addresses:

Hiroaki Kato: hkato@med.shimane-u.ac.jp

Mitsuhiro Shimizu: shimizum@chem.meisei-u.ac.jp

Takeshi Urano: turano@med.shimane-u.ac.jp

Corresponding author:

Hiroaki Kato

Department of Biochemistry, Shimane University School of Medicine

89-1 Enya-cho, Izumo 693-8501, Japan

Tel: +81-853-20-2127; Fax: +81-853-20-2125

hkato@med.shimane-u.ac.jp

Abstract [350 words]

Background:

Assessing the nucleosome-forming potential of specific DNA sequences is

important for understanding complex chromatin organization. Methods for

predicting nucleosome positioning include bioinformatics and biophysical

approaches. An advantage of bioinformatics methods, which are based on *in*

vivo nucleosome maps, is the use of natural sequences that may contain

previously unknown elements involved in nucleosome positioning *in vivo*. The

accuracy of such prediction attempts reflects the genomic coordinate resolution

of the nucleosome maps applied. Nucleosome maps are constructed using

micrococcal nuclease digestion followed by high-throughput sequencing

(MNase-seq). However, as MNase has a strong preference for A/T-rich

sequences, MNase-seq may not be appropriate for this purpose. In addition to

MNase-seq–based maps, base pair–resolution chemical maps of *in vivo*

nucleosomes from three different species (budding and fission yeasts, and mice)

are currently available. However, these chemical maps have yet to be integrated

into publicly available computational methods.

Results:

We developed a Bioconductor package (named nuCpos) to demonstrate the

superiority of chemical maps in predicting nucleosome positioning. The accuracy

of chemical map–based prediction in rotational settings was higher than that of

the previously developed MNase-seq–based approach. With our method,

predicted nucleosome occupancy reasonably matched *in vivo* observations and

was not affected by A/T nucleotide frequency. Effects of genetic alterations on

nucleosome positioning that had been observed in living yeast cells could also

be predicted. nuCpos calculates individual histone binding affinity (HBA) scores

for given 147-bp sequences to examine their suitability for nucleosome formation.

We also established local HBA as a new parameter to predict nucleosome

formation, which was calculated for 13 overlapping nucleosomal DNA

subsequences. HBA and local HBA scores for various sequences agreed well

with previous *in vitro* and *in vivo* studies. Furthermore, our results suggest that

nucleosomal subsegments that are disfavored in different rotational settings

contribute to the defined positioning of nucleosomes.

Conclusions:

Our results demonstrate that chemical map–based statistical models are

beneficial for studying nucleosomal DNA features. Studies employing nuCpos

software can enhance understanding of chromatin regulation and the

interpretation of genetic alterations and facilitate the design of artificial

sequences.

Keywords:

nucleosome, prediction of nucleosome positions, dHMM, HBA, local HBA,

translational and rotational settings, nucleosomal DNA subsegments

Background:

The nucleosome is a structure in which a ~147-base pair (bp) double-stranded

DNA forms a left-handed super-helix to wrap a histone octamer consisting of two

copies of histones H2A, H2B, H3, and H4 [1-3]. The positioning of nucleosomes

is defined by rotational and translational settings [4, 5]. Rotational settings define

the histone-DNA contact surfaces on the DNA string, whereas translational

position is the distinct position of the nucleosome (i.e., position of the dyad base)

with respect to the genomic coordinate. The positioning of nucleosomes along

genomic DNA is critical for various biological events, including transcription,

DNA replication and repair, and chromosome segregation [4]. In these events,

nucleosome positioning is combinatorially regulated by *trans*-acting factors (e.g.,

transcription factors and chromatin remodeling factors) and by the inherent

suitability of the DNA sequence for nucleosome formation [5-7]. Focusing on the

latter requirement, computational methods that assess the nucleosome-forming

potential of given DNA sequences have been developed [8].

These methods are classified into two groups: biophysical and

bioinformatics [8, 9]. Most biophysical methods are based on the structures of

reconstituted nucleosome core particles (NCPs) such as NCP147, which

contains a palindromic 147-bp sequence with intrinsic positional stability [3, 10,

11]. The use of the palindromic sequence optimized for homogeneous

nucleosome reconstitution and the highest resolution of the structure contributed

greatly to the development of prediction algorithms. However, relying on the

representative structure itself could also be a potential weakness, as there could

be unrecognized natural intra-nucleosomal DNA elements that mediate specific

physiologic functions in cells.

In contrast to the biophysical prediction methods, bioinformatics

methods are based on maps of *in vivo* nucleosomes determined using

high-throughput sequencing techniques. The use of complex, natural

nucleosomal sequences may cause difficulty in the interpretation of prediction

results; however, it may also lead to unexpected findings that enhance

understanding of nucleosome-based regulation.

A weakness of currently available bioinformatics methods is that the

statistical models are constructed based on *in vivo* nucleosome maps generated

using micrococcal nuclease (MNase) digestion followed by high-throughput

sequencing (MNase-seq). MNase digests linker DNA that connects NCPs,

allowing researchers to obtain nucleosomal DNA that has been protected from

enzymatic digestion. In MNase-seq experiments, DNA fragments of

mono-nucleosome size (~150 bp) are sequenced and mapped to the reference

genome. However, as MNase strongly prefers A/T-rich sequences, unwanted

digestion of A/T-rich nucleosomal sequences and less-effective digestion of

G/C-rich linkers may occur, leading to variation in nucleosome fragment size [7,

12]. Because of this enzymatic bias, precise determination of the genomic

coordinates of *in vivo* nucleosomes using MNase-seq (i.e., dyad base calling) is

challenging. In addition, although some correction methods for MNase-seq

coverage against enzymatic bias have been developed [13-15], no method that

corrects the deduced nucleosome position by shifting it to an appropriate

rotational position is available. Thus, the resolution of MNase-seq–based maps

with respect to the genomic coordinates is expected to be inevitably low.

One bioinformatics method considered successful in a thorough

comparative study [9] is NuPoP, which is built upon a duration hidden Markov

model (dHMM) that considers information regarding both the nucleosome and

linker DNA [16]. NuPoP enables the prediction of nucleosome positioning in nine

different species by rescaling the base composition of both the nucleosome and

linker models for the budding yeast *Saccharomyces cerevisiae* with those of

target organisms. The budding yeast models are generated using an

MNase-seq–based nucleosome map by considering the central base of each

nucleosomal fragment of varying length the dyad base [16]. This may particularly

affect construction of the time-dependent statistical model for the nucleosome

state, which ideally requires that the dyad bases be determined at bp resolution.

The construction of the linker model, which requires extraction of linker

sequences as genomic sequences not covered by nucleosomes, may also be

affected. Thus, although selecting the MNase-seq–based map was the best

practice at the time of the software's development, the prediction results output

provided by NuPoP can be biased by both the enzymatic preference of MNase

and the difficulty of dyad base calling.

A breakthrough in nucleosome mapping occurred with the development

of a method for site-directed chemical cleavage of nucleosomal DNA followed by

high-throughput sequencing [17, 18]. This technique, called chemical mapping,

was first applied to budding yeast and subsequently to the fission yeast

Schizosaccharomyces pombe and embryonic stem cells of the house mouse,

Mus musculus [17, 19-22]. In cells expressing histone H4 with an S47C amino

acid substitution, nucleosomal DNA is specifically cleaved near the nucleosome

center by site-directed chemical cleavage. By examining the cleavage sites,

previous studies determined the dyad positions of *in vivo* nucleosomes in the

three species at bp resolution [17, 19, 22]. The published maps contained whole

and representative sets of identified nucleosomes, which are referred to as

"redundant" and "unique" nucleosomes, respectively [17, 19, 22]. In addition to

the histone H4-S47C approach, chemical mapping of nucleosomes using a

histone H3 mutation (Q85C) determined the precise positions of −1 and +1

nucleosomes for protein-coding genes [12]. However, the bp-resolution chemical

maps have not yet been implemented in publicly available software for prediction

of nucleosome positioning.

One goal of nucleosome positioning prediction is *in silico* reproduction

of *in vivo* observations, preferably at the locus level. If this goal is achieved,

predicting the effects of genetic alterations on nucleosome positioning and

prediction-based engineering of nucleosome-forming and -depleted sequences

would also be practical. In this regard, budding and fission yeasts would serve

as preferable model organisms, because nucleosomes formed on native and

modified sequences *in vivo* have been precisely mapped [23-28]. These

nucleosomes are suitable for *in silico* reproduction of *in vivo* observations.

Importantly, for the currently available bioinformatics method NuPoP, the

reliability of locus-level nucleosome prediction has seldom been demonstrated,

despite its wide use in previous studies [29-38].

166   In this study, we developed a nucleosome positioning prediction

method that utilizes available chemical maps of three species for construction of

dHMMs. We show that the prediction accuracy of chemical map–based models

is higher with respect to the genomic coordinates than that of MNase-seq–based

models. The chemical map–based models appear to recognize appropriate

sides of DNA strings as histone-DNA contact surfaces. The effects of genetic

alterations on nucleosome positioning previously observed in living yeast cells

were reasonably reproduced *in silico*, suggesting that this software is useful for

chromatin engineering and interpretations of mutation effects. We found that the

prediction accuracy for mice was lower than that for yeasts, which should be

acknowledged by software users. Our software, nuCpos, is available at the

Bioconductor website (https://doi.org/doi:10.18129/B9.bioc.nuCpos).

Results:

Chemical map–based prediction of nucleosome positioning in yeasts

In order to determine whether the bp-resolution chemical maps are suitable for

predicting nucleosome positioning, we trained dHMMs with the nucleosome

maps generated with H4-S47C–dependent DNA cleavage (see **Methods** and

Table 1). The core dHMM algorithm implemented in our method (nuCpos) was

not altered from the original algorithm developed for the MNase-seq–based

method (NuPoP) to clarify the advantages of using the chemical maps.

187   First, we predicted nucleosome positioning in the genomes of budding

and fission yeasts (denoted 'Sc' and 'Sp', respectively, in **Figure 1**) with models

that were constructed with *in vivo* nucleosome maps of target species. Aiming to

evaluate the prediction accuracy with respect to the genomic coordinates,

prediction results were assessed by checking the matching status of predicted

nucleosomes with those experimentally determined by means of histone

H4-S47C–based chemical cleavage [17, 19]. Here, MNase-seq–based maps are

inappropriate to use as true-position references due to their lower resolution, as

described in the Background section. Ideally, prediction results should be

evaluated at 1-bp resolution, so that it will be possible to accurately discern the

rotational settings of given nucleosomes. As the resolution of the chemical maps

that we used for model construction could be 1-2 bp or lower, the resolution of

the prediction results would be expected to be at a similar level. Indeed, the

distance between chemically predicted nucleosomes and the nearest *in vivo*

nucleosomes was generally 0 bp (located at the same position) or 1 bp (located

at the juxtaposed nucleotide positions) (**Figure 1A**, right panel).

In order to determine the appropriate resolution for examining

prediction results, we calculated the matching rate between predicted and *in vivo*

nucleosomes with stepwise widening of the matching window (**Figure 1B**). Here,

we defined the matching window as the allowable distance between

nucleosomes regarded as matched. In chemical map–based predictions

(nuCpos), the matching rate doubled to over 50% when the matching window

was widened from 0 bp (1-bp resolution) to 1 bp (2-bp resolution) (**Figure 1B**,

red lines). However, the rate did not markedly increase upon further widening of

the matching window, suggesting that the chemical models distinguish the

histone-facing side of the DNA string.

In MNase-seq–based predictions (NuPoP), by contrast, the matching

rate increased gradually as the matching window was widened from 0 to 9 bp

(**Figure 1B**, blue lines). The matching rate drew a very loose anti-S shape,

suggesting that the MNase-seq–based model still recognized the histone-facing

side of the DNA string, but only to a small extent. We also noticed that a

substantial population of nucleosomes predicted with the MNase-seq–based

model was positioned approximately 5 nucleotides away from the *in vivo*

nucleosomes (**Figure 1A**, left panel, marked with a blue horizontal bar). These

observations suggest that the MNase-seq–based model has difficulty in

discriminating rotationally mispredicted nucleosomes. Considering these results,

we concluded that prediction results should be evaluated at 2-bp resolution, as

further widening of the matching window would increase the misprediction rate.

Figure 1C presents the evaluation of prediction results against

representative "unique" and whole "redundant" nucleosomes [17, 19] at 2-bp

resolution. When the budding yeast MNase-seq–based model (NuPoP) was

applied to the prediction of nucleosome positioning in the budding yeast genome

(Target: Sc, Model: Sc), only 3.2% and 12.6% of the nucleosomes on the most

probable (Viterbi) path matched *in vivo* unique (red) and redundant (orange)

nucleosomes, respectively. When the fission yeast model was applied to the

prediction of nucleosome positioning in the fission yeast genome (Target: Sp,

Model: Sp), the MNase-seq–based model scored 2.7% and 11.0%, respectively.

In contrast, when the budding yeast chemical map–based model (nuCpos) was

used for predictions (Target: Sc, Model: Sc), 23.5% and 50.9% of the

nucleosomes on the Viterbi path matched unique and redundant nucleosomes,

respectively. For the fission yeast (Target: Sp, Model: Sp), the corresponding

scores were 25.5% and 54.2%, respectively. Thus, with the chemical map–

based models, over half of the nucleosomes predicted as being on the Viterbi

path are present in living cells at exactly the same positions or at the juxtaposed

nucleotide positions.

In the chemical maps of budding and fission yeasts, *in vivo* unique

nucleosomes accounted for 19.6% and 17.8%, respectively, of all redundant

nucleosomes [17, 19]. Thus, in the MNase-seq–based predictions, enrichment

of unique nucleosomes on the Viterbi paths was only 1.3- and 1.4-fold,

respectively. In contrast, the chemical map–based predictions exhibited 2.4- and

2.6-fold enrichment, respectively. The higher enrichments suggest that chemical

map–based models better distinguish sequences favorable for nucleosome

formation [17, 19], compared with MNase-seq–based models.

Since the dHMMs were constructed using nucleosome and linker

information based solely on the unique nucleosomes (see **Methods**), detection

of portion of the redundant nucleosomes (from which unique nucleosomes have

been removed) is worthwhile. The nucleosomes in this category accounted for

27.4% and 28.7% of the Viterbi nucleosomes in the chemical-map–based

predictions for the budding and fission yeasts, respectively, and the proportions

were higher than the MNase-seq–based predictions (9.4% and 8.3%,

respectively). Giving these results, the chemical map–based models provide

better prediction outcomes with respect to the genomic coordinates than the

MNase-seq–based models.

Prediction of *in vitro* reconstituted nucleosome positions

The evaluations described above suggested that chemical map–based models

are superior in terms of the prediction of rotational settings. To confirm this

observation, we compared HBA scores [16] calculated along the original 282-bp

Widom 601 sequence [10] with the chemical map–based and MNase-seq–

based models (**Figure 2**). HBA scores are calculated in the dHMM algorithms in

order to examine the suitability of given 147-bp sequences for nucleosome

formation [16], and they are expected to predict the rotational setting of

nucleosomes if the resolution of the nucleosome map used for model

construction is sufficiently high. The original 282-bp Widom 601 sequence

contained primer sequences for the SELEX study at both ends and additional

sequences flanking the nucleosome-forming 147-bp sequence centered at

nucleotide position 154 [10]. When analyzed using the chemical map–based

model, discrete HBA peaks with an interval of approximately 10 bp were

observed at positions 124, 135, 143, 154, 164, and 175 (asterisks). This

periodicity indicated that nearly the same surface of the DNA string was

predicted to be in contact with the histone core in these potential translational

positions. Among these positions, the one with the highest HBA score

corresponds to the translational position of the *in vitro* stable nucleosome

(position 154, orange vertical lines). In contrast, the MNase-seq–based HBA

score was relatively low at position 154, and the peaks did not exhibit the

apparent 10-bp periodicity. Analyses of the mouse mammary tumor virus

(MMTV) 3'–long terminal repeat (LTR) sequence and *Xenopus borealis* 5S RNA

gene also showed the advantage of the chemical model (**Additional file 1:**

Figure S1). Collectively, the chemical map–based model reliably predicted the

rotational settings of the nucleosomes reconstituted with these suitable

sequences.

Prediction of *in vivo* nucleosome positions

Next, we calculated HBA scores around −1 and +1 nucleosomes for 5,542

protein-coding genes identified by means of histone H3-Q85C–based chemical

cleavage [12]. Due to the differences in detection methods, positions of

H3-Q85C–based nucleosomes are not always identical to those of the

corresponding H4-S47C–based nucleosomes [12]. In this regard, the H3-Q85C–

based map differed from the H4-S47C–based map that was used for the

statistical model construction. Thus, the H3-Q85C–based nucleosomes would

be a good reference for evaluation of prediction accuracy at bp-resolution.

As shown in **Figure 3A**, mean HBA scores calculated using both

models exhibited a periodicity with a 10-bp interval. However, the amplitude of

the chemical map–based HBA scores (red) around the tested nucleosomes was

larger than that of the MNase-seq–based HBA scores (blue). This suggested

that the chemical model could clearly distinguish the surface of the DNA string

that interacts with the histone core even when natural genomic sequences were

queried. We found that the chemical map–based HBA scores were highest at

the −1 and +1 nucleosome positions (0 bp). Chemical map–based HBA scores

for the three neighboring translational positions (10, 20, and 30 bp in distance)

relative to the nucleosome-depleted regions (marked with asterisks in **Figure**

3A) remained relatively high despite the gradual increase in the A/T frequency of

the tested sequences (**Figure 3B**). After the A/T frequency reached a plateau

(A/T≈0.64), the chemical map–based HBA scores began to decrease. In

contrast, MNase-seq–based HBA scores at the −1 and +1 nucleosome positions

(0 bp) were already at moderate levels, and they simply decreased as the A/T

frequency increased. Thus, the MNase-seq–based model was adversely

influenced by the enzymatic bias of MNase, whereas the chemical map–based

model was not.

The higher accuracy of chemical map–based models at the genomic

level prompted us to perform a locus-level evaluation of prediction results, using

the budding yeast *TRP1ARS1* minichromosome [25, 27]. On this well-studied,

circular, 1,465-bp DNA, three (numbered I to III) and four (IV to VII)

nucleosomes are interspersed with two nucleosome-depleted regions or

nuclease-hypersensitive regions designated *HSRA* and *HSRB*. *HSRA* includes

the DNA replication origin *ARS1*, whereas *HSRB* is located upstream of the

TRP1 marker gene. At a glance, the chemical map– and MNase-seq–based

model prediction results appeared very different (**Figure 3C**). Each model

suggested eight and seven representative nucleosomes, respectively. Seven of

the eight nucleosomes predicted by the chemical map–based model (**Figure 3C**,

nuCpos) reasonably matched the positions of *in vivo* nucleosomes determined

previously [27]. The only exception was a predicted nucleosome located in
HSRA, in which functional *ARS1* would be occupied by the ARS-binding factor
Abf1 and the origin recognition complex instead of the nucleosome to maintain
this circular DNA across generations [39]. This prediction was reasonable, as
DNA replication origins are known to be covered by nucleosomes when the
subunits of the origin recognition complex or Abf1 are perturbed *in vivo* [39-41].
In this native sequence, the chemical map–based HBA scores
exhibited a periodicity of an approximately 10-bp interval (**Additional file 1:**
Figure S2, nuCpos). As the dHMM considers the linker length distribution as a
major factor affecting nucleosome-linker state transition and does not allow
nucleosomes with shorter DNAs (<147 bp), nucleosome positions tend to be
selected in a limited manner. Although there were many high-HBA positions,
only a few were clearly selected by the dHMM as high-probability nucleosome
dyad positions. Thus, the predicted occupancy should be cautiously considered,
as it appears to provide only a rough image of chromatin state. Instead, it should
be kept in mind that periodic HBA scores may contribute to redundant
nucleosome positioning *in vivo*.
In contrast to the chemical map–based model, most of the nucleosome
locations predicted by the MNase-seq–based model were not in agreement with
their positions *in vivo* (**Figure 3C**, NuPoP); most of the *in vivo* nucleosome
centers (inverted triangles in figure) were located in the predicted linkers or
nucleosome ends. This result was consistent with the lower genome-wide
prediction accuracy (**Figure 1**). Importantly, MNase-seq–based HBA scores for
this sequence appeared to mirror the A/T-content over the X-axis (**Figure 3C**,
A/T); high-HBA positions were generally A/T-poor, and low-HBA positions were
A/T-rich. In addition, the local amplitude of MNase-seq–based HBA scores was
apparently lower compared with the chemical map–based model (**Additional**
file 1: Figure S2). These observations also demonstrated that MNase-seq–
based model prediction results are strongly influenced by the A/T preference of
MNase. A/T-content exhibited a similar effect in the MNase-seq–based
nucleosome positioning prediction around the fission yeast *ura4*+ gene
(**Additional file 1: Figure S3**) [42]. Given these results, we concluded that the

chemical map–based models output better genome-wide and locus-level

prediction results.

Prediction of effects of genetic alterations on nucleosome positioning

Next, we examined whether the effects of genetic alterations on nucleosome

positioning could be predicted. In **α** cells of budding yeast, the **a**-cell-specific

gene *BAR1* is repressed by *α2*-repressor–dependent positioning of the

nucleosome at its promoter, which masks the gene's *TATA* box [23, 26]. The

chemical map–based model predicted the positioning of this repressive

nucleosome with high accuracy (**Figure 4**, WT). The HBA score for this

translational position (0 bp), which is 82 bp away from the *α2*-operator, was very

high (HBA=8.47), suggesting that repression of the *BAR1* gene is assisted by

the intrinsic suitability of the promoter DNA for nucleosome formation. When a

36-bp sequence consisting of 12 repeats of CTG is inserted at the center of this

nucleosome, repression of *BAR1* and nucleosome positioning are not affected *in*

vivo [26]. Prediction results agreed that this insertion should not cause

nucleosome depletion (**Figure 4**, CTG_{12}). Other repeat sequences causing no

transcriptional derepression *in vivo* [26] were also predicted to not cause

nucleosome depletion (**Figure 4**, Sac_5 and Sac_6). It is also known that insertion

of a 30-bp A-stretch or a 10-bp CG (i.e., CpG) repeat sequence inhibits *in vivo*

nucleosome formation and causes derepression of *BAR1* [26]. Remarkably, the

effect of these genetic alterations on nucleosome positioning was reproduced *in*

silico with the chemical map–based model (**Figure 4**, A_{30} and CG_5). We noted

that the disruptive effect of fragment insertion on nucleosome positioning was

more pronounced in predictions than *in vivo* observations. Insertion of a 20-bp

A-stretch, which does not cause complete nucleosome depletion *in vivo*, was

clearly predicted to inhibit nucleosome formation (**Figure 4**, A_{20}). However, in the

case of CG repeat insertion, shortening of the repeat to 8 bp, which satisfies *in*

vivo nucleosome formation [26], was predicted to preserve the capability to form

nucleosomes (**Figure 4**, CG_4). We also confirmed that the inhibition of

nucleosome formation by telomeric DNA insertion demonstrated in a previous

study [43] was also largely predictable (**Additional file 1: Figure S4**). Thus,

overall, the effects of genetic alterations on nucleosome positioning can be

predicted with the chemical map–based model.

Local HBA for nucleosomal DNA subsegments

The successful increase in prediction accuracy and better discrimination of

rotationally favorable nucleosome positions with chemical map–based models

suggested that the chemical map–based HBA score for a given 147-bp

sequence is a good indicator of the suitability of that sequence for nucleosome

formation. Considering that nucleosomal DNA makes contact with histone

proteins at each superhelical location (SHL ±0.5-6.5), we defined a novel

parameter, designated "local HBA", to examine the suitability of

intra-nucleosomal DNA segments for each histone-DNA contact. In order to

implement this idea, a 147-bp whole nucleosomal segment was divided into 13

overlapping nucleosomal DNA subsegments, designated A through M

(**Additional file 1: Figure S5**, see **Methods**). Each 20- or 21-bp segment

corresponded to two histone-DNA contact sites, each of which was shared by

neighboring segments. For example, segment B corresponded to SHL −5.5 and

−4.5, which were shared by segments A and C, respectively. We took this

overlapping approach because it remains unclear how surrounding sequences

affect histone-DNA contacts. Calculating local HBA scores is conceptually the

same as that of HBA calculations, which consider the probability of a whole

147-bp sequence to be a nucleosome [16]. In local HBA calculation, for each

intra-nucleosomal segment of a given 147-bp DNA that wraps a histone octamer

to form a nucleosome, the probability of the selected 20- or 21-bp sequence to

be located at that part of the nucleosome and the probability of the same

sequence functioning as a linker are calculated. The log likelihood ratio of these

probabilities is defined as the local HBA score for that segment. Thus, the sum of

local HBA scores for the non-overlapping set of seven 21-bp segments, A, C, E,

G, I, K, and M (147 bp in total), would be nearly equal to the HBA score

calculated for the same 147-bp sequence.

In order to examine whether local HBA scores indicate the suitability of

nucleosomal subsegments for histone-DNA contacts, we used the 282-bp

Widom 601 sequence, which was used for the above HBA analysis (**Figure 2**),

as a test sequence. **Figure 5A** presents a heatmap of the calculated local HBA

scores. The topmost track, marked "A", represents local HBA scores for the A

segments of 147-bp sequences centered at the indicated dyad positions. At

potential translational positions with high HBA scores (**Figure 5A**, arrows), local

HBA scores for some segments (e.g., segments D and E of the 147-bp

sequence centered at position 154) were very high, as expected. However, each

147-bp sequence also contained segments with relatively low local HBA scores.

Interestingly, at translational positions with low HBA scores (those intervened by

the high-scoring HBA positions), some segments (e.g., segments F and G at

position 157) exhibited very low local HBA scores. This tendency was

particularly notable for the central segments E through I. The finding of these

low-scoring local HBA segments suggests that sub-sequences that are

disfavored in other rotational settings play a role in determining rotational

settings.

Next, we focused on the 147-bp Widom 601 sequence, of which the

dyad base is located at nucleotide position 154 of the original sequence (**Figure**

5A, red arrow). We chose this sequence because its derivatives are

well-characterized in reconstitution studies, as it forms a homogeneous

nucleosome, and because the left and right halves exhibit different features

[44-48]. In agreement with previous transcription studies [44, 45, 49], local HBA

scores for the 147-bp sequence were highest for segments D and E, both of

which share SHL −2.5, in which the high-affinity R3 element is located (**Figure**

5B, red dots). The region covered by segments D and E (SHL −3.5 to −1.5) is

also known for its very strong histone-DNA interaction [46]. These data indicate

that local HBA scores are useful for evaluating the suitability of nucleosomal

subsegments for histone-DNA contacts.

When the translational position was shifted along the 282-bp sequence

from nucleotide position 154 to surrounding nucleotide positions ranging from

150 to 158 (**Figure 5C**, −4 to +4 with respect to 154), the local HBA scores for

segments B through I changed dramatically (**Figure 5B**, black dots). However,

such shifting resulted in modest changes at segments J through M, with their

scores remaining at relatively high levels. These observations suggest that

segments J through M in the right-half sequence are generally suitable for

nucleosome formation, which may also be true even when the translational

position is shifted by several base pairs. In other words, segments J through M

do not seem to have intra-nucleosomal disfavored elements that strongly limit

the rotational setting. This nature might account for the observed poor crystal

development or lower homogeneity [46], as this 'slippery' sequence could disrupt

the uniformity of histone-DNA interactions.

Local HBA scores for modified nucleosome-forming sequences

The above testing with the sequence previously used in reconstitution studies

suggested that calculating local HBA scores would enhance understanding of

how DNA sequences characterize the nucleosomes formed *in vitro*. This

prompted us to examine whether we could apply local HBA calculations to the

evaluation of intracellular nucleosome formation. González et al. reported that

sequence-based determinants of nucleosome positioning are dispersed across

nucleosomes [42]. One gene that they thoroughly investigated was the fission

yeast gene, *ura4+*. This gene has six nucleosomes (designated +1 to +6) that

overlap the coding sequence, five of which were predictable with the chemical

map–based dHMM (**Additional file 1: Figure S3**, nuCpos). Although the

remaining nucleosome (+5) was not clearly observed in the predicted occupancy

plot, there was one position with a relatively high HBA score that reasonably

corresponded to the *in vivo* nucleosome (marked with an asterisk in the figure).

A heatmap of local HBA scores for this gene suggested that there is an element

disfavored for nucleosome formation near the +5 nucleosome (**Figure 6**, WT,

see the blueish slanted line spanning segments A through M around dyad

position 600-750). Thus, this element may have caused the lower predicted

occupancy of the +5 nucleosome.

González et al. generated three modified sequences, namely Dyad,

Linker, and Int, for the *ura4+* gene and studied nucleosome positioning on these

sequences *in vivo* [42]. In Dyad and Linker, 51-bp sequences centered to each

nucleosome dyad and each linker, respectively, were replaced with artificial

randomized sequences of the same length. In the sequence Int, two

intra-nucleosomal sequences of each nucleosome (positions from −51 to −24

and 24 to 51) were replaced. According to their report, the occupancy of the +1

nucleosome was markedly decreased in the Dyad sequence, that of the +2

nucleosome was slightly increased in the Dyad and Linker sequences, and the

position of the +3 nucleosome was shifted in the Int sequence [42]. However,

why these changes selectively occur on specific nucleosomes *in vivo* remains

unclear due to a lack of appropriate analytical methods.

A heatmap of local HBA scores for the Dyad sequence clearly showed

that the replacements in this sequence caused a marked decrease in local HBA

scores only around the position of the +1 nucleosome (**Figure 6**, Dyad, blue

blots not present in the WT map). In contrast, the 51-bp replacements at other

nucleosomal centers did not appear to dramatically change their suitability for

nucleosome formation. As to the +2 nucleosome, we did not observe any

supportive signatures that could explain why the Dyad and Linker replacements

slightly increased its occupancy (**Figure 6**, Dyad and Linker). We observed that

the Int replacements primarily affected local HBA scores around the position of

the +3 nucleosome (**Figure 6**, Int, see the blueish slanted line spanning

segments A through M). This suggests that the mutation specifically affects the

+3 nucleosome positioning and triggers a shift of it to a more suitable position *in*

vivo. Giving these results, local HBA score calculation for nucleosomal

subsegments is valuable for *in silico* assessment of modified

nucleosome-forming sequences.

Chemical map–based prediction of nucleosome positioning in mice

The observation that the chemical-map–based models successfully improved

the prediction accuracy of nucleosomes in both yeasts suggested that these

approaches would be useful for mice. We therefore examined nucleosome

positioning in the mouse genome using the MNase-seq–based and chemical

map–based models. Matching of the predicted nucleosomes on the Viterbi paths

at a 2-bp resolution demonstrated the better performance of the chemical map–

based model (**Figure 7A**). In the MNase-seq–based prediction, 1.4% and 11.0%

of predicted nucleosomes matched unique and redundant nucleosomes,

respectively. In contrast, the chemical map–based prediction scored 10.0% and

41.1%, respectively. Thus, the chemical map–based model is more suitable for

prediction of nucleosome positioning with respect to the genomic coordinates

than the MNase-seq–based model.

In order to determine whether the mouse chemical map–based model

could better predict the rotational setting of nucleosomes, as shown in the case

of yeasts, we calculated the matching rate between predicted and *in vivo*

nucleosomes with stepwise widening of the matching window (**Figure 7B**). As

expected, the matching rate for MNase-seq–based prediction increased

gradually as the matching window widened. In chemical map–based predictions,

a dramatic increase in matching rate was observed when the matching window

was widened from 0 to 1 bp. However, the matching rate also increased

gradually as the matching window was widened further (See **Figures 1B and 7B**

for comparison between species). The shapes of the matching rate curves

suggested the chemical map–based model still distinguished rotational settings

to some extent. However, the mouse chemical model did not appear to be as

good at discriminating rotationally mispredicted nucleosomes compared with the

yeast models.

Discussion:

We developed a chemical map–based computational method to predict

nucleosome positioning and assessed the prediction results. Training dHMMs

with chemical maps improved the prediction accuracy of nucleosome locations

with respect to the genomic coordinates in budding yeast, fission yeast, and the

house mouse, *Mus musculus* (**Figures 1 and 7**). A total of 41-54% of the

predicted nucleosomes on the Viterbi paths matched those of *in vivo*

nucleosomes at 2-bp resolution. Genome- and locus-level evaluations showed

that the software successfully predicted the positions of *in vivo* nucleosomes

(**Figures 3 and 4, and Additional file 1: Figures S2 and S3**). The *in vivo*

nucleosomes generally had high HBA scores at reasonably near genomic

positions. Furthermore, we demonstrated that perturbation of nucleosome

positioning associated with genetic alterations could also be predicted (**Figures**

4 and 6, and Additional file 1: Figure S4). Thus, we propose that the software

nuCpos can be used for predicting nucleosome positions and also for

engineering of nucleosome-forming sequences.

As expected, the use of chemical maps in prediction led to strict

recognition of rotational settings, which was demonstrated with *in vitro* and *in*

vivo nucleosome-forming sequences (**Figures 1, 2 and 7**). Indeed, chemical

map–based HBA scores for the original Widom 601 sequence indicated that the

147-bp sequence centered at nucleotide position 154 was suitable for

nucleosome formation (**Figure 2**). The *in vitro* nucleosome positions formed on

natural sequences were also predictable at a rotational level (**Additional file 1:**

Figure S1); however, the HBA amplitude along these sequences was smaller

than along the *in vitro*–optimized Widom sequence. In contrast to the chemical

map–based models, the MNase-seq–based models largely failed to recognize

rotational settings (**Figures 1, 2 and 7**). This may have been due to the A/T

preference of MNase and the difficulty of dyad base calling in the construction of

MNase-seq–based nucleosome maps [7, 12]. Indeed, the MNase-map–based

model simply predicted A/T-rich regions as nucleosome-depleted regions

(**Figure 3 and Additional file 1: Figure S3**). Therefore, instead of

lower-resolution MNase-seq–based maps, base-pair-resolution chemical maps

should be used in the development of bioinformatics methods to obtain better

prediction results.

In the field of synthetic biology, the MNase-seq–based method NuPoP

has contributed to the design of synthetic promoters and terminators [29, 30].

However, some studies reported that the histone binding affinity and predicted

nucleosome occupancy output by NuPoP do not correlate with synthetic

promoter and terminator activity [34, 35]. Differences in the DNA sequences of

the tested synthetic elements may of course account for this discrepancy, as

discussed elsewhere [34, 35]. Clearly, the A/T frequency does affect the

prediction results of NuPoP (**Figure 3 and Additional file 1: Figure S3**). As

demonstrated in the Results section, the chemical map–based method is more

accurate and not affected by the A/T frequency. Thus, we expect that nuCpos is

applicable to prediction-based engineering and more effective for synthesizing

functional elements. Similarly, nuCpos can also be used to examine whether

DNA sequences of interest are suitable for nucleosome formation, as previously

done with NuPoP [37, 38].

Dividing 147-bp nucleosomal DNA into subsegments and calculating

subsegment local HBA scores revealed unexpected landscapes of the tested

nucleosomes. Our study demonstrated that the Widom sequence had segments

that are both favored and disfavored for nucleosome formation (**Figure 5A**).

Importantly, segments with high local HBA scores in the Widom 601 sequence

matched the high-affinity histone-contacting regions (**Figure 5B**). Furthermore,

the Widom sequence had segments exhibiting the highest local HBA scores at

the *in vitro* rotational settings (**Figure 5B**). In contrast to the left-half sequence,

the right-half sequence contained segments exhibiting relatively high local HBA

scores that did not change markedly with different rotational settings. Thus,

these sequence-specific features may account for the different behavior of each

half in *in vitro* experiments [44-48]. Collectively, our data indicate that local HBA

scores for given sequences provide insights that enhance understanding of

nucleosome features.

Locus-level assessments of prediction results suggested that software

users should not place undue confidence in the predicted nucleosome

occupancy output from chemical map–based models. Indeed, there was a

misplaced nucleosome that occupied the replication origin *ARS1* (**Figure 3**). We

speculate that these discrepancies were due primarily to the fact that the models

do not consider involvement of functional DNA elements or transacting factors

that regulate nucleosome formation at specific genomic locations. In other words,

chromatin regions highlighted by the differences in nucleosome occupancy

between prediction and *in vivo* observations could be the targets of future

investigations aiming to uncover region-specific regulatory mechanisms.

We demonstrated that the prediction accuracy of the chemical map–

based mouse model (41%) was apparently lower than that of yeast models (over

50%) (**Figures 1 and 7**). A disadvantage of mouse prediction is that the

chemical map probably contains substantial numbers of pseudo-positive

nucleosomes, which could not be omitted due to the lower cleavage depth than

yeasts [22]. As a consequence, the density of nucleosomes on the mouse

genome is one nucleosome per 3 bp, which is about 10 times higher than yeast

values (see **Methods**). Thus, we speculate that the accuracy of mouse

nucleosome positioning prediction can be increased if other chemical maps of

higher quality are produced and used for model construction. In human

genomics, the relationship between various phenotypes and genetic variants

has been explored [50-53]. However, the effects of genetic variants on

nucleosome formation in this regard remain to be studied. Now that our study

showed that insertions and replacements that disrupt nucleosomes in yeast cells

can be predictable *in silico*, it is possible that nucleosome-disrupting human

mutations can be found with the aid of chemical map–based predictions.

Conclusions:

In this study, we demonstrated that the accuracy of dHMM-based nucleosome

positioning prediction can be substantially increased by using

base-pair-resolution nucleosome maps for model construction. Our prediction

results suggest that chemical map–based models are useful for predicting

nucleosome positioning in wild-type and modified sequences at the locus level.

We also demonstrated that strong histone-DNA contacts in a nucleosome and

their rotational settings can be predicted. Furthermore, as another advantage of

bioinformatics methods, our models indicate that the commonly used Widom

sequence contains subnucleosomal segments that are disfavored, or statistically

very rare in *in vivo* nucleosomes, at their nucleosomal positions in shifted

rotational settings. We expect that our prediction method will provide further

insights that will enhance understanding of nucleosome-based epigenetic

regulation.

Methods:

Software and data sets

Most analyses were performed in the GNU R environment

(https://www.r-project.org, ver. 3.6.1). R packages were obtained from CRAN

(https://cran.r-project.org/) and Bioconductor (http://bioconductor.org/). Chemical

maps for budding yeast, fission yeast, and house mouse (*Mus musculus*)

embryonic stem cells [17, 19, 22] were used for model construction and testing

the prediction accuracy. The budding yeast's chemical map in the sacCer2

coordinate was lifted over to the sacCer3 coordinate, as described elsewhere

[27]. The number of unique and redundant nucleosomes in the budding yeast

genome was 67,548 and 344,709, respectively; fission yeast, 75,828 and

425,653; mice, 10,677,016 and 850,701,275. Reference genomes of budding

yeast (R64-1-1), fission yeast (ASM294v2), and mice (mm9/NCBIM37.67) were

used. The original 282-bp Widom 601 sequence [10], the 485-bp MMTV 3'–LTR

sequence [54] and the somatic 5S RNA gene of *Xenopus borealis* [55] were

used as test sequences.

Construction of the nuCpos package

The codes for the construction of parameters used in the nuCpos package (ver.

1.2.0) are available online (**https://doi.org/10.5281/zenodo.3362065**). Genomic

regions covered with the 147-bp non-redundant (unique) chemically mapped

nucleosomes and uncovered were defined as nucleosome and linker regions,

respectively. DNA sequences of these regions were used to construct

parameters that were transferred to internal Fortran programs. Nucleosomes of

which dyads were located within 73 bp of the chromosomal ends were omitted.

For construction of the mouse model, hard-masked genomic sequences were

used, and nucleosome and linker regions containing N were omitted before

parameter construction to avoid potential prediction bias caused by repeat

elements. In total, 67,538 nucleosome regions and 50,622 linker regions were

obtained for the budding yeast genome (sacCer3); fission yeast (ASM294v2),

75,826 nucleosome and 46,557 linker regions; mice (mm9), 4,147,972

nucleosome and 2,484,347 linker regions.

We developed an R function designated *predNuCpos*, which predicts

nucleosome positioning based on a dHMM, as previously proposed by Xi et al.

[16]. Like its ancestral function *predNuPoP* in the NuPoP package

(https://doi.org/doi:10.18129/B9.bioc.NuPoP, ver. 1.34.0), *predNuCpos* receives

a DNA sequence of any length, invokes an internal Fortran program, and

outputs the prediction result either in the working directory or in the working

environment of R. In *predNuCpos*, construction of the dHMM is based on

chemical maps, as described below.

Parameters used in the *predNuCpos* function were constructed

according to the NuPoP paper [16] using the functionalities of the Biostrings

package (https://doi.org/doi:10.18129/B9.bioc.Biostrings, ver. 2.52.0). The

parameters were as follows: *freqL*, one-base frequencies for linker regions; *tranL*,

tranL2, *tranL3*, and *tranL4*, First- to 4th-order transition probabilities for linker

regions, respectively; *freqN4*, four-base frequencies at the first four nucleotide

positions of nucleosome regions; *tranN4*, time-dependent 4th-order transition

probabilities for nucleosome regions; *Pd*, linker length distribution that ranges

from 1 to 500 bp. Linker sequences of 7-500 bp in length were used for linker

model construction, as described elsewhere [16]. *freqL* and *freqN4* were

obtained using the *oligonucleotideFrequency* function of Biostrings; *tranL*,

tranL2, *tranL3*, *tranL4*, and *tranN4* were obtained using the

oligonucleotideTransitions function of Biostrings. Moving average smoothing

using the *SMA* function of the TTR package

(https://cran.r-project.org/package=TTR, ver. 0.23-4) at a 3-bp window was

applied to the 4th-order transition probability parameter *tranN4* and to the linker

length distribution parameter *Pd*. The parameters used in *predNuCpos* were

also used in another function, *mutNuCpos*, which predicts the effect of genetic

alterations on nucleosome positioning.

Xi et al. proposed the HBA score [16], which is also referred to as the

'nucleosome affinity score', as the log likelihood ratio of the probability for a

given 147-bp sequence to be a nucleosome versus a linker. According to their

definition, the HBA score for the 147-bp region x centering at position i (a_i) on a

given genomic sequence is,

$$a_i := log\left[\frac{P_N(x_{i-73},...,x_{i+73})}{G_L(x_{i-73},...,x_{i+73}|147)}\right],$$

where P_N and G_L represent the probability of observing the 147-bp sequence as

a nucleosome or a linker, respectively [16]. The probability of being a

nucleosome is calculated by referring to the parameters *freqN4* and *tranN4*,

which are derived from nucleosomal DNA sequences. Similarly, calculation of

the probability of being a linker is based on linker DNA sequences. As

nucleosomal and linker sequences do not overlap in terms of their genomic

coordinates, negativity of HBA does not directly mean that the tested sequence

is inappropriate for nucleosome formation. The *predNuCpos* function calculates

chemical map–based HBA scores along the input sequence and outputs them

as raw values as its default behavior. We developed an independent function

designated *HBA*, which only calculates the HBA score for a given 147-bp

sequence. The *HBA* function uses the abovementioned chemical parameters for

predNuCpos: *freqL*, *tranL*, *tranL2*, *tranL3*, *tranL4*, *freqN4*, and *tranN4*.

We defined 13 overlapping nucleosomal subsegments, A through M,

and developed a function designated *localHBA* that calculates "local" HBA

scores for each segment. Segment A corresponds to nucleosomal nucleotide

positions 1-21; B, 12-31; C, 22-42; D, 33-52; E, 43-63; F, 54-73; G, 64-84; H,

75-94; I, 85-105; J, 96-115; K, 106-126; L, 117-136; and M, 127-147. Similar to

the calculation of HBA [16], the local HBA score for segment A of the 147-bp

potential nucleosomal region x centering at position i (l_i) on a given genomic

sequence is calculated as,

$$l_i := log \left[\frac{P_N(x_{i-73},...,x_{i-53})}{G_L(x_{i-73},...,x_{i-53}|21)} \right],$$

where the probabilities of observing the 21-bp sequence as segment A of a

nucleosome and a linker are calculated. Local HBA scores for the other

segments are calculated in the same way, except that the considered nucleotide

positions are set appropriately. At the implementation level, four-base frequency

values for the first four nucleotide positions of each segment were prepared:

freqN4SA corresponds to nucleosome positions 1-4; *freqN4SB*, 12-15;

freqN4SC, 22-25; *freqN4SD*, 33-36; *freqN4SE*, 43-46; *freqN4SF*, 54-57;

freqN4SG, 64-67; *freqN4SH*, 75-78, *freqN4SI*, 85-88; *freqN4SJ*, 96-99;

freqN4SK, 106-109; *freqN4SL*, 117-120; and *freqN4SM*, 127-130. These

four-base frequency values were used to calculate the probability of the segment

as that part of nucleosomal DNA as done for HBA calculations [16].

Prediction of nucleosome positioning with nuCpos and NuPoP and

evaluation of the prediction results

The codes for prediction, evaluation, and figure drawing used in this study are

available online (**https://doi.org/10.5281/zenodo.4083950**). Unmasked

reference genomes for the budding yeast, fission yeast, and house mouse (*Mus*

musculus) were used as prediction target sequences. For the mouse

experiments, only the 19th chromosome (Chr19) was used for prediction

because its length (61,342,430 bp) was sufficient for evaluation, at

approximately five-times longer than the total length of yeast chromosomes

(budding yeast, 12,071,326 bp; fission yeast, 12,571,820 bp). Mouse Chr19

contained 249,210 unique and 19,965,481 redundant nucleosomes that had

been determined based on histone H4 S47C-dependent cleavage [22]. The

density of redundant nucleosomes on mouse Chr19 was one per 3.07 bp. With

this density, one in three randomly located nucleosomes in the chromosome

could be counted as a truly predicted nucleosome. Thus, we selected 2,044,747

nucleosomes with the highest NCP scores as true redundant nucleosomes for

the matching experiment. This selection yielded a density of one nucleosome

per 30.0 bp, which was comparable to that of budding yeast and fission yeast

(one nucleosome per 35.0 bp and 29.5 bp, respectively).

The *predNuPoP* function of NuPoP and the *predNuCpos* function of

nuCpos were used for dHMM-based predictions. In order to predict using

budding yeast models, the species arguments were set as "7" for *predNuPoP*

and "sc" for *predNuCpos*. Similarly, fission yeast ("9" and "sp") and mouse ("2"

and "mm") models were specified. We defined any nucleosomes on the Viterbi

path as predicted nucleosomes. Matching of predicted nucleosomes using *in*

vivo chemical nucleosomes was done by widening the genomic coordinates of *in*

vivo nucleosomes to both sides in a step-wise manner from zero to nine base

pairs. For instance, at 1-bp resolution, a predicted nucleosome was regarded as

"matched" when the genomic coordinate of its dyad was equal to that of the

nearest *in vivo* nucleosome. If the dyad of the predicted nucleosome was 1 bp

away from that of the nearest *in vivo* nucleosome, it was regarded as "not

matched."

Chemical map−based HBA scores were calculated using the

predNuCpos function or the *HBA* function of nuCpos, which yield the same

scores. The species argument was set as "sc" for budding yeast sequences and

sequences of *in vitro* reconstitution studies; for the fission yeast sequences, "sp."

The *predNuPoP* function of NuPoP smooths raw HBA scores using a moving

average of a 55-bp window before outputting the processed HBA scores. This

prevents software users from examining the actual HBA score for each genomic

coordinate to determine if the 147-bp sequence of interest is suitable for

nucleosome formation. To overcome this problem, non-smoothed MNase-seq−

based HBA scores were calculated using in-house functions that utilize the

parameters for the *predNuPoP* function of NuPoP. For comparison of *in vivo*

nucleosomes, 5,542 sets of −1 and +1 nucleosomes determined by Chereji et al.

[12] were used for HBA calculations. The 347-bp sequences around these

nucleosomes (from −173 to +173 nucleotide positions with respect to the dyads)

were extracted from the budding yeast genome. The HBA score was calculated

for each possible 147-bp segment along the sequences. A/T-frequency was

calculated using the *letterFrequency* function of Biostrings for each 147-bp

sequence. HBA and A/T-frequency scores were averaged at each coordinate

with respect to the −1 or +1 nucleosomes. Chemical map−based local HBA

scores were calculated using the *localHBA* function of nuCpos. Heatmaps of

local HBA scores were drawn using the *levelplot* function of the rasterVis

package (https://cran.r-project.org/package=rasterVis, ver. 0.46).

List of abbreviations:

dHMM: duration hidden Markov model

HBA: histone binding affinity

LTR: long terminal repeat

MMTV: mouse mammary tumor virus

MNase: micrococcal nuclease

NCP: nucleosome core particle

NDR: nucleosome depleted region

SELEX: Systematic Evolution of Ligands by EXponential enrichment

SHL: superhelical location

Declarations:

Ethics approval and consent to participate:

Not applicable.

Consent for publication:

Not applicable.

Availability of data and materials:

The datasets generated and analyzed during the current study are available
online (see **Methods** for the URLs).
819

820 **Competing interests:**
821 The authors declare that they have no competing interests.
822

823 **Funding:**
824 This study was supported by JSPS KAKENHI Grant Numbers JP18K05556 and
825 JP19H05264.
826

827 **Authors' contributions:**
828 H.K. designed the study, created the software, performed the experiments, and
829 was a major contributor in writing the manuscript. T.U. and M.S. were involved in
830 study design, data validation, and manuscript writing. All authors read and
831 approved the final manuscript.
832

833 **Acknowledgments:**
834 The authors thank Dr. K. Ura for providing the dinucleosome-forming sequence
835 and Dr. Y. Ichikawa for confirming the DNA sequences. The authors also thank
836 Drs. H. Kimura, H. Kurumizaka, J. Nakayama, and Y. Ohkawa for their helpful
837 comments and encouragement.
838

References:
1.   Luger K, Mader AW, Richmond RK, Sargent DF, Richmond TJ: **Crystal**
structure of the nucleosome core particle at 2.8 A resolution. *Nature*
1997, **389:**251-260.
2.   Luger K, Richmond TJ: **DNA binding within the nucleosome core.** *Curr*
Opin Struct Biol 1998, **8:**33-40.
3.   Richmond TJ, Davey CA: **The structure of DNA in the nucleosome**
core. *Nature* 2003, **423:**145-150.
4.   Lai WKM, Pugh BF: **Understanding nucleosome dynamics and their**
links to gene expression and DNA replication. *Nat Rev Mol Cell Biol*
2017, **18:**548-562.

5.   Struhl K, Segal E: **Determinants of nucleosome positioning.** *Nat Struct*
Mol Biol 2013, **20:**267-273.

6.   Lieleg C, Krietenstein N, Walker M, Korber P: **Nucleosome positioning**
in yeasts: methods, maps, and mechanisms. *Chromosoma* 2015,
124:131-151.

7.   Chereji RV, Clark DJ: **Major Determinants of Nucleosome Positioning.**
Biophys J 2018, **114:**2279-2289.

8.   Teif VB: **Nucleosome positioning: resources and tools online.** *Brief*
Bioinform 2016, **17:**745-757.

9.   Liu H, Zhang R, Xiong W, Guan J, Zhuang Z, Zhou S: **A comparative**
evaluation on prediction methods of nucleosome positioning. *Brief*
Bioinform 2014, **15:**1014-1027.

10.   Lowary PT, Widom J: **New DNA sequence rules for high affinity**
binding to histone octamer and sequence-directed nucleosome
positioning. *J Mol Biol* 1998, **276:**19-42.

11.   Pisano S, Galati A, Cacchione S: **Telomeric nucleosomes: forgotten**
players at chromosome ends. *Cell Mol Life Sci* 2008, **65:**3553-3563.

12.   Chereji RV, Ramachandran S, Bryson TD, Henikoff S: **Precise**
genome-wide mapping of single nucleosomes and linkers in vivo.
Genome Biol 2018, **19:**19.

13.   Chereji RV, Bryson TD, Henikoff S: **Quantitative MNase-seq accurately**
maps nucleosome occupancy levels. *Genome Biol* 2019, **20:**198.

14.   West JA, Cook A, Alver BH, Stadtfeld M, Deaton AM, Hochedlinger K,
Park PJ, Tolstorukov MY, Kingston RE: **Nucleosomal occupancy**
changes locally over key regulatory regions during cell
differentiation and reprogramming. *Nat Commun* 2014, **5:**4719.

15.   Mieczkowski J, Cook A, Bowman SK, Mueller B, Alver BH, Kundu S,
Deaton AM, Urban JA, Larschan E, Park PJ, et al: **MNase titration**
reveals differences between nucleosome occupancy and chromatin
accessibility. *Nat Commun* 2016, **7:**11485.

16.   Xi L, Fondufe-Mittendorf Y, Xia L, Flatow J, Widom J, Wang JP:
Predicting nucleosome positioning using a duration Hidden Markov
Model. *BMC Bioinformatics* 2010, **11:**346.

17. Brogaard K, Xi L, Wang JP, Widom J: **A map of nucleosome positions**
in yeast at base-pair resolution. *Nature* 2012, **486:**496-501.
18. Brogaard KR, Xi L, Wang JP, Widom J: **A chemical approach to**
mapping nucleosomes at base pair resolution in yeast. *Methods*
Enzymol 2012, **513:**315-334.
19. Moyle-Heyrman G, Zaichuk T, Xi L, Zhang Q, Uhlenbeck OC, Holmgren
R, Widom J, Wang JP: **Chemical map of Schizosaccharomyces**
pombe reveals species-specific features in nucleosome positioning.
Proc Natl Acad Sci U S A 2013, **110:**20158-20163.
20. Henikoff S, Ramachandran S, Krassovsky K, Bryson TD, Codomo CA,
Brogaard K, Widom J, Wang JP, Henikoff JG: **The budding yeast**
Centromere DNA Element II wraps a stable Cse4 hemisome in either
orientation in vivo. *Elife* 2014, **3:**e01861.
21. Ramachandran S, Zentner GE, Henikoff S: **Asymmetric nucleosomes**
flank promoters in the budding yeast genome. *Genome Res* 2015,
25:381-390.
22. Voong LN, Xi L, Sebeson AC, Xiong B, Wang JP, Wang X: **Insights into**
Nucleosome Organization in Mouse Embryonic Stem Cells through
Chemical Mapping. *Cell* 2016, **167:**1555-1570 e1515.
23. Shimizu M, Roth SY, Szent-Gyorgyi C, Simpson RT: **Nucleosomes are**
positioned with base pair precision adjacent to the alpha 2 operator
in Saccharomyces cerevisiae. *EMBO J* 1991, **10:**3033-3041.
24. Roth SY, Dean A, Simpson RT: **Yeast alpha 2 repressor positions**
nucleosomes in TRP1/ARS1 chromatin. *Mol Cell Biol* 1990,
10:2247-2260.
25. Thoma F, Bergman LW, Simpson RT: **Nuclease digestion of circular**
TRP1ARS1 chromatin reveals positioned nucleosomes separated by
nuclease-sensitive regions. *J Mol Biol* 1984, **177:**715-733.
26. Morohashi N, Yamamoto Y, Kuwana S, Morita W, Shindo H, Mitchell AP,
Shimizu M: **Effect of sequence-directed nucleosome disruption on**
cell-type-specific repression by alpha2/Mcm1 in the yeast genome.
Eukaryot Cell 2006, **5:**1925-1933.

27.   Fuse T, Katsumata K, Morohoshi K, Mukai Y, Ichikawa Y, Kurumizaka H,
916   Yanagida A, Urano T, Kato H, Shimizu M: **Parallel mapping with**
917   **site-directed hydroxyl radicals and micrococcal nuclease reveals**
918   **structural features of positioned nucleosomes in vivo.** *PLoS One*
919   2017, **12:**e0186974.

920   28.   Fuse T, Yanagida A, Shimizu M: **The Yeast Minichromosome System**
921   **Consisting of Highly Positioned Nucleosomes in Vivo.** *Biol Pharm*
922   *Bull* 2019, **42:**289-294.

923   29.   Curran KA, Crook NC, Karim AS, Gupta A, Wagman AM, Alper HS:
924   **Design of synthetic yeast promoters via tuning of nucleosome**
925   **architecture.** *Nat Commun* 2014, **5:**4002.

926   30.   Morse NJ, Gopal MR, Wagner JM, Alper HS: **Yeast Terminator**
927   **Function Can Be Modulated and Designed on the Basis of**
928   **Predictions of Nucleosome Occupancy.** *ACS Synth Biol* 2017,
929   **6:**2086-2095.

930   31.   Wight A, Yang D, Ioshikhes I, Makrigiannis AP: **Nucleosome Presence**
931   **at AML-1 Binding Sites Inversely Correlates with Ly49 Expression:**
932   **Revelations from an Informatics Analysis of Nucleosomes and**
933   **Immune Cell Transcription Factors.** *PLoS Comput Biol* 2016,
934   **12:**e1004894.

935   32.   Creamer KM, Job G, Shanker S, Neale GA, Lin YC, Bartholomew B,
936   Partridge JF: **The Mi-2 homolog Mit1 actively positions nucleosomes**
937   **within heterochromatin to suppress transcription.** *Mol Cell Biol* 2014,
938   **34:**2046-2061.

939   33.   Small EC, Xi L, Wang JP, Widom J, Licht JD: **Single-cell nucleosome**
940   **mapping reveals the molecular basis of gene expression**
941   **heterogeneity.** *Proc Natl Acad Sci U S A* 2014, **111:**E2462-2471.

942   34.   Portela RM, Vogl T, Kniely C, Fischer JE, Oliveira R, Glieder A: **Synthetic**
943   **Core Promoters as Universal Parts for Fine-Tuning Expression in**
944   **Different Yeast Species.** *ACS Synth Biol* 2017, **6:**471-484.

945   35.   MacPherson M, Saka Y: **Short Synthetic Terminators for Assembly of**
946   **Transcription Units in Vitro and Stable Chromosomal Integration in**
947   **Yeast S. cerevisiae.** *ACS Synth Biol* 2017, **6:**130-138.

36.   Mahmoudi T: **The BAF complex and HIV latency.** *Transcription* 2012,
3:171-176.
37.   Rafati H, Parra M, Hakre S, Moshkin Y, Verdin E, Mahmoudi T:
Repressive LTR nucleosome positioning by the BAF complex is
required for HIV latency. *PLoS Biol* 2011, **9:**e1001206.
38.   Gimeno-Valiente F, Riffo-Campos AL, Vallet-Sanchez A, Siscar-Lewin S,
Gambardella V, Tarazona N, Cervantes A, Franco L, Castillo J,
Lopez-Rodas G: **ZNF518B gene up-regulation promotes**
dissemination of tumour cells and is governed by epigenetic
mechanisms in colorectal cancer. *Sci Rep* 2019, **9:**9339.
39.   Lipford JR, Bell SP: **Nucleosomes positioned by ORC facilitate the**
initiation of DNA replication. *Mol Cell* 2001, **7:**21-30.
40.   Eaton ML, Galani K, Kang S, Bell SP, MacAlpine DM: **Conserved**
nucleosome positioning defines replication origins. *Genes Dev* 2010,
24:748-753.
41.   Rodriguez J, Lee L, Lynch B, Tsukiyama T: **Nucleosome occupancy as**
a novel chromatin parameter for replication origin functions.
Genome Res 2017, **27:**269-277.
42.   Gonzalez S, Garcia A, Vazquez E, Serrano R, Sanchez M, Quintales L,
Antequera F: **Nucleosomal signatures impose nucleosome**
positioning in coding and noncoding sequences in the genome.
Genome Res 2016, **26:**1532-1543.
43.   Ichikawa Y, Morohashi N, Nishimura Y, Kurumizaka H, Shimizu M:
Telomeric repeats act as nucleosome-disfavouring sequences in
vivo. *Nucleic Acids Res* 2014, **42:**1541-1552.
44.   Kulaeva OI, Gaykalova DA, Pestov NA, Golovastov VV, Vassylyev DG,
Artsimovitch I, Studitsky VM: **Mechanism of chromatin remodeling and**
recovery during passage of RNA polymerase II. *Nat Struct Mol Biol*
2009, **16:**1272-1278.
45.   Gaykalova DA, Kulaeva OI, Volokh O, Shaytan AK, Hsieh FK,
Kirpichnikov MP, Sokolova OS, Studitsky VM: **Structural analysis of**
nucleosomal barrier to transcription. *Proc Natl Acad Sci U S A* 2015,
112:E5787-5795.

46.   Chua EY, Vasudevan D, Davey GE, Wu B, Davey CA: **The mechanics**
behind DNA sequence-dependent properties of the nucleosome.
Nucleic Acids Res 2012, **40:**6338-6352.
47.   Vasudevan D, Chua EYD, Davey CA: **Crystal structures of**
nucleosome core particles containing the '601' strong positioning
sequence. *J Mol Biol* 2010, **403:**1-10.
48.   Tsunaka Y, Fujiwara Y, Oyama T, Hirose S, Morikawa K: **Integrated**
molecular mechanism directing nucleosome reorganization by
human FACT. *Genes Dev* 2016, **30:**673-686.
49.   Bondarenko VA, Steele LM, Ujvari A, Gaykalova DA, Kulaeva OI,
Polikanov YS, Luse DS, Studitsky VM: **Nucleosomes can form a polar**
barrier to transcript elongation by RNA polymerase II. *Mol Cell* 2006,
24:469-479.
50.   Li X, Kim Y, Tsang EK, Davis JR, Damani FN, Chiang C, Hess GT,
Zappala Z, Strober BJ, Scott AJ, et al: **The impact of rare variation on**
gene expression across tissues. *Nature* 2017, **550:**239-243.
51.   Consortium GT, Laboratory DA, Coordinating Center -Analysis Working G,
Statistical Methods groups-Analysis Working G, Enhancing Gg, Fund
NIHC, Nih/Nci, Nih/Nhgri, Nih/Nimh, Nih/Nida, et al: **Genetic effects on**
gene expression across human tissues. *Nature* 2017, **550:**204-213.
52.   Chakravorty S, Hegde M: **Inferring the effect of genomic variation in**
the new era of genomics. *Hum Mutat* 2018, **39:**756-773.
53.   Chakravorty S, Hegde M: **Gene and Variant Annotation for Mendelian**
Disorders in the Era of Advanced Sequencing Technologies. *Annu*
Rev Genomics Hum Genet 2017, **18:**229-256.
54.   Flaus A, Richmond TJ: **Positioning and stability of nucleosomes on**
MMTV 3'LTR sequences. *J Mol Biol* 1998, **275:**427-441.
55.   Ura K, Hayes JJ, Wolffe AP: **A positive role for nucleosome mobility in**
the transcriptional activity of chromatin templates: restriction by
linker histones. *EMBO J* 1995, **14:**3752-3765.

1011
1012

Figure legends:

Figure 1. Comparison of prediction accuracy between chemical map–

based and MNase-seq–based models for yeasts. (**A**) Distance between

predicted nucleosomes and nearest *in vivo* nucleosomes. Fraction of predicted

nucleosomes on the Viterbi path located at a particular distance from the nearest

in vivo nucleosomes is plotted. Species of the target genome and for model

construction were both *S. cerevisiae*. Magnified view is presented inside each

plot. Arrows point to the fractions of predicted nucleosomes that are 1-bp apart

from *in vivo* nucleosomes. (**B**) Matching rate between predicted and in vivo

nucleosomes with variable matching windows. The species of genomic

sequences used for prediction (Target) and the species of the chemical map

used for model construction (Model) are indicated. "Sc" stands for *S. cerevisiae*,

whereas "Sp" stands for *S. pombe*. Indicated R packages were used for

predictions. (**C**) Matching of nucleosomes on the Viterbi paths with *in vivo* unique

and redundant nucleosomes at 2-bp resolution. Note that the redundant

nucleosome datasets contain both unique nucleosomes (red), which are

representative, and non-representative nucleosomes (orange).

1030

Figure 2. HBA scores along the Widom 601 sequence. HBA scores along the

282-bp original Widom sequence were calculated using chemical map–based

and MNase-seq–based budding yeast models. The scores were normalized by

subtracting the mean value from the raw value on a per sequence basis. The

HBA score for a given 147-bp nucleosome sequence was assigned to its dyad

position. Orange vertical lines indicate the dyad positions of *in vitro* reconstituted

nucleosome: nucleotide position 154. Asterisks indicate high-scoring HBA

positions around the *in vitro* positioning site.

1039

Figure 3. Comparison between chemical map–based and MNase-seq–

based models on selected *in vivo* regions. (**A and B**) Chemical map–based

and MNase-seq–based HBA scores (**A**) and A/T nucleotide frequency (**B**) along

the sequences containing budding yeast −1 and +1 nucleosomes for

protein-coding genes. Gray vertical lines indicate the dyad positions of

respective nucleosomes. Nucleosomes of which the dyad base is located at 0 bp

are shown as ovals above the plots. Asterisks indicate positions with relatively

high HBA scores. NDR: nucleosome-depleted region. (**C**) Prediction results for

the budding yeast *TRP1ARS1* mini-chromosome. Schematic representation of *in*

vivo nucleosome positioning is shown above the plots. Nucleosomes labeled I

through VII are indicated as ovals. Note that this sequence is circularized *in vivo*

by being linked at the *Eco*RI sites. The top two panels show the prediction

results output by nuCpos, whereas the next two panels show NuPoP results.

The upper panel in each set shows predicted occupancy of the nucleosome

(Occup., gray polygons) and probabilities of the tested 147-bp sequences for

being in the nucleosome state (P-dyad, blue vertical lines). The lower panels

show HBA values for the tested 147-bp sequences calculated using the

indicated models. The very bottom panel shows the A/T-frequencies for the

tested 147-bp sequences. Horizontal lines at the bottom of each plot indicate a

unit of the circular *TRP1ARS1* mini-chromosome (colored in gray, nucleotide

positions 0 to 1,464), the coding region of *TRP1* (red, 115-879), and the *B3*

(black, 763-788), *B2* (purple, 810-820), *B1* (green, 847-859), and *ACS* (blue,

869-879) elements of *ARS1*. Inverted triangles indicate the histone H4

S47C-dependent cleavage centers determined by indirect end-labeling; orange

for mini-chromosome and light blue for genomic experiments [27].

1065

Figure 4. Predicting the effects of repeat insertion on nucleosome

positioning. Chemical map–based HBA scores and predicted nucleosomal

occupancy were calculated for original and modified *BAR1* promoter sequences.

The sequences were centered at the dyad position of the repressive

nucleosome, in which the indicated repeat sequences were inserted. Horizontal

lines at the bottom of each plot indicate the *α2*-operator (purple), the *TATA*

(blue) element, the coding region of *BAR1* (orange), and the insert (red). Note:

the sequence insertion caused a shift of the left of the three predicted

nucleosomes to the right to cover the *α2*-operator in the occupancy plots.

Overconfidence should not be placed in this shift, as it may simply reflect that the

dHMM tends to follow the linker length distribution, which generally does not

allow longer linkers.

1078

Figure 5. Local HBA analysis of the Widom 601 sequence. (**A**) Chemical

map–based local HBA scores along the 282-bp Widom 601 sequence. The

scores for each tested 147-bp sequence were heatmapped at the center position

of that sequence. The heatmap scale indicates the local HBA score. Black

arrows indicate high-HBA-score positions (124, 135, 143, 164, and 175). Red

arrow indicates the dyad position of the *in vitro* reconstituted nucleosome (154).

The region evaluated in (**B**) is indicated as a horizontal gray bar under the X-axis.

(**B**) Local HBA scores at (red) and around (black) the *in vitro* translational

position for each segment are plotted in columns. The scores for the M segment

are magnified to the left to show the indices on the X-axis. The relative positions

−4, 0, and +4 correspond to nucleotide positions 150, 154, and 158 of the

282-bp sequence. The location of the R3 element is indicated at the top. (**C**)

Schematic representation of shifting of translational positions in (**B**). Nucleotide

positions relative to the dyad of the central translated position (i=154) are

indicated. Only three (−4, 0, and +4) of the nine positions evaluated in (**B**) are

shown.

1095

Figure 6. Local HBA analysis of modified nucleosome-forming sequences.

Chemical map–based local HBA scores along the original and modified fission

yeast *ura4+* gene. The scores for each tested 147-bp sequence were

heatmapped at the center position of that sequence. The heatmap scale

indicates the local HBA score. Black arrows indicate the nucleosome centers

(nucleotide positions +17, +164, +332, +470, +629, and +806) determined by

MNase-seq [42].

1103

1104 **Figure 7. Comparison of prediction accuracy between chemical map–**

1105 **based and MNase-seq–based models for mice.** (**A**) Matching of nucleosomes

1106 on the Viterbi paths with *in vivo* unique and redundant nucleosomes. (**B**)

1107 Matching rate between predicted and *in vivo* nucleosomes with variable

1108 matching windows. The sequence of mouse Chr19 was used for prediction.

1109

1110 **Table 1. Functions of the software packages used in this study.**

NuPoP [16]	nuCpos	Description
*predNuPoP**	*predNuCpos***	Prediction of nucleosome positioning based on dHMMs
readNuPoP		Loading of prediction results
plotNuPoP		Plotting of prediction results
	HBA	Calculation of HBA scores
	localHBA	Calculation of local HBA scores
	mutNuCpos	Prediction of the effects of genetic alterations on nucleosome positioning

1111   *MNase-seq–based. HBA scores are smoothed before outputting the prediction

1112   results. **Chemical map–based. Smoothing of HBA scores is optional.

1113

Supplementary Figure Legends:

Additional file 1: Figure S1. HBA scores along *in vitro* nucleosome-forming

sequences. HBA scores along nucleosome-forming sequences were calculated

using chemical map–based and MNase-seq–based budding yeast models.

Sequences analyzed were the 3'-LTR of MMTV (**A**) and the *Xenopus borealis*

5S rDNA dinucleosome-forming sequence (**B**). The scores were normalized by

subtracting the mean value from the raw value on a per sequence basis. The

HBA score for a given 147-bp nucleosome sequence was assigned to its dyad

position. Orange vertical lines indicate the dyad positions of *in vitro* reconstituted

nucleosomes: nucleotide positions 139 and 335 (**A**) and 106 and 303 (**B**).

Asterisks indicate high-scoring HBA positions around the *in vitro* positioning

sites. Note that for the 5S rDNA sequence, two identical sequences are joined at

position 230, causing a difference in HBA scores around the left sides of the two

in vitro nucleosome positions.

1128

1129   **Additional file 1: Figure S2. Magnified view of prediction results for**

1130   ***TRP1ARS1.*** Prediction results for the budding yeast *TRP1ARS1*

1131   mini-chromosome output by nuCpos and NuPoP (**Figure 3C**) were magnified,

1132   being centered at nucleosome II.

1133

Additional file 1: Figure S3. Prediction results for the fission yeast *ura4*+

gene. Schematic representation of *in vivo* nucleosome positioning is shown

above the plots. Nucleosomes numbered +1 to +6 are indicated as ovals. The

top two panels show the prediction results output by nuCpos, whereas the next

two panels show NuPoP results. The upper panel in each set shows predicted

occupancy of nucleosomes (Occup., gray polygons) and probability that the

tested 147-bp sequences are in the nucleosome state (P-dyad, blue vertical

lines). Lower panels show HBA values for the tested 147-bp sequences

calculated using the indicated models. The very bottom panel shows the

A/T-frequency for the tested 147-bp sequences. Horizontal lines at the bottom of

each plot indicate the 5'- and 3'-untranslated regions (colored in gray, nucleotide

positions −151 to −1 and +795 to +986, respectively) and the protein-coding

region (red, 0 to +794) of the *ura4*+ gene. Inverted triangles indicate the

nucleosome centers determined by MNase-seq [42].

1148

Additional file 1: Figure S4. Prediction of the effects of repeat insertion in

the fourth nucleosome of the *TALS* minichromosome. Chemical map–based

HBA scores and predicted nucleosomal occupancy were calculated for original

and modified *TALS* sequences [43]. The sequences were centered at the dyad

position of the fourth nucleosome (0 bp), in which the indicated repeat

sequences were inserted. Note: the position for the fourth nucleosome was

predicted to shift to the right. Horizontal lines at the bottom of each plot indicate

the *α2*-operator (purple) and the insert (red). hTEL stands for human telomeric

repeat (5'-TTAGGG-3'); SI-A, sequence isomer-A (5'-TGTAGG-3'); SI-B,

sequence isomer-B (5'-TGTGAG-3'). When telomeric DNA fragments are

inserted at the center of the fourth nucleosome, shorter fragments (hTEL2 and

hTEL4) were predicted to not affect nucleosome positioning. In contrast, longer

fragments (hTEL12 and hTEL29) were predicted to severely compromise

nucleosome formation. In addition, sequence isomers of the telomeric repeat

(SI-A6, SI-A12, SI-B6, and SI-B12) were predicted to not inhibit nucleosome

formation. These prediction results agreed with previous *in vivo* observations

[43]. The hTEL6 insertion, which causes nucleosome depletion *in vivo*,

appeared not to affect nucleosome formation in the prediction, suggesting a

limitation of the prediction.

1168

Additional file 1: Figure S5. Thirteen overlapping nucleosomal DNA

subsegments for local HBA calculation. Segments A through M are colored in

orange on the nucleosome structure 1AOI [1]. The structures were drawn using

MacPyMOL (v1.8.4.1). The base pair length (red) and corresponding

superhelical locations (black) of each segment are indicated. The color of the

right-half segments (H-M) is faded because they are behind the front helix.

1175

1176

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

B

C

Figure 6

Figure 7

A

B