Antioxidant and antimicrobial activites on uropathogenic bacteria of Asteraceae medicinal plants

Trinh Canh Phan[1], Thao Thi Thanh Le[1], Ha Tran Viet Hoang[1] , TuAnh Nguyen[1]

[1] Department of Microbiology and Parasitology, Faculty of Pharmacy, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh 700000, Vietnam.

Correspondence should be addressed to Trinh C. Phan; phancanhtrinh@ump.edu.vn

Abstract

Antimicrobial and DPPH-scavenging assay were analyzed on nine Asteraceae species collected from Southest in Vietnam. Crude extracts from *Taraxacum officinale, Chrysanthemum morifolium, Ageratum conyzoides, Tagetes erecta* and essential oils from *Ageratum conyzoides, Helianthus annuus, Artemisia vulgaris* have antimicrobial activity on tested microorganisms. The essential oils of displayed effect on *Candida* spp. and Gram possitive bacteria, meanwhile, crude extracts expressed the inhibitory ability on bacteria. Ethyl acetate fraction from *Ageratum conyzoides* showed the most potent effect on urinary clinical strains (*E. coli* and *K. pneumoniae)* with MIC at 1.25-10 mg/ml and 5-12.5 mg/ml, respectively. All ethanol crude extracts exhibited strong reducing power indicating on the TLC plate. The results of the DPPH scavenging assay showed that *Tagetes erecta* extract had the lowest IC_{50} (17.280 μg/ml) and 2.5 times higher than vitamin C (7.321 μg/ml). This study revealed that *A. conyzoides* has good potential against uropathogenic *E. coli* and *K. pneumoniae*, and could therefore be applicable of prophylactic urinary tract infection.

1. Introduction

In recent years, the antibiotic resistance has become more complex, putting mankind into the post-antibiotic period. Many clinical *Enterobacteriaceae* strains such as *Escherichia coli* and *Klebsiella pneumoniae* have extended-spectrum beta-lactamase (ESBL) and carbapenem resistant Enterobacteriaceae (CRE) [1-3]. It is the important problem for both microbiologists and clinicians. Especially, polymyxin plasmid-mediated resistance gene (mcr-1) was observed in *E. coli* strains, which was isolated from animals and inpatients with infection during 2011-2014, in China [4]. Moreover, it is with deep concern that mcr-1 could be transferred to *K. pneumoniae* and *Pseudomonas aeruginosa* via transformation [4]. In 2016, the first report about mcr-1 gene in a patient with urinary infections in Pennsylvania, the United States was shown by Jennifer Abbasi [5]. In urinary infections, many reports show that *E. coli*, *K. pneumoniae* is the leading causative agents. In addition, *Staphylococcus saprophyticus, Enterococcus faecalis,* group B *Streptococcus, Proteus mirabitis, Pseudomonas aeruginosa, Staphylococcus aureus* and *Candida* spp. is also aetiology [6-9].

Herbal extracts and their essential oils have used as foods such as floral teas, beverages, functional foods, bakery products; and traditional medicines in many years, with minimal known "side-effects" on human health. Using herbal remedies might help to reduce dependence on antibiotic therapies and minimizing antibiotic resistance phenomenon [10].

The Asteraceae family (Compositae) is a widespread family of flowering plants, including 32,913 species names, belonging to 1,911 plant genera, distributed in 13 subfamilies [11]. The tropics, subtropics and temperate regions is the natural habitats of Asteraceae species [12]. As chemical components, they usually contain a large amount of essential oils, polyphenols, flavonoid compounds, which are often studied for antimicrobial and antioxidant activities [13-18].

Although there were many reports for antimicrobial and antioxidant effects of Asteraceae species, applications of these extracts in treating infectious diseases need an assessment on pathogenic bacteria strains isolated from clinical [10]. In the study, we had been screening antimicrobial and antioxidant activities of ethanol extracts and essential oils from 9 species of Asteraceae on 30 clinical strains causing urinary tract infection, collected from District 2 Hospital, Ho Chi Minh city, Vietnam. The target is seeking the best extract to apply for a healthcare serum to prevent recurrence UIT. Antioxidant activity might be a protection factor for urinary tract epithelium to avoid impact of oxidative stress.

2. Materials and Methods

We conducted the investigations antimicrobial activities of nine Asteraceae species collected from Southern in Vietnam following Figure 1. In particular, after pre-treating and extracting herbal samples to obtain crude extracts and essential oils, we evaluated antimicrobial effect applied diffusion method. The extracts which show activity were fractionated by different polar solvents; n - hexane, chloroform, ethyl acetate. The active fractions, which had been determined antimicrobial capacity by well agar diffusion method, were carried out MIC and MBC assays. As for essential oils, after antimicrobial screening assay, the active oils were evaluated MIC and MBC values. Simultaneously, antioxidant activity was also determined by DPPH free-radical scavenging assay.

Figure 1: Schematic representation of experimental layout

2.1. Plant Authentication and Preparation

Asteraceae-plant samples were collected from the Southern in Vietnam during March to May 2016. These samples were identification at the Botany Department, Faculty of Pharmacy, University of Medicine and Pharmacy at Ho Chi Minh city. These species are usually used as traditional medicines in Vietnam. Table 1 shows common name, traditional uses following the botanical nomenclature of nine species used in this study.

After harvesting, the samples were washed under running water to remove dust and rinse with distilled water to drain. Then they were dried in the shade. Afterwards, the dried plant materials were

fine grounded by mechanical grinders. The powder was stored in tightly closed glass containers in the dark at room temperature.

Table 1: The investigated plants

Nomenclature	Common name	Traditional uses*	Part tested[†]
Ageratum conyzoides	Billygoat-weed	Sinusitis, anti-inflammation.	Aerial parts, essential oils
Artemisia vulgaris	Mugwort	Skin ailments, wounds, ulcers.	Aerial parts, essential oils
Chrysanthemum coronarium	Crown daisy	Pain relief, fever, dysentery.	Aerial parts
Chrysanthemum morifolium	Florists chrysanthemum	Pimples, dermatitis, fevers.	Flowers
Helianthus annuus	Sunflower	Anti-inflammatory, malaria.	Flowers, essential oils
Tagetes erecta	Mexican marigold	Dysentery, asthma, ulcer.	Flowers
Taraxacum officinale	Dandelion	Hepatitis, bronchitis, pneumonia.	Aerial parts
Vernonia amygdalina	Bitter leaf	Fever, measles, parasites.	Leaves
Wedelia trilobata	Wedelia	Fever, infection.	Aerial parts

In this column there are presented only such traditional uses which can imply the presence of antimicrobial compounds. [†] Part tested: the part of plant used in this study.

2.2. Preparation of plant extracts

2.2.1. Preparation of essential oils

Plant samples after being washed, were cut into small pieces. Plant materials (100 g) were placed in a flask (1 L) together with distilled water. A continuous steam distillation extraction head was attached to the flask. After steam distillation (about 3 hours), the oil was isolated and dried over anhydrous sodium sulfate (Merck). The essential oils were used directly for antimicrobial assay.

2.2.2. Ethanol crude extracts

Plant materials (50 g) were extracted by cold soaking with 500 ml of 96% ethanol (Xilong Chemical) for 24 hours, at 10:1 solvent-to-sample ratio (v/w). Then, the mixtures were filtered through Whatman filter paper. The extracts were allowed to evaporate at a temperature of 45-50 °C. These step were repeated three times in order to achieve maximal extraction of compounds. Dried crude extracts were weighed and kept at -35 °C till further use (not more than one month). These extracts had been screened antimicrobial activity with well agar diffusion method to choose the good antimicrobial extracts for the next step.

2.2.3. Fractionation of the ethanol crude extracts

Ethanol crude extracts, which possess the strong antimicrobial effect, were carried out liquid/liquid extraction with *n* – hexane, chloroform and ethyl acetate, respectively. After evaporating solvent, each fractions was determined the antimicrobial activity.
Stock solutions of crude extracts and fractions were prepared at a concentration of 100 mg/ml, in 10% dimethyl sulfoxide (DMSO - Merck).

2.3. Phytochemical preliminary

The ethanol crude extracts had been analyzed phytochemical constituents for identification of various classes of compounds according to Maria, et al. [19].

2.4. Microorganism strains and culture conditions

Many bacterial and fungal strains from American Type Culture Collection (ATCC) were used in this study for preliminary antimicrobial assays, which included methicillin-sensitive *Staphylococcus*

aureus ATCC 25923 (SA), methicillin-resistant *S. aureus* ATCC 33591 (SR), *Enterococcus faecalis* ATCC 29212 (EF), *Escherichia coli* ATCC 25922 (EC), *Klebsiella pneumoniae* ATCC 35657 (KP), *Pseudomonas aeruginosa* ATCC 27853 (PA), *Candida albicans* ATCC 10231 (CA). Two clinical *non-albicans* strains being *Candida glabrata* ND31 (CG), *Candida tropicalis* PNT20 (CT) which were provided by Nguyen Dinh Nga et al. (2012) [20]. Clinical strains (15 *E. coli* and 15 *K. pneumoniae),* which were isolated from District 2 Hospital, Ho Chi Minh city, Vietnam, in 2016, were also applied for antimicrobial investigation on the potential extracts.

These strains were preserved in 25% glycerol at -80 °C. One strain tube thawed rapidly at 37 °C and cultured in 10 ml Brain heart infusion (BHI - Merck) at 37 °C over 24 hours. The bacteria were streaked on BHI agar (BHA - Merck) at 37 °C over 24 hours. One to five colonies were used to prepare bacterial suspension to match a 0.5 McFarland standard (1-1.5 x 10^8 CFU/ml). Mueller-Hinton agar medium (MHA - Merck) and MHA supplied 2% glucose medium (MHGA - Merck) were used for determination antibacterial and antifungal activity, respectively.

2.5. Antimicrobial diffusion method

The antimicrobial activity of the samples were initially evaluated by well agar diffusion assay for the crude extracts and disc diffusion assay for the essential oils [21]. Growth medium was poured into Petri dishes at 45-50 °C, approximately 4 mm depth, and they were left to solidify in laminar-flow hood. Subsequently, a sterile cotton swab was dipped into overnight bacterial or yeast suspensions of indicator strains (adjusted to turbidity of 0.5 McFarland standard). An agar plate was inoculated by evenly streaking cotton swab over the agar medium. And then follows wells with a diameter of 6 mm were cut in the medium with a sterile cork borer. Stock solutions of the samples were diluted in sterile distilled water to get 100 mg/ml concentration. The tested samples and controls (50 µl) were dispensed into the wells. 20 µl essential oils were applied on a 6 mm disc, then applied discs to agar surface. The plates were incubated at 37 °C for 24-48 hours. Then the diameters of growth inhibition zones were measured. Following control agents were used: positive control agents - ampicillin 20 µg/ml (for bacteria) and ketoconazole 20 µg/ml (for yeasts); negative control agent - 10% DMSO.

2.6. Determination of Minimum Inhibitory Concentration

Determination of minimum inhibitory concentrations (MIC) of the extracts and essential oils were done using agar dilution method [21]. Stock solutions were diluted with melted agar to a concentration range so that the following concentration equal to half the previous concentration. Pour the agar to Petri discs and wait for them to solidify in laminar-flow hood. Microorganism suspensions at 0.5 McFarland were diluted by 0.85% NaCl solution to reach 1-1,5 x 10^7 CFU/ml for bacteria and 1-5 x 10^6 CFU/ml for *Candida* spp.. These suspensions were spotted (1 µl) on the agar surface. Bacterial or yeast colonies growth at the spot after incubating at 37 °C for 24-48 hours indicated for the assay. The MIC is the lowest concentration of antimicrobial agent that completely inhibits growth. The experiment was replicated three times.

2.7. Determination of Minimum Bactericidal Concentration

In order to determine the minimal bactericidal concentration (MBC), the spots at MIC, 2MIC, 4MIC, and 8MIC were washed with 1 ml of 0.85% NaCl. The 100 µl of the washing suspension was spread evenly over BHA agar. After 24-48 hours of incubation at 37 ºC, the number of surviving bacteria was determined. The MBC was defined as the lowest extract concentration at which 99.9% of the bacteria have been killed compare with the microbial density which was used in MIC assay quantified before. The experiment was replicated three times.

2.8. DPPH free-radical scavenging activity assay

The 0.25 mg/ml 2,2-diphenyl-1-picryl-hydrazyl-hydrate (DPPH – Sigma Aldrich) solution in methanol (working solution) was used to determine the antioxidant capacity. The stock solution was prepared at a sample concentration of 10 mg/ml. The preliminary test was carried out on TLC Silica gel 60 F_{254} (Merck). After impregnating 2 µl/spot of stock solution on the TLC plate, the plate was dipped into DPPH working solution and incubated at the room temperature 30 minutes. The positive test indicated the yellow on the violet background [21].

Reaction mixtures consisted of stock solution, 2 ml DPPH working solution and methanol as solvent to have a sample concentration from 0.1 to 0.5 mg/ml. The negative controls had only solvent instead of test solution. The mixtures were incubated for 30 minutes at 37 ºC in dark. The decrease in the absorbance at 517 nm was measured (A_E) [21]. The experiment was carried out in triplicate. Samples and positive control ascorbic acid were tested in triplicate over the same range of sample concentrations. Radical scavenging activity was calculated using the following formula:

$$SC \% = 100\% \text{ x } (A_B\text{-}A_E)/A_B \text{ (\%)}.$$

where A_B: absorbance of the blank sample, and A_E: absorbance of the plant extract.

The antioxidant activity was expressed as the IC_{50} value. This value was determined from the plotted graphs of scavenging activity against the concentration of the sample.

3. Results and Discussion
3.1. Results
3.1.1. Phytochemical preliminary

The Table 2 depicts various classes of phytoconstituents presenting in testing herbs. In general, tannins, flavonoids, phenolics were found in all testing herbs. *A. conyzoides*, *H. annuus,* and *A. vulgaris* possess the large amount of essential oils.

Table 2: Phytochemical profile of ethanol crude extracts

Group of compounds	TO	CM	AC	CC	HA	VA	AV	TE	WT
Carotenoid	-	-	-	-	-	-	-	-	-
Free triterpenoids	+	-	-	-	-	-	-	+	-
Alkaloids	-	-	-	-	+	+	+	+	-
Coumarins	-	-	-	-	-	-	-	-	-
Anthraglycosides	+	-	-	-	-	+	-	-	-
Flavonoids	++	+++	++	+	++	+	+	+++	+
Heart glycolysis	-	-	-	-	-	-	-	-	-
Tannins	+++	++	+++	+	++	++	++	+++	++
Phenolics	+++	+++	+++	++	++	++	++	+++	+
Saponins	+	+	-	+	-	+	+	+	+
Organic acids	-	-	-	-	-	-	-	-	-
Reducing agent	+	+	+	+	+	+	+	+	+

TO: *T. officinale*; CM: *C. morifolium*; AC: *A. conyzoides*; CC: *C. coronarium*; HA: *H. annuus*; VA: *V. amygdalina*; AV: *A. vulgaris*; TE: *T. erecta*; WT: *W. trilobata*.

3.1.2. Antimicrobial screening assay

Among ethanol crude extracts tested, there were four plant species included *Ageratum conyzoides*, *Chrysanthemum morifolium*, *Tagetes erecta*, and *Taraxacum officinale* indicated the antibacterial activities from one to four standard bacteria strains. None of them had the anti-yeast activity on three yeast strains listed in Table 3. To be more specific, *A. conyzoides* and *T. erecta* expressed the broad-spectrum antimicrobial effect on Gram positive (20-22 mm on MSSA and MRSA) and Gram negative bacteria (11-18 mm), meanwhile *T. erecta* the inhibitory capacity on *P. aeruginosa* (13 mm). *C. morifolium* and *T. officinale* indicated the activity only on methicillin-sensitive *S. aureus*.

Table 3: The inhibitory zone diameter (IZD, mm) of the ethanol crude plant extracts determined by agar well diffusion assay

Plant species	Inhibitory Zone Diameter (mm)								
	SA	SR	EF	KP	EC	PA	CA	CG	CT
A. conyzoides	22	20	-	18	11	-	-	-	-
T. erecta	21	20	-	16	-	13	-	-	-
C. morifolium	14	-	-	-	-	-	-	-	-
T. officinale	10	-	-	-	-	-	-	-	-
A. vulgaris	-	-	-	-	-	-	-	-	-
C. coronarium	-	-	-	-	-	-	-	-	-
H. annuus	-	-	-	-	-	-	-	-	-
V. amygdalina	-	-	-	-	-	-	-	-	-
W. trilobata	-	-	-	-	-	-	-	-	-

- no inhibitory zone

The ethanol crude extracts, which showed antimicrobial activity (active extracts), were decanted with different polarization solvents included *n* - hexane, chloroform, ethyl acetate to obtain fractions. Diffusion method was conducted to determine antimicrobial activity of these fractions on microbial strains. Noticeably, antimicrobial agents usually distribute in moderate polarity fractions such as chloroform and ethyl acetate (Table 4). The ethyl acetate fraction of *A. conyzoides* displayed the IZD on MSSA (23 mm), MRSA (21 mm), *K. pneumoniae* (15 mm), and *E. coli* (14 mm). Similarly, *T. erecta*'s fraction of ethyl acetate shows IZD from 11-18 mm on MSSA, MRSA, *K. pneumoniae* and *P. aeruginosa*.

Table 4: IZD (mm) of the fractions determined by well agar diffusion assay

Plant species	Part tested	Fractions	IZD (mm)				
			SA	SR	KP	EC	PA
A. conyzoides	Aerial parts	*n* – hexane	-	-	-	-	-
		CHCl₃	11	11	-	-	-
		EtOAc	23	21	15	14	-
		EtOH	9	-	10	-	-
T. erecta	Flower	*n* – hexane	11	-	11	-	-
		CHCl₃	11	-	10	-	-
		EtOAc	17	12	18	-	11
		EtOH	-	-	-	-	-
C. morifolium	Flower	*n* – hexane	-	-	-	-	-
		CHCl₃	13	-	-	-	-
		EtOAc	14	-	-	-	-
		EtOH	-	-	-	-	-
T. officinale	Aerial parts	*n* – hexane	-	-	-	-	-
		CHCl₃	-	-	-	-	-
		EtOAc	11	-	-	-	-
		EtOH	-	-	-	-	-

- no inhibitory zone

The essential oils of *A. conyzoides* (aerial parts)*, A. vulgaris* (aerial parts)*,* and *H. annuus* (flowers) were demonstrated the antimicrobial activity (Table 5). Although, ethanol crude extracts of *A. vulgaris* and *H. annuus* do not have inhibitory capacity on testing microorganisms, this test reveals the effect of their essential oils on *S. aureus* and *Candida* spp.

Table 5: Antimicrobial activity of the essential oils determined by disc diffusion assay

Essential oils	IZD (mm)								
	SA	SR	EF	KP	EC	PA	CA	CG	CT
A. conyzoides	15	-	-	-	-	-	-	8	10
A. vulgaris	23	10	-	-	-	-	14	15	16
H. annuus	10	-	-	-	-	-	-	-	11

- no inhibitory zone

3.1.3. MIC values of active extracts and essential oils on standard strains

MIC and MBC values of the selected four plant parts: *Ageratum conyzoides*, *Chrysanthemum morifolium*, *Tagetes erecta*, and *Taraxacum officinale* crude extracted and four fractions were determined (Table 6).

Table 6: MIC and MBC values (mg/ml) of crude extracts and fractions of four selected plant materials

Plant species	Part tested	Fractions	MIC/MBC (mg/ml)				
			SA	SR	KP	EC	PA
A. conyzoides	Aerial parts	Cru.Ext	2.5/2.5	10/20	5/5	25/25	-
		CHCl₃	20/40	20/40	-	-	-
		EtOAc	2.5/5	5/10	2.5/2.5	10/10	-
		EtOH	20/20	-	10/20	-	-
C. morifolium	Flower	Cru.Ext	30/30	-	-	-	-
		CHCl₃	15/30	-	-	-	-
		EtOAc	15/15	-	-	-	-
T. erecta	Flower	Cru.Ext	0.78/1.56	3.13/6.25	1.25/1.25	-	2.5/2.5
		n – hexan	1.56/3.12	-	5/5	-	-
		CHCl₃	1.56/3.12	-	5/5	-	-
		EtOAc	3.13/3.13	6.25/12.5	2.5/2.5	-	5/5
T. officinale	Aerial parts	Cru.Ext	20/20	-	-	-	-
		EtOAc	10/10	-	-	-	-

Cru.Ext: ethanol crude extract; - no inhibitory zone.

MIC and MBC values of the three essential oils, *A. conyzoides, A. vulgaris,* and *H. annuus,* were show in Table 7.

Table 7: Determination of minimum inhibitory concentration (MIC) of essential oils

Essential oils	MIC/MBC (µl/ml)				
	SA	SR	CA	CG	CT
A. conyzoides	3.75/7.50	-	-	7.5/15	10/20
A. vulgaris	2.50/2.50	6.25/6.25	6.25/12.5	5/10	5/10
H. annuus	3.75/7.50	-	-	-	10/20

3.1.4. Antimicrobial activity on uropathogenic strains of *E. coli* and *K. pneumoniae*

Through these results, we found the anti-Gram-negative-bacteria effect of extracts from *A. conyzoides* and *T. erecta* on *K. pneumoniae* and *E. coli* being 2 main pathogenic bacteria of urinary tract infection. In order to seek the best extract to prevent recurrence UIT, we evaluated the crude extracts and fractions of *A. conyzoides* and *T. erecta* on 15 *E. coli* and 15 *K. pneumoniae* strains isolated from urine specimens at District 2 Hospital, Ho Chi Minh city.

The well agar diffusion assay was used for analysing the activity of the investigated extracts. The results showed that only the ethyl acetate fraction from *A. conyzoides* had antibacterial activity against tested strains. MIC values of the ethyl acetate fraction of *A. conyzoides* was determined by agar dilution assay on the strains of *E. coli* and *K. pneumoniae* that had minimum and maximum diameter of growth inhibition zone as well as on ESBL-producing strains. The diameter of inhibition zones and minimum inhibitory concentration (MIC) showed in Table 8. Specifically, the IZD on *E. coli* and *K. pneumoniae* indicated from 10.83 mm (E3) to 23.72 mm (E72), and from 9.33 mm (K18) to 19.73 mm (K17), respectively. *A. conyzoid*'s ethyl acetate fraction displayed MIC values on E3, E72, K17, and K18 being 12.5, 5, 10, and 1.25 mg/ml, respectively. ESBL-producing strains, E68 and K26 expressed MIC 6.25 and 2.5 mg/ml, respectively.

Table 8. Antimicrobial activity of the ethyl acetate fraction of *A. conyzoides*
determined by agar well diffusion assay

Strains	IZD (mm)	Strains	IZD (mm)
E3	10.83	K18	9.33
E7	11.43	K27	10.03

E25	12.17	K19	11.23
E36	13.33	K20	11.37
E77	13.43	K15	11.67
E38	13.67	K23	11.83
E27	14.47	K14	11.93
E39	14.47	K21	12.10
E84	14.97	K28	12.37
E68*	15.43	K25	13.23
E63	20.37	K22	13.27
E51	21.40	K16	14.20
E42	22.37	K29	17.67
E94	22.37	K26*	19.27
E72	23.27	K17	19.73

ESBL-producing strain

3.1.5. Free radical scavenging activity

DPPH screening test showed the good antioxidant effect of all crude extracts and essential oils. Free radical scavenging activity of total ethanol extracts was quantitatively determined using a DPPH assay. IC_{50} value represents the concentration of tested extract where the inhibition of test activity reached 50% (Figure 2). The results were graphed by Microsoft Excel 2016. The IC_{50} values of essential oils were too large (3000 - 6000 µg/ml), which were outside the linear range.

T. erecta shows the significant scavenging effect for free radical with IC_{50} = 17.3 µg/ml, 2.4 folds comparing to ascorbic acid.

Figure 2: (A) DPPH screening test on TLC and IC_{50} values. (B) The plotted graphs of scavenging activity against the concentration of the ethanol crude extracts and ascorbic acid

3.2. Discussions

Among the tested extracts and essential oils, there were 4 crude extracts (*Ageratum conyzoides*, *Chrysanthemum morifolium*, *Tagetes erecta*, and *Taraxacum officinale)* and 3 essential oils (*A. conyzoides, A. vulgaris,* and *H. annuus*) that had the antimicrobial activity against 9 bacteria and 3 yeast strains. While the crude extracts only had effect on bacteria, the essential oils had effect on both bacteria and yeast strains. There are many reports about antiseptic, antimicrobial, antioxidant, and insecticidal activities of essential oils [22]. Chemical constituents represent in essential oils usually drived from terpenes, phenolic compounds, and aromatic or aliphatic acid esters, which can partition into the lipids of bacterial and mictochondrial membrane resulting in disturbing the cell

structures. The death of cell caused by leakage of a large amount of essential molecules and ions from the bacterial cell [23].

Tagetes erecta flower extract had the most potent effect on bacteria including MSSA, MRSA, *K. pneumoniae* and *P. aeruginosa* with the lowest MIC compare to other extracts. Padalia and Chanda [24] reported antimicrobial activities of aceton extract of *T. erecta* on *K. pneumoniae* and *P. aeruginosa* with MIC values 62 µg/ml and 78 µg/ml, respectively [24].

Ageratum conyzoides aerial part extracts had effect on MSSA, MRSA, *K. pneumoniae,* and *E. coli,* but the MIC values were higher than *T. erecta* extracts. Amadi, et al. [25] demonstrated ethanol extract show the significant inhibitory zone on *Streptococcus mutans*. Following the report of Akinyemi, et al. [26], ethanol extract from *A. conyzoides* indicated MIC and MBC on MRSA of 43 µg/ml and 63.2 µg/ml, respectively. Kouame, et al. [27] studied on esential oils extracted from flower and stem of *A. conyzoides*, the results displayed antibacterial activities of with MIC in the range of 64 to 256 µg/ml on both Gram positive and Gram negative bacteria [27]. In this study, the MIC values were higher than previous reports [26, 27], this differences can relate to the dissimilarity of the time of sampling and extraction condition.

The ethyl acetate fractions had the most potent effect on the bacteria strains. But the MIC values still higher than the crude extracts. It suggested that the combination of every components in the crude extract give the best antibacterial effect. When tested on clinical strains, only ethyl acetate fractions of *Ageratum conyzoides* were active. Especially, it was active against *E. coli* and *K. pneumoniae* producing ESBL – the strains that were highly resistant in clinical infection. Hence, *Ageratum conyzoides* is a good candidate for anti-urologic bacteria.

A. conyzoides, A. vulgaris, and *H. annuus* essential oils had antimicrobial activity against both bacteria (MSSA, MRSA) and yeasts (*C. albicans, C. glabrata, C. tropicalis*). Especially, *A. vulgaris* essential oils were active against MRSA and *Candida* spp. This showed that the potential of *A.vulgaris* essential oils, suggested further research into its antifungal activity.

DPPH is a free radical, stable at room temperature, which produces a violet solution in methanol. It is reduced in the presence of an antioxidant molecule, making the color of the solutions turn to yellow. The use of DPPH provides an easy and rapid way to evaluate antioxidants.

All 9 ethanol extracts were capable of capturing DPPH radicals. *T. erecta* extracts had the lowest IC_{50} value, 17.3 µg/ml, which is only 2.5 times higher than the IC_{50} value of ascorbic acid. This result is due to the presence of many phenolic compounds in *T. erecta*. In previous studies, marigolds have been shown to contain quercetagetin, glucoside of quercetagetin, phenolics, syringic acid, methyl 3,5-dihydroxy-4-methoxy benzoate, quercetin, thienyl, and ethyl gallate [28]. Quercetin and ethyl gallate are compounds for potent antioxidant effects in both *in vivo* and *in vitro* [29-31]. Containing many compounds with strong free radical scavenging effect of *T. erecta* showed the ability for treatment of diseases caused by free radicals such as cancer, diabetes, cardiovascular, etc. However, this capacity needs to be evaluated more accurately by *in vivo* tests.

4. Conclusions

The fraction of ethyl acetat extracted from *A. conyzoides* possesses antimicrobial activities on uropathogenic *E. coli* and *K. pneumoniae* collected from District 2 Hospital, Ho Chi Minh city, Vietnam. This is the potential extract to apply for healthcare serum in prophylactic recurrence UIT.

5. Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

6. Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

7. Funding Statement

This study was financially supported by Univerisity of Medicine and Pharmacy at Ho Chi Minh city, through the grant No. 240-17.

8. Acknowledgments

The authors would like to express their gratitude to Associate Professor Nguyen Dinh Nga for providing the clinical microorganisms.

9. References

1. M. H. Legese, G. M. Weldearegay and D. Asrat, "Extended-spectrum beta-lactamase and carbapenemase-producing Enterobacteriaceae among Ethiopian children," *Infection and drug resistance*, vol. 10, pp. 27, 2017.

2. L. K. Logan and R. A. Weinstein, "The epidemiology of carbapenem-resistant Enterobacteriaceae: the impact and evolution of a global menace," *The Journal of infectious diseases*, vol. 215, no. suppl_1, pp. S28-S36, 2017.

3. R. F. Potter, A. W. D'souza and G. Dantas, "The rapid spread of carbapenem-resistant Enterobacteriaceae," *Drug Resistance Updates*, vol. 29, pp. 30-46, 2016.

4. Y.-Y. Liu, Y. Wang, T. R. Walsh, L.-X. Yi, R. Zhang, J. Spencer, Y. Doi, G. Tian, B. Dong and X. Huang, "Emergence of plasmid-mediated colistin resistance mechanism mcr-1 in animals and human beings in China: a microbiological and molecular biological study," *The Lancet infectious diseases*, vol. 16, no. 2, pp. 161-168, 2016.

5. J. Abbasi, "Infectious disease expert sees threat from colistin-resistant superbug," *JAMA*, vol. 316, no. 8, pp. 806-807, 2016.

6. B. Foxman, "Urinary tract infection syndromes: occurrence, recurrence, bacteriology, risk factors, and disease burden," *Infectious disease clinics of North America*, vol. 28, no. 1, pp. 1-13, 2014.

7. K. A. Kline, D. J. Schwartz, W. G. Lewis, S. J. Hultgren and A. L. Lewis, "Immune activation and suppression by group B streptococcus in a murine model of urinary tract infection," *Infection and immunity*, vol. 79, no. 9, pp. 3588-3595, 2011.

8. G. R. Nielubowicz and H. L. Mobley, "Host-pathogen interactions in urinary tract infection," *Nature Reviews Urology*, vol. 7, no. 8, pp. 430, 2010.

9. A. Ronald, "The etiology of urinary tract infection: traditional and emerging pathogens," *The American journal of medicine*, vol. 113, no. 1, pp. 14-19, 2002.

10. H. C. Voon, R. Bhat and G. Rusul, "Flower extracts and their essential oils as potential antimicrobial agents for food uses and pharmaceutical applications," *Comprehensive Reviews in Food Science and Food Safety*, vol. 11, no. 1, pp. 34-55, 2012.

11. J. M. Kalwij, "Review of "The Plant List, a working list of all plant species"," *Journal of Vegetation Science*, vol. 23, no. 5, pp. 998-1002, 2012.

12. M. B. Theodore, B. Luc and L. S. John, "Asteraceae in Flora of North America," Ed., http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=10074, 2018.

13. M. J. Abad, L. M. Bedoya and P. Bermejo, "Essential oils from the Asteraceae family active against multidrug-resistant bacteria," in *Fighting multidrug resistance with herbal extracts, essential oils and their components*, Ed., pp. 205-221, Elsevier, 2013.

14. I. Jallali, Y. Zaouali, I. Missaoui, A. Smeoui, C. Abdelly and R. Ksouri, "Variability of antioxidant and antibacterial effects of essential oils and acetonic extracts of two edible halophytes: *Crithmum maritimum* L. and *Inula crithmoïdes* L," *Food chemistry*, vol. 145, pp. 1031-1038, 2014.

15. A. Jamalian, M. Shams-Ghahfarokhi, K. Jaimand, N. Pashootan, A. Amani and M. Razzaghi-Abyaneh, "Chemical composition and antifungal activity of *Matricaria recutita* flower essential oil against medically important dermatophytes and soil-borne pathogens," *Journal de mycologie medicale*, vol. 22, no. 4, pp. 308-315, 2012.

16. H. Sebai, M.-A. Jabri, A. Souli, K. Rtibi, S. Selmi, O. Tebourbi, J. El-Benna and M. Sakly, "Antidiarrheal and antioxidant activities of chamomile (*Matricaria recutita* L.) decoction extract in rats," *Journal of ethnopharmacology*, vol. 152, no. 2, pp. 327-332, 2014.

17. A. A. Shahat, A. Y. Ibrahim and M. S. Elsaid, "Polyphenolic content and antioxidant activity of some wild Saudi Arabian asteraceae plants," *Asian Pacific Journal of Tropical Medicine*, vol. 7, no. 7, pp. 545-551, 2014.

18. N. Stanković, T. Mihajilov-Krstev, B. Zlatković, V. Stankov-Jovanović, V. Mitić, J. Jović, L. Čomić, B. Kocić and N. Bernstein, "Antibacterial and antioxidant activity of traditional medicinal plants from the Balkan Peninsula," *NJAS - Wageningen Journal of Life Sciences*, vol. 78, pp. 21-28, 2016.

19. R. Maria, M. Shirley, C. Xavier, S. Jaime, V. David, S. Rosa and D. Jodie, "Preliminary phytochemical screening, total phenolic content and antibacterial activity of thirteen native species from Guayas province Ecuador," *Journal of King Saud University-Science*, vol. 30, no. 4, pp. 500-505, 2018.

20. Nguyễn Thị Thúy Anh, Trần Cát Đông and N. Đ. Nga, "Khảo sát mức độ nhạy cảm của *Candida spp.* với fluconazole," *Y Học TP. Hồ Chí Minh*, vol. 16, no. 1, pp. 83-85, 2012.

21. N. K. Sethiya, M. M. M. Raja and S. H. Mishra, "Antioxidant markers based TLC-DPPH differentiation on four commercialized botanical sources of Shankhpushpi (A Medhya Rasayana): A preliminary assessment," *Journal of advanced pharmaceutical technology & research*, vol. 4, no. 1, pp. 25, 2013.

22. S. Burt, "Essential oils: their antibacterial properties and potential applications in foods - a review," *International journal of food microbiology*, vol. 94, no. 3, pp. 223-253, 2004.

23. S. Chouhan, K. Sharma and S. Guleria, "Antimicrobial activity of some essential oils - present status and future perspectives," *Medicines*, vol. 4, no. 3, pp. 58, 2017.

24. H. Padalia and S. Chanda, "Antimicrobial efficacy of different solvent extracts of *Tagetes erecta* L. flower, alone and in combination with antibiotics," *Applied Microbiology: Open Access*, vol. 1, pp. 106, 2015.

25. E. Amadi, C. Oyeka, R. Onyeagba, O. Ugbogu and I. Okoli, "Antimicrobial screening of *Breynia nivosus* and *Ageratum conyzoides* against dental caries organisms," *Journal of Biological Sciences*, vol. 7, no. 2, pp. 354-358, 2007.

26. K. O. Akinyemi, O. Oladapo, C. E. Okwara, C. C. Ibe and K. A. Fasure, "Screening of crude extracts of six medicinal plants used in South-West Nigerian unorthodox medicine for anti-methicillin

resistant *Staphylococcus aureus* activity," *BMC complementary and alternative medicine*, vol. 5, no. 1, pp. 6, 2005.

27. B. K. F. P. Kouame, D. Toure, L. Kablan, G. Bedi, I. Tea, R. Robins, J. C. Chalchat and F. Tonzibo, "Chemical constituents and antibacterial activity of essential oils from flowers and stems of *Ageratum conyzoides* from Ivory coast," *Records of Natural Products*, vol. 12, no. 2, pp. 160, 2018.

28. F. Nikkon, M. R. Habib, Z. A. Saud, M. R. Karim, A. K. Roy and S. Zaman, "Toxicological evaluation of chloroform fraction of flower of *Tagetes erecta* L. on rats," *International Journal of Drug Development and Research*, vol. 1, no. 1, pp. 161 - 165, 2009.

29. A. B. Bentz, "A review of quercetin: Chemistry, antioxidant properties, and bioavailability," *Journal of young investigators*, vol. 19, no. 10, 2009.

30. T. Kalaivani, C. Rajasekaran and L. Mathew, "Free radical scavenging, cytotoxic, and hemolytic activities of an active antioxidant compound ethyl gallate from leaves of *Acacia nilotica* (L.) wild. Ex. Delile sub sp. Indica (Benth.) Brenan," *Journal of food science*, vol. 76, no. 6, pp. T144-T149, 2011.

31. H. Le Son and N. P. Anh, "Phytochemical composition, in vitro antioxidant and anticancer activities of quercetin from methanol extract of *Asparagus cochinchinensis* (Lour.) Merr. tuber," *Journal of Medicinal Plants Research*, vol. 7, no. 46, pp. 3360-3366, 2013.