A Novel Chloroplast Super-Complex Consisting of the ATP Synthase and Photosystem I Reaction Center

Satarupa Bhaduri[1]† [*], Sandeep K Singh[1, ††] [*], Whitaker Cohn[2], S. Saif Hasan[1, †††], Julian P. Whitelegge[2], and William A. Cramer.[1#]

[1]Department of Biological Sciences, Purdue University, West Lafayette, IN 47907;
[2]Pasarow Mass Spectrometry Laboratory, The Jane and Terry Semel Institute for Neuroscience and Human Behavior, David Geffen School of Medicine, University of California/Los Angeles, CA 90024
[*]authors contributed equally; [#]to whom correspondence should be addressed, (waclab@purdue.edu)

Keywords: ATP synthase, mass spectrometry, native-gel electrophoresis, photosystem I, supercomplex

Abstract. Several 'super-complexes' of individual hetero-oligomeric membrane protein
complexes, whose function is to facilitate intra-membrane electron and proton transfer and
harvesting of light energy, have been previously characterized in the mitochondrial cristae and
chloroplast thylakoid membranes. The latter membrane is reported here to also be the location of
an intra-membrane super-complex which is dominated by the ATP-synthase and photosystem I
(PSI) reaction-center complexes, defined by mass spectrometry, clear-native PAGE and Western
Blot analyses. This is the first documented presence of ATP synthase in a super-complex with the
PSI reaction-center located in the non-appressed stromal domain of the thylakoid membrane.

INTRODUCTION

Super-complexes of hetero-oligomeric electron transport membrane protein complexes, formed

in the fluid mosaic membrane environment [1,2] have been described in mitochondria [3-6] where

their formation has been shown to be facilitated by anionic lipid [3, 7-12]. The chloroplast thylakoid

membrane contains four large hetero-oligomeric energy-transducing protein complexes (**Fig. 1**),

for which molecular structure information has been obtained: (i) the photosystem II (PSII) reaction-

center complex, responsible for the primary photochemistry associated with water splitting [13];

(ii) the cytochrome b_6f complex [14] which provides the quinol acceptor, electron transfer

connection, and trans-membrane proton translocation pathway [15, 16] between the PSII and

photosystem I (PSI) reaction centers; (iii) the 561 kDa plant photosystem I (PSI) complex [17, 18]

which is responsible for reduction of ferredoxin in the pathway of cyclic electron transport and

reduction of NADP$^+$ utilized for CO_2 fixation; and (iv) the 560 kDa ATPase/synthase [19,20]. No

super-complex has previously been documented which consists of complexes located in the non-

appressed thylakoid region.

Fig 1. **Distribution of the major intra-membrane energy transducing complexes**
associated with oxygenic photosynthesis in the appressed (grana) and non-
appressed (stroma) thylakoid membranes. Photosystem II (PSII), photosystem I (PSI),
cytochrome b_6f complex, and ATP synthase, with molecular weights of approximately 350,
530, 270 (with FNR), and 560 kDa, respectively. A supercomplex of ATP synthase and
photosystem I is implied in the figure by their proximity. Inter-membrane distances are
obtained from the references [32, 41].

Bipartite Thylakoid Membrane Structure; Residence of Energy-Transducing Complexes. The

chloroplast thylakoid membrane contains the energy transducing light-harvesting and electron-

transport protein complexes arranged in two structurally distinct membrane domains, (i) the

appressed grana membranes, and (ii) the non-appressed stroma or stroma-facing membrane

domain (Fig. 1). The membrane locations of the PSI, PSII, b_6f, and ATPase protein complexes

have been discussed in the context of long-range redox communication [21], and the locations in

grana and stromal compartments of the redox proteins that function in the 'non-cyclic' electron

transport pathway from water to NADP$^+$, and the PSI-linked redox proteins responsible for cyclic

electron transport and linked phosphorylation, described [21]. The discovery of super-complexes

in thylakoid membranes, which include complexes of the reaction centers with associated light

harvesting chlorophyll proteins [22-30] and/or the b_6f complex [31-33] and the electron

transport/water-splitting structure [13] has led to an improved perspective on the organization of

the membrane protein complexes which support the light harvesting and electron transport-proton

translocation functions central to photosynthetic membrane energy transduction.

Super-complexes isolated from thylakoid membranes thus far are: (**i**) LHCII: PSII reaction center

complexes [22-28]; (**ii**) nano-domains surrounding the PSII reaction center described by atomic

force microscopy which are populated by the b_6f complex [34] present in appressed and non-

appressed membrane domains [32]; (**iii**) cytochrome b_6f complex from *C. reinhardtii,* to which is

bound light harvesting complexes LHCI and LHCII and ferredoxin-NADP+ reductase (FNR),

which was obtained under conditions ('state 1- state 2 transition') favoring light energy transfer

from the antenna pigment proteins to the PSI reaction center complex [32,33]; (**iv**) the b_6f complex

with PSI reaction center complex in *Arabidopsis* and *Chlamydomonas* [32,33]; (**v**) a super-

complex of the PSI reaction center and the cytochrome b_6f complex isolated from cold-adapted

C. reinhardtii [35]; (**vi**) the structures described of the PSI reaction center complex and in

association with LHCs from plant sources [29,30]; (**vii**) NDH-associated supercomplexes [32, 36].

The presence of the PSI reaction center and ATPase in the edge of the unstacked grana

membranes has been documented [37-39].

The present study concerns a super-complex that combines the two hetero-oligomeric

membrane protein complexes of the non-appressed thylakoid domain, ATP synthase and PSI

reaction center, the first membrane protein super-complex in which ATP synthase has a major

presence. This result has been documented by the use of high-resolution Orbitrap mass

spectrometry to analyze the contribution to the super-complex of multiple major and minor protein

components, and clear-native-PAGE, SDS-PAGE, and immuno-blot analysis to identify protein

subunits of the component complexes.

MATERIALS AND METHODS

Purification of (i) Un-crosslinked Super-Complex (Preparations 1 and 2). Spinach

leaves macerated in darkness in HEPES-KOH pH 7.5, 150 mM NaCl, 5 mM EDTA at 4 °C and

sedimented at 4000 x g, were re-suspended in HEPES (4-(2-hydroxyethyl)-1-piperazine-

ethanesulfonic acid) -KOH pH 7.5, 15 mM NaCl, 5 mM EDTA, centrifuged at 4,000 x g, and re-

suspended in the same medium containing 5 mM NaCl. The chlorophyll concentration of thylakoid

membranes was adjusted to 1 mg/ ml, extracted with 1 % DDM, centrifuged at 15,000 x g, the

supernatant loaded on a 4-40% sucrose density gradient, and centrifuged at 200,000 x g.

Purification of (ii) Crosslinked Super-Complex (Preparation 3). Spinach leaves

macerated in darkness in 50 mM Tris-HCl, pH 7.5, 0.1M NaCl, 0.3 M sucrose, and 1 mM EDTA,

at 4 °C and sedimented at 6000 x g, were re-suspended in Tris-HCl, pH 7.5, 0.1 M NaCl, and 1

mM EDTA. The suspension was lysed for a second time using the French Press at a pressure of

8000 psi. The lysate was centrifuged at 10,000 x g, followed by ultracentrifugation at 160,000 x g

for 30 min to separate stacked and unstacked grana thylakoids. The fractions were washed with

1 mM EDTA before the chlorophyll concentration was measured and adjusted to 2.5 mg/ml

(chlorophyll a/b ratio = 5.4). Membrane proteins were extracted with 1 % β-DDM (β-dodecyl-

maltoside), centrifuged at 17,000 x g, the supernatant loaded on a 4-40% sucrose density

gradient, and centrifuged at 200,000 x g. The samples were crosslinked by 0.2% glutaraldehyde

in the 40% sucrose solution used to make the 10-40% gradient.

Electrophoretic Analysis. Protein bands from the sucrose gradient were fractionated, run on 4-

12 % clear-native PAGE, and eight resolved complexes excised for mass spectrometric analysis.

The largest band of approximately 1 MDa on the native gel (**Fig. 2B**) is inferred to contain the

ATP synthase-PSI super-complex.

Western Blotting*. Isolated super-complex was assayed on SDS (data not shown) and clear-

native PAGE, and transferred to a nitrocellulose membrane, the blot probed with anti-PsaA of PSI,

anti-ATP synthase α-subunit, and anti-FNR antibodies obtained from PhytoAb, Inc. (**Fig. 2C,D**

and E).

Mass Spectroscopy. Mass spectrometry was performed as described previously using nano-

liquid chromatography and a bench top Orbitrap mass spectrometer (QE-Plus, Thermo-Fisher

Scientific) operated in positive ion mode (nLC-MSMS) [40]. Proteins in native gel-bands were

alkylated, digested with trypsin and extracted for nLC-MSMS. The Mascot Search Algorithm

(Matrix Science, London, UK) was utilized to identify the significant presence of peptides, using

10 ppm mass tolerance on high-resolution precursor MS1 scans, and 20 milli-mass unit tolerance

on high-resolution product MS2 scans, obtaining an overall < 5% peptide false discovery rate

based upon a reversed sequence decoy search [41]. A spinach database was downloaded from

Uniprot (UP000054095). The top twenty Mascot protein scores for each sample were presented

representing the most abundant proteins in the sample and minimizing low scoring proteins that

are more likely to represent noise (Mascot 'protein' score is the summed score for unique peptides

within the sample analysis). A 'complex' score for each of the four major thylakoid protein

complexes was generated by adding together the protein scores for the top four highest scoring

proteins (or fewer than four if there were fewer than four proteins from that complex in the twenty

highest scoring proteins). The top twenty proteins were chosen as a cutoff because proteins below

this threshold have protein scores around 10% or less of the highest scoring proteins and thus

are more likely to be in the 'noise' fraction [32].

Assay of ATPase Activity. The "ADP-Glo" kit (Promega, Wisconsin) was used to measure

hydrolysis of ATP to ADP and inorganic phosphate by ATPase. ATP (0.4 mM) was added to the

super-complex fraction in the presence of buffer (50 mM Tris-HCl, 5 mM Mg^{2+}, and BSA) at room

temperature, incubated for 15 min, followed by the addition of "ADP-Glo reagent" and incubation

at room temperature (40 min) to consume unused ATP. The reaction mixture was then incubated

with the 'detection reagent' to convert ADP to ATP, measured by the reaction of luciferase/luciferin,

and the luminescence recorded in a spectral band centered at 560 nm, utilizing a Spectramax

Luminometer (Molecular Devices, CA).

Fluorescence Emission Spectroscopy. Fluorescence measurements were made with a

FluoroMax 3 Fluorimeter (Horriba-Yvon Inc.), using a 3 x 3 mm quartz microcell with 0.3 ml

volume. FAD containing the super-complex fraction in 10 mM HEPES buffer, pH 7.5, 0.5 mM

EDTA, 2 mM DDM, 10 % sucrose. Fluorescence was excited at 460 nm, and emission spectra

collected in a 500-580 nm spectral interval, with a stepping increment of 1 nm and a 0.1 sec

integration time. Excitation spectra were measured from 420 – 500 nm using an emission band

interval centered at 530 nm. Difference spectra were obtained by measuring the fluorescence in

the presence and absence of sodium dithionite, the latter a reductant with a redox potential

negative enough to fully reduce the flavin in FNR, thus effectively eliminating its absorbance and

resulting fluorescence.

RESULTS

A unique super-complex from thylakoid membranes; electrophoretic and immuno-blot

analysis*.* The super-complex was isolated from spinach thylakoid membranes by detergent

extraction followed by sucrose density gradient centrifugation (**Fig. 2A**, marked as "SC"), and the

approximate size was determined on a 4–12 % clear-native gel (**Fig. 2B**).

The presence of ATP synthase, PS I and FNR was confirmed by Western blot analyses

using antibodies raised against the *Arabidopsis* ATP synthase α-subunit (55 kDa) (**Fig. 2C**), the

60 kDa PsaA subunit of PSI (**Fig. 2D**), and FNR (36 kDa) (**Fig. 2E**).

Fig 2. **Purification and characterization of the PSI-ATP synthase super-complex**. **(A)**
4-40 % sucrose density gradient of spinach thylakoid membrane extract. Super-complex
(SC), marked by the arrow, is separated for further characterization. **(B)** 4-12 % Clear
Native PAGE of the super-complex fraction isolated from the sucrose gradient and stained
with Coomassie Brilliant Blue R-250. (**C, D, E**) Western blots of super-complex using 12
% SDS-PAGE probed with antibodies raised against **(C)** α-subunit (55 kDa) of
Arabidopsis ATP synthase, **(D)** PsaA subunit (60 kDa) of PSI, and **(E)** ferredoxin-NADPH
oxidoreductase (FNR).

Mass Spectrometry. High-resolution mass spectrometry of the isolated 1 MDa super-complex

from spinach chloroplast thylakoid membranes (**Table 1**) revealed the presence of ATP synthase

(0.56 MDa), Photosystem I (0.53 MDa). Polypeptide components derived from the 1MDa band in

Fig. 2B, were identified after in-gel digestion of the native gel-band and tandem mass

spectrometry with database searching constrained by a specified high statistical confidence

threshold, as described in *Methods*, and summarized in **Table 1A, B**. Presence of ATP synthase

and Photosystem I were identified with high statistical confidence by multiple polypeptides. Notably absent were Photosystem II (trace amounts of PsbB were detected in 1 preparation, assumed to represent contamination), and cytochrome b_6f complex polypeptides. Other components that were detected consistently were the polypeptides FtsH and FtsH2, Lhca3.1 (a known PSI chlorophyll a/b binding protein) and ferredoxin-NADPH oxidoreductase (FNR) that was confirmed by Western blot (**Fig. 2E**). A calcium-sensing receptor was found in 2/3 preparations and rubisco activase was found with a high score after crosslinking. Because of differential recovery of larger versus smaller subunits and peptides derived from them it is not possible to infer polypeptide stoichiometry from these measurements.

Table 1A. Identification of Twenty Highest Scoring Proteins in Three Super-Complex Preparations

Preparation 1	Score	Preparation 2	Score	Preparation3 (crosslinked)	Score
AtpB	2844	AtpB	2106	FtsH	4138
PsaB	2055	PsaB	2006	Rubisco activase [p]	4065
AtpA	1295	PsaD	1343	PsaB	3707
PsaD[a]	1261	AtpA	861	AtpB	3573
FtsH[b]	753	Lhca3.1	694	FtsH2	3281
Lhca3.1[c]	677	FNR	604	NadH1[q]	2548
Lhca6A[d]	664	PsbB	517	PsaD	2135
FNR	652	PsaF	492	NdhB2	1892
AtpG[e]	634	PsaA	471	Lhca3.1	1848
FtsH2[f]	613	PsaC	450	AtpA	1829
AtpE	591	AtpD	449	PsaG	1713
NdhB2[g]	559	FtsH2	439	Calcium-sensing receptor	1584
PsaC	503	PsaE[m]	435	PsaA	1337

PsaA	493	Lhca(7)	431	PsaE	1094
Lhca(7)[h]	449	PsaG	349	AtpG	1074
AtpD[i]	444	Lhca6A	348	ClpA_CD4B[r]	1066
PsaF[j]	442	AtpE	334	NADH1_13B[s]	1012
calcium-sensing recep.[k]	422	FtsH	330	□_carbonic anhydr. 1[t]	1005
RbLrubisco	409	Lhca4[n]	310	Lhca4	912
PsaG[l]	371	P31853.2[o]	301	Lhca6A	808

By BLAST homology, PsaD, tr|A0A0K9R886|A0A0K9R886_SPIOL

By BLAST homology, FtsH, tr|A0A0K9RFL8|A0A0K9RFL8_SPIOL

By BLAST homology, Lhca3.1, tr|A0A0K9R1K7|A0A0K9R1K7_SPIOL

By BLAST homology, Lhca6A, tr|A0A0K9RRT4|A0A0K9RRT4_SPIOL

By BLAST homology, AtpG, tr|A0A0K9QPJ7|A0A0K9QPJ7_SPIOL

By BLAST homology, FtsH2, tr|A0A0K9RD58|A0A0K9RD58_SPIOL

By BLAST homology, NdhB2, tr|A0A0K9QKT8|A0A0K9QKT8_SPIOL

By BLAST homology, Lhca(7), tr|A0A0K9RU56|A0A0K9RU56_SPIOL

By BLAST homology, AtpD, tr|A0A0K9RXF7|A0A0K9RXF7_SPIOL

By BLAST homology, PsaF, tr|A0A0K9RVH9|A0A0K9RVH9_SPIOL

By BLAST homology, calcium-sensing receptor, tr|A0A0K9R5V6|A0A0K9R5V6_SPIOL

By BLAST homology, PsaG, tr|A0A0K9RHQ2|A0A0K9RHQ2_SPIOL

By BLAST homology, PsaE, tr|A0A0K9RHN1|A0A0K9RHN1_SPIOL

By BLAST homology, Lhca4, tr|A0A0K9R772|A0A0K9R772_SPIOL

By BLAST homology, ATPase b', tr|A0A0K9QQW2|A0A0K9QQW2_SPIOL

By BLAST homology, Rubisco activase, tr|A0A0K9Q765|A0A0K9Q765_SPIOL

By BLAST homology, NadH1, tr|A0A0K9QED7|A0A0K9QED7_SPIOL

By BLAST homology, ATP-dependent Clp protease ATP-binding subunit ClpA homolog CD4B,

tr|A0A0K9QTC9|A0A0K9QTC9_SPIOL

By BLAST homology, NadH1_13B, tr|A0A0K9Q7H2|A0A0K9Q7H2_SPIOL

By BLAST homology, □_carbonic anhydr. 1, tr|A0A0K9RNJ7|A0A0K9RNJ7_SPIOL

Table 1B. Major Thylakoid Complexes in Supercomplex.

4 highest scores for each complex component added together.

Complex	Prep. 1	Prep.2	Prep.3 (crosslinked)
ATPase	5364	3750	6476
PSI	4312	4312	8892
b_6f	0	0	0
PSII*	0	517[c]	0

191   *Minor contamination in by PSII.

193   The presence of FNR was confirmed through Western blot analysis using an anti-FNR antibody

194   probe (**Fig. 2E**), and determination of fluorescence excitation and emission spectra that are

195   characteristic of FAD [42] (**Fig. 3**). No polypeptides from the cytochrome b_6f complex were

196   detected at the stringency used.

Figure 3. Difference fluorescence emission (top) and excitation (bottom) spectra of the 'super-complex;' measured with a FluoroMax 3 Fluorimeter (Horiba-Yvon Inc.). (**A**) Emission spectra measured between 500 – 600 nm. Stepping increment, 1 nm; Integration time, 0.1 sec; Excitation, 466 nm; bandwidth 5 nm for excitation and emission. (**B**) Excitation spectra measured in 400–500 nm range, measuring the emission at its 530 nm peak. Spectra of undiluted sample were measured in a quartz microcell, 3 x 3 mm, volume, 0.3 ml (Starna Cells) before and after addition of dithionite grains. Difference spectra were obtained by subtraction of spectra, i. e., 'oxidized–reduced,' measured before and after addition of dithionite.

Presence of FNR in the Super-Complex. The presence of FNR in the spinach super complex was confirmed from fluorescence emission and excitation spectra produced by the isoalloxazine moiety of FAD (**Figs. 3A, B**). The super-complex, suspended in 10 mM HEPES buffer, pH 7.5, and 0.5 mM EDTA, 2 mM DDM, 10 % sucrose, was excited at 460 nm and emission spectra were acquired over the wavelength range 500-600 nm. The difference emission spectrum, calculated with and without the strong reductant, sodium dithionite, showed an emission maximum in the oxidized state at 530 nm, which is characteristic of the presence of flavin [42]. A similar conclusion was reached from the difference excitation spectra measured with and without sodium dithionite

over the 400-500 nm range using emission at 530 nm. The excitation spectrum showed maximum

at 460 nm, characteristic of oxidized flavin [42]. Spectra are normalized to unit emission intensity.

ATPase Activity. The F1-ATPase component in the super-complex demonstrated *in vitro* ATP

hydrolysis, as described in *Methods*. The "ADP-Glo kit" (Promega) was used to provide a

qualitative measure of ATP hydrolysis, which was dependent on the presence of the PSI-ATPase-

FNR super-complex (**Table 2**). ATP hydrolysis was measured by addition of 0.4 mM ATP to the

super-complex fraction in the buffer containing 50 mM Tris-HCl, 5 mM Mg^{2+}, and 0.1 mg/ml BSA).

Table 2. ATPase Activity of the Super-Complex. As described in *Methods*, kinase activity was
measured qualitatively through luciferase emission in the positive control (buffer and 0.4 mM
ADP); negative control (buffer and ATP (0.4 mM) and the purified super-complex in buffer and 0.4
mM ATP.

Sample	Chemiluminescent Signal (cpm)
+ADP (positive control)	$(3.1 \pm 2) \times 10^6$
+ATP (negative control)	$(5.5 \pm 1.2) \times 10^2$
+ Supercomplex	$(4.4 \pm 0.6) \times 10^5$

DISCUSSION

Several super-complexes of PSI with thylakoid complexes have previously been described

[17,18, 29, 30], but a complex with ATP synthase has thus far been absent. These authors defined

the super-complex based upon the ability to visualize the complex using cryo-electron

microscopy. In the present case it has not been possible to preserve the PSI-ATP synthase

complex during the freezing process for cryo-EM. Consequently, we do not include an EM image

of this super-complex in this study. Alternatively, classical biochemical techniques show this

super-complex running as a discrete band in sucrose-gradient centrifugation and as a ~1 MDa

band on a native gel combined with a modern description of the major components of the super-

complex by high-resolution mass spectrometry.

The identity of the PSI-ATP synthase super-complex is supported by the facts that (i) the

super-complex band is the densest, implying a higher protein/lipid content than other thylakoid

super-complexes consistent with the presence of the ATP synthase which contains no pigments;

(ii) the molecular weight observed by native gel electrophoresis of ~1 MDa is consistent with an

equimolar mixture of PSI [17,18] (MW= 561 kDa) and ATP synthase (MW = 560 kDa) [19,20], and

no other complex.

Modern mass spectrometry using high-resolution precursor- and production scans in

tandem MS protein identification underscores current proteomics [44]. In-gel analysis of bands on

native gels is relatively uncommon compared to bands on SDS gels. In fact, the number of tryptic

peptides derived from the subunits of a super-complex, far exceeds that expected from trypsin

treatment of a single protein. The modern mass spectrometer combined with reverse-phase

chromatography provides a powerful tool to analyze a heterogenous mixture of peptides to

determine the super-complex. Integral membrane proteins with large extrinsic domains yield

abundant peptides that are easily recoverable in the in-gel procedure yielding very high relative

abundance 'scores' where these subunits are present. Thus, ATP synthase and PSI with very

high scores are identified, based on the four highest scoring polypeptides underlying our

conclusion that these two complexes are the only complexes within this super-complex (**Table**

1B). PSII and cytochrome b_6f components are easily identified in other bands from the sucrose

gradient (not shown) but not in the densest band representing the super-complex as described.

Small subunits with a large proportion of transmembrane sequence are poorly represented in this

type of analysis [45] and were not seen in this analysis. Intact protein analysis is better suited for

full subunit coverage, as reported in 2002 [46] and would be well suited to detect all the subunits

in the supercomplex. Native intact supercomplex mass spectrometry would be most desirable as

it addresses stoichiometry [47], but it as yet unavailable for this preparation.

Mass spectrometry also showed the presence of other proteins that might be harder to define

by cryo-EM, reliably identifying FtsH and FtsH2, Lhca3A and FNR. Initial studies on these

components have documented the presence of FNR.

This is the first instance of detection of a photosynthetic super-complex which contains the

~560 kDa ATP synthase protein complex. Localized coupling of the electron transport system and

ATP synthesis by super-complex formation in mitochondria has been proposed [48]. It has been

suggested that the oxidative phosphorylation system in respiratory systems can operate in two

states: via utilization of (i) the bulk phase trans-membrane electrochemical potential gradient [48],

or (ii) protons localized in the membrane that are accessible to the active site chemistry [49]. In

the present study, the occurrence of ATP synthase in association with the PSI reaction-center

complex, suggests the possibility of a relatively localized coupling between (i) ATP synthesis and

the transmembrane electrical potential generated in the PSI reaction center, and/or (ii) a localized

domain of carbon dioxide fixation.

The isolated complex dominated by the presence of PSI and ATP synthase is also

associated with a number of proteins like the ATP dependent, membrane anchored zinc

metalloprotease FtsH, FNR and Rubisco Activase. FtsH has been implicated in chloroplast

biogenesis and repair of photosystem II under high light conditions [50]. It has been demonstrated

that FtsH, localized in the stroma lamellae, is critical for the biogenesis of PSI complex under

normal light conditions [51].   In bacteria FtsH is responsible for regulating ATP synthase

biogenesis [52]. However, such a function has not yet been documented in the plant system.

Previous studies have shown that a Chl a/b ratio of 4.6 in the stroma lamellae isolated from

spinach leaves, had a PSI/PSII ratio of ~3 [53]. In the current preparation, a chlorophyll a/b ratio

of 5.4 was measured in the membrane fraction, which indicates that the super-complex PSI-

ATPase containing has been derived from a PSI enriched stroma lamellae membrane. A high

relative abundance of Lhca 3.1, a light harvesting protein associated with PSI [54], as determined

by mass spectrometry, also confirms a major presence of photosystem I in the super-complex.

The flavoenzyme, ferredoxin-NADP$^+$ reductase (FNR), found in both soluble and thylakoid

membrane associated forms, has been proposed to define the photosynthetic electron transport

pathway [55]. PSI bound FNR or soluble stromal FNR facilitates linear electron flow, while under

reduced conditions, FNR associates with the cytochrome b_6f complex and promotes cyclic

electron flow [56]. The presence of FNR in the super-complex has been confirmed by mass

spectrometry (**Table1A**), Western Blot (**Fig. 2E**) and difference fluorescence spectroscopy (**Fig.

3**). A third protein, rubisco activase, found associated with the high molecular weight crosslinked

band containing PSI and ATP synthase (Preparation 3), interacts with Rubisco to release the

bound inhibitors in an ATP dependent fashion and hence, initiates the CO_2 fixation cycle [57].

AUTHOR INFORMATION

Corresponding Author

*e-mail: waclab@purdue.edu; telephone: (765)-494-4956

Present Addresses

†Department of Cellular and Developmental Biology and Department of Biosciences, University
of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093; †† Department of Biological
Chemistry & Molecular Pharmacology, Harvard Medical School, 240 Longwood Avenue, Boston
MA, 02115; ††† Department of Biochemistry and Molecular Biology, University of Maryland
School of Medicine, 108 N. Greene St., Baltimore, MD 21201 AND Center for Biomolecular
Therapeutics (CBT), 9600 Gudelsky Drive, Rockville, MD 20850

Author Contributions. Project proposed by SKS and WAC. Super-complex isolation and
biochemical characterization done by SKS and SB. Mass spectrometry analyses done by JPW.
SSH contributed to the experimental design. Manuscript written by SB, JPW, and WAC.

ABBREVIATIONS. CN- and SDS-PAGE, clear native- and sodium dodecyl sulfate-
polyacrylamide gel electrophoresis; cpm, counts/min; cyt, cytochrome; DDM, n-dodecyl-β-D-
malto-pyranoside; FNR, ferredoxin-NADP$^+$ reductase; EM, electron microscopy; FtsH,
filamentous, temperature-sensitive ATP-dependent protease; HRP, horse radish peroxidase;
LHC, light harvesting complex; Mr, relative molecular weight; PAGE, polyacrylamide gel
electrophoresis; pdb, protein data bank; PSI, PSII, photosystems I, II.
323

ACKNOWLEDGMENTS.
We thank S. D. Zakharov, S. Puthiyaveetil, R. Morgan-Kiss, and I. Kalra for helpful discussions,
and S. Naurin for technical support. Studies were supported by Dept. of Energy/Basic Energy
Sciences/Photosynthetic Systems grant DE-SC00118238, the Henry Koffler Distinguished
Professorship (WAC), and NIHDK-063491 (JPW).

REFERENCES

1.   Singer SJ and Nicholson GL. The fluid mosaic model of the structure of cell membranes.
Science 1972;175: 720-731.
2.   Engelman DM. Membranes are more mosaic than fluid, Nature 2005; 438: 578-580.
3.   Pfeiffer K, Gohil V, Stuart RA, Hunte C, Brandt U, Greenberg ML, Schägger H. Cardiolipin
stabilizes respiratory chain supercomplexes, Journal of Biological Chemistry 2003; 278: 52873-
52880.
4.   Genova ML, Lenaz G. Functional role of mitochondrial respiratory supercomplexes,
Biochimica et Biophysica Acta (BBA) - Bioenergetics 2014;1837: 427-443.
5.   Sousa JS, D'Imprima E, Vonck J. Mitochondrial respiratory chain complexes, in *membrane*
protein complexes: structure and function (Harris, J. R., and Boekema, E. J., Eds.) 2018; pp. 167-
227, Springer Singapore, Singapore.
6.   Davies KM, Blum TB, Kühlbrandt W. Conserved in situ arrangement of complex I and III$_2$
in mitochondrial respiratory chain supercomplexes of mammals, yeast, and plants. Proceedings
of the National Academy of Sciences 2018; 115: 3024-3029.
7.   Wenz T, Hielscher R, Hellwig P, Schagger H, Richers S, Hunte C. Role of phospholipids
in respiratory cytochrome *bc*$_1$ complex catalysis and supercomplex formation Biochim. Biophys.
Acta 2009; 1787: 609-616.
8.   Mileykovskaya E, Dowhan W. Cardiolipin-dependent formation of mitochondrial
respiratory supercomplexes, Chemistry and Physics of Lipids 2014; 179: 42-48.
9.   Arnarez C, Marrink SJ, Periole X. Molecular mechanism of cardiolipin-mediated assembly
of respiratory chain supercomplexes, Chemical Science 2016; 7: 4435-4443.
10.   Blaza JN, Serreli R, Jones AJ, Mohammed K, Hirst J. Kinetic evidence against partitioning
of the ubiquinone pool and the catalytic relevance of respiratory chain supercomplexes.
Proceedings of the National Academy of Sciences of the United States of America 2014; 111:
15735-15740.
11.   Bottinger L, Horvath SE, Kleinschroth T, Hunte C, Daum G, Pfanner N, Becker T.
Phosphatidyl-ethanolamine and cardiolipin differentially affect the stability of mitochondrial
respiratory chain supercomplexes, J Mol Biol 2012; 423: 677-686.
12.   Arnarez C, Marrink SJ, Periole X. Identification of cardiolipin binding sites on cytochrome
c oxidase at the entrance of proton channels, Sci Rep 2013; 3: 1263.
13.   Umena Y, Kawakami K, Shen JR, Kamiya, N. Crystal structure of oxygen-evolving
photosystem II at a resolution of 1.9 A, Nature 2011; 473: 55-60.

14. Hasan SS, Cramer WA. Internal lipid architecture of the hetero-oligomeric cytochrome b_6f complex. Structure 2014; 22: 1008-1015.

15. Hasan SS, Yamashita E, Baniulis D, Cramer WA. Quinone-dependent proton transfer pathways in the photosynthetic cytochrome b_6f complex. Proc Natl Acad Sci. 2013; 110: 4297–4302.

16. Shikanai T, Yamamoto H. Contribution of cyclic and pseudo-cyclic electron transport to the formation of proton motive force in chloroplasts. Molecular Plant 2017; 10: 20-29.

17. Mazor Y, Borovikova A, Caspy I, Nelson N. Structure of the plant photosystem I supercomplex at 2.6 A resolution. Nature Plants 2017; 3: 17014.

18. Caspy I, Nelson N. Structure of the plant photosystem I. Biochemical Society Transactions 2018; 46(2): 285-294.

19. Hahn A, Vonck J, Mills DJ, Meier T, Kühlbrandt W. Structure, mechanism, and regulation of the chloroplast ATP synthase, Science 2018; 360: aat4318.

20. Kühlbrandt W. Structure and mechanisms of f-type ATP synthases. Annual Review of Biochemistry 2019; 88: 515-549.

21. Anderson JM. Lateral heterogeneity of plant thylakoid protein complexes: early reminiscences. Philos Trans R Soc Lond B Biol Sci 2012; *367:* 3384-3388.

22. Joliot P, Joliot GN. Regulation of cyclic and linear electron flow in higher plants, Proceedings of the National Academy of Sciences of the United States of America 2011; 108 13317-13322.

23. Crepin A, Santabarbara S, Caffarri S. Biochemical and spectroscopic characterization of highly stable photosystem ii supercomplexes from *Arabidopsis*, The Journal of biological chemistry 2016; 291: 19157-19171.

24. van Bezouwen LS, Caffarri S, Kale RS, Kouril R, Thunnissen AWH, Oostergetel GT, Boekema EJ. Subunit and chlorophyll organization in the plant photosystem II supercomplex, Nature Plants 2017; 3: 17080.

25. Wei X, Su X, Cao P, Liu X, Chang W, Li M, Zhang X, Liu Z. Structure of spinach photosystem II–LHCII supercomplex at 3.2 Å resolution, Nature 2016; 534: 69.

26. Su X, Ma J, Wei X, Cao P, Zhu D, Chang W, Liu Z, Zhang X, Li, M. Structure and assembly mechanism of plant $C_2S_2M_2$-type PSII-LHCII supercomplex. Science 2017; 357: 815.

27. Sheng X, Liu X, Cao P, Li M, Liu Z. Structural roles of lipid molecules in the assembly of plant PSII-LHCII supercomplex. Biophys Rep 2018; 4: 189-203.

28. Pi X, Zhao S, Wang W, Liu D, Xu C, Han G, Kuang T, Sui SF, Shen JR. The pigment-protein network of a diatom photosystem II–light-harvesting antenna supercomplex, Science 2019; 365, eaax4406.

29. Suga M, Qin X, Kuang T, Shen JR. Structure and energy transfer pathways of the plant photosystem I-LHCI supercomplex. Current Opinion in Structural Biology 2016; 39: 46-53.

30. Pan X, Ma, J, Su X, Cao P, Chang W, Liu Z, Zhang X, Li M.Structure of the maize photosystem I supercomplex with light-harvesting complexes I and II. Science 2018; 360: 1109.

31. Iwai M, Takizawa K, Tokutsu R, Okamuro A, Takahashi Y, Minagawa J. Isolation of the supercomplex that drives cyclic electron flow in photosynthesis. Nature 2010; 464: 1210-1213.

32. Yadav KN, Semchonok DA, Nosek L, Kouril R, Fucile G, Boekema EJ, Eichacker LA. Supercomplexes of plant photosystem I with cytochrome b_6f, light-harvesting complex II and NDH, Biochim Biophys Acta 2017; 1858*:* 12-20.

33. Steinbeck J, Ross IL, Rothnagel R, Gäbelein P, Schulze S, Giles N, Ali R, Drysdale R, Sierecki E, Gambin Y, Stahlberg H, Takahashi Y, Hippler M, Hankamer B. Structure of a PSI–LHCI–cyt b_6f supercomplex in *Chlamydomonas reinhardtii* promoting cyclic electron flow under anaerobic conditions. Proceedings of the National Academy of Sciences 2018; 115: 10517-10522.

34.   Johnson MP, Vasilev C, Olsen JD, Hunter CN. Nanodomains of cytochrome b_6f and photosystem II complexes in spinach grana thylakoid membranes, Plant Cell 2014; 26: 3051-3061.

35.   Szyszka-Mroz B, Pittock P, Ivanov AG, Lajoie G, Huner NP. The Antarctic psychrophile *Chlamydomonas sp*. UWO 241 preferentially phosphorylates a photosystem I-cytochrome b_6f supercomplex, Plant physiology 2015; 169: 717-736.

36.   Kato Y, Sugimoto K, Shikanai T. NDH-PSI supercomplex assembly precedes full assembly of the ndh complex in chloroplast, Plant Physiology 2018;176: 1728.

37. Kaftan D, Brumfeld V, Nevo R, Scherz A, Reich Z. From chloroplasts to photosystems: in situ scanning force microscopy on intact thylakoid membranes. The EMBO journal 2002; 21: 6146-6153.

38.   Puthiyaveetil S, Tsabari O, Lowry T, Lenhert S, Lewis RR, Reich Z, Kirchhoff H. Compartmentalization of the protein repair machinery in photosynthetic membranes, Proceedings of the National Academy of Sciences of the United States of America 2014; 111: 15839-15844.

39.   Kirchhoff H, Li M, Puthiyaveetil S. Sublocalization of cytochrome b_6f complexes in photosynthetic membranes. Trends in Plant Science 2017; 22: 574-582.

40.   Ding L, Xie F, Zhao M, Wang S, Xie J, Xu G. Rapid quantification of sucrose esters in oriental tobacco by liquid chromatography-ion trap mass spectrometry, Journal of Separation Science 2007; 30: 35-41.

41.   Elias JE, Gygi SP.Target-decoy search strategy for mass spectrometry-based proteomics, Methods Mol Biol 2010; 604: 55-71.

42.   Maeda M, Hamada D, Hoshino M, Onda Y, Hase T, Goto Y. Partially folded structure of flavin adenine dinucleotide-depleted Ferredoxin-NADP$^+$ Reductase with residual NADP$^+$ binding domain. Journal of Biological Chemistry 2002; 277: 17101-17107.

43.   Vossen KM, Fried MG. Sequestration stabilizes lac repressor–DNA complexes during gel electrophoresis. Analytical Biochemistry 1997; 245: 85-92.

44.   Mann M,Kelleher NL.Precision proteomics: the case for high resolution and high mass accuracy, Proceedings of the National Academy of Sciences of the United States of America 2008; 105: 18132-18138.

45.   Souda P, Ryan CM, Cramer WA, Whitelegge J. Profiling of integral membrane proteins and their post translational modifications using high-resolution mass spectrometry, Methods 2011; 55: 330-336.

46.   Whitelegge JP, Zhang H, Aguilera R, Taylor RM, Cramer WA. Full subunit coverage liquid chromatography electrospray ionization mass spectrometry (lcms+) of an oligomeric membrane protein, cytochrome b_6f complex from spinach *and the* cyanobacterium *Mastigocladus Laminosus*. Mol. Cell Proteomics 2002; *1*, 816-827.

47. Robinson CV. Mass spectrometry: From plasma proteins to mitochondrial membranes. Proc Natl Acad Sci U S A. 2019 ;116(8): 2814-2820.

48.   Mitchell P. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. Nature 1961; 191: 144-148.

49.   Williams RJP. Possible functions of chains of catalysts, Journal of Theoretical Biology 1961;1: 1-17.

50.   Kato Y, Sakamoto W. FtsH protease in the thylakoid membrane: physiological functions and the regulation of protease activity, Frontiers in Plant Science 2018; 9: 855-855.

51.   Järvi S, Suorsa M, Tadini L, Ivanauskaite A, Rantala S, Allahverdiyeva Y, Leister D, Aro EM. Thylakoid-bound FtsH proteins facilitate proper biosynthesis of photosystem I, Plant physiology 2016;171: 1333-1343.

52.   Deckers-Hebestreit G. Assembly of the *Escherichia coli*; F_0F_1ATP synthase involves distinct subcomplex formation, Biochemical Society Transactions 2013; 41: 1288.

53.   Danielsson R, Albertsson PÅ, Mamedov F, Styring S. Quantification of photosystem I and
II in different parts of the thylakoid membrane from spinach, Biochimica et Biophysica Acta (BBA)
- Bioenergetics 2004; 1608: 53-61.

54.   Ozawa SI, Bald T, Onishi T, Xue H, Matsumura T, Kubo R, Takahashi H, Hippler M,
Takahashi Y.Configuration of Ten Light-Harvesting Chlorophyll a/b Complex I Subunits in
Chlamydomonas reinhardtii Photosystem I, Plant Physiology 2018; 178: 583-595.

55.   Mosebach, L, Heilmann C, Mutoh R, Gäbelein P, Steinbeck J, Happe T, Ikegami T, Hanke
G, Kurisu G, Hippler M. Association of Ferredoxin:NADP+ oxidoreductase with the photosynthetic
apparatus modulates electron transfer in *Chlamydomonas reinhardtii*, Photosynthesis Research
2017; 134: 291-306.

56.   Joliot P, Joliot A. Cyclic electron flow in C3 plants, Biochimica et Biophysica Acta (BBA) -
Bioenergetics 2006;1757: 362-368.

57.   Keown JR, Griffin MD, Mertens HD, Pearce FG. Small oligomers of ribulose-bisphosphate
carboxylase/oxygenase (rubisco) activase are required for biological activity, Journal of Biological
Chemistry 2013; 288: 20607-20615.

Stroma

Lumen

Aprressed Membrane

21 nm

2 nm

Non-Aprressed Membrane

Photosystem I

ATP Synthase

PQH$_2$

Photosystem II

Cytochrome b_6f

PC

A

10%

← SC

40%

Sucrose Density Gradient

B

Supercomplex

Mol.Wt. (kDa)

CN-PAGE

C

Mol.Wt. (kDa)

Western ATPα

250
150
100
75

Western-ATP synthase

D

Western PsaA

Western-PSI

E

Western FNR

Western-FNR