Temperature-dependence of metabolic rate in tropical and temperate aquatic insects: support for

the Climate Variability Hypothesis in mayflies but not stoneflies

Alisha A. Shah[1,2], H. Arthur Woods[2], Justin C. Havird[3], Andrea C. Encalada[4], Alexander S. Flecker[5], W.

Chris Funk[1,6], Juan M. Guayasamin[4], Boris C. Kondratieff[1,6,7], N. LeRoy Poff[1,6,8], Steven A. Thomas[9],

Kelly R. Zamudio[5], Cameron K. Ghalambor[1,6,10]

Affiliations:

[1] Department of Biology, Colorado State University, Fort Collins, CO, USA

[2] Division of Biological Sciences, University of Montana, Missoula, MT, USA

[3] Department of Integrative Biology, University of Texas, Austin, TX, USA

[4] Colegio de Ciencias Biológicas y Ambientales COCIBA, Instituto BÍOSFERA-USFQ, Campus

Cumbayá, Universidad San Francisco de Quito USFQ, Quito, Ecuador

[5] Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY, USA

[6] Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO, USA

[7] Department of Bioagricultural Sciences and Pest Management, Colorado State University, Fort Collins,

CO, USA

[8] Institute for Applied Ecology, University of Canberra, ACT, Australia

[9] School of Natural Resources, University of Nebraska, Lincoln, NE, USA

[10] Department of Biology, Centre for Biodiversity Dynamics, Norwegian University of Science and

Technology (NTNU), Trondheim, Norway

Correspondence:

Alisha A. Shah, Division of Biological Sciences, University of Montana, Missoula, MT, USA; Email:

alisha.shah@mso.umt.edu; Phone: (512) 694-7532; ORCID: 0000-0002-8454-7905

Running Head:

Aquatic insect sensitivity to climate change

Abstract:

A fundamental gap in climate change vulnerability research is an understanding of the relative thermal

sensitivity of ectotherms. Aquatic insects are vital to stream ecosystem function and biodiversity but

insufficiently studied with respect to their thermal physiology. With global temperatures rising at an

unprecedented rate, it is imperative that we know how aquatic insects respond to increasing temperature

and whether these responses vary among taxa, latitudes, and elevations. We evaluated the thermal

sensitivity of standard metabolic rate in stream-dwelling baetid mayflies and perlid stoneflies across a

~2,000 m elevation gradient in the temperate Rocky Mountains in Colorado, U.S.A., and the tropical

Andes in Napo, Ecuador. We used temperature-controlled water baths and microrespirometry to estimate

changes in oxygen consumption. Tropical mayflies generally exhibited greater thermal sensitivity in

metabolism compared to temperate mayflies; tropical mayfly metabolic rates increased more rapidly with

temperature and the insects more frequently exhibited behavioral signs of thermal stress. By contrast,

temperate and tropical stoneflies did not clearly differ. Varied responses to temperature among baetid

mayflies and perlid stoneflies may reflect differences in evolutionary history or ecological roles as

herbivores and predators, respectively. Our results show that there is physiological variation across

elevations and species and that low elevation tropical mayflies may be especially imperiled by climate

warming. Given such variation among species, broad generalizations about the vulnerability of tropical

ectotherms should be made more cautiously.

Keywords:

Metabolic rate, thermal tolerance, latitude, elevation, aquatic insects, climate change vulnerability,

thermal performance curve

Introduction:

Metabolism integrates an organism's existing energy supplies and underlies fitness-related

activities like movement, growth, tissue maintenance, and development (Dillon, Wang, & Huey, 2010;

Gillooly, Brown, West, Savage, & Charnov, 2001). In ectotherms, metabolic rates are sensitive to body

temperature, which can influence not only individual fitness, but also geographic distributions and

abundances (Angilletta, 2009; Terblanche & Chown, 2007; Vannote & Sweeney, 1980). A major

challenge in global change biology is to understand how temperature alters metabolism and energy

budgets and, in turn, the ability of species to cope with climate change (Colwell, Brehm, Cardelús,

Gilman, & Longino, 2008; Dillon et al., 2010; Parmesan & Yohe, 2003; Walther et al., 2002). To date,

attempts to measure vulnerability to climate change have largely assessed species responses to

temperature challenges, such as lethal and critical thermal limits (Deutsch et al., 2008; Pinsky, Eikeset,

McCauley, Payne, & Sunday, 2019). Despite their utility, these metrics do not capture the continuous or

incremental sub-critical effects of temperature on an organism's energy budget. Such effects are

particularly important in ectotherms, because body temperature and metabolic rate increase with

environmental temperature, resulting in higher energy expenditure. This response is likely to trade off

against other demands on energy budgets, such as growth, reproduction, and locomotor performance

(Kozłowski, 1992; Pörtner et al., 2006; Wieser, 1994). For example, in aquatic ectotherms, increased

metabolic demand resulting from higher temperature can outpace the supply of oxygen from the

environment causing decreased performance and lowered tolerance to heat stress (Pörtner, Bock, & Mark,

2017). Thus, understanding how metabolic responses to temperature vary across different thermal

environments can help us predict geographic variation in response to global warming (Dillon et al., 2010).

In this study, we measured metabolic rate profiles in juveniles (nymphs) of two major groups of

stream insects – mayflies and stoneflies – in temperate and tropical streams across a paired elevation

gradient. Mayflies and stoneflies spend most of their life cycles as aquatic nymphs, during which they

feed, grow, and develop eggs and sperm before emerging to breed as short-lived flying adults (Brittain,

1990). Water temperature is therefore a major driver of the life cycle in these taxa. We aimed to elucidate

differences in patterns of species' thermal sensitivity and thus better evaluate potential vulnerabilities of

mayflies and stoneflies to projected climate warming (Chown, Duffy, & Sørensen, 2015).

Much like other types of thermal performance curves (TPCs), oxygen-consumption profiles (used

to measure metabolic rate) vary with increasing temperature and display three identifiable regions

(Pörtner, 2002; Schulte, 2015): 1) an ascending phase; 2) a peak, which indicates the onset of stress (Huey

& Stevenson, 1979; Pörtner & Knust, 2007); and 3) a descending phase of metabolic depression (Guppy

& Withers, 1999). When coupled with other measures of performance and critical thermal limits, changes

in oxygen consumption provide a robust picture of how temperature shapes the ecology of an organism

(Angilletta, 2009; Pörtner et al., 2017) (Fig. 1).

Figure 1. Expectations for change in standard metabolic rate (oxygen consumption) and performance in response to temperature in an aquatic ectotherm. The thermal performance curve (TPC, solid green line, left panel) refers to any functional performance trait (e.g., swimming performance), where higher performance is better. T_{OPT}, where performance is highest, reflects the animal's preferred temperature. The oxygen consumption curve (dashed line) is a special case of TPC. The peak in this case (T-MR$_{PEAK}$) could represent the temperature at which standard metabolic rate (SMR) is so high that it essentially equals the maximum metabolic rate (MMR) that the ectotherm can achieve (circle, A). T-MR$_{PEAK}$ may therefore reflect stress (red) and aligns with a decline in the functional performance trait. The general location of behaviorally determined critical thermal limits (CT$_{MIN}$ and CT$_{MAX}$) are also shown (blue line, left panel). In this study we focus only on SMR. We predict that SMR will rise steeply in tropical species (B) and be stressful when temperatures exceed the typical narrow range of temperatures they experience (gray rectangle). In comparison, temperate species should exhibit a more gradual rise in SMR across the same temperatures (C), reflecting the broad range of environmental temperatures they experience (grey rectangle).

Although TPCs for traits like locomotion and metabolic rate are expected to have similar shapes,

a metabolic rate curve is a special type of TPC (Pörtner, 2002; Schulte, Healy, & Fangue, 2011), with

different implications for organismal fitness (Kellermann et al., 2019). The performance traits used in

many TPCs—running speed, feeding rates, or growth—are often interpreted as components of fitness,

such that *more is better*. For these traits, peaks indicate the location of thermal optima (T_{OPT}, Fig. 1) and

suggest that performance is optimized at that temperature (Angilletta, Niewiarowski, & Navas, 2002;

Huey & Kingsolver, 1989; Huey & Stevenson, 1979) or nearby (Martin & Huey, 2008). For metabolism,

however, more is not necessarily better, and peak rates of standard oxygen consumption represent

elevated minimum costs, typically occurring at warmer temperatures than peak performance traits

(Pörtner, 2001). For example, aerobic scope is maximized where the difference between standard

metabolic rate (SMR) and maximum metabolic rate (MMR) is greatest (Fig. 1A; often the location of

peak performance T_{OPT}) because maintenance costs are minimized relative to the peak demand (Clark,

Sandblom, & Jutfelt, 2013). At higher temperatures, when SMR meets MMR, a metabolic depression

often follows as organismal functions decline (Fig. 1; also see Pörtner, 2002). Presumably, these

temperatures lie above the maximum temperatures typically experienced by the organism and are likely to

be stressful. Thus, the metabolic peak should lie to the right of T_{OPT} (i.e., at a higher temperature than

T_{OPT}), align with decreasing performance on the TPC, and should be interpreted as the onset of stress

(Fig. 1; Pörtner, 2001). We refer to this peak as $T-MR_{PEAK}$.

The specific shapes of metabolic rate curves found for different species, populations, and thermal

experiences (e.g., acclimation; DeLong et al., 2018; Havird et al., 2020) could reflect either the acute

effects of temperature on biological rates (Dillon et al., 2010; Payne & Smith, 2017) or the evolutionary

response to past climatic conditions (Dell, Pawar, & Savage, 2011; Deutsch et al., 2008). If the shape of

the metabolic rate TPC simply reflects the thermodynamics of biological reactions, we would predict that

organisms of equivalent size experiencing a similar range of temperatures should exhibit similar

metabolic rate curves regardless of their evolutionary history (Payne & Smith, 2017).

Alternatively, the Climate Variability Hypothesis posits that if stable climatic regimes, such as

those found in the tropics, expose organisms to only a narrow range of temperatures and favor the

evolution of thermal specialists (*sensu* Huey & Kingsolver, 1989), then we predict that those organisms

should have lower SMR within the range of temperatures they experience to maximize the allocation of

energy to growth, reproduction, and aerobic performance (Fig. 1B). These thermal specialists should also

be more sensitive to temperature, i.e., have metabolic rates that change rapidly in response to temperature

outside their normal range. Climates characterized by large seasonal temperature variation, such as those

found at temperate latitudes, should select for thermal generalists capable of functioning effectively over

a wide range of temperatures (Angilletta et al., 2002; Gilchrist, 1995; Huey & Kingsolver, 1993). Thermal

generalists are predicted to have relatively low SMRs across a wider range of temperatures (Fig. 1C) and

be less sensitive to increasing temperature to keep their minimum energy demands low. The generality of

these contrasting hypotheses remains unresolved because few comparative studies have measured the

thermal sensitivity of metabolic rates in species from temperate and tropical regions under similar

experimental conditions.

To test predictions generated by these alternative hypotheses, we compared standard metabolic

rates of temperate and tropical mayflies and stoneflies across different elevations. Because stream

temperatures at a given elevation were similar at the time of the experiments, we predicted that the

metabolic rate curves should look similar between temperate and tropical species if they were strictly

determined by thermodynamic effects on metabolic processes (Payne & Smith, 2017). Alternatively, if

metabolic rates reflect adaptation to climate, we predicted greater sensitivity to temperature in the tropical

species (Fig. 1B) compared to temperate ones (Fig. 1C; Angilletta et al., 2002; Huey & Kingsolver,

1993).

Methods

 Site and species descriptions

Temperate sites were located in the Cache La Poudre river drainage of the southern Rocky Mountains in

Colorado, USA, and tropical sites in the Papallacta-Quijos river drainage of the Andes Mountains in Napo

Province, Ecuador. We chose small, relatively pristine streams where the mayflies and stoneflies used in this

study are typically found. Sites at both latitudes were matched by elevation, where high >3,000 m, mid = 2,500 –

3,000 m, and low = 1,000 – 2,500 m. Streams within each elevation pair were approximately the same

temperature at the time of collection (Table 1). Thus, we were able to make latitudinal comparisons of insect

metabolic rate for each elevation pair under the same thermal acclimation conditions.

Ephemeroptera (mayflies) and Plecoptera (stoneflies) do not share genera between tropical and temperate

sites. Therefore, we selected representative species from the same family (i.e., mayfly family Baetidae and

stonefly family Perlidae) at each location to contrast their metabolic rate. Based on our previous DNA-barcoding

work (Gill et al., 2016; Polato et al., 2018; Shah et al., 2017b), we could identify taxa to species using

morphological descriptions of individuals from these sites. Ultimately, we chose these taxa because 1) they

represent the most closely related phylogenetic temperate and tropical relatives, 2) they are ecologically and

morphologically very similar to each other, and 3) they tend to be common in streams around the world. Results

from our work could therefore have general implications.

In the Rockies, we collected 313 mayflies from the genus *Baetis* (*Baetis bicaudatus* at high and mid

elevations and *Baetis tricaudatus* at low elevations) and 107 stoneflies (*Hesperoperla pacifica* at low and mid-

elevations). In the Andes, we collected 245 mayflies from the genus *Andesiops*. Earlier work has shown cryptic

diversity in this genus, such that *Andesiops* collected at each elevation were putatively different species (Gill et

al., 2016). We collected 76 stoneflies from the genus *Anacroneuria*, which comprises *Anacroneuria rawlinsi* at

low elevation and *Anacroneuria guambiana* at mid-elevation. Although baetid mayflies were easily collected

across all elevations, perlid stoneflies were much harder to find and were altogether absent at high elevation, so

latitudinal comparisons between stoneflies are at mid and low elevations only. We conducted experiments

between January and April of 2013 in the Andes and June to August of 2013 in the Rockies.

Table 1. Average stream temperatures with standard errors and acclimation temperatures for each
temperate (Rocky Mountain) and tropical (Andes) stream from which insects were collected.

Latitude	Site ID	Type	Elevation (m)	Stream temperature (ºC)	Acclimation temperature (ºC)
Temperate	COP 2212	Low	2212	12.5 ± 3.73	14.0
	COP 2590	Mid	2590	10.8 ± 4.68	10.0
	COP 2798	Mid	2798	7.6 ± 1.41	10.0
	COP 3166	High	3166	6.2 ± 3.56	6.0
Tropical	ECP 1845	Low	1845	13.5 ± 0.72	14.0
	ECP 2003	Low	2003	14.0 ± 0.38	14.0
	ECP 2694	Mid	2694	10.0 ± 0.79	10.0
	ECP 3683	High	3683	8.6 ± 0.89	6.0
	ECP 3898	High	3898	6.6 ± 1.74	6.0

Lab acclimation

Field-caught nymphs were brought back to the lab (~1,600 m at both latitudes) and placed in

containers of filtered stream water from their native streams without food for 48 h. During this time, high

oxygen levels were maintained by bubbling air into the containers, and water was circulated by small

pumps, with temperatures approximating the insects' home stream temperatures (Table 1). Specifically,

acclimation temperatures were chosen to closely match the average temperature of the temperate and

tropical stream pair at each elevation. Temperate and tropical high elevation insects were acclimated to

6.0 °C, mid-elevation insects to 10.0 °C, and low elevation insects to 14.0 °C (Table 1). Acclimation

temperatures did not exactly match average stream temperatures (Table 1) due to the difficultly of

maintaining precise temperatures in large water baths with our heating and cooling equipment, especially

in the tropics. Notwithstanding these limitations, the difference in acclimation versus average stream

temperature was between 0 - 2.6 °C at any given site. We selected a paired elevation design because

tropical insects failed to acclimate to temperatures outside average temperatures of their native streams

(see also Shah, Funk, & Ghalambor, 2017a). For example, we often detected evidence of stress (e.g.,

jerking movements, unnatural swimming patterns) and even mortality, especially when tropical mayflies

from high and low elevations were warmed for 48 h outside native stream temperature (i.e., 12 °C and 10

°C, respectively). Therefore, the use of a single acclimation temperature across all elevations was not

possible. Because acclimation temperatures differed among elevations but were the same for each

elevation pair, we restricted our statistical comparisons to the same elevations between the two latitudes.

Metabolic rate

Oxygen consumption rates (metabolic rates, hereafter) were measured using closed glass micro-

respirometry chambers (Unisense A/S, Aarhus, Denmark). After the 48 h acclimation period, we

introduced insects randomly to one of nine test temperatures (5, 7.5, 10, 12.5, 15, 17.5, 20, 22.5, or 25 °C)

by allowing temperature to rise naturally or by slowly cooling (0.5 °C · min^{-1}) to prevent thermal shock.

On average, 16 individuals per population for mayflies (min = 3, max = 22) and 6 per population for

stoneflies (min = 3, max = 12) were tested at each temperature (Tables S1, S2). Each individual was

tested at only one temperature. Once at the test temperature, insects were moved individually into glass

chambers filled with stream water that had been treated with UV light to reduce background microbial

respiration and immersed in a water bath held at the test temperature (Fig. S1). Mayflies were placed in 4

mL chambers, and stoneflies, which are larger, were placed in 50 mL chambers. Chambers were fitted

with a glass-coated magnetic stir bar and a plastic mesh to which the insect could cling (Fig. S1). We

acclimated insects to the chamber space for an additional hour before sealing each chamber with a glass

stopper to prevent further introduction of atmospheric oxygen. Each stopper had a central capillary hole,

allowing us to insert an oxygen microelectrode (500 μm diam.) connected to a picoammeter (Unisense

OXY-Meter, Aarhus, Denmark) to measure change in oxygen concentration. We measured background

respiration rates by placing a control chamber (without an insect) in each experiment. The microelectrode

was first lowered into the control chamber at the start of the experiment. After reading oxygen

concentration continuously for 2 mins, it was moved to the first chamber that contained an animal, where

data were recorded for another 2 mins (Fig. S1), and so on. Experiments were terminated when oxygen

levels reached 80% of the air-saturated values because natural dissolved oxygen levels in our study sites

rarely fall below this value. Consequently, we made 3-6 readings for each chamber – fewer readings at

the higher temperatures due to the combined effect of lower oxygen concentrations and faster oxygen

consumption by insects – which lasted around 128 minutes. Because there were typically 16 chambers in

each experiment, the average time interval between readings was 25 mins. While recording oxygen

concentration, we also observed insects inside the chambers, noting any visible behavioral stress

responses or mortality. Insects were then removed from chambers, euthanized in ethanol, dried, and

weighed on a laboratory scale with readability to 0.01mg.

Statistical Analyses

We first calculated average rate of oxygen consumption for each insect or control (blank)

chamber by measuring change in oxygen concentration across all measurements taken in a given chamber

(see Supplementary Materials and Fig. S2 for an explanation of this calculation). We then averaged the

control values for each temperature such that we calculated a control rate for each combination of test

temperature, elevation, and latitude. The mean control rates were subtracted from individual insect

consumption rates to correct for background microbial respiration at that temperature. We conducted all

statistical analyses on these 'control-corrected oxygen consumption rates' (hereafter, oxygen

consumption) in R version 3.4.0 (R Core Team 2017). For each taxon, latitudinal contrasts were

conducted separately for each elevation category. Thus, for mayflies, we compared temperate and tropical

baetids in three statistical comparisons: low, mid, and high elevation. Similarly, for stoneflies, we ran two

separate analyses for low and mid elevation.

Because we were interested in the thermal sensitivity of temperate and tropical insect metabolic

rates between elevation pairs (i.e., the thermal reaction norm), we first compared metabolic rates during

the ascending phase of the TPC by fitting a linear model with oxygen consumption as the response

variable for each elevation type (i.e., high, mid, and low) separately. For this analysis, we used only the

temperatures 5, 7.5, 10, 12.5, 15 and 17.5 °C because these temperatures were within the ascending phase

of the SMR curves for all populations in this study. The model included fixed effects of test temperature

and latitude, and their interaction, with body mass as a covariate (Packard & Boardman, 1999). A

significant interaction would denote a difference in thermal sensitivity (i.e., slope) between temperate and

tropical species.

We conducted a similar analysis for the descending phase using data from the 17.5 and 20 °C

treatments as these two temperatures encompassed the descending phase of the curve. At the highest

temperatures (22.5 and 25 °C), we observed stress responses (e.g., unnatural movement, frenzied attempts

to escape from the chamber and in some cases elevated metabolic rates) in tropical mayflies. We therefore

did not include SMR at 22.5 and 25 °C in this analysis but instead calculated separate Q_{10} values for these

temperatures (see below).

To compare the maximum SMR at each elevation between temperate and tropical taxa, we

examined the latitude x elevation combination individually to find the physiologically relevant peak (i.e.,

before CT_{MAX} and when the first signs of stress were apparent, if observed). This peak was typically at

17.5 °C in our study because in most cases, across both latitudes, SMR was highest at this temperature

even if not always statistically different from 15 °C. We then compared these values using an ANOVA

for each elevation pair using latitude as the fixed effect and body mass as a covariate. For all analyses, we

plotted the least square means to control for the effect of body mass on standard metabolic rates.

Finally, we calculated Q_{10}, a measure of sensitivity to temperature, using the formula

267
$$Q_{10} = \left(\frac{R_2}{R_1}\right)^{\left(\frac{10}{T_2 - T_1}\right)}$$

where R_1 and R_2 are reaction (oxygen consumption) rates measured at two different temperatures, T_1 and

T_2, respectively. Q_{10}s were calculated across cooler, 'non-stressful' temperatures ($T_1 = 7.5$ and $T_2 = 15$ °C)

to ensure measurements were within the exponential phase of the metabolic curve and warmer, 'stressful'

temperatures ($T_1 = 15$ and $T_2 = 25$ °C) to capture the relative decline in rates.

Results

Behavioral observations

Tropical mayflies showed visible signs of stress (jerking movements, loss of righting response) at

several test temperatures outside the range typically experienced (e.g., 22.5 and 25 °C; Fig. 2 red bars,

Table S1; Fig. S3A); many of these visibly stressed individuals died at the end of the experiment (Table

S. 1; Fig. S3B). Low elevation tropical mayflies also experienced some mortality at 5 °C, which is much

colder than the minimum temperatures they encounter in their home streams (Fig. 2F). Temperate

mayflies, by contrast, did not show visible signs of stress at any experimental temperature (Table S2). We

also did not detect any overt signs of stress in any of the tropical or temperate stoneflies in this study.

Figure 2. Plots of   flies from high (A, B), mid
(C,D), and low (E   t panel) streams. Red bars
indicate temperatu   ith yellow arrows for each
population. Gray r   e measured in each stream.

CT$_{MIN}$ and CT$_{MAX}$

Figure 3. Plots of metabolic rate least square means in perlid stoneflies from mid (A, B) and low (C,D) elevations in temperate (left panel) and tropical (right panel) streams. Stoneflies were not found at high elevation. T-MR$_{PEAK}$ is shown with yellow arrows for each population. Gray rectangles represent the approximate stream temperature range measured in each stream. We did not detect overt stress in stoneflies in this study. CT$_{MIN}$ and CT$_{MAX}$ determined in a previous study are shown.

Thermal sensitivity of standard metabolic rates and reaction norms

SMR profiles differed between temperate and tropical mayflies and stoneflies at many elevations (Figs. 2, 3). In general, metabolic rates of mayflies from the two latitudes were lowest at temperatures approximating those in their native streams. Outside these temperatures, however, tropical mayflies exhibited a greater increase in metabolic rate (Fig. 2). In all mayfly populations across both latitudes, a secondary peak was observed at 22.5 and 25 °C that corresponded with visible stress during experiments. We denoted the initial peak in SMR as T-MR $_{PEAK}$ (shown as yellow arrows on Figs. 2 and 3) or the onset of stress, which occurred at temperatures higher than maximum stream temperature, as expected.

Secondary peaks were viewed as an acute stress response because they were followed by death in some

cases. As indicated in the Methods, we excluded these stressful test temperatures from the analyses of the

ascending and descending phases of metabolic rate profiles and dealt with them separately in an analysis

of Q_{10}.

In the ascending phase, metabolic rates of tropical mayflies were more sensitive to temperature

increases than their temperate counterparts at high elevation ($F_{(1,105)} = 13.79$, $p < 0.001$, Fig. 4A) and mid

elevation ($F_{(1,105)} = 11.95$, $p = 0.001$, Fig. 4B). At low elevation, despite high mortality and visible signs

of stress in tropical but not temperate mayflies, we did not find a significant interaction between tropical

and temperate groups ($F_{(1,159)} = 1.505$, $p = 0.222$, Fig. 4C). In stoneflies, ascending phases did not differ

between temperate and tropical populations (mid elevation: $F_{(1,79)} = 2.363$, $p = 0.128$; Fig. 5A and low

elevation: $F_{(1,61)} = 0.136$, $p = 0.714$; Fig. 5B)

When comparing descending phases, temperate mayflies had a marginally steeper decrease in

metabolic rate at high elevation (Fig. 4A; $F_{(1,38)} = 4.044$, $p = 0.051$) but tropical mayflies had a much

steeper decrease at low elevation (Fig. 4C; $F_{(1,28)} = 9.464$, $p = 0.005$). At mid elevation, there was no

significant interaction ($F_{(1,35)} = 2.477$, $p = 0.125$; Fig. 4B). However, a Student's t-test revealed that

oxygen consumption values were significantly higher in mid-elevation tropical mayflies compared to

temperate mayflies at 17.5 °C ($t = 3.946$, $p = 0.001$), but did not differ between latitudes at 20 °C ($t = -$

1.006, $p = 0.33$). This suggests that the descending phase was steeper for tropical mayflies compared to

temperate ones. Descending phases of stoneflies did not differ between tropical and temperate populations

(Figs. 5 A & B).

Finally, maximum SMR was higher in tropical compared to temperate mayflies at all elevations

(Fig. 6). Stoneflies showed an opposite pattern where tropical species had lower peaks at both measured

elevations, although this was significant only at mid elevation (Fig. 6).

Figure 4. A comparison of the ascending (left panel) and descending (right panel) phases of metabolic profiles in baetid mayflies at high (A), mid (B) and low (C) elevations. Plotted values are the least square means of metabolic rate in mayflies from tropical (orange circles) and temperate (blue circles) streams, after accounting for body mass in the model. Vertical bars are standard errors.

Figure 5. A comparison of the ascending (left panel) and descending (right panel) phases of metabolic profiles in perlid stoneflies at mid (A), and low (B) elevations. Plotted values are the least square means of metabolic rate in stoneflies from tropical (orange circles) and temperate (blue circles) streams, after accounting for body mass in the model. Vertical bars are standard errors.

Thermal sensitivity based on Q_{10}

When comparing across 'stressful' and 'non-stressful' temperatures, we found Q_{10}s for both taxa to be higher at non-stressful temperatures than at stressful temperatures (Table 2). Across latitude, tropical mayflies showed generally greater sensitivity to temperature (higher Q_{10}s) at both non-stressful and stressful temperatures than their temperate counterparts (Table 2; Fig. 7). Notably, at mid and low elevations, tropical mayflies were highly sensitive to temperature even at non-stressful temperatures (Q_{10}

= 8.40 and 5.17, respectively). In contrast, tropical stoneflies were less sensitive than their temperate

relatives for stressful and non-stressful temperatures (Fig. 7).

Figure 6. Peak SMRs (i.e., the highest metabolic rate) are compared between temperate and tropical
baetid mayflies from high (A), mid (B), and low (C) elevations and perlid stoneflies from mid (D) and
low (E) elevations. Plotted values are the least square means of metabolic rate for tropical (orange circles)
and temperate (blue circles) insects, after accounting for body mass in the model. Vertical bars are
standard errors and p values from an ANOVA for each elevation pair are shown.

Table 2. Q_{10} values calculated for temperate and tropical mayfly and stonefly metabolic rates between typically non-stressful temperatures (7.5°C to 15°C) and typically stressful temperatures (15°C to 25°C).

		Mayflies		*Stoneflies*	
	Elevation	Temperate	Tropical	Temperate	Tropical
	High	3.63	3.15	n/a	n/a
Non-stressful	Mid	3.15	5.17	6.76	2.49
	Low	2.36	8.40	6.66	5.78
	High	1.07	1.46	n/a	n/a
Stressful	Mid	0.73	1.59	6.10	0.87
	Low	1.34	1.67	1.25	0.42

Figure 7. Comparison of average Q_{10}s for mayflies and stoneflies during the ascending phase of the metabolic rate profiles.

Discussion

Understanding the influence of temperature on the minimum energy demands of ectotherms is critical to predicting how global warming will impact different species (Huey & Kingsolver, 2019). Previous comparative studies of thermal tolerance (critical thermal limits) have generally found tropical ectotherms to be adapted to narrower ranges of temperatures compared to their temperate counterparts, suggesting that tropical species are more vulnerable to global warming (Deutsch et al., 2008; Gutiérrez-Pesquera et al., 2016; Shah et al., 2017b; Sunday, Bates, & Dulvy, 2011, 2012), a finding consistent with the CVH. Payne and Smith (2017) recently challenged such conclusions by arguing that narrower thermal breadth may simply reflect the acute effects of temperature on biological rates of organisms living in warmer environments. Resolving this debate requires an understanding of the acute effects of temperature on continuous biological rates like metabolism (Havird et al., 2020). Here, using a paired-elevation design, we compared the thermal sensitivity of standard metabolic rate (SMR) in closely related temperate and tropical aquatic insect nymphs. This study design enabled a comparison of closely-related

temperate and tropical species under a common range of field, acclimation, and test temperatures. Our

results did not support Payne and Smith's (2017) model, but instead supported the CVH in mayflies, but

not stoneflies.

Despite originating in streams with similar average environmental temperatures at the time of

collection, tropical and temperate insects had different metabolic rate profiles (Figs. 2, 3). These results

indicate that the variation between temperate and tropical species in our study does not simply reflect the

acute effects of temperature on biological rates, but rather that there are physiological differences between

taxa from these regions. Tropical baetid mayflies had metabolic rates that were more sensitive to

temperature, and they exhibited stress at lower temperatures compared to their temperate counterparts. By

contrast, tropical and temperate perlid stoneflies showed no strong divergence in their thermal sensitivity.

Although the ecological implications of greater thermal sensitivity remain unknown for most ectotherms,

in the absence of compensatory changes in productivity and energy expenditure, elevated metabolic rates

could negatively impact the energy budget and capacity for aerobic performance in tropical mayflies by

increasing baseline energetic costs (Dillon et al., 2010; Huey & Kingsolver, 2019).

Comparison of metabolic rate profiles

In most cases, metabolic rate rose with increasing temperature, peaked, and then fell. This

pattern, however, varied among taxa and populations and across elevations. As a group, the tropical

mayflies are more thermally sensitive than their temperate counterparts (Figs. 2, 4, 6 and 7). The

metabolic rate profiles of tropical mayflies had overall steeper ascending and descending phases, and

higher Q_{10}s. The particularly high Q_{10}s for mid and low elevation tropical mayflies were similar to those

of North Atlantic stenothermal fishes when experiencing temperatures away from an optimum (Johnston,

Clarke, & Ward, 1991). By contrast, high elevation tropical mayflies had similar Q_{10}s to their temperate

relatives (Table 2), which may reflect the higher diel variation (i.e., variation across 24 h) experienced in

streams above tree line in the Andes (higher standard error for stream temperature Table 1). In seasonally

stable streams, such diel effects may play a potentially important role in regulating metabolic activity of

aquatic insects (Gilchrist, 1995), but so far, no studies have explored the relative effects of diel and

seasonal thermal variation in freshwater insects. In general, however, our results support the prediction

that tropical mayflies, which experience more stable stream temperatures year-round (Shah et al., 2017b),

are more sensitive to temperatures outside the ranges they normally experience.

Given that mayflies were paired by elevations having similar mean environmental and

acclimation temperatures, these differences cannot be attributed to the acute or passive response of

metabolic rate to higher temperatures alone (Havird et al., 2020). Instead, they appear to reflect

divergence in thermal physiology as shown by the greater thermal sensitivity of metabolism in tropical

mayflies. In contrast to tropical mayflies, temperate mayflies exhibited more modest increases in

metabolism with temperature and had similar metabolic rate profiles across elevations (Fig. 2). We note

that although there was no statistically significant interaction for the ascending limb between temperate

and tropical mayflies at low elevation (Fig. 4), possibly driven by very low SMR at 7 ℃, tropical

mayflies at low elevation exhibited strong stress responses to moderate temperatures (Fig. 2). Further,

they have previously shown sensitivity to temperature with lower critical maximum temperatures and

reduced acclimation responses (Shah et al., 2017a, 2017b). For both temperate and tropical mayflies,

metabolic rates tended to be lower within the range of environmental temperatures they naturally

experienced, suggesting there may be selection to reduce the minimum energetic costs associated with

maintenance (Artacho & Nespolo, 2009; Bochdansky, Grønkjær, Herra, & Leggett, 2005; Brown,

Marquet, & Taper, 1993). For example, even a moderate increase in stream temperature outside the

normal range (Fig. 2, right edge of grey rectangles) is associated with peaks in metabolism, which has

potentially cascading effects on energy budgets and possible chronic stress (Fig. 2). If selection acts to

minimize SMR under typical stream temperatures (Brown et al., 1993), these results indicate that the

narrower, more stable tropical stream temperatures make tropical mayflies more stenothermic and

underlie their greater thermal sensitivity, as predicted by the CVH.

Such an interpretation is consistent with other aspects of temperate and tropical mayfly thermal

physiology. First, thermal breadths (measured as the difference between critical thermal limits) reported

in a parallel study were narrower for tropical (Ecuador) versus temperate (U.S.A) mayfly populations

(Shah et al., 2017b). Second, in this study, we observed signs of stress in the metabolic rate profiles of

tropical but not temperate mayflies. Specifically, after the decline in metabolic rate, secondary peaks

occurred at 22.5 °C and 25 °C for most groups, which corresponded to observed signs of behavioral stress

and mortality in tropical but not temperate mayflies. Secondary peaks are relatively rare in metabolic rate

measurements but can occur, often near the critical thermal limits (Havird et al., 2020). Mayflies showed

signs of behavioral stress at these high temperatures, including leg spasms, loss of righting, release from

substrate, and mortality (see Supplemental Information Table S1, Figs. S3A, 3B). Collectively, these

results support the idea that temperate mayflies are thermal generalists, with broad thermal breadths and

reduced metabolic sensitivity to temperature, whereas tropical mayflies are thermal specialists, with

narrower thermal breadths and increased metabolic sensitivity to temperature (c.f. Huey & Hertz, 1984).

The differences observed between temperate and tropical mayflies were mostly unambiguous and

consistent with prior work in this system. Similarly, the *lack* of divergence between temperate and

tropical stoneflies is also consistent with previous research on other physiological traits (Shah et al.,

2017a, 2017b). Although there are differences in metabolic rate profiles between temperate and tropical

stoneflies (e.g., higher metabolic rates at warmer temperatures in temperate but not tropical species), these

differences are far less striking than those observed in mayflies. Ascending and descending phases did not

differ between temperate and tropical stoneflies, but absolute maximum metabolic rates and Q_{10}s across

non-stressful and stressful temperatures were higher in the temperate groups. Shah et al. (2017a) also

found a lack of latitudinal differences in stoneflies when comparing acclimation abilities. One of the few

other studies on the respiration rates of tropical stoneflies reported similar Q_{10}s to those reported here for

non-stressful temperatures in mid-elevation stoneflies (Jacobsen & Brodersen, 2008), corroborating our

conclusions of lower thermal sensitivity in at least the *Anacroneuria* genus. Though many tropical species

may be imperiled by warming (Deutsch et al., 2008; Dillon et al., 2010), the overall mixed support for

this prediction in stoneflies cautions against generalizing about the vulnerability across tropical species.

Numerous hypotheses could explain the lack of divergence between temperate and tropical

stoneflies. First, stoneflies as a group have a temperate origin and invaded tropical streams relatively

recently (Baumann & Kondratieff, 1991; Stewart, Baumann, & Stark, 1974). Studies show that stoneflies

are still adapting to tropical habitats (Polato et al., 2018), which may explain why thermal physiological

traits in present-day temperate and tropical populations are similar.

Second, perlid stoneflies and baetid mayflies have different life histories, which may contribute to

differential sensitivity to temperature. Stoneflies live much longer in streams as nymphs than mayflies

(which have multiple generations per year), and may experience years of unpredictable variation in

stream temperature (Clifford, 1982). In fact, many stoneflies enter diapause, unlike mayflies (Brittain,

1982; Hynes, 1976), giving them a greater ability to withstand adverse thermal conditions (Brittain,

1990). Once emerged, adult stoneflies often remain close to natal streams, whereas adult mayflies can

disperse more widely in tropical (Finn, Encalada, & Hampel, 2016) and temperate (Polato et al., 2018)

regions. Therefore, stoneflies rely more heavily on the nymph stage for dispersal (Brittain, 1990) and may

experience more thermal variation while moving up- or downstream than juvenile mayflies. This

increased likelihood of encountering thermal variation may favor the evolution of decreased thermal

sensitivity (Muñoz & Bodensteiner, 2019).

Third, perlid stoneflies are important predators of baetid mayflies and midges (Peckarsky, 1980;

Tamaris-Turizo, Turizo-Correa, & del Carmen Zúñiga, 2007) and they actively pursue prey in the wild

(Peckarsky, 1980) and in laboratory settings (Sjöstrom, 1985; A.A.Shah *pers. obs*.). A reliance on aerobic

metabolism and the need to maintain a high metabolic scope for active hunting (Kordas, Harley, &

O'Connor, 2011) could constrain physiological divergence between temperate and tropical stoneflies.

Unfortunately, few comparative studies of metabolic profiles are available to test the generality of such

hypotheses. Nevertheless, higher maximum metabolic rates and higher Q_{10}s suggest some differences

between temperate and tropical populations that could have ecological consequences when pursuing prey

across a broad range of temperatures. Differences in thermal sensitivity between these predators and their

prey may have community-level consequences under climate change (Gilman, Urban, Tewksbury,

Gilchrist, & Holt, 2010; Grigaltchik, Ward, & Seebacher, 2012; Kordas et al., 2011; Pincebourde &

Casas, 2019; Shah, Dillon, Hotaling, & Woods, 2020b; Vucic-Pestic, Ehnes, Rall, & Brose, 2011).

Comparison of metabolic rates, critical thermal maxima, and performance curves as indicators of

vulnerability

One should expect CT_{MAX}, a commonly used metric for thermal sensitivity (Diamond et al., 2012;

Lutterschmidt & Hutchison, 1997; Piyaphongkul, Pritchard, & Bale, 2012; Ribeiro, Camacho, & Navas,

2012), to coincide with the descending limb of the metabolic rate profiles (Fig. 1). Yet, previously

measured values of CT_{MAX} in temperate mayflies occur at much higher temperatures (Fig.1A, B, C). For

example, CT_{MAX} was ~29 °C in high elevation temperate baetid mayflies (Shah et al., 2017b) but they

began to show elevated metabolic rates and behavioral signs of stress at lower temperatures (~17.5 °C).

The large difference between CT_{MAX} and the onset of stress in this study is likely due to methodology, as

rapid ramping or high starting temperatures can inflate estimated values of CT_{MAX} (Rezende, Tejedo, &

Santos, 2011; Terblanche, Deere, Clusella-Trullas, Janion, & Chown, 2007). In addition, the discrepancy

could also arise because CT_{MAX} and metabolic rates are governed by fundamentally different processes.

CT_{MAX} is the result of changes in fluidity of cell membranes and whole-organism physiology in response

to heat stress (Cossins, Friedlander, & Prosser, 1977; Hochachka & Somero, 2002), whereas standard

metabolic rates arise from different activation energies of enzymes and sensitivity of biochemical

processes to temperature (Rolfe & Brown, 1997). Thus, using CT_{MAX} in conjunction with other measures

of thermal sensitivity such as metabolic rate provides a more complete assessment of thermal tolerance.

Another commonly measured indicator of thermal sensitivity is locomotor performance, in which

T_{OPT} should lie at cooler temperatures than $T\text{-}MR_{PEAK}$ (Fig. 1). However, in a separate study of swimming

performance across temperature in the same mayfly species, we found no clear T_{OPT}. Instead, TPCs for

burst swimming were relatively flat in mayflies, with the exception of low elevation populations (Shah,

Bacmeister, Rubalcaba, & Ghalambor, 2020a). Although it remains unclear why performance does not

vary with temperature, one possible explanation is that selection favors a generalist strategy for

swimming performance in all mayflies, perhaps because there is more thermal variation in streams than

expected (Gilchrist, 1995). Alternatively, TPCs may not vary because burst swimming is less sensitive to

temperature than more aerobically demanding activities like sustained swimming. Unlike most freshwater

fishes, baetid mayflies do not engage in sustained swimming, but instead, walk and cling to rocks where

their flattened bodies decrease shear stress from the water current (Weissenberger, Spatz, Emanns, &

Schwoerbel, 1991). Thus, the relationship between temperature, SMR, and the allocation of energy to

locomotion and other demands such as growth, reproduction, and maintenance remains an important area

for future research.

Climate change in Andean and Rocky Mountain Streams

Mountain stream temperatures are already increasing worldwide (Isaak & Rieman, 2013; Niedrist

& Füreder, 2020). Temperatures of shallow mountain streams, such as those in this study, are also likely

to show concomitant increases with air temperature (Birrell et al., 2020; Mohseni & Stefan, 1999;

Pilgrim, Fang, & Stefan, 1998). Projected mean air temperature increases of 2-3 °C in the Andes and 3-4

°C in the northern Rocky Mountains by the 2050s (Nogués-Bravo, Araújo, Errea, & Martínez-Rica, 2007;

Rangecroft, Suggitt, Anderson, & Harrison, 2016) are therefore troubling. However, previous analyses

indicate that aquatic insects of the southern Rocky Mountains are relatively robust to warming because

they may currently occur at suboptimal temperatures (Pyne & Poff, 2017). Moreover, mesocosm and field

common garden experiments showed that temperate baetid mayflies grew faster when subjected to

warmer water (A. Landeira-Dabarca et al.; A. Rugenski et al.; *unpubl. data*). By contrast, Andean mayfly

growth declined with increasing temperature in mesocosms (A. Landeira-Dabarca et al.; *unpubl. data*)

further supporting our conclusion that a warming environment will have greater negative impacts on

tropical mayflies, especially at low elevation.

The effects of increasing temperatures may also lead to mismatches between dissolved oxygen

supply and demand (Pörtner, 2001; Verberk, Bilton, Calosi, & Spicer, 2011; Verberk et al., 2016b). For

many aquatic insects, the ability to maintain an adequate rate of oxygen uptake decreases at higher

temperatures (Jacobsen & Brodersen, 2008; Rotvit & Jacobsen, 2013), and reduces energy available for

activities like movement, growth, and egg development. Thus, even small increases in stream temperature

can negatively impact insect metabolism and growth. This is illustrated in a field experiment, in which a

moderate 2 °C temperature increase in a stream resulted in declines in abundance of Eurasian mayflies,

presumably as a result of decreased oxygen in the stream (Verberk, Durance, Vaughan, & Ormerod,

2016a).

Finally, between the aquatic nymph stage and the flying adult stage there is potential for dispersal

to mitigate the effects of future warming. Thermally sensitive tropical mayflies may seek cooler

temperatures at higher elevations (Jacobsen, 2020), but whether the taxa in our study can move upstream

as nymphs is debatable given the low dispersal rates of aquatic insects in these streams (Polato et al.,

2018). Further, it is not known whether aquatic insects can mitigate the effects of warming by seeking out

microhabitats that supply oxygen at high rates, i.e., those with relatively high oxygen concentrations or

faster flows. Future work should therefore also focus on whether aquatic insects can move upstream to

cooler waters or seek out high flow microhabitats to counteract the effects of warming in their native

streams (Shah et al., 2020b; Sheldon, 2012).

Acknowledgements

We thank the U.S. National Science Foundation for funding through the Dimensions of Biodiversity grant

DEB-1046408; DEB-1045960, and DEB-1045991; and a Graduate Research Fellowship awarded to

A.A.S under DGE-1321845. This research would not have been possible without the hard work of many

field assistants. We especially thank M. Rojas, J. Fajardo, G. Waneka, B. Choat, L. Granizo, and L. Nagle

for joining us over multiple field seasons. We are extremely grateful to J. Schreckinger, N. Quiroz, M.

Celinscak, and E.J.G. Orejuela in Ecuador and K. L. Casner in the U.S. for help with field logistics. The

Ministerio del Ambiente of Ecuador provided collection permits (Permit nos. 56-IC- FAU/FLO-DPN/MA

and MAE-DNB- CM-2015-0017).

Conflicts of Interest

NONE

Author contributions:

A.A.S. and C.K.G. designed the experiments. A.A.S. conducted fieldwork and experiments, analyzed

results, and wrote the manuscript with contributions from all authors.

Data sharing and accessibility

Data generated in this study are openly available in the supporting information files of this article and on

the public repository GitHub (https://github.com/Alisha-Shah/Standard-metabolic-rates.git).

References

Angilletta, M. J. (2009). *Thermal adaptation: a theoretical and empirical synthesis.* New York: Oxford

University Press.

Angilletta, M. J., Niewiarowski, P. H., & Navas, C. A. (2002). The evolution of thermal physiology in

ectotherms. *Journal of Thermal Biology*, *27*(4), 249–268. https://doi.org/10.1016/S0306-

4565(01)00094-8

Artacho, P., & Nespolo, R. F. (2009). Natural selection reduces energy metabolism in the garden snail,

Helix Aspersa (Cornu Aspersum). *Evolution*, *63*(4), 1044–1050. https://doi.org/10.1111/j.1558-

5646.2008.00603.x

Baumann, R. W., & Kondratieff, B. C. (1991). The stonefly fauna of northern Mexico. In J. Alba-

Tercedor & A. Sanchez-Ortega (Eds.), *Overview and Strategies of Ephemeroptera and Plecoptera*.

Gainesville, Florida.: Sandhill Crane Press, Inc.

Birrell, J. H., Shah, A. A., Hotaling, S., Giersch, J. J., Williamson, C. E., Jacobsen, D., & Woods, H. A.

(2020). Insects in high-elevation streams: life in extreme environments imperiled by climate change.

Global Change Biology, in press. https://doi.org/10.1111/gcb.15356

Bochdansky, A. B., Grønkjær, P., Herra, T. P., & Leggett, W. C. (2005). Experimental evidence for

selection against fish larvae with high metabolic rates in a food limited environment. *Marine*

Biology, *147*(6), 1413–1417. https://doi.org/10.1007/s00227-005-0036-z

Brittain, J. E. (1982). Biology of mayflies. *Annual Review of Entomology*, *27*(1), 119-147.

https://doi.org/10.1146/annurev.en.27.010182.001003

Brittain, J. E. (1990). Life history strategies in Ephemeroptera and Plecoptera. In *Mayflies and Stoneflies:*

Life Histories and Biology (pp. 1–12). https://doi.org/10.1007/978-94-009-2397-3_1

Brown, J. H., Marquet, P. A., & Taper, M. L. (1993). Evolution of body size: consequences of an

energetic definition of fitness. *The American Naturalist*, *142*(4), 573–584.

Chown, S. L., Duffy, G. A., & Sørensen, J. G. (2015). Upper thermal tolerance in aquatic insects. *Current*

Opinion in Insect Science, *11*, 78–83. https://doi.org/10.1016/j.cois.2015.09.012

Clark, T. D., Sandblom, E., & Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of

climate change: respirometry, relevance and recommendations. *Journal of Experimental*

Biology, *216*(15), 2771-2782. https://doi.org/10.1242/jeb.084251

Clifford, H. F. (1982). Life cycles of mayflies(Ephemeroptera), with special reference to voltinism.

Quaestiones Entomologicae, *18*.(1), 15–90.

Colwell, R. K., Brehm, G., Cardelús, C. L., Gilman, A. C., & Longino, J. T. (2008). Global warming,

elevational range shifts, and lowland biotic attrition in the wet tropics. *Science*, *322*(5899), 258–261.

https://doi.org/10.1126/science.1162547

Cossins, A. R., Friedlander, M. J., & Prosser, C. L. (1977). Correlations between behavioral temperature

adaptations of goldfish and the viscosity and fatty acid composition of their synaptic membranes.

Journal of Comparative Physiology, *120*(2), 109–121.

Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature dependence of

physiological and ecological traits. *Proceedings of the National Academy of Sciences 108*(26),

10591–10596. https://doi.org/10.1073/pnas.1015178108

DeLong, J. P., Bachman, G., Gibert, J. P., Luhring, T. M., Montooth, K. L., Neyer, A., & Reed, B.

(2018). Habitat, latitude and body mass influence the temperature dependence of metabolic rate.

Biology Letters, *14*(8). https://doi.org/10.1098/rsbl.2018.0442

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin,

P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of*

the National Academy of Sciences, *105*(18), 6668–6672. https://doi.org/10.1073/pnas.0709472105

Diamond, S. E., Sorger, D. M., Hulcr, J., Pelini, S. L., Toro, I. Del, Hirsch, C., … Dunn, R. R. (2012).

Who likes it hot? A global analysis of the climatic, ecological, and evolutionary determinants of

warming tolerance in ants. *Global Change Biology*, *18*(2), 448–456.

https://doi.org/10.1111/j.1365-2486.2011.02542.x

Dillon, M. E., Wang, G., & Huey, R. B. (2010). Global metabolic impacts of recent climate warming.

Nature, *467*(7316), 704–706. https://doi.org/10.1038/nature09407

Finn, D. S., Encalada, A. C., & Hampel, H. (2016). Genetic isolation among mountains but not between

stream types in a tropical high-altitude mayfly. *Freshwater Biology*, *61*(5), 702–714.

https://doi.org/10.1111/fwb.12740

Gamboa, M., Chacón, M. M., & Segnini, S. (2009). Diet composition of the mature larvae of four

Anacroneuria species (Plecoptera: Perlidae) from the Venezuelan Andes. *Aquatic Insects*, *31*(1),

409–417 https://doi.org/10.1080/01650420802643537

Gilchrist, G. W. (1995). Specialists and generalists in changing environments. Fitness landscapes of

thermal sensitivity. *American Naturalist*, *146*(2), 252–270.

Gill, B. A., Kondratieff, B. C., Casner, K. L., Encalada, A. C., Flecker, A. S., Gannon, D. G., … Funk, W.

C. (2016). Cryptic species diversity reveals biogeographic support for the 'mountain passes are

higher in the tropics' hypothesis. *Proceedings of the Royal Society B: Biological Sciences*,

283(1832). https://doi.org/10.1098/rspb.2016.0553

Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., & Charnov, E. L. (2001). Effects of size and

temperature on metabolic rate. *Science*, *293*(5538), 2248–2251. DOI: 10.1126/science.1061967

Gilman, S. E., Urban, M. C., Tewksbury, J., Gilchrist, G. W., & Holt, R. D. (2010). A framework for

community interactions under climate change. *Trends in Ecology and Evolution*, *25*(6), 325–331.

https://doi.org/10.1016/j.tree.2010.03.002

Grigaltchik, V. S., Ward, A. J. W., & Seebacher, F. (2012). Thermal acclimation of interactions:

differential responses to temperature change alter predator–prey relationship. *Proceedings of the*

Royal Society B: Biological Sciences, *279*(1744), 4058–4064.

https://doi.org/10.1098/rspb.2012.1277

Guppy, M., & Withers, P. (1999). Metabolic depression in animals⸱: physiological perspectives and

biochemical generalizations. *Biological Reviews*, *74*, 1–40.

https://doi.org/10.1111/j.1469-185X.1999.tb00180.x

Gutiérrez-Pesquera, L. M., Tejedo, M., Olalla-Tárraga, M. Á., Duarte, H., Nicieza, A., & Solé, M. (2016).

Testing the climate variability hypothesis in thermal tolerance limits of tropical and temperate

tadpoles. *Journal of Biogeography*, *43*(6), 1166–1178. https://doi.org/10.1111/jbi.12700

Havird, J. C., Neuwald, J. L., Shah, A. A., Mauro, A., Marshall, C. A., & Ghalambor, C. K. (2020).

Distinguishing between active plasticity due to thermal acclimation and passive plasticity due to Q10

effects: Why methodology matters. *Functional Ecology*, 1–14. https://doi.org/10.1111/1365-

2435.13534

Hochachka, P. W., & Somero, G. N. (2002). Mechanism and process in physiological evolution. In

Biochemical Adaptation. New York: Oxford University Press.

Huey, R. B., & Hertz, P. E. (1984). Is a Jack-of-all-temperatures a master of none? *Evolution*, *38*(2), 441–

444. https://doi.org/10.2307/2408502

Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm performance.

Trends in Ecology & Evolution, *4*(5), 131–135. https://doi.org/10.1016/0169-5347(89)90211-5

Huey, R. B., & Kingsolver, J. G. (1993). Evolution of resistance to high temperature in ectotherms. *The*

American Naturalist, *142*.

Huey, R. B., & Kingsolver, J. G. (2019). Climate warming, resource availability, and the metabolic

meltdown of ectotherms. *The American Naturalist*, *194*(6), E140–E150.

https://doi.org/10.1086/705679

Huey, R. B., & Stevenson, R. D. (1979). Integrating thermal physiology and ecology of ectotherms: a

discussion of approaches. *American Zoologist*, *19*(1), 357–366. https://doi.org/10.1093/icb/19.1.357

Hynes, H. B. N. (1976). Biology of Plecoptera. *Annual Review of Entomology*, *21*(1), 135–153.

https://doi.org/10.1146/annurev.en.21.010176.001031

Isaak, D. J., & Rieman, B. E. (2013). Stream isotherm shifts from climate change and implications for

distributions of ectothermic organisms. *Global Change Biology*, *19*(3), 742–751.

https://doi.org/10.1111/gcb.12073

Jacobsen, D. (2020). The dilemma of altitudinal shifts: caught between high temperature and low oxygen.

Frontiers in Ecology and the Environment, 18(4), 211-218. https://doi.org/10.1002/fee.2161

Jacobsen, D., & Brodersen, K. P. (2008). Are altitudinal limits of equatorial stream insects reflected in

their respiratory performance? *Freshwater Biology*, *53*(11), 2295–2308.

https://doi.org/10.1111/j.1365-2427.2008.02050.x

Johnston, I. A., Clarke, A., & Ward, P. (1991). Temperature and metabolic rate in sedentary fish from the

Antarctic, North Sea and Indo-West Pacific Ocean. *Marine Biology*, *109*(2), 191–195.

https://doi.org/10.1007/BF01319386

Kellermann, V., Chown, S. L., Schou, M. F., Aitkenhead, I., Janion-Scheepers, C., Clemson, A., … Sgrò,

C. M. (2019). Comparing thermal performance curves across traits: How consistent are they?

Journal of Experimental Biology, *222*(11). https://doi.org/10.1242/jeb.193433

Kordas, R. L., Harley, C. D. G., & O'Connor, M. I. (2011). Community ecology in a warming world: The

influence of temperature on interspecific interactions in marine systems. *Journal of Experimental*

Marine Biology and Ecology, *400*(1–2), 218–226. https://doi.org/10.1016/j.jembe.2011.02.029

Kozłowski, J. (1992). Optimal allocation of resources to growth and reproduction: Implications for age

and size at maturity. *Trends in Ecology and Evolution*, *7*(1), 15–19. https://doi.org/10.1016/0169-

5347(92)90192-E

Lutterschmidt, W. I., & Hutchison, V. H. (1997). The critical thermal maximum: history and critique.

Canadian Journal of Zoology, *75*(10), 1561–1574. https://doi.org/10.1139/z97-783

Martin, T. L., & Huey, R. B. (2008). Why "suboptimal" is optimal: Jensen's Inequality and ectotherm

thermal preferences. *American Naturalist*, *171*(3), 102–118. https://doi.org/10.1086/527502

Mohseni, O., & Stefan, H. G. (1999). Stream temperature/air temperature relationship: A physical

interpretation. *Journal of Hydrology*, *218*(3–4), 128–141. https://doi.org/10.1016/S0022-

1694(99)00034-7

Muñoz, M. M., & Bodensteiner, B. L. (2019). Janzen's hypothesis meets the Bogert Effect: Connecting

climate variation, thermoregulatory behavior, and rates of physiological evolution. *Integrative*

Organismal Biology, *1*(1), 1–12. https://doi.org/10.1093/iob/oby002

Niedrist, G. H., & Füreder, L. (2020). Real‐time warming of alpine streams: (re)defining invertebrates'

temperature preferences. *River Research and Applications*, https://doi.org/10.1002/rra.3638

Nogués-Bravo, D., Araújo, M. B., Errea, M. P., & Martínez-Rica, J. P. (2007). Exposure of global

mountain systems to climate warming during the 21st Century. *Global Environmental Change*,

17(3–4), 420–428. https://doi.org/10.1016/j.gloenvcha.2006.11.007

Packard, G. C., & Boardman, T. J. (1999). The use of percentages and size-specific indices to normalize

physiological data for variation in body size: wasted time, wasted effort? *Comparative Biochemistry*

and Physiology Part A: Molecular & Integrative Physiology, *122*(1), 37–44.

https://doi.org/10.1016/S1095-6433(98)10170-8

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across

natural systems. *Nature*, *421*(6918), 37. https://doi.org/10.1038/nature01286

Payne, N. L., & Smith, J. A. (2017). An alternative explanation for global trends in thermal tolerance. *Ecology Letters*, *20*(1), 70–77. https://doi.org/10.1111/ele.12707

Peckarsky, B. L. (1980). Predator-prey interactions between stoneflies and mayflies: Behavioral observations. *Ecology*, *61*(4), 932–943. https://doi.org/10.2307/1936762

Pilgrim, J. M., Fang, X., & Stefan, H. G. (1998). Stream temperature correlations with air temperature in Minnesota: Implications for climate warming. *Journal of the American Water Resources Association*, *34*(5), 1109–1121. https://doi.org/10.1111/j.1752-1688.1998.tb04158.x

Pincebourde, S., & Casas, J. (2019). Narrow safety margin in the phyllosphere during thermal extremes. *Proceedings of the National Academy of Sciences*, *116*(12), 5588–5596. https://doi.org/10.1073/pnas.1815828116

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, *569*(7754), 108. https://doi.org/10.1038/s41586-019-1132-4

Piyaphongkul, J., Pritchard, J., & Bale, J. (2012). Can tropical insects stand the heat? A case study with the brown planthopper Nilaparvata lugens (Stål). *PLoS One*, *7*(1), e29409. https://doi.org/10.1371/journal.pone.0029409

Polato, N. R., Gill, B. A., Shah, A. A., Gray, M. M., Casner, K. L., Barthelet, A., … Zamudio, K. R. (2018). Narrow thermal tolerance and low dispersal drive higher speciation in tropical mountains. *Proceedings of the National Academy of Sciences*, *115*(49), 12471–12476. https://doi.org/10.1073/pnas.1809326115

Pörtner, H. O. (2001). Climate change and temperature-dependent biogeography: Oxygen limitation of

thermal tolerance in animals. *Naturwissenschaften*, *88*(4), 137–146.

https://doi.org/10.1007/s001140100216

Pörtner, H. O. (2002). Climate variations and the physiological basis of temperature dependent

biogeography: systemic to molecular hierarchy of thermal tolerance in animals. *Comparative*

Biochemistry and Physiology Part A, *132*, 739–761. https://doi.org/10.1016/S1095-6433(02)00045-

4

Pörtner, H. O., Bennett, A. F., Bozinovic, F., Clarke, A., Lardies, M. A., Lucassen, M., … Stillman, J. H.

(2006). Trade-offs in thermal adaptation: the need for a molecular to ecological integration.

Physiological and Biochemical Zoology, *79*(2), 295–313. https://doi.org/10.1086/499986

Pörtner, H. O., Bock, C., & Mark, F. C. (2017). Oxygen- & capacity-limited thermal tolerance: Bridging

ecology & physiology. *Journal of Experimental Biology*, *220*(15), 2685–2696. https://doi.org/

10.1242/jeb.134585

Po□rtner, H. O., & Knust, R. (2007). Climate change affects marine fishes through the oxygen limitation

of thermal tolerance. *Science*, *315*(5808), 95–97. https://doi.org/ 10.1126/science.1135471

Pyne, M. I., & Poff, N. L. (2017). Vulnerability of stream community composition and function to

projected thermal warming and hydrologic change across ecoregions in the western United States.

Global Change Biology, *23*(1), 77–93. https://doi.org/10.1111/gcb.13437

Rangecroft, S., Suggitt, A. J., Anderson, K., & Harrison, S. (2016). Future climate warming and changes

to mountain permafrost in the Bolivian Andes. *Climatic Change*, *137*, 231–243.

https://doi.org/10.1007/s10584-016-1655-8

Rezende, E. L., Tejedo, M., & Santos, M. (2011). Estimating the adaptive potential of critical thermal

limits: methodological problems and evolutionary implications. *Functional Ecology*, *25*(1), 111–

121. https://doi.org/10.1111/j.1365-2435.2010.01778.x

Ribeiro, P. L., Camacho, A., & Navas, C. A. (2012). Considerations for assessing maximum critical

temperatures in small ectothermic animals: Insights from leaf-cutting ants. *PLoS ONE*, *7*(2), e32083.

https://doi.org/10.1371/journal.pone.0032083

Rolfe, D. F. S., & Brown, G. C. (1997). Cellular energy utilization and molecular origin of standard

metabolic rate in mammals. *Physiological Reviews*, *77*(3), 731–758.

https://doi.org/10.1152/physrev.1997.77.3.731

Rotvit, L., & Jacobsen, D. (2013). Temperature increase and respiratory performance of

macroinvertebrates with different tolerances to organic pollution. *Limnologica*, *43*(6), 510–515.

https://doi.org/10.1016/j.limno.2013.04.003

Schulte, P. M. (2015). The effects of temperature on aerobic metabolism: towards a mechanistic

understanding of the responses of ectotherms to a changing environment. *Journal of Experimental*

Biology, *218*(12), 1856–1866. https://doi.org/10.1242/jeb.118851

Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal performance curves, phenotypic

plasticity, and the time scales of temperature exposure. *Integrative and Comparative Biology*, *51*(5),

691–702. https://doi.org/10.1093/icb/icr097

Shah, A. A., Bacmeister, E. M. S., Rubalcaba, J. G., & Ghalambor, C. K. (2020a). Divergence and

constraint in the thermal sensitivity of aquatic insect swimming performance. *Current Zoology*,

https://doi.org/10.1093/cz/zoaa040.

Shah, A. A., Dillon, M. E., Hotaling, S., & Woods, H. A. (2020b). High elevation insect communities

face shifting ecological and evolutionary landscapes. *Current Opinion in Insect Science*, 41, 1–6.

https://doi.org/10.1016/j.cois.2020.04.002

Shah, A. A., Funk, W. C., & Ghalambor, C. K. (2017a). Thermal acclimation ability varies in temperate

and tropical aquatic insects from different elevations. *Integrative and Comparative Biology*, *57*(5),

977–987. https://doi.org/10.1093/icb/icx101

Shah, A. A., Gill, B. A., Encalada, A. C., Flecker, A. S., Funk, W. C., Guayasamin, J. M., … Ghalambor, C. K. (2017b). Climate variability predicts thermal limits of aquatic insects across elevation and latitude. *Functional Ecology*, *31*(11), 2118–2127. https://doi.org/10.1111/1365-2435.12906

Sheldon, A. L. (2012). Possible climate-induced shift of stoneflies in a southern Appalachian catchment. *Freshwater Science*, *31*(3), 765–774. https://doi.org/10.1899/11-135.1

Sjöstrom, P. (1985). Hunting behaviour of the perlid stonefly nymph Dinocras cephalotes (Plecoptera) under different light conditions. *Animal Behavior*, *33*, 534–540. https://doi.org/10.1016/S0003-3472(85)80076-2

Stewart, K. W., Baumann, R. W., & Stark, B. P. (1974). The distribution and past dispersal of southwestern United States Plecoptera. *Transactions of the American Entomological Society*, *99*(4), 507–546.

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1713), 1823–1830. https://doi.org/10.1098/rspb.2010.1295

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global redistribution of animals. *Nature Climate Change*, *2*(9), 686–690. https://doi.org/10.1038/nclimate1539

Tamaris-Turizo, C. E., Turizo-Correa, R. R., & del Carmen Zúñiga, M. (2007). Spatial-temporal distribution and feeding in nymphs of Anacroneuria (Insecta: Plecoptera: Perlidae) in the Gaira River (Sierra Nevada de Santa Marta, Colombia). *Caldasia*, 375–385.

Terblanche, J. S., & Chown, S. L. (2007). The effects of temperature, body mass and feeding on metabolic rate in the tsetse fly Glossina morsitans centralis. *Physiological Entomology*, *32*(2), 175–

180. https://doi.org/10.1111/j.1365-3032.2006.00549.x

Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C., & Chown, S. L. (2007). Critical thermal

limits depend on methodological context. *Proceedings of the Royal Society B: Biological Sciences*,

274(1628), 2935–2943. https://doi.org/10.1098/rspb.2007.0985

Vannote, R. L., & Sweeney, B. W. (1980). Geographic analysis of thermal equilibria: A conceptual model

for evaluating the effect of natural and modified thermal regimes on aquatic insect communities. *The*

American Naturalist, *115*(5), 667–695.

Verberk, W. C. E. P., Bilton, D. T., Calosi, P., & Spicer, J. I. (2011). Oxygen supply in aquatic

ectotherms: Partial pressure and solubility together explain biodiversity and size patterns. *Ecology*,

92(8), 1565–1572. https://doi.org/10.1890/10-2369.1

Verberk, W. C. E. P., Durance, I., Vaughan, I. P., & Ormerod, S. J. (2016a). Field and laboratory studies

reveal interacting effects of stream oxygenation and warming on aquatic ectotherms. *Global Change*

Biology, *22*(5), 1769–1778. https://doi.org/10.1111/gcb.13240

Verberk, W. C. E. P., Overgaard, J., Ern, R., Bayley, M., Wang, T., Boardman, L., & Terblanche, J. S.

(2016b). Does oxygen limit thermal tolerance in arthropods? A critical review of current evidence.

Comparative Biochemistry and Physiology -Part A : Molecular and Integrative Physiology, *192*,

64–78. https://doi.org/10.1016/j.cbpa.2015.10.020

Vucic-Pestic, O., Ehnes, R. B., Rall, B. C., & Brose, U. (2011). Warming up the system: higher predator

feeding rates byt lwer energetic differences. *Global Change Biology*, *17*(3), 1301–1310.

https://doi.org/10.1111/j.1365-2486.2010.02329.x

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., … Bairlein, F. (2002).

Ecological responses to recent climate change. *Nature*, *416*(6879), 389.

https://doi.org/10.1038/416389a

Weissenberger, J., Spatz, H.-C., Emanns, A., & Schwoerbel, J. (1991). Measurement of lift and drag

forces in the m N range experienced by benthic arthropods at flow velocities below 1.2 m s^{-1}.

Freshwater Biology, *25*(1), 21–31. https://doi.org/10.1111/j.1365-2427.1991.tb00469.x

Wieser, W. (1994). Cost of growth in cells and organisms: General rules and comparative aspects.

Biological Reviews of the Cambridge Philosophical Society, *69*(1), 1–33. 10.1111/j.1469-

185x.1994.tb01484.x