Temperature-dependence of metabolic rate in tropical and temperate aquatic insects:

support for the Climate Variability Hypothesis in mayflies but not stoneflies.

Alisha A. Shah[1,2], H. Arthur Woods[2], Justin C. Havird[3], Andrea C. Encalada[4], Alexander S. Flecker[5], W. Chris Funk[1,6], Juan M. Guayasamin[4], Boris C. Kondratieff[1,6,7], N. LeRoy Poff[1,6,8], Steven A. Thomas[9], Kelly R. Zamudio[5], Cameron K. Ghalambor[1,6]

Affiliations:

[1] Department of Biology, Colorado State University, Fort Collins, CO, USA

[2] Division of Biological Sciences, University of Montana, Missoula, MT, USA

[3] Department of Integrative Biology, University of Texas, Austin, TX, USA

[4] Colegio de Ciencias Biológicas y Ambientales COCIBA, Instituto BÍOSFERA-USFQ, Campus Cumbayá, Universidad San Francisco de Quito USFQ, Quito, Ecuador

[5] Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY, USA

[6] Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO, USA

[7] Department of Bioagricultural Sciences and Pest Management, Colorado State University, Fort Collins, CO, USA

[8] Institute for Applied Ecology, University of Canberra, ACT, Australia

[9] School of Natural Resources, University of Nebraska, Lincoln, NE, USA

Corresponding Author:

Alisha A. Shah, Division of Biological Sciences, University of Montana, Missoula, MT, USA;

Email: alisha.shah@mso.umt.edu; Phone: (512) 694-7532

Running Head:

Climate vulnerability of aquatic insects

Keywords:

Metabolic rate, thermal tolerance, latitude, elevation, mayflies, stoneflies, vulnerability to

climate change, thermal performance curve

Abstract:

A fundamental gap in climate change vulnerability research is an understanding of the relative

thermal sensitivity of tropical and temperate organisms. Aquatic insects are vital to stream

ecosystem function and biodiversity. With global temperatures rising at an unprecedented rate, it

is imperative that we understand how sensitive aquatic insects are to increasing temperature and

whether sensitivities vary among taxa across latitude and elevation. We measured thermal

sensitivity of standard metabolic rate in stream-dwelling mayflies and stoneflies across a ~2,000

m elevation gradient in the temperate Rocky Mountains in Colorado, U.S.A., and the tropical

Andes in Napo, Ecuador. Mayflies generally adhered to the predictions of the Climate

Variability Hypothesis, such that tropical species exhibited greater thermal sensitivity than did

temperate species. For stoneflies, by contrast, there was no evidence in support of our

predictions; temperate stoneflies showed greater thermal sensitivity than tropical relatives.

Differences among mayflies and stoneflies may reflect the evolutionary history of these insect

clades or different ecological roles of mayflies (often herbivorous) and stoneflies (often

predatory). Collectively our results suggest that there is physiological variation across elevations

and species and low elevation tropical mayflies may be the most imperiled by climate warming.

However, broad generalizations about the vulnerability of tropical ectotherms should be made

with caution.

Introduction:

Metabolism is crucial to all life. It integrates an organism's existing energy supplies and

materials to fuel fitness-related activities like movement, growth, tissue maintenance, and

development (Dillon, Wang, & Huey, 2010; Gillooly, Brown, West, Savage, & Charnov, 2001).

In ectotherms, metabolic rates are sensitive to temperature, which can influence not only

individual fitness, but distribution and abundance as well (Angilletta, 2009; Terblanche &

Chown, 2007; Vannote & Sweeney, 1980). A major challenge in global change biology is to

understand how temperature alters metabolism and energy budgets of different species, and in

turn, impacts the ability of species to cope with climate change (Colwell, Brehm, Cardelús,

Gilman, & Longino, 2008; Parmesan, Yohe, & Andrus, 2003; Walther et al., 2002). So far, most

attempts to measure vulnerability to climate change have assessed species response to

temperature challenges, such as lethal and critical thermal limits (Deutsch et al., 2008; Pinsky,

Eikeset, McCauley, Payne, & Sunday, 2019). Despite their utility, these metrics address only the

responses of organisms to extreme temperatures and do not capture the continuous or

incremental effects of temperature on an organism's energy budget. Such effects are particularly

important in ectotherms, because body temperature and metabolic rate increase with

environmental temperature, resulting in higher energy expenditure. This response is likely to

trade-off against other competing demands such as growth, reproduction, and locomotor

performance (Pörtner et al., 2006). For example, in aquatic ectotherms, increased metabolic

demand can outpace supply of oxygen from the environment and result in decreased

performance and lowered tolerance to heat stress (Pörtner, Bock, & Mark, 2017). Thus,

understanding metabolic responses to temperature in species occupying different thermal

environments can help us predict geographic variation in response to global warming (Dillon et

al., 2010). In this study, we measured metabolic rate profiles in juveniles (nymphs) of two major

groups of stream insects – mayflies and stoneflies – in temperate and tropical streams across a

paired elevation gradient. At this aquatic stage, nymphs spend much of their time feeding,

growing, and developing eggs and sperm packets (Brittain, 1990). Water temperature is therefore

a major driver of life history traits in these taxa. We aimed to elucidate differences in patterns of

thermal sensitivity and thus better evaluate potential vulnerabilities of mayflies and stoneflies to

projected climate warming for which, currently, very little data exist (Chown, Duffy, &

Sørensen, 2015).

82   Much like other types of thermal performance curves (TPCs), oxygen-consumption

profiles (used to measure metabolic rate) vary with increasing temperature and display three

identifiable regions (Pörtner, 2002; Schulte, 2015): an ascending phase; a peak, which indicates

the onset of stress (Huey & Stevenson, 1979; Pörtner & Knust, 2007); and a descending phase of

metabolic depression (Guppy & Withers, 1999). When coupled with other measures of

performance and the critical limits, changes in oxygen consumption provide a more complete

picture of how temperature shapes the ecology of an organism (Angilletta, 2009; Pörtner et al.,

2017) (Fig. 1).

90   Although TPCs for traits like locomotion and metabolic rate have similar shapes, a

metabolic rate curve is a special type of TPC (Pörtner, 2002; Schulte, Healy, & Fangue, 2011),

with different implications for organismal fitness (Kellermann et al., 2019). The performance

traits used in many TPCs—running speed, feeding rates, growth, or reproduction—are often

interpreted as components of fitness, such that *more* is *better*. For these traits, peaks indicate the

location of thermal optima (T_{OPT}, Fig. 1) and suggest that performance is optimized at that

temperature (Angilletta, Niewiarowski, & Navas, 2002; Huey & Kingsolver, 1989; Huey &

Stevenson, 1979) or something close to it (Martin & Huey, 2008). For metabolism, in contrast,

more is not necessarily better, and peak standard rates of oxygen consumption typically occur at

warmer temperatures than do peak performance traits (Pörtner, 2001). For example, aerobic

scope is maximized when the difference between standard metabolic rate (SMR) and maximum

metabolic rate (MMR) is greatest (Fig. 1A; often the location of peak performance T_{OPT})—not

when SMR itself is at its peak. At higher temperatures, when SMR meets MMR, a metabolic

depression often follows as organismal functions decline (Fig. 1; see also Pörtner, 2002). Thus,

the metabolic peak should lie to the right of T_{OPT} (i.e. at higher temperatures than T_{OPT}) and align

with decreasing performance on the TPC (Fig. 1; Pörtner, 2001). We refer to this peak as T-

MR_{PEAK}.

The shapes of metabolic rate curves found for different species, populations, and thermal

experience (e.g., acclimitization; DeLong et al., 2018) could either reflect the acute effects of

temperature on biological rates (Dillon et al., 2010; Payne & Smith, 2017) or the evolutionary

response to past climatic conditions (Dell, Pawar, & Savage, 2011; Deutsch et al., 2008). If the

shape of the metabolic rate TPCs simply reflects thermodynamics of biological reactions, we

would predict that organisms of equivalent size experiencing a similar range of temperatures

should exhibit similar metabolic rate curves (Payne & Smith, 2017).

Alternatively, the Climate Variability Hypothesis posits that if stable climatic regimes,

such as those found in the tropics, expose organisms to only a narrow range of temperatures and

favor the evolution of thermal specialists (*sensu* Huey & Kingsolver, 1989), we predict that those

organisms should have lower SMR within the range of temperatures they experience to

maximize the allocation of energy to growth, reproduction, and aerobic performance (Fig.1B).

These thermal specialists should also be more sensitive to temperature, i.e. have metabolic rates

that change rapidly in response to temperature outside their normal range. Climates characterized

by large seasonal temperature variation, such as those found at temperate latitudes, should select

for thermal generalists capable of functioning effectively over a wide range of temperatures

(Angilletta et al., 2002; Huey & Kingsolver, 1993). Thermal generalists are predicted to have

relatively low SMRs across a wide range of temperatures (Fig. 1C) and be less sensitive to

increasing temperature to keep their minimum energy demands low. The generality of these

contrasting hypotheses remains open to debate because few comparative studies have measured

metabolic rates in species from temperate and tropical regions under similar experimental

conditions.

To test predictions generated by the alternative hypotheses detailed above, we examined

metabolic rates of mayflies and stoneflies from temperate and tropical regions across different

elevations. Because stream temperatures at a given elevation were similar at the time of the

experiments, we predicted that the metabolic rate curves should look similar between temperate

and tropical species if they were strictly determined by physicochemical effects of temperature

on metabolic processes (Payne & Smith, 2017). But, if metabolic rates reflect adaptation to

climate, we predicted greater sensitivity to temperature in the tropical species (i.e. larger Q_{10}

values; Angilletta et al., 2002; Huey & Kingsolver, 1993); (Fig. 1B) compared to temperate ones

(Fig. 1C). We also expected to find differences in SMR *within* each latitude, across elevations,

such that insects experiencing greater thermal variation (e.g., in low elevation temperate streams)

would be less sensitive to temperature than insects experiencing reduced thermal variation (e.g.,

high elevation temperate streams).

Figure 1. Expectations for change in standard metabolic rate (oxygen consumption) and
performance in response to temperature in an aquatic ectotherm. The thermal performance curve
(TPC, solid green line, left panel) could refer to any functional performance trait (e.g. swimming
performance), where higher performance is better. T_{OPT}, where performance is highest, reflects
the animal's preferred temperature. The oxygen consumption curve (dashed line) is a special
case of TPC. The peak in this case (T-MR$_{PEAK}$) could represent the temperature at which
standard metabolic rate (SMR) is so high that essentially equals the maximum metabolic rate
(MMR) that the ectotherm can achieve (circle, right panel **A**). T-MR$_{PEAK}$ may therefore reflect
stress (red color) and aligns with a decline in the functional performance trait. The general
location of behaviorally determined critical thermal limits (CT$_{MIN}$ and CT$_{MAX}$) are also shown
(blue line, left panel). In this study we focus only on SMR. We predict that SMR will rise steeply
in tropical species (right panel **B**) and be stressful when temperatures exceed the typical narrow
range of temperatures they experience (gray rectangle). In comparison, temperate species should
exhibit a gradual rise in SMR across the same temperatures (right panel **C**), reflecting the broad
range of environmental temperatures they experience (grey rectangle).

Methods

Site description

Temperate sites were located in the Cache La Poudre river drainage of the southern

Rocky Mountains in Colorado, USA, and tropical sites in the Papallacta-Quijos river drainage of

the Andes Mountains in Napo Province, Ecuador. We chose small, relatively pristine streams

where focal mayflies and stoneflies are typically found. Sites at both latitudes were matched by

elevation, where high >3000 m, mid = 2000 – 3000 m, and low = 1000 – 2000 m. Each elevation

pair was approximately the same temperature at the time of collection (Table 1). Thus, we were

able to make latitudinal comparisons of insect metabolic rate for each elevation pair under the

same thermal conditions. Ephemeroptera (mayflies) and Plecoptera (stoneflies) do not share

genera between tropical and temperate sites. Therefore, we selected representative genera from

the same family (mayfly family Baetidae and stonefly family Perlidae) at each location to

contrast their metabolic rate. In the Rockies, we collected a total of 313 individual mayflies

(genus *Baetis*) and 107 stoneflies (genus *Hesperoperla*). In the Andes, we collected 245 mayflies

(genus *Andesiops*) and 76 stoneflies (genus *Anacroneuria*). All insects were identified to genus

by morphology in the field. Although mayflies were easily collected across all elevations,

stoneflies were much harder to find and were altogether absent at high elevation, so comparisons

between stoneflies are at mid and low elevations only. We conducted experiments between

January and April of 2013 in the Andes and June to August of 2013 in the Rockies.

181   **Table 1**. Average stream temperatures with standard errors and acclimation temperatures for
182   each temperate (Rocky Mountain) and tropical (Andes) stream from which insects were
183   collected.
184

Latitude	Site ID	Type	Elevation (m)	Stream temperature (°C)	Acclimation temperature (°C)
Temperate	COP2212	Low	2212	12.5 ± 3.73	14
	COP2590	Mid	2590	10.8 ± 4.68	10
	COP2798	Mid	2798	7.6 ± 1.41	10
	COP3166	High	3166	6.2 ± 3.56	6
Tropical	ECP1845	Low	1845	13.5 ± 0.72	14
	ECP2003	Low	2003	14.0 ± 0.38	14
	ECP2694	Mid	2798	10.0 ± 0.79	10
	ECP3683	High	3683	8.6 ± 0.89	6
	ECP3898	High	3898	6.6 ± 1.74	6

Lab Acclimation

Field-caught nymphs were brought back to the lab (~1600 m at both latitudes) and placed in containers of filtered stream water from their native streams without food for 48 h. During this time, high oxygen levels were maintained by bubbling air into the bath, and water was circulated by small pumps, with temperatures approximating the insects' home stream temperatures (Table 1). Temperate and tropical high elevation insects were acclimated to 6 °C, mid-elevation insects to 10 °C, and low elevation insects to 14 °C. We selected this paired design because tropical insects failed to acclimate to temperatures outside typical temperatures of their native streams (see also Shah, Funk, & Ghalambor, 2017a). For example, we often detected evidence of stress (e.g., jerking movements, unnatural swimming patterns) and even mortality, when tropical mayflies from high and low elevations were warmed outside native stream temperature (i.e., 12 °C and 10 °C, respectively). Therefore, the use of a single acclimation temperature across all elevations was not possible. Because acclimation temperatures differed among elevations but

were the same for each elevation pair, we restricted our comparisons to the same elevations

between the two latitudes to control for the background acclimation temperature.

Metabolic Rate

Oxygen consumption rates (metabolic rates, hereafter) were measured using closed glass

micro-respirometry chambers (Unisense A/S, Aarhus, Denmark). Mayflies were placed in 4 mL

chambers, and stoneflies, which are larger, were placed in 50 mL chambers. After the 48 h

acclimation period, we introduced insects randomly to one of nine test temperatures (5, 7.5, 10,

12.5, 15, 17.5, 20, 22.5, or 25 °C) by allowing temperature to rise naturally or by slowly cooling

(0.5 °C-min) to prevent thermal shock. On average, 16 individuals per population for mayflies and

6 per population for stoneflies were tested at each temperature (Tables S1, S2). Once at the test

temperature, insects were moved individually into glass chambers filled with stream water that

had been treated with UV light to reduce background microbial respiration, and immersed in a

water bath held at test temperature (Fig. S1). Chambers were fitted with a glass-coated magnetic

stir bar and a plastic mesh to which the insect could cling (Fig. S1). We allowed insects to

acclimate to the chamber space for an additional hour before sealing each chamber with a glass

stopper to prevent further introduction of atmospheric oxygen. Each stopper had a central

capillary hole, allowing us to insert an oxygen microelectrode (500 μm diam.) connected to a

picoammeter (Unisense OXY-Meter, Aarhus, Denmark) to measure change in oxygen

concentration. To measure background respiration rates, a control chamber (without an insect)

was included in each experiment. The microelectrode was first lowered into the control chamber

at the start of the experiment. After reading oxygen concentration continuously for 2 minutes, it

was moved to the first chamber that contained an animal, where data were recorded for another 2

minutes (Fig. S1), and so on. Experiments were terminated when oxygen levels reached 80% of

the air-saturated values because natural dissolved oxygen levels in our study sites rarely fall

below this value. Consequently, we made 3-6 readings for each chamber – fewer readings at the

higher temperatures compared to the lower temperatures – which lasted around 128 minutes.

Because there were typically 16 chambers running simultaneously, the average time interval

between readings was 25 mins. While recording oxygen concentration, we also observed insects

inside the chambers, noting any stress responses or mortality. Insects were then removed from

chambers, euthanized in ethanol, dried, and weighed.

Statistical Analyses

We first calculated average rate of oxygen consumption for each insect or control (blank)

chamber by measuring change in oxygen concentration across all measurements taken in a given

chamber (see Supplemental Materials and Fig. S2 for an explanation of this calculation). We

then averaged the control values for each temperature such that we calculated a control rate for

each combination of test temperature, elevation, and latitude. Control rates were subtracted from

insect consumption rates to correct for background microbial respiration. We conducted all

statistical analyses on these 'control-corrected oxygen consumption rates' (hereafter, oxygen

consumption) in R version 3.4.0 (R Core Team 2017).

240   Because we were interested in the thermal sensitivity of temperate and tropical metabolic

rates between elevation pairs (i.e. the thermal reaction norm), we first compared metabolic rates

during the ascending phase of the TPC by fitting a linear model in R with oxygen consumption

as the response variable for each elevation type (i.e. high, mid, and low) separately. For this

analysis, we used only the temperatures 5, 7.5, 10, 12.5, 15 and 17.5 °C because these

temperatures were within the ascending phase of the SMR curves for most populations in this

study. The model included fixed effects of test temperature and latitude, and their interaction,

with body mass as a covariate (Packard & Boardman, 1999). A significant interaction would

denote a difference in thermal sensitivity (i.e. slope) between temperate and tropical species.

249   We conducted a similar analysis for the descending phase using data from the 17.5 and

250   20 °C treatments as these two temperatures encompassed the descending phase of the curve. At

251   the highest temperatures (22.5 and 25 °C), we observed stress responses (e.g. unnatural

252   swimming patterns, frenzied attempts to escape from the chamber and in some cases elevated

253   metabolic rates) in tropical mayflies. We therefore did not include SMR at 22.5 and 25 °C in this

254   analysis but instead calculated separate Q_{10} values for these temperatures (see below).

255   To compare the maximum SMR at each elevation between temperate and tropical taxa,

we examined latitude x elevation combination individually to find the physiologically relevant

peak (i.e., before CT_{MAX} and when the first signs of stress were apparent, if observed). This peak

was typically at 17.5 °C in our study because in most cases, across both latitudes, SMR was

highest at this temperature even if not always statistically different from 15 °C. We then

compared these values using an ANOVA for each elevation pair using latitude as the fixed effect

and body mass as a covariate.

262   Finally, we calculated Q_{10}, a measure of sensitivity to temperature, using the formula

263
$$Q_{10} = \left(\frac{R_2}{R_1}\right)^{\left(\frac{10}{T_2 - T_1}\right)}$$

where R_1 and R_2 are reaction (oxygen consumption) rates measured at two different

temperatures, T_1 and T_2, respectively. Q_{10}s were calculated across cooler 'non-stressful'

temperatures ($T_1 = 7.5$ and $T_2 = 15$ °C) and warmer 'stressful' temperatures ($T_1 = 15$ and $T_2 = 25$

°C).

Results

Behavioral observations

Tropical mayflies showed visible signs of stress (jerking movements, loss of righting

response) at several test temperatures outside the range they typically experience (e.g., 22.5 and

25 °C; Fig. 2 red bars, Tables S1; Fig. S3 A); many of the stressed individuals died at the end of

the experiment (Table S1; Fig. S3 B). Low elevation tropical mayflies also experienced some

mortality at 5 °C, which is much colder than the minimum temperatures they encounter in their

home streams (Fig. 2F). Temperate mayflies, in contrast, experienced little-to-no visible signs of

stress at any experimental temperature (Table S2). We also did not detect any overt signs of

stress in any of the stoneflies in this study.

Thermal sensitivity based on standard metabolic rates and reaction norms

SMR profiles differed between temperate and tropical mayflies and stoneflies at all

elevations (Figs. 2 and 3). In general, metabolic rates of mayflies from the two latitudes were

lowest at temperatures approximating those in their native streams. Outside these temperatures,

however, tropical mayflies exhibited a greater increase in metabolic rate (Fig. 2). In all mayfly

populations across both latitudes, a secondary peak was observed at 22.5 and 25 °C that

corresponded with visible stress during experiments. We denoted the initial peak in SMR as T-

MR PEAK (shown as yellow arrows on Figs. 2 and 3) or the onset of stress, and the secondary

peak as an acute stress response because it was followed by death in some cases. As indicated in

the Methods, we excluded these stressful test temperatures from the analyses of the ascending

and descending phases of metabolic rate profiles and dealt with them separately in an analysis of

Q_{10}.

Figure 2. Plots of standard metabolic rate least square means in mayflies from high (A, B), mid
(C,D), and low (E,F) elevations in temperate (left panel) and tropical (right panel) streams. Red
bars indicate temperatures at which we observed stress. T-MR$_{PEAK}$ is indicated by yellow arrows
for each population. Gray rectangles represent the approximate stream temperature range
measured in each stream. CT$_{MIN}$ and CT$_{MAX}$ determined in a previous study are shown.

Figure 3. Plots of metabolic rate least square means in predatory stoneflies from mid (A, B) and low (C,D) elevations in temperate (left panel) and tropical (right panel) streams. Stoneflies were not found at high elevation. T-MR$_{PEAK}$ is indicated by yellow arrows for each population. Gray rectangles represent the approximate stream temperature range measured in each stream. We did not detect overt stress in stoneflies in this study. CT$_{MIN}$ and CT$_{MAX}$ determined in a previous study are shown.

In the ascending phase, metabolic rates in tropical mayflies across all elevations were more sensitive to temperature increases than were their temperate counterparts ($F_{(5,371)} = 8.57$, $p < 0.001$). Focusing on each elevation separately, we found tropical species to have greater increases in metabolic rate at high elevation ($F_{(5,97)} = 3.27$, $p = 0.009$, Fig. 4A), mid elevation ($F_{(4,98)} = 4.17$, $p = 0.003$, Fig. 4B), and at low elevation ($F_{(4,152)} = 6.77$, $p < 0.001$, Fig. 4C). In stoneflies, ascending phases did not differ between temperate and tropical populations (Figs. 5 A and B)

Temperate mayflies had a steeper decrease in metabolic rate at high elevation (Fig. 4A; $F_{(1,1)} = 4.04$, $p = 0.05$) and at low elevation (Fig. 4C; $F_{(1,1)} = 6.86$, $p = 0.01$). At mid elevation, there was no significant interaction (Fig. 4B). However, a Student's T-test revealed that oxygen

consumption values were significantly higher in tropical mayflies compared to temperate ones at

17.5°C ($t = 3.946$, $p = 0.001$), but rates were equally low between latitudes at 20°C ($t = -1.006$, p

$= 0.33$). This suggests that the descending phase was steeper for tropical mayflies compared to

temperate ones. Descending phases of stoneflies did not differ between tropical and temperate

populations.

Figure 4. A comparison of the ascending (left panel) and descending (right panel) phases of
metabolic profiles at high (A), mid (B) and low (C) elevations. Values are the least square means
of metabolic rate in mayflies from tropical (white circles) and temperate (black circles) streams,
after accounting for body mass in the model. Vertical bars are standard errors and p values are
shown.

Figure 5. A comparison of the ascending (left panel) and descending (right panel) phases of metabolic profiles at mid (A), and low (B) elevations. Values are the least square means of metabolic rate in stoneflies from tropical (white circles) and temperate (black circles) streams, after accounting for body mass in the model. Vertical bars are standard errors and p values are shown.

Finally, the maximum SMR was higher in tropical compared to temperate mayflies at all

elevations (Fig. 6). Stoneflies showed an opposite pattern where tropical species had lower peaks

at both measured elevations, although this was significant only at mid elevation (Fig. 6).

Figure 6. Peak SMRs (i.e. the highest metabolic rate) are compared between temperate and
tropical mayflies from high (A), mid (B), and low (C) elevations and stoneflies from mid (D) and
low (E) elevations. Vertical bars are standard errors and p values are shown.

Thermal Sensitivity based on Q_{10}

Our Q_{10} calculations revealed contrasting results for stoneflies, but not mayflies. When

comparing Q_{10}s across 'stressful' and 'non-stressful' temperatures, we found Q_{10}s for both taxa

to be higher at non-stressful temperatures than at stressful temperatures (Table 2). Across

latitude, tropical mayflies showed greater sensitivity to temperature (higher Q_{10} values) at both

non-stressful and stressful temperatures than their temperate counterparts (Table 2; Fig. 7).

Notably, at mid and low elevations, tropical mayflies were highly sensitive to temperature even

at non-stressful temperatures (Q_{10} = 8.40 and 5.17, respectively). In contrast, tropical stoneflies

were less sensitive than their temperate relatives for stressful and non-stressful temperatures

(Fig. 7).

Table 2. A comparison of Q_{10} values calculated for temperate and tropical mayfly metabolic
rates between typically non-stressful temperatures (7.5°C to 15°C) and typically stressful
temperatures (15°C to 25°C).

		Mayflies		*Stoneflies*	
	Elevation	**Temperate**	**Tropical**	**Temperate**	**Tropical**
Non-stressful	High	3.63	3.15	n/a	n/a
	Mid	3.15	5.17	6.76	2.49
	Low	2.36	8.40	6.66	5.78
Stressful	High	1.07	1.46	n/a	n/a
	Mid	0.73	1.59	6.10	0.87
	Low	1.34	1.67	1.25	0.42

361
362

Figure 7. Comparison of average Q_{10} for mayflies and stoneflies during the ascending phase of
the metabolic rate curve.

Discussion

Comparative studies of critical thermal limits have generally found tropical ectotherms to

be adapted to narrower ranges of temperatures compared to their temperate counterparts,

suggesting that tropical species are more vulnerable to global warming (Deutsch et al., 2008;

Gutiérrez-Pesquera et al., 2016; Shah et al., 2017b; Sunday, Bates, & Dulvy, 2011, 2012). Payne

and Smith (2017) recently challenged such conclusions by arguing that narrower thermal breadth

may simply reflect the acute effects of temperature on biological rates of organisms living in

warmer environment. Here, using a paired-elevation design, we compared the thermal sensitivity

of SMR in closely related temperate and tropical aquatic insect nymphs. This study design

enabled a comparison of temperate and tropical species under a common range of environmental,

acclimation, and test temperatures. Despite originating in streams with similar environmental

temperatures at the time of collection, tropical and temperate insects had different metabolic rate

profiles (Figs. 2, 3). These results suggest that variation between temperate and tropical species

does not reflect simple, direct effects of temperature on biological rates, but instead support the

view that there are physiological differences between taxa from these regions. Tropical mayflies,

in particular, exhibited greater sensitivity to temperature in terms of metabolic rates and the onset

of stress compared to their temperate counterparts. By contrast, stoneflies showed little-to-no

divergence in their thermal sensitivity. The ecological implications of greater thermal sensitivity

remain unresolved for most ectotherms, but in the absence of compensatory changes in

productivity, elevated metabolic rates could negatively impact the energy budget and capacity

for aerobic performance in tropical mayflies (Dillon et al., 2010; Huey & Kingsolver, 2019).

Comparison of metabolic rate profiles

In most cases, metabolic rate generally rose with increasing temperature, peaked, and then fell.

This pattern, however, varied considerably among taxa, populations, and elevations. As a group,

tropical mayflies were more thermally sensitive than their temperate counterparts (Figs. 2, 4, 6

and 7). The metabolic rate curves of tropical mayflies had overall steeper ascending and

descending phases, and higher Q_{10}s, across all three elevations. The particularly high Q_{10} for mid

and low elevation tropical mayflies were similar to those of North Atlantic stenothermal fishes

when experiencing temperatures away from an optimum (Johnston, Clarke, & Ward, 1991).

These results support the prediction that tropical mayflies, which experience more stable stream

temperatures (Shah et al., 2017b), are more sensitive to temperature.

Given that mayflies were paired by elevations that had similar environmental and

acclimation temperatures, these differences cannot be attributed simply to the passive response

of metabolic rate to higher temperatures. Instead, the greater thermal sensitivity of metabolism in

tropical mayflies largely reflects a rapid increase and then decline outside the relatively narrow

range of environmental temperatures that they experience (Figs. 2, 4, 6, 7). In contrast, temperate

mayflies exhibited more modest increases in metabolism in response to temperature and had

similar metabolic rate profiles across elevations (Fig. 2). For both temperate and tropical

mayflies, metabolic rates tended to be lower within the range of environmental temperatures they

experience (Fig. 2), suggesting there may be selection to reduce the minimum energetic costs

associated with maintenance (Artacho & Nespolo, 2009; Bochdansky, Grønkjær, Herra, &

Leggett, 2005; Brown, Marquet, & Taper, 1993). For example, even a moderate increase in

stream temperature outside the normal range (Fig. 2, right edge of grey rectangles) is associated

with peaks in metabolism, which has potentially cascading effects on energy budgets and

possible chronic stress (Fig. 2). If selection acts to minimize SMR under typical stream

temperatures (Brown et al., 1993), these results would support the hypothesis that the narrower

and more stable stream temperatures experienced by tropical mayflies underlie their greater

thermal sensitivity. Such an interpretation is consistent with other aspects of temperate and

tropical mayfly thermal physiology. First, thermal breadth (difference between critical thermal

limits) measured in a parallel study were the narrowest for tropical mayfly populations (Shah et

al., 2017b). Second, in this study, we observed visible signs of stress in the metabolic rate

profiles of tropical but not temperate mayflies. Specifically, after the decline in metabolic rate,

secondary peaks occurred at 22.5 °C and 25 °C for most groups, which corresponded to visible

signs of stress and mortality in tropical but not temperate mayflies. Signs of stress at these high

temperatures included leg spasms, loss of righting, release from substrate, and mortality (see

Supplemental Information Table S1, Figs. S3 A, B). Other aquatic ectotherms, such as brown

bullhead catfish (Heath, 1973) and harvester ants (Lighton & Turner, 2004), also show similar

postmortem peaks in metabolism associated with heat stress. Collectively, these results are

consistent with previous interpretations that temperate mayflies are thermal generalists, with

broad thermal breadths and reduced metabolic sensitivity to temperature, while tropical mayflies

are thermal specialists, with a narrower thermal breadth and increased metabolic sensitivity to

temperature (c.f. Huey & Hertz, 1984).

The differences observed between temperate and tropical mayflies were generally

unambiguous and supported by prior work in this system. In stoneflies, evidence in support of

our predictions was less straightforward. However, the similarity between temperate and tropical

stoneflies is consistent with previous research on other physiological traits (Shah et al., 2017a,

2017b). While there are some differences in metabolic rate profiles between temperate and

tropical stoneflies (e.g. higher metabolic rates at warmer temperatures in temperate, but not

tropical species), these differences are far less striking than those observed in mayflies (Fig. 2 vs.

3). Overall, the ascending and descending phases did not differ between temperate and tropical

stoneflies (Figs. 3, 5), but absolute maximum metabolic rates and Q_{10}s across non-stressful and

stressful temperatures were higher in the temperate species (Fig. 7). This lack of latitudinal

differences in stoneflies was also found when comparing acclimation ability (Shah et al., 2017a).

Another rare study of respiration rate in tropical stoneflies (Jacobsen & Brodersen, 2008) reports

similar Q_{10} values to those reported here for non-stressful temperatures in mid-elevation

stoneflies, confirming our conclusions of lower thermal sensitivity in at least this group of perlid

stoneflies. Though many species in the tropics will be imperiled by warming (Deutsch et al.,

2008; Dillon et al., 2010), the overall mixed support for this prediction in stoneflies a cautionary

tale for generalizations about the vulnerability of tropical species.

Numerous hypotheses could explain the lack of divergence between temperate and

tropical stoneflies. First, stoneflies as a group have a temperate origin and invaded tropical

streams only recently (Baumann & Kondratieff, 1991; Stewart, Baumann, & Stark, 1974).

Studies show that stoneflies are still adapting to tropical habitats (Polato et al., 2018), which may

explain why thermal physiological traits in present-day temperate and tropical populations are

similar. Second, perlid stoneflies and baetid mayflies have different life histories, which may

contribute to differential sensitivity to temperature. As nymphs, semivoltine stoneflies live much

longer than multivoltine mayflies, and may need to cope with multiple years of unpredictable

variation in stream temperature (Clifford, 1982). In fact, many stoneflies enter diapause, unlike

mayflies (Brittain, 1982; Hynes, 1976), giving them a greater ability to withstand adverse

thermal conditions (Brittain, 1990). Once emerged, adult stoneflies often remain close to natal

streams, whereas adult mayflies in tropical (Finn, Encalada, & Hampel, 2016) and temperate

(Polato et al., 2018) regions can disperse more widely. Therefore, stoneflies rely more heavily on

the nymph stage for dispersal (Brittain, 1990) and as a consequence, may experience more

thermal variation moving up-or- downstream than juvenile mayflies. Finally, stoneflies are

important predators of mayflies and midges (Peckarsky, 1980; Tamaris-Turizo, Turizo-Correa, &

del Carmen Zúñiga, 2007). Perlid stoneflies actively pursue prey in the wild (Peckarsky, 1980)

as well as in laboratory settings (Sjöstrom, 1985; A.A.Shah *pers. obs*.). A reliance on aerobic

metabolism and the need to maintain a high metabolic scope for active hunting (Kordas, Harley,

& O'Connor, 2011) could constrain physiological divergence between temperate and tropical

stoneflies, but too few comparative studies of metabolic profiles are available to test the

generality of such hypotheses. Nevertheless, higher maximum metabolic rates (Fig. 6) and higher

Q_{10}s (Fig. 6) do suggest some differences between temperate and tropical populations that could

have ecological consequences when pursuing prey across a broad range of temperatures. Because

perlid stoneflies and baetid mayflies interact as predator and prey in temperate and tropical

streams (Gamboa, Chacón, & Segnini, 2009; Peckarsky, 1980), differential sensitivities to

temperature may have community-level consequences (Gilman, Urban, Tewksbury, Gilchrist, &

Holt, 2010; Grigaltchik, Ward, & Seebacher, 2012; Kordas et al., 2011; Pincebourde & Casas,

2019; Vucic-Pestic, Ehnes, Rall, & Brose, 2011). Our results underscore the importance of

assessing community-level responses to global warming by explicitly testing interactions under

different temperatures (Barton & Schmitz, 2009) and how these responses differ across latitude

(Matassa & Trussell, 2015)

Comparison of metabolic rates and critical thermal maxima as indicators of vulnerability

Some of the results of the present study are surprising because we expected CT$_{MAX}$, a commonly used metric for thermal sensitivity (Diamond et al., 2012; Lutterschmidt & Hutchison, 1997; Piyaphongkul, Pritchard, & Bale, 2012; Ribeiro, Camacho, & Navas, 2012), to coincide with the descending limb of the metabolic rate profiles (Fig. 1). Yet, previously measured values of CT$_{MAX}$ in temperate mayflies occur at much higher temperatures (Fig.1A, B, C). For example, CT$_{MAX}$ was ~29 °C in high elevation temperate mayflies (Shah et al., 2017b) but they began to show elevated metabolic rates and behavioral signs of stress at lower temperatures (~17.5 °C) and an elevated metabolism. The large difference between CT$_{MAX}$ and the onset of stress in this study is likely due to methodology, as rapid ramping or high starting temperatures can inflate estimated values of CT$_{MAX}$ (Rezende, Tejedo, & Santos, 2011; Terblanche, Deere, Clusella-Trullas, Janion, & Chown, 2007). The discrepancy could also arise because CT$_{MAX}$ and metabolic rates are governed by fundamentally different processes. CT$_{MAX}$ is the result of changes in fluidity of cell membranes and whole-organism physiology in response to heat stress (Cossins, Friedlander, & Prosser, 1977; Hochachka & Somero, 2002). Standard metabolic rates arise from different activation energies of enzymes and sensitivity of biochemical processes to temperature (Rolfe & Brown, 1997). Thus, using CT$_{MAX}$ in conjunction with other measures of thermal sensitivity such as metabolic rate provides a more complete assessment of thermal tolerance.

Climate change in Andean and Rocky Mountain Streams

Mountain streams are already increasing in temperature worldwide (Isaak & Rieman, 2013). Temperatures of small mountain streams, such as those in this study, are likely to show

concomitant increases with air temperature (Mohseni & Stefan, 1999; Pilgrim, Fang, & Stefan,

1998). Projections of mean air temperature increases of 2-3 °C in the Andes and 3-4 °C in the

northern Rocky Mountains by the 2050s are therefore troubling (Nogués-Bravo, Araújo, Errea,

& Martínez-Rica, 2007; Rangecroft, Suggitt, Anderson, & Harrison, 2016). However, evidence

suggests that insects of the southern Rocky Mountains are relatively robust to warming because

they may currently occur at suboptimal temperatures (Pyne & Poff, 2017). Moreover, mesocosm

and field common garden experiments showed that temperate baetid mayflies grow faster when

subjected to warmer water (N.L. Poff et al., *unpubl. data*). Given the wide elevation distribution

and broad dispersal ability of these insects (Gill et al., 2016), they may be able to move across

the landscape to find thermally suitable habitats as their environment warms. In contrast to

mayflies of the Rockies, Andean mayfly growth declined with increasing temperature in

mesocosms (N.L. Poff et al., *unpubl. data*) further supporting our conclusion that a warming

environment will negatively affect tropical mayflies. Increased temperatures may also be locally

exacerbated in some Andean streams where riparian forests are being cleared for human

activities. The resulting exposure to increased solar radiation can alone raise stream temperature

(Moore, Spittlehouse, & Story, 2005). The tropical mayflies in our study will experience the

effects of these increasing temperatures leading to a possible mismatch of dissolved oxygen

supply and demand (Pörtner, 2001; Verberk, Bilton, Calosi, & Spicer, 2011; Verberk et al.,

2016b). The ability to maintain an adequate rate of oxygen uptake decreases at higher

temperatures for many aquatic insects (Jacobsen & Brodersen, 2008; Rotvit & Jacobsen, 2013),

thus reducing energy available for activities like movement, growth, and egg development. Thus,

streams need not even reach temperatures as high as CT$_{MAX}$ values to have a negative impact on

insects. This is illustrated in a field experiment, where a moderate 2 °C temperature increase in

the stream resulted in declines in abundance of Eurasian mayflies, presumably as a result of

decreased oxygen in the stream (Verberk, Durance, Vaughan, & Ormerod, 2016a). Whether

tropical mayflies in our study will move upstream to track cooler temperatures is debatable given

the low dispersal rates of aquatic insects in these streams (Gill et al., 2016). Future work should

therefore also focus on whether aquatic insects can move upslope to counteract the effects of

warming in their native streams (Sheldon, 2012).

Acknowledgements

This work would not have been possible without the hard work of many field assistants. We

especially thank M. Rojas, J. Fajardo, T. Waneka, B. Choat, L. Granizo, L. Nagle for joining us

over multiple field seasons. We thank J. Schreckinger, N. Quiroz, M. Celinscak, and E.J.G.

Orejuela in Ecuador and K. L. Casner in the U.S. for help with field logistics. The Ministerio del

Ambiente of Ecuador provided collection permits (Permit nos. 56-IC- FAU/FLO-DPN/MA and

MAE-DNB- CM-2015-0017).

Funding

This research was funded by a U.S. National Science Foundation Dimensions of Biodiversity

grant DEB-1046408; DEB-1045960, and DEB-1045991; and a Graduate Research Fellowship

awarded to A.A.S under Grant No. 1321845.

Author contributions:

A.A.S. and C.K.G. designed the experiments. A.A.S. conducted fieldwork and experiments,

analyzed results, and wrote the manuscript with contributions from all authors.

References

Angilletta, M. J. (2009). *Thermal adaptation: a theoretical and empirical synthesis*. New York:

Oxford University Press.

Angilletta, M. J., Niewiarowski, P. H., & Navas, C. A. (2002). The evolution of thermal

physiology in ectotherms. *Journal of Thermal Biology*, *27*(4), 249–268.

Artacho, P., & Nespolo, R. F. (2009). Natural selection reduces energy metabolism in the garden

snail, Helix Aspersa (Cornu Aspersum). *Evolution*, *63*(4), 1044–1050.

Barton, B. T., & Schmitz, O. J. (2009). Experimental warming transforms multiple predator

effects in a grassland food web. *Ecology Letters*, *12*(12), 1317–1325.

Baumann, R. W., & Kondratieff, B. C. (1991). The stonefly fauna of northern Mexico. In J.

Alba-Tercedor & A. Sanchez-Ortega (Eds.), *Overview and Strategies of Ephemeroptera*

and Plecoptera. Gainesville, Florida.: Sandhill Crane Press, Inc.

Bochdansky, A. B., Grønkjær, P., Herra, T. P., & Leggett, W. C. (2005). Experimental evidence

for selection against fish larvae with high metabolic rates in a food limited environment.

Marine Biology, *147*(6), 1413–1417.

Brittain, J. E. (1982). Biology of mayflies. *Annual Review of Entomology*, *27*(1), 119-147.

Brittain, J. E. (1990). Life history strategies in Ephemeroptera and Plecoptera. In *Mayflies and*

Stoneflies: Life Histories and Biology (pp. 1–12).

Brown, J. H., Marquet, P. A., & Taper, M. L. (1993). Evolution of body size: consequences of an

energetic definition of fitness. *The American Naturalist*, *142*(4), 573–584.

Chown, S. L., Duffy, G. A., & Sørensen, J. G. (2015). Upper thermal tolerance in aquatic insects.

Current Opinion in Insect Science, *11*, 78–83.

Clifford, H. F. (1982). Life cycles of mayflies(Ephemeroptera), with special reference to

voltinism. *Quaestiones Entomologicae*, *18*.(1), 15–90.

Colwell, R. K., Brehm, G., Cardelús, C. L., Gilman, A. C., & Longino, J. T. (2008). Global

warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science*,

322(5899), 258–261.

Cossins, A. R., Friedlander, M. J., & Prosser, C. L. (1977). Correlations between behavioral

temperature adaptations of goldfish and the viscosity and fatty acid composition of their

synaptic membranes. *Journal of Comparative Physiology*, *120*(2), 109–121.

Dell, A. I., Pawar, S., & Savage, V. M. (2011). *Systematic variation in the temperature*

dependence of physiological and ecological traits. *108*(26), 10591–10596.

DeLong, J. P., Bachman, G., Gibert, J. P., Luhring, T. M., Montooth, K. L., Neyer, A., & Reed,

B. (2018). Habitat, latitude and body mass influence the temperature dependence of

metabolic rate. *Biology Letters*, *14*(8).

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., &

Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude.

Proceedings of the National Academy of Sciences, *105*(18), 6668–6672.

Diamond, S. E., Sorger, D. M., Hulcr, J., Pelini, S. L., Toro, I. Del, Hirsch, C., … Dunn, R. R.

(2012). Who likes it hot? A global analysis of the climatic, ecological, and evolutionary

determinants of warming tolerance in ants. *Global Change Biology*, *18*(2), 448–456.

Dillon, M. E., Wang, G., & Huey, R. B. (2010). Global metabolic impacts of recent climate

warming. *Nature*, *467*(7316), 704–706.

Finn, D. S., Encalada, A. C., & Hampel, H. (2016). Genetic isolation among mountains but not

between stream types in a tropical high-altitude mayfly. *Freshwater Biology*, *61*(5), 702–

714.

Gamboa, M., Chacón, M. M., & Segnini, S. (2009). Diet composition of the mature larvae of

four Anacroneuria species (Plecoptera: Perlidae) from the Venezuelan Andes. *Aquatic*

Insects, *31*, 409–417.

Gill, B. A., Kondratieff, B. C., Casner, K. L., Encalada, A. C., Flecker, A. S., Gannon, D. G., …

Funk, W. C. (2016). Cryptic species diversity reveals biogeographic support for the

'mountain passes are higher in the tropics' hypothesis. *Proceedings of the Royal Society B:*

Biological Sciences, *283*(1832).

Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., & Charnov, E. L. (2001). Effects of

size and temperature on metabolic rate. *Science*, *293*(5538), 2248–2251.

Gilman, S. E., Urban, M. C., Tewksbury, J., Gilchrist, G. W., & Holt, R. D. (2010). A framework

for community interactions under climate change. *Trends in Ecology and Evolution*, *25*(6),

325–331.

Grigaltchik, V. S., Ward, A. J. W., & Seebacher, F. (2012). Thermal acclimation of interactions:

differential responses to temperature change alter predator–prey relationship. *Proceedings*

of the Royal Society B: Biological Sciences, *279*(1744), 4058–4064.

Guppy, M., & Withers, P. (1999). Metabolic depression in animals : physiological perspectives

and biochemical generalizations. *Biological Reviews*, *74*, 1–40.

Gutiérrez-Pesquera, L. M., Tejedo, M., Olalla-Tárraga, M. Á., Duarte, H., Nicieza, A., & Solé,

M. (2016). Testing the climate variability hypothesis in thermal tolerance limits of tropical

and temperate tadpoles. *Journal of Biogeography*, *43*(6), 1166–1178.

Heath, A. G. (1973). Ventilatory responses of teleost fish to exercise and thermal stress.

American Zoologist, *13*(2), 491–503.

Hochachka, P. W., & Somero, G. N. (2002). Mechanism and process in physiological evolution.

In *Biochemical Adaptation*. New York: Oxford University Press.

Huey, R. B., & Hertz, P. E. (1984). Is a Jack-of-all-temperatures a master of none? *Evolution*,

38(2), 441–444.

Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm

performance. *Trends in Ecology & Evolution*, *4*(5), 131–135.

Huey, R. B., & Kingsolver, J. G. (1993). Evolution of resistance to high temperature in

ectotherms. *The American Naturalist*, *142*.

Huey, R. B., & Kingsolver, J. G. (2019). Climate warming, resource availability, and the

metabolic meltdown of ectotherms. *The American Naturalist*, *194*(6), E140–E150.

Huey, R. B., & Stevenson, R. D. (1979). Integrating thermal physiology and ecology of

ectotherms: a discussion of approaches. *American Zoologist*, *19*(1), 357–366.

Hynes, H. B. N. (1976). Biology of Plecoptera. *Annual Review of Entomology*, *21*(1), 135–153.

Isaak, D. J., & Rieman, B. E. (2013). Stream isotherm shifts from climate change and

implications for distributions of ectothermic organisms. *Global Change Biology*, *19*(3),

742–751.

Jacobsen, D., & Brodersen, K. P. (2008). Are altitudinal limits of equatorial stream insects

reflected in their respiratory performance? *Freshwater Biology*, *53*(11), 2295–2308.

Johnston, I. A., Clarke, A., & Ward, P. (1991). Temperature and metabolic rate in sedentary fish

from the Antarctic, North Sea and Indo-West Pacific Ocean. *Marine Biology*, *109*(2), 191–

195.

Kellermann, V., Chown, S. L., Schou, M. F., Aitkenhead, I., Janion-Scheepers, C., Clemson, A.,

… Sgrò, C. M. (2019). Comparing thermal performance curves across traits: How

consistent are they? *Journal of Experimental Biology*, *222*(11).

Kordas, R. L., Harley, C. D. G., & O'Connor, M. I. (2011). Community ecology in a warming

world: The influence of temperature on interspecific interactions in marine systems. *Journal*

of Experimental Marine Biology and Ecology, *400*(1–2), 218–226.

Lighton, J. R. B., & Turner, R. J. (2004). Thermolimit respirometry: An objective assessment of

critical thermal maxima in two sympatric desert harvester ants, Pogonomyrmex rugosus and

P. californicus. *Journal of Experimental Biology*, *207*(11), 1903–1913.

Lutterschmidt, W. I., & Hutchison, V. H. (1997). The critical thermal maximum: history and

critique. *Canadian Journal of Zoology*, *75*(10), 1561–1574.

Martin, T. L., & Huey, R. B. (2008). Why "suboptimal" is optimal: Jensen's Inequality and

ectotherm thermal preferences. *American Naturalist*, *171*(3), 102–118.

Matassa, C. M., & Trussell, G. C. (2015). Effects of predation risk across a latitudinal

temperature gradient. *Oecologia*, *1*, 775–784.

Mohseni, O., & Stefan, H. G. (1999). Stream temperature/air temperature relationship: A

physical interpretation. *Journal of Hydrology*, *218*(3–4), 128–141.

Moore, R. D., Spittlehouse, D. L., & Story, A. (2005). Riparian microclimate and stream

temperature response to forest harvesting: A review. *Journal of the American Water*

Resources Association, *41*(4), 813–834.

Nogués-Bravo, D., Araújo, M. B., Errea, M. P., & Martínez-Rica, J. P. (2007). Exposure of

global mountain systems to climate warming during the 21st Century. *Global*

Environmental Change, *17*(3–4), 420–428.

Packard, G. C., & Boardman, T. J. (1999). The use of percentages and size-specific indices to

normalize physiological data for variation in body size: wasted time, wasted effort?

Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology,

122(1), 37–44.

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, *421*(6918), 37.

Payne, N. L., & Smith, J. A. (2017). An alternative explanation for global trends in thermal

tolerance. *Ecology Letters*, *20*(1), 70–77.

Peckarsky, B. L. (1980). Predator-prey interactions between stoneflies and mayflies: Behavioral

observations. *Ecology*, *61*(4), 932–943.

Pilgrim, J. M., Fang, X., & Stefan, H. G. (1998). Stream temperature correlations with air

temperature in Minnesota: Implications for climate warming. *Journal of the American*

Water Resources Association, *34*(5), 1109–1121.

Pincebourde, S., & Casas, J. (2019). Narrow safety margin in the phyllosphere during thermal

extremes. *Proceedings of the National Academy of Sciences*, *116*(12), 5588–5596.

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater

vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, *569*(7754), 108.

Piyaphongkul, J., Pritchard, J., & Bale, J. (2012). Can tropical insects stand the heat? A case

study with the brown planthopper Nilaparvata lugens (Stål). *PLoS One*, *7*(1), e29409.

Polato, N. R., Gill, B. A., Shah, A. A., Gray, M. M., Casner, K. L., Barthelet, A., … Zamudio, K.

R. (2018). Narrow thermal tolerance and low dispersal drive higher speciation in tropical

mountains. *Proceedings of the National Academy of Sciences*, *115*(49), 12471–12476.

Pörtner, H. O. (2001). Climate change and temperature-dependent biogeography: Oxygen

limitation of thermal tolerance in animals. *Naturwissenschaften*, *88*(4), 137–146.

Pörtner, H. O. (2002). Climate variations and the physiological basis of temperature dependent

biogeography: systemic to molecular hierarchy of thermal tolerance in animals.

Comparative Biochemistry and Physiology Part A, *132*, 739–761.

Pörtner, H. O., Bennett, A. F., Bozinovic, F., Clarke, A., Lardies, M. A., Lucassen, M., … Stillman, J. H. (2006). Trade-offs in thermal adaptation: the need for a molecular to ecological integration. *Physiological and Biochemical Zoology*, *79*(2), 295–313.

Pörtner, H. O., Bock, C., & Mark, F. C. (2017). Oxygen- & capacity-limited thermal tolerance: Bridging ecology & physiology. *Journal of Experimental Biology*, *220*(15), 2685–2696.

Pörtner, H. O., & Knust, R. (2007). Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science*, *315*(5808), 95–97.

Pyne, M. I., & Poff, N. L. (2017). Vulnerability of stream community composition and function to projected thermal warming and hydrologic change across ecoregions in the western United States. *Global Change Biology*, *23*(1), 77–93.

Rangecroft, S., Suggitt, A. J., Anderson, K., & Harrison, S. (2016). Future climate warming and changes to mountain permafrost in the Bolivian Andes. *Climatic Change*, *137*, 231–243.

Rezende, E. L., Tejedo, M., & Santos, M. (2011). Estimating the adaptive potential of critical thermal limits: methodological problems and evolutionary implications. *Functional Ecology*, *25*(1), 111–121.

Ribeiro, P. L., Camacho, A., & Navas, C. A. (2012). Considerations for assessing maximum critical temperatures in small ectothermic animals: Insights from leaf-cutting ants. *PLoS ONE*, *7*(2), e32083.

Rolfe, D. F. S., & Brown, G. C. (1997). Cellular energy utilization and molecular origin of standard metabolic rate in mammals. *Physiological Reviews*, *77*(3), 731–758.

Rotvit, L., & Jacobsen, D. (2013). Temperature increase and respiratory performance of macroinvertebrates with different tolerances to organic pollution. *Limnologica*, *43*(6), 510–

515.

Schulte, P. M. (2015). The effects of temperature on aerobic metabolism: towards a mechanistic

understanding of the responses of ectotherms to a changing environment. *Journal of*

Experimental Biology, *218*(12), 1856–1866.

Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal performance curves, phenotypic

plasticity, and the time scales of temperature exposure. *Integrative and Comparative*

Biology, *51*(5), 691–702.

Shah, A. A., Funk, W. C., & Ghalambor, C. K. (2017a). Thermal acclimation ability varies in

temperate and tropical aquatic insects from different elevations. *Integrative and*

Comparative Biology, *57*(5), 977–987.

Shah, A. A., Gill, B. A., Encalada, A. C., Flecker, A. S., Funk, W. C., Guayasamin, J. M., …

Ghalambor, C. K. (2017b). Climate variability predicts thermal limits of aquatic insects

across elevation and latitude. *Functional Ecology*, *31*(11), 2118–2127.

Sheldon, A. L. (2012). Possible climate-induced shift of stoneflies in a southern Appalachian

catchment. *Freshwater Science*, *31*(3), 765–774.

Sjöstrom, P. (1985). Hunting behaviour of the perlid stonefly nymph Dinocras cephalotes

(Plecoptera) under different light conditions. *Animal Behavior*, *33*, 534–540.

Stewart, K. W., Baumann, R. W., & Stark, B. P. (1974). The distribution and past dispersal of

southwestern United States Plecoptera. *Source: Transactions of the American*

Entomological Society, *99*(4), 507–546.

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and

latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1713),

1823–1830.

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global

redistribution of animals. *Nature Climate Change*, *2*(9), 686–690.

Tamaris-Turizo, C. E., Turizo-Correa, R. R., & del Carmen Zúñiga, M. (2007). Spatial-temporal

distribution and feeding in nymphs of Anacroneuria (Insecta: Plecoptera: Perlidae) in the

Gaira River (Sierra Nevada de Santa Marta, Colombia). *Caldasia*, 375–385.

Terblanche, J. S., & Chown, S. L. (2007). The effects of temperature, body mass and feeding on

metabolic rate in the tsetse fly Glossina morsitans centralis. *Physiological Entomology*,

32(2), 175–180.

Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C., & Chown, S. L. (2007). Critical

thermal limits depend on methodological context. *Proceedings of the Royal Society B:*

Biological Sciences, *274*(1628), 2935–2943.

Vannote, R. L., & Sweeney, B. W. (1980). Geographic analysis of thermal equilibria: A

conceptual model for evaluating the effect of natural and modified thermal regimes on

aquatic insect communities. *The American Naturalist*, *115*(5), 667–695.

Verberk, W. C. E. P., Bilton, D. T., Calosi, P., & Spicer, J. I. (2011). Oxygen supply in aquatic

ectotherms: Partial pressure and solubility together explain biodiversity and size patterns.

Ecology, *92*(8), 1565–1572.

Verberk, W. C. E. P., Durance, I., Vaughan, I. P., & Ormerod, S. J. (2016a). Field and laboratory

studies reveal interacting effects of stream oxygenation and warming on aquatic ectotherms.

Global Change Biology, *22*(5), 1769–1778.

Verberk, W. C. E. P., Overgaard, J., Ern, R., Bayley, M., Wang, T., Boardman, L., &

Terblanche, J. S. (2016b). Does oxygen limit thermal tolerance in arthropods? A critical

review of current evidence. *Comparative Biochemistry and Physiology -Part A : Molecular*

and Integrative Physiology, *192*, 64–78.

Vucic-Pestic, O., Ehnes, R. B., Rall, B. C., & Brose, U. (2011). Warming up the system: higher

predator feeding rates byt lwer energetic differences. *Global Change Biology*, *17*(3), 1301–

1310.

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., … Bairlein, F.

(2002). Ecological responses to recent climate change. *Nature*, *416*(6879), 389.