Single cell transcriptome of CD8+ T cells in multiple cancers reveals comprehensive exhaustion

associated mechanisms

Yunmeng Bai[1], Zixi Chen[1], Xiaoshi Chen[1], Ziqing He[1], Jie Long[2,3,4], Shudai Lin[1], Hongli Du[1*]

[1] School of Biology and Biological Engineering, South China University of Technology, Guangzhou 51

0006, China

[2] Department of General Surgery, Guangzhou Digestive Disease Center, Guangzhou First People's

Hospital, School of Medicine, South China University of Technology, Guangzhou, Guangdong, 510180,

China.

[3] Chronic Disease Laboratory, Institutes for Life Sciences and School of Medicine, South China

University of Technology, Guangzhou, 510006, China.

[4] Institute of Immunology and School of Life Sciences, University of Science and Technology of China,

Hefei, Anhui, 230027, China.

[*]Correspondence: hldu@scut.edu.cn; Tel.: +86-020-3938-0667

Abstract

T cell exhaustion is one of the mechanisms that cancer cells get rid of control from the immune system. Single-cell RNA sequencing has showed superiority on immunity mechanism in recent studies. Here, we collected more than 6000 single CD8+ T cells from three cancers including CRC, HCC and NSCLC, and identified five clusters of each cancer. We obtained 71 and 159 DEGs for pre_exhausted or exhausted vs. effector comparison in all three cancers, respectively. Specially, we found some key genes including the four exhaustion associated genes of *PDCD1*, *HAVCR2*, *TIGIT* and *TOX,* and two vital genes of *CD69* and *JUN* in the interaction network. Additionally, we identified the gene *SAMSN1* which highly expressed in the exhausted cells had a poor overall survival and played a negative role in immunity. We summarized the putative interrelated mechanisms of above key genes identified in this study by integrating the reported knowledge. Furthermore, we explored the heterogeneous and preference of exhausted CD8+ T cells in each patient and found only one exhausted sub-cluster existed in the most of patients, especially in CRC and HCC. As far as we know, this is the first time to study the mechanism of T cell exhaustion with the data of single-cell RNA sequencing of multiple cancers. Our study may facilitate the understanding of the mechanism of T cell exhaustion, and provide a new way for functional research of single-cell RNA sequencing data across cancers.

Keywords: Single-cell RNA sequencing; multiple cancers; T cell exhaustion; tumor heterogeneous

1. Introduction

T cell exhaustion (Tex), a hyporesponsive state of T cells, was originally described in CD8+ T cells during chronic lymphocytic choriomeningitis virus (LCMV) of mice[1]. In recent years, the phenomenon of Tex has also been found in cancer[2, 3]. It has been reported that exhausted T cells in

cancers share many similarities with that in chronic infection[4], and play a significant role in

tumorigenesis[5]. The main reasons leading to Tex in chronic infection and cancer are as following[6]:

long-term and persistent explosion to antigens[7]; the upregulation of inhibitory receptors (IRs)

including PDCD1, CLTA4, HAVCR2, LAG3 and 2B4[8]; complex effects of soluble factors such as

cytokines IL-10 and transforming growth factor-β (TGFβ) [9] ; the expression of transcription factor

including T-bet and Eomes[10] and corresponding epigenetic regulation[11].However, at present, to reveal

molecular mechanisms of exhausted T cells, most researches are still focusing on chronic infected

mouse models by microarray or RNA sequencing[8, 12]. Tex is a general trend of tumor, which can be

used as one of the main targets of immunosuppression therapy to save T cell from exhaustion and

reactivate the cytotoxicity of T cells, providing a new opportunity for clinical immunotherapy[8].

Nevertheless, due to the complexity and heterogeneity of cancer, the concrete mechanism of Tex in

cancer is still unclear. Thus, a deeper understanding of this mechanism is urgently needed.

Currently, single-cell RNA sequencing (scRNA-seq) has clearly revealed some new mechanisms and

phenomena of cancer with the advantages of high accuracy and reproducibility[13-15].   Using single cell

transcriptome profiling, we can identify new types of immune cells which can't be revealed at the

original tissue level and can construct a developmental trajectory for immune cells which can reveal the

random heterogeneity[16]. These new findings are useful to better understand the immune system and its

mechanism of action on tumors. Notably, this technology makes it possible to explore complicated

tumor microenvironment including tumor-infiltrating lymphocytes (TILs) in melanoma, head and neck

cancer, breast cancer and glioblastoma cancer[17-20]. Thus, using advantage of scRNA-seq to analyze T

cells and obtain the hallmarks of exhausted T cells can bring a new therapeutic strategy on clinical

cancer treatment.

In the present study, due to the vital role of CD8+ T cells in eliciting antitumor responses[21], we

integrated single cell sequencing data from colorectal cancer (CRC), liver cancer (HCC), and non-small

cell lung cancer (NSCLC) to analyze CD8+ T cells in various cancers. By classifying cells into

different clusters with the unique markers, we identified five clusters of each cancer. Furthermore, we

compared the pre_exhausted or exhausted cells with effector cells to make a deeper understanding of

exhaustion mechanism. Additionally, the sub-clusters of exhausted cells in each patient revealed the

individual differences and preferences. Overall, the findings of molecules changed in pre_exhasuted

and exhausted clusters may provide the potential targets to anticancer therapy, and the different

exhausted sub-clusters of each patient help the understanding of exhaustion heterogeneity.

2. Materials and methods

2.1 Data Resources and Preprocessing

Single cell transcriptome profiling of human T cells in three cancers including CRC, HCC and

NSCLC was obtained from the GEO database (GSE108989, GSE99254, GSE98638), including raw

read count and TPM data. According to the annotations, we isolated CD8+ T cells from peripheral

blood (PTC), adjacent normal (NTC) and tumor tissues (TTC). To filtered out the low-quality cells, we

excluded cells with fewer than 3000 detected genes and the expression of the housekeeping gene

($\log_2(\text{TPM}_{ACTB}/10+1)$) below 2.5[22].

2.2 Unsupervised Clustering

For each dataset, we selected the genes with top n highest variance where n changed from 500,

1000, 1500, 2000, 2500, 3000. Then the expression data of these genes was used to perform clustering

by single-cell consensus clustering (SC3)[23]. To obtain the reliable result, we set the number of clusters

(k) from 2 to 10. Only the genes with p value < 0.05 and auroc > 0.75 could be considered as markers.

For each SC3 run, we focused on three indicators - the silhouette, the consensus matrix and the cluster

specific genes, which could help us to fix the suitable n and k. When determined the clusters, we

mapped the markers to the CellMarker databases[24] to find out the characteristics of each cluster.

2.3 Trajectory Analysis

To explore the potential functional changes of CD8+ T cell of different clusters for each cancer,

we performed development trajectory analysis by Monocle[25] with the top 100 cluster-specific genes of

each cluster. The differentially expressed genes (DEGs) between each two cluster pairs were identified

by R package edgeR[26] using the criteria as following: (1) the mean of CPM value was greater than 1;

(2) false discovery rate (FDR) < 0.05; (3) the absolute value of logFC > 1. We defined the

cluster-specific genes in each cluster of each cancer based on the condition that its expression was

upregulated when compared with those in the other clusters.

2.4 Differentially Expressed Genes, Functional Annotation and Transcription Network Analysis

To further get known the potential mechanisms that leading to the exhausted CD8+ T cells, we

compare genes between pre_exhausted or exhausted cluster in tumor and effector cluster in peripheral

blood. The cell numbers of the two groups participating in the comparison were set as m and n, and the

smaller one was defined as p. A gene was defined as exhaustion associated DEG if (1) the number of

expressed cells > p; (2) FDR < 0.05; (3) the absolute value of logFC > 1. To find the common

regulation mechanisms of Tex, we identified the DEGs with the same regulatory status in three cancers.

Next, the DEGs were subjected to the Gene Ontology(GO), including biological process (BP), cell

component (CC), molecular function (MF), and the Kyoto Encyclopedia of Genes and Genomes

(KEGG) pathway enrichment analysis by clusterProfiler[27] with adjusted p value < 0.05.

To find out the interaction among these DEGs, Search Tool for the Retrieval of Interacting Genes

database (STRING) was used to construct the protein-protein interaction (PPI) network, which was

visualized by Cytoscape[28], based on the combined score of 0.4[29]. Additionally, the Molecular

Complex Detection (MCODE) as a Cytoscape plugin was performed to explore the closely connected

regions with default parameter[30]. Furthermore, to investigate the potential upstream regulators that

affect gene differentially expression, we integrated three transcription factor targets databases,

including ChIP-X enrichment analysis (ChEA)[31], Encyclopedia of DNA Elements (ENCODE)[32] from

Harmonizonme[33] and Transcriptional Regulatory Relationships Unraveled by Sentence-based

Text-mining (TRRUST)[34] to reveal regulatory relationships among DEGs.

2.5 Survival analysis

The TCGA data[35] including COAD (colon adenocarcinoma), LIHC (liver hepatocellular

carcinoma), LUAD (lung adenocarcinoma), LUSC (lung squamous cell carcinoma) was used to

explore the relationship between gene expression levels of the key genes and patients' overall survival.

LUAD and LUSC were integrated as NSCLC. The mRNA expression data and corresponding clinical

annotation files were obtained from TCGA. To correct the effector of CD8+ T cells[36], the expression

of genes were divided by that of the geometric mean of *CD3D*, *CD3E*, *CD3G*, *CD8A* and *CD8B*. The

threshold was defined as the median of relative expression. The samples with relative expression below

the threshold were classified as ''low expression'' group, and the samples with relative expression

above the threshold were classified as "high expression" group. R package survival was used to

perform survival analysis.

2.6 Sub-cluster of exhausted cells

We further clustered the exhausted cells into more specific sub-clusters using the same pipeline of

133 SC3 as mentioned above. Especially, we displayed the distribution of each sub-cluster of each patient.

134 The differential genes expression analysis between exhausted sub-clusters and effector cells was

135 performed by edgeR with the same criteria as mentioned above. To further explore the function of

136 each sub-cluster, we performed gene set variation analysis (GSVA)[37], a sensitive technique that can

137 detect underlying pathways over a sample population in an unsupervised manner, with the DEGs of

138 the exhausted sub-clusters vs. effector comparison and the KEGG pathways in canonical gene sets

139 (C2) from the Molecular Signature Database (MSigDB)[38] as the reference gene set. In addition,

140 comparisons of single-cell enrichment scores were performed using the R package limma[39].

141 Differentially enriched pathways were defined as the number of genes >10 and FDR adjusted p value

142 < 0.05 as described previously[40].

144 **3. Results**

145 **3.1 Single CD8+ T cell transcriptome landscape and processing**

146  We obtained single-cell transcriptome data of human T cells from GEO database, including 12

147 patients from CRC, 6 patients from HCC and 14 patients from NSCLC. For CD8+ T cells, low-quality

148 cells were filtered out, remaining 2967, 1214, 2053 cells, respectively (Table 1). We noticed that the

149 cells in NSCLC were filtered out more than half.

151   Table 1. The number of CD8+ T cells of different tissues in three cancers

	N	P	T
CRC	798(1072)	810(1132)	1359(1722)
HCC	262(412)	345(563)	607(777)

	NSCLC	351(934)	524(1323)	1178(2182)

152   Outside the brackets means the number of post-filtered cells, inside means pre-filtered cells

3.2 CD8+ T cell clustering

155   To reveal the potential functional sub-clusters of CD8+ T cells, we performed unsupervised

clustering by SC3. As for CRC, finally we chose the top 2500 genes with highest variance to divide

cells into 5 clusters (Figure 1A). Similarly, the top 1000 genes were used to divide into 5 clusters in

HCC and NSCLC (Figure 1B, 1C).

159   To identify each cluster of three cancers, we compared the markers of each 15 clusters with the

CellMarker database. For example, the 9 of 10 markers (*CCR7, LEF1, SELL, LDLRAP1, NOSIP,*

PRKCQ−AS1, TXK, TMEM123, NELL2) in CRC_C4 were found in the cell markers of naïve CD8+ T

cell, then we identified it as naïve CD8+ T cell. Taken together, we defined the clusters for each cancer

with known markers: CRC_C4, HCC_C5, NSCLC_C4 were identified as naïve T cells, CRC_C1,

HCC_C3, NSCLC_C1 were identified as effector T cells, CRC_C3, HCC_C2, NSCLC_C3 were

identified as exhausted T cells. CRC_C2 was identified as natural killer T (NKT) cells and HCC_C4

was identified as mucosal-associated invariant T (MAIT) cells (Table S1). While in NSCLC, C2

couldn't be identified as a defined cluster according to the CellMarker, in which *ZNF683* was one of

the markers belonging to natural killer T (NKT) cell. Thus, we called CRC_C2, HCC_C4, NSCLC_C2

as unconventional T cells[41] temporarily which were excluded in the following analysis. We noted that

there was a common cluster with a same marker named *GZMK* that couldn't be well-defined, either. To

figure out the potential biological functions of these clusters, we performed trajectory analysis by

monocle (Figure 1D, 1E, 1F). Except for the known clusters, we could see that CRC_C5, HCC_C1,

NSCLC_C5 located between effector and exhausted CD8+ T cells and appeared a transition state, thus

them were defined as "pre_exhausted" CD8+ T cells. In order to check the classification effect, we

used the t-Distributed Stochastic Neighbor Embedding (t-SNE), a non-linear dimensionality reduction

technique, by Rtsne package with all the markers of all clusters of each cancer to visualize the cell

population clusters in two dimensions (Figure 1G, 1H, 1I). We could see that in CRC, five clusters with

different functions were mostly clustered in different parts with obvious boundaries. In addition, the

same clustering phenomenon could also be observed in HCC and NSCLC. We combined clusters of

three cancers with their all markers and found that the same functional cells from the different cancer

types were clustered into the same clusters (Figure 1J).

3.3 Differentially expressed genes associated with exhaustion

The proportion of different clusters varied greatly in different tissues (Figure 2A). Peripheral

blood had most naïve and effector CD8+ T cells in all three cancers, tumor tissues had most

pre_exhausted and exhausted CD8+ T cells, while there was no regular pattern in normal tissues. The

difference in normal tissue may be related to tissue specificity. Since the most of effector cells existed

in peripheral blood in three cancers and there showed no difference among peripheral blood, normal

and tumor tissues (Figure S1), we compared pre_exhausted or exhausted cells in tumor tissues with

effector cells in peripheral blood to explore the exhaustion associated mechanism.

In the pre_exhausted vs. effector comparison, there were 73 DEGs existed in all three cancers, 71

of whom were in the same regulatory status (Figure 2B) including the known exhaustion marker

PDCD1 (programmed cell death□1) and *CD27* (CD27 molecule). The top 15 significant GO and

KEGG terms based on the p value were shown as Figure 2D and 2E. We could find that most terms

were related to the production of chemokine and response to tumor necrosis factor which can facilitate

the immunoregulation effects[42, 43]. Apart from that, we found other terms associated with immunity.

For example, there were 7 upregulated DEGs, including *GPR171* (G protein-coupled receptor 171),

TNFAIP3 (TNF alpha induced protein 3), *CD96* (CD96 molecule), *SAMSN1* (SAM domain, SH3

domain and nuclear localization signals 1, also known as *HACS1*), *ZFP36* (ZFP36 ring finger protein),

DUSP1 (dual specificity phosphatase 1), *PDCD1,* enriched in the negative regulation of immune

system process.

In the exhausted vs. effector comparison, there were 159 DEGs existed in all three cancers (Figure

2C) including multiple known exhaustion markers such as *PDCD1*, *HAVCR2* (hepatitis A virus cellular

receptor 2, also known as T cell immunoglobulin and mucin-domain containing-3, Tim-3), *TIGIT* (T

cell immunoglobulin and ITIM domain), *TOX* (thymocyte selection associated high mobility group

box). The top 15 significant GO and KEGG terms were shown as Figure 2F and 2G. Several terms

related to glycolytic process which was the main metabolism pathway of T cells to sustain effector

function after antigen encounter[44] were enriched. Besides, pathways related to virus infection,

NF-kappa B signaling pathway were also enriched. Correspondingly, there were 14 DEGs, including 9

upregulated genes *PDCD1*, *TIGIT*, *HAVCR2*, *CCL3* (C-C Motif Chemokine Ligand 3), *DUSP1*,

TNFAIP3, *PAG1* (phosphoprotein membrane anchor with glycosphingolipid microdomains 1), *CDK6*

(cyclin dependent kinase 6), *SAMSN1*, participated in the negative regulation of immune system

process.

Interestingly, we found several common terms between pre-exhausted and exhausted cells,

including cell chemotaxis, leukocyte migration, chemokine signaling pathway, cytokine-cytokine

receptor interaction and so on. Additionally, the pathways existed only in pre_exhausted cells included

cytolytic granule and interleukin-2 production, which could represent the effector functions of CD8+ T

cells, showing the features in transition of pre_exhausted cells. The exhausted cells most enriched in

the negative regulation of cytokine production, showing the further dysfunction of T cells. The results

of functional analysis of DEGs was shown in Table S2 and S3.

It was obvious that as the cell state changed from pre-exhausted to exhausted cells, the number of

DEGs also increased, especially the exhaustion associated markers. Moreover, the expression level of

the common markers increased from pre_exhausted to exhausted cells (Figure S2). Interestingly, we

compared the genes existed in both pre_exhausted and exhausted cells and filtered those with greater

fold change in the exhausted cells (Figure 3A). We found *SAMSN1* was dysregulated during exhaustion

state, which participated in the negative regulation of immune system process in both pre_exhausted

and exhausted cells, showed poor survival in the validation based on TCGA data except for CRC

(Figure 3B).

Moreover, we performed the PPI network analysis of all 193 DEGs in pre_exhausted or exhausted

vs. effector comparison and found seven significant functional modules (Figure 3C and Figure S3). The

module with the highest score consisted of 15 nodes and 101 edges, in which *CD69* was the seed gene

and linked with 26 genes.

To further explore the factors affecting the expression of DEGs, we integrated the transcription

factor (TF) from CHEA, ENCODE and TRRUST databases, and found the TFs with dysregulated

expression were ATM (ATM serine/threonine kinase), RORA (RAR Related Orphan Receptor A),

TBX21 (T-Box Transcription Factor 21), JUNB (JunB Proto-Oncogene, AP-1 Transcription Factor

Subunit), RBPJ (Recombination Signal Binding Protein For Immunoglobulin Kappa J Region),

TNFAIP3 and JUN (Jun proto-oncogene, AP-1 transcription factor subunit) (Figure 3D, Figure S4).

ATM, RORA, TBX21, which were the TFs of *DUSP1*, *TUBA1B* (tubulin alpha 1b), *CXCR3* (C-X-C

motif chemokine receptor 3) and *IFNG* (Interferon Gamma) respectively, were significantly

downregulated in exhausted cells. Among the upregulated factors, JUN regulated more than 100 DEGs

and may play a critical role in the occurrence and development of exhausted CD8+ T cells. Therefore,

these TFs may lead to the dysregulated expression of DEGs during T cell exhaustion.

3.4 Sub-cluster of exhausted cells

Exhausted T cell populations are heterogeneous[45]. In the further classification of exhausted cells,

the top 1500 genes with highest variance were used to divide exhausted cells into 2 sub-clusters in

CRC, the top 3000 genes in HCC to 2 sub-clusters, and the top 500 genes in NSCLC to 2 sub-clusters

(Figure 5A, 5B, 5C). In the visualization of exhausted sub-clusters, the distribution of exhausted cells

of each patient tended to appear in one sub-cluster(Figure 5D, 5E, 5F), for instance, P0413, who was

diagnosed at stage III of CRC, mostly exhausted in sub-cluster I, while P0909, who was diagnosed at

stage I of CRC, mostly exhausted in sub-cluster II, illustrating the individual differences.

Since different sub-clusters existed in each cancer, we tried to characterize the features of

sub-clusters across cancers. We combined the six sub-clusters with re-clustering and found that most

cells of sub-cluster I could be distinguished in three cancers (Figure 6A) and the cells of different

cancers were clustered relatively closely, illustrating the tumor specificity, while the cells of sub-cluster

II were clustered disorderly and couldn't distinguish the cells from different cancers, showing no tumor

specificity.

In order to find out the potential different mechanisms of each sub-cluster, we compared the

sub-cluster of exhausted cells from each tumor with effector cells in peripheral blood using the

aforementioned criteria. The numbers of DEGs and intersection of each group were shown by

UpSetR[46] (Figure 6B). There were 63 common DEGs among the six sub-clusters, including 27

upregulated and 36 downregulated genes in all three cancers, in which 60 DEGs appeared in the above

common DGEs of the exhausted vs. effector comparisons except 3 genes including *PLAC8* (placenta

associated 8), *TAPT1-AS1* (transmembrane anterior posterior transformation 1- antisense RNA 1) and

GZMB (Granzyme B).

In the KEGG pathways enriched by GSVA (Figure 7A-F), there was one common pathway in 6

sub-clusters: cytokine-cytokine receptor interaction, which also existed in the above analysis of

pre_exhausted or exhausted clusters compared with effector cluster. Additionally, the widely enriched

pathways, including pathways in cancer, natural killer cell mediated cytotoxicity and ribosome, were

found in 5 sub-clusters. What's more, the unique pathways belonging to each exhausted sub-cluster

might illustrate the specific difference for each exhausted sub-cluster, which were shown in Table S5.

For example, Wnt signaling pathway existed only in the sub-cluster I of CRC; apoptosis pathway

existed only in the sub-cluster I of NSCLC. These differences may contribute to the various

mechanisms of sub-clusters.

4. Discussion

T cell exhaustion is a dysfunctional state and is possible to be used in immunotherapy because of

its reversibility[47]. Combining with single-cell RNA sequencing, which can facilitate detecting the

transcriptome on the level of single cell[48], we can shed light on the complication of tumor-infiltrating

T cells. In order to study the common mechanisms of multiple cancers, we performed transcriptomic

analysis of single CD8+ T cells isolated from three cancers including CRC, HCC, NSCLC. As expected,

we identified some known genes related to T cell exhaustion, such as *PDCD1*, *HAVCR2*, *TIGIT*, *TOX*.

Moreover, two vital genes *CD69* and *JUN* in the interaction network and the gene *SAMSN1* which

highly expressed in the exhausted cells were also identified. Furthermore, the distribution of exhausted

T cells for each patient tended to appear in one exhausted sub-cluster, which indicated that the

exhaustion mechanism in each individual might have homogeneity.

In the functional enrichment analysis, several DEGs played a negative regulation of immune

system process and led to the pre-exhausted or exhausted state of CD8+ T cells. To be specific, the

gene *PDCD1* encodes an inhibitory receptor and functions as an immune checkpoint. In the current

study, *PDCD1* was significantly upregulated in the pre_exhausted and exhausted cells, which was

consistent with previous studies[49, 50]. The upregulation of *PDCD1* played a crucial role in the

regulation of CD8+ T cell exhaustion and blockade of the interaction between PDCD1 and its ligand

PD-L1 could restore CD8+ effector function[49, 50]. It has been reported that the combination of PDCD1

and PD-L1 can recruit phosphatases[51], dampen TCR signaling[52], inhibit the PI3K signaling

pathway[53], suppress proliferation and cytokine production and finally result in CD8+ T cell exhaustion.

Besides, the gene *HAVCR2*, which encodes an inhibitory receptor, was upregulated in the exhausted

cells in the present study. The previous studies indicated that the binding of its ligand Galectin 9 can

mediate the release of BAT3 via tyrosine phosphorylation in the HAVCR2 tail and lead to inhibit the

PI3K signaling pathway[54, 55]. In the models of mouse and human tumors, using the antibodies of

HAVCR2 can rescue the exhausted T cells[56, 57], revealing that *HAVCR2* was associated with

exhaustion of CD8+ T cell. Moreover, the inhibitory receptor encoded by gene *TIGIT* acts as T cell

intrinsic inhibitor with its ligand CD155 by restraining effector cell activation and proliferation, leading

to the decreased effector function and cytokine production[58], and the knockdown of *TIGIT* can revert

the decreased production of cytokines[59]. *TIGIT* was upregulated in the exhausted cells according to

our study, showing its potential key role in the exhaustion of CD8+ T cells. Except for the inhibitory

gene, the critical regulator of exhausted CD8+ T cells[60], *TOX*, was upregulated in the exhausted cells

in the current study. As previously reported, TOX-knockout cells showed low expression of *PDCD1*,

ENTPD1, *TIGIT*, *HAVCR2* as well as high expression of the transcription factors genes *TCF7*, *LEF1*

and *ID3*[60, 61], while TOX-overexpressed cells resulted in reduced TNF and IFNγ expression as well as

increased *PDCD1*, *TIGIT*, *LAG3* levels[62], which supported that *TOX* could drive T cell exhaustion.

The genes described above were all upregulated in pre-exhausted or exhausted T cells, which could

result in exhaustion of CD8+ T cell.

Apart from that, we found some other crucial genes. *CD69* is an early activation marker of T

lymphocytes[63] and acts as a negative regulator of anti-tumor responses[64]. In the PPI network analysis,

CD69 was a hub gene of the module with the highest score and was upregulated in both pre_exhausted

and exhausted cells. CD69 can mediate the cell retention via the interaction with S1PR1

(Sphingosine-1-Phosphate Receptor 1) which acts as a central mediator of lymphocyte output[65]. *CD69*

was upregulated and *S1PR1* was downregulated significantly in the exhausted cells according to our

study (Figure 3A). Thus we deduced that upregulated CD69 binds with downregulated S1PR1, leading

to the impairment of lymphocyte output during an immune response[65]. The upregulated CD69

expressed on tumor-infiltrating T cells might interact with unknown ligands in the tumor to lead to the

persistent antigen stimulation, which resulted in T cell exhaustion with the overexpression of inhibitory

genes such as *PDCD1* and *HAVCR2*[66]. In the murine breast cancer model, *Cd69*$^{-/-}$ mice can reduce

tumor growth and metastasis, increase the number of tumor-infiltrating lymphocytes, enhance IFNγ

production and attenuate the T cell exhaustion[67]. In light of these researches, the upregulation of *CD69*

and downregulation of *S1PR1* might promote the development of T cell exhaustion. Additionally, *JUN*

was a proto-oncogene and involved in the JNK signaling pathway which played a vital role in the

proliferation and differentiation of CD8+ T cells[68]. Based on our findings, *JUN* was upregulated in

both pre_exhausted and exhausted cells and regulated over 100 DEGs. According to the previous

researches, *JUN* was a positive regulator of T cell activation[69, 70] and can promote T cell proliferation

and the production of cytokines[71]. It is reported that one of the reasons of T cell exhaustion is the

continuous stimulation after the initial cell activation[45, 72], and the upregulation of *JUN* both in the

pre_exhausted and exhausted cells showed the lasting role to regulate cell activation, which might lead

to T cell exhaustion finally. The ENCODE database showed that JUN is a transcription factor of

PDCD1 and *TIGIT*, thus we speculated that *JUN* may affect T cell exhaustion through transcriptional

regulation of the exhaustion markers. However, the mechanisms of T cell exhaustion regulated by *JUN*

needed further study. In addition, *SAMSN1* is an immunoinhibitory adaptor and the knockdown of

SAMSN1 can enhance the adaptive immunity in mice[73]. In the comparison of the common DEGs

between pre_exhausted or exhausted and effector cells, *SAMSN1* showed greater upregulation in the

exhausted cells and was associated with poor prognosis in HCC and NSCLC based on TCGA data. The

expression of *SAMSN1* increased in B cells once stimulated by IL-4 through both PI3k/PKC/NF-κB

and STAT6 pathway[74, 75]. *SAMSN1* might be associated with T cell exhaustion according to our results,

however, further study is needed to explore the exact relationship between *SAMSN1* and T cell

exhaustion.

By integrating above previous researches and our results, we summarized the key genes related to

T cell exhaustion in Figure 4, including the upregulated genes of *CD69*, *PDCD1*, *HAVCR2*, *TIGIT*,

TOX and *JUN,* as well as a downregulated gene *S1PR1*. The more specific mechanisms needed to be

explored in the future study.

In the further classification of exhausted cluster to explore the heterogeneity, we found that each

patient tended to have one exhausted sub-cluster. Although the sub-clusters of each cancer may be

regulated by different hub genes, we found that these sub-clusters were dysregulated in the

cytokine-cytokine receptor interaction. Different kinds of cytokine played various and complex effects

on T cells, for instance, the decreased expression of effector cytokines such as IL-2 and TNF-α as well

as the increased expression of inhibitory cytokines such as IL-10 and TGF-β were reported to be

associated with T cell exhaustion[76]. Additionally, the upregulation of pathways in cancer was

consistent with the immune evasion of tumor caused by T cell exhaustion[77]. What's more, there were

two downregulated pathways including natural killer cell mediated cytotoxicity and ribosome. There is

a reciprocal relationship between NK and T cells, and most NK cell receptors are found on T cells[78].

Besides, there are similarities of NK cell and CD8+ T cell on the mechanisms of cell killing. They can

release cytotoxic granules to the target cell such as tumor cells, and induce programmed cell death[79].

The dysfunctional cytotoxicity might attenuate the killing effector of CD8+ T cells on tumor cells.

Additionally, the pathway of ribosome was also downregulated. Ribosome were essential for

production in all living cells and regulated the cell growth and proliferation[80, 81]. During T cell

exhaustion, the cells were characterized by the dysfunctional activity and bioenergetic deficiencies[82],

which might be related to the damaged functions of ribosome. In addition to the universal pathways,

there existed different pathways owing to each sub-cluster. In the CRC_S1 pathways, wnt signaling

pathway was upregulated in our study. It has been reported that wnt signaling pathway contributes to

tumor progression[83] and its blockade can improve the generation of effector cells[84], thus the

upregulation of wnt signaling pathway might contribute to T cell exhaustion and facilitate the growth

of tumor; In the NSCLC_S1 pathways, the upregulated apoptosis may lead to the impaired function of

T cells[85].

In the current study, we integrated three cancers and found some key genes associated with T cell

exhaustion, including the known markers *PDCD1*, *HAVCR2*, *TIGIT*, *TOX*. In the gene interaction

network, we found significant genes *CD69* and *JUN*. Besides, we firstly identified a potential

exhaustion associated gene, *SAMSN1*, which enriched in the negative regulation of immunity and

showed poor survival. We summarized the putative interrelated mechanisms of above key genes

identified in this study by integrating the previous knowledge. Furthermore, we explored the

heterogeneous and preference of exhausted CD8+ T cells in each patient and found only one exhausted

sub-cluster existed in most of patients, especially in CRC and HCC. Our study provides a universal

molecular mechanism on T cell exhaustion and a new way for functional research of single-cell RNA

sequencing data, which might promote the further realization of immunotherapy.

Acknowledgements

This work was supported by the National Key R&D Program of China (2018YFC0910200), the Key

R&D Program of Guangdong Province (2019B020226001), and the Science and the Technology

Planning Project of Guangzhou (201704020176).

References

[1]   MOSKOPHIDIS D, LECHNER F, PIRCHER H, et al. Virus persistence in acutely infected

immunocompetent mice by exhaustion of antiviral cytotoxic effector T cells [J]. Nature, 1993,

362(6434): 758-61.

[2]   CATAKOVIC K, KLIESER E, NEUREITER D, et al. T cell exhaustion: from

pathophysiological basics to tumor immunotherapy [J]. Cell Communication & Signaling Ccs,

2017, 15(1): 1.

[3]   LEE P P, YEE C, SAVAGE P A, et al. Characterization of circulating T cells specific for

tumor-associated antigens in melanoma patients [J]. Nat Med, 1999, 5(6): 677-85.

[4]   PAUKEN K E, WHERRY E J. Overcoming T cell exhaustion in infection and cancer [J].

Trends Immunol, 2015, 36(4): 265-76.

[5]   SCHIETINGER A, PHILIP M, KRISNAWAN V E, et al. Tumor-Specific T Cell Dysfunction

Is a Dynamic Antigen-Driven Differentiation Program Initiated Early during Tumorigenesis [J].

Immunity, 2016, 45(2): 389-401.

[6]   PAUKEN K E, WHERRY E J. Overcoming T cell exhaustion in infection and cancer [J].

Trends in Immunology, 2015, 36(4): 265-76.

[7]   BUCKS C M, NORTON J A, BOESTEANU A C, et al. Chronic antigen stimulation alone is

sufficient to drive CD8+ T cell exhaustion [J]. J Immunol, 2009, 182(11): 6697-708.

[8]   BLACKBURN S D, SHIN H, HAINING W N, et al. Coregulation of CD8+ T cell exhaustion

by multiple inhibitory receptors during chronic viral infection [J]. Nature immunology, 2009,

10(1): 29.

[9]   BAITSCH L, BAUMGAERTNER P, DEVEVRE E, et al. Exhaustion of tumor-specific

CD8(+) T cells in metastases from melanoma patients [J]. J Clin Invest, 2011, 121(6):

2350-60.

[10] MCLANE L M, ABDEL-HAKEEM M S, WHERRY E J. CD8 T Cell Exhaustion During

Chronic Viral Infection and Cancer [J]. Annual Review of Immunology, 2019, 37(1): 457-95.

[11] SEN D R, KAMINSKI J, BARNITZ R A, et al. The epigenetic landscape of T cell

exhaustion [J]. Science, 2016, 354(6316): 1165-9.

[12] CRAWFORD A, ANGELOSANTO J M, KAO C, et al. Molecular and transcriptional basis

of CD4$^+$ T cell dysfunction during chronic infection [J]. Immunity, 2014, 40(2): 289-302.

[13] WAGNER A, REGEV A, YOSEF N. Revealing the vectors of cellular identity with

single-cell genomics [J]. Nature Biotechnology, 2016, 34(11): 1145.

[14] WINTERHOFF B J, MAILE M, MITRA A K, et al. Single cell sequencing reveals

heterogeneity within ovarian cancer epithelium and cancer associated stromal cells [J].

Gynecol Oncol, 2017, 144(3): 598-606.

[15] WU H, YU J, LI Y, et al. Single-cell RNA sequencing reveals diverse intratumoral

heterogeneities and gene signatures of two types of esophageal cancers [J]. Cancer Lett,

2018, 438(133-43.

[16] PAPALEXI E, SATIJA R. Single-cell RNA sequencing to explore immune cell

heterogeneity [J]. Nature Reviews Immunology, 2017, 18(1): 35.

[17] AZIZI E, CARR A J, PLITAS G, et al. Single-Cell Map of Diverse Immune Phenotypes in

the Breast Tumor Microenvironment [J]. Cell, 2018, 174(S0092867418307232-.

[18] DARMANIS S, SLOAN S A, CROOTE D, et al. Single-Cell RNA-Seq Analysis of

Infiltrating Neoplastic Cells at the Migrating Front of Human Glioblastoma [J]. Cell Reports,

2017, 21(5): 1399.

[19] PURAM S V, TIROSH I, PARIKH A S, et al. Single-Cell Transcriptomic Analysis of

Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer [J]. Cell, 2017, 171(7):

1611-24 e24.

[20] TIROSH I, IZAR B, PRAKADAN S M, et al. Dissecting the multicellular ecosystem of

metastatic melanoma by single-cell RNA-seq [J]. Science, 2016, 352(6282): 189-96.

[21] HUANG A C, POSTOW M A, ORLOWSKI R J, et al. T-cell invigoration to tumour burden

ratio associated with anti-PD-1 response [J]. Nature, 2017, 545(7652): 60-5.

[22] VENTEICHER A S, TIROSH I, HEBERT C, et al. Decoupling genetics, lineages, and

microenvironment in IDH-mutant gliomas by single-cell RNA-seq [J]. Science, 2017,

355(6332):

[23] KISELEV V Y, KIRSCHNER K, SCHAUB M T, et al. SC3: consensus clustering of

single-cell RNA-seq data [J]. Nature Methods, 2017, 14(483-6.

[24] ZHANG X, LAN Y, XU J, et al. CellMarker: a manually curated resource of cell markers in

human and mouse [J]. Nucleic Acids Research, 2018, 47(D1): D721-D8.

[25] TRAPNELL C, CACCHIARELLI D, GRIMSBY J, et al. The dynamics and regulators of cell

fate decisions are revealed by pseudotemporal ordering of single cells [J]. Nat Biotechnol,

2014, 32(4): 381-6.

[26] ROBINSON M D, MCCARTHY D J, SMYTH G K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data [J]. Bioinformatics, 2010, 26(1):

139-40.

[27] YU G, WANG L G, HAN Y, et al. clusterProfiler: an R package for comparing biological

themes among gene clusters [J]. OMICS, 2012, 16(5): 284-7.

[28] SHANNON P, MARKIEL A, OZIER O, et al. Cytoscape: a software environment for

integrated models of biomolecular interaction networks [J]. Genome Res, 2003, 13(11):

2498-504.

[29] VON MERING C, HUYNEN M, JAEGGI D, et al. STRING: a database of predicted

functional associations between proteins [J]. Nucleic Acids Res, 2003, 31(1): 258-61.

[30] BADER G D, HOGUE C W. An automated method for finding molecular complexes in

large protein interaction networks [J]. BMC Bioinformatics, 2003, 4(2.

[31] LACHMANN A, XU H, KRISHNAN J, et al. ChEA: transcription factor regulation inferred

from integrating genome-wide ChIP-X experiments [J]. Bioinformatics, 2010, 26(19): 2438-44.

[32] CONSORTIUM E P. The ENCODE (ENCyclopedia Of DNA Elements) Project [J].

Science, 2004, 306(5696): 636-40.

[33] ROUILLARD A D, GUNDERSEN G W, FERNANDEZ N F, et al. The harmonizome: a

collection of processed datasets gathered to serve and mine knowledge about genes and

proteins [J]. Database (Oxford), 2016, 2016(

[34] HAN H, CHO J W, LEE S, et al. TRRUST v2: an expanded reference database of human

and mouse transcriptional regulatory interactions [J]. Nucleic Acids Res, 2018, 46(D1):

D380-D6.

[35] TOMCZAK K, CZERWINSKA P, WIZNEROWICZ M. The Cancer Genome Atlas (TCGA):

an immeasurable source of knowledge [J]. Contemp Oncol (Pozn), 2015, 19(1A): A68-77.

[36] ZHENG C, ZHENG L, YOO J K, et al. Landscape of Infiltrating T Cells in Liver Cancer

Revealed by Single-Cell Sequencing [J]. Cell, 2017, 169(7): 1342-56 e16.

[37] HANZELMANN S, CASTELO R, GUINNEY J. GSVA: gene set variation analysis for

microarray and RNA-seq data [J]. BMC Bioinformatics, 2013, 14(7.

[38] LIBERZON A, BIRGER C, THORVALDSDOTTIR H, et al. The Molecular Signatures

Database (MSigDB) hallmark gene set collection [J]. Cell Syst, 2015, 1(6): 417-25.

[39] RITCHIE M E, PHIPSON B, WU D, et al. limma powers differential expression analyses

for RNA-sequencing and microarray studies [J]. Nucleic Acids Res, 2015, 43(7): e47.

[40] LI R, QIAN J, WANG Y Y, et al. Long noncoding RNA profiles reveal three molecular

subtypes in glioma [J]. CNS Neurosci Ther, 2014, 20(4): 339-43.

[41] XIYING F, ALEXANDER.Y R. Hallmarks of Tissue-Resident Lymphocytes [J]. Cell, 2016,

164(1198-211.

[42] MEHTA A K, GRACIAS D T, CROFT M. TNF activity and T cells [J]. Cytokine, 2018,

101(14-8.

[43] HARLIN H, MENG Y, PETERSON A C, et al. Chemokine expression in melanoma

metastases associated with CD8+ T-cell recruitment [J]. Cancer Res, 2009, 69(7): 3077-85.

[44] SUKUMAR M, LIU J, JI Y, et al. Inhibiting glycolytic metabolism enhances CD8+ T cell

memory and antitumor function [J]. J Clin Invest, 2013, 123(10): 4479-88.

[45] WHERRY E J K, M. Molecular and cellular insights into T cell exhaustion [J]. Nat Rev

Immunol, 2015, 15(486–99.

[46] CONWAY J R, LEX A, GEHLENBORG N. UpSetR: an R package for the visualization of

intersecting sets and their properties [J]. Bioinformatics, 2017, 33(18): 2938-40.

[47] KALLIES A, ZEHN D, UTZSCHNEIDER D T. Precursor exhausted T cells: key to

successful immunotherapy? [J]. Nat Rev Immunol, 2019,

[48] POLLEN A A, NOWAKOWSKI T J, SHUGA J, et al. Low-coverage single-cell mRNA

sequencing reveals cellular heterogeneity and activated signaling pathways in developing

cerebral cortex [J]. Nat Biotechnol, 2014, 32(10): 1053-8.

[49] WHERRY E J. T cell exhaustion [J]. Nat Immunol, 2011, 12(6): 492-9.

[50] BARBER D L, WHERRY E J, MASOPUST D, et al. Restoring function in exhausted CD8

T cells during chronic viral infection [J]. Nature, 2006, 439(7077): 682-7.

[51] CHEMNITZ J M, PARRY R V, NICHOLS K E, et al. SHP-1 and SHP-2 associate with

immunoreceptor tyrosine-based switch motif of programmed death 1 upon primary human T

cell stimulation, but only receptor ligation prevents T cell activation [J]. J Immunol, 2004,

173(2): 945-54.

[52] RIELLA L V, PATERSON A M, SHARPE A H, et al. Role of the PD-1 pathway in the

immune response [J]. American journal of transplantation : official journal of the American

Society of Transplantation and the American Society of Transplant Surgeons, 2012, 12(10):

2575-87.

[53] PATSOUKIS N, BROWN J, PETKOVA V, et al. Selective effects of PD-1 on Akt and Ras

pathways regulate molecular components of the cell cycle and inhibit T cell proliferation [J]. Sci

Signal, 2012, 5(230): ra46.

[54] KOGUCHI K, ANDERSON D E, YANG L, et al. Dysregulated T cell expression of TIM3 in

multiple sclerosis [J]. J Exp Med, 2006, 203(6): 1413-8.

[55] DAS M, ZHU C, KUCHROO V K. Tim-3 and its role in regulating anti-tumor immunity [J].

Immunol Rev, 2017, 276(1): 97-111.

[56] NGIOW S F, VON SCHEIDT B, AKIBA H, et al. Anti-TIM3 antibody promotes T cell

IFN-gamma-mediated antitumor immunity and suppresses established tumors [J]. Cancer Res,

2011, 71(10): 3540-51.

[57] FOURCADE J, SUN Z, BENALLAOUA M, et al. Upregulation of Tim-3 and PD-1

expression is associated with tumor antigen-specific CD8+ T cell dysfunction in melanoma

patients [J]. J Exp Med, 2010, 207(10): 2175-86.

[58] JOLLER N, KUCHROO V K. Tim-3, Lag-3, and TIGIT [J]. Curr Top Microbiol Immunol,

2017, 410(127-56.

[59] KONG Y, ZHU L, SCHELL T D, et al. T-Cell Immunoglobulin and ITIM Domain (TIGIT)

Associates with CD8+ T-Cell Exhaustion and Poor Clinical Outcome in AML Patients [J]. Clin

Cancer Res, 2016, 22(12): 3057-66.

[60] SCOTT A C, DUNDAR F, ZUMBO P, et al. TOX is a critical regulator of tumour-specific T

cell differentiation [J]. Nature, 2019, 571(7764): 270-4.

[61] KHAN O, GILES J R, MCDONALD S, et al. TOX transcriptionally and epigenetically

programs CD8(+) T cell exhaustion [J]. Nature, 2019, 571(7764): 211-8.

[62] ALFEI F, KANEV K, HOFMANN M, et al. TOX reinforces the phenotype and longevity of

exhausted T cells in chronic viral infection [J]. Nature, 2019, 571(7764): 265-9.

[63] ZIEGLER S F, RAMSDELL F, HJERRILD K A, et al. Molecular characterization of the

early activation antigen CD69: a type II membrane glycoprotein related to a family of natural

killer cell activation antigens [J]. Eur J Immunol, 1993, 23(7): 1643-8.

[64] ESPLUGUES E, SANCHO D, VEGA-RAMOS J, et al. Enhanced antitumor immunity in

mice deficient in CD69 [J]. The Journal of experimental medicine, 2003, 197(9): 1093-106.

[65] CYSTER J G, SCHWAB S R. Sphingosine-1-phosphate and lymphocyte egress from

lymphoid organs [J]. Annu Rev Immunol, 2012, 30(69-94.

[66] KIMURA M Y, HAYASHIZAKI K, TOKOYODA K, et al. Crucial role for CD69 in allergic

inflammatory responses: CD69-Myl9 system in the pathogenesis of airway inflammation [J].

Immunol Rev, 2017, 278(1): 87-100.

[67] MITA Y, KIMURA M Y, HAYASHIZAKI K, et al. Crucial role of CD69 in anti-tumor

immunity through regulating the exhaustion of tumor-infiltrating T cells [J]. Int Immunol, 2018,

30(12): 559-67.

[68] ARBOUR, N. c-Jun NH2-Terminal Kinase (JNK)1 and JNK2 Signaling Pathways Have

Divergent Roles in CD8+ T Cell-mediated Antiviral Immunity [J]. Journal of Experimental

Medicine, 195(7): 801-10.

[69] BEIQING L, CHEN M, WHISLER R L. Sublethal levels of oxidative stress stimulate

transcriptional activation of c-jun and suppress IL-2 promoter activation in Jurkat T cells [J]. J

Immunol, 1996, 157(1): 160-9.

[70] QIAO X, PHAM D N T, LUO H, et al. Ran Overexpression Leads to Diminished T Cell

Responses and Selectively Modulates Nuclear Levels of c-Jun and c-Fos [J]. Journal of

Biological Chemistry, 285(8): 5488-96.

[71] UCHIDA M, SAWA H. [The SWI/SNF chromatin-remodeling complex regulates

asymmetric cell division in C. elegans] [J]. Tanpakushitsu Kakusan Koso Protein Nucleic Acid

Enzyme, 2005, 50(6 Suppl): 569-74.

[72] MOGNOL G P, SPREAFICO R, WONG V, et al. Exhaustion-associated regulatory

regions in CD8(+) tumor-infiltrating T cells [J]. Proc Natl Acad Sci U S A, 2017, 114(13):

E2776-E85.

[73] WANG D, STEWART A K, ZHUANG L, et al. Enhanced adaptive immunity in mice lacking

the immunoinhibitory adaptor Hacs1 [J]. FASEB J, 2010, 24(3): 947-56.

[74] YAN Y, ZHANG L, XU T, et al. SAMSN1 is highly expressed and associated with a poor

survival in glioblastoma multiforme [J]. PLoS One, 2013, 8(11): e81905.

[75] ZHU Y X, BENN S, LI Z H, et al. The SH3-SAM adaptor HACS1 is up-regulated in B cell

activation signaling cascades [J]. J Exp Med, 2004, 200(6): 737-47.

[76] HE Q F, XU Y, LI J, et al. CD8+ T-cell exhaustion in cancer: mechanisms and new area

for cancer immunotherapy [J]. Briefings in functional genomics, 2019, 18(2): 99-106.

[77] CATAKOVIC K, KLIESER E, NEUREITER D, et al. T cell exhaustion: from

pathophysiological basics to tumor immunotherapy [J]. Cell Commun Signal, 2017, 15(1): 1.

[78] VIVIER E, NUNES J A, VELY F. Natural killer cell signaling pathways [J]. Science, 2004,

306(5701): 1517-9.

[79] GALVIN J P, SPAENY-DEKKING L H, WANG B, et al. Apoptosis induced by granzyme

B-glycosaminoglycan complexes: implications for granule-mediated apoptosis in vivo [J].

Journal of immunology (Baltimore, Md : 1950), 1999, 162(9): 5345-50.

[80] CATEZ F, DALLA VENEZIA N, MARCEL V, et al. Ribosome biogenesis: An emerging

druggable pathway for cancer therapeutics [J]. Biochem Pharmacol, 2019, 159(74-81.

[81] TAN T C J, KNIGHT J, SBARRATO T, et al. Suboptimal T-cell receptor signaling

compromises protein translation, ribosome biogenesis, and proliferation of mouse CD8 T cells

[J]. Proc Natl Acad Sci U S A, 2017, 114(30): E6117-E26.

[82] WHERRY E J, HA S J, KAECH S M, et al. Molecular signature of CD8+ T cell exhaustion

during chronic viral infection [J]. Immunity, 2007, 27(4): 670-84.

[83] POLAKIS P. Wnt signaling in cancer [J]. Cold Spring Harbor perspectives in biology, 2012,

4(5):

[84] SCHINZARI V, TIMPERI E, PECORA G, et al. Wnt3a/beta-Catenin Signaling Conditions

Differentiation of Partially Exhausted T-effector Cells in Human Cancers [J]. Cancer Immunol

Res, 2018, 6(8): 941-52.

[85] BARATHAN M, GOPAL K, MOHAMED R, et al. Chronic hepatitis C virus infection triggers

spontaneous differential expression of biosignatures associated with T cell exhaustion and

apoptosis signaling in peripheral blood mononucleocytes [J]. Apoptosis, 2015, 20(4): 466-80.

Figure legends

Figure 1. Clustering of tumor-infiltrating CD8+ T cells.

A-C. Cell clustering using the selected genes of the top n ones with highest variance in CRC (A,

n=2500), HCC (B, n=1000), NSCLC (C, n=1000). The colors of the upper bar highlight the different

clusters.

D-F. The trajectory analysis of CD8+ T cell (excluding conventional cells) state transition inferred by

Monocle in CRC (D), HCC (E), NSCLC (F). Each dot represents an individual cell, colored according

to the clusters.

G-I. t-SNE plots of CD8+ T cells show distinct clusters predominantly determined by cell types in

CRC (G), HCC (H), NSCLC (I). Each dot represents an individual cell, colored according to the

clusters.

J. t-SNE plots of the total CD8+ T cells of three cancers show the uniformity according to the cell types,

regardless of the type of different cancers.

Figure 2. The Differentially expressed genes associated with exhausted state.

A. The fractions of five clusters of CD8+ T cells across peripheral blood (P), adjacent normal (N), and

tumor tissues (T). Bars were colored according to the clusters.

B-C. Venn diagram of DEGs in pre_exhausted (B) or exhausted (C) vs. effector comparison.

D-E. GO and KEGG enrichment analysis of DEGs in pre_exhausted vs. effector comparison. The top

15 significantly enriched GO terms (D) and KEGG pathways (E).

F-G. GO and KEGG enrichment analysis of DEGs in exhausted vs. effector comparison. The top 15

significantly enriched GO terms (F) and KEGG pathways (G).

Figure 3. The key DEGs associated with exhausted state.

A. The DEGs with higher fold change in the exhausted cells. The number represents the logFC value.

B. The survival analysis of *SAMSN1* based on TCGA data. Red corresponds to the group of high

expression, blue corresponds to the group of low expression.

C. The PPI network module with the highest score. Colors correspond to state of DEGs (red: up, blue:

down), shapes correspond to the types of DEGs (triangle: the specific genes existed in only

pre_exhausted cells, ellipse: the common genes existed in both exhausted and pre_exhausted cells,

diamond: the specific genes existed in only exhausted cells).

D. The number of DEGs regulated by their corresponding TFs. Colors correspond to state of TFs (red:

up, blue: down), shapes correspond to the types (ellipse: the dysregulated TFs, diamond: the number

DEGs regulated by corresponding TFs).

Figure 4. The putative interrelated mechanisms of key genes related to CD8+ T cell exhaustion.

A. CD69, which can combine with S1PR1 and inhibit the function of S1PR1 to regulate cell retention,

expressed on tumor-infiltrating T cells and may interact with unknown ligands in the tumor cell to lead

to the persistent antigen stimulation.

B-D. The mechanism of genes encoding inhibitory receptor related to CD8+ T cell exhaustion. The

upregulation of all three inhibitory receptors, *PDCD1, HAVCR2* and *TIGIT,* would result in the

inhibition of PI3K/Akt signaling pathway and lead to reduced survival and proliferation.

E-F. The transcription factor related to CD8+ T cell exhaustion. (E)*TOX* can promote the expression of

inhibitory receptors including *PDCD1*, *HAVCR2*, *TIGIT* to mediate T cell exhaustion program. (F) *JUN*

is proposed to regulate the expression of exhaustion associated *PDCD1* and *TIGIT* to play vital role in

T cell exhaustion at present study.

The red fonts represent upregulated expression, while the blue fonts represent downregulated

expression.

Figure 5. Clustering of exhausted cells.

A-C. Cell clustering using the selected genes of the top n ones with highest variance in the exhausted

cluster of CRC (A, n=1500), HCC (B, n=3000), NSCLC (C, n=500). The colors of the upper bar

highlight the different sub-clusters.

D-F. t-SNE plots of the exhausted sub-clusters show heterogeneous and preference in each patient of

CRC (D), HCC (E) and NSCLC (F). Each dot represents an individual cell, colored according to the

sub-clusters, gray means the non-exhausted cells.

Figure 6. The DEGs of exhausted sub-clusters.

A. Heatmap of six sub-clusters across three cancers. The colors of the upper bar highlight the different

sub-clusters.

B. Upset plot of DEGs between exhausted sub-clusters and effector cells. The numbers of exhausted

sub-clusters are shown on the left of the bar. The red line linked with dots show the common genes

across six sub-clusters.

Figure 7. The differential pathways between exhausted sub-clusters and effector cells by GSVA with

more than 10 DEGs and FDR < 0.05.

Supplements

Table S1. The number of markers in each cluster of each cancer and corresponding genes in
CellMarker database.
667

Table S2. DEGs of pre_exhausted or exhausted vs. effector comparison.

Table S3. GO and KEGG pathways in pre_exhausted or exhausted vs. effector comparison.

Table S4. DEGs of exhausted sub-clusters vs. effector comparison.

Table S5. GSVA pathway analysis results in exhausted sub-clusters vs. effector comparison.

Figure S1. The distribution of effector cells among peripheral blood, normal and tumor tissues in CRC

(A), HCC (B), NSCLC (C).

Figure S2. Violin plots of the key genes in pre_exhausted or exhausted vs. effector comparison.

A. The expression of key genes in the effector, pre_exhausted and exhausted cells of different cancers.

The black dot represents the median of expression value of genes ($\log_2$(TPM/10+1)).

B-D. The expression of key genes in each patient of CRC (B), HCC (C) and NSCLC (D).

Figure S3. The PPI network of DEGs in pre_exhausted or exhausted vs. effector comparison. The top

network consists of overall DEGs, and the remaining six networks are the module identified by plugin

MCODE of Cytoscape. Colors correspond to state of DEGs (red: up, blue: down), shapes correspond to

the types of DEGs (triangle: the specific genes existed in only pre_exhausted cells, ellipse: the common

genes existed in both exhausted and pre_exhausted cells, diamond: the specific genes existed in only

exhausted cells).

Figure S4. The network of DEGs and their corresponding TFs. Colors correspond to state of TFs (red:

up, blue: down), shapes correspond to the types (ellipse: the dysregulated TFs, diamond: the target

genes of the TFs).

A

B