H3K27me3 natural variation selectively marks genes predicted to be important for
differentiation in unicellular algae

Xue Zhao[1,3], Achal Rastogi[1†], Anne Flore Deton Cabanillas[1], Ouardia Ait Mohamed[1], Catherine
Cantrel[1], Berangère Lombard[2], Omer Murik[1¥], Auguste Genovesio[1], Chris Bowler[1], Daniel
Bouyer[1], Damarys Loew[2], Xin Lin[1 §], Alaguraj Veluchamy[4], Fabio Rocha Jimenez Vieira[1] and
Leila Tirichine[1,3*]

8
[1]Institut de Biologie de l'ENS (IBENS), Département de biologie, École normale supérieure,
CNRS, INSERM, Université PSL, 75005 Paris, France

[2]Institut Curie, PSL Research University, Centre de Recherche, Laboratoire de Spectrométrie
de Masse Protéomique, 26 rue d'Ulm 75248 Cedex 05 Paris, France

[3]CNRS UMR6286, UFIP UFR Sciences et Techniques, Université de Nantes, 2 rue de la
Houssinière 44322, Nantes Cedex 03

[4]Laboratory of Chromatin Biochemistry, BESE Division Building 2, Level 3, Office B2-3327,
4700 King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900,
Kingdom of Saudi Arabia

[†]Present address: Corteva Agriscience, Madhapur, Hyderabad 500 081, Telangana, India

[¥]Present address: Medical Genetics Institute, Shaare Zedek Medical Center, Jerusalem, Israel

[§]Present address: State Key Laboratory of Marine Environmental Science, Xiamen University

25
26

Abstract

In multicellular organisms H3K27me3 has been shown to be deposited by Polycomb Repressive Complex 2 (PRC2) to establish and maintain gene silencing, critical for cell fate and developmentally regulated processes. PRC2 complex is absent in both yeasts *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe,* which initially suggested that PRC2 arose with the emergence of multicellularity. However, its discovery in several unicellular species including microalgae questions its role in unicellular eukaryotes. Here, we show in the model diatom *Phaeodactylum tricornutum* (Pt), using mutants in the homologue of the catalytic subunit of PRC2, *enhancer of zeste E(z)*, that Pt *E(z)* is responsible for di and tri-methylation of lysine 27 of histone H3. H3K27me3 depletion abolishes cell morphology in Pt providing evidence for a role of H3K27me3 in cell differentiation in unicellular species. Genome wide profiling of H3K27me3 in fusiform and triradiate cells further revealed genes that may specify cell identity. These results suggest a role for PRC2 and its associated histone mark in cell differentiation in unicellular species and highlights their ancestral function in a broader evolutionary context than is currently appreciated.

Introduction

Tri-methylation of lysine 27 of histone H3 (H3K27me3) is a mark deposited by Polycomb Repressive Complex 2 (PRC2), which mediates silencing of gene expression during differentiation and development in both animals and plants [1-3]. PRC2 is comprised of four core proteins, highly conserved among multicellular organisms: the histone methyltransferase (HMTase) enhancer of zeste *E(z)*, the WD40 domain containing polypeptide Extra Sex Comb *Esc*, the C2H2 type zinc finger protein Suppressor of zeste 12 *Su(z)12* and the Nucleosome remodeling factor 55 kDa subunit *Nurf-55* [4,5]. The absence of PRC2 in the unicellular yeast models *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* initially led to suggestions that it arose to regulate cell differentiation in multicellular organisms [6]. This hypothesis has recently been questioned because components of PRC2 and the associated mark H3K27me3 are found in several unicellular species that belong to different lineages including, but not only, Stramenopiles, Alveolates and Rhizaria (SAR)[7-9] (Supplementary Fig. 1a-d), thus questioning the function of such a widespread complex in single celled organisms.

Attempts to understand the role of H3K27me3 in the unicellular green alga *Chlamydomonas reinhardtii* were not conclusive because tri-methylation of lysine 27 could not be assessed reliably due to its nominal mass which was found similar to acetylation of lysine 27 of the same histone[7]. However, we have previously identified H3K27me3 by mass spectrometry and mapped its localization genome-wide in the pennate diatom *Phaeodactylum tricornutum*[9] (Pt), which belongs to the stramenopile group of eukaryotes, only distantly related to the animal (Opisthokonta) and plant (Archaeplastida) eukaryotic crown groups. Pt has different morphotypes, fusiform (FM hereafter), which is the most prevailing morphology among the sampled accessions known so far, triradiate (TM hereafter), oval (OM hereafter) and cruciform (CM hereafter) [10-13] (Fig. 1a). Each morphotype can switch reversibly into a different morphology in response to several growth and environmental cues[10]. FM is the most stable morphotype while switching is more prominent in TM, CM and OM, which tend to convert to FM in the growth conditions used in this study[10,11].

Interestingly, western blot analysis using a monoclonal antibody against the mark in FM, TM, CM and OM cells revealed a strong correlation between the complexity of the morphology (branching of the cell) and the absolute quantity of H3K27me3, which is higher in both CM and TM cells compared to FM and OM cells (Supplementary Fig. 1e), suggesting that PRC2 activity controls cell differentiation in *P. tricornutum*. Cell differentiation is often orchestrated by H3K27me3-mediated silencing that underlies the establishment and maintenance of cellular

identity in multicellular model species [14]. These results prompted us to investigate the putative

role of H3K27me3 in cell differentiation in a unicellular model. In silico annotation of

polycomb complex members and identification of H3K27me3 by mass spectrometry[15] in a well-

developed experimental model such as *P. tricornutum* (Supplementary Fig. 1a-c) might give

new insights for the unsolved questions in the study of PRC2 complex in multicellular species,

especially the availability of different morphotypes which is unique to *P. tricornutum* among

stramenopiles, present an opportunity to decipher its role in single celled organisms with respect

to its potential contribution to establish morphotype switches as well as its function in an

evolutionary context.

To gain insights into the function of *E(z)* and its associated H3K27me3 mark in Pt, we generated

two CRISPR/cas9 knockouts of the gene in each of the three morphotypes (FM, TM and CM)

leading to putative loss of function mutations including premature stop codon and frameshifts

(Fig. 1b). Light microscopy analysis of *E(z)* knockouts shows a change in cell morphology

which becomes shorter in the FM background. Both triradiate and cruciform morphologies were

abolished in TM and CM, respectively while transgenic lines carrying cas9 control vectors in

each morphotype remain unchanged (Fig. 1c). Whereas *P. tricornutum* transgenic lines with

the Cas9 control vector show similar H3K27me3 enrichment than the wild type (Supplementary

Fig. 2a). knockout of *E(z)* led to an overall depletion of H3K27me3 shown by western blot

using a monoclonal antibody against the mark (Fig. 1d), suggesting that the loss or diminution

of *E(z)* activity, and hence H3K27me3, causes the observed changes in cell morphology (Fig.

1c; Supplementary Fig. 2a; Table S1). Overall, this suggests that *E(z)* and its associated mark

are required for morphotype switch/to establish specific cell identity.

Mass spectrometry analysis of histones extracted from both wild type and *E(z)* knockout

confirmed the loss of H3K27me3 and revealed a depletion of di-methylation of H3K27,

corroborating the role of *E(z)* in di and tri-methylation of lysine 27 of histone H3

(Supplementary Fig. 2b,c,d). This is similar to fungi [16] and mammals[17] but different from *A.*

thaliana where PRC2 loss of function leads to specific depletion of H3K27me3[18], although in

vitro assays with reconstituted *A. thaliana* PRC2 components showed mono, di and tri-

methylation of lysine 27 of histone H3[19,20]. Western blot analysis of *E(z)* knockout mutants

using a monoclonal antibody against H3K27me2 (Supplementary Fig. 2e) confirmed the

depletion of the mark from the mutants, supporting further the mass spectrometry analysis and

the role of *E(z)* in di-methylation of lysine 27 of histone H3 in *P. tricornutum*.

Similar to animals, plants and *Neurospora crassa*, H3K27me3 appears not to be essential for

cell survival in *P. tricornutum*, as indicated by the overall growth of *E(z)* knockout lines, which

are only slightly retarded compared to wild type lines (Supplementary Fig. 2f).

To further investigate the role of H3K27me3 and its targets in different morphotypes, we carried

out a genomic approach and performed Chromatin Immuno-Precipitation (ChIP) on two

biological replicates of TM morphotypes using an antibody against H3K27me3 followed by

DNA sequencing (ChIP-Seq) to generate a map of H3K27me3 distribution, which we compared

to the one previously generated in FM[21]. ChIP-Seq data analysis revealed a similar H3K27me3

enrichment profile between TM and FM that localizes principally on transposable elements

(TEs), with 58% and 60% of the reads overlapping with TE annotations for FM[21] and TM,

respectively (Fig. 2a). The mark was found to occupy, on average, ~11.6% of the genome in

FM cells, targeting approximately 15% of genes (consistent with [15]) and ~13.2% of the genome

within TM, targeting 19% of genes in agreement with the absolute amount of H3K27me3 to be

elevated in TM compared to FM as detected by western blot (Supplementary Fig. 1e). Indeed,

more genes are marked by H3K27me3 in TM than in FM (Fig. 2a, Supplementary Fig. 3a),

although most of the PRC2 targets are shared between both morphotypes (Fig. 2a) and exhibit

globally broad coverage over the annotation (Supplementary Fig. 3b). Among the PRC2 target

genes, 635 and 297 genes are found to be specifically marked by H3K27me3 in TM and FM,

respectively (Fig. 2a). We used ChIP followed by quantitative PCR (ChIP-qPCR) to validate

H3K27me3 enrichment over specifically marked loci in both backgrounds, which corroborated

the genome wide data for most of the tested genes (Fig. 2b, c). Common marked loci and

unmarked loci were tested as internal controls (Supplementary Fig. 3c). To test for the loss of

H3K27me3 in *E(z)* knockouts, we performed ChIP-qPCR in FM and TM as well as the

respective mutants, which confirmed the depletion of H3K27me3 in these mutants (Fig. 1e,f).

To gain insights into the functional categories enriched in H3K27me3 target genes that are

shared between the two morphotype or specific, we applied to the updated Phatr3 annotation a

gene ontology analysis using DAMA[22] and CLADE[23] which is a machine learning methodology

that uses pHMMs, positive score matrix and support vector machines to infer the corresponding

most probable GO category to genes. DAMA and Clade allow a more sensitive remote

homology that permits to assign to genes with no or poor domain conservation the

corresponding GO categories that would have been missed by other methods. Are considered

statistically significant, only GO classes that are represented by at least 3 standard deviations

above the average of observed entries.

Out of 1,640 H3K27me3 marked genes, 753 could not be assigned to a more specific GO
category and are therefore marked as unknown. The genes that are marked in both morphotypes
show top enrichment in RNA related biological processes such as RNA-dependent DNA
biosynthetic process, RNA phosphodiester bond hydrolysis and RNA-DNA hybrid
ribonuclease activity. Genes marked by H3K27me3 specifically in TM displayed top
enrichment exclusively in (1) glycoprotein biosynthetic processes involved in the transfer of
sugar moieties that might determine different sugar composition of the cell wall which is known
to be sugar rich in *P. tricornutum*[24], (2) Peptidyl tyrosine dephosphorylation processes with
Ankyrin repeats proteins known to act as scaffold for connecting molecular interactions, likely
important for development of the numerous signaling pathways associated generally to more
complex multicellular organisms[25] (Fig. 2d; Supplementary Table S2). Genes that are
specifically marked in FM cells exhibit enrichment in categories such as peptidyl-tyrosine
phosphorylation containing genes with central roles as modulators of cell differentiation and
cell fate decisions[26] (Fig. 2d; Supplementary Table S2). Interestingly, additional genes
specifically marked in each FM or TM share categories with predicted functions in positive
regulation of (1) GTPase activity with a role in cell morphology changes, and neurite outgrowth
and guidance[27] as well as the differentiation of many cell types, including neurons, T
lymphocytes and myocytes[28]; (2) protein ubiquitination shown to play a role in the complex
regulation of the levels and function of many proteins and signaling pathways involved in
determining cell fate[29]. Overall, the genes that are specifically marked in the TM or FM
morphotypes reflect processes related to cell growth, proliferation and differentiation.

We have reported previously that H3K27me3 marked genes in FM are characterized by low
expression[15], consistent with the role of H3K27me3 as a repressive mark. Interestingly, when
genes are marked by H3K27me3 in both FM and TM, their expression is lower compared to
the genes that are uniquely marked in FM and to a lesser extent in TM (Fig. 3a,b). This suggests
that specifically marked genes are kept under less stringent and tight repression which might be
due to their putative role in morphotype switch, which is known to be a dynamic process[10].

Considering the conserved role of H3K27me3 in repression, we tested the effect of *E(z)*
knockout on gene expression. Therefore, RNA sequencing (RNA-seq) of two biological
replicates of the *E(z)* mutant (Del6) was carried out and compared to previously generated
RNA-seq in the wild type (FM). Around 1/4 of all genes are (23%, 2795 out of 12152
annotations) differentially expressed in the *E(z)* mutant (P-value. < 0.05), (Supplementary Fig.
3d; Supplementary Table S2), indicating an essential role in gene regulation by PRC2 in Pt.

We further monitored by RT-qPCR the expression of 27 specifically marked genes in the *E(z)*

knockout of the TM and found that 18 genes out of 27 showed a gain of expression in the mutant

compared to the TM background, demonstrating further that depletion of H3K27me3 likely

releases the repression of target genes and correlates with the loss of the triradiate morphology

(Fig. 3c). Although the remaining genes showed no change or even a gain in expression, these

genes can be targets of other repressive or active marks as shown previously with co-occurrence

of several repressive marks over genic regions[9]. The analysis of the R value[30] which reflects

the entropy, and therefore the variability in expression of genes between FM, TM and loci

marked in both morphotypes, showed a higher value in specifically marked genes compared to

commonly marked ones (Fig. 3d). This supports further the finding that specific enrichment in

each of the morphotypes are less silenced and potentially more dynamic compared to genes

marked by H3K27me3 in both TM and FM cells whereas commonly H3K27me3 marked genes

are globally silenced.

To substantiate the assumption that phenotypic plasticity and morphotype switch are regulated

by PRC2 in *P. tricornutum*, we took advantage of the lack of stability of the TM phenotype and

its tendency to switch to FM. Specifically, we used clonal cell samples with FM and TM

morphologies from the same genetic background (TM), which switches habitually to fusiform

and therefore contains a mixture of FM and TM cells. We reasoned that the activity of *E(z)*

should correlate with H3K27me3 levels in the following way: (1) a pure triradiate population

isolated from TM-N (named here TM-T1): highest level of H3K27me3), (2) a population of

cells from TM after N generations (N is 60 ± 5) of cell division containing a mixture of triradiate

and fusiform morphotypes (TM-N): medium level of H3K27me3) and (3) fusiform cells

isolated from the triradiate background TM (TM-Fusi): lowest level of H3K27me3 (Fig. 4a).

E(z) transcript levels show a clear decrease in TM-N and TM-Fusi compared to TM-T1 (Fig.

4b) which correlates with the switch from TM to FM, reflecting a lower activity of E(z) and

H3K27me3 levels. We then asked whether specifically H3K27me3-marked loci in TM lose the

mark upon cell switching to FM after multiple generations of sub-culturing leading to TM-N

and in transformed fusiform cells (TM-Fusi). As expected, ChIP-qPCR showed clearly a loss

of the mark in a population containing a mixture of fusiform and triradiate cells (TM-N) as well

as in TM-Fusi compared to TM-T1 (Fig. 4c,d), which contains only triradiate cells, thus

correlating the morphology with the level of enrichment in H3K27me3 over specific genes.

In summary, we have demonstrated in this study the role in *P. tricornutum* of *E(z)* as a histone

methyltransferase responsible for di and tri-methylation of lysine 27 of histone H3. Knockout

of *E(z)* causes H3K27me3 depletion and loss of triradiate cell shape maintenance, providing

evidence for the involvement of *E(z)* and its associated mark in establishing and/or maintaining

cell morphology in unicellular species. We showed the dynamic nature of the mark, depending

on the specific morphology between and within *P. tricornutum* accessions that correlate with

the level of H3K27me3 enrichment. We showed differential marking in two different

accessions of *P. tricornutum*, FM versus TM, which identified genes related to cell fate

decisions compared to commonly marked genes. This is the first evidence of the involvement

of H3K27me3 in cell differentiation in unicellular eukaryotes only distantly related to animals

and plants. Our study points to the emerging function of PRC2 and its H3K27me3 associated

mark as a determinant of the establishment and maintenance of cell morphology in single celled

species such as *P. tricornutum* that shows signs of differentiation of the cell into diverse

morphologies. This same function likely diversified with the emergence of multicellularity with

PRC2 orchestrating development in plants and animals.

Methods

Strains and growth conditions

Phaeodactylum tricornutum Bohlin Clone Pt1 8.6 (CCMP2561) (referred as FM) and Clone

Pt8Tc (referred as TM) cells were grown as described previously [31].

Isolation and immunoprecipitation of chromatin

Chromatin isolation and immunoprecipitation were performed as described previously [32]. The

following antibodies were used for immunoprecipitation: H3K27me3 (07-449) from Millipore

and H3K27me3 from cell signaling technology. qPCR on recovered DNA was performed as

described previously [32]

CRISPR/Cas9 plasmid construction

hCAS9n (Cas9 from *Streptococcus pyogenes*, adapted to human codon usage, fused to SV40

nuclear localization sequence, and containing a D10A mutation) was amplified from

pcDNA3.3-TOPO-hCAS9n (kindly received from Dr. Yonatan B. Tzur, Hebrew University of

Jerusalem), using the primers 5'-CAC CAT GGA CAA GAA GTA CTC-3' and 5'- TCA CAC

CTT CCT CTT CTT CTT-3'. The PCR product was first cloned into pENTR using pENTR/D-

TOPO cloning kit (ThermoFisher Scientific), and then sub-cloned into a PT pDest, containing

an N-terminal HA-tag [31], following the manufacturer's protocol, which was named pDest-HA-
hCAS9n.

For the sgRNA vector we first cloned the snRNA U6 promoter[33] from *P. tricornutum* genomic
DNA using the primers 5'- AAA CGA CGG CCA GTG AAT TCT CGT TTC TGC TGT CAT
CAC C-3' and 5'- TCT TTA ATT TCA GAA AAT TCC GAC TTT GAA GGT GTT TTT TG-
3'. PU6::unc-119_sgRNA (kindly received from Dr. Yonatan B. Tzur) backbone was amplified
using the primers 5'-CAA AAA ACA CCT TCA AGT CGG AAT TTT CTG AAA TTT AAA
GA-3' and 5'-GGT GAT GAC AGC AGA AAC GAG AAT TCA CTG GCC GTC GTT T-3'.
The two PCR products were used as template for a second round fusion PCR reaction as
described in [34]. We further transformed the resulting product into *E. coli*, and extracted the
ligated plasmid. The terminator sequence of the P. tricornutum U6 was amplified using the
primers   5'-CATTCTAGAAGAACCGCTCACCCATGC-3'   and   5'-
GTTAAGCTTGAAAAGTTCGTCGAGACCATG-3', digested by XbaI/HindIII and ligated
into XbaI/HindIII digested pU6::unc-119. The resulting vector, ptU6::unc-119-sgRNA, was
used as template to replace the target sequence to *E(Z)* target by PCR using primers
32817TS12fwd GTG TCG GAG CCC GCC ATA CCG TTT TAG AGC TAG AAA TAG C
and 32817TS12rev GGT ATG GCG GGC TCC GAC ACC GAC TTT GAA GGT GTT TTT
TG. Target sequences were picked using PhytoCRISP-Ex [35].

Transformation of *P. tricornutum* cells and screening for mutants

Wild type cells of the reference strain FM and the TM were transformed with three plasmids
(pPhat1, Cas9 and guide RNA with the target sequence) as described previously [36]. Positive
transformants were validated by triple PCR screen for pPhaT1 shble primers (ACT GCG
TGCACTTCGTGGC/TCGGTCAGTCCTGCTCCTC),   sgRNA
(GAGCTGGAAATTGGTTGTC/GACTCGGTGCCACTTTTTCAAGTT)   and   CAS9n
(GGGAGCAGGCAGAAAACATT/TCACACCTTCCTCTTCTTCTT). For each colony, a
rapid DNA preparation was performed as described previously and fragment of 400 bp was
amplified with primers flanking the target sequence in the *E(z)* gene. The forward primer used
is   5'-TAAGATGGAGTATGCCGAAATTC-3'   and   reverse   primer   is   5'-
AGGCATTTATTCGTGTCTGTTCG-3' PCR product was run in 1% agarose gel and a single
band was extracted using Machery Nagle kit and according to the manual manufacturer. PCR
product was sequenced using the primer 5'-AGCCACCCTGCGTTAACTGAAAAT-3'.

To make sure that the fusiform cells originating from the switch of TM are not contaminants

from the FM, each of the TM-T1, TM-Fusi and TM-N were checked for their genetic

background whether it is FM or TM using a molecular marker designed around a 400 bp

insertion in the FM background (Supplementary Fig. 4g), identified from genome sequencing

of FM and TM strains of *P. tricornutum*[37]. The PCR check confirmed that all the cell samples

described above are in the TM genetic background.

Validation of enrichment and expression of target genes

276 **qPCR:** Total RNA was extracted from TM and FM cells as described previously [31] and cDNA

277 was synthesized with cDNA high yield synthesis kit according to the manufacturer user manual.

278 Quantitative PCR was performed as described previously [31] using the primer list in

279 Supplementary Table S3

Proteomics and PRM Measurements

Three independent histone purifications recovered from FM wild type cells as well as *E(z)*

knockout mutant Del6 in FM genetic background were simultaneously separated by SDS-

PAGE and stained with colloidal blue (LabSafe Gel Blue GBiosciences). Three gel slices were

excised for each purification and in-gel digested by using trypsin/LysC (Promega). Peptide

extracted from each set were pooled and analyzed by nanoLC-MS/MS using an Ultimate 3000

system (Thermo Scientific) coupled to a TripleTOFTM 6600 mass spectrometer (AB Sciex).

Peptides were first trapped onto a C18 column (75 μm inner diameter × 2 cm; nanoViper

Acclaim PepMapTM 100, Thermo Scientific) with buffer A (2/98 MeCN/H2O in 0.1% formic

acid) at a flow rate of 2.5 μL/min over 4 min. Separation was performed on a 50 cm x 75 μm

C18 column (nanoViper C18, 3 μm, 100Å, Acclaim PepMapTM RSLC, Thermo Scientific)

regulated to 50°C and with a linear gradient from 1% to 30% buffet B (100 MeCN in 0.085%

formic acid) at a flow rate of 400 nL/min over 90 min. The mass spectrometer was operated in

PRM top30 high sensitivity mode with 100 ms acquisition time for MS1 and MS2 scans

respectively with included precursor mass list for 600 sec (see Supplementary Table S4)

PRM Data Analysis

The PRM data were analyzed using Skyline version 3.7.0.11317 MacCoss Lab Software,

Seattle, WA; https://skyline.ms/project/home/software/Skyline/begin.view, fragment ions for

each targeted mass were extracted and peak areas were integrated. The peptide areas were log2

transformed and the mean log2- area was normalized by the mean area of peptide STDLLIR

using software R version 3.1.0. On each peptide a linear model was used to estimate the mean

fold change between the conditions, its 97.5% confidence interval and the p-value of the two

sided associated t-test. The p-values were adjusted with the Benjamini-Hochberg procedure[38].

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE [1] partner repository with the dataset identifier PXD012347.

Western blot analysis: Chromatin was extracted from wild type as well as mutants of both TM

and FM cells and western blot performed as described previously [32].

Sequencing and computational data analysis

ChIP-Seq: Chromatin Immunoprecipitation (ChIP) was done with monoclonal cell cultures

grown using single triradiate cell from Pt8 population (referred as TM). CHIP-Seq was

performed as described previously [15,39]. Two replicates were performed and showed a good

Pearson correlation (Supplementary Fig. 5). Raw reads were filtered and low quality read-pairs

were discarded using FASTQC with a read quality (Phred score) cutoff of 30. Using the genome

assembly published in 2008 as reference (Pt1 8.6), we performed reference-assisted mapping

of filtered reads using BOWTIE. We then performed the processing and filtering of the

alignments using SAMTOOLS and BEDTOOLS. SICER [40] was then used to identify

significant enriched H3K27me3 peaks by comparing it with the INPUT. Differential

H3K27me3 peak enrichment analysis between FM and TM backgrounds was also done using

SICER-df plugin. Peaks with Padj < 0.05 differential enrichment or depletion were considered

significant. Functional inferences were obtained by overlapping the differentially enriched

peaks over structural annotations from Phatr3 genome annotation[41].

RNA sequencing (RNA-Seq): Total RNA was extracted from FM, TM, and FM Ez-KO (Del6)

cell lines. RNA expression and differential gene expression analysis was performed using

Eoulsan version 1.2.2 with default parameters [42]. Genes having at least 2 folds expression

change with P-value < 0.05 were considered as significant different expressed genes (DEGs).

Whole genome sequencing (WGS): Whole genome sequencing was performed using DNA

extracted from monoclonal cell cultures grown using single triradiate cell taken from Pt8 and

Pt1 accession, referred to as Pt8tc and Pt1 8.6, respectively. At least 6 µg of genomic DNA

from each accession was used to construct a sequencing library following the manufacturer's

instructions (Illumina Inc.). Paired-end sequencing libraries with a read size of 100 bp and an insert size of approximately 400 bp were sequenced on an Illumina HiSeq 2000 sequencer at Berry Genomics Company (China) and Fasteris for Pt1 8.6 and Pt8tc, respectively. Low quality read-pairs were discarded using FASTQC with a read quality (Phred score) cutoff of 30. Using the genome assembly published previously[43], we performed reference-assisted assembly of all the accessions. We used BOWTIE (-n 2 –X 400) for mapping the high quality NGS reads to the reference genome followed by the processing and filtering of the alignments using SAMTOOLS and BEDTOOLS. For estimating the genetic diversity between Pt1 8.6 and Pt8Tc genome, GATK [44] configured for diploid genomes, was used for variant calling, which included single nucleotide polymorphisms (SNPs), small insertions and deletions ranging between 1 and 300 base pairs (bp). The genotyping mode was kept default (genotyping mode = DISCOVERY), Emission confidence threshold (-stand_emit_conf) was kept 10 and calling confidence threshold (-stand_call_conf) was kept at 30. The minimum number of reads per base to be called as a high quality SNP was kept at 4 (i.e., read-depth >=4x). SNPEFF was used to annotate the functional nature of the polymorphisms.

GO enrichment analysis: GO categories were grouped by 3 different levels of expression according to a simple density clustering algorithm (also confirmed by iterative k-means clustering).

Data Availability:

All data are available through NCBI Sequence Read Archive with accession number PRJNA565539.

References

1   Aldiri, I. & Vetter, M. L. PRC2 during vertebrate organogenesis: a complex in transition. *Dev Biol* **367**, 91-99, doi:10.1016/j.ydbio.2012.04.030 (2012).

2   Fragola, G. *et al.* Cell reprogramming requires silencing of a core subset of polycomb targets. *PLoS Genet* **9**, e1003292, doi:10.1371/journal.pgen.1003292 (2013).

3   Surface, L. E., Thornton, S. R. & Boyer, L. A. Polycomb group proteins set the stage for early lineage commitment. *Cell Stem Cell* **7**, 288-298, doi:10.1016/j.stem.2010.08.004 (2010).

4   Schwartz, Y. B. & Pirrotta, V. A new world of Polycombs: unexpected partnerships and emerging functions. *Nature reviews. Genetics* **14**, 853-864, doi:10.1038/nrg3603 (2013).

5   Martinez-Balbas, M. A., Tsukiyama, T., Gdula, D. & Wu, C. Drosophila NURF-55, a WD repeat protein involved in histone metabolism. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 132-137 (1998).

6   Kohler, C. & Villar, C. B. Programming of gene expression by Polycomb group proteins. *Trends*
Cell Biol **18**, 236-243, doi:10.1016/j.tcb.2008.02.005 (2008).
7   Shaver, S., Casas-Mollano, J. A., Cerny, R. L. & Cerutti, H. Origin of the polycomb repressive
complex 2 and gene silencing by an E(z) homolog in the unicellular alga Chlamydomonas.
Epigenetics **5**, 301-312 (2010).
8   Mikulski, P., Komarynets, O., Fachinelli, F., Weber, A. P. M. & Schubert, D. Characterization of
the Polycomb-Group Mark H3K27me3 in Unicellular Algae. *Front Plant Sci* **8**, 607,
doi:10.3389/fpls.2017.00607 (2017).
9   Veluchamy, A. *et al.* An integrative analysis of post-translational histone modifications in the
marine diatom Phaeodactylum tricornutum. *Genome biology* **16**, 102, doi:10.1186/s13059-
015-0671-8 (2015).
10   De Martino, A. *et al.* Physiological and Molecular Evidence that Environmental Changes Elicit
Morphological Interconversion in the Model Diatom Phaeodactylum tricornutum. *Protist*
162, 462-481, doi:S1434-4610(11)00006-X [pii]

10.1016/j.protis.2011.02.002 (2011).
11   De Martino, A. M., A. Juan Shi, K.P. Bowler, C. Genetic and phenotypic characterization of
Phaeodactylum tricornutum (Bacillariophyceae) accessions. *J. Phycol.* **43**, 992–1009 (2007).
12   He, L., Han, X. & Yu, Z. A rare Phaeodactylum tricornutum cruciform morphotype: culture
conditions, transformation and unique fatty acid characteristics. *PLoS one* **9**, e93922,
doi:10.1371/journal.pone.0093922 (2014).
13   Borowitzka, M. A., Volcani, B.E. The polymorphic diatom Phaeodactylum tricornutum:
Ultrastructure of its morphotypes. *J. Phycolo.* **14**, 10-21 (1978).
14   Margueron, R. & Reinberg, D. The Polycomb complex PRC2 and its mark in life. *Nature* **469**,
343-349, doi:10.1038/nature09784 (2011).
15   Veluchamy, A. *et al.* An integrative analysis of post-translational histone modifications in the
marine diatom Phaeodactylum tricornutum. *Genome biology* **16**, 102, doi:10.1186/s13059-
015-0671-8 (2015).
16   Jamieson, K., Rountree, M. R., Lewis, Z. A., Stajich, J. E. & Selker, E. U. Regional control of
histone H3 lysine 27 methylation in Neurospora. *Proceedings of the National Academy of*
Sciences of the United States of America **110**, 6027-6032, doi:10.1073/pnas.1303750110
(2013).
17   Ferrari, K. J. *et al.* Polycomb-dependent H3K27me1 and H3K27me2 regulate active
transcription and enhancer fidelity. *Molecular cell* **53**, 49-62,
doi:10.1016/j.molcel.2013.10.030 (2014).
18   Lafos, M. *et al.* Dynamic regulation of H3K27 trimethylation during Arabidopsis
differentiation. *PLoS Genet* **7**, e1002040, doi:10.1371/journal.pgen.1002040 (2011).
19   Jacob, Y. *et al.* ATXR5 and ATXR6 are H3K27 monomethyltransferases required for chromatin
structure and gene silencing. *Nat Struct Mol Biol* **16**, 763-768, doi:10.1038/nsmb.1611
(2009).
20   Schmitges, F. W. *et al.* Histone methylation by PRC2 is inhibited by active chromatin marks.
Molecular cell **42**, 330-341, doi:10.1016/j.molcel.2011.03.025 (2011).
21   Veluchamy, A. *et al.* An integrative analysis of post-translational histone modifications in the
marine diatom Phaeodactylum tricornutum. *Genome Biol* **16**, 102 (2015).
22   Bernardes, J. S., Vieira, F. R., Zaverucha, G. & Carbone, A. A multi-objective optimization
approach accurately resolves protein domain architectures. *Bioinformatics* **32**, 345-353,
doi:10.1093/bioinformatics/btv582 (2016).
23   Bernardes, J., Zaverucha, G., Vaquero, C. & Carbone, A. Improvement in Protein Domain
Identification Is Reached by Breaking Consensus, with the Agreement of Many Profiles and
Domain Co-occurrence. *PLoS Comput Biol* **12**, e1005038, doi:10.1371/journal.pcbi.1005038
(2016).

24   Le Costaouëc, T., Unamunzaga, C., Mantecon, L. and Helbert, W. New structural insights into
the cell-wall polysaccharide of the diatom Phaeodactylum tricornutum. *Algal research* **26**
172-179 (2017).
25   Marcotte, E. M., Pellegrini, M., Yeates, T. O. & Eisenberg, D. A census of protein repeats. *J*
Mol Biol **293**, 151-160, doi:10.1006/jmbi.1999.3136 (1999).
26   Yu, J. S. & Cui, W. Proliferation, survival and metabolism: the role of PI3K/AKT/mTOR
signalling in pluripotency and cell fate determination. *Development* **143**, 3050-3060,
doi:10.1242/dev.137075 (2016).
27   Etienne-Manneville, S. & Hall, A. Rho GTPases in cell biology. *Nature* **420**, 629-635,
doi:10.1038/nature01148 (2002).
28   Bryan, B. A., Li, D., Wu, X. & Liu, M. The Rho family of small GTPases: crucial regulators of
skeletal myogenesis. *Cell Mol Life Sci* **62**, 1547-1555, doi:10.1007/s00018-005-5029-z (2005).
29   Thompson, S. J., Loftus, L. T., Ashley, M. D. & Meller, R. Ubiquitin-proteasome system as a
modulator of cell fate. *Curr Opin Pharmacol* **8**, 90-95, doi:10.1016/j.coph.2007.09.010 (2008).
30   Maheswari, U. *et al.* Digital expression profiling of novel diatom transcripts provides insight
into their biological functions. *Genome biology* **11**, R85, doi:gb-2010-11-8-r85 [pii]

10.1186/gb-2010-11-8-r85 (2010).
31   Siaut, M. *et al.* Molecular toolbox for studying diatom biology in Phaeodactylum tricornutum.
Gene **406**, 23-35, doi:S0378-1119(07)00275-2 [pii]

10.1016/j.gene.2007.05.022 (2007).
32   Lin, X., Tirichine, L. & Bowler, C. Protocol: Chromatin immunoprecipitation (ChIP)
methodology to investigate histone modifications in two model diatom species. *Plant*
methods **8**, 48, doi:10.1186/1746-4811-8-48 (2012).
33   Rogato, A. *et al.* The diversity of small non-coding RNAs in the diatom Phaeodactylum
tricornutum. *BMC genomics* **15**, 698, doi:10.1186/1471-2164-15-698 (2014).
34   Hobert, O. PCR fusion-based approach to create reporter gene constructs for expression
analysis in transgenic C. elegans. *Biotechniques* **32**, 728-730 (2002).
35   Rastogi, A., Murik, O., Bowler, C. & Tirichine, L. PhytoCRISP-Ex: a web-based and stand-alone
application to find specific target sequences for CRISPR/CAS editing. *BMC bioinformatics* **17**,
261, doi:10.1186/s12859-016-1143-1 (2016).
36   Falciatore, A., Casotti, R., Leblanc, C., Abrescia, C. & Bowler, C. Transformation of
nonselectable reporter genes in marine diatoms. *Mar Biotechnol (NY)* **1**, 239-251, doi:MBT30
[pii] (1999).
37   Rastogi, A. V. F., Deton-Cabanillas AF, Veluchamy A., Cantrel, C., Wang, G., Vanormelingen,
P., Bowler, C., Piganeau, G., Hu, H. and Leila Tirichine. A genomics approach reveals the
global genetic polymorphism, structure and functional diversity of ten accessions of the
marine model diatom Phaeodactylum tricornutum

ISME J (2019).
38   Benjamini, Y., and Yekutieli, D. . The control of the false discovery rate in multiple testing
under dependency. *Annals of Statistics* **29**, 1165–1188 (2001).
39   Veluchamy, A. *et al.* Insights into the role of DNA methylation in diatoms by genome-wide
profiling in Phaeodactylum tricornutum. *Nat Commun* **4**, doi:10.1038/ncomms3091 (2013).
40   Zang, C. *et al.* A clustering approach for identification of enriched domains from histone
modification ChIP-Seq data. *Bioinformatics* **25**, 1952-1958,
doi:10.1093/bioinformatics/btp340 (2009).
41   Rastogi, A. *et al.* Integrative analysis of large scale transcriptome data draws a
comprehensive landscape of Phaeodactylum tricornutum genome and evolutionary origin of
diatoms. *Sci Rep* **8**, 4834, doi:10.1038/s41598-018-23106-x (2018).

42   Jourdren, L., Bernard, M., Dillies, M. A. & Le Crom, S. Eoulsan: a cloud computing-based
framework facilitating high throughput sequencing analyses. *Bioinformatics* **28**, 1542-1543,
doi:10.1093/bioinformatics/bts165 (2012).
43   Bowler, C. *et al.* The Phaeodactylum genome reveals the evolutionary history of diatom
genomes. *Nature* **456**, 239-244, doi:nature07410 [pii]

10.1038/nature07410 (2008).
44   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res* **20**, 1297-1303, doi:10.1101/gr.107524.110
(2010).

Acknowledgements

Hanhua Hu from the Chinese Academy of Science is acknowledged for the gift of CM

morphotype.). LT acknowledges funds from the CNRS and the region of Pays de la Loire

(ConnecTalent EPIALG project). CB acknowledges funding from the ERC Advanced Award

'Diatomite. XZ was supported by a PhD fellowship from the Chinese Scholarship Council

(CSC-201604910722). AR was supported by an International PhD fellowship from the MEMO

LIFE Program.

Competing interests

The authors declare no competing interests.

 Figure legends

 Figure 1. *Phaeodactylum tricornutum* morphotypes and enhancer of zeste knockout

 mutants. (a) *P. tricornutum* morphotypes (top left fusiform, scale bar = 3µm, top right oval,

 bottom left triradiate, scale bar = 2µm, bottom right cruciform). Scale bars correspond to 1 µm

 in OM and CM. **(b)** Sequence chromatograms of PCR product from WT cells and CRISPR cas9

 mutants of enhancer of zeste showing the different indels in FM, TM and CM. **(c)** Light

 microscopy images of WT, E(z) KO and Cas9 control cells. Empty vector controls containing

 Cas9 and Shble antibiotic resistance gene show no loss of H3K27me3 and retain the wild type

 morphology suggesting that morphology distortion is not due to the transformation but to the

 absence of H3K27me3. **(d)** Western blots of WT and two E(z)KOs from each morphotype

 using a monoclonal antibody against H3K27me3. Histone H4 was used as a loading control.

 ChIP-QPCR enrichment levels of H3K27me3 on genes in WT and E(z) KO in FM **(e)** and TM

 (f) backgrounds.

 Figure 2. Genomic features of H3K27me3 targets in FM and TM cells. **(a)** Venn diagrams

 showing the number of common and specific genomic features [Genes, Transposable elements

 (TEs), and Intergenic Regions (IGRs)] targeted by H3K27me3 in TM (orange circles) and FM

 (blue circles). ChIP-QPCR validation of H3K27me3 specifically marked genes in **(b)** TM and

 (c) FM morphotypes. **(d)** Distribution of the most frequent GO terms on genes marked with

 H3K27me3. The distribution was sub-divided into different categories, where TM and FM

 represent the GOs observed exclusively on triradiate and fusiform genes, respectively. A third

 category (Both) also presents a GO distribution for genes observed on both morphotypes.

 Figure 3. Gene expression profiles of 3HK27me3 specific and common targets in FM and

 TM. **(a)** The box plot represents mean enrichment of H3K27me3 (Y-axis), with log10 scaling,

 over genes marked specifically in FM (blue), TM (orange), and also on genes marked in both

 (commonly marked) morphotypes. The enrichment profile is generated using number of genes

 marked by H3K27me3 specifically in each morphotype and also in both. The significant

 H3K27me3 enrichment difference between specifically marked and commonly marked is

 estimated using two-tailed t-test with P value < 0.0001, denoted by "****". **(b)** Expression of

genes marked specifically in FM (blue), TM (orange), and in both phenotypic backgrounds with same principle aesthetics and categorical genes, used in 2f. The significance/non-significance of the variability of expression between specifically and commonly marked genes is estimated using two-tailed t-test with P value = 0.0433, as denoted by "*". "ns" denote non-significant. (**c**) Relative expression level of H3K27me3 targets genes in the TM and enhancer of zeste knock out M2-11. In the plot, fold change log2(PtM2-11/Pt8Tc) values are shown (**d**) Boxplot showing the entropy value distribution of H3K27me3 marked genes. Entropy values measures the differential expression of genes under different experimental conditions. Entropy values are derived from expression data (fragments per kilobase of exon per million fragments mapped) under five different experimental conditions. Genes marked specifically by H3K27me3 in TM (Triradiate) shows higher variation in expression followed by FM (Fusiform) specific H3K27me3 marked genes. Horizontal lines represent the median entropy values. The significant of FM/TM-specific H3K27me3 marked genes and genes marked on both conditions are estimated using two-tailed t-test with P value < 0.0001.

Figure 4. H3K27me3 enrichment levels and morphotype switch. (**a**) Schematic diagram showing how TM-N, TM-T1 and TM-Fusi were generated. After generations of culture in lab growth condition (ASW media, 19°C, 12h/12h light dark period), some triradiate cells switch to fusiform forming a mixture of FM and TM cells, named TM-N. A single fusiform cell from TM-N was picked and propagated clonally giving rise to TM-Fusi. Similarly, single triradiate cell was isolated from TM-N and its clonal propagation gave a population of pure triradiate cells named TM-T1. (**b**) Relative expression level of Enhancer of Zest in FM, TM-N, TM-T1 and TM-Fusi respectively, two pairs of primer were designed at N-terminal of Enhancer of Zeste gene, and at the CXC domain respectively. (**c**) ChIP-QPCR enrichment levels of H3K27me3 in TM-T1 and TM-Fusi. (**d**) ChIP-QPCR showing enrichment levels of H3K27me3 in TM-T1 and TM-N.

Supplementary Table 1. Cell counts of different morphologies and cell size measurements in each wild type morphotype and knock out of E(z)

Supplementary Table 2. List of genes marked by H3K27me3 with their annotation and GOs.

Supplementary Table 3. List of the primers used in this study

Supplementary Table 4. Mass spectrometry quantification of Di- and Tri-methylation of H3K27me3

Supplementary Figure 1. Phylogeny **(a-c)**. **(d)** Western blots using a monoclonal antibody against H3K27me3 on protein or chromatin extracts of different species representative of the super SAR lineage (1: *Bigelowiella_natans,* 2: *Gymnophora dimorpha,* 3: *Skeletonema marinoi,* 4: *Thalassiosira pseudonana,* 5: *Raphoneis sp,* 6: *Synedra sp,* 7: *Asterionellopsisglacialis,* 8: *Thalassiosira rotula,* 9: *Phaeodactylum tricornutum,* 10: *Isocrhrysis lutea,* 11: *Amhedinium klebselii,* 12: *Amhedinium carteri).* **(e)** Western blot on chromatin extracts from OM, FM, TM and CM with a monoclonal antibody against H3K27me3 showing a significant difference in enrichment levels of H3K27me3 in TM and CM which are higher compared to FM and OM.

Supplementary Figure 2. **(a)** Western blot of chromatin extracts of Cas9 control lines from each of FM, TM and CM. H4 histone antibody was used as a loading control. **(b, c)** Mass spectrometry quantification of di and tri-methylation of lysine 27 of histone H3 in both wild type and enhancer of zeste knockout mutant. MS/MS spectrum of the $[M + 2H]^{2+}$ precursor ion of histone H3 (27–36 residue peptide) tri-methylated or di-methylated on K27. Broken bonds above and below sequence denote b and y ions, respectively, that were annotated from the spectrum. **(d)** Abundance of H3 K27 di- and tri-methylated KSAPATGGVK peptide. Y axis shows normalized log2 (WT/Mutant) of the di-methylated and tri-methylated peptides. All measurements have been performed in triplicate, and error bars indicated the 97.5% confidence interval (see supplementary Table 1). **(e)** Western blot of chromatin extracts from wild type FM and TM as well as E(z) knockouts in both backgrounds with a monoclonal antibody against H3K27me2. H4 histone antibody was used as loading control. **(f)** Growth curves of wild type, enhancer of zeste mutants and cas9 control line in each of FM, TM and CM. **(g)** Gel picture of a molecular marker distinguishing FM and TM and amplifying an insertion in one allele present in FM but absent in TM.

Supplementary Figure 3.

(a)Total genome coverage of H3K27me3 within TM (orange) and FM (blue) showing a higher

mapping of H3K27me3 in TM compared to FM. **(b)** mean distribution of H3K27me3 over 500

bp upstream, gene body, and 500 bp downstream region of all the gene targets in TM (orange

line) and FM (blue line). **(c)** Unmarked genes (J01910, J31617), commonly marked genes

(J34600, J44413) were chosen as internal controls. FM (EG00164) and TM specifically marked

(J49062) genes were used as controls for the reproducibility of independent ChIP-QPCR

results. (**d**) violin plot represents the mean fold change of gene expression in Ez(KO) lines in

TM compared to the wild-type (WT). The significant expression difference between Ez(KO)

and WT is estimated using two-tailed t-test with P value < 0.0001

Supplementary Figure 4.   Scatter plots with Pearson correlation coefficient displaying the

relationship between TM and E(z) knock out RNA Seq replicates.

Supplementary Figure 5.   Scatter plots with Pearson correlation coefficient displaying the

relationship between TM ChIP-Seq replicates

593
594

Supplementary Table 1. Cell counts of different morphologies and cell size measurements in

each wild type morphotype and knock out of E(z)

Supplementary Table 2. List of genes marked by H3K27me3 with their annotation and GOs.

Supplementary Table 3. List of the primers used in this study

Supplementary Table 4. Mass spectrometry quantification of Di- and Tri-methylation of

H3K27me3

Supplementary Figure 1. Phylogeny **(a-c)**. **(d)** Western blots using a monoclonal antibody

against H3K27me3 on protein or chromatin extracts of different species representative of the

super SAR lineage (1: *Bigelowiella_natans,* 2: *Gymnophora dimorpha,* 3: *Skeletonema*

marinoi, 4: *Thalassiosira pseudonana,* 5: *Raphoneis sp,* 6: *Synedra sp,* 7:

Asterionellopsisglacialis, 8: *Thalassiosira rotula,* 9: *Phaeodactylum tricornutum,* 10:

Isocrhrysis lutea, 11: *Amhedinium klebselii,* 12: *Amhedinium carteri).* **(e)** Western blot on

chromatin extracts from OM, FM, TM and CM with a monoclonal antibody against H3K27me3

showing a significant difference in enrichment levels of H3K27me3 in TM and CM which are

higher compared to FM and OM.

Supplementary Figure 2. **(a)** Western blot of chromatin extracts of Cas9 control lines from

each of FM, TM and CM. H4 histone antibody was used as a loading control. **(b, c)** Mass

spectrometry quantification of di and tri-methylation of lysine 27 of histone H3 in both wild

type and enhancer of zeste knockout mutant. MS/MS spectrum of the $[M + 2H]^{2+}$ precursor ion

of histone H3 (27–36 residue peptide) tri-methylated or di-methylated on K27. Broken bonds

above and below sequence denote b and y ions, respectively, that were annotated from the

spectrum. **(d)** Abundance of H3 K27 di- and tri-methylated KSAPATGGVK peptide. Y axis

shows normalized log2 (WT/Mutant) of the di-methylated and tri-methylated peptides. All

measurements have been performed in triplicate, and error bars indicated the 97.5% confidence

interval (see supplementary Table 1). **(e)** Western blot of chromatin extracts from wild type FM

and TM as well as E(z) knockouts in both backgrounds with a monoclonal antibody against

H3K27me2. H4 histone antibody was used as loading control. **(f)** Growth curves of wild type,

enhancer of zeste mutants and cas9 control line in each of FM, TM and CM. **(g)** Gel picture of

a molecular marker distinguishing FM and TM and amplifying an insertion in one allele present

in FM but absent in TM.

Supplementary Figure 3.

(a) Total genome coverage of H3K27me3 within TM (orange) and FM (blue) showing a higher

mapping of H3K27me3 in TM compared to FM. **(b)** mean distribution of H3K27me3 over 500

bp upstream, gene body, and 500 bp downstream region of all the gene targets in TM (orange

line) and FM (blue line). **(c)** Unmarked genes (J01910, J31617), commonly marked genes

(J34600, J44413) were chosen as internal controls. FM (EG00164) and TM specifically marked

(J49062) genes were used as controls for the reproducibility of independent ChIP-QPCR

results. **(d)** violin plot represents the mean fold change of gene expression in Ez(KO) lines in

TM compared to the wild-type (WT). The significant expression difference between Ez(KO)

and WT is estimated using two-tailed t-test with P value < 0.0001

Supplementary Figure 4.   Scatter plots with Pearson correlation coefficient displaying the

relationship between TM and E(z) knock out RNA Seq replicates.

Supplementary Figure 5.   Scatter plots with Pearson correlation coefficient displaying the

relationship between TM ChIP-Seq replicates

Figure 1

Figure 2

Figure 3

Figure 4

a

Figure S1

b

Figure S1

c

Amphiprora_sp
Amphora_coffeaeformis
Phaeodactylum_tricornutum
Ditylum_brightwellii
Extubocellulus_spinifer
Minutocellus_polymorphus
Odontella_sp
Cyclophora_tenuis
Thalassionema_nitzschioides
Asterionellopsis_glacialis
Stephanopyxis_turris
Proboscia_inermis
Proboscia_alata
Aulacoseira_subarctica
Kryptoperidinium_foliaceum
Cyclotella_meneghiniana
Detonula_confervacea
Thalassiosira_miniscula
Thalassiosira_weissflogii
Skeletonema_dohrnii
Nitzschia_sp.
Pseudo_nitzschia_delicatissima
Pseudo_nitzschia_australis
Nitzschia_punctata
Heterosigma_akashiwo
Pinguiococcus_pyrenoidosus
Fibrocapsa_japonica
Guillardia_theta
Hanusia_phi
Rhodomonas_sp
Chroomonas_mesostigmatica
Protocruzia_adherens
Pseudopedinella_elastica
Phaeomonas_parva
Isochrysis_galbana
Phaeocystis_sp
Eutreptiella_gymnastica
Percolomonas_cosmopolitus
Bolidomonas_pacifica
Cryptomonas_curvata
Vannella_sp
Tiarina_fusus
Phaeocystis_antarctica
Picochlorum_oklahomensis
Picochlorum_sp
Alexandrium_temarense
Bicosoecid_sp

Figure S1

d

	1	2	3	4	5	6	7	8	9	10	11	12

αH3K27me3

Rhizaria   Diatoms   Haptophyte   Alveolates

e

αH3K27me3

αH4

OM   FM   TM   CM

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Table 1. Cell counts of different morphologies and cell size measurements in each wild type morphotype and knock out of *E(z)*

	FM	TM	CM	OM	Majority	Percentage	Size
FM	107	0	0	0	FM	100%	26.58μm
TM	0	100	0	0	TM	100%	L:13.59μm; W:11.48μm
CM	4	31	66	0	CM	67.34%	-
OM	0	0	0	100	OM	100%	-
TM-N	63	55	0	0	FM	53.38%	-
T1	0	219	0	0	TM	100%	-
DEL6	74	0	0	0	FM	100%	19.91μm
TS12	41	0	0	0	FM	100%	-
M2-10	65	0	0	0	FM	100%	-
M2-11	74	0	0	0	FM	100%	21.04μm
M20	32	0	0	0	FM	100%	23.59μm

Table 2. Mass spectrometry quantification of Di- and Tri-methylation of H3K27me3

peptide sequence	Protein Modification (without Nterm Met)	peptide used for normalization	log2(WT/Mutant)	CI 2.5%	CI 97.5%	adjusted p value
KSAPATGGVK	Dimethyl (K:27)	STDLLIR	1.52038639	1.09023167	1.95054112	2.10E-07
KSAPATGGVK	Trimethyl (K:27)	STDLLIR	3.5091078	2.7625983	4.25561731	7.82E-10