Opponent regulation of action performance and timing by striatonigral and striatopallidal pathways

Authors:

Konstantin I. Bakhurin[1*], Xiaoran Li [1*], Alexander D. Friedman[1], Nicholas A. Lusk[1], Glenn D.R. Watson[1], Namsoo Kim[1], Henry H. Yin[1,2]

Author affiliations:

1. Department of Psychology and Neuroscience, Duke University, Durham, NC, 27708, USA.

2. Department of Neurobiology, Duke University School of Medicine, Durham, NC, 27708, USA.

* These authors contributed equally.

Corresponding author:

Henry H. Yin, Ph.D., hy43@duke.edu

Keywords:

Interval timing, basal ganglia, striatum, striatonigral, striatopallidal, licking, dopamine

Abstract

The basal ganglia have been implicated in action selection and timing, but the relative

contributions of the striatonigral (direct) and striatopallidal (indirect) pathways to these functions

remain unclear. This study investigated the effects of optogenetic stimulation of D1+ (direct) and

A2A+ (indirect) neurons in the ventrolateral striatum in head-fixed mice on a fixed time

reinforcement schedule. Direct pathway stimulation initiates licking. In contrast, indirect

pathway stimulation suppresses licking and results in rebound

licking following stimulation. Moreover, direct and indirect pathways also play distinct roles in

timing. Direct pathway stimulation produced a resetting of the internal timing

process, whereas indirect pathway stimulation transiently paused timing, and proportionally

delayed the next bout of licking. Our results provide evidence for the continuous and opposing

contributions of the direct and indirect pathways in the production and timing of reward-guided

behavior.

Introduction

The striatum is the major input nucleus of the basal ganglia (BG) and contains two major

populations of medium spiny projection neurons (Gerfen et al 1990, Mink 1996). Striatonigral

(direct pathway) neurons express D1 receptors and project to the substantia nigra pars reticulata

(SNr) and other BG output nuclei. Striatopallidal (indirect pathway) neurons express D2

receptors and project to the external globus pallidus, which in turn inhibits the SNr. Ever since

the connectivity of these pathways was defined, there has been controversy on their contributions

to behavior. Given their opposite effects on BG output, an influential model of the striatum casts

the direct and indirect pathways in opposing roles with the direct pathway promoting behavior

and the indirect pathway suppressing behavior (Albin et al 1989, Freeze et al 2013, Kravitz et al

2010). More recently, however, this view has been questioned by studies that described

concurrent activation of these two populations during behavior, suggesting complementary

contributions of these two pathways (Cui et al 2013, Isomura et al 2013).

The lateral or sensorimotor striatum receives topographically organized innervation from

the cortex and thalamus (Hintiryan et al 2016). Micro-stimulation of different striatal regions

results in movements of different body parts (Alexander & DeLong 1985). In rats, the

ventrolateral striatum (VLS) has been implicated in orofacial behaviors, including licking

(Mittler et al 1994, Pisa 1988a, Pisa 1988b). Although licking behavior is generated by brain

stem pattern generators (Travers et al 1997), BG output can provide top-down regulation of the

pattern generators, either directly or via the nigrotectal projections (Deniau & Chevalier 1992,

Redgrave et al 1992, Rossi et al 2016, Shammah-Lagnado et al 1992, Toda et al 2017).

Stimulation or enhancing dopaminergic signaling in the VLS can generate orofacial behavior, in

some cases dyskinetic movements, while dopamine depletion can abolish such behaviors (Delfs

& Kelley 1990, Jicha & Salamone 1991, Kelley et al 1988). However, the detailed pathway-

specific mechanisms underlying such observations remain largely unknown.

One important characteristic of self-initiated behaviors is that they can be precisely timed

relative to some anticipated event like a reward (Gibbon et al 1984, Pavlov 1927, Roberts 1981).

The BG have been implicated in interval timing in the seconds to minutes range (Buhusi & Meck

2005, Merchant et al 2013). Manipulation of dopamine activity can bidirectionally influence

estimation of the passage of time (Buhusi & Meck 2005, Meck 1996) and lesions of the striatum

also impair timing behavior (Meck 2006). Furthermore, striatal activity has also been shown to

represent time intervals (Bakhurin et al 2017, Gouvea et al 2015). Although the role of D1-

expressing and D2-expressing neurons in timing has been studied using systemic

pharmacological manipulations (Frederick & Allen 1996), it remains unclear how the two

pathways of the BG interact to mediate these behaviors.

In this study, we used optogenetic stimulation to examine the contributions of the VLS in

both action generation and timing. Using a fixed-time (FT) schedule of reinforcement, combined

with occasional peak probe trials, we showed that mice rapidly acquired highly predictable

licking patterns that reflects reliable internal timing processes around reward delivery (Toda et al

2017). Using temporally and spatially precise manipulations of neural activity with

optogenetics, we studied how direct and indirect pathways regulate the initiation, maintenance,

and timing of self-initiated licking.

Results

To study the roles of the direct and indirect pathways during self-initiated behavior, we

manipulated the activity of cell populations that originate these pathways in the orofacial region

of the striatum (Figures 1A & B). We bilaterally injected a Cre-dependent AAV5-DIO-ChR2

virus in the VLS region of the striatum in D1-Cre (Figure 1C) and A2A-Cre (Figure 1D) mice

and implanted chronic optic fibers just above this area. Five additional A2A-Cre mice received

AVV5-DIO-eYFP virus. We tracked licking behavior in head-fixed mice performing an interval

timing task (Toda et al 2017). Mice received a 5 μL drop of 10% sucrose solution every 10

seconds (Figure 1E). Voluntary licking behavior in all mice consisted bouts of rhythmic

protrusion and retractions of the tongue at a relatively fixed rate (4-8 Hz). Well-trained mice

reliably generated anticipatory licking behavior before reward delivery (Figure 1F). As a result,

the average licking rate gradually ramps up in expectation of reward, and then peaks following

reward and shows a sharper decline after consumption (Figure 1G).

We tested the effects of the direct pathway in the VLS on licking at three different time

points during the interval. We stimulated the direct pathway for 1 second before mice normally

initiate anticipatory licking, starting at 5 seconds before reward delivery (Figure 2A). Though

licking was sparse at this time point (Figure 2B, top), activation of direct pathway MSNs induced

licking (Figure 2B, bottom and Figure 2C, Video S1). We determined the change in lick rate that

resulted from laser stimulation by calculating the lick rate during the 1 second laser stimulation

period for D1-Cre mice and a control group that received eYFP virus. The normal lick rate

obtained for the same time point during trials without laser stimulation was subtracted from this

value to quantify the stimulation-induced change in lick rate (Figure 2D). A two-way ANOVA

(stimulation frequency x experimental group) revealed a change of licking rate resulting from

VLS direct pathway stimulation (main effect of group, $F = 56.4$, p < 0.0001) and a significant

effect of laser frequency ($F =8.38$, p < 0.01); there was a also an interaction between frequency

and group ($F = 8.76$, p < 0.01) . Post-hoc tests revealed that lick rate was significantly increased

by 10 Hz (p < 0.01), 25 Hz, (p < 0.001), and 50 Hz (p < 0.001) stimulation of the direct pathway.

We next stimulated the direct pathway at 1 second prior to reward delivery (Figure 2E), during

the time of maximal anticipatory licking (Figure 2F, top). Although there was a ceiling effect on

licking at this time point (Figure 2F, bottom), we regularly observed a potentiation of the licking

rate during stimulation when compared to trials without stimulation (Figures 2G). A two-way

ANOVA revealed significant increase in licking as a result of VLS stimulation (Figure 2H, main

effect of group, $F = 54.11$, p < 0.0001), but no effect of frequency ($F = 4.37$, p = 0.16) nor a

significant interaction between frequency and group ($F = 4.37$, p = 0.3). Stimulation at the time

of reward delivery (Figure 2I) had the most modest impact on licking rate (Figures 2J and 2K).

However, a two-way ANOVA detected a significant effect of stimulation on licking even during

consumption (Figure 2L, $F = 16.43$, p < 0.01), but did not show a significant effect of

stimulation frequency ($F = 7.23$, p = 0.3), nor a significant interaction ($F = 9.88$, p = 0.18). We

directly compared the effects of stimulation on licking as a function of time of laser stimulation

during the task. A two-way ANOVA revealed significant effects of both the time of stimulation

in the task (Figure 2M, main effect of time, $F = 17.65$, p < 0.0001) and laser frequency ($F =$

31.22, p < 0.0001), with no significant interaction ($F = 3.82$, p = 0.44). Together, these results

show for the first time that direct pathway stimulation in the orofacial striatum can result in the

generation of licking behavior. The effects of direct pathway stimulation were dependent on the

on-going behavioral state of the animals, which can be explained by a ceiling effect.

Remarkably, we observed that it was possible for licking to exceed the frequencies

normally produced by mice (Figure 3A). To explore this effect, we analyzed licking patterns in

the frequency domain. Licking during laser presentation could be shifted rightward toward a

frequency of 10 Hz (Video S2). This appeared to be the upper limit of the central pattern

generator for licking, as naturally generated licking showed a predominant frequency of 5-8 Hz.

We quantified power spectral density (PSD) distributions for licking and determined the licking

frequency that corresponded to the peak of the PSD during laser delivery and during naturally

generated licking without stimulation (Figure 3B). Mice displayed two distinct licking

frequency patterns during the task, depending on whether licking occurred prior to or following

reward. Laser stimulation further increased the licking frequency to around 10 Hz, which is

rarely reached naturally (Figure 3C). Indeed, we found that the peak power of licking was

potentiated: there was an interaction between the effects of stimulation frequency and licking

pattern on the peak of the PSD function (Figure 3D, two-way, mixed ANOVA, $F = 12.92$, p <

0.05; main effect of pattern, $F = 48.01$, p <0.0001; main effect of frequency, $F = 0.88$, p = 0.7).

Post-hoc tests showed that peaks of PSDs of laser-evoked licking were significantly greater than

those detected during anticipatory licking at 10 Hz (p < 0.001), 25 Hz (p < 0.001), and 50 Hz (p

< 0.001) stimulation. Peaks of PSDs of laser-evoked licking were also significantly greater than

licking following reward at 25 Hz (p < 0.001) and 50 Hz (p < 0.01) stimulation. While we found

a significant effect of licking pattern on licking frequency (Figure 3E, two-way mixed ANOVA,

$F = 31.62$, p < 0.0001), there was no effect of stimulation frequency on licking frequency ($F =$

8.31, p = 0.2), and no significant interaction between frequency and the pattern of licking ($F =$

4.82, p = 0.23).

Stimulation in the middle of the interval at 5 seconds also resulted in a significant

interaction between the effects of stimulation frequency and licking pattern on peak power

(Figure 3F, $F = 13.28$, p < 0.01). Although there was no main effect of frequency on peak power

($F = 6.61$, p < 0.2), we found a significant effect of licking pattern ($F = 25.9$, p < 0.0001). Post-

hoc tests revealed that the peak of the PSD of licking frequency during direct pathway

stimulation increased relative to licking following reward at 10 Hz (p < 0.01), 25 Hz (p < 0.001),

and 50 Hz (p < 0.01) stimulation. The peak of the PSD of laser-evoked licking frequency

increased relative to licking prior to reward at 10 Hz (p < 0.01) and 25 Hz (p < 0.001)

stimulation. We also observed a significant effect of stimulation on licking frequency (Figure

3G, $F = 43.22$, p < 0.0001), but no effect of frequency ($F = 2.19$, p = 0.6) and no significant

interaction between stimulation frequency and lick pattern ($F = 5.34$, p = 0.1). We even observed

a significant effect of licking pattern on the peak of the PSD during licking after reward while

reward was being consumed (Figure 3H, $F = 44.1$, p < 0.0001). While there was no significant

effect of stimulation frequency ($F = 6.31$, p = 0.09), we found a significant interaction between

the effects of licking pattern and stimulation frequency on peak PSD power during consumption

licking ($F = 9.76$, p < 0.05). Post-hoc tests revealed that there were significant differences

between peak PSDs of laser-evoked licking and anticipatory licking during stimulation at 10 Hz

(p<0.01), 25 Hz (p < 0.001), and 50 Hz (p < 0.001). Post-hoc tests also found significant

differences between peak PSDs of laser-evoked licking and consumption licking during 25 Hz

(p<0.01), and 50 Hz (p < 0.001) stimulation. In addition, there was an effect of licking pattern on

the licking frequency (Figure 3I, $F = 62.35$, p < 0.0001), but no significant effects of stimulation

frequency on licking frequency ($F = 0.19$, p = 0.98) and no significant interaction between

stimulation frequency and licking pattern ($F = 2.15$, p < 0.26). Interestingly, 5 Hz stimulation did

not reduce the peak licking frequency. Regardless of stimulation frequency, light delivery to the

VLS resulted in a positive shift in the frequency domain, above the naturally occurring states that

mice normally generate. In addition, direct pathway stimulation could increase the amount of

licking at a given frequency, reflected in the peak of the PSD function. Together, these results

suggest a role for the BG in regulating the duty cycle of the licking oscillator.

When stimulating the direct pathway at 5 seconds prior to reward, we observed that

licking resulting from stimulation only began after a certain delay (Figure 4A), and that this

delay was reduced with higher stimulation frequencies (Figure 4B). A one-way ANOVA

revealed a significant effect of stimulation frequency on the latency to initiate licking following

the onset of laser stimulation (Figure 4C; $F = 4.92$, p < 0.01). The lick bout that resulted from

stimulation also lasted longer than the 1 second stimulation period. We counted the number of

licks that occurred over the course of one second following the offset of stimulation and

compared that value with the number of licks normally detected during the same period in non-

stimulation trials. Stimulation would result in some additional licking even after offset of

stimulation, but this was not related to the stimulation frequency (Figure 4D; two-way mixed

ANOVA; main effect of stimulation, $F = 12.23$, p < 0.001; main effect of frequency, $F = 9.58$, p

= 0.30; interaction, $F = 3.88$, p = 0.15). These results show that greater stimulation of the direct

pathway reduces licking initiation latency, and that licking continues for a short time beyond the

immediate period of laser stimulation.

In contrast to D1 activation, indirect pathway activation in the VLS in well-trained mice

resulted in a pronounced suppression of licking. When stimulation occurred 5 seconds prior to

reward delivery (Figure 5A), we observed a reduction in licking rate (Figures 5B and 5C). We

found a significant effect of experimental group on the change in licking rate (Figure 5D, two-

way ANOVA, $F = 18.56$, $p < 0.01$), but no effect of frequency ($F = 9.18$, $p = 0.12$) nor an

interaction between group and frequency ($F = 9.93$, $p = 0.1$). Although we did regularly observe

a suppression of licking activity, the likelihood of licking at this period in the interval was low,

reflecting a potential floor effect. When activating the indirect pathway 1 second prior to reward

(Figure 5E), during peak anticipatory licking, we also observed a significant impact on licking by

indirect pathway stimulation (Figures 5F and 5G, Video S3). In addition to a strong suppression

of licking by indirect pathway stimulation (Figure 5H, two-way ANOVA; significant effect of

experimental group $F = 25.93$, $p < 0.0001$), we observed a significant effect of stimulation

frequency on licking ($F = 19.47$, $p < 0.0001$), and a significant interaction between group and

frequency ($F = 13.83$, $p < 0.01$). Post-hoc tests revealed significant suppression of licking by 25

Hz ($p < 0.001$) and 50 Hz ($p < 0.001$) stimulation. We were also able to suppress consummatory

licking that occurred following reward (Figures 5I, 5J, and 5K). There was a significant effect of

experimental group on the change in lick rate (Figure 5L, two-way ANOVA, $F = 8.37$, $p < 0.05$),

a significant effect of frequency ($F = 15.52$, $p < 0.05$), and a significant interaction between

group and frequency ($F = 13.08$, $p < 0.05$). Post-hoc tests showed that the interaction was

mediated by significant suppression of licking by 25 Hz ($p < 0.05$) and 50 Hz ($p < 0.05$)

stimulation. Like with the direct pathway, the effect of indirect pathway stimulation was

dependent on the time of stimulation. A two-way ANOVA revealed significant effects of both

time of stimulation in the task (Figure 5M, $F = 3.98$, $p < 0.05$) and laser frequency ($F = 40.05$, p

< 0.0001), with no significant interaction. Together, these results reveal a frequency-dependent

effect of indirect pathway stimulation on licking behavior. Furthermore, indirect pathway

activation shows less of an effect on licking occurring during consumption than on licking during

anticipation.

When we activated the indirect pathway at 5 seconds prior to reward delivery, when mice

are least likely to lick, we observed a striking rebound of licking that followed the offset of

stimulation (Figures 6A and B, Video S4). Licking rates during this rebound were higher during

the second following laser offset as compared to the same time point in the absence of

stimulation (Figure 6C, two-way mixed ANOVA, main effect of stimulation, $F = 11.73$, $p <$

0.01), but there was no main effect of frequency ($F = 1.29$, $p = 0.9$) and no interaction between

stimulation frequency and the rebound licking rate ($F = 1.16$, p = 0.78). Indirect pathway

stimulation also resulted in a reduced latency to initial lick onset following stimulation offset

when compared to non-stimulation trials (Figure 6D, two-way mixed ANOVA, $F = 16.05$, p <

0.001; main effect of frequency, $F = 11.09$, p = 0.05). There was a significant interaction

between trial type and the stimulation frequency, with higher frequencies resulting in lower onset

latencies ($F = 16.82$, p < 0.01). Post-hoc tests showed that the interaction was mediated by

shorter rebound latencies following 25 Hz (p < 0.05) and 50 Hz (p < 0.001) stimulation. Varying

stimulation frequencies resulted in differing effects on the regularity of rebound latencies,

indicated by measuring the variance of lick onset (Figure 6E, two-way mixed ANOVA, main

effect of frequency, $F = 20.66$, p < 0.01; main effect of stimulation, $F = 3.27$, p = 0.08;

interaction between stimulation frequency and trial type, $F = 25.63$, p < 0.001). Post-hoc tests

showed greater variance of lick onset time following 5 Hz stimulation (p < 0.05), and

significantly lower variance following 50 Hz stimulation (p < 0.01).

We observed that the effects of laser stimulation would be altered over the course of the

behavioral session, as a function of motivational state. For example, the effects of direct pathway

stimulation at 5 seconds prior to reward (Figure 7A) on changes in lick rate were often dependent

on how many trials the mice received (Figure 7B). At higher frequencies, this gradual reduction

of laser-generated licking rate was not as evident (Figure 7C). We divided the session into

quartiles and quantified the increase of licking rate by stimulation as a function of trial quartile

(Figure 7D). A two-way, mixed ANOVA revealed a significant interaction between frequency

and quartile ($F = 3.51$, p < 0.01) in addition to significant effects of frequency ($F = 45.05$, p <

0.001) and trial quartile ($F = 1.69$, p < 0.05). Post-hoc tests showed significant reductions in lick

rate between the 4_{th} quartile and the 1_{st} (p < 0.001), 2_{nd} (p < 0.001), and the 3_{rd} (p < 0.01)

quartiles under 10 Hz stimulation. We also found that the latency to stimulation-evoked lick

onset increased over time. A two-way, mixed ANOVA revealed a significant effect of trial

quartile on lick onset latency (Figure 7E, $F = 3.03$, p < 0.05) and a significant effect of frequency

($F = 35.34$, p < 0.01), but no interaction between trial quartile and stimulation frequency ($F =$

1.08, p = 0.9), suggesting a uniform increase in latency across all parameters. Like the effects of

stimulation of the direct pathway on lick rate, the latency to rebound licking following

stimulation at 5 seconds was affected by motivation (Figure 7F). The latency of rebound licking

increased as the mouse became sated in the course of a session, but this was most evident at

higher stimulation frequencies (Figure 7G, two-way mixed ANOVA, interaction between trial

quartile and stimulation frequency $F = 8.21$, $p < 0.05$; main effect of frequency, $F = 18.18$, $p <$

0.05; main effect of trial quartile $F = 5.12$, $p < 0.05$). Post-hoc tests showed that latency to

rebound during the 4_{th} quartile was significantly greater than during the 1_{st} and 2_{nd} quartiles

under 25 Hz ($p < 0.05$) and 50 Hz stimulation ($p < 0.01$). These results suggest that engagement

of lower-level licking generators by the BG can be modulated by motivational signals.

Our task allowed for the interrogation of the roles of the BG pathways in timing, as mice

will readily anticipate the arrival of the reward (Figure 8A). We incorporated probe trials on

which the reward was omitted to assess internal timing (Figure 8B). All mice generated

anticipatory licking bouts during probe trials. Averaging across many trials results in a

distribution of licking rates whose peak reflects the expected time of reward delivery, in this task

at approximately 10 s. On a fraction of the probe trials we stimulated direct and indirect

pathways to examine their contributions to timing.

We found that stimulation of the direct pathway during the probe trial could shift the

peak rightward by entire interval-duration. Thus, stimulation of the direct pathway concurrently

with reward did not result in an obvious peak shift (Figure 9A, *left*). However, as we moved the

stimulation time rightward by 3 or 5 seconds, the peak would move rightward proportionally

(Figure 9A, *middle* and *right*). Indeed, we found that the peak shift is a function of the start time

of direct pathway stimulation (Figure 9B, one-way, RM ANOVA; $F = 21.68$, $p < 0.001$).

Control mice did not display any shifts in peak timing (data not shown; one-way RM ANOVA; F

$= 0.7$, $p = 0.5$). We also quantified other characteristics of the peak distributions that have been

shown to be affected by pharmacological manipulations of the dopaminergic system. We found

that direct pathway stimulation reduced the peak width, but we did not observe a significant

effect of stimulation time on the width (Figure 9C, two-way mixed ANOVA; main effect of

stimulation, $F = 20.05$, $p < 0.0001$; no main effect of stimulation time, $F = 3.77$, $p = 0.8$; no

interaction between stimulation time and stimulation, $F = 2.42$, $p = 0.3$). Direct pathway

stimulation did not significantly impact the skew of the peak distributions (Figure 9D). These

results reveal a critical role of direct pathway activation in resetting the internal timekeeping

mechanism. This is the first report of a specific neural correlate of timer resetting. Interestingly,

stimulation prior to reward did not impact the peak, suggesting that normally reward receipt may

be involved in resetting the timing mechanism.

We also tested the effect of indirect pathway stimulation during probe trials. As observed

previously, stimulation of the indirect pathway at 5 seconds after reward resulted in a rebound in

licking following the termination of stimulation. This was also the case during peak probe trials.

Mice showed different types of responses following rebound licking. First, rebound licking

would result in a brief delay in licking which was followed by a bout of licking ('recovery'

pattern, Figure 10A). Moreover, rebound licking could also initiate the peak, so that the peak

lasts longer than usual ('initiation' pattern, Figure 10B). Occasionally mice also generate a peak

without showing a clear rebound following stimulation ('no-rebound' pattern, Figure 10C).

These different patterns (recovery, initiation, and no-rebound) were equally common (Figure

10D, n = 5, one-way RM ANOVA; $F = 0.2$, p > 0.05). Interestingly, peak analysis on these three

licking patterns revealed distinct effects of indirect pathway stimulation on interval timing. For

each pattern of licking, we compared the mean peak time with the peak time on trials in the same

session that did not contain laser stimulation. During recovery-pattern trials, we found a large

rightward shift of the peak that averaged approximately 2 seconds (Figure 10E, two-tailed t-test,

p < 0.01). In contrast, we found that when licking was initiated by the rebound licking bout, the

peak was not affected. On trials that lacked rebound licking, we found a rightward peak shift of

approximately 750 ms (two-tailed t-test, p < 0.05). Although initiation-type trials did not show a

change in peak timing, we found that these peaks showed increased duration (Figure 10F, two-

tailed t-test, p < 0.05). We did not find a significant impact on the skew of the peaks in any trial

type (Figure 10G).

Finally, we tested whether behavior following indirect pathway stimulation was altered as

the session progressed. The probability of the recovery pattern decreased as the session

progressed, whereas the likelihood of detecting no-rebound patterns increased (Figure 10H, two-

way, mixed ANOVA; interaction of the effects of trial type and session half, $F = 25.61$, p $<$

0.01). We did not detect a change in the probability of detecting the initiation-type pattern.

Discussion

We found for the first time that optogenetic activation of direct-pathway neurons in the

VLS can initiate licking (**Figure 2**), whereas activation of the indirect pathway in this region can

terminate licking (**Figure 5**). These results are in accord with previous work implicating the

VLS in orofacial behaviors (Mittler et al 1994, Pisa 1988a, Salamone et al 1993). The observed

effects are distinct from those following activation of the dorsolateral striatum, which produces

locomotion and turning behavior (Bartholomew et al 2016, Kravitz et al 2010, Tecuapetla et al

2014a). The acute effects on licking is not surprising given the topographical organization of the

sensorimotor striatum and our target in the ventrolateral orofacial region.

Although BG projections to lower brainstem pattern generators are not required for

reflexive orofacial and tongue movements (Grill & Norgren 1978), intact BG are required for

goal-directed food seeking behavior (Bignall & Schramm 1974). It has been shown that a subset

of GABAergic projections from SNr to the superior colliculus is critical for the initiation and

termination of self-paced licking behavior (Redgrave et al 1992, Rossi et al 2016, Toda et al

2017). A pause in licking can be caused by increased activity in nigrotectal output (Rossi et al

2016, Toda et al 2017). The GABAergic output from the SNr can also directly innervate the

reticular formation (Deniau & Chevalier 1992), which contains regions that are known to be

involved in the generation of licking behavior (Travers et al 1997). Direct pathway activation

may reduce SNr output and indirect pathway activation may increase SNr output. Consequently,

brainstem or tectal targets that receive GABAergic SNr projections may be disinhibited by direct

pathway activation and suppressed by indirect pathway activation. We previously showed that

opponent classes of SNr projection neurons command downstream position control systems in

the midbrain and brainstem to move the body in different directions (Barter et al 2015b, Fan et al

2012, Yin 2014a, Yin 2014c, Yin 2017). The present results suggest a potential role of the direct

and indirect pathways for modulating the output channels for control of the orofacial modalities,

which interact with distinct lower level circuitry.

Top-down regulation of lick pattern generator

A variety of evidence suggests that the output nuclei of the BG may function as an

integrator (Barter et al 2015a, Yin 2014b, Yin 2017). The rate of change in the integrator output

is proportional to the input magnitude. With continuous control of position, a larger striatal

signal (spikes per time window) is converted to faster rate of change in position control

command (higher movement velocity) as well as greater movement amplitude (Bartholomew et

al 2016). For licking behavior, we assume that the basal ganglia output also produces a top-

down command from an integrator that influences the operation of the brainstem pattern

generator. However, when the behavior in question is generated by pattern generators with

relatively fixed innate rhythms, the mechanism is different. In this case, the BG output does not

modulate the lick frequency in a continuous manner (Rossi et al. 2016), and it is unlikely that the

BG output is directly responsible for the generation of each lick. With continuous modulation,

we would expect lick frequency to linearly scale with direct pathway stimulation frequency

(**Figure 11A**), but this was not observed. Once activated, the pattern generator appears to

operate in several discrete frequency settings, analogous to a fan with discrete settings, such as

off, low, and high. Normally, the lick pattern generator oscillates at 5-6 Hz for anticipatory

licking and 6-8 Hz for consummatory licking. When the direct pathway was stimulated, licking

could reach a new regime (~10 Hz) which appears to be the limit of the lick pattern generator

and possibly also the biomechanical constraints of the orofacial musculature. When stimulation

frequency exceeded 10 Hz, the licking frequency cannot increase further.

BG output may regulate lick-generating centers' activation state around a certain

threshold, and direct pathway stimulation is sufficient to bring activity above that threshold. The

integrator is filled by direct pathway activation. Because direct pathway projections are

inhibitory, the integrator is filled by inhibiting GABAergic nigral neurons, which results in a

disinhibition of downstream structures. Greater activation of the direct pathway produces longer

suprathreshold activation of the pattern generator (**Figure 11B**). This could be responsible for the

increasing licking frequency levels that we observed (**Figure 11C**), which reach the maximum of

roughly 10 Hz. Evidence for the presence of an integrator is reflected in the increased power of

licking in a given frequency band, indicating higher duty cycle of the oscillator (**Figures 11A** &

11D). We also observed decreasing latencies to lick onset during higher stimulation frequencies,

and that licking could be sustained for some time after the end of stimulation (**Figure 4**).

Artificially driving the direct pathway efficiently fills the integrator, and this results in near

saturation even at relatively low stimulation frequencies, leading to animals achieving near-

maximum licking rates. However, mice rarely enter this regime on their own, perhaps due to the

concurrent activity of the indirect pathway.

The indirect pathway may therefore play an important role in the system by discharging

the integrator, preventing the instantaneous licking frequency from reaching its maximum level

in natural licking. Selective activation of the indirect pathway is equivalent to rapidly reversing

the BG output signal, bringing or keeping the pattern generators below their activation thresholds

and pausing behavior. It is therefore only under conditions of artificially activating the direct and

indirect pathways separately that behavior can be described as operating in an all-or-none, go/no-

go fashion. Prior observations of coincident activity of the two pathways may reflect coordinated

regulation of behavior to keep output within a certain range (DeLong 1990, Tecuapetla et al

2014b). The balanced relationship between these two pathways may be responsible for the

continuous adjustments to specific kinematic details of ongoing movement.

We propose that the BG provide top-down modulation of central pattern generators in the

form of duty cycle regulation. Coordinated activity of the direct and indirect pathways are

combined to produce a BG output that modulates how long a relatively stereotyped pattern

generator is activated. This form of top-down regulation is distinct from the more continuous

regulation of position controllers (Barter et al 2015a, Yin 2014c).

We observed a reliable rebound of licking following indirect pathway stimulation in the

middle of the interval (**Figure 6**). This was only possible to detect because mice rarely lick at

that time point in our task, and the effect was not masked by the delivery of reward or the

presence of consumption licking. In addition, higher frequency stimulation resulted in shorter

latency to rebound licking as well as more uniform timing in the start of rebound licking.

Indirect pathway stimulation may result in more active nigral output, which results in a large

suppression of downstream structures. The release of this inhibition then gives rise to a burst of

rebound activity responsible for lick generation, resulting in fast reactivation of licking CPGs

above the threshold, and a more consistent latency of the rebound licking.

The effects of direct and indirect pathway stimulation were sensitive to motivational

state, as the latencies to licking, overall licking rates, and the latency to rebound all changed over

the course of the session (**Figure 7**). This could not be explained as simply a reduction in the

efficacy of laser stimulation over the course of a single session because stimulation of the direct

pathway at higher frequencies could override the effects of reducing motivation. These results

can also be explained by the possibility that, over time, the activation threshold for licking

increases.

The effects of stimulation on licking varied in a manner that was dependent on the time in

the task when the stimulation occurred. Direct pathway stimulation had the greatest impact on

licking when mice were not licking, in the middle of the interval. On the other hand, stimulation

during ongoing licking had a smaller effect. This could be explained by a ceiling effect, as mice

could only boost their licking frequency so much once they were already performing the

behavior. The opposite trend was observed with indirect pathway stimulation: stimulation at 5

seconds prior to reward showed a lower change in rate than stimulation during anticipatory

licking.

Striatal contribution to timing

Our results also shed light on the distinct contributions of direct and indirect pathways to

timing. The use of the fixed-interval schedule allowed us to incorporate peak probe trials that

measure interval timing in mice.

Some have suggested that the brain's ability to time intervals is an emergent property of

sequential neural population dynamics (Bakhurin et al 2017, Crowe et al 2014, Mauk &

Buonomano 2004, Mello et al 2015). However, these models do not explain why specific neural

circuits are critical for timing; nor do they incorporate detailed anatomical organization such as

the direct and indirect pathways, which are shown to have distinct roles in timing. Our

observations support classic pacemaker-accumulator models (Gibbon et al 1984), which contain

specific mechanisms that can be paused or reset (Buhusi & Meck 2002). In fact, the proposed

BG integrator mechanism is quite similar to the pacemaker-accumulator model, which also

makes use of an integrator.

We found that stimulation of the direct pathway reset the interval timing mechanism and

activation of the indirect pathway could pause its operation (**Figures 9 & 10**). During peak probe

trials, mice show peak responding at the trained interval of 10 seconds. When stimulating the

direct pathway, we found that the peak would appear approximately 10 seconds following the

onset of stimulation. The interval was maintained, but the activation of the direct pathway

appeared to reinitiate the accumulation of the 10 second interval, just like resetting some

'internal clock'.

On the other hand, indirect pathway stimulation did not result in a resetting effect,

suggesting that resetting is specific to direct pathway activation and not a consequence of any

distracting neural perturbation. This also makes it unlikely that the rebound effect is mediated by

excessive direct pathway activity. We found an interaction between the timing system and the

presence of rebound licking immediately following indirect pathway stimulation. In general, the

chief effect of indirect pathway activation appears to be a pause in the timer, similar to what has

been observed previously with nigrotectal stimulation (Toda et al 2017). Driving the indirect

pathway paused the 'clock,' and the presence of rebound licking could extend the duration of the

pause. This impact on timing was only observed occasionally, as sometimes rebound licking

simply initiated peak responding. These trials were also noteworthy in that they resulted in a

peak of longer duration, but not one with a significant leftward shift in the peak time. Thus only

the precision of timing was affected. The distinct types of trials observed could be explained by

variable rates of filling the accumulator: faster accumulation on a given trial would allow the

rebound licking to initiate the peak, whereas with slower accumulation the indirect pathway

activity could interrupt or pause this process. In partial support of this idea are our observations

of the motivational effects on behavioral responses to indirect pathway stimulation. As rebound

licking latency increases over time, the proportion of peak-probe trials that contained rebound

licking also decreased.

The presence of distinct mechanisms for resetting and pausing an internal clock could be

related to the interaction of the direct and indirect pathways to regulate the integrator in the BG

output nuclei. The opponent impact that direct and indirect pathways have on licking is due to

their opponent roles in the accumulation process.

Conclusions

In summary, we show that direct pathway stimulation in the VLS can initiate licking,

whereas indirect pathway stimulation can suppress ongoing licking, and that these pathways can

work in concert to continuously regulate ongoing licking. In addition, the distinct effects of

direct and indirect stimulation on timing behavior suggest that each pathway differentially

interacts with the internal time-keeping system: direct pathway stimulation can reset the timer,

whereas indirect pathway activation can only pause or interfere with it. Together these results

suggest for the first time a uniform underlying mechanism that can explain the role of the BG

circuits in action generation as well as interval timing.

Some have argued that the accumulator is distributed in the cortex (Wang et al 2018).

Our results, however, suggest that it resides in the BG output circuit, though the underlying

mechanisms for integration remain unclear. Future work will be necessary to elucidate the

neural implementation of this integrator and how the BG output influence and modulate targets

in the midbrain and thalamus to regulate performance and timing of actions.

Finally, a few limitations of the current study should be noted. We cannot rule out that

the same striatal units may be involved in multiple different behaviors depending on context. As

the head-fixed preparation makes it impossible for the other behaviors to be expressed, the

trained behavior, namely licking, may become the prioritized action under this condition. It

would be important in future studies to examine the role of the VLS in unrestrained animals, and

the relationship between orofacial behavior and other components of natural appetitive

behaviors.

Methods

Subjects: 8 adult D1-Cre and 13 A2A-Cre mice of both sexes were used (Jackson Labs, Bar

Harbor, ME). These mice expressed Cre-recombinase only in D1 or A2A striatal neurons,

respectively. All experiments were conducted during the light phase of the 12:12 light cycle.

Mice had unrestricted access to food in their home cages, but were water deprived and received a

10% sucrose solution during behavioral sessions. Body weight was maintained at 85% body

weight and was monitored on a daily basis, with additional water given after experimental

training as needed. All experimental procedures were approved by the Duke University

Institutional Animal Care and Use Committee.

Viral vectors: AAV5-EF1α-DIO-hChR2(H134R)-eYFP was injected into 8 D1-Cre and 8 A2A-

Cre mice. AAV5-EF1α-DIO-eYFP was injected into 5 A2A-Cre mice for control experiments.

Vectors were obtained from the University of North Carolina Vector Core.

Surgery: Mice were placed in a stereotaxic frame and anesthetized with ~1-1.5% isoflurane.

Small holes were drilled bilaterally at AP: +0.2 mm and ML: +/- 2.5 mm relative to bregma.

Stereotaxically guided injections of (200 nL per hemisphere) were made at AP: +0.2 and ML:

2.5 and DV: -4.6 & -4.4 relative to bregma using a Nanoject III microinjector (Drummond

Scientific, Broomall, PA). 100 µm-core optic fibers (Precision Fiber Products, Chula Vista, CA)

were implanted with the same AP and ML coordinates and at DV: -4.2 in order to target cell

bodies of the VLS. Fibers were secured to the skull with screws and dental cement along with a

head post for head fixation.

System for behavioral training and recording: Mice were allowed 7-14 days to recover from

surgery and then were water deprived in their home cages. The interval timing task is the same as

previously described (Toda et al 2017). Briefly, behavioral experiments were conducted in a

sound-proof chamber with the mice perched in a custom-made elevated tunnel platform. For

head fixation, an implanted steel head post was clamped on both sides of the head. A metal

drinking spout was positioned directly in front of the mouth so that the mouse only needs to

protrude its tongue to access the spout. A 10% sucrose solution was gravity fed to the spout and

its delivery was controlled by the opening of a solenoid valve. The spout was connected to a

capacitance touch-sensor (MPR121, Adafruit, New York, NY) coupled to an Arduino Leonardo

(www.arduino.cc) that reported the time of each contact with the spout by the tongue. The task

was controlled using custom Matlab (version 2014b, Mathworks, Natick, MA) programs

interfaced with a Blackrock Cerebrus recording system (Blackrock Microsystems, Salt Lake

City, UT) to generate digital signals to control reward delivery, and analog commands to drive

laser stimulation. The Cerebrus system was simultaneously used for recording these signals in

addition to the digital timestamps received from the lickometer. All timestamps were saved for

offline analysis.

Fixed-interval timing task: Initial training began with water-deprived mice and involved

habituation to head-fixation and training mice to collect experimenter-delivered water drops

from the spout. Once reliable licking was established for drop delivery after a few days, training

began on a fixed-interval version of the task. During the task, mice received a 5 µL drop of

reward every 10 seconds. No external stimuli were presented. White noise was played inside of

the chamber to mask the sound of the solenoid (Rossi & Yin 2015). Water was presented at fixed

intervals for 200 trials a day until mice showed anticipatory licking behavior.

Peak probe trials: Once mice displayed consistent anticipatory licking in during fixed-interval

sessions, sessions began to incorporate peak probe trials. During peak trials, no reward is

presented in order to quantify the internal representation of time. Peak probe trials occurred with

a 60% possibility after 3 consecutive rewards. No indication was given as to when a peak trial

would occur. Peak trials lasted 30 seconds plus a random duration sampled from a gamma

distribution (Matlab function *gamrnd* using shape parameter = 2.5 and scale parameter = 4). The

consecutive reward counter was then reset with a reward delivery. Mice that struggled to learn

this version of the peak procedure were run on a schedule that used a 30% probability of peak

probe trial occurrence after 3 consecutive rewards.

Optogenetic stimulation: Optical stimulation occurred both in rewarded and probe trials to

prevent mice from predicting the probe trials with the sensation of the stimulation. Stimulation

on a given trial was determined using random sampling from a uniform distribution. Following 3

fixed-interval trials without laser delivery, the program entered a decision point. There was a

30% chance of laser delivery during a normal trial, a 30% chance of laser delivery during a peak

trial, and a 30% chance of a peak trial without laser. The remaining 10% of cases resulted in a

new decision point. This resulted in a 10-15% of all trials being a peak trial without stimulation,

a peak trial with stimulation, or a normal. Once head fixed in the testing chamber, mice were

connected bilaterally to a 473-nm DPSS laser (BL473T3, Shanghai Laser, Shanghai, China) via

fiber optic cables. A fiber splitter (TM105R5F1A, ThorLabs, Newton, NJ) divided the beam

(50:50) for bilateral stimulation. Stimulation was pulsed (0.8-2mW; 5-50 Hz, 10 ms square pulse

width, 1 s duration). Stimulation onset was either at -1 s from reward, was concurrent with the

time of reward (0s condition) or occurred 5 s prior to the reward. During peak probe trials,

stimulation was delivered at -1 s from reward, with the time of reward, or 5 seconds following

reward delivery. Mice received multiple sessions with stimulation in order to account for the

effects of stimulation frequency and timing relative to reward on licking, meaning that animals

received at least 12 (3 times in the trial x 4 stimulation frequencies) sessions of stimulation, in

addition to additional tests to investigate the effects on timing. Stimulation parameters

(stimulation frequency and timing relative to reward) were consistent within a session, but the

order of stimulation was semi-randomized between mice. Stimulation could be repeated over

multiple consecutive sessions without detriment to timing (data not shown).

Quantification and Statistical Analysis: Behavioral data was analyzed with NeuroExplorer (Nex

Technologies, Colorado Springs, CO) and Matlab. Licking timestamps were filtered to exclude

events occurring less than 8 ms apart. Unless otherwise stated, all analyses were performed on

the first half of trials to exclude motivational effects. Statistical tests were performed in

GraphPad Prism (GraphPad Software, San Diego, CA). Two-tailed or one-tailed parametric tests

were used. Post-hoc tests were done using Bonferroni correction. Power spectrum analysis was

performed in NeuroExplorer.

Peak analysis: Peak-detection and quantification was performed on a single-trial basis using a

custom-written Matlab algorithm. For each peak probe trial, licking time stamps were aligned to

the reward that initiated the trial. An analysis window extending 0 to 20 seconds following the

reward was used to detect peaks. Licking within this window was binned using 100ms time bins

to produce a continuous licking rate estimate and smoothed using a gaussian filter with a width

of 1 second. The Matlab function *findchangepts* was then applied to the smoothed lick rate signal

to detect steep transitions in licking rate. These change points were classified as rising or falling

transitions based on the slope of the lick rate signal around them. The first increasing change

point in the analysis window and the next falling change point were determined to represent the

start and end of the peak, respectively. The peak was fit with a gaussian function, the maximum

of which was taken as the peak for that trial. Peak duration had to be at least 2 seconds in

duration and licking rate had to exceed 3 Hz to be included. In cases of failures to identify a

peak, the analysis window was shifted by +500 milliseconds and the procedure was applied

again up to a maximum of 5 shifts, after which the trial would be discarded upon failure to detect

a valid peak. To detect peaks occurring after direct pathway stimulation at 3 or 5 seconds, the 20

second analysis window was shifted right by 3 or 5 seconds, respectively, following the reward

to exclude stimulation-evoked licking as a peak.

Lick onset latency analysis: Lick onsets was defined as the first lick occurring after all inter-lick-

intervals of 1 second or greater. For determining the latency to lick onset, the first lick onset

occurring during the 1 second laser stimulation was counted for each stimulation trial. For

determining rebound licking onset time, we measured the time until the first lick onset that

occurred within a window of 0 to 3 seconds following the offset of laser termination. The same

window was applied to non-laser trials for within-subject comparisons. Trials without licking

were excluded from analysis.

Post rebound peak trial classification: Peak trials that contained indirect pathway activation at 5

seconds post reward were classified by the temporal relationships of lick onsets relative to

rebound licking. Trials that contained a lick onset in a 2 second period following stimulation

onset were considered having a rebound. Otherwise, the trial was classified as lacking a rebound.

For trials classified as a rebound-containing peak trial, a second lick onset timestamp had to

occur within 8 seconds following initiation of rebound licking. In this case, the trial was

classified as a rebound recovery trial. Otherwise, if no lick onset timestamp followed the

rebound, the trial was classified as a rebound absorption trial. To detect peaks during rebound

recovery trials, the analysis window was shifted by 6 seconds from reward delivery in order to

exclude the rebound from being counted as a peak.

Author contributions:

Author contributions: XL and HHY designed research; KIB, XL, ADF, NAL, GDRW performed research; KIB, XL, ADF, NAL, and HHY analyzed data; and KIB, and HHY wrote the paper.

Acknowledgements:

This work was supported by NIH grants DA040701, NS094754 and MH112883 (HHY). We thank Dr. Fengxia Allen for technical assistance.

Competing interests:

The authors declare no competing financial interests.

References:

Albin RL, Young AB, Penney JB. 1989. The functional anatomy of basal ganglia disorders. *Trends Neurosci* 12: 366-75

Alexander GE, DeLong MR. 1985. Microstimulation of the primate neostriatum. II. Somatotopic organization of striatal microexcitable zones and their relation to neuronal response properties. *J Neurophysiol* 53: 1417-30

Bakhurin KI, Goudar V, Shobe JL, Claar LD, Buonomano DV, Masmanidis SC. 2017. Differential Encoding of Time by Prefrontal and Striatal Network Dynamics. *J Neurosci* 37: 854-70

Barter JW, Li S, Sukharnikova T, Rossi MA, Bartholomew RA, Yin HH. 2015a. Basal ganglia outputs map instantaneous position coordinates during behavior. *Journal of Neuroscience* 35: 2703–16

Barter JW, Li S, Sukharnikova T, Rossi MA, Bartholomew RA, Yin HH. 2015b. Basal ganglia outputs map instantaneous position coordinates during behavior. *J Neurosci* 35: 2703-16

Bartholomew RA, Li H, Gaidis EJ, Stackmann M, Shoemaker CT, et al. 2016. Striatonigral control of movement velocity in mice. *The European journal of neuroscience* 43: 1097-110

Bignall KE, Schramm L. 1974. Behavior of chronically decerebrated kittens. *Exp Neurol* 42: 519-31

Buhusi CV, Meck WH. 2002. Differential effects of methamphetamine and haloperidol on the control of an internal clock. *Behav Neurosci* 116: 291-7

Buhusi CV, Meck WH. 2005. What makes us tick? Functional and neural mechanisms of interval timing. *Nature reviews. Neuroscience*

Crowe DA, Zarco W, Bartolo R, Merchant H. 2014. Dynamic representation of the temporal and sequential structure of rhythmic movements in the primate medial premotor cortex. *J Neurosci* 34: 11972-83

Cui G, Jun SB, Jin X, Pham MD, Vogel SS, et al. 2013. Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature* 494: 238-42

Delfs J, Kelley A. 1990. The role of D 1 and D 2 dopamine receptors in oral stereotypy induced by dopaminergic stimulation of the ventrolateral striatum. *Neuroscience* 39: 59-67

DeLong MR. 1990. Primate models of movement disorders of basal ganglia origin. *Trends in neurosciences* 13: 281-85

Deniau JM, Chevalier G. 1992. The lamellar organization of the rat substantia nigra pars reticulata: distribution of projection neurons. *Neuroscience* 46: 361-77

Fan D, Rossi MA, Yin HH. 2012. Mechanisms of action selection and timing in substantia nigra neurons. *J Neurosci* 32: 5534-48

Frederick DL, Allen JD. 1996. Effects of selective dopamine D1- and D2-agonists and antagonists on timing performance in rats. *Pharmacol Biochem Behav* 53: 759-64

Freeze BS, Kravitz AV, Hammack N, Berke JD, Kreitzer AC. 2013. Control of basal ganglia output by direct and indirect pathway projection neurons. *J Neurosci* 33: 18531-9

Gerfen CR, Engber TM, Mahan LC, Susel Z, Chase TN, et al. 1990. D1 and D2 dopamine receptor-regulated gene expression of striatonigral and striatopallidal neurons. *Science* 250: 1429-32

Gibbon J, Church RM, Meck WH. 1984. Scalar timing in memory. *Ann N Y Acad Sci* 423: 52-77

Gouvea TS, Monteiro T, Motiwala A, Soares S, Machens C, Paton JJ. 2015. Striatal dynamics explain duration judgments. *Elife* 4

Grill HJ, Norgren R. 1978. The taste reactivity test. II. Mimetic responses to gustatory stimuli in chronic thalamic and chronic decerebrate rats. *Brain Res* 143: 281-97

Hintiryan H, Foster NN, Bowman I, Bay M, Song MY, et al. 2016. The mouse cortico-striatal projectome. *Nature neuroscience* 19: 1100-14

Isomura Y, Takekawa T, Harukuni R, Handa T, Aizawa H, et al. 2013. Reward-modulated motor information in identified striatum neurons. *J Neurosci* 33: 10209-20

Jicha GA, Salamone JD. 1991. Vacuous jaw movements and feeding deficits in rats with ventrolateral striatal dopamine depletion: possible relation to parkinsonian symptoms. *J Neurosci* 11: 3822-9

Kelley AE, Lang CG, Gauthier AM. 1988. Induction of oral stereotypy following amphetamine microinjection into a discrete subregion of the striatum. *Psychopharmacology* 95: 556-9

Kravitz AV, Freeze BS, Parker PR, Kay K, Thwin MT, et al. 2010. Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature* 466: 622-6

Mauk MD, Buonomano DV. 2004. The neural basis of temporal processing. *Annual review of neuroscience* 27: 307-40

Meck WH. 1996. Neuropharmacology of timing and time perception. *Brain Res Cogn Brain Res* 3: 227-42

Meck WH. 2006. Neuroanatomical localization of an internal clock: a functional link between mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Res* 1109: 93-107

Mello GB, Soares S, Paton JJ. 2015. A scalable population code for time in the striatum. *Current biology : CB* 25: 1113-22

Merchant H, Harrington DL, Meck WH. 2013. Neural basis of the perception and estimation of time. *Annual review of neuroscience* 36: 313-36

Mink JW. 1996. The basal ganglia: focused selection and inhibition of competing motor programs. *Prog Neurobiol* 50: 381-425

Mittler T, Cho J, Peoples LL, West MO. 1994. Representation of the body in the lateral striatum of the freely moving rat: single neurons related to licking. *Exp Brain Res* 98: 163-7

Pavlov I. 1927. *Conditioned Reflexes*. Oxford: Oxford University Press.

Pisa M. 1988a. Motor functions of the striatum in the rat: critical role of the lateral region in tongue and forelimb reaching. *Neuroscience* 24: 453-63

Pisa M. 1988b. Motor somatotopy in the striatum of rat: manipulation, biting and gait. *Behavioural brain research* 27: 21-35

Redgrave P, Marrow L, Dean P. 1992. Topographical organization of the nigrotectal projection in rat: evidence for segregated channels. *Neuroscience* 50: 571-95

Roberts S. 1981. Isolation of an internal clock. *J Exp Psychol Anim Behav Process* 7: 242-68

Rossi MA, Li HE, Lu D, Kim IH, Bartholomew RA, et al. 2016. A GABAergic nigrotectal pathway for coordination of drinking behavior. *Nature neuroscience*

Rossi MA, Yin HH. 2015. Elevated dopamine alters consummatory pattern generation and increases behavioral variability during learning. *Frontiers in integrative neuroscience* 9: 37

Salamone JD, Mahan K, Rogers S. 1993. Ventrolateral striatal dopamine depletions impair feeding and food handling in rats. *Pharmacol Biochem Behav* 44: 605-10

Shammah-Lagnado SJ, Costa MS, Ricardo JA. 1992. Afferent connections of the parvocellular reticular formation: a horseradish peroxidase study in the rat. *Neuroscience* 50: 403-25

Tecuapetla F, Matias S, Dugue GP, Mainen ZF, Costa RM. 2014a. Balanced activity in basal ganglia projection pathways is critical for contraversive movements. *Nat Commun* 5: 4315

Tecuapetla F, Matias S, Dugue GP, Mainen ZF, Costa RM. 2014b. Balanced activity in basal ganglia projection pathways is critical for contraversive movements. *Nature communications* 5

Toda K, Lusk NA, Watson GDR, Kim N, Lu D, et al. 2017. Nigrotectal Stimulation Stops Interval Timing in Mice. *Current biology : CB* 27: 3763-70 e3

Travers JB, Dinardo LA, Karimnamazi H. 1997. Motor and premotor mechanisms of licking. *Neuroscience & Biobehavioral Reviews* 21: 631-47

Wang J, Narain D, Hosseini EA, Jazayeri M. 2018. Flexible timing by temporal scaling of cortical responses. *Nature neuroscience* 21: 102-10

Yin HH. 2014a. Action, time and the basal ganglia. *Philos Trans R Soc Lond B Biol Sci* 369: 20120473

Yin HH. 2014b. Action, time and the basal ganglia. *Philosophical Transactions of the Royal Society B: Biological Sciences* 369

Yin HH. 2014c. How basal ganglia outputs generate behavior. *Advances in neuroscience* 2014: 768313

Yin HH. 2017. The Basal Ganglia in Action. *Neuroscientist* 23

Figure 1

Figure 2

Figure 3

Figure 4

A

Laser on

-5 -4   0

B

Trials with 5Hz laser

-1   0   1   2   3
Time from laser onset (s)

Trials with 25Hz laser

-1   0   1   2   3
Time from laser onset (s)

C

Lick onset latency (s)

0   20   40   60
Stimulation frequency (Hz)

D

● Direct stim
● No stim

Post stim lick count

0   20   40   60
Stimulation frequency (Hz)

Figure 5

Figure 6

Figure 7

A

Direct stim

B

D

C

E

Indirect stim

F

G

Figure 8

<u>**Figure 9**</u>

Figure 10

Figure 11

A

D1 stim-evoked licking frequency

Hypothetical linear model

D1 stim-evoked licking power

B

5 Hz direct stim

SNr activity

C

Lick pattern generator

← *Physical limit*

Low activation threshold

D

⊞ *Low lick frequency*
⊞ *High lick frequency*

Lower duty cycle

10 Hz direct stim

SNr activity

Lick pattern generator

← *Physical limit*

High activation threshold

25 Hz direct stim

SNr activity

Lick pattern generator

Higher duty cycle

Figure legends

Figure 1. Stimulation of the direct and indirect pathways of the orofacial striatum during a licking-based interval timing task

A) *Left*: Strategy of injection of AAV-ChR2 virus to orofacial area of the striatum in D1-Cre or A2A-Cre mice and implantation of chronic optical fibers over the injection site. **B**) Serial section diagrams showing target zone for orofacial striatum. **C**) Representative image of eYFP-expression in ventrolateral striatum in a D1-Cre+ mouse injected with AAV-DIO-ChR2-eYFP. Scale bar = 500 µm. **D**) Representative image of eYFP-expression in an A2A-Cre+ mouse injected with AAV-DIO-ChR2-eYFP virus. Scale bars = 500 µm. (Abbreviations: *ic*, internal capsule; *CP*, caudoputamen; *GPe*, globus pallidus, external segment.) **E**) Diagram illustrating the fixed-time interval task for head-fixed mice. Mice receive a 10% sucrose reward (5 µL) every 10 seconds. **F**) Example lick raster of a well-trained mouse during normal trials aligned to reward delivery. Tick marks represent individual licks. **G**) Mean licking rate calculated from the raster in *E*. Error denotes SEM.

Figure 2. Stimulation of direct pathway in VLS generates licking

A) Depiction of stimulation strategy 5 seconds prior to reward in D1-cre mice. **B**) *Top*: Peri-event lick raster diagrams of representative trials demonstrating normal anticipatory licking patterns in trials without laser. *Bottom*: 10 Hz stimulation at 5 seconds prior to reward results in large increases in licking activity. Both rasters reflect licking from the same mouse in the same session. **C**) Mean lick rate calculated from the example session shown in *B* for trials with and without laser stimulation. **D**) Mean increases in licking rate during laser delivery resulting from stimulation of D1 MSNs relative to control at 5 seconds increased as a function of stimulation frequency (n = 8 D1-ChR2 mice, n = 5 control mice; two-way ANOVA; main effect of experimental group, $F = 56.4$, $p < 0.0001$; main effect of frequency, $F = 8.38$, $p < 0.01$; interaction between frequency and group, $F = 8.76$, $p < 0.01$). **E**) Depiction of stimulation strategy 1 second prior to reward. **F**) *Top*: Peri-event lick raster diagrams of representative non-laser trials. *Bottom*: Raster diagram of trials containing 10 Hz stimulation during anticipatory licking. Both rasters reflect licking from the same mouse in the same session. **G**) Mean lick rate calculated from the example session shown in *F* for trials with and without laser stimulation. **H**) Stimulation of D1 MSNs at 1 second prior to reward increased the rate of anticipatory licking (two-way ANOVA, main effect of group, $F = 54.11$, $p < 0.0001$). **I**) Depiction of stimulation strategy coinciding with reward delivery. **J**) *Top*: Representative lick raster diagrams of licking behavior during trials without laser stimulation. *Bottom*: Raster diagram of trials from the same animal containing 10 Hz stimulation during consumption licking. Both rasters reflect behavior recorded in the same session. **K**) Mean licking rate calculated from the example session shown in *J* for trials with and without stimulation. **L**) Stimulation of D1 MSNs was capable of increasing consumption licking rate (two-way ANOVA, main effect of group, $F = 16.43$, $p < 0.01$). **M**) The time of stimulation during the interval influenced the change in licking rate (two-way ANOVA, main effect of time of stimulation, $F = 17.65$, $p < 0.0001$; main effect of frequency, $F = 31.22$, $p < 0.0001$). Error denotes SEM. *X* symbol reflects a significant interaction between factors.

Figure 3. Direct pathway boosts licking frequency and modulates licking duty cycle

A) *Left*: Example lick rasters from two mice showing licking frequency potentiation by direct pathway stimulation 1 second before reward. Note increase of licking to upwards of 10 Hz. Blue tick-marks denote laser pulse times. *Right*: Example lick raster from the same sessions showing normal licking activity around reward. **B**) Demonstration of spectral density analyses of licking during laser stimulation, anticipation (1 second intervals beginning prior to reward), and consumption (1 second intervals immediately following reward) periods. Analysis of data from the sessions represented in *A* is shown. **C**) Power spectral density distributions of licking during stimulation at increasing laser frequencies. Note consistent increase in power. **D**) Effect of stimulation frequency on peak power of the PSD when stimulation occurred during anticipatory licking. Peak licking power was related to stimulation frequency (n = 8 mice; two-way mixed ANOVA; main effect of lick pattern, $F = 48.01$, p < 0.0001; interaction lick pattern vs frequency, $F = 12.92$, p < 0.05). **E**) Mean licking frequency corresponding to the peak of the PSD as a function of laser frequency for stimulation during anticipatory licking. Stimulation increased the licking frequency (two-way mixed ANOVA; $F = 31.62$, p < 0.0001). **F**) Effect of stimulation frequency on peak power of licking when stimulation occurred 5 seconds before reward. Peak power was related to stimulation frequency (two-way mixed ANOVA; main effect of lick pattern, $F = 25.9$, p < 0.0001; interaction lick pattern vs frequency, $F = 13.28$, p < 0.01). **G**) Mean licking frequency corresponding to the peak of the PSD as a function of laser stimulation frequency delivered 5 seconds prior to reward. Stimulation increased the licking frequency (two-way mixed ANOVA; $F = 43.22$, p < 0.0001). **H**) Effect of stimulation frequency on peak power of licking when stimulation occurred during consumption licking. Peak licking power was related to stimulation frequency (two-way mixed ANOVA; main effect of lick pattern, $F = 44.1$, p < 0.0001; interaction lick pattern vs frequency, $F = 9.76$, p < 0.05). **I**) Mean licking frequency corresponding to the peak of the PSD as a function of laser frequency for stimulation during consumption licking. Stimulation increased the licking frequency (two-way mixed ANOVA; $F = 62.35$, p < 0.0001). Error bars reflect SEM. *X* symbol reflects a significant interaction between factors.

Figure 4. Direct pathway activates modulates onset latency and duration of evoked licking bout

A) *Top*: The effect of stimulation frequency on evoked licking was measured for sessions with stimulation occurring at 5 seconds prior to reward. *Bottom*: Example raster plot showing longer latency to lick and shorter bout duration during 5 Hz direct pathway stimulation. **B**) Stimulation of the direct pathway at 25 Hz resulted in rapid licking onset and sustained licking following offset of laser stimulation. **C**) Latency to lick bout onset as a function of the frequency of direct pathway stimulation (n = 8; one-way ANOVA; $F = 4.92$, p < 0.01). **D**) Number of licks that were counted during the 1 second following direct pathway stimulation offset. Blue points reflect counts during trials with laser stimulation at 5 seconds before reward. Black points reflect the baseline count number observed during trials without stimulation for the same time period of the interval (two-way mixed ANOVA; main effect of stimulation: $F = 12.23$, p < 0.001; interaction stimulation vs. frequency, $F = 3.88$, p = 0.15). Error bars denote SEM.

Figure 5. Stimulation of the indirect pathway in the orofacial striatum suppresses licking

A) Depiction of stimulation strategy 5 seconds prior to reward in A2A-cre mice. **B**) *Top*: Peri-event lick raster diagrams of representative trials demonstrating normal anticipatory licking patterns in trials without laser. *Bottom*:10 Hz stimulation at 5 seconds prior to reward results in a pause in licking. Both rasters reflect licking from the same mouse in the same session. **C**) Mean lick rate calculated from the example session shown in *B* for trials with and without laser stimulation. **D**) Mean reduction of licking rate during stimulation of the indirect pathway at 5 seconds (n = 8 A2A-ChR2 mice, n = 5 control mice; two-way ANOVA; main effect of experimental group, $F = 18.56$, $p < 0.01$). **E**) Depiction of stimulation strategy 1 second prior to reward. **F**) *Top*: Peri-event lick raster diagrams of representative non-laser trials. *Bottom*: Raster diagram of trials containing 10 Hz stimulation of the indirection pathway during anticipatory licking. Both rasters show licking from the same mouse in the same session. **G**) Mean lick rate calculated from the example session shown in *F* for trials with and without laser stimulation. **H**) Increasing stimulation frequency of the indirect pathway at 1 second prior to reward reduced the rate of anticipatory licking in A2A-Cre mice (two-way ANOVA; main effect of group, $F = 25.93$, $p < 0.0001$; main effect of frequency, $F = 19.47$, $p < 0.0001$; interaction group vs frequency, $F = 13.83$, $p < 0.01$). **I**) Depiction of stimulation strategy coinciding with reward delivery. **J**) *Top*: Representative lick raster diagrams of licking behavior during trials without laser stimulation. *Bottom*: Raster diagram of trials from the same mouse containing 10 Hz stimulation during consumption licking. Both rasters reflect behavior recorded in the same session. **K**) Mean licking rate calculated from the example session shown in *J* for trials with and without stimulation. **L**) Indirect pathway stimulation at the time of reward reduced licking rate (two-way ANOVA; main effect of group, $F = 8.37$, $p < 0.05$; main effect of frequency, $F = 15.52$, $p < 0.05$; interaction group vs frequency, $F = 13.08$, $p < 0.05$). **M**) The time of stimulation during the interval influenced reductions in licking rate by stimulation (two-way ANOVA, main effect of time of stimulation, $F = 3.98$, $p < 0.05$; main effect of Frequency, $F = 40.05$, $p < 0.0001$). Error reflects SEM. *X* symbol reflects a significant interaction between factors.

Figure 6. Rebound licking following stimulation of the indirect pathway

A) *Top:* Schematic showing stimulation of A2A-Cre mice 5 seconds prior to reward. *Middle*: Example raster plot showing the resulting rebound licking following the termination of stimulation of the indirect pathway. Orange markers indicate the onset time of discrete lick bouts. *Bottom*: Raster plot of licking during no-stimulation trials from the same mouse, aligned to the same time point in the interval as above. Note the predominant alignment of lick onset times following laser stimulation. **B**) *Top*: Example of another mouse showing rebound licking after indirect pathway stimulation. *Bottom*: Raster plot of licking during non-stimulation trials from the same mouse in the same session. **C**) Comparison of licking rate during stimulation and non-stimulation trials. Indirect pathway stimulation resulted in an increase in licking rate following laser termination when compared to the same time period in non-stimulation trials (n = 8 mice; two-way mixed ANOVA, effect of stimulation $F = 11.73$, $p < 0.01$). **D**) Reduction of lick onset latency following stimulation termination compared to mean latency for the same time period in non-stimulation trials (two-way mixed ANOVA, main effect of stimulation: $F = 16.05$, $p < 0.001$; interaction stimulation vs trial type, $F = 16.82$, $p < 0.01$). **E**) Reduction of lick onset

latency variance following stimulation offset compared to the variance of licking initiation for the same time period in non-stimulation trials (two-way mixed ANOVA, main effect of frequency, $F = 20.66$, $p < 0.01$; interaction frequency vs. trial type, $F = 25.63$, $p < 0.001$). Error bars denote SEM. *X* symbol reflects a significant interaction between factors.

Figure 7. The effect of motivation on licking affected by direct and indirect pathway stimulation

A) The effect of motivation on licking related to direct and indirect pathway manipulation was measured for sessions with stimulation occurring at 5 seconds prior to reward. **B**) Example raster plot showing progressive reduction of licking evoked by 10 Hz direct pathway stimulation at 5 seconds before reward. Trial numbers reflect the number of laser presentations delivered with the number of rewards the animal received in parentheses. **C**) Example raster plot showing a reduced influence of motivation on evoked licking with higher frequency stimulation of the direct pathway. Data is from the same mouse but a different session as shown in *B*. Trial numbers reflect the number of laser presentations delivered with the number of rewards the animal received in parentheses. **D**) Grouped bar plot showing the reduced impact of laser stimulation at 5 s before reward on evoked licking as a function of trial quartile (n = 8; two-way mixed ANOVA; effect of quartile, $F = 1.69$, $p < 0.05$; effect of frequency, $F = 31.13$, $p < 0.01$; interaction quartile vs frequency, $F = 3.51$, $p < 0.01$). **E**) Grouped bar plot showing an increase of lick onset latency as a result of direct pathway stimulation as a function of trial quartile (two-way mixed ANOVA, effect of quartile, $F = 3.03$, $p < 0.05$; effect of frequency, $F = 35.34$, $p < 0.01$; interaction quartile vs frequency, $F = 1.08$, $p = 0.93$). **F**) Example raster plot showing the increase in rebound latency following 50Hz stimulation of the indirect pathway over the course of the experimental session. Trial numbers reflect the number of laser presentations delivered with the number of rewards the animal received in parentheses. **G**) Grouped bar plot showing the increase of lick onset latency following laser stimulation offset as a function of trial quartile (n = 8, two-way mixed ANOVA, effect of quartile, $F = 5.21$, $p < 0.05$; effect of frequency, $F = 18.18$, $p < 0.05$; interaction quartile vs frequency, $F = 8.21$, $p < 0.05$). Error bars reflect SEM. *X* symbol reflects a significant interaction between factors.

Figure 8. Peak probe trials reveal the operation of an internal timing mechanism

A) *Top*: Example raster plot of licking during consecutive 10 second fixed-time trials. Licking is aligned to the first of two consecutive reward delivery times. *Bottom*: Mean licking rate for the session shown in the above raster. **B**) *Top*: Example raster plot showing licking during peak probe trials. During peak trials, reward is delivered then withheld 10 seconds later, resulting in a discrete bout of licking in the absence of any stimuli. *Bottom*: The average licking rate for the session shown in the above raster. Averaging across probe trials results in a characteristic peak in licking. Data shown for both trial types were recorded from the same mouse and session. Error reflects SEM.

Figure 9. Direct pathway stimulation resets the internal clock

A) *Left*: Mean licking rate across subjects during peak probe trials with and without laser stimulation concurrent with reward delivery. Scale bars reflect the population mean peak times

for probe trials with (blue) and without (green) laser stimulation. *Middle*: Mean licking rate across subjects during probe trials delivered in the presence and absence of laser stimulation at 3 seconds post reward. Scale bars reflect the population mean peak times for probe trails with (blue) and without (green) laser stimulation. *Right*: Mean licking rate across subjects during normal probe trials and peak probe trials containing laser stimulation at 5 seconds following reward. Scale bars reflect the population mean peak times for probe trials with (blue) and without (green) laser stimulation. **B**) Magnitudes of peak shifts in seconds as a function of time of direct pathway stimulation. The y-axis reflects the difference between the mean peak time occurring during laser trials subtracted by the mean peak time during trials without stimulation (n = 4; One-way RM ANOVA; $F = 21.68$, p < 0.001). Peak analyses were performed on the entire behavioral session to maximize statistical power. **C**) Quantification of the peak duration during normal peak probe trials and those with laser stimulation. Stimulation resulted in a reduction of the peak width (n = 4; two-way mixed ANOVA; effect of stimulation, $F = 20.05$, p < 0.0001). **D**) There were no changes in the skewness of the peak distributions. Error bars denote SEM.

Figure 10. Indirect pathway stimulation pauses the internal clock

A) *Top*: Peri-event lick raster of representative peak probe trials without laser delivery. *Middle*: 25 Hz stimulation at 5 seconds following reward during trials in which rebound licking causes a delay and a recovery in peak licking. Both rasters reflect licking from the same mouse in the same session. *Bottom*: Mean lick rate calculated from the example session shown above for peak probe trials without laser stimulation and stimulation trials showing a "recovery" pattern. **B**) *Top*: Peri-event lick raster diagrams of representative peak probe trials without laser delivery. *Middle*: 25 Hz stimulation at 5 seconds following reward during trials in which rebound licking initiates peak licking. Both rasters show licking from the same mouse in the same session. *Bottom*: Mean lick rate calculated from the example session shown above for peak probe trials without laser stimulation and stimulation trials showing an "initiation" pattern. **C**) *Top*: Peri-event lick raster of a third subject's representative peak probe trials produced in the absence of laser delivery. *Middle*: 25 Hz stimulation at 5 seconds following reward during trials in which no rebound licking was detected. *Bottom*: Mean lick rate calculated from the example session shown above for peak probe trials without laser stimulation and stimulation trials showing a "no rebound" pattern. **D**) The three types of patterns occurred with equal probability throughout the behavioral session (n = 5; one-way RM ANOVA, $F = 0.08$, p = 0.92). **E**) Peak shifts were measured by subtracting the mean peak time of each pattern by the mean peak time without stimulation. Recovery and no rebound trials showed significant positive peak shifts (two-tailed t-tests, p < 0.05), whereas initiation trials trended toward negative peak shifts (two-tailed t-test, p = 0.1). **F**) Initiation trials showed significant increases in the duration of the peak (two-tailed t-test, p < 0.05). **G**) There were no changes in skewness of licking distributions for any trial type. **H**) Fraction of peak trials showing each pattern during the first and second halves of the behavioral session. On the whole, the recovery pattern was gradually replaced with no rebound pattern (two-way, mixed ANOVA, interaction trial type x session half, $F = 25.61$, p < 0.01). Error bars denote SEM. *X* symbol reflects a significant interaction between factors.

Figure 11. Direct pathway modulates licking CPG activity via integration
A) *Left*: Laser-evoked licking frequency corresponding to the peak of the power spectral density distribution as a function of direct pathway laser stimulation frequency. Data for stimulation

frequencies 5-25 Hz are shown. *Middle*: Hypothetical expected linear laser-evoked licking frequencies if direct pathway stimulation directly drove each lick. Note that in this scenario, 5 Hz stimulation would result in licking at 5 Hz and 25 Hz stimulation would result in licking at 25 Hz. *Right*: Peak power of laser-evoked licking as a function of direct pathway laser stimulation frequency. **B**) Proposed mechanism translating direct pathway stimulation to SNr output activity via integration. Higher frequency direct pathway stimulation results in faster filling of the integrator that leads to a faster rate of change as well as longer-lasting output. **C**) Increasing SNr output brings lower-level licking centers above several different activity thresholds, corresponding to varying licking frequencies. CPG output is capped at 10Hz. **D**) Greater filling of the integrator results in a more sustained licking output at a given frequency. This is reflected in the increasing power in a given frequency band with increasing frequency of stimulation, suggesting a role for the modulation of licking duty cycle by the BG.