Vineyard ecosystems are structured and distinguished by fungal communities impacting

the flavour and quality of wine

Di Liu[1], Qinglin Chen[1], Pangzhen Zhang[1], Deli Chen[1] and Kate S. Howell[1,*]

[1] School of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, University of
Melbourne Parkville 3010 Australia

[*]Corresponding author. khowell@unimelb.edu.au

Abstract

The flavours of foods and beverages are formed by the agricultural environment where the plants are

grown. In the case of wine, the location and environmental features of the vineyard site imprint the

wine with distinctive aromas and flavours. Microbial growth and metabolism play an integral role in

wine production from the vineyard to the winery, by influencing grapevine health, wine fermentation,

and the flavour, aroma and quality of finished wines. The mechanism by which microbial distribution

patterns drive wine metabolites is unclear and while flavour has been correlated with bacterial

composition for red wines, bacterial activity provides a minor biochemical conversion in wine

fermentation. Here, we collected samples across six distinct winegrowing areas in southern Australia

to investigate regional distribution patterns of both yeasts and bacteria and how this corresponds with

wine aroma compounds. Results show that soil and must microbiota distinguish winegrowing regions

and are related to wine chemical profiles. We found a strong relationship between microbial and wine

metabolic profiles, and this relationship was maintained despite differing abiotic drivers (soil

properties and weather/ climatic measures). Notably, fungal communities played the principal role in

shaping wine aroma profiles and regional distinctiveness. We found that the soil microbiome is a

potential source of grape- and must-associated fungi, and therefore the weather and soil conditions

could influence the wine characteristics via shaping the soil fungal community compositions. Our

study describes a comprehensive scenario of wine microbial biogeography in which microbial

diversity responds to surrounding environments and ultimately sculpts wine aromatic characteristics.

These findings provide perspectives for thoughtful human practices to optimise food and beverage

flavour and composition through understanding of fungal activity and abundance.

Keywords: wine regionality, microbial biogeography, fungal diversity, climate, soil

Introduction

Regional distinctiveness of wine traits, collectively known as "*terroir*", can be measured by chemical composition and sensory attributes (Gonzálvez et al 2009, Pereira et al 2005, Robinson et al 2012), and this variation has been related to the physiological response of grapevines to local environments, such as soil properties (e.g., soil type, texture, nutrient availability), climate (temperature, precipitation, solar radiation), topography and human-driven agricultural practices (Gladstones 1992, van Leeuwen and Seguin 2006, Vaudour et al 2015). Wines made from the same grape cultivar but grown in different regions are appreciated for their regional diversity, increasing price premiums and the market demand (van Leeuwen and Seguin 2006). However, which factors from the vineyard to the winery drive wine quality traits have remained elusive.

Microbial biogeography contributes to the expression of wine regionality. Microorganisms, including yeasts, filamentous fungi and bacteria, originate in the vineyard and are impacted upon by the built environment (winery) and play a decisive role in wine production and quality of the final product (Barata et al 2012, Stefanini and Cavalieri 2018). The fermentative conversion of grape must (or juice) into wine is a complex and dynamic process, involving numerous transformations by multiple microbial species (Swiegers et al 2005). The majority of fermentations involve *Saccharomyces* yeasts conducting alcoholic fermentation (AF) and lactic acid bacteria (LAB) for malolactic fermentation (MLF), but many other species are present and impact the chemical composition of the resultant wine (Goddard 2008, Heard and Fleet 1985). Recent studies pose the existence of non-random geographical dispersion patterns of microbiota in grapes and wines (Bokulich et al 2014, Gayevskiy and Goddard 2012, Morrison-Whittle and Goddard 2018, Pinto et al 2015, Portillo and Mas 2016, Stefanini et al 2016, Taylor et al 2014). Few studies have explored the associations between microbial communities and wine chemical composition (Bokulich et al 2016, Knight et al 2015). Knight et al. (2015) showed empirically that regionally distinct *S. cerevisiae* populations drove different metabolites present in the resultant wines. Bokulich et al. (2016) suggested that wine metabolite profiles were more closely associated with bacterial profiles of wines undergone both AF and MLF. However, both numerically

and biochemically, yeasts dominate wine fermentation, so it is unclear why bacteria should be

dominant.

The vineyard soil has long been attributed to fundamental wine characteristics and flavour. Vineyard

soil provides the grapevine with water and nutrients and soil type and properties profoundly affect

vine growth and development (van Leeuwen and Seguin 2006). Soil-borne microbiota associate with

grapevines in a beneficial, commensal, or pathogenic way, and determine soil quality, host growth

and health. For example, soil microbes can mineralize soil organic matter, trigger plant defence

mechanisms, and thus influence flavour and quality of grapes and final wines (Müller et al 2016,

Rillig et al 2017). Alternatively, serving as a reservoir for grapevine-associated microbiota, the soil

microbiome can be recruited from the soil to the grapevine and transferred to the grape and must

(Morrison-Whittle and Goddard 2018, Zarraonaindia et al 2015). Some of these microbes can

influences the fermentation and contribute to final wine characteristics (Gilbert et al 2014, Stefanini

and Cavalieri 2018). Geographic areas can constrain the vineyard soil microbiota (Burns et al 2015,

Coller et al 2019, Knight et al 2019, Mezzasalma et al 2018, Zarraonaindia et al 2015), but whether

the biogeographic patterns can link to resulting wine attributes has not been clearly shown.

Limited but increasing evidence reveals that environmental heterogeneity conditions microbial

biogeography in wine production (recently reviewed by Liu et al (2019), Bokulich et al 2014, Burns et

al 2015, Jara et al 2016, Martiny et al 2006, Zarraonaindia et al 2015), along with differentiations by

geographic distances (Burns et al 2015, Martiny et al 2006, Miura et al 2017). Local climatic

conditions significantly correlate with microbial compositions in grape musts, for example

precipitation and temperatures that associate with the abundance of filamentous fungi (for example

Cladosporium and *Penicillium* spp.) and ubiquitous bacteria (for example, the *Enterobacteriaceae*

family) (Bokulich et al 2014), as well as yeast populations (particularly *Hanseniaspora* and

Metschnikowia spp.) (Jara et al 2016). Dispersion of soil microbiota is driven by soil physicochemical

properties such as soil texture, soil pH, carbon (C) and nitrogen (N) pools (Burns et al 2015, Colle et

al 2019, Zarraonaindia et al 2015), with some influences from topological characteristics (for example

orientation of the vineyard) (Burns et al 2015, Portillo et al 2016). Soil microbiome/ bacteria may

colonise grapes by physical contact (moving by rain splashes, dust, winds) (Zarraonaindia et al 2015)

or migration through the plant (xylem/phloem) from the rhizosphere to the phyllosphere (Compant et

al 2008, Mandl et al 2015). Insects help the movement and dispersal of microbes in the vineyard and

winery ecosystem, for example, honey bees, social wasps and *Drosopholid* flies can vector yeasts

among different niches (Goddard et al. 2010, Lam and Howell, 2015; Stefanini et al. 2012). The

vineyard microbes enter the winery associated with grapes or must, so the effects of environmental

conditions are finally reflected on microbial consortia in wine fermentation. How environmental

conditions modulate microbial ecology from the vineyard to the winery and shape regional

distinctiveness of wine is still largely unknown.

Here, we sampled microbial communities from the vineyard to the winery across six geographically

separated wine-producing regions in southern Australia to provide insights into microbial

biogeography, growing environments and wine regionality. We evaluated the volatile chemicals of

wines made with Pinot Noir grapes to validate that these different regions have differently flavoured

wines. Using next-generation sequencing (NGS) to profile bacterial and fungal communities, we

demonstrate that the soil and must microbiota exhibit distinctive regional patterns, and this can be

correlated to the wine metabolome. Network analysis supports the interaction of microbial community

abundance with abiotic factors (weather and soil properties) to drive wine regionality both directly

and indirectly. The important contributions arise from fungal diversity. We investigated a potential

route of wine-related fungi from the soil to the grapes by isolating yeasts from the xylem/phloem of

grapevines. Using vineyards, grapes and wine as a model food system, we have related the regional

identity of an agricultural commodity to biotic components in the growing system to show the

importance of conserving regional microbial diversity to produce distinctive foods and beverages.

Materials and methods

Sample sites and weather parameters

15 *Vitis vinifera* cv. Pinot Noir vineyards were selected in 2017 from Geelong, Mornington Peninsula (Mornington), Macedon Ranges, Yarra Valley, Grampians and Gippsland in southern Australia, ranging from 5 km to 400 km between vineyards (Supplementary Figure S1). In 2018, the sampling from the five vineyards in Mornington Peninsula (all < 20 km apart) was repeated to elucidate the influence of sampling year (vintage) on microbial patterns and wine profiles. Each site's GPS coordinates (longitude, latitude, altitude) were utilised to extract weekly weather data from the dataset provided by Australian Water Availability Project (AWAP). Variables were observed by robust topography resolving analysis methods at a resolution of $0.05° \times 0.05°$ (approximately 5 km x 5 km) (Jones et al 2009). Retrospective weekly measurements for all vineyards were extracted for mean solar radiation (MS), mean high temperature (MHT), mean low temperature (MLT), maximum temperature (MaxT), minimum temperature (MinT), mean temperature (MT), mean precipitation, mean relative soil moisture, mean evaporation (soil + vegetation), and mean transpiration (MTrans) in growing seasons (October 2016/ 2017 - April 2017/ 2018).

Collection of soil and plant materials

In each vineyard, soil samples were collected from three sites at harvest March-April 2017 (n = 45) and 2018 (n = 15), at a depth of 0 - 15 cm, 30 – 50 cm distant from the grapevine. Comprehensive vineyard samples were collected from two vineyards 5 km apart in the 2018 vintage. The vineyards were managed by a single winery, and the viticultural management was very similar. Five replicate Pinot Noir vines in each vineyard were selected from the top, middle, and bottom of the dominant slope, covering topological profiles of the vineyard. For each grapevine, five different sample types were collected: soil (0 - 15 cm deep, root zone), roots, xylem/ phloem sap, leaves, and grapes were collected at harvest in March 2018. Xylem sap (n = 10) was collected from the shoots using a centrifugation method in aseptic conditions (López-Millán et al 2000). Details of xylem sap collection, nutrient composition analysis, and yeast isolation were provided in supplementary

materials. Samples (n = 50) were stored in sterile bags, shipped on ice, and stored at -80 °C until

processing.

Longitudinal samples to study microbial abundance during fermentation were collected at five time

points: must (destemmed, crushed grapes prior to fermentation), at early fermentation (AF, less than

10% of the sugar is fermented), at middle of fermentation (AF-Mid, around 50% of sugar fermented),

at the end of fermentation (AF-End, following pressing but prior to barrelling), at the end of

malolactic fermentation (MLF-End, in barrels). Grapes and must sampled in this study were

fermented in the wineries following similar fermentation protocols of three days at a cool temperature

(known as cold-soaking) followed by warming up the must, so fermentation could commence.

Fermentations were conducted startedwithout addition of commercial yeasts and bacteria. Two

fermentations from Grampians and Mornington did not complete and were excluded from analysis,

giving wine samples from 13 vineyards in the 2017 vintage. Triplicate sub-samples from tanks or

barrels were collected and mixed as composite samples (n = 90). All samples were shipped on ice and

stored at -20 °C until processing.

Soil analysis

Edaphic factors were analysed to explore the effects of soil properties on wine-related microbiota and

aroma profiles. Soil pH was determined in a 2:5 soil/water suspension. Soil carbon (C), nitrogen (N),

nitrate and ammonium were analysed by Melbourne Trace Analysis for Chemical, Earth and

Environmental Sciences (TrACEES) Soil Node, at the University of Melbourne. Total C and N were

determined using classic Dumas method of combustion (Rayment and Lyons 2011) on a LECO

TruMac CN at a furnace temperature of 1350°C (LECO Corporation, Michigan, USA). Nitrate and

ammonium were extracted with 2 M KCl and determined on a Segmented Flow Analyzer (SAN++,

Skalar, Breda, Netherlands) (Rayment and Lyons 2011).

Wine volatile analysis

Volatile compounds of wine samples (MLF-End) were collected to represent the wine aroma. Wine

volatiles were determined using headspace solid-phase microextraction gas-chromatographic mass-

spectrometric (HS-SPME–GC/MS) method (Liu et al 2016, Zhang et al 2015) with some

modifications. Analyses were performed with Agilent 6850 GC system and a 5973 mass detector

(Agilent Technologies, Inc., CA, USA), equipped with a PAL RSI 120 autosampler (CTC Analytics

AG, Switzerland). Briefly, 10 mL wine was added to a 20 ml glass vial with 2 g of sodium chloride

and 20 µL of internal standard (4-Octanol, 100 mg/L), then equilibrated at 35 °C for 15 min and

immersed by a polydimethylsiloxane/divinylbenzene (PDMS/DVB, Supelco) 65 µm SPME fibre for

10 min at 35°C with agitation. The fibre was desorbed in the GC injector for 4 min at 220 °C.

Volatiles were separated on an Agilent J&W DB-Wax Ultra Inert capillary GC column (30 m × 0.25

mm × 0.25 µm), with helium carrier gas at a flow rate of 0.7 mL/min. The column temperature

program was as follows: holding 40 °C for 10 min, increasing at 3.0 °C/min to 220 °C and holding at

this temperature for 10 min. The temperature of transfer line of GC and MS was set at 240 °C. The

ion source temperature was 230 °C. The electron impact (EI) at 70 eV scanning over a mass

acquisition range of 35–350 m/z. Raw data were analysed with Agilent ChemStation Software for

qualification and quantification (Liu et al 2015). Volatile compounds (n = 88) were identified in wine

samples according to retention indices reference standards and mass spectra matching with NIST11

library. 13 successive levels of standard solution in model wine solutions (12% v/v ethanol saturated

with potassium hydrogen tartrate and adjusted to pH 3.5 using 40% w/v tartaric acid) were analysed

by the same protocol as wine samples to establish the calibration curves for quantification. Peak areas

of volatile compounds were integrated via target ions model. The concentrations of volatile

compounds were calculated with the calibration curves and used for downstream data analysis.

DNA extraction and sequencing

Genomic DNA was extracted from botanical and soil samples using PowerSoil™ DNA Isolation kits

(QIAgen, CA, USA). DNA extraction of soil (0.25 g) and xylem sap (0.5 mL) samples was directly

followed the kit protocol. Wine fermentation samples were thawed and recovered the biomass by

centrifugation at 4,000 × g for 15 min, washed three times in ice-cold phosphate buffered saline

(PBS) with 1% polyvinylpolypyrrolidone (PVPP) and centrifuged at 10,000 × g for 10 min (Bokulich

et al 2014). The obtained pellets were used for DNA extraction following the kit protocol. For

grapevine samples, roots, leaves, grapes (removed seeds and stems), were ground into powder under

the protection of liquid nitrogen with 1% PVPP, and isolated DNA following the kit protocol

afterward. DNA extracts were stored at -20 °C until further analysis.

Genomic DNA was submitted to Australian Genome Research Facility (AGRF) for amplification and

sequencing. To assess the bacterial and fungal communities, 16S rRNA gene V3-V4 region and

partial fungal internal transcribed spacer (ITS) region were amplified using the universal primer pairs

341F/806R (Yu et al 2005) and ITS1F/2 (Gardes and Bruns 1993), respectively. The primary PCR

reactions contained 10 ng DNA template, 2× AmpliTaq Gold® 360 Master Mix (Life Technologies,

Australia), 5 pmol of each primer. A secondary PCR to index the amplicons was performed with

TaKaRa Taq DNA Polymerase (Clontech). Amplification were conducted under the following

conditions: for bacteria, 95 °C for 7 min, followed by 30 cycles of 94 °C for 30 s, 50 °C for 60 s, 72

°C for 60 s and a final extension at 72 °C for 7 min; for fungi, 95 °C for 7 min, followed by 35 cycles

of 94 °C for 30 s, 55 °C for 45 s, 72 °C for 60 s and a final extension at 72 °C for 7 min. PCR

products were purified, quantified and pooled at the same concentration (5 nM). This was followed by

300 bp paired-end sequencing on an Illumina MiSeq (San Diego, CA, USA).

Raw sequences were processed using QIIME v1.9.2 (Caporaso et al., 2010). Low quality regions (Q <

20) were trimmed from the 5' end of the sequences, and the paired ends were joined using FLASH

(Magoč and Salzberg 2011). Primers were trimmed and a further round of quality control was

conducted to discard full length duplicate sequences, short sequences (< 100 nt), and sequences with

ambiguous bases. Sequences were clustered followed by chimera checking using UCHIME algorithm

from USEARCH v7.1.1090 (Edgar et al., 2011). Operational taxonomic units (OTUs) were assigned

using UCLUST open-reference OTU-picking workflow with a threshold of 97% pairwise identity

(Edgar 2010). Singletons or unique reads in the resultant data set were discarded, as well as the

chloroplast and mitochondrion related reads. Taxonomy was assigned to OTUs in QIIME using the

Ribosomal Database Project (RDP) classifier (Wang et al 2007) against GreenGenes bacterial 16S

rRNA database (v13.8) (DeSantis et al 2006) or the UNITE fungal ITS database (v7.2) (Kõljalg et al

2005), respectively.

Data analysis

Raw data are publicly available in the National Centre for Biotechnology Information Sequence Read

Archive under the bioproject PRJNA594458 (bacterial 16S rRNA sequences) and PRJNA594469

(fungal ITS sequences).

Microbial alpha-diversity was calculated using the Shannon index in R (v3.5.0) with the "vegan"

package (Oksanen et al 2007). One-way analysis of variance (ANOVA) was used to determine

whether sample classifications (e.g., region, fermentation stage) contained statistically significant

differences in the diversity after Bonferroni correction with the "ggsignif" package (Ahlmann-Eltze

2017). Principal coordinate analysis (PCoA) was performed to evaluate the distribution patterns of

wine metabolome and wine-related microbiome based on beta-diversity calculated by the Bray–Curtis

distance with the "labdsv" package (Roberts 2007). Permutational multivariate analysis of variance

(PERMANOVA) with 999 permutations was conducted within each sample classification to

determine the statistically significant differences with "adonis" function in "vegan" (Oksanen et al

2007).

Significant taxonomic differences of wine microbiome between fermentation stages were tested using

linear discriminant analysis (LDA) effect size (LEfSe) analysis (Segata et al 2011)

(https://huttenhower.sph.harvard.edu/galaxy/). The OTU table was filtered to include only OTUs >

0.01% relative abundance to reduce LEfSe complexity. This method applies the factorial Kruskal–

Wallis sum-rank test ($\alpha = 0.05$) to identify taxa with significant differential abundances between

categories (using all-against-all comparisons), followed by the logarithmic LDA score (threshold =

2.0) to estimate the effect size of each discriminative feature. Significant taxa were used to generate

taxonomic cladograms illustrating differences between sample classes.

The co-occurrence/interaction patterns between wine metabolome and microbiota in the soil and must

were explored in network analysis using Gephi (v0.9.2) (Bastian et al 2009). Only top 100 OTUs

(both bacteria and fungi) based on the relative abundance were used to construct the network. A

correlation matrix was calculated with all possible pair-wise Spearman's rank correlations between

volatile compounds and OTUs. Correlations with a Spearman correlation coefficient $\rho \geq 0.8$ and a $p <$

0.01 were considered statistically robust and displayed in the networks (Junker and Schreiber 2008).

A Random forest supervised-classification model (Breiman 2001) was conducted to identify the main

predictors of wine regionality among the following variables: must and soil microbial diversity

(Shannon index), soil properties, and weather. The importance of each predictor is determined by

evaluating the decrease in prediction accuracy (that is, increase in the mean square error (MSE)

between observations and out-of-bag predictions) when the data is randomly permuted for the

predictor. This analysis was conducted with 5,000 trees using the "randomForest" package in R (Liaw

and Wiener 2002). The significance of the model and the cross-validated R^2 were assessed using the

"A3" package (ntree = 5000) (Fortmann-Roe 2013). The structural equation model (SEM) (Grace

2006) was used to evaluate the direct and indirect relationships between must and soil microbial

diversity, soil properties, climate and wine regionality. SEM is an *a priori* approach partitioning the

influence of multiple drivers in a system to help characterize and comprehend complex networks of

ecological interactions (Eisenhauer et al 2015). An *a priori* model was established based on the

known effects and relationships among these drivers of regional distribution patterns of wine aroma to

manipulate the data before modelling. Weather and soil properties were used as composite variables

(both Random Forest and SEM) to collapse the effects of multiple conceptually-related variables into

a single-composite effect, thus aiding interpretation of model results (Grace 2006). A path coefficient

describes the strength and sign of the relationship between two variables (Grace 2006). The good fit

of the model was validated by the χ2-test ($p > 0.05$), using the goodness of fit index (GFI > 0.90) and

the root MSE of approximation (RMSEA < 0.05) (Schermelleh-Engel et al 2003). The standardised

total effects of each factor on the wine regionality pattern were calculated by summing all direct and

indirect pathways between two variables (Grace 2006). All the SEM analyses were conducted using

AMOS v25.0 (AMOS IBM, NY, USA).

SourceTracker was used to track potential sources of wine-related fungi within the vineyards (Knights

et al 2011). SourceTracker is a Bayesian approach that treats each give community (sink) as a mixture

of communities deposited from a set of source environments and estimates the proportion of taxa in

the sink community that come from possible source environments. When a sink contains a mixture of

taxa that do not match any of the source environments, that portion of the community is assigned to an

"unknown" source (Knights et al 2011). In this model, we examined musts (n = 2) and vineyard

sources (n = 50) including grapes, leaves, xylem sap, roots, and soils. The OTU tables were used as

data input for modelling using the "SourceTracker" R package

(https://github.com/danknights/sourcetracker).

Results

Chemical composition/ aroma profiles separate wines based on geographic location

Using GC-MS, we analysed the volatile compounds of Pinot Noir wine samples (MLF-End)

representing distinct wine growing regions. These were finished wines, which are being prepared for

sale to consumers and enabled the metabolite profiles to be compared directly to the microbial

communities inhabiting the musts from which these wines were fermented. In all, 88 volatile

compounds were identified in these wines. We used α- and β-diversity measures to elucidate wine

complexity and regionality, respectively. In 2017, α-diversity varied with regional origins (ANOVA,

$p < 0.001$), with higher Shannon indices (2.17 ± 0.05) were observed in the wines from regions of

Mornington, Yarra Valley and Gippsland than others (1.94 ± 0.03) (Figure 1A). PCoA showed that

74.5% of the variance was explained by the first two principal coordinates, and that wines within

regions grouped together (Bray–Curtis, ADONIS, $p < 0.001$) (Figure 1B). Overall, wine aroma

profiles display highly significant regional differentiation across both vintages (ADONIS, $R^2 = 0.566$,

$p < 0.001$) and the clustering patterns become more distinct and regression coefficients improve when

comparing regional differences in 2017 (ADONIS, $R^2 = 0.703$, $p < 0.001$) (Supplementary Table S1).

Microbial ecology from the vineyard to the winery

To elucidate how microbial ecology drives regional traits of wine from the vineyard to the winery,

150 samples covering soils, musts and fermentations were collected to analyse wine-related

microbiota. A total of 11,508,480 16S rRNA and 12,403,610 ITS high-quality sequences were

generated from all the samples, which were clustered into 13,689 bacterial and 8,373 fungal OTUs

with a threshold of 97% pairwise identity, respectively.

The dominant bacterial taxa across all soil samples were *Actinobacteria*, *Proteobacteria*,

Acidobacteria, *Chloroflexi*, *Verrucomicrobia*, *Bacteroidetes*, *Gemmatimonadetes*, *Firmicutes*,

Planctomycetes and *Nitrospirae* (Supplementary Figure S2A). Compared with bacteria, soil fungal

communities were less diverse. *Ascomycota* was the most abundant and diverse phylum of fungi,

accounting for 72% of reads, followed by *Basidiomycota*, *Mortierellomycota*, *Chytridiomycota* and

Olpidiomycota (Supplementary Figure S2B). The microbiome richness, α-diversity (Shannon index)

varied significantly with regions for both bacteria and fungi (ANOVA, $p < 0.01$) (data not shown).

Soil community structures widely varied across different grape growing regions, exerting significant

impacts on both bacterial and fungal taxonomic dissimilarity (Bray–Curtis, ADONIS, $p < 0.001$)

independent of vintage, with clearer differences within a single vintage (Supplementary Table S1). In

2017, soil fungal communities could discriminate growing regions (except Yarra Valley and

Gippsland) (Figure 2B), whereas regional separation of bacteria was weaker with overlap between

two or three regions (Figure 2A).

In grape musts, bacterial communities across six wine growing regions in both vintages consisted of

ubiquitous bacteria *Enterobacteriales*, *Rhizobiales*, *Burkholderiales*, *Rhodospirillales*,

Actinomycetales, *Sphingomonadales*, *Pseudomonadales*, *Saprospirales*, and *Xanthomonadales*, which

do not participate in wine fermentations or spoilage (Barata et al 2012). The LAB *Lactobacillales*,

responsible for malolactic fermentation, was present in low abundance (average = 0.4%) in the must

(Figure 3A). Fungal profiles were dominated by filamentous fungi, mostly of the genera

Aureobasidium, *Cladosporium*, *Botrytis*, *Epicoccum*, *Penicillium*, *Alternaria*, and *Mycosphaerella*,

with notable populations of yeasts, including *Saccharomyces*, *Hanseniaspora*, and *Meyerozyma*, as

well as the *Basidiomycota* genus *Rhodotorula* (Figure 3B). Pinot Noir musts exhibited significant

regional patterns for fungal communities across vintages 2017 and 2018 (Bray–Curtis, ADONIS, $p <$

0.001) but no significant differences for bacterial communities across both vintages (Bray–Curtis,

ADONIS, $p = 0.152$) (Supplementary Table S1). Within the 2017 vintage, both bacteria and fungi in

the must showed distinctive structures and composition based on the region (Bray–Curtis, ADONIS, p

< 0.001), with more distinct trend and improved regression coefficient for fungi ($R^2_{2017, 2018} = 0.292$,

$R^2_{2018} = 0.565$) (Figure 3A, B) (Supplementary Table S1). Notably, the relative abundance of

Saccharomyces yeasts varied significantly from 1.3% (Macedon Range) to 65.6% (Gippsland)

between regions (ANOVA, $p < 0.01$) (Figure 3B). As the wine fermentation proceeded, fermentative

populations including yeasts and LAB grew and dominated, thus reshaping the community diversity

(Supplementary Figure S3A, B) and composition (Supplementary Figure S3C, D). Fungal species

diversity collapsed as alcoholic fermentation progressed (ANOVA, $p < 0.001$) (Supplementary Figure

S3B), while the impact of the fermentation on bacteria was less than fungi, with slight decreased

diversity at early stages and significant recovery at the end (ANOVA, $p < 0.05$) (Supplementary

Figure S3A). Linear discriminant analysis (LDA) effect size (LEfSe) analysis further identified

differentially abundant taxa (Kruskal–Wallis sum-rank test, $\alpha < 0.05$) associated with fermentation

stages (Figure 3). For fungal populations, *Dothideomycetes* (including *Aureobasidium* and

Cladosporium), *Debaryomycetaceae* (notably yeast *M. guilliermondii*), *Penicillium corylophilum* and

Filobasidium oeirense were significantly abundant in the grape must, with *Leotiomycetes*,

Sarocladium and *Vishniacozyma victoriae* in early fermentations (AF), *Saccharomycetes* yeasts

(notably *S. cerevisiae*) in mid fermentations (AF-Mid), and *Tremellales* at the end of fermentation

(AF-End) (Figure 3D). For bacterial communities, *Acidobacteriia* (spoilage), *Chloroflexi*,

Deltaproteobacteria, *Sphingobacteriia*, *Cytophagia*, *Planococcaceae* and *Rhizobiaceae* were

observed with higher abundances in the must, *Proteobacteria* (including *Burkholderiaceae* and

Tremblayales, spoilage) and *Micrococcus* in the AF, *Burkholderia* spp. in the AF-Mid,

Rhodobacterales and *Pseudonocardiaceae* in the AF-End, LAB *Leuconostocaceae* (notably

Oenococcus) in the MLF-End (Figure 3C). Regional differences in microbial profiles were not

significant in the finished wines (ADONIS, $R^2_{bacteria} = 0.149$, $p = 0.321$; $R^2_{fungi} = 0.109$, $p = 0.205$)

(Supplementary Table S1).

To uncover the impact of growing season (vintage) on wine regionality and related microbiota, five

vineyards in Mornington were sampled in 2017 and 2018 to compare within and between vintages.

Within these five vineyards alone, both microbial patterns and resulting wine aroma profiles see a

significant vintage influence (ADONIS, $p < 0.001$) (Supplementary Table S1). However, vintage only

weakly impacts microbial patterns and wine regionality, in particular an insignificant influence on

fungal patterns when comparing all must samples (ADONIS, $p = 0.066$) (Supplementary Table S1).

We used 2017 vintage data to further explore microbial biogeography and wine regionality in the

following analyses.

Microbial patterns correlate to wine aroma profiling

Network analysis was used to explore connections between regional microbial and wine metabolic

patterns. We selected 57 volatile compounds and 246 OTUs based on strong correlation coefficients

(Spearman correlation coefficient, $\rho \geq 0.8$; $p < 0.01$), forming the co-occurrence patterns. These

variables showed a sophisticated internal structure, consisting of 303 nodes and 610 edges (average

degree = 4.026), with mostly positive correlations (90.66%) (Figure 4A). Both must (average degree =

3.791) and soil microbiota (average degree = 3.333) presented high connectivity with wine

metabolites (Figure 4B, C). Fungal OTUs (n = 69) more densely connected with volatile compounds

than bacteria (n = 42) in the soil (Figure 4B). This indicated that soil microbial communities,

especially fungi, play an important role in wine regionality. The most densely connected volatile

compounds were assigned to groups of monoterpenes (C18, *p*-cymene; C46, *α*-terpineol),

phenylpropanoids (C71, benzaldehyde; C74, phenethyl acetate), and acetoin (C20) (Figure 4A).

Correspondingly, high-connectivity nodes with these compounds were OTUs of bacterial taxa

Enterobacteriales, *Rhizobiales*, *Burkholderiales* and *Xanthomonadales*, and fungal taxa *Penicillium*,

Rhodotorula and *Neocatenulostroma*, from soil and must (Figure 4B, C). The group of esters, which

are yeast-driven products (Swiegers et al 2005), were highly associated with fungal OTUs (in

particular *Hypocreales*, *Alternaria*, *Cryptococcus and Mortierella*) from the vineyard soil, for

example, ethyl hexanoate (C16) and ethyl dihydrocinnamate (C87) (Figure 4B). Fatty acids (including

C47, 4-hydroxybutanoic acid; C77, octanoic acid) were mostly negatively connected with fungi

(including *Acremonium*, *Penicillium*) from the must (Figure 4C).

Multiple drivers modify wine regionality in the vineyard

To disentangle the role of microbial ecology on wine regionality expression, we used Random Forest

modelling (Breiman 2001) to identify important predictors (weather/ climatic conditions, soil

properties, microbial diversity of soil and must) driving wine regionality, and structural equation

modelling (SEM) (Grace 2006) to testify whether the relationship between microbial diversity and

wine regionality can be maintained when accounting for multiple drivers simultaneously. The

Random Forest model demonstrated that fungal diversities were major predictors of wine regionality.

Not surprisingly, must fungal diversity played an important role (increase in MSE) than soil (Figure

5). The SEM explained 94% of the variance found in the pattern of wine regionality (Figure 6A).

Weather drove wine aroma profiles directly (especially MT, MLT, MinT and MS) and indirectly by

powerful effects on soil and must microbial diversity, in particular strong effects on soil fungal

diversity (Figure 6A). Must fungal diversity had the highest direct positive effect on wine aroma

characteristics, with direct influences from soil fungal diversity (Figure 6A). Weather and climate

could impact soil nutrient pools primarily through MS, MLT, MinT, and MTrans. Soil properties

showed strong effects on soil microbial diversity and must bacterial diversity, but weak effects on

must fungal diversity (Figure 6A). Must bacterial diversity had a weak effect on wine aroma profiles,

as well as soil bacteria. Overall, must fungal diversity is the most important driver of wine

characteristics, followed by soil fungal diversity (Figure 6B), with influences from weather and soil

properties both directly and indirectly (Figure 6A).

Source tracking wine-related fungi within vineyard

As we showed in the SEM, soil fungal diversity had a direct positive influence on must fungal

diversity and thus indirectly contributed to wine aroma profiles (Figure 6). Given that soil is a

potential source of fungi associated with wine production (Morrison-Whittle and Goddard 2018), here

we attempt to uncover the mechanism whereby soil fungi could be transported from soil to the grapes.

We sampled fungal communities from grapevines and soil and hypothesised that the xylem/ phloem

was the internal mechanism to transport microbes. A total of 2,140,820 ITS high-quality sequences

were generated from soil and grapevine samples (grape, leaf, xylem sap, root), which were clustered

into 4,050 fungal OTUs with 97% pairwise identity. Using SourceTracker (Knights et al 2011), fungal

communities in the must were matched to multiple sources from the belowground to the aboveground.

Results showed that grape and xylem sap were primary sources of must fungi, with 32.6% and 41.9%

contributions, respectively (Figure 7A). Xylem sap showed similar fungal structure with must (Figure

S4A). Further source tracking revealed that the root and soil contributed 90.2% of fungal OTUs of

xylem sap which contributed 67.9% of the fungi of grapes (Figure 7B, C).

Notably, *S. cerevisiae* were found shared between niches of soil, root, xylem sap, grape and must,

with the highest (1.2%) and lowest (0.04%) abundance in the root and soil, respectively (Figure S4B).

To explore whether xylem/phloem sap could be a translocation pathway of *S. cerevisiae*, we

attempted to culture yeasts from the collected xylem sap. Chemical analysis of nutrient compositions

showed that xylem sap contained nine carbohydrates (predominantly glucose, fructose and sucrose),

15 amino acids (mainly arginine, aspartic acid and glutamic acid) and six organic acids (primarily

oxalic acid), which could be utilised as carbon sources and support yeast growth (Figure S4E, F, G)

(Yadeta and Thomma 2013). No *S. cerevisiae* yeasts were isolated, but distinct isolates of the

Basidiomycota yeasts of *Cryptococcus* spp. (primarily *C. saitoi*) and *Rhodotorula slooffiae* were

found (Figure S4C). Similar species of yeasts were isolated when xylem/phloem sap was collected

from grapevines grown in the glasshouse (Figure S4D).

Discussion

Microbial ecology can influence grapevine health and growth, fermentation, wine quality and style and flavour characteristics (Bokulich et al 2014, Gayevskiy and Goddard 2012, Knight et al 2015). Here we show that wine regionality is dependent on fungal ecology and is sensitive to local weather, climate and soil properties. A new mechanism to transfer fungi from the soil to grapes and must via xylem/ phloem sap was investigated. We systematically investigated the microbiome from the soil to wine to show that soil and grape must microbiota exhibit regional patterns and that these patterns can be translated into wine metabolites.

A microbial component of wine *terroir*

Biogeographic patterns are clearly present in the soil and grape must microbiota. The most abundant bacterial phyla in the soil were *Actinobacteria*, *Proteobacteria* and *Acidobacteria*, which are known dominant and ubiquitous in the vineyard soil worldwide (Burns et al 2015, Coller et al 2019, Fierer et al 2012, Zarraonaindia et al 2015). For fungi, the community composition and structure were more diverse within regions we studied and compared with studies in other major wine-producing areas in the world. Overall, we recovered 14 phyla, 30 classes, 65 orders, 125 families and 216 genera of fungi, recording a higher diversity than found in previous studies (Coller et al 2019, Knight et al 2019, Morrison-Whittle et al 2017). *Glomeromycota* and *Blastocaldiomycota* were reported as abundant phyla across vineyards in Marlborough, New Zealand (Morrison-Whittle and Goddard 2017). In our study, however, *Glomeromycota* was only recovered with low frequency in soils coming from Mornington and Macedon Ranges, and *Blastocaldiomycota* was not detected (data not shown). In the must, both principal fermentation drivers (*S. cerevisiae* and LAB) and innocent species (those that neither ferment nor spoil, such as *Enterobacteriales* and *Aureobasidium*) were present in different abundances among regions (Figure 3A, B, ANOVA, $p < 0.01$). The order *Lactobacillales*, representing LAB, was present 0.4% across regions, compared to 29.7% found in California, US (Bokulich et al 2014) and 14% in Catalonia, Spain (Portillo et al 2016).

Environmental factors (such as weather and climate) and geographic characteristics (in particular

latitude) drive microbial diversity and biogeography across various habitats, such as soil, ocean, and

plant ecosystems (Bokulich et al 2014, Fierer and Jackson 2006, Martiny et al 2006, Tedersoo et al

2014). In this study, we demonstrate that microbial biogeographic communities are distinct in both

vineyard soils and grape musts at the scale of 400 km and this is not dependent on the growing

season/ vintage. This aligns with previous studies on wine microbial biogeography and provide

further evidence for microbial *terroir* (reviewed by Liu et al. 2019; Bokulich et al 2014, Burns et al

2015, Morrison-Whittle and Goddard 2018, Taylor et al 2014). Soil bacteria can distinguish wine

growing regions, with impacts from soil properties (Figure 6A) and this is supported by previous

work in this field (Burns et al 2015, Mezzasalma et al 2018, Zarraonaindia et al 2015). An interesting

finding is that the must bacterial diversity is strongly influenced by soil properties, in particular C: N.

Previous work has shown that structures of must and soil communities are similar and some species

Enterobacteriales and *Actinomycetales* originate from the soil (Pinto et al 2015, Stefanini and

Cavalieri 2018). As C: N can be manipulated by composting and cover crops (Burns et al 2016) there

is an opportunity to manipulate wine microbiota by a focus on vineyard management (Liu et al 2019).

The bacterial microflora is recognised as important for plant growth processes more broadly (Berg et

al 2009), but fungal diversity beyond endosymbiotic mycorrhizae has not been systematically

investigated for grapevines. Here, we show that the soil fungal communities are distinct for a region.

Our modelling suggests that soil properties (C: N) and weather (MS) strongly affect soil fungal

diversity, which was in line with large-scale studies in which climatic (especially precipitation) and

edaphic factors (especially C: N) are the best predictors of soil fungal richness and community

composition (Lauber et al 2008, Tedersoo et al 2014). Must fungal diversity is also shaped by weather

and soil properties indirectly via the soil fungi to ultimately affect wine aroma profiles.

It is noteworthy that the drivers of microbial patterns change during wine fermentation. Microbial

diversity decreased as alcoholic fermentation proceeds, with a clear loss of microorganisms including

filamentous fungi, non-*Saccharomyces* yeasts (for example *M. guilliermondii*), spoilage bacteria

Acidobacteria and *Proteobacteria*, and other bacteria with unknown fermentative functions (for

example *Chloroflexi*), and thus the biogeographic trend was lost by the stage of MLF-End

(Supplementary Table S1). This trend was observed more distinct in fungal communities compared

with bacteria. This is not unexpected as it is clear that fermentation more strongly affects fungal

populations compared with bacteria, due to increasing fermentation rate, temperature and ethanol

concentration induced by *S. cerevisiae* growth (Bokulich et al 2012, Goddard 2008). In this case,

fermentation conditions, such as chemical environments and interactions/competition within the

community, reshape the microbial patterns. Despite the complex change of microbial ecosystem

during fermentation, we show that biogeographic patterns in the must can be translated into regional

metabolic profiling of wine. Our modelling indicates that the indirect influences of weather and soil

properties via shaping soil and must microbial diversity are more powerful than the direct influences

on wine aroma profiles (Figure 6). Co-occurrence networks demonstrate that soil and must microbiota

closely correlate to the wine metabolome. These interactions also indicate some potential modulations

of soil and must microbiota on wine chemistry, in particular that are numerically dominant taxa in the

early fermentation. For example, bacteria *Enterobacteriales* positively correlated with some grape-

derived volatile compounds, thus potentially enhancing variety aroma in wine (Swiegers et al 2005).

Enterobacteriales have been also previously related to wine fermentation rate by Bokulich et al.

(2016). Fungal genera abundance (for example *Penicillium*) negatively correlated with medium-chain

fatty acid octanoic acid, which can contribute to mushroom flavour and also alter *Saccharomyces*

metabolism (Bisson 1999). These microorganisms correlate with wine metabolites and/or can change

chemical environments and fermentation behaviours and finally exert substantial effects on wine

metabolite profiles (Swiegers et al 2015, Bokulich et al 2016). It is particularly important to inform

farming practices to structure regional microbial communities that can benefit soil quality, hence, crop

productivity.

Fungal communities distinguish wine quality and style

In grape musts, bacterial and fungal communities exhibit different responses to site-specific and

environmental effects. Bacterial regional patterns were not as distinct as fungi, and significantly

impacted by vintage (Supplementary Table S1). Although profound responses to soil properties (for

example C: N), affects wine fermentation, must bacteria show insignificant influences on wine aroma

profiles (Figure 6). In contrast, fungal communities display discriminant distribution patterns at the

regional scale and are consistent between the years studied in this work than the bacterial

communities, aligning with results presented by Bokulich et al (2014). Soil fungal communities are

less diverse than bacterial communities (Figure S2; Fierer et al 2017), but of more importance to

resulting wine regionality (see Figure 4, 5, 6). Must fungi, in particular the fermentative yeasts,

conduct alcoholic fermentations and provide aroma compounds to influence wine flavour (Swiegers et

al 2005). As indicated by SEM, soil fungal communities are affected by local soil properties and

weather and exert impacts on must fungal communities (Figure 6). Co-occurrence relationships

between wine metabolome and soil microbiota highlight potential contributions from fungal taxa

(Figure 4). One explanation is that grapevines filter soil microbial taxa selecting for grape and must

consortia (Deveau et al 2018, Panke-Buisse et al 2015). Beyond fungi, plant fitness is linked strongly

to the responses of soil microbial communities to environmental conditions (Lau and Lennon 2012).

More sensitive responses of vineyard soil fungi might improve grapevine fitness to local

environments thus benefiting the expression of regional characteristics of grapes and wines.

How could yeasts present in the soil be transported to the berry surface? Soil is a reservoir of

grapevine-associated microbiome (Figure 7) and supported by results presented by Grady et al 2019,

Mezzasalma et al 2018, Morrison-Whittle and Goddard 2018, Zarraonaindia et al 2015). As well as

transporting water and minerals absorbed by roots to the photosynthetic organs, xylem sap is also a

niche for microbes that can bear its nutritional environment (Yadeta and Thomma 2013). Here we

investigated xylem sap as a conduit to shape the microbiota on the grape surface by enrichment and

recruited by roots and transported by xylem sap to grape berries (Berg 2009, Compant et al 2008,

Zarraonaindia et al 2015). No *S. cerevisiae* yeasts were cultured from collected sap in vines in the

vineyard or in the glasshouse. We were able to isolate and identify yeasts belonging to *Cryptococcus*

and *Rhodotorula* genera, which shows that this environment is not sterile and can potentially transport

yeasts to the phyllosphere.

Fungi, while not as ecologically diverse as bacteria, are direct contributors to wine aroma. In the soil, fungi can distinguish wine regions in this study have relatively stable populations between years. While we were unable to find fermentative yeasts in the sap of grapevines, yeasts were present and thus may be transported in the grapevine as well as making their way to the phyllosphere through other mechanisms (water splashes, insect vectors). Previous studies have shown that fermentative yeasts are persistent in vineyards (Börlin et al 2016, Valero et al 2005) and can be transported by grapevines (Mandl et al 2015). We can thus conclude that fungi are the principal drivers of consistent expression of regionality in wine production.

Our study illustrates that microbes are absolute contributors to wine aroma and that this comes from the environment in which they are grown and provides a basis to explain how wine may have a distinctive flavour from a particular region. Fungi plays a crucial role in interrelating these biotic and abiotic elements in vineyard ecosystem and could be transported by the internal. Climate and soil properties profoundly drive microbial patterns from the soil to the grape must, which ultimately sculpt wine metabolic profiling. We do not yet know how grapevines recruit their microbiome to maximise physiological development and microbial diversity under local conditions.

Acknowledgements

The authors give their sincere thanks to the vignerons who kindly allowed access, enabled sampling

and provided wine samples. DL acknowledges support from a Ph.D. scholarship and funding from

Wine Australia (AGW Ph1602) and a Melbourne Research Scholarship from the University of

Melbourne.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

References

Ahlmann-Eltze C (2017). ggsignif: Significance Brackets for 'ggplot2'. *R package version 04 0*.

570
571   Austin CN, Grove GG, Meyers JM, Wilcox WF (2011). Powdery mildew severity as a function of
572   canopy density: Associated impacts on sunlight penetration and spray coverage. *American journal of*
573   *enology and viticulture* **62:** 23-31.

574
575   Barata A, Malfeito-Ferreira M, Loureiro V (2012). The microbial ecology of wine grape berries.
576   *International journal of food microbiology* **153:** 243-259.

577
578   Berg G (2009). Plant–microbe interactions promoting plant growth and health: perspectives for
579   controlled use of microorganisms in agriculture. *Applied microbiology and biotechnology* **84:** 11-18.

580
581   Bisson LF (1999). Stuck and sluggish fermentations. *American Journal of Enology and Viticulture*
582   **50:** 107-119.

583
584   Bokulich NA, Joseph CL, Allen G, Benson AK, Mills DA (2012). Next-generation sequencing
585   reveals significant bacterial diversity of botrytized wine. *PloS one* **7:** e36357.

586
Bokulich NA, Thorngate JH, Richardson PM, Mills DA (2014). Microbial biogeography of wine
grapes is conditioned by cultivar, vintage, and climate. *Proceedings of the National Academy of*
Sciences **111:** E139-E148.

Bokulich NA, Collins TS, Masarweh C, Allen G, Heymann H, Ebeler SE *et al* (2016). Associations
among Wine Grape Microbiome, Metabolome, and Fermentation Behavior Suggest Microbial
Contribution to Regional Wine Characteristics. *mBio* **7:** e00631-00616.

Börlin M, Venet P, Claisse O, Salin F, Legras J-L, Masneuf-Pomarede I (2016). Cellar-associated
Saccharomyces cerevisiae population structure revealed high diversity and perennial persistence in
Sauternes wine estates. *Applied and environmental microbiology*: AEM. 03627-03615.

Breiman L (2001). Random forests. *Machine learning* **45:** 5-32.

Burns KN, Kluepfel DA, Strauss SL, Bokulich NA, Cantu D, Steenwerth KL (2015). Vineyard soil
bacterial diversity and composition revealed by 16S rRNA genes: Differentiation by geographic
features. *Soil Biology and Biochemistry* **91:** 232-247.

Coller E, Cestaro A, Zanzotti R, Bertoldi D, Pindo M, Larger S *et al* (2019). Microbiome of vineyard
soils is shaped by geography and management. *Microbiome* **7:** 140.

Compant S, Kaplan H, Sessitsch A, Nowak J, Ait Barka E, Clément C (2008). Endophytic
colonization of Vitis vinifera L. by Burkholderia phytofirmans strain PsJN: from the rhizosphere to
inflorescence tissues. *FEMS Microbiology Ecology* **63:** 84-93.

DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K *et al* (2006). Greengenes, a
chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and
environmental microbiology* **72:** 5069-5072.

Deveau A, Bonito G, Uehling J, Paoletti M, Becker M, Bindschedler S *et al* (2018). Bacterial–fungal
interactions: ecology, mechanisms and challenges. *FEMS microbiology reviews* **42:** 335-352.

Edgar RC (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26:**
2460-2461.

Eisenhauer N, Bowker MA, Grace JB, Powell JR (2015). From patterns to causal understanding:
structural equation modeling (SEM) in soil ecology. *Pedobiologia* **58:** 65-72.

Fierer N, Jackson RB (2006). The diversity and biogeography of soil bacterial communities.
Proceedings of the National Academy of Sciences of the United States of America **103:** 626-631.

Fierer N, Leff JW, Adams BJ, Nielsen UN, Bates ST, Lauber CL *et al* (2012). Cross-biome
metagenomic analyses of soil microbial communities and their functional attributes. *Proceedings of
the National Academy of Sciences* **109:** 21390-21395.

Fortmann-Roe S (2013). Accurate, Adaptable, and Accessible Error Metrics for Predictive. *R package
version 09* **2**.

Gayevskiy V, Goddard MR (2012). Geographic delineations of yeast communities and populations
associated with vines and wines in New Zealand. *The ISME journal* **6:** 1281-1290.

637
638 Gephi: an open source software for exploring and manipulating networks. *Third international AAAI*
639 *conference on weblogs and social media*.

640
641 Gilbert JA, van der Lelie D, Zarraonaindia I (2014). Microbial terroir for wine grapes. *Proceedings of*
642 *the National Academy of Sciences* **111:** 5-6.

643
644 Gladstones J (1992). *Viticulture and environment*. Winetitles.

645
646 Goddard MR (2008). Quantifying the complexities of Saccharomyces cerevisiae's ecosystem
647 engineering via fermentation. *Ecology* **89:** 2077-2082.

648
649 Gonzálvez A, Llorens A, Cervera M, Armenta S, De la Guardia M (2009). Elemental fingerprint of
650 wines from the protected designation of origin Valencia. *Food Chemistry* **112:** 26-34.

651
652 Grace JB (2006). *Structural equation modeling and natural systems*. Cambridge University Press.

653
654 Grady KL, Sorensen JW, Stopnisek N, Guittar J, Shade A (2019). Assembly and seasonality of core
655 phyllosphere microbiota on perennial biofuel crops. *bioRxiv*: 446369.

656
657 Heard GM, Fleet GH (1985). Growth of natural yeast flora during the fermentation of inoculated
658 wines. Appl Environ Microbiol 50: 727-728.
659
660 Jackson D, Lombard P (1993). Environmental and management practices affecting grape composition
661 and wine quality-a review. *American Journal of Enology and Viticulture* **44:** 409-430.

662
663 Jara C, Laurie VF, Mas A, Romero J (2016). Microbial Terroir in Chilean Valleys: Diversity of Non-
664 conventional Yeast. *Frontiers in microbiology* **7**.

665
666 Jones DA, Wang W, Fawcett R (2009). High-quality spatial climate data-sets for Australia. *Australian*
667 *Meteorological and Oceanographic Journal* **58:** 233.

668
669 Junker BH, Schreiber F (2008). *Analysis of biological networks*, vol. 2. Wiley Online Library.

670
671 Knight S, Klaere S, Fedrizzi B, Goddard MR (2015). Regional microbial signatures positively
672 correlate with differential wine phenotypes: evidence for a microbial aspect to terroir. *Scientific*
673 *reports* **5:** 14233.

674
675 Knight SJ, Karon O, Goddard MR (2019). Small scale fungal community differentiation in a vineyard
676 system. *Food Microbiology*: 103358.

677
678 Knight SJ, Klaere S, Morrison☐Whittle P, Goddard MR (2018). Fungal diversity during fermentation
679 correlates with thiol concentration in wine. *Australian journal of grape and wine research* **24:** 105-
680 112.

681
682 Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG *et al* (2011). Bayesian
683 community-wide culture-independent microbial source tracking. *Nature methods* **8:** 761.

684
685
686 Kõljalg U, Larsson KH, Abarenkov K, Nilsson RH, Alexander IJ, Eberhardt U *et al* (2005). UNITE: a
687 database providing web□based methods for the molecular identification of ectomycorrhizal fungi.
688 *New Phytologist* **166:** 1063-1068.

689
690 Lauber CL, Strickland MS, Bradford MA, Fierer N (2008). The influence of soil properties on the
691 structure of bacterial and fungal communities across land-use types. *Soil Biology and Biochemistry*
692 **40:** 2407-2415.

693
694 Lau JA, Lennon JT (2012). Rapid responses of soil microorganisms improve plant fitness in novel
695 environments. *Proceedings of the National Academy of Sciences* **109:** 14058-14062.

696
697 Liaw A, Wiener M (2002). Classification and regression by randomForest. *R news* **2:** 18-22.

698
699 Liu D, Gao Y, Li X-X, Li Z, Pan Q-H (2015). Attenuated UV Radiation Alters Volatile Profile in
700 Cabernet Sauvignon Grapes under Field Conditions. *Molecules* **20:** 16946-16969.

701
702 Liu D, Xing R-R, Li Z, Yang D-M, Pan Q-H (2016). Evolution of volatile compounds, aroma
703 attributes, and sensory perception in bottle-aged red wines and their correlation. *European Food*
704 *Research and Technology***:** 1-12.

705
706 Liu D, Zhang P, Chen D, Howell KS (2019). From the vineyard to the winery: how microbial ecology
707 drives regional distinctiveness of wine. *Frontiers in Microbiology* **10:** 2679.

708
709 Mandl K, Schieck J, Silhavy-Richter K, Alexander P, Schneider V, Schmidt H-P (2015). Vines take
710 up yeasts from soil and transport them through the vine to the stem and skins of grapes. *Ithaka J***:** 349-
711 355.

712
713 Magoč T, Salzberg SL (2011). FLASH: fast length adjustment of short reads to improve genome
714 assemblies. *Bioinformatics* **27:** 2957-2963.

715
716 Martiny JBH, Bohannan BJ, Brown JH, Colwell RK, Fuhrman JA, Green JL *et al* (2006). Microbial
717 biogeography: putting microorganisms on the map. *Nature Reviews Microbiology* **4:** 102.

718
719 Mezzasalma V, Sandionigi A, Guzzetti L, Galimberti A, Grando MS, Tardaguila J *et al* (2018).
720 Geographical and cultivar features differentiate grape microbiota in Northern Italy and Spain
721 vineyards. *Frontiers in Microbiology* **9:** 946.

722
723 Miura T, Sánchez R, Castañeda LE, Godoy K, Barbosa O (2017). Is microbial terroir related to
724 geographic distance between vineyards? *Environmental Microbiology Reports* **9:** 742-749.

726   Morrison-Whittle P, Goddard MR (2018). From vineyard to winery: a source map of microbial
727   diversity driving wine fermentation. *Environmental microbiology*.

728
729   Müller DB, Vogel C, Bai Y, Vorholt JA (2016). The plant microbiota: systems-level insights and
730   perspectives. *annual review of genetics* **50:** 211-234.

731
732   Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ *et al* (2007). The vegan
733   package. *Community ecology package* **10:** 631-637.

734
735   Panke-Buisse K, Poole AC, Goodrich JK, Ley RE, Kao-Kniffin J (2015). Selection on soil
736   microbiomes reveals reproducible impacts on plant function. *The ISME journal* **9:** 980.

737
738   Pereira GE, Gaudillere J-P, Van Leeuwen C, Hilbert G, Lavialle O, Maucourt M *et al* (2005). 1H
739   NMR and chemometrics to characterize mature grape berries in four wine-growing areas in Bordeaux,
740   France. *Journal of agricultural and food chemistry* **53:** 6382-6389.

741
742   Pinto C, Pinho D, Cardoso R, Custódio V, Fernandes J, Sousa S *et al* (2015). Wine fermentation
743   microbiome: a landscape from different Portuguese wine appellations. *Frontiers in microbiology* **6:**
744   905.

745
746   Portillo MdC, Franquès J, Araque I, Reguant C, Bordons A (2016). Bacterial diversity of Grenache
747   and Carignan grape surface from different vineyards at Priorat wine region (Catalonia, Spain).
748   *International journal of food microbiology* **219:** 56-63.

749
750   Portillo MdC, Mas A (2016). Analysis of microbial diversity and dynamics during wine fermentation
751   of Grenache grape variety by high-throughput barcoding sequencing. *LWT-Food Science and*
752   *Technology* **72:** 317-321.

753
754   Rayment GE, Lyons DJ (2011). *Soil chemical methods: Australasia*, vol. 3. CSIRO publishing.

755
756   Reynolds AG, Heuvel JEV (2009). Influence of Grapevine Training Systems on Vine Growth and
757   Fruit Composition: A Review. *American Journal of Enology and Viticulture* **60:** 251-268.

758
759   Rillig MC, Lehmann A, Lehmann J, Camenzind T, Rauh C (2017). Soil Biodiversity Effects from
760   Field to Fork. *Trends in plant science*.

761
762   Roberts DW (2007). labdsv: Ordination and multivariate analysis for ecology. *R package version* **1**.

763
764   Robinson AL, Adams DO, Boss PK, Heymann H, Solomon PS, Trengove RD (2012). Influence of
765   geographic origin on the sensory characteristics and wine composition of Vitis vinifera cv. Cabernet
766   Sauvignon wines from Australia. *American journal of enology and viticulture***:** ajev. 2012.12023.

767
Robinson AL, Boss PK, Solomon PS, Trengove RD, Heymann H, Ebeler SE (2014). Origins of grape
and wine aroma. Part 1. Chemical components and viticultural impacts. *American Journal of Enology*
and Viticulture **65:** 1-24.

Sabate J, Cano J, Esteve-Zarzoso B, Guillamón JM (2002). Isolation and identification of yeasts
associated with vineyard and winery by RFLP analysis of ribosomal genes and mitochondrial DNA.
Microbiological research **157:** 267-274.

775
776   Schermelleh-Engel K, Moosbrugger H, Müller H (2003). Evaluating the fit of structural equation
777   models: Tests of significance and descriptive goodness-of-fit measures. *Methods of psychological*
778   *research online* **8:** 23-74.

779
780   Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS *et al* (2011). Metagenomic
781   biomarker discovery and explanation. *Genome biology* **12:** R60.

782
783   Stefanini I, Albanese D, Cavazza A, Franciosi E, De Filippo C, Donati C *et al* (2016). Dynamic
784   changes in microbiota and mycobiota during spontaneous 'Vino Santo Trentino'fermentation.
785   *Microbial biotechnology* **9:** 195-208.

786
787   Swiegers J, Bartowsky E, Henschke P, Pretorius I (2005). Yeast and bacterial modulation of wine
788   aroma and flavour. *Australian Journal of grape and wine research* **11:** 139-173.

789
790   Taylor MW, Tsai P, Anfang N, Ross HA, Goddard MR (2014). Pyrosequencing reveals regional
791   differences in fruit☐associated fungal communities. *Environmental microbiology* **16:** 2848-2858.

792
793   Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R *et al* (2014). Global diversity
794   and geography of soil fungi. *science* **346:** 1256688.

795
796   Van Leeuwen C, Friant P, Chone X, Tregoat O, Koundouras S, Dubourdieu D (2004). Influence of
797   climate, soil, and cultivar on terroir. *American Journal of Enology and Viticulture* **55:** 207-217.

798
799   Van Leeuwen C, Seguin G (2006). The concept of terroir in viticulture. *Journal of Wine Research* **17:**
800   1-10.

801
802   Vaudour E, Costantini E, Jones G, Mocali S (2015). An overview of the recent approaches to terroir
803   functional modelling, footprinting and zoning. *Soil* **1:** 287.

804
805   Valero E, Schuller D, Cambon B, Casal M, Dequin S (2005). Dissemination and survival of
806   commercial wine yeast in the vineyard: A large-scale, three-years study. *FEMS Yeast Research* **5:**
807   959-969.

808
809   Wang Q, Garrity GM, Tiedje JM, Cole JR (2007). Naive Bayesian classifier for rapid assignment of
810   rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* **73:** 5261-5267.

811
812   Yadeta K, Thomma B (2013). The xylem as battleground for plant hosts and vascular wilt pathogens.
813   *Frontiers in plant science* **4:** 97.

814
815   Zarraonaindia I, Owens SM, Weisenhorn P, West K, Hampton-Marcell J, Lax S *et al* (2015). The soil
816   microbiome influences grapevine-associated microbiota. *MBio* **6:** e02527-02514.

818   Zhang P, Barlow S, Krstic M, Herderich M, Fuentes S, Howell K (2015). Within-vineyard, within-
819   vine, and within-bunch variability of the rotundone concentration in berries of Vitis vinifera L. cv.
820   Shiraz. *Journal of agricultural and food chemistry* **63:** 4276-4283.

Figures

Figure 1 Wine metabolome shows regional variation across six wine growing regions. Shown are α-diversity (Shannon index) (A) and PCoA comparisons of Bray-Curtis dissimilarity (B) of wine volatile profiles.

Figure 2 Vineyard soil microbial communities demonstrate distinct regional patterns. Bray-Curtis distance PCoA of bacterial communities (A) and fungal communities (B).

Figure 3 Microbiota exhibits regional differentiation in musts for both bacterial and fungal profiles. Stage of fermentation influences microbial composition of bacteria and fungi. (A) Must bacterial taxa in the order level with greater than 1.0% relative abundance, and *Lactobacillales*. (B) Must fungal taxa in the genus level with greater than 1.0% relative abundance. (C and D) LDA effect size taxonomic cladogram comparing all musts and wines categorised by fermentation stage. Significantly discriminant taxon nodes are coloured and branch areas are shaded according to the highest-ranked stage for that taxon. For each taxon detected, the corresponding node in the taxonomic cladogram is coloured according to the highest-ranked group for that taxon. If the taxon is not significantly differentially represented between sample groups, the corresponding node is coloured yellow.

Figure 4 Statistically significant and strong co-occurrence relationships between wine volatile compounds and bacterial and fungal OTUs in vineyard soils and musts across six wine growing regions. Network plots for wine aroma with soil and must microbiota (A), and simplified network plots for wine aroma with soil microbiota (B) and with must microbiota (C), separately. Circle nodes represent assigned volatile compounds, bacterial and fungal OTUs, with different colours. Direct connections between compounds and OTUs indicate strong correlations (Spearman correlation coefficient, $\rho \geq 0.8$; $p < 0.01$). The colour of edges describes the positive correlation (pink) or the negative correlation (green). The size of nodes is proportioned to the inconnected degree.

Figure 5 Main predictors of wine regionality. Shown is the Random Forest mean predictor importance (% of increase in MSE) of climate, soil properties and microbial diversity (Shannon index) on wine regionality. Soil bac. div., soil bacterial diversity; Soil fung. div., soil fungal diversity; Must bac. div., must bacterial diversity; Must fung. div., must fungal diversity. Significance levels: *$p < 0.05$, **$p < 0.01$.

Figure 6 Direct and indirect effects of climate, soil properties and microbial diversity (Shannon index) on wine regionality. Structural equation model (SEM) fitted to the diversity of wine aroma profiles (A) and standardized total effects (direct plus indirect effects) derived from the model (B). Climate and soil properties are composite variables encompassing multiple observed parameters (see materials and methods for the complete list of factors used to generate this model). Numbers adjacent to arrows are path coefficients and indicative of the effect size of the relationship. The width of arrows is proportional to the strength of path coefficients. R^2 denotes the proportion of variance explained. (a) (0.75* MT) (0.67* MLT) (0.69* MinT) (-0.88** MS). (b) (0.75* MinT) (0.77* MLT) (-0.68* MS) (-0.74* MHT) (-0.84** MaxT). C, soil carbon; N, soil nitrogen; C:N, soil carbon nitrogen ratio; MS, mean solar radiation; MT, mean temperature; MLT, mean low temperature; MHT, mean high temperature; MinT, minimum temperature; MaxT, maximum temperature; MTrans, mean transpiration. Significance levels: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 7 Fungal communities in musts emerge from multiple sources in the vineyard, but primarily from grape and xylem sap. Percent composition estimate for the contributions of possible sources in the vineyard to the wine-related fungal community for sourcetracking must (A), grape (B) and xylem sap (C).

Supplementary Table and Figures

Table S1 Permuted multivariate analysis of variance (PERMANOVA) of region and vintage effects on microbial patterns and wine profiles (beta-diversity)

Sample Group	Vintage	Factor	Soil Bacteria R2	p	Soil Fungi R2	p	Must Bacteria R2	p	Must Fungi R2	p	Wine (MLF-End) Bacteria R2	p	Wine (MLF-End) Fungi R2	p	Wine Aroma MLF-End R2	p
All	Both	Region	0.318	0.001	0.254	0.001	0.108	0.152	0.292	0.001	N/A	N/A	N/A	N/A	0.566	0.001
All	2017	Region	0.392	0.001	0.419	0.001	0.248	0.013	0.565	0.001	0.149	0.321	0.109	0.205	0.703	0.001
All	Both	Vintage	0.102	0.001	0.086	0.001	0.186	0.001	0.049	0.066	N/A	N/A	N/A	N/A	0.267	0.001
Mornington	Both	Vintage	0.155	0.001	0.151	0.001	0.305	0.001	0.143	0.001	N/A	N/A	N/A	N/A	0.355	0.001

Figure S1 Map of 15 sampling vineyards from six Pinot Noir wine-producing regions in southern Australia, spanning 400 km (E-W).

Figure S2 Vineyard soil microbial community structures across all samples from six grape growing regions. Microbial community compositions in soil samples are characterized to the phylum level: (A) dominant bacterial taxa with greater than 1.0% relative abundance (top 10 phyla); (B) fungal phyla.

Figure S3 Stage of fermentation influences microbial diversities. Bacterial (A) and fungal (B) α-diversity (Shannon index) changes during wine fermentation. Bray-Curtis distance PCoA of bacterial communities (C) and fungal communities (D) according to the fermentation stage.

Figure S4 Xylem sap is a potential translocation medium of yeasts from the soil to the grapevine. (A) Fungal taxa in the genus level of xylem sap from the vineyard. (B) *S. cerevisiae* shared between niches of soil, root, xylem sap, grape, and must. Cultured yeast species from the xylem sap from vineyard (C) and glasshouse (D). Nutrient compositions of xylem sap: carbohydrates (E), total amino acids (protein and free) (F), and organic acids (G).

Supplementary Materials and Methods

Xylem sap collection

Xylem sap (n = 10) was collected from the vineyard using a centrifugation method in aseptic conditions (López-Millán et al 2000). Briefly, the shoots were peeled the outer bark and phloem layer, sterilised with 70% ethanol and cut to fit into 10-ml sterile centrifuge tubes that had 10 glass beads at the bottom. Xylem sap was centrifuged out the shoot pieces at $15,000 \times g$ for 10 min at 4°C and collected at the bottom (~ 2.0 mL). Additional xylem sap (n = 5) were sampled from *Vitis vinifera* Shiraz grapevines grown in the glasshouse, the University of Melbourne. Xylem fluid was extracted with a pressure cylinder apparatus (similar with Scholander pressure chamber) (Schurr 1998). Grapevines were uprooted and carefully removed the soil from roots. In an aseptic bench, the main stems were cut 5-10 cm above roots with a sterile blade, girdled to remove phloem tissue, sterilised by 70% ethanol, and immediately inserted into the pressure cylinder. The cylinder was applied 60-70 kPa pressure for two hours to extract xylem sap (~ 4.0 mL). Xylem sap was divided into two subsamples, one for yeast culture experiment immediately, one for next-generation sequencing with flash frozen in liquid nitrogen and stored at -80°C until DNA extraction.

Chemical analysis of xylem sap composition

Carbohydrates of xylem sap were determined using enzymatic methods by Megazyme assay kits (Megazyme, Ireland) following the manufacturer's protocol. Amino acids (free and proteins) were determined using pre-column derivatisation with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate followed by separation and quantification with an ACQUITY Ultra Performance LC (UPLC; Waters, MA, USA) system by Australian Proteome Analysis Facility. The column employed was an ACQUITY UPLC BEH C18 column (1.7 μm × 2.1 mm × 5 mm) with detection at 260 nm (UV) and a flow rate of 0.7 mL/min at 57 – 60°C. Identification and quantitation were using a set of prepared standards, with DL-Norvaline as the internal standard (Cohen and Michaud 1993, Cohen 2001). Organic acids were determined using a Waters High-performance liquid chromatography (HPLC) (Waters, MA, USA) based on Andersen et al (1989) with modification. 20 μL xylem sap was injected

through a Synergi™ Hydro-RP LC Column (250 mm × 4.6 mm × 4 μm; Phenomenex Inc, CA, USA) at 60°C, with detection at 210 nm (UV). Mobile phases at a flow rate of 1.0 mL/min, with 20 mM potassium phosphate buffer (A, pH = 1.5) and 100% methanol (B) following a gradient programme: (0-2.5) min, 100% A; (2.5-2.9) min, linear ramp to 30% B; (2.9-8.0) min, 30% B; (8.0-8.5) min, linear ramp to 100% A; (8.5-10) min, 100% A. Identification and quantitation were using a set of prepared standards.

Isolation and identification of yeasts from xylem sap

Yeasts were isolated and identified from xylem sap to explore the potential translocation mechanism of yeasts in the vineyard ecosystem. Xylem sap was serially diluted and plated (0.1 mL) onto the yeast extract peptone dextrose (YPD) medium that was supplemented with 34 mg/mL chloramphenicol and 25 mg/mL ampicillin to inhibit bacterial growth. Plates were incubated at 28°C for 2-3 days in aerobic conditions. Single colonies with different morphological types were streaked onto Wallerstein Nutrient (WLN) agar media to obtain pure cultures, which were stored in 15% glycerol at -80°C. DNA was recovered from pure colonies using the MasterPure™ Yeast DNA Purification Kit (Epicentre, Madison, WI) following the manufacturer's instructions. The 26S rDNA D1/D2 domain was amplified using primers NL1/4 (Kurtzman and Robnett 1998) for sequencing by Australian Genome Research Facility (AGRF). Sequences were trimmed, aligned and analysed using BLAST at NCBI (http://blast.ncbi.nlm.nih.gov/Blast.cgi).

References

Andersen PC, Brodbeck BV, Mizell RF (1989). Metabolism of amino acids, organic acids and sugars extracted from the xylem fluid of four host plants by adult Homalodisca coagulata. *Entomologia Experimentalis et Applicata* **50:** 149-159.

Cohen SA, Michaud DP (1993). Synthesis of a fluorescent derivatizing reagent, 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate, and its application for the analysis of hydrolysate amino acids via high-performance liquid chromatography. *Analytical biochemistry* **211:** 279-287.

Cohen SA (2001). Amino acid analysis using precolumn derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. *Amino Acid Analysis Protocols*. Springer. pp 39-47.

Kurtzman CP, Robnett CJ (1998). Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie van Leeuwenhoek* **73:** 331-371.

López-Millán AF, Morales F, Abadía A, Abadía J (2000). Effects of iron deficiency on the composition of the leaf apoplastic fluid and xylem sap in sugar beet. Implications for iron and carbon transport. *Plant Physiology* **124:** 873-884.

Schurr U (1998). Xylem sap sampling—new approaches to an old topic. *Trends in Plant Science* **3:** 293-298.