Title: Vineyard ecosystems are structured and distinguished by fungal communities

impacting the flavour and quality of wine

Running title: Environmental fungi determine the flavour of wine

Di Liu, Qinglin Chen, Pangzhen Zhang, Deli Chen and Kate S. Howell*

School of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, University of

Melbourne Parkville 3010 Australia

*Corresponding author. khowell@unimelb.edu.au

Abstract

The flavours of foods and beverages are formed by the agricultural environment where the plants are grown. In the case of wine, the location and environmental features of the vineyard site imprint the wine with distinctive aromas and flavours. Microbial growth and metabolism play an integral role in wine production from the vineyard to the winery, by influencing grapevine health, wine fermentation, and the flavour, aroma and quality of finished wines. The mechanism by which microbial distribution patterns drive wine metabolites is unclear and while flavour has been correlated with bacterial composition for red wines, bacterial activity provides a minor biochemical conversion in wine fermentation. Here, we collected samples across six distinct winegrowing areas in southern Australia to investigate regional distribution patterns of both fungi and bacteria and how this corresponds with wine aroma compounds. Results show that soil and must microbiota distinguish winegrowing regions and are related to wine chemical profiles. We found a strong relationship between microbial and wine metabolic profiles, and this relationship was maintained despite differing abiotic drivers (soil properties and weather/ climatic measures). Notably, fungal communities played the principal role in shaping wine aroma profiles and regional distinctiveness. We found that the soil microbiome is a potential source of grape- and must-associated fungi, and therefore the weather and soil conditions could influence the wine characteristics via shaping the soil fungal community compositions. Our study describes a comprehensive scenario of wine microbial biogeography in which microbial diversity responds to surrounding environments and ultimately sculpts wine aromatic characteristics. These findings provide perspectives for thoughtful human practices to optimise food and beverage flavour and composition through understanding of fungal activity and abundance.

Keywords: wine regionality, microbial biogeography, fungal diversity, climate, soil

Introduction

Regional distinctiveness of wine traits, collectively known as "*terroir*", can be measured by chemical composition and sensory attributes (1-3) , and this variation has been related to the physiological response of grapevines to local environments, such as soil properties (e.g., soil type, texture, nutrient availability),

climate (temperature, precipitation, solar radiation), topography and human-driven agricultural practices

(4-6). Wines made from the same grape cultivar but grown in different regions are appreciated for their

regional diversity, increasing price premiums and the market demand (5). However, which factors from

the vineyard to the winery drive wine quality traits have remained elusive.

Microbial biogeography contributes to the expression of wine regionality. Microorganisms, including

yeasts, filamentous fungi and bacteria, originate in the vineyard and are impacted upon by the built

environment (winery) and play a decisive role in wine production and quality of the final product (7, 8).

The fermentative conversion of grape must (or juice) into wine is a complex and dynamic process,

involving numerous transformations by multiple microbial species (9). The majority of fermentations

involve *Saccharomyces* yeasts conducting alcoholic fermentation (AF) and lactic acid bacteria (LAB) for

malolactic fermentation (MLF), but many other species are present and impact the chemical composition

of the resultant wine (10, 11). Recent studies propose the existence of non-random geographical

dispersion patterns of microbiota in grapes and wines (12-18). Few studies have explored the associations

between microbial communities and wine chemical composition (19, 20). Knight et al. (20) showed

empirically that regionally distinct *S. cerevisiae* populations drove different metabolites present in the

resultant wines. Bokulich et al. (19) suggested that wine metabolite profiles were more closely associated

with bacterial profiles of wines undergone both AF and MLF. However, both numerically and

biochemically, yeasts dominate wine fermentation, so it is unclear why bacteria should be dominant.

The vineyard soil has long been attributed to fundamental wine characteristics and flavour. Vineyard soil

provides the grapevine with water and nutrients and soil type and properties profoundly affect vine

growth and development (5). Soil-borne microbiota associate with grapevines in a beneficial, commensal,

or pathogenic way, and determine soil quality, host growth and health. For example, soil microbes can

mineralize soil organic matter, trigger plant defence mechanisms, and thus influence flavour and quality

of grapes and final wines (21, 22). Alternatively, serving as a reservoir for grapevine-associated

microbiota, the soil microbiome can be recruited from the soil to the grapevine and transferred to the

grape and must (14, 23). Some of these microbes can influence the fermentation and contribute to final

wine characteristics (8, 24). Overall, biogeographic boundaries can constrain the vineyard soil microbiota

(23, 25-28), but correlations between geographical structure of soil microbiota and wine attributes are

weak.

Limited but increasing evidence reveals that environmental heterogeneity conditions microbial

biogeography in wine production on different spatial scales [recently reviewed by Liu et al. (29); (12, 23,

25, 30, 31). Local climatic conditions significantly correlate with microbial compositions in grape musts,

for example precipitation and temperature that correlate with the abundance of filamentous fungi (for

example *Cladosporium* and *Penicillium* spp.) and ubiquitous bacteria (for example, the

Enterobacteriaceae family) (12), as well as yeast populations (particularly *Hanseniaspora* and

Metschnikowia spp.) (30). Dispersal of soil microbiota is driven by soil physicochemical properties such

as soil texture, soil pH, carbon (C) and nitrogen (N) pools (23, 25, 26), with some influences from

topological characteristics (for example orientation of the vineyard) (25, 32). Soil microbiome/ bacteria

may colonise grapes by physical contact (moving by rain splashes, dust, winds) (23) or migration through

the plant (xylem/phloem) from the rhizosphere to the phyllosphere (33, 34). Insects help the movement

and dispersal of microbes in the vineyard and winery ecosystem, for example, honey bees, social wasps

and *Drospholid* flies can vector yeasts among different microhabitats (35-37). The vineyard microbes

enter the winery associated with grapes or must, so the effects of environmental conditions are finally

reflected on microbial consortia in wine fermentation. How environmental conditions modulate microbial

ecology from the vineyard to the winery and shape regional distinctiveness of wine is still largely

unknown.

Here, we sampled microbial communities from the vineyard to the winery across six geographically

separated wine-producing regions in southern Australia to provide insights into microbial biogeography,

growing environments and wine regionality. We evaluated the volatile chemicals of wines made with

Pinot Noir grapes to validate that these different regions have differently flavoured wines. Using next-

generation sequencing (NGS) to profile bacterial and fungal communities, we demonstrate that the soil

and must microbiota exhibit distinctive regional patterns, and this correlated to the wine metabolome.

Network analysis supports microbial community composition that correlated with abiotic factors (weather

and soil properties) drives wine regionality both directly and indirectly. The important contributions arise

from fungal diversity. We investigated a potential route of wine-related fungi from the soil to the grapes

by isolating yeasts from the xylem/phloem of grapevines. Using vineyards, grapes and wine as a model

food system, we have related the regional identity of an agricultural commodity to biotic components in

the growing system to show the importance of conserving regional microbial diversity to produce

distinctive foods and beverages.

Materials and methods

Sample sites and weather parameters

15 *Vitis vinifera* cv. Pinot Noir vineyards were selected in 2017 from Geelong, Mornington Peninsula

(Mornington), Macedon Ranges, Yarra Valley, Grampians and Gippsland in southern Australia, ranging

from 5 km to 400 km between vineyards (Supplementary Figure S1). In 2018, the sampling from the five

vineyards in Mornington Peninsula (all < 20 km apart) was repeated to elucidate the influence of

sampling year (vintage) on microbial patterns and wine profiles. Each site's GPS coordinates (longitude,

latitude, altitude) were utilised to extract weekly weather data from the dataset provided by Australian

Water Availability Project (AWAP). Variables were observed by robust topography resolving analysis

methods at a resolution of $0.05° \times 0.05°$ (approximately 5 km x 5 km) (38). Weekly measurements for all

vineyards were extracted for mean solar radiation (MS), mean high temperature (MHT), mean low

temperature (MLT), maximum temperature (MaxT), minimum temperature (MinT), mean temperature

(MT), precipitation, mean relative soil moisture, mean evaporation (soil + vegetation), and mean

transpiration (MTrans) in growing seasons (October 2016/ 2017 - April 2017/ 2018).

Collection of soil, plant, must and ferment samples

In each vineyard, soil samples were collected from three sites at harvest March-April 2017 (n = 45) and

2018 (n = 15), at a depth of 0 - 15 cm, 30 – 50 cm from the grapevine into the interrow (Supplementary

Table S1A). To further investigate fungal ecology in the vineyard, comprehensive vineyard samples (n =

50) were collected from two vineyards 5 km apart in the 2018 vintage (Supplementary Table S1B). These

two vineyards were managed by the same winery, and the viticultural management was very similar, for

example, grapevines were under vertical shoot positioning trellising systems and the same sprays were

applied at the same time of year. Five replicate Pinot Noir vines in each vineyard were selected from the

top, middle, and bottom of the dominant slope, covering topological profiles of the vineyard. For each

grapevine, five different sample types were collected: soil (0 - 15 cm deep, root zone), roots, xylem/

phloem sap, leaves, and grapes were collected at harvest in March 2018. Xylem sap (n = 10) was

collected from the shoots using a centrifugation method in aseptic conditions (39) (Supplementary Table

S1B). Details of xylem sap collection, nutrient composition analysis, and yeast isolation were provided in

supplementary materials. Samples were stored in sterile bags, shipped on ice, and stored at -80 °C until

processing.

Longitudinal samples to study microbial communities during fermentation were collected at five time

points: must (destemmed, crushed grapes prior to fermentation), at early fermentation (AF, less than 10%

of the sugar is fermented), at middle of fermentation (AF-Mid, around 50% of sugar fermented), at the

end of fermentation (AF-End, following pressing but prior to barrelling), at the end of malolactic

fermentation (MLF-End, in barrels) (Supplementary Table S1A). Grapes and must sampled in this study

were fermented in the wineries following similar fermentation protocols of three days at a cool

temperature (known as cold-soaking) followed by warming up the must, so fermentation could

commence. Fermentations were conducted without addition of commercial yeasts and bacteria. Two

fermentations from Grampians and Mornington did not complete and were excluded from analysis, giving

wine samples from 13 vineyards in the 2017 vintage. Triplicate sub-samples from tanks or barrels were

collected and mixed as composite samples. All samples (n = 90) were shipped on ice and stored at -20 °C

until processing.

Soil analysis

Edaphic factors were analysed to explore the effects of soil properties on wine-related microbiota and

aroma profiles. Soil pH was determined in a 2:5 soil/water suspension. Soil carbon (C), nitrogen (N),

nitrate and ammonium were analysed by Melbourne Trace Analysis for Chemical, Earth and

Environmental Sciences (TrACEES) Soil Node, at the University of Melbourne. Total C and N were

determined using classic Dumas method of combustion (40) on a LECO TruMac CN at a furnace

temperature of 1350°C (LECO Corporation, Michigan, USA). Nitrate and ammonium were extracted with

2 M KCl and determined on a Segmented Flow Analyzer (SAN++, Skalar, Breda, Netherlands) (40).

Wine volatile analysis

To represent the wine aroma, volatile compounds of MLF-End samples were determined using headspace

solid-phase microextraction gas-chromatographic mass-spectrometric (HS-SPME–GC-MS) method (41,

42) with some modifications. Analyses were performed with Agilent 6850 GC system and a 5973 mass

detector (Agilent Technologies, Santa Clara, CA, USA), equipped with a PAL RSI 120 autosampler

(CTC Analytics AG, Switzerland). Briefly, 10 mL wine was added to a 20 ml glass vial with 2 g of

sodium chloride and 20 μL of internal standard (4-Octanol, 100 mg/L), then equilibrated at 35□°C for

15□min. A polydimethylsiloxane/divinylbenzene (PDMS/DVB, Supelco) 65 μm SPME fibre was

immersed in the headspace for 10 min at 35°C with agitation. The fibre was desorbed in the GC injector

for 4 min at 220 °C. Volatiles were separated on an Agilent J&W DB-Wax Ultra Inert capillary GC

column (30 m × 0.25 mm × 0.25 μm), with helium carrier gas at a flow rate of 0.7 mL/min. The column

temperature program was as follows: holding 40 °C for 10 min, increasing at 3.0 °C/min to 220 °C and

holding at this temperature for 10 min. The temperature of the transfer line of GC and MS was set at 240

°C. The ion source temperature was 230 °C. The electron impact (EI) at 70 eV scanning over a mass

acquisition range of 35–350 *m/z*. Raw data were analysed with Agilent ChemStation Software for

qualification and quantification (43). Volatile compounds (n = 88) were identified in wine samples

according to retention indices reference standards and mass spectra matching with NIST11 library. 13

successive levels of standard solution in model wine solutions (12% v/v ethanol saturated with potassium

hydrogen tartrate and adjusted to pH 3.5 using 40% w/v tartaric acid) were analysed by the same protocol

as wine samples to establish the calibration curves for quantification. Peak areas of volatile compounds

were integrated via target ions model. The concentrations of volatile compounds were calculated with the

calibration curves and used for downstream data analysis.

DNA extraction and sequencing

Genomic DNA was extracted from plant and soil samples using PowerSoil™ DNA Isolation kits

(QIAgen, CA, USA). DNA extraction from soil and xylem sap followed the kit's protocols. Wine

fermentation samples were thawed and biomass was recovered by centrifugation at $4,000 \times g$ for 15 min,

washed three times in ice-cold phosphate buffered saline (PBS) with 1% polyvinylpolypyrrolidone

(PVPP) and centrifuged at $10,000 \times g$ for 10 min (12). The obtained pellets were used for DNA

extraction following the kit protocol. For grapevine samples, roots, leaves, grapes (removed seeds and

stems), were ground into powder under the protection of liquid nitrogen with 1% PVPP, and isolated

DNA following the kit protocol afterward. DNA extracts were stored at -20 °C until further analysis.

Genomic DNA was submitted to Australian Genome Research Facility (AGRF) for amplification and

sequencing. To assess the bacterial and fungal communities, 16S rRNA gene V3-V4 region and partial

fungal internal transcribed spacer (ITS) region were amplified using the universal primer pairs 341F/806R

(44) and ITS1F/2 (45), respectively. The primary PCR reactions contained 10 ng DNA template, 2×

AmpliTaq Gold® 360 Master Mix (Life Technologies, Australia), 5 pmol of each primer. A secondary

PCR to index the amplicons was performed with TaKaRa Taq DNA Polymerase (Clontech).

Amplification were conducted under the following conditions: for bacteria, 95 °C for 7 min, followed by

30 cycles of 94 °C for 30 s, 50 °C for 60 s, 72 °C for 60 s and a final extension at 72 °C for 7 min; for

fungi, 95 °C for 7 min, followed by 35 cycles of 94 °C for 30 s, 55 °C for 45 s, 72 °C for 60 s and a final

extension at 72 °C for 7 min. PCR products were purified, quantified and pooled at the same

concentration (5 nM). The resulting amplicons were cleaned again using magnetic beads, quantified by

fluorometry (Promega Quantifluor) and normalised. The equimolar pool was cleaned a final time using

magnetic beads to concentrate the pool and then measured using a High-Sensitivity D1000 Tape on an

Agilent 2200 TapeStation. The pool was diluted to 5nM and molarity was confirmed again using a High-

Sensitivity D1000 Tape. This was followed by 300 bp paired-end sequencing on an Illumina MiSeq (San

Diego, CA, USA).

Raw sequences were processed using QIIME v1.9.2 (46). Low quality regions (Q < 20) were trimmed

from the 5′ end of the sequences, and the paired ends were joined using FLASH (47). Primers were

trimmed and a further round of quality control was conducted to discard full length duplicate sequences,

short sequences (< 100 nt), and sequences with ambiguous bases. Sequences were clustered followed by

chimera checking using UCHIME algorithm from USEARCH v7.1.1090 (48). Operational taxonomic

units (OTUs) were assigned using UCLUST open-reference OTU-picking workflow with a threshold of

97% pairwise identity (48). Singletons or unique reads in the resultant data set were discarded, in

addition, chloroplast and mitochondrion related reads were removed from the OTU dataset for 16S rRNA.

Taxonomy was assigned to OTUs in QIIME using the Ribosomal Database Project (RDP) classifier (49)

against GreenGenes bacterial 16S rRNA database (v13.8) (50) or the UNITE fungal ITS database (v7.2)

(51), respectively. To avoid/ reduce biases generated by varying sequencing depth, sequences were

rarefied to an even depth per sample (the lowest sequencing depth of each batch, that is for the soil, must

and wine, soil and plant samples) prior to downstream analysis. Raw data are publicly available in the

National Centre for Biotechnology Information Sequence Read Archive under the bioproject

PRJNA594458 (bacterial 16S rRNA sequences) and PRJNA594469 (fungal ITS sequences).

Data analysis

Microbial alpha-diversity was calculated using the Shannon index in R (v3.5.0) with the "vegan" package

(52). One-way analysis of variance (ANOVA) was used to determine whether sample classifications (e.g.,

region, fermentation stage) contained statistically significant differences in the diversity. Principal

coordinate analysis (PCoA) was performed to evaluate the distribution patterns of wine metabolome and

wine-related microbiome based on beta-diversity calculated by the Bray–Curtis distance with the "labdsv"

package (53). Permutational multivariate analysis of variance using distance matrices with 999

permutations was conducted within each sample classification to determine the statistically significant

differences with "adonis" function in "vegan" (52).

Significant differences of wine microbiome between fermentation stages were tested based on taxonomic

classification using linear discriminant analysis (LDA) effect size (LEfSe) analysis (54)

(https://huttenhower.sph.harvard.edu/galaxy/). The OTU table was filtered to include only OTUs > 0.01%

relative abundance to reduce LEfSe complexity. This method applies the factorial Kruskal–Wallis sum-

rank test ($\alpha = 0.05$) to identify taxa with significant differential abundances between categories (using all-

against-all comparisons), followed by the logarithmic LDA score (threshold = 2.0) to estimate the effect

size of each discriminative feature. Significant taxa were used to generate taxonomic cladograms

illustrating differences between sample classes.

The co-occurrence/interaction patterns between wine metabolome and microbiota in the soil and must

were explored in network analysis using Gephi (v0.9.2) (55). Only top 100 OTUs (both bacteria and

fungi) based on the relative abundance were used to construct the network. A correlation matrix was

calculated with all possible pair-wise Spearman's rank correlations between volatile compounds and

selected OTUs. Correlations with a Spearman correlation coefficient $\rho \geq 0.8$ and a $p < 0.01$ were

considered statistically robust and displayed in the networks (56).

A Random forest supervised-classification model (57) was conducted to identify the main predictors of

wine regionality among the following variables: must and soil microbial diversity (Shannon index), soil

properties, and weather. The importance of each predictor is determined by evaluating the decrease in

prediction accuracy (that is, increase in the mean square error (MSE) between observations and out-of-

bag predictions) when the data is randomly permuted for the predictor. This analysis was conducted with

5,000 trees using the "randomForest" package in R (58). The significance of the model and the cross-

validated R^2 were assessed using the "A3" package (ntree = 5000) (59). The structural equation model

(SEM) (60) was used to evaluate the direct and indirect relationships between must and soil microbial

diversity, soil properties, climate and wine regionality. SEM is an *a priori* approach partitioning the

influence of multiple drivers in a system to help characterize and comprehend complex networks of

ecological interactions (61). An *a priori* model was established based on the known effects and

relationships among these drivers of regional distribution patterns of wine aroma to manipulate the data

before modelling. Weather and soil properties were used as composite variables (both Random Forest and

SEM) to collapse the effects of multiple conceptually-related variables into a single-composite effect, thus

aiding interpretation of model results (60). A path coefficient describes the strength and sign of the

relationship between two variables (60). The good fit of the model was validated by the χ^2-test ($p > 0.05$),

using the goodness of fit index (GFI > 0.90) and the root MSE of approximation (RMSEA < 0.05) (62).

The standardised total effects of each factor on the wine regionality pattern were calculated by summing

all direct and indirect pathways between two variables (60). All the SEM analyses were conducted using

AMOS v25.0 (AMOS IBM, NY, USA).

SourceTracker was used to track potential sources of wine-related fungi within the vineyards (63).

SourceTracker is a Bayesian approach that treats each give community (sink) as a mixture of

communities deposited from a set of source environments and estimates the proportion of taxa in the sink

community that come from possible source environments. When a sink contains a mixture of taxa that do

not match any of the source environments, that portion of the community is assigned to an "unknown"

source (63). In this model, we examined musts (n = 2) and vineyard sources (n = 50) including grapes,

leaves, xylem sap, roots, and soils. The OTU tables were used as data input for modelling using the

"SourceTracker" R package (https://github.com/danknights/sourcetracker).

Results

Chemical composition/ aroma profiles separate wines based on geographic location

Using GC-MS, we analysed the volatile compounds of Pinot Noir wine samples (MLF-End) to represent

wine metabolite profiles coming from different growing regions and compared directly to the microbial

communities inhabiting the musts from which these wines were fermented. In all, 88 volatile compounds

were identified in these wines, containing 48 regionally differential compounds (see Table S2 in the

supplementary material). Here we used α- and β-diversity measures to further elucidate wine complexity

and regionality, respectively. In 2017, α-diversity varied with regional origins (ANOVA, F = 36.021, $p <$

0.001), with higher Shannon indices observed in the wines from regions of Mornington, Yarra Valley and

Gippsland (H = 2.17 ± 0.05) than others (H = 1.94 ± 0.03) (Figure 1A). Overall, wine aroma profiles

displayed significant regional differentiation across both vintages based on Bray–Curtis dissimilarity

(ADONIS, R^2 = 0.566, $p < 0.001$) and the clustering patterns became more distinct and the coefficient of

determination (R^2) improved when comparing regional differences in 2017 vintage (ADONIS, R^2 =

0.703, $p < 0.001$) (Supplementary Table S3). PCoA showed that 74.5% of the variance was explained by

the first two principal coordinates in 2017, and on PCo1 there were some wines within regions grouped

together (Figure 1B).

Microbial ecology from the vineyard to the winery

To elucidate how microbial ecology drives regional traits of wine from the vineyard to the winery, 150

samples covering soils, musts and fermentations were collected to analyse wine-related microbiota. A

total of 11,508,480 16S rRNA and 12,403,610 ITS high-quality sequences were generated from all the

samples, which were clustered into 13,689 bacterial and 8,373 fungal OTUs with a threshold of 97%

pairwise identity, respectively.

The dominant bacterial taxa across all soil samples were *Actinobacteria*, *Proteobacteria*, *Acidobacteria*,

Chloroflexi, *Verrucomicrobia*, *Bacteroidetes*, *Gemmatimonadetes*, *Firmicutes*, *Planctomycetes* and

Nitrospirae (Supplementary Figure S2A). Compared with bacteria, soil fungal communities were less

diverse (Supplementary Table S4). *Ascomycota* was the most abundant and diverse phylum of fungi,

accounting for 72% of reads, followed by *Basidiomycota*, *Mortierellomycota*, *Chytridiomycota* and

Olpidiomycota (Supplementary Figure S2B). The microbiome richness (α-diversity, Shannon index)

varied significantly with regions for both bacteria and fungi (ANOVA, $F_{bacteria}$ = 4.645, $p < 0.01$; F_{fungi} =

4.913, $p < 0.01$). Soil microbial communities varied widely across different grape growing regions,

exerting significant impacts on both bacterial and fungal taxonomic dissimilarity based on Bray-Curtis

distances matrices at OTU level (ADONIS, $R^2_{bacteria} = 0.318$, $p < 0.001$; $R^2_{fungi} = 0.254$, $p < 0.001$), with

clearer differences within a single vintage (ADONIS, $R^2_{bacteria\ 2017} = 0.392$, $p < 0.001$; $R^2_{fungi\ 2017} = 0.419$, p

< 0.001) (Supplementary Table S3). In 2017, soil fungal communities could discriminate growing regions

(except Yarra Valley and Gippsland) (Figure 2B), whereas regional separation of bacteria was weaker

with overlap between regions (Figure 2A).

In grape musts, bacterial communities across six wine growing regions in both vintages consisted of

ubiquitous bacteria *Enterobacteriales*, *Rhizobiales*, *Burkholderiales*, *Rhodospirillales*, *Actinomycetales*,

Sphingomonadales, *Pseudomonadales*, *Saprospirales*, and *Xanthomonadales*, which do not participate in

wine fermentations or spoilage (7). The LAB *Lactobacillales*, responsible for malolactic fermentation,

were present in low abundance (0.4% on average) in the must (Figure 3A). Fungal profiles were

dominated by filamentous fungi, mostly of the genera *Aureobasidium*, *Cladosporium*, *Botrytis*,

Epicoccum, *Penicillium*, *Alternaria*, and *Mycosphaerella*, with notable populations of yeasts, including

Saccharomyces, *Hanseniaspora*, and *Meyerozyma*, as well as the *Basidiomycota* genus *Rhodotorula*

(Figure 3B). Pinot Noir musts exhibited significant regional patterns for fungal communities across

vintages 2017 and 2018 based on Bray-Curtis dissimilarity at OTU level (ADONIS, $R^2_{fungi} = 0.292$, $p <$

0.001) but no significant differences for bacterial communities across both vintages (ADONIS, $R^2_{bacteria} =$

0.108, $p = 0.152$) (Supplementary Table S3), as well as regarding community richness (ANOVA, $F_{bacteria} =$

1.567, $p = 0.374$; $F_{fungi} = 5.142$, $p < 0.01$) (Supplementary Table S4). Within the 2017 vintage, both

bacteria and fungi in the must showed distinctive composition based on the region (ADONIS, $R^2_{bacteria} =$

0.342, $p < 0.001$; $R^2_{fungi} = 0.565$, $p < 0.001$) (Figure 3A, B) (Supplementary Table S3), with more distinct

trend and the improved R^2 coefficient for fungi. Notably, the relative abundance of *Saccharomyces* yeasts

varied significantly from 1.3% (Macedon Range) to 65.6% (Gippsland) between regions (ANOVA, F

$=26.053$, $p < 0.001$) (Figure 3B). As the wine fermentation proceeded, fermentative populations including

yeasts and LAB grew and dominated, thus reshaping the community diversity (Supplementary Figure

S3A, B) and composition (Supplementary Figure S3C, D). Fungal species diversity collapsed as alcoholic

fermentation progressed (ANOVA, $F = 6.724$, $p < 0.01$) (Supplementary Figure S3B), while the impact of

the fermentation on bacterial diversity was insignificant (ANOVA, F = 1.307, p = 0.301), with slight

decrease at early stages and recovery at fermentation end (Supplementary Figure S3A). Linear

discriminant analysis (LDA) effect size (LEfSe) analysis further identified differentially abundant taxa

(Kruskal–Wallis sum-rank test, $\alpha < 0.05$) associated with fermentation stages (Figure 3). For fungal

populations, *Dothideomycetes* (including *Aureobasidium* and *Cladosporium*), *Debaryomycetaceae*

(notably yeast *M. guilliermondii*), *Penicillium corylophilum* and *Filobasidium oeirense* were significantly

abundant in the grape must, with *Leotiomycetes*, *Sarocladium* and *Vishniacozyma victoriae* in early

fermentations (AF), *Saccharomycetes* yeasts (notably *S. cerevisiae*) in mid fermentations (AF-Mid), and

Tremellales at the end of fermentation (AF-End) (Figure 3D). For bacterial communities, *Acidobacteriia*

(spoilage), *Chloroflexi*, *Deltaproteobacteria*, *Sphingobacteriia*, *Cytophagia*, *Planococcaceae* and

Rhizobiaceae were observed with higher abundances in the must, *Proteobacteria* (including

Burkholderiaceae and *Tremblayales*, spoilage) and *Micrococcus* in the AF, *Burkholderia* spp. in the AF-

Mid, *Rhodobacterales* and *Pseudonocardiaceae* in the AF-End, LAB *Leuconostocaceae* (notably

Oenococcus) in the MLF-End (Figure 3C). Regional differences in microbial profiles were not significant

in the finished wines (ADONIS, $R^2_{bacteria}$ = 0.149, p = 0.321; R^2_{fungi} = 0.109, p = 0.205) (Supplementary

Table S3).

To uncover the impact of growing season (vintage) on wine regionality and related microbiota, five

vineyards in Mornington were sampled in 2017 and 2018 to compare within and between vintages.

Within these five vineyards alone, both microbial communities and wine aroma saw significant influence

from vintage effects (Supplementary Table S3). When comparing all samples in the large scale, vintage

only weakly impacted microbial and wine aroma profiles, in particular an insignificant influence on fungi

(ADONIS, R^2_{fungi} = 0.049, p = 0.066) (Supplementary Table S3). We used 2017 vintage data to further

explore microbial biogeography and wine regionality in the following analyses.

Microbial patterns correlate to wine aroma profiling

Network analysis was used to explore connections between regional microbial and wine metabolic

patterns. 57 volatile compounds and 246 OTUs were screened out based on strong correlation coefficients

(Spearman correlation coefficient, $\rho \geq 0.8$; $p < 0.01$), forming the co-occurrence patterns. These variables

showed a sophisticated internal structure, consisting of 303 nodes and 610 edges (average degree =

4.026), with mostly positive correlations (90.66%) (Figure 4A). Both soil (average degree = 3.333; Figure

4B) and must microbiota (average degree = 3.791; Figure 4C) presented high connectivity with wine

metabolites (Figure 4B, C). Soil fungal OTUs (69 of 5610 OTUs) more densely connected with volatile

compounds than bacteria (42 of 12117 OTUs) (Figure 4B). This indicated that soil microbial

communities, especially fungi, might play a role in wine properties. The most densely connected volatile

compounds were assigned to groups of monoterpenes (C18, *p*-cymene; C58, *α*-terpineol),

phenylpropanoids (C40, benzaldehyde; C69, phenethyl acetate), and acetoin (C20) (Figure 4A).

Correspondingly, high-connectivity nodes with these compounds were OTUs of bacterial taxa

Enterobacteriales, *Rhizobiales*, *Burkholderiales* and *Xanthomonadales*, and fungal taxa *Penicillium*,

Rhodotorula and *Neocatenulostroma* from soil and must (Figure 4B, C). The group of esters, which are

yeast-driven products (9), were highly associated with fungal OTUs (in particular *Hypocreales*,

Alternaria, *Cryptococcus and Mortierella*) from the vineyard soil, for example, ethyl hexanoate (C16)

and ethyl dihydrocinnamate (C75) (Figure 4B). Fatty acids (including C49, 4-hydroxybutanoic acid; C80,

octanoic acid) were mostly negatively connected with fungi (including *Acremonium*, *Penicillium*) from

the must (Figure 4C).

Multiple drivers modify wine regionality in the vineyard

Alongside regional patterns in soil and must microbiota, environmental measures of the wine growing

regions displayed significant differences, such as C and N of soil properties, solar radiation and

temperature of weather/ climatic conditions during the growing season (October 2017 – April 2018) (see

Supplementary Table S5 for a complete list). To disentangle the role of microbial ecology on wine

regionality, we used Random Forest modelling (57) to identify biotic (soil and must microbial diversity)

and abiotic predictors (soil and weather parameters) structuring wine regionality, and structural equation

modelling (SEM) (60) to testify whether the relationship between microbial diversity and wine regionality

can be maintained when accounting for multiple drivers simultaneously. The Random Forest model (R^2 =

0.451, $p < 0.01$) demonstrated that fungal diversity was a major predictor of wine regionality. Not

surprisingly, must fungal diversity showed higher importance on the model (increase in MSE) than soil

(Figure 5). The SEM explained 93.8% of the variance found in the pattern of wine regionality (Figure

6A). Weather drove wine aroma profiles directly (especially MT, MLT, MinT and MS) and indirectly by

powerful effects on soil and must microbial diversity, in particular strong effects on soil fungal diversity

(Figure 6A). Must fungal diversity had the highest direct positive effect on wine aroma characteristics,

with direct influences from soil fungal diversity (Figure 6A). Weather and climate could impact soil

nutrient pools primarily through MS, MLT, MinT, and MTrans. Soil properties showed strong effects on

soil microbial diversity and must bacterial diversity, but weak effects on must fungal diversity (Figure

6A). Must bacterial diversity had a weak effect on wine aroma profiles, as well as soil bacteria. Overall,

must fungal diversity was the most important driver of wine characteristics, followed by soil fungal

diversity (Figure 6B), with influences from weather and soil properties both directly and indirectly

(Figure 6A).

Source tracking wine-related fungi within vineyard

As we showed in the SEM, soil fungal diversity had a direct positive influence on must fungal diversity

and thus indirectly contributed to wine aroma profiles (Figure 6). Given that soil is a potential source of

fungi associated with wine production (14), here we attempt to uncover the mechanism whereby soil

fungi could be transported from soil to the grapes. We sampled fungal communities from grapevines and

soil and hypothesised that the xylem/ phloem was the internal mechanism to transport microbes. A total

of 2,140,820 ITS high-quality sequences were generated from soil and grapevine samples (grape, leaf,

xylem sap, root), which were clustered into 4,050 fungal OTUs with 97% pairwise identity. Using

SourceTracker (63), fungal communities in the must were matched to multiple sources from below the

ground to above the ground. Results showed that grape and xylem sap were primary sources of must

fungi, with 32.6% and 41.9% contributions, respectively (Figure 7). Xylem sap showed similar fungal

structure with must (Figure S4A). Further source tracking revealed that the root and soil contributed

90.2% of fungal OTUs of xylem sap, the latter contributed 67.9% of the fungi of grapes (Figure 7).

Notably, *S. cerevisiae* were found shared between microhabitats of soil, root, xylem sap, grape and must,

with the highest (1.22%) and lowest (0.038%) relative abundance in the root and soil, respectively (Figure

S4B). Could xylem vessels be a possible translocation pathway of *S. cerevisiae* from roots to the

aboveground? Chemical analysis of nutrient compositions showed that xylem sap contained nine

carbohydrates (predominantly glucose, fructose and sucrose), 15 amino acids (mainly arginine, aspartic

acid and glutamic acid) and six organic acids (primarily oxalic acid), which could be utilised as carbon

sources and support yeast growth (Figure S4E, F, G) (64). However, no *S. cerevisiae* yeasts were isolated,

distinct isolates of the *Basidiomycota* yeasts of *Cryptococcus* spp. (primarily *C. saitoi*) and *Rhodotorula*

slooffiae were found instead (Figure S4C). The exclusive existence of these species was validated by

isolation from xylem/phloem sap coming from grapevines grown in the glasshouse (Figure S4D).

Discussion

Microbial ecology can influence grapevine health and growth, fermentation, flavour characteristics, and

wine quality and style (12, 13, 20). We systematically investigated the microbiome from the soil to wine

to show that soil and grape must microbiota exhibit regional patterns and that these patterns correlate with

wine metabolites and thus could affect wine quality and style. Here we show that wine regionality is

dependent on fungal ecology and is sensitive to local weather, climate and soil properties. A new

mechanism to transfer fungi from the soil to grapes and must via xylem sap was investigated.

A microbial component of wine *terroir*

Regional spatial patterns have been proposed for soil and grape must microbiota (12, 25-28). The most

abundant bacterial phyla in the vineyard soils in our study were *Actinobacteria*, *Proteobacteria* and

Acidobacteria, which are known to be dominant and ubiquitous in vineyard soil (23, 25, 26, 65). For

fungi, we recovered 14 phyla, 30 classes, 65 orders, 125 families and 216 genera of fungi, recording a

higher diversity than reported in other wine-producing areas in the world (26, 27, 66). *Glomeromycota*,

the phylum of arbuscular mycorrhizal fungi reported to positively affect grapevine growth, was reported

as abundant in New Zealand vineyards (66). In our study, which used amplicon sequences at the ITS

region (rather than the D1/D2 region in (66)), *Glomeromycota* was only recovered with low frequency

from Mornington and Macedon Ranges vineyards. Clearly, there are differences based on the barcoding

region but geographic location may also affect distribution (67), as Coller et al. (2019) using the same

ITS1F/2 primers as in our study, retrieved *Glomeromycota* as a core phylum member from vineyards in

Italy (26). In the must, both principal fermentation drivers (*S. cerevisiae* and LAB) and innocent species

(those that neither ferment nor spoil, such as *Enterobacteriales* and *Aureobasidium*) were present in

different abundances among regions (Figure 3A, B). The order *Lactobacillales*, representing LAB, was

present 0.4% across regions, compared to 29.7% found in California, US (12) and 14% in Catalonia,

Spain (32).

Environmental factors (such as weather and climate) and geographic features structure microbial diversity

and biogeography across various habitats in the soil and plant ecosystems (12, 25, 68-70). In this study,

we demonstrate that microbial biogeographic communities were distinct in both vineyard soils and grape

musts in southern Australia regardless of the growing season/ vintage. This aligns with previous studies

on wine microbial biogeography and provides further evidence for microbial *terroir* [reviewed by Liu et

al.(29); (12, 14, 18, 25)]. Soil bacteria can distinguish wine growing regions, with impacts from soil

properties (Figure 6A) and this is supported by previous work in this field (23, 25, 28). An interesting

finding is that the must bacterial diversity is strongly influenced by soil properties, in particular C: N.

Previous work has shown that structures of must and soil communities are similar and some species

Enterobacteriales and *Actinomycetales* originate from the soil (8, 15). As C: N can be manipulated by

composting and cover crops (71), there is an opportunity to manipulate wine microbiota by a focus on

vineyard management (29). Soil bacterial microflora is recognised as important for plant growth

processes more broadly (72), but fungal diversity beyond endosymbiotic mycorrhizae has not been

systematically investigated for grapevines. Here, we show that the soil fungal communities are distinct for

a region. Our modelling suggests that soil properties and weather strongly affect soil fungal diversity,

which was in line with large-scale studies in which climatic (especially precipitation) and edaphic factors

(especially C: N) are the best predictors of soil fungal richness and community composition (70, 73).

Must fungal diversity is also shaped by weather and soil properties indirectly via soil fungi that ultimately

affect wine aroma profiles.

It is noteworthy that the drivers of microbial patterns change during wine fermentation. Microbial

diversity decreased as alcoholic fermentation proceeds, with a clear loss of microorganisms including

filamentous fungi, non-*Saccharomyces* yeasts (for example *M. guilliermondii*), spoilage bacteria

Acidobacteria and *Proteobacteria*, and other bacteria with unknown fermentative functions (for example

Chloroflexi) (Figure 3C, D), and thus the biogeographic trend was lost by the stage of MLF-End

(Supplementary Table S3). This trend was observed more distinctly in fungal communities compared

with bacteria (Figure S3A, B). This is not unexpected as it is clear that fermentation more strongly affects

fungal populations compared with bacteria, due to increasing fermentation rate, temperature and ethanol

concentration induced by *S. cerevisiae* growth (10, 74). In this case, fermentation conditions, such as the

chemical environment and interactions and/or competition within the community (10, 75), reshape the

microbial patterns observed. Despite the complex change of microbial ecosystem during fermentation, we

show that biogeographic patterns in the must could be reflected in the regional metabolic profiling of

wine. Our modelling indicates that indirect influences of weather and soil properties via shaping soil and

must microbial diversity are more powerful than the direct influences on wine aroma profiles (Figure 6).

In the resulting wines, most volatile compounds were alcohols, esters, acids and aldehydes, some of

which were likely microbial products. Some compunds are grape-derived and are modified by microbial

metabolism during fermentation (9). Co-occurrence networks demonstrate that the soil and must

microbiota correlate closely to the wine metabolome. These interactions also indicate potential

modulations of soil and must microbiota on wine chemistry, in particular those taxa that are numerically

dominant early in the fermentation. For example, bacteria *Enterobacteriales* positively correlated with

some grape-derived volatile compounds (monoterpenes and phenylpropanoids), thus potentially

enhancing varietal aroma in wine (9). *Enterobacteriales* have been also previously related to wine

fermentation rate by Bokulich et al. (19). Fungal genera abundance (for example *Penicillium*) negatively

correlated with the medium-chain fatty acid octanoic acid, which can contribute to mushroom flavour and

also alter *Saccharomyces* metabolism (Bisson 1999). These microorganisms correlate with wine

metabolites and/or can change chemical environments and fermentation behaviours and finally exert

substantial effects on wine metabolite profiles (9, 19). It is particularly important to inform farming

practices to structure regional microbial communities that can benefit soil quality, and thus, crop

productivity.

Fungal communities distinguish wine quality and style

In grape musts, bacterial and fungal communities exhibit different responses to site-specific and

environmental effects. Bacterial regional patterns were not as distinct as fungi, and significantly impacted

by vintage (Supplementary Table S3). Although profound responding to soil properties (for example C:

N) and affecting wine fermentation, must bacteria show insignificant influences on wine aroma profiles

(Figure 6). In contrast, fungal communities display discriminating distribution patterns at the regional

scale and are weakly or insignificantly impacted by vintage in this study, aligning with results presented

by Bokulich et al. (2014) (12). Soil fungal communities are less diverse than bacterial communities

(Figure S2; ref. 67), but of more importance to the resultant wine regionality (see Figure 4, 5, 6). Must

fungi, in particular the fermentative yeasts, conduct alcoholic fermentations and provide aroma

compounds to influence wine flavour (9). As indicated by SEM, soil fungal communities are affected by

local soil properties and weather and exert impacts on must fungal communities (Figure 6). Co-

occurrence relationships between wine metabolome and soil microbiota highlight potential contributions

from fungal taxa (Figure 4). One explanation is that grapevines filter soil microbial taxa selecting for

grape and must consortia (76, 77). Beyond fungi, plant fitness is linked strongly to the responses of soil

microbial communities to environmental conditions (78). More sensitive responses of vineyard soil fungi

might improve grapevine fitness to local environments thus benefiting the expression of regional

characteristics of grapes and wines.

How could yeasts present in the soil be transported to the grape berry? Soil is a reservoir of grapevine-

associated microbiome (Figure 7) that is supported by previous publications (14, 23, 28, 79). As well as

transporting water and minerals absorbed by roots to the photosynthetic organs, xylem sap is also a

microhabitat for microbes that can bear its nutritional environment (64). Here we investigated xylem sap

as a conduit to shape the microbiota in the grape by enrichment of the microbes recruited by roots and

transported by xylem sap to the grape berries (23, 33, 72). The isolated yeasts belong to the *Cryptococcus*

and *Rhodotorula* genera, indicating that the xylem sap environment is not sterile and can potentially

transport yeasts to the phyllosphere. The endophyte *Burkholderia phytofirmans* strain PsJN has been

shown to colonise grapevine roots from the rhizosphere and spread to inflorescence tissues through the

xylem (33, 80). While we were unable to find fermentative yeasts in the sap of grapevines, other yeasts

and/or spores were present and thus may also be transported within the grapevine as well as making their

way to the phyllosphere through other mechanisms (water splashes, insect vectors). As previous studies

show, fermentative yeasts are persistent in vineyards (81, 82) and can be transported by grapevines (34).

We can thus conclude that fungi are the principal drivers of consistent expression of regionality in wine

production.

Our study illustrates that microbes are absolute contributors to wine aroma and that this comes from the

environment in which they are grown and provides a basis to explain how wine may have a distinctive

flavour from a particular region. Fungi play a crucial role in interrelating these biotic and abiotic elements

in vineyard ecosystems and could potentially be transported internally. Climate and soil properties

profoundly structure microbial patterns from the soil to the grape must, which ultimately sculpt wine

metabolic profiling. We do not yet know how grapevines recruit their microbiome to maximise

physiological development and encourage and maintain microbial diversity under local conditions.

Acknowledgements

The authors give their sincere thanks to the vignerons who kindly allowed vineyard access, enabled

sampling and provided wine samples. DL acknowledges support from a Ph.D. scholarship and funding

from Wine Australia (AGW Ph1602) and a Melbourne Research Scholarship from the University of

Melbourne.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

References

References

1.   Gonzálvez A, Llorens A, Cervera M, Armenta S, De la Guardia M. Elemental fingerprint of wines
from the protected designation of origin Valencia. Food Chemistry. 2009;112(1):26-34.
2.   Pereira GE, Gaudillere J-P, Van Leeuwen C, Hilbert G, Lavialle O, Maucourt M, et al. 1H NMR and
chemometrics to characterize mature grape berries in four wine-growing areas in Bordeaux, France.
Journal of agricultural and food chemistry. 2005;53(16):6382-9.
3.   Robinson AL, Adams DO, Boss PK, Heymann H, Solomon PS, Trengove RD. Influence of
geographic origin on the sensory characteristics and wine composition of Vitis vinifera cv. Cabernet
Sauvignon wines from Australia. American journal of enology and viticulture. 2012:ajev. 2012.12023.
4.   Gladstones J. Viticulture and environment: Winetitles; 1992.
5.   van Leeuwen C, Seguin G. The concept of terroir in viticulture. Journal of Wine Research.
2006;17(1):1-10.
6.   Vaudour E, Costantini E, Jones G, Mocali S. An overview of the recent approaches to terroir
functional modelling, footprinting and zoning. Soil. 2015;1(1):287.
7.   Barata A, Malfeito-Ferreira M, Loureiro V. The microbial ecology of wine grape berries.
International journal of food microbiology. 2012;153(3):243-59.
8.   Stefanini I, Cavalieri D. Metagenomic Approaches to Investigate the Contribution of the
Vineyard Environment to the Quality of Wine Fermentation: Potentials and Difficulties. Frontiers in
Microbiology. 2018;9.
9.   Swiegers J, Bartowsky E, Henschke P, Pretorius I. Yeast and bacterial modulation of wine aroma
and flavour. Australian Journal of grape and wine research. 2005;11(2):139-73.
10.   Goddard MR. Quantifying the complexities of Saccharomyces cerevisiae's ecosystem
engineering via fermentation. Ecology. 2008;89(8):2077-82.
11.   Heard GM, Fleet GH. Growth of natural yeast flora during the fermentation of inoculated wines.
Appl Environ Microbiol. 1985;50(3):727-8.
12.   Bokulich NA, Thorngate JH, Richardson PM, Mills DA. Microbial biogeography of wine grapes is
conditioned by cultivar, vintage, and climate. Proceedings of the National Academy of Sciences.
2014;111(1):E139-E48.
13.   Gayevskiy V, Goddard MR. Geographic delineations of yeast communities and populations
associated with vines and wines in New Zealand. The ISME journal. 2012;6(7):1281-90.
14.   Morrison-Whittle P, Goddard MR. From vineyard to winery: a source map of microbial diversity
driving wine fermentation. Environmental microbiology. 2018.
15.   Pinto C, Pinho D, Cardoso R, Custódio V, Fernandes J, Sousa S, et al. Wine fermentation
microbiome: a landscape from different Portuguese wine appellations. Frontiers in microbiology.
2015;6:905.
16.   Portillo MdC, Mas A. Analysis of microbial diversity and dynamics during wine fermentation of
Grenache grape variety by high-throughput barcoding sequencing. LWT-Food Science and Technology.
2016;72:317-21.
17.   Stefanini I, Albanese D, Cavazza A, Franciosi E, De Filippo C, Donati C, et al. Dynamic changes in
microbiota and mycobiota during spontaneous 'Vino Santo Trentino'fermentation. Microbial
biotechnology. 2016;9(2):195-208.
18.   Taylor MW, Tsai P, Anfang N, Ross HA, Goddard MR. Pyrosequencing reveals regional
differences in fruit-associated fungal communities. Environmental microbiology. 2014;16(9):2848-58.
19.   Bokulich NA, Collins TS, Masarweh C, Allen G, Heymann H, Ebeler SE, et al. Associations among
Wine Grape Microbiome, Metabolome, and Fermentation Behavior Suggest Microbial Contribution to
Regional Wine Characteristics. mBio. 2016;7(3):e00631-16.
20.   Knight S, Klaere S, Fedrizzi B, Goddard MR. Regional microbial signatures positively correlate
with differential wine phenotypes: evidence for a microbial aspect to terroir. Scientific reports.
2015;5:14233.

21. Müller DB, Vogel C, Bai Y, Vorholt JA. The plant microbiota: systems-level insights and
perspectives. annual review of genetics. 2016;50:211-34.
22. Rillig MC, Lehmann A, Lehmann J, Camenzind T, Rauh C. Soil Biodiversity Effects from Field to
Fork. Trends in plant science. 2017.
23. Zarraonaindia I, Owens SM, Weisenhorn P, West K, Hampton-Marcell J, Lax S, et al. The soil
microbiome influences grapevine-associated microbiota. MBio. 2015;6(2):e02527-14.
24. Gilbert JA, van der Lelie D, Zarraonaindia I. Microbial terroir for wine grapes. Proceedings of the
National Academy of Sciences. 2014;111(1):5-6.
25. Burns KN, Kluepfel DA, Strauss SL, Bokulich NA, Cantu D, Steenwerth KL. Vineyard soil bacterial
diversity and composition revealed by 16S rRNA genes: Differentiation by geographic features. Soil
Biology and Biochemistry. 2015;91:232-47.
26. Coller E, Cestaro A, Zanzotti R, Bertoldi D, Pindo M, Larger S, et al. Microbiome of vineyard soils
is shaped by geography and management. Microbiome. 2019;7(1):140.
27. Knight SJ, Karon O, Goddard MR. Small scale fungal community differentiation in a vineyard
system. Food Microbiology. 2019:103358.
28. Mezzasalma V, Sandionigi A, Guzzetti L, Galimberti A, Grando MS, Tardaguila J, et al.
Geographical and cultivar features differentiate grape microbiota in Northern Italy and Spain vineyards.
Frontiers in Microbiology. 2018;9:946.
29. Liu D, Zhang P, Chen D, Howell KS. From the vineyard to the winery: how microbial ecology
drives regional distinctiveness of wine. Frontiers in Microbiology. 2019;10:2679.
30. Jara C, Laurie VF, Mas A, Romero J. Microbial Terroir in Chilean Valleys: Diversity of Non-
conventional Yeast. Frontiers in microbiology. 2016;7.
31. Miura T, Sánchez R, Castañeda LE, Godoy K, Barbosa O. Is microbial terroir related to geographic
distance between vineyards? Environmental Microbiology Reports. 2017;9(6):742-9.
32. Portillo MdC, Franquès J, Araque I, Reguant C, Bordons A. Bacterial diversity of Grenache and
Carignan grape surface from different vineyards at Priorat wine region (Catalonia, Spain). International
journal of food microbiology. 2016;219:56-63.
33. Compant S, Kaplan H, Sessitsch A, Nowak J, Ait Barka E, Clément C. Endophytic colonization of
Vitis vinifera L. by Burkholderia phytofirmans strain PsJN: from the rhizosphere to inflorescence tissues.
FEMS Microbiology Ecology. 2008;63(1):84-93.
34. Mandl K, Schieck J, Silhavy-Richter K, Alexander P, Schneider V, Schmidt H-P. Vines take up
yeasts from soil and transport them through the vine to the stem and skins of grapes. Ithaka J.
2015:349-55.
35. Goddard MR, Anfang N, Tang R, Gardner RC, Jun C. A distinct population of Saccharomyces
cerevisiae in New Zealand: evidence for local dispersal by insects and human-aided global dispersal in
oak barrels. Environmental microbiology. 2010;12(1):63-73.
36. Lam SS, Howell KS. Drosophila-associated yeast species in vineyard ecosystems. FEMS
microbiology letters. 2015;362(20).
37. Stefanini I, Dapporto L, Legras J-L, Calabretta A, Di Paola M, De Filippo C, et al. Role of social
wasps in Saccharomyces cerevisiae ecology and evolution. Proceedings of the National Academy of
Sciences. 2012;109(33):13398-403.
38. Jones DA, Wang W, Fawcett R. High-quality spatial climate data-sets for Australia. Australian
Meteorological and Oceanographic Journal. 2009;58(4):233.
39. López-Millán AF, Morales F, Abadía A, Abadía J. Effects of iron deficiency on the composition of
the leaf apoplastic fluid and xylem sap in sugar beet. Implications for iron and carbon transport. Plant
Physiology. 2000;124(2):873-84.
40. Rayment GE, Lyons DJ. Soil chemical methods: Australasia: CSIRO publishing; 2011.
41. Liu D, Xing R-R, Li Z, Yang D-M, Pan Q-H. Evolution of volatile compounds, aroma attributes, and
sensory perception in bottle-aged red wines and their correlation. European Food Research and
Technology. 2016:1-12.

42. Zhang P, Barlow S, Krstic M, Herderich M, Fuentes S, Howell K. Within-vineyard, within-vine, and
within-bunch variability of the rotundone concentration in berries of Vitis vinifera L. cv. Shiraz. Journal of
agricultural and food chemistry. 2015;63(17):4276-83.
43. Liu D, Gao Y, Li X-X, Li Z, Pan Q-H. Attenuated UV Radiation Alters Volatile Profile in Cabernet
Sauvignon Grapes under Field Conditions. Molecules. 2015;20(9):16946-69.
44. Yu Y, Lee C, Kim J, Hwang S. Group-specific primer and probe sets to detect methanogenic
communities using quantitative real-time polymerase chain reaction. Biotechnology and bioengineering.
2005;89(6):670-9.
45. Gardes M, Bruns TD. ITS primers with enhanced specificity for basidiomycetes-application to the
identification of mycorrhizae and rusts. Molecular ecology. 1993;2(2):113-8.
46. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows
analysis of high-throughput community sequencing data. Nature methods. 2010;7(5):335.
47. Magoč T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome
assemblies. Bioinformatics. 2011;27(21):2957-63.
48. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. UCHIME improves sensitivity and speed of
chimera detection. Bioinformatics. 2011;27(16):2194-200.
49. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of rRNA
sequences into the new bacterial taxonomy. Appl Environ Microbiol. 2007;73(16):5261-7.
50. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes, a chimera-
checked 16S rRNA gene database and workbench compatible with ARB. Applied and environmental
microbiology. 2006;72(7):5069-72.
51. Kõljalg U, Larsson KH, Abarenkov K, Nilsson RH, Alexander IJ, Eberhardt U, et al. UNITE: a
database providing web-based methods for the molecular identification of ectomycorrhizal fungi. New
Phytologist. 2005;166(3):1063-8.
52. Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ, et al. The vegan package.
Community ecology package. 2007;10:631-7.
53. Roberts DW. labdsv: Ordination and multivariate analysis for ecology. R package version.
2007;1(1).
54. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic biomarker
discovery and explanation. Genome biology. 2011;12(6):R60.
55. Bastian M, Heymann S, Jacomy M, editors. Gephi: an open source software for exploring and
manipulating networks. Third international AAAI conference on weblogs and social media; 2009.
56. Junker BH, Schreiber F. Analysis of biological networks: Wiley Online Library; 2008.
57. Breiman L. Random forests. Machine learning. 2001;45(1):5-32.
58. Liaw A, Wiener M. Classification and regression by randomForest. R news. 2002;2(3):18-22.
59. Fortmann-Roe S. Accurate, Adaptable, and Accessible Error Metrics for Predictive. R package
version 09. 2013;2.
60. Grace JB. Structural equation modeling and natural systems: Cambridge University Press; 2006.
61. Eisenhauer N, Bowker MA, Grace JB, Powell JR. From patterns to causal understanding:
structural equation modeling (SEM) in soil ecology. Pedobiologia. 2015;58(2-3):65-72.
62. Schermelleh-Engel K, Moosbrugger H, Müller H. Evaluating the fit of structural equation models:
Tests of significance and descriptive goodness-of-fit measures. Methods of psychological research
online. 2003;8(2):23-74.
63. Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG, et al. Bayesian
community-wide culture-independent microbial source tracking. Nature methods. 2011;8(9):761.
64. Yadeta K, Thomma B. The xylem as battleground for plant hosts and vascular wilt pathogens.
Frontiers in plant science. 2013;4:97.
65. Fierer N, Leff JW, Adams BJ, Nielsen UN, Bates ST, Lauber CL, et al. Cross-biome metagenomic
analyses of soil microbial communities and their functional attributes. Proceedings of the National
Academy of Sciences. 2012;109(52):21390-5.

66.   Morrison-Whittle P, Goddard M. Fungal communities are differentially affected by conventional
and biodynamic agricultural management approaches in vineyard ecosystems. Agriculture, Ecosystems
and Environment. 2017.
67.   Öpik M, Vanatoa A, Vanatoa E, Moora M, Davison J, Kalwij J, et al. The online database MaarjAM
reveals global and ecosystemic distribution patterns in arbuscular mycorrhizal fungi (Glomeromycota).
New Phytologist. 2010;188(1):223-41.
68.   Fierer N, Jackson RB. The diversity and biogeography of soil bacterial communities. Proceedings
of the National Academy of Sciences of the United States of America. 2006;103(3):626-31.
69.   Martiny JBH, Bohannan BJ, Brown JH, Colwell RK, Fuhrman JA, Green JL, et al. Microbial
biogeography: putting microorganisms on the map. Nature Reviews Microbiology. 2006;4(2):102.
70.   Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, et al. Global diversity and
geography of soil fungi. science. 2014;346(6213):1256688.
71.   Burns KN, Bokulich NA, Cantu D, Greenhut RF, Kluepfel DA, O'Geen AT, et al. Vineyard soil
bacterial diversity and composition revealed by 16S rRNA genes: Differentiation by vineyard
management. Soil Biology and Biochemistry. 2016;103:337-48.
72.   Berg G, Smalla K. Plant species and soil type cooperatively shape the structure and function of
microbial communities in the rhizosphere. FEMS microbiology ecology. 2009;68(1):1-13.
73.   Lauber CL, Strickland MS, Bradford MA, Fierer N. The influence of soil properties on the
structure of bacterial and fungal communities across land-use types. Soil Biology and Biochemistry.
2008;40(9):2407-15.
74.   Bokulich NA, Joseph CL, Allen G, Benson AK, Mills DA. Next-generation sequencing reveals
significant bacterial diversity of botrytized wine. PloS one. 2012;7(5):e36357.
75.   Liu Y, Rousseaux S, Tourdot-Maréchal R, Sadoudi M, Gougeon R, Schmitt-Kopplin P, et al. Wine
microbiome: a dynamic world of microbial interactions. Critical reviews in food science and nutrition.
2017;57(4):856-73.
76.   Deveau A, Bonito G, Uehling J, Paoletti M, Becker M, Bindschedler S, et al. Bacterial–fungal
interactions: ecology, mechanisms and challenges. FEMS microbiology reviews. 2018;42(3):335-52.
77.   Panke-Buisse K, Poole AC, Goodrich JK, Ley RE, Kao-Kniffin J. Selection on soil microbiomes
reveals reproducible impacts on plant function. The ISME journal. 2015;9(4):980.
78.   Lau JA, Lennon JT. Rapid responses of soil microorganisms improve plant fitness in novel
environments. Proceedings of the National Academy of Sciences. 2012;109(35):14058-62.
79.   Grady KL, Sorensen JW, Stopnisek N, Guittar J, Shade A. Assembly and seasonality of core
phyllosphere microbiota on perennial biofuel crops. bioRxiv. 2019:446369.
80.   Compant S, Reiter B, Sessitsch A, Nowak J, Clément C, Barka EA. Endophytic colonization of Vitis
vinifera L. by plant growth-promoting bacterium Burkholderia sp. strain PsJN. Applied and
Environmental Microbiology. 2005;71(4):1685-93.
81.   Börlin M, Venet P, Claisse O, Salin F, Legras J-L, Masneuf-Pomarede I. Cellar-associated
Saccharomyces cerevisiae population structure revealed high diversity and perennial persistence in
Sauternes wine estates. Applied and environmental microbiology. 2016:AEM. 03627-15.
82.   Valero E, Schuller D, Cambon B, Casal M, Dequin S. Dissemination and survival of commercial
wine yeast in the vineyard: A large-scale, three-years study. FEMS Yeast Research. 2005;5(10):959-69.
83.   Schurr U. Xylem sap sampling—new approaches to an old topic. Trends in Plant Science.
1998;3(8):293-8.
84.   Cohen SA. Amino acid analysis using precolumn derivatization with 6-aminoquinolyl-N-
hydroxysuccinimidyl carbamate. Amino Acid Analysis Protocols: Springer; 2001. p. 39-47.
85.   Cohen SA, Michaud DP. Synthesis of a fluorescent derivatizing reagent, 6-aminoquinolyl-N-
hydroxysuccinimidyl carbamate, and its application for the analysis of hydrolysate amino acids via high-
performance liquid chromatography. Analytical biochemistry. 1993;211(2):279-87.
86.   Andersen PC, Brodbeck BV, Mizell RF. Metabolism of amino acids, organic acids and sugars
extracted from the xylem fluid of four host plants by adult Homalodisca coagulata. Entomologia
Experimentalis et Applicata. 1989;50(2):149-59.

87.   Kurtzman CP, Robnett CJ. Identification and phylogeny of ascomycetous yeasts from analysis of
nuclear large subunit (26S) ribosomal DNA partial sequences. Antonie van Leeuwenhoek.
1998;73(4):331-71.

Figures

Figure 1 Wine metabolome shows regional variation across six wine growing regions in 2017. Shown

are α-diversity (Shannon index) (A) and PCoA based on Bray-Curtis dissimilarity obtained from

comparing volatile profiles (B).

Figure 2 Vineyard soil microbial communities demonstrate regional patterns. Bray-Curtis distance PCoA

of bacterial communities (A) and fungal communities (B).

Figure 3 Microbiota exhibit regional differentiation in musts for both bacterial and fungal profiles. Stage

of fermentation influences microbial composition of bacteria and fungi. Must bacterial taxa (A) in the

order level with greater than 1.0% relative abundance, and *Lactobacillales*. Must fungal taxa (B) in the

genus level with greater than 1.0% relative abundance. Linear discriminant analysis (LDA) effect size

(LEfSe) taxonomic cladogram comparing all musts and wines categorised by fermentation stage.

Significantly discriminant taxon nodes (C, bacteria; D, fungi) are coloured and branch areas are shaded

according to the highest-ranked stage for that taxon. For each taxon detected, the corresponding node in

the taxonomic cladogram is coloured according to the highest-ranked group for that taxon. If the taxon is

not significantly differentially represented between sample groups, the corresponding node is coloured

yellow.

Figure 4 Statistically significant and strong co-occurrence relationships between wine volatile

compounds and bacterial and fungal OTUs in vineyard soils and musts across six wine growing regions.

Network plots for wine aroma with soil and must microbiota (A), and simplified network plots for wine

aroma with soil microbiota (B) and with must microbiota (C), separately. Circle nodes represent assigned

volatile compounds, bacterial and fungal OTUs, with different colours. Direct connections between

compounds and OTUs indicate strong correlations (Spearman correlation coefficient, $\rho \geq 0.8$; $p < 0.01$).

778   The colour of edges describes the positive correlation (pink) or the negative correlation (green). The size

779   of nodes is proportioned to the inconnected degree.

Figure 5 Main predictors of

wine regionality. Shown is the Random Forest mean predictor importance by % of increase in mean

square error (MSE) of climate, soil properties and microbial diversity (Shannon index) on wine

regionality. Soil bac. div., soil bacterial diversity; Soil fung. div., soil fungal diversity; Must bac. div.,

must bacterial diversity; Must fung. div., must fungal diversity. Significance levels: *$p < 0.05$, **$p <$

0.01.

Figure 6 Direct and indirect effects of climate, soil properties and microbial diversity (Shannon index)

on wine regionality. Structural equation model (SEM) fitted to the diversity of wine aroma profiles (A)

and standardized total effects (direct plus indirect effects) derived from the model (B). Climate and soil

properties are composite variables encompassing multiple observed parameters (see materials and

methods for the complete list of factors used to generate this model). Numbers adjacent to arrows are path

coefficients and indicative of the effect size of the relationship. The width of arrows is proportional to the

strength of path coefficients. R^2 denotes the proportion of variance explained. (a) (0.747* MT) (0.666*

MLT) (0.686* MinT) (-0.875** MS). (b) (0.753* MinT) (0.772* MLT) (-0.683* MS) (-0.737* MHT) (-

0.843** MaxT). C, soil carbon; N, soil nitrogen; C:N, soil carbon nitrogen ratio; MS, mean solar

radiation; MT, mean temperature; MLT, mean low temperature; MHT, mean high temperature; MinT,

minimum temperature; MaxT, maximum temperature; MTrans, mean transpiration. Significance levels:

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 7 Fungal communities in musts emerge from multiple sources in the vineyard, but primarily from

grape and xylem sap. Percent composition estimate for the contributions of possible sources in the

 Supplementary Tables and Figures

 Table S1 Number of soil, must and ferments, and plant samples collected in this study

(A) Soil, must and ferment samples

Vintage	Region	Vineyard (V)	Sample Fraction					
			Soil	Must[1]	AF[1]	AF-Mid[1]	AF-End[1]	MLF-End[1,2]
2017	Geelong	V1	3	1	1	1	1	1
		V2	3	1	1	1	1	1
	Mornington	V3	3	1	1	1	1	1
		V4	3	1	1	1	1	1
		V5	3	1	1	1	1	1
		V6	3	1	1	1	1	1
		V7	3	N/A	N/A	N/A	N/A	N/A
	Macedon Ranges	V8	3	1	1	1	1	1
		V9	3	1	1	1	1	1
	Yarra Valley	V10	3	1	1	1	1	1
		V11	3	1	1	1	1	1
	Grampians	V12	3	1	1	1	1	1
		V13	3	N/A	N/A	N/A	N/A	N/A
	Gippsland	V14	3	1	1	1	1	1
		V15	3	1	1	1	1	1
2018	Mornington	V3	3	1	1	1	1	1
		V4	3	1	1	1	1	1
		V5	3	1	1	1	1	1
		V6	3	1	1	1	1	1
		V7	3	1	1	1	1	1

[1] For microbial analysis: triplicates from tanks/ barrels were collected and mixed as one composite sample which was analysed by NGS

[2] For wine aroma analysis: triplicates were analysed by GC-MS

N/A: not applicable as spontaneous fermentations did not complete and were excluded from analysis

(B) Soil and plant samples

Vintage	Region	Vineyard (V)	Vine (v)	Sample Fraction				
				Soil	Root	Xylem sap	Leaf	Grape
2018	Mornington	V5	v1	1	1	1	1	1
			v2	1	1	1	1	1
			v3	1	1	1	1	1
			v4	1	1	1	1	1
			v5	1	1	1	1	1
		V6	v6	1	1	1	1	1
			v7	1	1	1	1	1
			v8	1	1	1	1	1
			v9	1	1	1	1	1
			v10	1	1	1	1	1

827   **Table S2 ANOVA results of wine volatile compounds among wine growing regions in 2017**

	Retention time (min)	Volatile Compound	CAS No*	$F_{(5, 33)}$	p	Pr (>F)
C1	3.123	Ethyl Acetate	141-78-6	5.506	0.001	**
C2	4.256	Ethyl propanoate	105-37-3	0.102	0.986	
C3	5.326	Isobutyl acetate	110-19-0	5.664	8.58E-04	***
C4	5.938	Ethyl butanoate	105-54-4	6.896	2.16E-04	***
C5	7.265	Hexanal	66-25-1	0.640	0.618	
C6	8.245	2-Methyl-1-propanol	78-83-1	44.540	5.42E-13	***
C7	8.734	3-Methylbutyl acetate	123-92-2	1.657	0.176	
C8	9.253	Ethyl pentanoate	539-82-2	0.345	0.765	
C9	9.865	1-Butanol	71-36-3	52.260	6.58E-14	***
C10	11.300	Methyl hexanoate	106-70-7	26.870	3.09E-10	***
C11	11.500	Isoamyl propanoate	105-68-0	2.015	0.078	
C12	12.012	*D*-Limonene	5989-27-5	14.420	0.00000032	***
C13	12.457	*(E)*-2-Hexenal	6728-26-3	14.870	2.33E-07	***
C14	12.689	3-Methyl-1-butanol	123-51-3	1.046	0.274	
C15	12.894	4-Octanone	589-63-9	2.145	0.075	
C16	13.345	Ethyl hexanoate	123-66-0	10.950	0.0000046 3	***
C17	14.105	Styrene	100-42-5	0.370	0.865	
C18	14.523	*p*-Cymene	99-87-6	22.128	8.91E-10	***
C19	14.923	Hexyl acetate	142-92-7	2.711	0.039	*
C20	15.245	Acetoin	513-86-0	4.806	0.002	**
C21	17.312	2-Heptanol	543-49-7	8.030	0.0000669	***
C22	17.523	Ethyl heptanoate	106-30-9	4.648	0.003	**
C23	17.646	6-Methyl-5-heptene-2-one	110-93-0	1.784	0.221	
C24	18.123	Ethyl lactate	97-64-3	1.085	0.389	
C25	18.726	1-Hexanol	111-27-3	14.430	3.18E-07	***
C26	19.123	3-Hexanol	623-37-0	0.444	0.868	
C27	19.356	Heptyl acetate	112-06-1	2.332	0.067	
C28	20.463	3-Octanol	589-98-0	8.009	0.0000683	***
C29	20.912	2-Hexanol	626-93-7	0.925	0.566	
C30	21.312	*(E)*-2-Octenal	2548-87-0	0.460	0.747	
C31	21.925	Ethyl octoate	106-32-1	1.918	0.121	
C32	22.702	Acetic acid	64-19-7	7.994	0.0000693	***
C33	22.813	1-Octen-3-ol	3391-86-4	16.700	6.89E-08	***
C34	22.900	Isopentyl hexanoate	2198-61-0	3.164	0.021	*
C35	23.103	1-Heptanol	111-70-6	24.680	8.49E-10	***
C36	23.383	6-Methyl-5-hepten-2-ol	1569-60-4	5.329	0.001	**
C37	23.713	Octyl acetate	112-14-1	0.152	0.856	
C38	24.125	Ethyl 6-heptenoate	25118-23-4	2.216	0.079	
C39	24.500	2-Ethyl-1-hexanol	104-76-7	10.020	0.0000103	***
C40	25.103	Benzaldehyde	100-52-7	5.842	7.00E-04	***

C41	25.725	2-Nonanol	628-99-9	14.080	4.06E-07	***
C42	26.114	Ethyl nonanoate	123-29-5	1.223	0.322	
C43	26.806	Linalool	78-70-6	16.480	7.95E-08	***
C44	27.212	1-Octanol	111-87-5	7.832	0.0000816	***
C45	27.513	2-Methyl-propanoic acid	79-31-2	10.880	0.00000491	***
C46	28.012	2,3-Butanediol	513-85-9	20.140	8.87E-09	***
C47	28.426	Methyl decanoate	110-42-9	0.915	0.485	
C48	28.634	Terpinen-4-ol	562-74-3	1.287	0.296	
C49	29.123	4-Hydroxybutanoic acid	591-81-1	6.407	3.69E-04	***
C50	29.405	Ethyl 2-furylcarboxylate	614-99-3	1.254	0.258	
C51	29.716	Benzeneacetaldehyde	122-78-1	1.887	0.126	
C52	30.023	Butyric acid	107-92-6	0.105	0.891	
C53	30.321	Ethyl decanoate	110-38-3	0.644	0.668	
C54	30.926	3-Methylbutyl octanoate	2035-99-6	0.957	0.459	
C55	31.216	1-Nonanol	143-08-8	9.500	0.0000165	***
C56	31.605	Diethyl succinate	123-25-1	3.197	0.020	*
C57	32.213	Ethyl 9-decenoate	67233-91-4	0.397	0.847	
C58	32.405	α-Terpineol	98-55-5	3.861	0.008	**
C59	33.012	Methionol	505-10-2	15.600	1.42E-07	***
C60	33.426	Benzyl acetate	140-11-4	2.225	0.071	
C61	34.738	Methyl salicylate	119-36-8	6.335	4.00E-04	***
C62	33.616	Naphthalene, 1,2-dihydro-1,1,6-trimethyl	30364-38-6	0.145	0.899	
C63	35.012	1-Decanol	112-30-1	4.640	0.003	**
C64	35.209	Citronellol	106-22-9	1.673	0.172	
C65	35.529	Ethyl benzeneacetate	101-97-3	2.097	0.093	
C66	36.123	Ethyl salicylate	118-61-6	5.335	0.001	**
C67	36.312	Nerol	106-25-2	4.026	0.007	**
C68	36.404	Methyl dodecanoate	111-82-0	0.011	0.956	
C69	36.503	Phenethyl acetate	103-45-7	21.302	2.91E-10	***
C70	36.613	β-Damascenone	23726-93-4	26.070	4.44E-10	***
C71	37.723	Ethyl dodecanoate	106-33-2	0.121	0.987	
C72	38.002	Hexanoic acid	142-62-1	3.879	0.008	**
C73	38.303	Guaiacol	90-05-1	2.391	0.061	
C74	38.716	Benzyl alcohol	100-51-6	7.588	1.00E-07	***
C75	38.913	Ethyl dihydrocinnamate	2021-28-5	846.000	<2E-16	***
C76	39.812	Ethyl 3-methylbutyl succinate	28024-16-0	8.985	0.0000265	***
C77	40.023	Phenylethyl Alcohol	60-12-8	10.080	0.00000977	***
C78	40.712	β-Ionone	127-41-3	1.053	0.451	
C79	44.625	Ethyl tetradecanoate	124-06-1	0.139	0.982	
C80	45.217	Octanoic acid	124-07-2	3.437	0.014	*
C81	46.813	Ethyl cinnamate	103-36-6	35.660	9.44E-12	***
C82	48.125	Eugenol	97-53-0	0.285	0.847	
C83	48.526	4-Ethyl-phenol	123-07-9	0.156	0.457	
C84	48.608	Nonanoic acid	112-05-0	1.728	0.079	

C85	51.112	Ethyl hexadecanoate	628-97-7	1.061	0.401	
C86	51.710	Methyl dihydrojasmonate	24851-98-7	0.945	0.243	
C87	51.801	n-Decanoic acid	334-48-5	3.785	0.009	**
C88	52.866	2,4-bis(1,1'-dimethylethyl)phenol	96-76-4	4.387	0.004	**

*CAS Registry number

 Table S3 Permutational multivariate analysis of variance using distance matrices of category
 effects on microbial and wine aroma diversity

Sample Group	Vintage	Factor	Soil Bacteria R^2	p	Soil Fungi R^2	p	Must Bacteria R^2	p	Must Fungi R^2	p	Wine (MLF-End) Bacteria R^2	p	Wine (MLF-End) Fungi R^2	p	Wine Aroma MLF-End R^2	p
All	Both	Region	0.318	0.001	0.254	0.001	0.108	0.152	0.292	0.001	N/A	N/A	N/A	N/A	0.566	0.001
All	2017	Region	0.392	0.001	0.419	0.001	0.342	0.001	0.565	0.001	0.149	0.321	0.109	0.205	0.703	0.001
All	Both	Vintage	0.102	0.001	0.086	0.001	0.186	0.001	0.049	0.066	N/A	N/A	N/A	N/A	0.267	0.001
orningto n	Both	Vintage	0.155	0.001	0.151	0.001	0.305	0.001	0.143	0.001	N/A	N/A	N/A	N/A	0.355	0.001

ta-diversity calculated by the Bray–Curtis distance; Microbial diversity was tested at OTU level.

834 **Table S4 α-diversity (Shannon index) of soil and must microbial communities from six wine**

835 **growing regions in 2017 and 2018**

Vintage	Region	Soil		Must	
		Bacteria	Fungi	Bacteria	Fungi
2017	Geelong	5.486 ± 0.058	3.869 ± 0.181	1.838 ± 0.145	0.938 ± 0.125
	Mornington	6.178 ± 0.074	4.472 ± 0.139	2.216 ± 0.242	1.840 ± 0.217
	Macedon Ranges	4.317 ± 0.047	2.654 ± 0.148	2.424 ± 0.425	1.582 ± 0.246
	Yarra Valley	6.201 ± 0.062	3.355 ± 0.238	1.756 ± 0.314	2.560 ± 0.123
	Grampians	6.238 ± 0.120	3.615 ± 0.191	1.966 ± 0.125	1.297 ± 0.086
	Gippsland	6.416 ± 0.036	4.080 ± 0.227	1.855 ± 0.245	2.200 ± 0.361
2018	Mornington	6.559 ± 0.029	4.187 ± 0.104	1.995 ± 0.357	1.694 ± 0.128

838 **Table S5 ANOVA results of soil properties and weather conditions among wine growing regions in**
839 **2017**

		$F_{(5,39)}$	p	Pr (>F)
Soil	pH	2.300	0.067	
	C	5.891	5.37E-04	***
	N	3.981	0.006	**
	C:N	6.309	3.27E-04	***
	Nitrate	11.640	1.56E-06	***
	Ammonium	1.780	0.144	
Weather	Mean solar radiation	631.700	<2e-16	***
	Mean high temperature	22.100	1.14E-09	***
	Mean low temperature	762.600	<2e-16	***
	Maximum temperature	45.580	7.09E-14	***
	Minimum temperature	3823.000	<2e-16	***
	Mean temperature	18.290	1.12E-08	***
	Precipitation	172.300	<2e-16	***
	Mean relative soil moisture	0.901	0.492	
	Mean evaporation	0.729	0.607	
	Mean transpiration	188.100	<2e-16	***

Figure S1 Map of 15 sampling vineyards from six Pinot Noir wine-producing regions in southern

Australia, spanning 400 km (E-W).

Figure S2 Vineyard soil microbial community compositions across all samples from six grape growing

regions. Shown are average percentages of taxa (characterised to the phylum level) across sites in each

region: (A) dominant bacterial phyla with greater than 1.0% relative abundance; (B) fungal phyla.

Figure S3 Stage of fermentation influences microbial diversities. Bacterial (A) and fungal (B) α-diversity

(Shannon index) changes during wine fermentation. Bray-Curtis distance PCoA of bacterial communities

(C) and fungal communities (D) according to the fermentation stage.

Figure S4 Xylem sap is a potential translocation medium of yeasts from the soil to the grapevine. (A)

Fungal taxa in the genus level of xylem sap from the vineyard. (B) Relative abundances of *S. cerevisiae* in

the soil, root, xylem sap, grape, must. Cultured yeast species from the xylem sap from vineyard (C) and

glasshouse (D). Nutrient compositions of xylem sap: carbohydrates (E), total amino acids (protein and

free) (F), and organic acids (G).

Supplementary Materials and Methods
870

Xylem sap collection

Vine shoots (n = 10) were harvested from the vineyard (Table S1) under aseptic conditions and then

centrifuged (39). The shoots were peeled to remove the outer bark and phloem layer, sterilised with 70%

ethanol and cut to fit into 10-ml sterile centrifuge tubes that had 10 sterile glass beads at the bottom. The

shoots were centrifuged at $15,000 \times g$ for 10 min at 4°C and the xylem sap collected giving ~ 2.0 mL per

sample. Additional xylem sap (n = 5) were sampled from *Vitis vinifera* Shiraz grapevines grown in a

glasshouse at the University of Melbourne. Xylem fluid was extracted with a pressure cylinder apparatus

(similar to a Scholander pressure chamber) (83). Grapevines were uprooted and the soil was carefully

removed from the roots. On an aseptic bench, the main trunk was cut 5-10 cm above the roots with a

sterile blade, girdled to remove phloem tissue, sterilised by immersion in 70% ethanol, and immediately

inserted into a pressure cylinder. The cylinder applied 60-70 kPa pressure for two hours to extract the

xylem sap (~ 4.0 mL). Xylem sap was divided into two subsamples, one used immediately to isolate

living yeasts and the other flash frozen with liquid nitrogen and stored at -80°C for DNA extraction, next-

generation sequencing and chemical analysis.

Chemical analysis of xylem sap composition

Carbohydrates of xylem sap were determined using enzymatic methods by Megazyme assay kits

(Megazyme, Ireland) following the manufacturer's protocol. Amino acids (free and protein- bound) were

determined using pre-column derivatisation with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate

followed by separation and quantification with the ACQUITY Ultra Performance LC (UPLC; Waters,

MA, USA) system at the Australian Proteome Analysis Facility. The column was an ACQUITY UPLC

BEH C18 column (1.7 μm × 2.1 mm × 5 mm) with detection at 260 nm (UV) and a flow rate of 0.7

mL/min at 57 – 60°C. Identification and quantitation of the amino acids was performed against a set of

prepared standards, with DL-norvaline as the internal standard (84, 85). Organic acids were determined

using a Waters High-performance liquid chromatography (HPLC) (Waters, MA, USA) based on

Andersen et al (1989) (86) with modification. Here, 20 μL xylem sap was injected through a Synergi™

Hydro-RP LC Column (250 mm × 4.6 mm × 4 μm; Phenomenex Inc, CA, USA) at 60°C, with detection

at 210 nm (UV). Mobile phases at a flow rate of 1.0 mL/min, with 20 mM potassium phosphate buffer (A,

pH = 1.5) and 100% methanol (B) following a gradient programme: (0-2.5) min, 100% A; (2.5-2.9) min,

linear ramp to 30% B; (2.9-8.0) min, 30% B; (8.0-8.5) min, linear ramp to 100% A; (8.5-10) min, 100%

A. Identification and quantitation of the compounds was performed using a set of prepared standards.

903   Isolation and identification of yeasts from xylem sap

Yeasts were isolated and identified from xylem sap to explore the potential translocation mechanism of

yeasts in the vineyard. Xylem sap was serially diluted and plated (0.1 mL) onto solid yeast extract

peptone dextrose (YPD) medium that was supplemented with 34 mg/mL chloramphenicol and 25 mg/mL

ampicillin to inhibit bacterial growth. Plates were incubated at 28°C for 2-3 days in aerobic conditions.

Single colonies with different morphological types were streaked onto Wallerstein Nutrient (WLN) agar

media to obtain pure cultures. DNA was recovered from pure colonies using the MasterPure™ Yeast

DNA Purification Kit (Epicentre, Madison, WI) following the manufacturer's instructions. The 26S

rDNA D1/D2 domain was amplified using primers NL1/4 (87) for sequencing by Australian Genome

Research Facility (AGRF). Sequences were trimmed, aligned and analysed using BLAST at NCBI

(http://blast.ncbi.nlm.nih.gov/Blast.cgi). Sequence data was uploaded to Genbank with accession numbers

MN847682 - MN847692.