DNA methylation across the genome in aged human skeletal muscle tissue and

muscle stem cells: The role of HOX genes and physical activity

Turner DC[#1,2,3], Gorski PP[#1,3], Maasar MF[4], Seaborne RA[2,3,6], Baumert P[4,5], Brown AD[2],

Kitchen MO[3], Erskine RM[4,7], Dos-Remedios I[8], Voisin S[9], Eynon N[9], Sultanov RI[10], Borisov

OV[10,11], Larin AK[10], Semenova EA[10], Popov DV[12], Generozov EV[10],

Stewart CE[2], Drust B[14], Owens DJ[2,4], Ahmetov II[$4,10,13], Sharples AP*[1,2,3].

[1] Department for Physical Performance, Norwegian School of Sport Sciences (NiH), Oslo, Norway.

[2] Stem Cells, Ageing and Molecular Physiology Unit, Exercise Metabolism and Adaptation Research

Group, Research Institute for Sport and Exercise Sciences, Liverpool John Moores University,

Liverpool, United Kingdom.

[3] Institute for Science and Technology in Medicine (ISTM), School of Pharmacy & Bioengineering,

Keele University, Staffordshire, United Kingdom.

[4] Exercise Metabolism and Adaptation Research Group, Research Institute for Sport and Exercise

Sciences, Liverpool John Moores University, Liverpool, United Kingdom.

[5] Exercise Biology Group, Faculty of Sport and Health Sciences, Technical University of Munich,

Munich, Germany.

[6] Centre for Genomics and Child Health, Blizard Institute, Barts and the London School of Medicine

and Dentistry, Queen Mary University of London, London, United Kingdom.

[7] Institute of Sport, Exercise and Health, University College London, London, United Kingdom

[8] Orthopedics Department, University Hospitals of the North Midlands, Staffordshire, UK.

[9] Institute for Health and Sport (iHeS), Victoria University, Footscray, Victoria, Australia.

[10] Department of Molecular Biology and Genetics, Federal Research and Clinical Center of Physical-

Chemical Medicine of Federal Medical Biological Agency, Moscow, Russia.

[11] Institute for Genomic Statistics and Bioinformatics, University Hospital Bonn, Bonn, Germany

[12] Laboratory of Exercise Physiology, Institute of Biomedical Problems of the Russian Academy of

Sciences, Moscow, Russia.

[13] Laboratory of Molecular Genetics, Kazan State Medical University, Kazan, Russia.

[14] School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences,

University of Birmingham.

[#]These authors contributed equally to the work.

*Corresponding author for Skeletal muscle tissue, stem cells and DNA / HOX methylation

a.p.sharples@googlemail.com

[$] Corresponding author for physical activity and HOX methylation

I.Ahmetov@ljmu.ac.uk

Abstract

Introduction: Skeletal muscle tissue and stem cells show global hypermethylation with aging. However, it is unknown whether the methylomes of muscle stem cells overlap that of skeletal muscle tissue and how age modulates this overlap. **Methods:** Using high coverage DNA methylation arrays (850K CpG sites) we compared the methylomes of young/aged human skeletal muscle to that of young/aged muscle stem cells at several time points of differentiation (0, 72 hrs, 7, 10 days). **Results:** Aged muscle tissue was hypermethylated compared with young tissue, enriched for; 'pathways-in-cancer' (that combines; focal adhesion, MAPK signaling, PI3K-Akt-mTOR signaling, p53 signaling, Jak-STAT signaling, TGF-beta and notch signaling), 'rap1-signaling', 'axon-guidance' and 'hippo-signalling'. Aged muscle stem cells also demonstrated a hypermethylated profile in pathways; 'axon-guidance', 'adherens-junction' and 'calcium-signaling' pathways, particularly at later timepoints of myotube formation, corresponding with reduced morphological differentiation and reductions in MyoD/Myogenin gene expression compared with young cells. While young cells showed less variability in DNA methylation during differentiation, aged cells demonstrated extensive and significantly varied DNA methylation at 7 days of differentiation in pathways; 'regulation-of-localisation', 'regulation-of-cell-communication' and 'regulation-of-signaling'. We also identified CpG sites whose hypermethylated state with age was retained in muscle cells *in-vitro* once isolated from tissue *in-vivo* (on genes KIF15, DYRK2, FHL2, MRPS33, ABCA17P). Differential chromosomal region analysis identified 3 regions with 6-8 CpGs in the Homeobox (HOX) transcription factor family of genes that were altered with age in muscle tissue and cells. With, HOXD10, HOXD9, HOXD8, HOXA3, HOXC9, HOXB1, HOXB3, HOXC-AS2 and HOXC10 all significantly differentially hypermethylated in aged tissue. In aged cells the same HOX genes (and additionally HOXC-AS3) also displayed the most varied methylation at 7 days of differentiation versus young cells. With HOXD8, HOXC9, HOXB1 and HOXC-AS3 hypermethylated and HOXC10 and HOXC-AS2 hypomethylated. We also determined inverse relationships between mRNA expression and DNA methylation in HOXB1, HOXA3 and HOXC-AS3. Finally, we were able to demonstrate that increased physical activity in young adults was associated with oppositely regulating HOXB1, HOXA3 and HOXC10 compared with age. **Conclusion:** For the first time, we demonstrate HOX genes were differentially epigenetically regulated in aged human skeletal muscle and stem cells and increased physical activity may help prevent the age-related epigenetic changes observed in the HOX genes.

88
89 **Introduction**
90
Maintaining skeletal muscle mass and function into older age is fundamental for human
health-span and quality of life [1]. Five to ten percent of older humans have sarcopenia [2],
which is characterized by reductions in muscle mass and strength [3]. This loss of muscle
mass and strength leads to frailty, increased incidence of falls, hospitalization and morbidity
[3-10]. Annual costs of fragility are estimated to be 39/32 billion (Euros/USD) for European
and USA fragility fractures respectively, with the cost of sarcopenia estimated to be £2 billion
in the UK [11]. With an advancing population these costs are only likely to increase with time.
98
A primary hallmark of ageing is the alteration of the epigenetic landscape. Epigenetics
encompasses the interaction between lifestyle/environmental factors and modifications to
DNA that affect gene function, without changes to the inherited DNA sequence [12, 13]. DNA
methylation is the most studied epigenetic modification, which involves the addition of a
covalent methyl group to the 5' position of the pyrimidine ring of a cytosine (5mC). Increased
methylation (hypermethylation) to cytosine-guanine (C-G) pairings (CpG sites), especially in
CpG-rich regions such as gene promoters, typically leads to reduced capacity for the
transcriptional apparatus to bind to these regions, suppressing gene expression [13].
Methylated CpG islands in promoters also lead to a tight compaction of adjacent chromatin
via the recruitment of chromatin modifying protein/protein-complexes, further silencing gene
expression. In contrast, reduced methylation (hypomethylation) provides a more favorable
DNA landscape for the transcriptional apparatus to bind to these regions, as well as more
'relaxed' chromatin, enabling gene expression to occur. Generally, albeit not in every
scenario, methylation levels therefore are inversely correlated with gene expression, where
hypermethylation is associated with reduced gene expression and hypomethylation with
increased gene expression.
115
DNA methylation in aged skeletal muscle occurs at tissue-specific genes [14]. However,
aged muscle also has the smallest overlap with other aged tissue types, suggesting skeletal
muscle may be unique in comparison with other tissues in its epigenetic aging processes
[14]. Indeed, it has recently been demonstrated that the methylation status of approximately
200 CpG sites can accurately predict chronological age in skeletal muscle tissue [15]. But
that this muscle 'clock' only shares 16 of these CpG's with the original 353 CpG pan-tissue
Horvath clock [15, 16]. Further, using DNA methylation arrays with coverage of ~450,000
CpG sites [17], Alibegovic *et al.* demonstrated that compared with young human skeletal
muscle, aged skeletal muscle is hypermethylated at a considerable number of CpG sites
across the genome. This suggests that aging may globally suppress gene expression in
skeletal muscle, and therefore may underpin impaired muscle maintenance into older age.
However, corresponding gene expression changers were not measured in this particular
paper and therefore warrant future investigation. Moreover, our group have demonstrated
that when mouse skeletal muscle stem cells were exposed to a high dose inflammatory
stress in early proliferative life, the cells retained hypermethylation of muscle specific
regulatory factor, MyoD, 30 population doublings later [18]. Suggesting that inflamed
proliferative aging in muscle stem cells leads to a retained accumulation of DNA methylation.
Finally, it is also worth suggesting that increased physical activity [19] or resistance exercise
has been reported to predominantly hypomethylate the genome in young skeletal muscle
[20, 21] which is in contrast to the hypermethylation observed with aging, suggestive that

exercise may provide a stimulus to reverse age-related changes in DNA methylation. Albeit,
this hypothesis warrants further attention.
138
Given skeletal muscle fibers are post-mitotic, as they contain terminally differentiated/fused
nuclei (myonuclei), repair and regeneration of skeletal muscle tissue is mediated by a
separate population of resident stem cells, called satellite cells, that can undergo mitosis.
Once activated these cells proliferate and migrate to site of injury to differentiate and fuse
with the existing fibers to enable repair. Indeed, altered DNA methylation of target genes
have been identified during the differentiation of muscle cells into myotubes *in-vitro* [22]. This
has included varied methylation of MyoD [23], Myogenin [24] and Six1 [25]. Furthermore,
muscle stem cell derived from aged individuals demonstrate similar proliferative capacity and
time to senescence as young adult cells [26, 27], but generally demonstrate impaired
differentiation and fusion into myotubes [28-44]. There have been a small number of studies
to the contrary, that suggested no effect of age on the differentiation capacity of isolated cells
[26, 45, 46]. However, the majority of studies demonstrate an impaired differentiation
phenotype in aged cells, yet despite this, there has been only one study that assessed DNA
methylation across the genome (450K CpG sites) in aged versus young adult muscle stem
cells [26]. This study identified that there was increased (hyper) methylation across the
genome in these cells as well as aged tissue [26].
155
To date, no investigation has compared DNA methylation across the genome during the
entire time-course of muscle cell differentiation, particularly using array technology which
permits greater coverage (850K CpG arrays) in aged muscle stem cells. Furthermore, to the
authors knowledge no one has investigated the differences in gene ontology, associated
enriched pathways, differentially methylated chromosomal regions or specific temporal
profiles of DNA methylation in aged muscle tissue or across differentiation compared with
young adults. Therefore, the objectives of the current study were: 1) To investigate the DNA
methylome of human aged and young adult skeletal muscle tissue and primary differentiating
muscle derived stem cells over an extensive time-course of; 0 h (30 minutes post transfer to
differentiation media), 72 h (hours), 7 d (days) and 10 d of differentiation (using 850K CpG
arrays) to identify aged associated epigenetic changes. 2) Identify methylation patterns that
are retained *in-vitro* by muscle stem cells from the aging tissue *in-vivo* niche. Finally, 3) to
identify if any of the most significant changes observed in DNA methylation with age in
muscle tissue and stem cells were also influenced by physical activity levels.
170
171 **Methods**
172
173 *Skeletal muscle biopsies and primary cell isolations*
174
For young adults (total n = 9, male, age 27 ± 4.4 years), skeletal muscle tissue (~150 mg)
was obtained from the vastus lateralis via a conchotome biopsy. Consent and ethical
approval were granted for the collection of muscle tissue under NREC, UK approval
(16/WM/010) or LJMU, UK local ethics committee approvals [H19/SPS/028 & H15/SPS/031].
Six (out of 9) of the young adult's tissue (male, 28 ± 5.3 years) baseline (at rest) array data
was derived from Seaborne *et al.* (2018). This is because we used this baseline (at rest)
tissue to derived cells (detailed below) for DNA methylation analysis experiments in the
present study. We also further added 3 new baseline (rested) young adult tissue biopsies
(male, 27 ± 2.5) for new DNA array analyses in the present study. For older adults (n = 5, 2

male/3 female, 83 ± 4 years), tissue biopsies were obtained during elective orthopedic
surgeries from University Hospitals of the North Midlands, from the vastus lateralis (knee
surgery, n = 2) or gluteus medius muscles (hip surgery, n = 3), under consent and ethical
approval 18/WM/0187. DNA and RNA were isolated from these young and aged tissue
samples. DNA samples from all 9 young adult and 5 aged adults were analysed for DNA
methylation arrays (detailed below), and a subset were analysed for gene expression (young
adult n = 4, aged adult n = 5). Primary skeletal muscle cells were derived from a subset of
young adult and aged tissue samples, and isolated as per our previous work [20, 21, 47-49].
Briefly, approximately 100 mg biopsy tissue was immediately (~10-30 mins) transferred to a
sterile class II biological safety cabinet in pre-cooled (4°C) transfer media (Hams F-10, 2% hi-
FBS, 100 U/ml penicillin, 100 µg/ml streptomycin and 2.5 µg/ml amphotericin-B). Any visible
connective and adipose tissue were removed using sterile scalpels and muscle tissue was
thoroughly washed 2 × in sterile PBS (containing 100 U/ml penicillin, 100 µg/ml streptomycin,
2.5 µg/ml amphotericin-B). PBS was removed and the muscle tissue was minced in the
presence of 0.05% Trypsin/0.02% EDTA and all contents (tissue and trypsin) were
transferred to a magnetic stirring platform at 37°C for 10 minutes. The addition of 0.05%
Trypsin/0.02% EDTA and titration was repeated on any remaining tissue. The supernatant
was collected from both procedures and horse serum (HS) was added at 10% of the total
supernatant volume to neutralize the trypsin. The supernatant was centrifuged at 340 g for 5
minutes where the supernatant and cell pellet were both plated in separate pre-gelatinised
(0.2% gelatin) T25 flasks containing 7.5 ml fresh pre-heated growth media/GM (GM; Ham's
F10 nutrient mix supplemented with 10% hi-NCS, 10% hi-FBS, 100 U/ml penicillin, 100 µg/ml
streptomycin, 2.5 µg/ml amphotericin B and 5 mM L-glutamine). Once confluent, cells were
trypsinised and reseeded into larger T75's to expand the cell population. Human derived
muscle cells (HMDCs) were seeded onto gelatin coated 6 well plates, at a density of 9×10^{5}
cells/ml in 2 ml of GM for until ~90% confluency was attained (~48 h). GM was removed, and
cells underwent 3 × PBS washes before switching to low serum differentiation media (DM;
Ham's F10 nutrient mix supplemented with 2% hiFBS, 100 U/ml penicillin, 100 µg/ml
streptomycin, 2.5 µg/ml amphotericin B and 5 mM L-glutamine). HMDCs were differentiated
for a total of 10 days (d) which received a 1 ml top up at 72 h and 7 d timepoints. Cells were
lysed for DNA and RNA at 0 h (30 minutes in DM), 72 h, 7 d and 10 d. All experiments were
carried out below passage 10 to prevent senescence. We undertook methylation arrays on
DNA isolated from: 0 h young (n = 7), 0 h aged (n = 4), 72 h young (n = 7), 72 h aged (n = 4),
7 d young (n = 6), 7d aged (n = 3), 10 d young (n = 2) and 10 d aged (n= 4). Gene
expression was analysed using young (n = 4, 7 d) and aged (n = 3, 7 d) cells.
219
Myogenicity and morphology measurements

Following isolations, attached single cells were fixed prior to staining and fluorescent
immunocytochemistry analysis to determine myogenicity (via desmin positivity) of the
isolated young and aged muscle derived cultures. Briefly, approximately 2×10^{4} cells were
seeded onto 3 × wells of a 6-well plate and were incubated for 24 h in GM. Existing media
was removed and cells were washed 3 × in PBS before fixation using the graded
methanol/acetone method (50:25:25 TBS:methanol:acetone for 15 minutes followed by 50:50
methanol:acetone) after which cells were permeabilised in 0.2% Triton X-100 and blocked in
5% goat serum (Sigma-Aldrich, UK) in TBS for 30 minutes. Cells were washed 3 × in TBS
and incubated overnight (4 °C) in 300 µl of TBS (with 2% goat serum and 0.2% Triton X-100)
containing primary anti-desmin antibody (1:50; ab15200, Abcam, UK). After overnight

incubation, cells were washed 3 × in TBS and incubated at RT for 3 h in 300 µl of secondary
antibody solution (TBS, 2% goat serum and 0.2% Triton X-100) containing the secondary
antibody, anti-rabbit TRITC (1:75; T6778, Sigma-Aldrich, UK) to counterstain desmin. Finally,
cells were washed again 3 × in TBS, prior to counterstaining nuclei using 300 µl of DAPI
solution at a concentration of (300 nM; D1306, Thermo Fisher Scientific, UK) for 30 minutes.
Immunostained cells were then visualised using a fluorescent microscope (Nikon, Eclipse Ti-
S, Japan or Olympus IX83, Japan) and imaged using corresponding software (Nikon, NIS
Elements and Olympus FV10-ASW 4.2). Myoblasts, myotubes and nuclei were visualized
using TRITC (Desmin, Excitation: 557 nm, Emission: 576 nm) DAPI (Excitation: 358 nm,
Emission: 461 nm) filter cubes. All immunostained samples were imported to Fiji/ImageJ
(version 2.0.0) software for subsequent calculations. Note, there was no difference (p = 0.86)
in myogenic cell proportions in aged (35 ± 5%) versus young cells (34 ± 9%). Therefore, the
aged and young muscle derived cells were matched for their desmin positivity (% myoblasts)
prior to differentiation experiments and downstream DNA methylation and gene expression
analysis. To determine myotube differentiation and formation at each time point, cells were
imaged using light microscopy (Olympus, CKX31, Japan) at 0, 72 h, 7 and 10 d of
differentiation.

DNA isolation and bisulfite conversion
249
Prior to DNA isolation, tissue samples were homogenized for 45 seconds at 6,000 rpm × 3 (5
minutes on ice in between intervals) in lysis buffer (180 µl buffer ATL with 20 µl proteinase K)
provided in the DNeasy spin column kit (Qiagen, UK) using a Roche Magnalyser instrument
and homogenization tubes containing ceramic beads (Roche, UK). Cells were lysed in 180 µl
PBS containing 20 µl proteinase K**. The DNA was then bisulfite converted using the EZ DNA
Methylation Kit (Zymo Research, CA, United States) as per manufacturer's instructions.

256
257 *Infinium MethylationEPIC BeadChip Array*
258
All DNA methylation experiments were performed in accordance with Illumina manufacturer
instructions for the Infinium Methylation EPIC BeadChip Array. Methods for the amplification,
fragmentation, precipitation and resuspension of amplified DNA, hybridisation to EPIC
beadchip, extension and staining of the bisulfite converted DNA (BCD) can be found in detail
in our open access methods paper [48]. EPIC BeadChips were imaged using the Illumina
iScan® System (Illumina, United States).

265
266 *DNA methylation analysis, CpG enrichment analysis (GO and KEGG pathways), differentially*
267 *modified region analysis and Self Organising Map (SOM) profiling*
268
Following DNA methylation quantification via MethylationEPIC BeadChip array, raw .IDAT
files were processed using Partek Genomics Suite V.7 (Partek Inc. Missouri, USA) and
annotated using the MethylationEPIC_v-1-0_B4 manifest file. We first checked the average
detection p-values. The highest average detection p-value for the samples was 0.0023
(Suppl. Figure **1a**), which is well below that recommended in the Oshlack workflow, of 0.01
[50]. We also produced density plots of the raw intensities/signals of the probes (Suppl.
Figure **1b**). These demonstrated that all methylated and unmethylated signals were over
11.5 (mean was 11.52 and median was 11.8), and the difference between median
methylation and median unmethylated signal was 0.56 [50]. Upon import of the data into

Partek Genomics Suite we removed probes that spanned X and Y chromosomes from the
analysis due to having both males and females in the study design, and although the
average detection p-value for each samples was on average very low (no higher than
0.0023) we also excluded any individual probes with a detection p-value that was above 0.01
as recommended in [50]. Out of a total of 865,860 probes, removal of the X&Y chromosome
probes and those with a detection p-value above 0.01 reduced the probe number to 846,233
(removed 19,627 probes). We also filtered out probes located in known single-nucleotide
polymorphisms (SNPs) and any known cross-reactive probes using previously defined SNP
and cross-reactive probe lists identified in earlier 850K validation studies [51]. This resulted
in a final list of 793,200 probes to be analysed. Following this, background normalisation was
performed via functional normalisation (with noob background correction), as previously
described [52]. Following functional normalisation, we also undertook quality control
procedures of principle component analysis (PCA), density plots by lines as well as box and
whisker plots of the normalised data for tissue samples (Suppl. Figures **1c, d, e** respectively)
and cell samples (Suppl. Figures 1 **f, g, h** respectively). Any outlier samples were detected
using PCA and analysing the normal distribution of β-values. Outliers were then removed if
they fell outside 2 standard deviations (SDs) (e.g. Suppl. Figure **1c**) using ellipsoids and or
demonstrated different distribution patterns to the samples of the same condition. Only one
young adult male tissue sample was removed due to being outside 2 standard deviations
outside samples from the same condition (Suppl. Figure **1c**; sample with a strikethrough
line). Following normalisation and quality control procedures, we undertook differential
methylation analysis by converting β-values to M-values (M-value = log2(β / (1 - β)), as M-
values show distributions that are more statistically valid for the differential analysis of
methylation levels [53]. We undertook a 1-way ANOVA for comparisons of young and aged
skeletal muscle tissue. For primary human muscle cells, we undertook a 2-way ANOVA for
age (young adult cells and aged cells) and time (0, 72 h, 7 and 10 d) and also explored the
ANOVA main effects for both age and time independently. We also performed planned
contrasts within the differentiation time-course in both young adult cells alone and aged cells
alone e.g. 0 h vs. 72 h, 0 h vs. 7d 0 h vs. 10 d. As well as planned contrasts for young versus
aged cells at each time point of differentiation (0 h vs. 0 h, 72 h vs. 72 h, 7 d vs. 7 d, 10 d vs.
10 d in aged vs. young adult cells respectively). From these ANOVA's and contrasts we then
generated CpG lists based on a statistical cut-off that used the False Discovery Rate (FDR)
of at least ≤ 0.05. Because in some analysis, an extremely large number of significantly
differentially methylated CpG sites were identified (e.g. over 100,000) with an FDR of ≤ 0.05,
on these occasions we also provide the data for a more stringent FDR of ≤ 0.01 or 0.001 or
FDR ≤ 0.05 'a change (difference in M-value) in methylation greater than 2'. This therefore
produced a smaller list of the most highly significant sites and also subsequently enabled
sensible pathway analysis depicting the most pertinent and significant CpGs/genes, even if
this made the sheer number of significant CpG sites less impressive. We specify when a
more stringent FDR than 0.05 was used in the methods text. On these significantly
differentially methylated CpG lists we then undertook CpG enrichment analysis within gene
ontology pathways and KEGG pathways [54-56] using Partek Genomics Suite and Partek
Pathway. We also undertook differentially methylated region analysis (e.g. identifies where
several CpG's are differentially methylated within a short chromosomal location/region) using
the Bioconductor package DMRcate (DOI: 10.18129/B9.bioc.DMRcate). Finally, in order to
plot temporal changes in methylation across the time-course of muscle cell differentiation we
undertook Self Organising Map (SOM) profiling of the change in mean methylation within
each condition using Partek Genomics Suite.

326
327
328
329 *RNA isolation, primer design & gene expression analysis*
330
Skeletal muscle tissue muscle was homogenised in tubes containing ceramic beads (MagNA
Lyser Green Beads, Roche, Germany) and 1 ml Tri-Reagent (Invitrogen, Loughborough, UK)
for 45 seconds at 6,000 rpm × 3 (and placed on ice for 5 minutes at the end of each 45
second homogenization) using a Roche Magnalyser instrument (Roche, Germany). Human
cells from differentiation experiments were also lysed in 300 µl Tri-Reagent for 5 minutes at
RT mechanically dissociated/lysed using a sterile scraper. RNA was then isolated as per
Invitrogen's manufacturer's instructions for Tri-reagent as described in our previous work
(REFS). Then a one-step RT-PCR reaction (reverse transcription and PCR) was performed
using QuantiFastTM SYBR$^®$ Green RT-PCR one-step kits on a Rotorgene 3000Q. Each
reaction was setup as follows; 4.75 µl experimental sample (7.36 ng/µl totalling 35 ng per
reaction), 0.075 µl of both forward and reverse primer of the gene of interest (100 µM stock
suspension), 0.1 µl of QuantiFast RT Mix (Qiagen, Manchester, UK) and 5 µl of QuantiFast
SYBR Green RT-PCR Master Mix (Qiagen, Manchester, UK). Reverse transcription was
initiated with a hold at 50°C for 10 minutes (cDNA synthesis) and a 5-minute hold at 95°C
(transcriptase inactivation and initial denaturation), before 40-50 PCR cycles of; 95°C for 10
sec (denaturation) followed by 60°C for 30 sec (annealing and extension). Primer sequences
for genes of interest and reference genes are included in Table 1. All genes demonstrated no
unintended targets via BLAST search and yielded a single peak after melt curve analysis
conducted after the PCR step above. All relative gene expression was quantified using the
comparative Ct (^{ΔΔ}Ct) method [57]. For human cell differentiation analysis via measurement
of myoD and myogenin, a pooled mean Ct for the 0 h young adult control samples were used
as the calibrator when comparing aged vs. young adult cells. This approach demonstrated a
low % variation in C_t value of 9.5 and 8.5% for myoD and myogenin, respectively. For HOX
gene analysis between young and aged tissue and for the 7 d aged cells vs. 7 d young adult
cells, the mean Ct of the young adult cells were used as the calibrator. The average,
standard deviation and variations in Ct value for the B2M reference gene demonstrated low
variation across all samples (mean ± SD, 13.12 ± 0.98, 7.45% variation) for the analysis of
myoD and myogenin. For HOX gene analysis, the RPL13a reference gene variation was low
in the human tissue (17.77 ± 1.71, 9.6% variation) and cell (15.51 ± 0.59, 3.82% variation)
experiments. The average PCR efficiencies of myoD and myogenin were comparable (94.69
± 8.9%, 9.4% variation) with the reference gene B2M (89.45 ± 3.76%, 4.2% variation). The
average PCR efficiencies of the all genes of interest in the tissue analysis of the HOX genes
were also comparable (90.87 ± 3.17%, 3.39% variation) with the human reference gene
RPL13a (92 ± 2.67%, 2.9% variation). Similarly, for the cell analysis, HOX genes of interest
efficiencies were comparable (89.59 ± 4.41%, 4.93% variation 4.93%) with the reference
gene RPL13a (89.57 ± 3.55%, 3.97% variation).
367
368 *Physical Activity and DNA methylation*
369
The human association studies involved 30 physically active and endurance- oriented men of
Eastern European descent (age 32.9 ± 9.9 years). The study was approved by the Ethics
Committee of the Physiological Section of the Russian National Committee for Biological
Ethics and Ethics Committee of the Federal Research and Clinical Center of Physical-

chemical Medicine of the Federal Medical and Biological Agency of Russia. Written informed
consent was obtained from each participant. The study complied with the guidelines set out
in the Declaration of Helsinki and ethical standards in sport and exercise science research.
Physical activity was assessed using questionnaire. Participants were classified as mildly
active (1-2 training sessions per week, n=6), moderately active (3-4 training sessions per
week, n=8) and highly active (5-7 sessions per week, n=16). DNA methylation of CpG-sites
in the muscle fibres of the vastus lateralis was evaluated using Infinium MethylationEPIC
BeadChip Array (Illumina, USA). Bisulfite conversion of genomic DNA (500 ng of DNA was
taken into the reaction) was performed using the EpiMark® Bisulfite Conversion Kit in
accordance with the manufacturer's instructions. For EPIC BeadChips imaging the Illumina
iScan® System (Illumina, United States) was used. Preprocessing of raw *.idat files was
performed with the R/Bioconductor package RnBeads v 2.0.0. BMIQ algorithm was used for
correcting the bias of type-2 probe values. Greedycut algorithm was used for filtering out low-
quality probes. All probes with known SNPs and probes on a previously published blacklist
of cross-hybridized were also filtered [58]. The final analyses included 796,180 of 868,565
(92%) probes passed QC procedures. QC for samples was performed with Greedycut
algorithm from RnBeads package. Outliers were detected by PCA (exclusion criteria: an
absolute value of z-score of PC1 or PC2 > 5). All samples passed the QC. The methylation
level of each CpG-site after normalization and filtering processes was represented as a β-
value ranging from 0 (unmethylated) to 1 (fully methylated) in order to undertake multiple
regression (as regression analysis performed better with finite values. Ideally, if values range
from 0 to 1 with beta-values satisfying this criteria). Statistical analyses were conducted
using PLINK v1.90, R (version 3.4.3) and GraphPad InStat (GraphPad Software, Inc., USA)
software. All data are presented as mean (standard deviation). Multiple regression was used
for testing associations between the CpG-methylation level and physical activity adjusted for
age. P values < 0.05 were considered statistically significant to test the 54 HOX CpG sites
identified.
401
402
403 **Results**
404
405 *Aged human skeletal muscle tissue is hypermethylated compared with young adult tissue*
406
At an FDR ≤ 0.05 there were 17,994 CpG sites significantly differentially methylated CpGs in
aged versus young adult tissue (Suppl. File **1a**), with a more stringent FDR of ≤ 0.01 there
were 6,828 CpG sites (Suppl. File **1b**). Out of the most highly significant 6,828 CpG sites in
aged muscle, the majority (6,384) were hypermethylated with only 444 hypomethylated
versus young adult muscle. A finding that can be visualised via hierarchical clustering of the
aged and young adult tissue methylation data (Figure **1a)**. Furthermore, these CpG sites
were mainly located in CpG islands (5,018 out of 6,828 CpG's) with the remaining in N-
shores (354), S-Shelf (48), S-Shores (341), S-Shelf (69) and 'other' (998). Indeed, 4,976
CpG's out of the 5,018 CpG sites within CpG islands were also significantly hypermethylated
in aged compared with young adult tissue (Suppl. File **1c**). After gene ontology analysis, we
also confirmed that the majority of the CpG's enriched within the three overarching gene
ontology terms: 'Biological process' (Figure **1b**), 'cellular component;' (Figure **1c**) and
'molecular function' (Figure **1d**) demonstrated a predominantly hypermethylated profile in
aged compared with young adult muscle. Within the GO terms that include the search term
'muscle', the most significantly enriched was, 'regulation of muscle system process' (Figure

1e, CpG list Suppl. Figure **1e**). Within this GO term was also the term 'regulation of muscle
contraction' (CpG list Suppl. file **1f**). Once again, the predominant number of CpG's within
these muscle specific gene ontologies demonstrated a hypermethylated versus
hypomethylated profile in aged tissue compared with young adult tissue (e.g. 96 out of 110
CpG's and 70 out of 83 CpG's were hypermethylated in 'regulation of muscle system
process' and 'regulation of muscle contraction' respectively). Further, enrichment analyses of
methylation within KEGG pathways between aged and young adult tissue, identified the top
significantly differentially enriched pathways, as: 'Pathways in cancer', 'Rap1 signaling',
'Axon guidance', 'Hippo signaling' (Suppl. file **1g**). These enriched KEGG pathways also
displayed predominantly hypermethylated profiles in aged tissue. For example, within the top
enriched pathway, 'pathways in cancer', 266 out of 277 CpG's were hypermethylated and
only the remaining 11 hypomethylated in aged compared with young adult muscle tissue
(Figure **2**; Suppl. File **1h**). Further, following analysis of most significantly differentially
methylated CpG loci list for differentially methylated chromosomal regions in aged versus
young adult tissue (Suppl. File **1i**), the top 5 regions included: chr8:22422678-22423092 (415
bp) within a CpG island of the SORBS3 gene with 6 CpG's that were all hypermethylated in
aged versus young adult tissue. Also, chr6:30653732-30655720 (1989 bp) within a CpG
island of the PPPR1 gene contained 8 CpG's that were all hypermethylated compared with
young adult muscle. Chr20:13200939-13202437 (1499 bp) within a CpG island of the
promoter of gene ISM, contained 8 CpG sites were also all hypermethylated. Similarly, the
gene PDE4D1P on Chr1:144931376-144932480 (1105 bp) contained 6 CpG sites that
spanned its promoter within a CpG island, once again demonstrating hypermethylation. The
only gene demonstrating the opposite direction in methylation status (hypomethylation) in
aged tissue was the gene C1orf132, also on chromosome 1 (Chr1:207990896-207991936;
1041 bp), where 5 CpG's within this region were hypomethylated in opposition to the majority
of gene regions that were hypermethylated in aged versus young tissue.
448
449 *Aged primary human muscle stem cells displayed more varied DNA methylation signatures*
450 *versus young adults during the differentiation time-course*
451
We analysed DNA methylation from differentiating human muscle stem cells at 0 h (30
minutes post dropping to 2% serum), 72 h, 7 and 10 d. Both young and aged adult cells
morphologically demonstrated increasing differentiation and myotube formation across the
time-course (Figure **3a**). This was confirmed by increases in gene expression in myoD and
myogenin in both young adult and aged cells as differentiation progressed (Figure **3b**).
Indeed, myotube formation appeared not to be as extensive in the elderly cells, despite the
number of starting myogenic cells being the same in both aged and young cells. This
reduced capacity for differentiation in the aged cells was confirmed at the molecular level by
reduced myoD and myogenin gene expression at 72 h and 10 d in elderly cells compared
with the young adult cells (Figure **3b**). Indeed, as well as reduced myogenin in aged cells at
72h, any increases in myogenin gene expression were also delayed in aged cells compared
with young cells (Figure **3b**), with this previously been shown to be associated with impairing
the fusion process and myotube formation [31, 32]. Indeed, the interaction for a 2-way
ANOVA (Age x Time) generated a list of 40,854 CpG sites that were significantly altered with
both age and across the time-course of differentiation (FDR ≤ 0.05, Suppl. File **2a**). With a
more stringent FDR ≤ 0.01 and ≤ 0.001 there were still 9,938 and 2,020 CpG's significantly
differentially methylated respectively in aged cells versus young cells across all time points

(Suppl. File **2b** and **2c** respectively). Overall demonstrating that there were highly significant
differences in DNA methylation between aged and young cells during differentiation.
471
After determining these changes with both age and time. We next wished to assess if there
were changes in DNA methylation within each cell type (aged and young adult) across
differentiation, to identify if differentiation itself was changing DNA methylation profiles in the
aged or young adult cells. In order to do this, we further analysed the main effect for TIME (0,
72 h, 7 d, 10 d) however, this did not demonstrate any significant changes in DNA
methylation with an FDR ≤ 0.05. For example, even with a less stringent unadjusted p-value
of 0.001 there were only a small number of CpG's (333) considered significant across all
differentiation time points (Suppl. File **2d**). This suggested, that DNA methylation was not
considerably changing over the time course of muscle stem cell differentiation itself.
However, in order to investigate this more closely within aged and young cells at each time
point, we performed contrasts across differentiation, first in young adult cells (FDR ≤ 0.05)
only, and confirmed there was little to no changes in DNA methylation as the young cells
differentiated, with only 1 and 14 CpG's significantly modified at 72 h and 10 d versus
baseline levels at 0 h respectively, and with no CpG's significantly different at 7 d vs 0 h
(FDR ≤ 0.05). In aged cells similar to young cells, there were no differences observed early
after the induction of differentiation in DNA methylation profiles, with no CpG sites
significantly different at 72 h vs. 0 h (FDR ≤ 0.05). However, at the later time point of
differentiation, 7d, within aged cells, 2,785 CpG's were significant vs. 0 h (Suppl. file **2e**) and
404 CpG's were significantly different at 10 d vs. 0 h (FDR ≤ 0.05; Suppl. File **2f**). Even with
a higher stringency (FDR ≤ 0.05 and change/difference in methylation greater than 2) at 7
days of differentiation there were 1,229 significantly different CpG sites versus 0 h in aged
cells (Suppl. File **2g**). In this 1,229 CpG list there were balanced number of CpG's that were
hypo and hypermethylated (compared with, for example, the aged vs. young tissue analysis
above, where aged tissue demonstrated a predominantly profile hypermethylated versus
young cells). Indeed, there were slightly more CpG's hypomethylated (676) versus
hypermethylated (553) in the aged cells at 7 days of differentiation versus baseline levels at
0 h, where out of these CpG's, 73 were located in islands and 41 associated with promotors.
This analysis of time, therefore suggested, that between 0-72 h in both young and aged cells
there was little change in DNA methylation profiles. However, by 7 days in aged cells, DNA
methylation (both hypo and hypermethylation) was significantly varied, whereas in young
cells it remained unchanged. Overall, this may suggest a more dysfunctional differentiation
program in aged human muscle cells that is associated with the impaired differentiation and
myotube formation we observed morphologically (together with reduced myoD and delayed
myogenin expression) in comparison with the young adult cells. Conducting gene ontology
analyses at this 7 day time point between young and aged cells (Suppl. File **2h**), it suggested
that this varied methylation response in aged cells was enriched in GO terms: 'regulation of
localisation', 'regulation of cell communication' and 'regulation of signaling' (Figure **3c**; Suppl.
File **2i, j, k** respectively). CpG's within these ontologies also confirmed that there was a
similar hypo and hypermethylation profile in aged cells at 7 days. Further, KEGG pathway
analysis suggested the top enriched pathways at 7 days in aged cells versus 0 h were: 'axon
guidance', cholinergic synapse', 'adrenergic signaling in cardiomyocytes' and 'circadian
entrainment' (Suppl. File **2l**). Finally, differential region analysis between young and aged
cells at 7 days of differentiation identified two regions on 2 genes that had 5 or more CpG
sites significantly differentially methylated (Suppl. File **2m**). The first being Chr3:155394096-
155394303 (209 bp) located on gene PLCH1 that had 5 of its CpG's hypomethylated. The

second being a region non-annotated in location Chr2: 119615888-119617128 (1241 bp) that
was hypermethylated on 5 CpG's. Suggesting that enriched and varied methylation of these
genes occurred at 7 d differentiation in aged cells that was not detected at 7 days in young
adult cells.
521
522 *Aged primary human muscle cells demonstrate hypermethylated signatures versus young*
523 *adults across differentiation, particularly at 7 days.*
524
We next wished to identify the differences in DNA methylation in aged versus young cells
across differentiation at each time point. A one-way ANOVA with the main effect for 'age',
that analysed aged cells versus young cells alone (yet included all time points 0, 72 h, 7, 10
d of differentiation), generated a significant differentially methylated (FDR ≤ 0.05) CpG list of
269,898 CpG sites that significantly varied in aged cells versus young cells across all time
points. Even with a more stringent cut-off (FDR ≤ 0.01) there were still 159,235 sites
significantly modified (Suppl. File **3a**). Increasing the stringency to a change of greater than 2
while keeping an FDR ≤ 0.01 identified 2,719 CpGs with the most highly significant and
demonstrated the largest differences between aged and young cells (Suppl. File **3b**). As with
the aged versus young skeletal muscle tissue analysis, the majority of these significantly
modified CpG sites in aged cells were hypermethylated (2,017 out of 2,719) versus
hypomethylated (702 out of 2,719) compared with young cells, visualised in hierarchical
clustering of these CpG sites (Figure **4a**). Two hundred and eleven out of these 2,719 CpG's
were located in islands and 97 were promoter associated. Gene ontology identified that aged
cells demonstrated this significantly enriched hypermethylated profile in GO terms:
'developmental process', 'anatomical structure development', 'anatomical structure
morphogenesis' (Suppl. File **3c)** and also in KEGG pathways 'axon guidance', adherens
junction', 'calcium signaling', 'focal adhesion' and 'protein digestion and absorption' (Suppl.
File **3d).** Analysis of the significantly differentially methylated regions using the 2,719 CpG list
above (Suppl. File **3e**), containing the most CpG sites within a given region, identified that a
non-coding location on chr12:115134344-115135576 (1233 bp) that contained 13 CpG's that
were hypermethylated in aged cells versus young cells. Further, there were 7 CpG's that
were hypermethylated in the region of the gene LY6G5C (Chr6:31650736-31651158, 423
bp). There was also region containing 8 CpG's of the HOXC10 gene just upstream of
HOXC6 and MIR196 (Chr12:54383692-54385621, 1933 bp) that all demonstrated a
hypomethylated profile in aged cells vs. young cells. Interestingly, on the same chromosome
just upstream of the HOXC10 gene (Chr12:54376020-54377868, 1849 bp), within the
lncRNA HOXC-AS3, there were another 6 CpG's that were hypomethylated in aged cells
versus young cells. The concentration of hypomethylated CpG's in aged versus young cells
in HOXC genes is interesting given the majority of CpG's, more generally, were
hypermethylated in aged cells versus young cells. A finding that is analysed further in the
results below.
557
Given these changes, predominantly favouring hypermethylation (except the HOXC genes
identified above) in aged cells versus young cells, we also performed contrasts between
aged and young adult cells at each time point of differentiation. When we compared 0 h aged
cells with young adult cells at 0 h, there were significant changes in 1,569 CpG sites (FDR ≤
0.05, with a difference/change greater than 2), this included 738 significantly different CpG
sites. Out of these 738 CpG's, 582 were hypermethylated and only 156 hypomethylated in
aged versus young cells (Suppl. File **3f**). Gene ontology also confirmed that this

hypermethylated DNA profile in aged cells versus young adult cells at 0 hrs (Suppl. File **3g**)
was located in enriched gene ontology terms including: 'cytoskeletal protein binding' (Figure
4b; Suppl. File **3h**), 'developmental process', 'cell junction' and 'cytoskeleton' and 'actin
binding' (Suppl. File **3i, j, k, l** respectively). With KEGG analysis of 0 h aged versus 0 hr
young adult cells (Suppl. File **3m**) also suggesting that there were significantly enriched
hypermethylated pathways for 'Axon Guidance' (Figure **4c**; CpG List Suppl. File **3n**), 'Insulin
secretion', 'Phospholipase D signaling', 'cAMP signaling pathway' and 'Aldosterone synthesis
and secretion' (Suppl. Files **3o, p, q, r** respectively).
573
At 72 h after the induction of differentiation in aged primary human muscle cells versus
young adult cells, there were 1,418 significantly differentially methylated CpG sites (FDR ≤
0.05, with a difference/change greater than 2) there were 645 significant CpG sites between
aged and young adult cells (Suppl. File **4a**) again with a greater number hypermethylated
(478 out of 645) versus hypomethylated (167 out of 645) in aged vs. young cells at this time
point. Once again, as with the GO enrichment in the 0 h analysis above, at 72 h this analysis
also identified (Suppl. File **4b**); 'cytoskeletal protein binding' (Suppl. File **4c**), 'actin binding'
(Figure **4d**, Suppl. File **4d**), 'cytoskeleton' and 'development process' as most significantly
enriched, followed by 'regulation of signaling' (Suppl. Files **4e, f, g** respectively). Again, these
GO terms included CpG sites that were predominantly hypermethylated in aged muscle cells
at 72 h versus young adult cells at the same time point. Similarly, to the 0 h KEGG pathways
analysis, at 72 hrs (Suppl. File **4h**) this analysis also identified: 'Insulin secretion' (Suppl. File
4i), 'aldosterone synthesis and secretion' (Suppl. File 4j) and 'cAMP signaling pathway'
(Figure **4e**, Suppl. Files **4k**) as significantly enriched. Again, demonstrating overall a
predominately hypermethylated profile in aged cells at 72 h of differentiation versus young
cells at the same timepoint.
590
For 7 d aged vs. 7 d differentiated young cells (FDR ≤ 0.05, with a difference/change greater
than 2) there was the largest number of CpG sites significantly differentially methylated,
5,524 CpG sites (Suppl. File **5a**). This was considerably more than that identified between
aged and young cells at the other time points of 0, 72 h (above) and 10 d (below). This
included 4,061 (out of 5,524 CpG's) hypermethylated versus only 1,463 (out of 5,524 CpG's)
hypomethylated. Gene ontology suggested that this increased hypermethylation in aged cells
at 7 d versus young adult cells at 7 d (Suppl. File **5b**), were enriched within terms:
'developmental process' (Figure **5a**, Suppl. File. **5c**) 'anatomical structure morphogenesis'
(Suppl. File **5d**), 'neuron part', 'cytoskeletal protein binding' and 'cell junction' (Suppl. File **5d,**
e, f, g). KEGG analysis (Suppl. File **5h**) identified: 'Focal adhesion' (Figure **5b**; Suppl. file **5i**),
'adherens junction' (Suppl. File **5j**), 'regulation of actin cytoskeleton' (Suppl. File **5k**), 'cGMP-
PKG signaling pathway' Suppl. File **5l**), 'rap1 signaling pathway' (Suppl. File **5m**), as well as
the well-known muscle differentiation pathway of 'PI3K-Akt signaling pathway' (Suppl. File
5n) to be predominantly hypermethylated in aged cells at 7 d versus young cells at the same
time point. Analysis of the significantly differentially methylated regions using this 5,524 CpG
list at 7 d in aged cells versus 7 d young adult cells (Suppl. File **5o**) also identified that a
region of the a gene within the HOX family (identified in earlier analysis above), HOXB1,
located in Chr17:46607104-46608568, contained 8 CpG's that were hypermethylated with
this 1465 bp region. Also, that a location within Chr3:113160184-113160624 spanning 441
bp also contained 6 CpG's that were hypermethylated spanning the gene WDR52. As well as
a gene already identified to be hypermethylated in differentiation in aging cells only at 7 days,
the gene LY6G5C, this time spanning a slightly larger region Chr6:31650736-31651362, of

627 bp (vs. Chr6:31650736-31651158, 423 bp in the above analysis), confirming that 6 of its
CpG's were hypermethylated at 7 d compared with young adult cells at the same time point.
Finally, CTD-2035E11.3 and CTD-2201E18.3/ RPS23P5 located on Chr5:43019772-
43020628 (857 bp) and the gene NLRP1 (Chr17:5487376-5488048, 673 bp) were both
hypomethylated on 6 of their CpG's within those regions.
618
For 10 d aged vs. 10 d differentiated young cells with ((FDR ≤ 0.05 with a change greater
than 2) there were 288 significantly differentially methylated CpG sites (Suppl. File **6a**). Of
which 255 were hypermethylated and only 33 hypomethylated. Gene ontology of the 288
CpG list, suggested that this increased hypermethylation in aged cells at 10 d versus young
adult cells at 10 d (Suppl. File **6b**), were enriched within terms 'GDP binding' (Figure **5c**,
Suppl. File **6c**), 'phosphotransferase activity', 'positive regulation of antigen receptor',
'mesoderm morphogenesis' and 'actin binding' (Suppl. File **6d, e, f, g** respectively). KEGG
analysis (Suppl. File **6h**) identified 'Regulation of actin cytoskeleton' (Figure **5d**; Suppl. File
6i), 'ErbB signaling pathway' (Suppl. File **6j**) and 'Selenocompound metabolism' (Suppl. File
6k) to be predominantly hypermethylated in aged cells at 10 days versus young cells at the
same time point.
630
631   *SOM profiling of aged muscle stem cells confirms a varied methylation profile*
632
With the greatest number of significantly differentially methylated CpG's at 7 d in aged cells
versus the young. In order to visualise the temporal changes in methylation over the time
course of differentiation in aged versus young cells, we conducted SOM profiling analysis of
the 2,719 significantly differentially methylated CpG list generated above (those highly
significant for 'age', across all time-points of differentiation). SOM analysis averages the
methylation values for the group of samples within each condition to identify temporal profile
changes in significantly differentially methylated CpG's between aged and young cells over
the time-course of differentiation. Confirming the data above, it was also clear using this
analysis that aged cells demonstrated a more varied (and perhaps dysfunctional) methylation
profile earlier in the time-course of differentiation (between 72 h and 7d) compared with
young adult cells (Figure **6a**). When looking at aged cells temporal change individually
compared with young cells across the differentiation time-course, out of the 2,719 CpG list, in
aged cells 1,504 of them were hypomethylated at 7 days and 956 hypermethylated at 7 days
(confirming the aged cells 'time' main effect analysis above) (Figure **6a**). However, in
contrast, only a small number of genes demonstrated this altered profile at 7 days in young
cells (284 hypo and 110 hyper out of the 2,719 CpG list).
649
Finally, in order to identify common CpG changes across both tissue and cells and to
determine if there were any CpG sites demonstrating similar methylation profiles once
isolated *in-vitro* compared with the tissue they were derived from *in-vivo,* the significantly
differentially methylated CpG lists from young cells vs. aged cells across all time were
overlapped (2,719 CpG list), the 0 h aged vs. 0 h young (738 CpG list), 72 h aged vs. young
72 h (645 CpG list), 7 d aged vs. 7 d young (5,524 CpG list), and the 10 d aged vs 10d young
(288 CpG list) (Venn diagram, Figure **6b**). There were 24 genes that were identified across
both tissue and cell significantly differentially methylated CpG lists (Suppl. File **7a**). They
included 24 CpG's on 16 genes: TSPAN9, RBM22, UBAP1, CAPZB, ZNF549, MBNL2,
RMI1, CHRM5, RAB4A, C19orf21, MOBKL1A, ANAPC11, GAS7, PBX1, ELOVL2, FGGY.
Furthermore, generating a SOM profile of temporal change in methylation over the time-

course of differentiation in these 24 CpG's, visualised in (Figure 6**c**), also demonstrated that
the majority of these CpG's in aged cells at 7 days (16 out of 24 CpG's) demonstrated varied
methylated at 7 days, including 11 out of 16 CpG's for annotated genes: TSPAN9, RBM22,
UBAP1, CHRM5, C19orf21, ELOVL2, MOBKL1A, ANAPC11, PBX1, CAPZB, FGGY (all
detailed in Suppl. File **7a).**

Distinguishing differentiation specific CpG sites in aging cells versus those altered as a
consequence of age alone.

The data above suggest that aged cells across all time demonstrate hypermethylation versus
young adult cells. Also, that aged cells significantly altered their methylation profiles during
differentiation itself, particularly in a large number of CpG's at 7 days of differentiation versus
young cells. Therefore, we conducted further analysis on the overlap of differentially
methylated CpG's within aging cells at 7 days of differentiation (from the 'time' analysis
above) and those that were changed as a consequence of age alone at 7 days (aged cells at
7 days versus young cells at 7 d). This enabled the identification of which methylation sites
were altered, but also shared in both aged cell differentiation alone and as a consequence of
age. Or, alternatively the sites that were simply changed with age and not differentiation
process and vice versa. Indeed, overlapping the aged cells 0 h vs. 7 d (1,229 CpG list) with
the 7 d aged vs. 7 d young significant 5,524 CpG list, there were only 334 (206
hypermethylated, 128 hypomethylated) CpG's that were shared (Suppl. File. **8a**). This
suggested that differentiation itself modified only 334 CpG's (out of 1,229) in aged cells that
were also changed as a consequence of age alone (i.e. in aged vs. young cells at 7 days).
The remaining 895 CpG's (1,229 - 334 CpG's; Suppl. File **8b**) were differentiation specific to
aged cells. Out of this 895 CpG list, an equal number were hypo and hypermethylated (458
hypo and 437 hypermethylated). Therefore, this overlap analysis also confirmed that data
above, where over the time-course of aged cell differentiation itself the methylome is both
hypo and hyper methylated on a similar number of CpG's, whereas aging alone
predominantly hypermethylates (where out of 5,524 changed at 7 d young vs. aged, 4061
were hypermethylated and only 1,463 hypomethylated). This also suggested that the
remaining 5,190 (5,524 minus 334 CpG list, equalling 5,190 CpG's with 3,910
hypermethylated vs. 1,280 hypomethylated) were as a consequence of aging alone (Suppl.
File **8c**).

Hypermethylation for a small number of CpG sites is retained in aged muscle cells in-vitro
from the in-vivo tissue niche.

We next examined whether methylated profiles were maintained from the tissue once
isolated in an *in-vitro* setting. Indeed, 6 CpG's that were identified in the 6,828 significantly
differentially methylated CpG list from the aged versus young tissue analysis, as well as
highlighted in the 2,719 list 'age' cells (across all time) analysis (Figure **7**). These included:
KIF15 (2 CpG's Cg00702638 & cg24888989), DYRK2 (cg09516963), FHL2 (cg22454769),
MRPS33 (cg26792755), ABCA17P (cg02331561). Importantly, all of these genes were
hypermethylated in the tissue analysis as well as the cells, highlighting that the cells were
perhaps retaining their *in-vivo* hypermethylation one these particular CpG's once isolated *in-*
vitro. Given that aged cells demonstrated the most varied methylation at 7 days versus
young cells. When comparing 7 d aged vs. 7 d young cell list (5,524 list), 4 CpG's (out of the
6 CpG sites identified above) were also identified in the 6,828 tissue CpG list, including:

MGC87042 (cg00394316), C2orf70 (cg23482427), ABCA17P (cg02331561) and
cg27209395 (not on an annotated gene). Once more, all of these CpG's (with the exception
of C2orf70, cg23482427) were hypermethylated in the tissue analysis as well as the cells.
With ABCA17P (cg02331561) highlighted across all gene lists (Figure **6**).
713
714
715
716 *Varied methylation in the HOX family of genes in aging skeletal muscle tissue and stem cells*
717
In the above analyses in aged versus young cells (all time points) there were 8 CpG's with
altered methylation within the region of the HOXC10 gene just upstream of HOXC6 and
MIR196 (Chr12:54383692-54385621, 1933 bp). Also, on the same chromosome just
upstream of the HOXC10 gene (Chr12:54376020-54377868, 1849 bp) within lnRNA HOXC-
AS3 there were another 6 CpG's that were altered in aged cells versus young cells. Similarly,
using the 5,524 CpG list at 7 d in aged cells versus 7 d young adult cells (Suppl. File **5o**) we
also identified that a region of the HOXB1, located in Chr17:46607104-46608568 contained 8
CpG's that were hypermethylated with the 1465 bp region. Therefore, we further analysed all
the HOX gene changes in the significantly differentially methylated aged vs. young tissue
(6,828 CpG list) and identified that CpG's within HOXD10, HOXD9, HOXD8, HOXA3,
HOXC9, HOXB1, HOXB3, HOXC-AS2 and HOXC10 were significantly differentially
methylated across all analyses, including the aged vs. young tissue (6828 CpG list; Suppl.
File **9a**), aged versus young adult cells (all time) 2,719 CpG list (Suppl. File **9b**) as well as
the 7 d aged versus 7 d young cell analysis 5,524 CpG list (Suppl. File **9c**). The timepoint
that showed the largest number of significant changes during the time course of
differentiation. A Venn diagram analysis depicted the overlap in common gene symbols for
these three gene lists (Figure **8a,** Suppl. File **9d**). It was also demonstrated that the majority
of these HOX genes were hypermethylated in aged tissue (Suppl. File. **9a**; Figure **8b**). In the
cell analysis across all time, these HOX genes also displayed the most varied methylation at
7 days of differentiation in aged cells versus young cells, therefore confirming the varied
temporal profile in methylation described above at 7 d was also the case for these HOX
genes identified. Finally, when SOM-profiling these 9 HOX genes by symbol (17 CpG's as
some HOX genes contained more than one CpG site), over the time-course of differentiation
based on the 'age' cells (all-time) 2,719 significantly differentially methylated CpG list. Eight
CpG sites across HOX family genes: HOXD8 (cg18448949, cg00035316), HOXB1
(cg04904318, cg02497558, cg22660933, cg10558129), HOXC9 (cg02227188) and HOXB3
(cg09952002) were hypermethylated at 7 days, whereas and 7 CpG's across HOXC10
(cg20402783, cg20403938, cg16898193, cg16791250), HOXB3 (cg23014425, cg04800503),
HOXC-AS2 (cg09605287) were hypomethylated at 7 days (Figure **8c;** Suppl. File **9e**). This
meant distinct genes were hypermethylated (HOXD8, HOXC9, HOXB1) and hypomethylated
(HOXC10, HOXC-AS2) at 7 days in aged cells versus young adult cells, except for one of
these genes, HOXB3, that contained 1 CpG that was hypermethylated versus 2 that were
hypomethylated.
751
752 The HOX genes that changed in both the tissue and cell analysis (by gene symbol) at the
753 mRNA expression level (HOXD8, HOXA3, HOXC9, HOXB1, HOXB3, HOXC-AS2 and
754 HOXC10) were further analysed in order to associate the significant changes in DNA
755 methylation with corresponding gene expression. Interestingly, in aged tissue, despite
756 displaying hypermethylation in all these HOX genes versus young tissue, the genes were not

suppressed at the gene expression level, which may have been expected, yet all significantly elevated (Figure **9a**). However, in the aged cells when analysing these genes at the expression level at 7 days of differentiation (including HOXD8, HOXA3, HOXC9, HOXB1, HOXB3, HOXC-AS2 and HOXC10, as well as HOXC-AS3 identified in the above in the cell analysis only) (Figure **9b**), we identified that there was significantly reduced gene expression in gene HOXB1 (Figure **9b**), that corresponded with HOXB1 hypermethylation. Where in the 5,524 CpG list at 7 d in aged cells versus 7 d young adult cells (Suppl. File **5o**), we previously identified that a region of the HOXB1 located in Chr17:46607104-46608568, contained 8 CpG's that were hypermethylated, as well as HOXB1 Cg's: cg04904318, cg02497558, cg22660933, cg10558129 being identified as hypermethylated in the 2,719 significant main effect for 'age' cell CpG list above (Figure **8c**, Suppl. File **3b**). Further, there was significantly increased gene expression for HOXC-AS3, with this gene identified earlier in the analysis as being with hypomethylated. Indeed, hypomethylation occurred in 6 CpG's in a region upstream of the HOXC10 gene (Chr12:54376020-54377868, 1849 bp) within the HOXC-AS3 gene. Interestingly, HOXC10 also demonstrated an average increase in gene expression, however, it was not statistically significant (Figure **9b**). Finally, HOXA3 also demonstrated significantly reduced expression (Figure **9b**) with corresponding hypermethylation (yet at 10 not 7 days of differentiation) in aged cells (see Figure **8c**). Overall, HOXB1, HOXC-AS3 and HOXA3 demonstrated an inverse relationship with CpG methylation and gene expression in aged versus young cells.

Effect of physical activity on methylation status of HOX family genes

Next, given that aging generally hypermethylated the genome, we have tested the hypothesis that physical activity may be associated with a reverse these epigenetic changes in the *HOX* family of genes. We thus performed a multiple regression analysis using methylation data of 54 sites of *HOX* family genes (those across Suppl. File 9 a,b,c) and the level of physical activity of 30 endurance-trained men. We determined that highly active men had hypomethylated *HOXB1* (cg10558129, $P = 0.032$) and *HOXA3* (cg16406967, $P = 0.038$), *HOXA13* (cg04428700, $P = 0.017$) genes and hypermethylated *HOXC10* gene (cg20403938, $P = 0.045$) compared to less active men (adjusted for age). Given that we identified the opposite trend in aged muscle and cells, where that particularly HOXB1 and HOXA3 were hypermethylated in aged tissue and cells with reduced gene expression in aged cells, and HOXC10 was hypomethylated in aged tissue and cells and the average gene expression increased. These findings suggest that increase exercise is associated with reversed methylation status in these HOX genes compared to age-related changes.

Discussion

In the present study we aimed to investigate the methylome in aged skeletal muscle tissue and differentiating primary muscle stem cells compared with young adults, in order to identify important epigenetically regulated genes in both aging skeletal muscle tissue and stem cells. As with previous studies ([17, 26], and by using more recent, higher coverage, array technology, we identified that aged skeletal muscle tissue demonstrated a considerably hypermethylated profile compared with young adult tissue. We also demonstrated that these hypermethylated profiles in aged tissue were enriched in gene ontology pathways including, 'regulation of muscle system process' and KEGG pathways 'pathways in cancer', a pathway that incorporates previously well described molecular pathways in the regulation of skeletal

muscle such as; focal adhesion, MAPK signaling, PI3K-Akt-mTOR signaling, p53 signaling,
Jak-STAT signaling, TGF-beta and Notch signaling, as well as the other significantly
enriched pathways of 'rap1 signaling', 'axon guidance', and 'hippo signaling'. This was also
the first study to profile DNA methylation over the entire time-course of skeletal muscle
differentiation (0, 72 h, 7 and 10 d) using the highest coverage 850K methylation assays. In
primary cell cultures, isolated from aged and young adults matched for the proportion of
myogenic cells, we identified that (as observed at the tissue level) aged muscle stem cells
also demonstrated hypermethylated profiles versus young adult cells. This hypermethylation
was enriched in: 'axon guidance', 'adherens junction' and 'calcium signaling' pathways.
Furthermore, we identified that the process of cellular differentiation itself did not significantly
affect DNA methylation in young cells, however aged cells demonstrated varied methylation
profiles particularly at 7 d of differentiation in GO terms: 'regulation of localisation', 'regulation
of cell communication', and 'regulation of signaling'. Furthermore, in the majority of different
CpG sites that were altered during the process of differentiation alone, aged cells
demonstrated significantly hypermethylated profiles during differentiation when compared
with young adult cells, again, specifically at 7 d of differentiation including CpG's located
within: 'focal adhesion', 'adherens junction' and 'regulation of actin cytoskeleton' pathways,
as well as the well-known muscle differentiation pathway, 'PI3K-Akt signaling'. This
corresponded with reduced differentiation and myotube formation observed morphologically
as well as associated reductions in myoD and myogenin and delayed increases in myogenin
gene expression in aged compared with young adult cells. Furthermore, we were able to
identify that a small number of CpG sites hypermethylated in aged tissue were also
hypermethylated in aged cells, with CpG's located on genes: KIF15, DYRK2, FHL2,
MRPS33, ABCA17P. This suggested, that these CpG's retained their methylation status *in-
vitro* after being isolated from the *in-vivo* niche. However, it is worth noting that this was a
very small number of CpG's compared with the thousands of CpG sites significantly
differentially methylated in the aged tissue versus young adult tissue, and those also
observed to be significantly different in the aged versus young muscle cells. This was
suggestive that the majority of the hypermethylated CpG's observed at the aged tissue level
were generally not retained on the same CpG sites in the isolated aged muscle stem cells *in-
vitro*. This therefore also perhaps indicates that hypermethylation of DNA within myonuclei
(the predominant source of DNA in the tissue) maybe therefore unique to that observed in
the isolated aged muscle cells. Indeed, retention of methylation during aging has been
previously observed in artificially aged muscle cells [18]. In skeletal muscle tissue, retention
of DNA methylation has been observed after skeletal muscle growth, even after a
subsequent period of muscle loss, suggesting an epigenetic memory at the DNA methylation
level [20, 21, 48]. However, the relative contribution of methylation from myonuclei or satellite
cells (or other resident cell types in muscle) to this epigenetic memory effect, and how these
retained profiles can last (e.g. past 22 weeks in the Seaborne *et al.*, 2018 study) has not
been determined. However, interestingly based on the present studies data, we could
hypothesise that myonuclear hypermethylation in the tissue with age is quite different to the
hypermethylation observed in satellite cells (and daughter progeny). Overall, suggesting that
environmental stimuli and aging could affect methylation profiles in the myonuclei differently
than those in satellite cells. A hypothesis that requires further investigation, perhaps using
single-cell methylome analysis of the different cell populations resident in cells derived from
skeletal muscle tissue biopsies.
851

Importantly, in both tissue and stem cell analysis, we also identified that the homeobox
(HOX) family of genes were significantly enriched in differentially methylated region analysis,
showing several (e.g. 6-8) CpG's to be methylated within narrow chromosomal regions on
these genes, as well as being methylated in the same direction (hyper or hypomethylated) in
aged compared with young adults. In particular, we identified: HOXC10 (just upstream of
HOXC6) and HOXB1 as having several CpG's differentially methylated. Therefore, closer
analysis of all HOX gene associated CpG changes across both tissue and cell differentiation
data identified that CpG's located within: HOXD10, HOXD9, HOXD8, HOXA3, HOXC9,
HOXB1, HOXB3, HOXC-AS2 and HOXC10 were all significantly differentially methylated
across these analyses. In aged tissue the majority of these HOX genes were
hypermethylated. In the cell analysis, these HOX genes displayed the most varied
methylation at 7 days of differentiation in aged cells versus young cells. Furthermore, distinct
HOX genes were hypermethylated (HOXD8, HOXC9, HOXB1) and hypomethylated
(HOXC10, HOXC-AS2) at 7 days in aged cells versus young adult cells. In aged cells versus
young cells, gene expression analysis also demonstrated an inverse relationship with
hypermethylation of HOXB1 and HOXA3 and reduced gene expression, as well as
hypomethylation of HOXC-AS3 together with increased gene expression.
869
This data is interesting given that the HOX genes are evolutionary conserved members of
the homeobox superfamily, with 39 HOX genes found in the mammalian genome. They are
'homeotic' genes coding for transcription factors, with a fundamental role in the determination
of cellular identity. They were first shown to be important in embryogenesis in drosophila
melanogaster (fruit fly) [59]. In muscle they have been described to morphologically identify
the hindlimb tissues during development [60-63], but have also been demonstrated to be
activated in satellite cells [64, 65], and as markers of hindlimb derived myoblasts [64]. In
particular HOXC10, demonstrated 8 CpG's (just upstream of HOXC6 and MIR196;
Chr12:54383692-54385621, 1933 bp) that all demonstrated a hypomethylated signature in
aged versus young muscle stem cells. There were also 4 CpG sites hypomethylated,
particularly at 7 days of differentiation in aged versus young cells. Indeed, HOXC10 has been
identified to determine hindlimb identity [60, 62]. Together with HOXC10 hypomethylation, we
also demonstrated average (yet not significant) increases in HOXC10 gene expression at 7
days in aged cells versus young cells. Counterintuitively, to our data, previously HOXC10
upregulation has been associated with satellite cell activation in skeletal muscle in response
to Roux-en-Y gastric bypass (RYGB) surgery [66], as well as being a marker for hindlimb
specific satellite cells. Interestingly, there was lower expression of HOXC10 observed in
exercised rats [67], which is perhaps more intuitive with the data provided here, where aged
cells demonstrated an increase. However, HOXC10 requires more experimentation to define
its mechanistic role in aging skeletal muscle, with HOXC10 and physical exercise being
discussed below. Interestingly, the hypomethylation of the HOXC10 (Chr12:54383692-
54385621, 1933 bp) occurred in 8 CpG's and on the same chromosome just upstream of the
HOXC10 gene (Chr12:54376020-54377868, 1849 bp), within the lncRNA HOXC-AS3, where
there were another 6 CpG's that were hypomethylated in aged cells versus young cells (See
Suppl. Figure **2** for visualisation of HOXC10 and HOXC-AS3 and their genomic location and
methylation). HOXC-AS3 is a long coding RNA (lcRNA) with currently no known data in
skeletal muscle, but some literature within cancer and mesenchymal stem cell (MSC) fields
[68]. Indeed, MSC's administered with silencing HOXC-AS3 prevented bone loss by
enhancing HOXC10 expression [69], and HOXC-AS3 upregulation has also been linked with
aggressive cancer [70]. Interestingly, we were able to identify that together with associated

hypomethylation, in a region close to the lcRNA HOXC-AS3, there was significantly increased gene expression of HOXC-AS3 (and an average, yet not significant increase in HOXC10 gene expression) in aged muscle cells at 7 days of differentiation. Given the data above in bone and cancer, HOXC-AS3 upregulation appears to be pro-growth and linked with expression of HOXC10, therefore their increase in the current study maybe hypothesised to be a co-operative and compensatory drive to maintain aged muscle. However, this hypothesis is speculative, and more work needs to be conducted as to the role of HOXC10 and HOXC-AS3 and their potential cooperative mechanisms of action in aged skeletal muscle. We also identified that HOXB1 was hypermethylated with increased gene expression in aged cells at 7 days. HOXB1 has been demonstrated to be hypermethylated in inflamed muscle of children with Juvenile Dermatomyositis (JDM) [71]. This is interesting given aged skeletal muscle is known to be chronically inflamed [72, 73], where we also demonstrate this hypermethylated profile in methylation in aged cells. Finally, HOXA3 was hypermethylated with reduced gene expression in aged cells. However, there is currently little to no work on this gene in skeletal muscle [63] and therefore requires future investigation.

Finally, with aging evoking a hypermethylated signature in tissue and aged muscle stem cells, it was also interesting to speculate that physical exercise, that has been shown to hypomethylate the genome [19-21], could therefore be 'anti-ageing' at the epigenetic level. Indeed, this hypothesis was first supported indirectly in the present study and by previous literature, where the aged tissue analysis in the present study identified the top significantly enriched KEGG pathway as, 'pathways in cancer'. A pathway that incorporates well known pathways important in skeletal muscle, including: Focal adhesion, MAPK, PI3K-Akt, mTOR, p53 signaling, Jak-STAT, TGF-beta and Notch signaling. Furthermore, this pathway was also the top enriched hypomethylated pathway after acute and chronic resistance exercise [21]. Overall, suggesting exercise (resistance exercise in this case) could perhaps reverse the hypermethylated profiles in these pathways in aged muscle. Therefore, in the present study, given that we identified the HOX family of genes to be extensively differentially methylated in aged tissue and stem cells, we went on to determine that higher physical activity levels (endurance exercise in this case) in healthy young adults was associated with changes in HOXB1, HOXC10 and HOXA3 DNA methylation in the opposite direction to the changes we observed in aged tissue and stem cells. This also provided evidence to suggest that increased physical activity could perhaps reverse the age-related epigenetic changes in the HOX genes. However, more research into the effect of exercise in an aged population and the changes in HOX methylation status will be required in the future to confirm these findings.

Conclusion
Overall, for the first time, we demonstrate that altered methylation of a large number of the HOX genes are epigenetically differentially regulated in aged human skeletal muscle tissue and during impaired differentiation in aged muscle stem cells. In particular HOXB1, HOXA3 and HOXC-AS3 (and to a certain extent, HOXC10) also demonstrated significant changes in gene expression in aged cells. Importantly, that increased physical activity may help prevent the age-related epigenetic changes observed in these HOX genes.

Acknowledgments

These data were supported by a North Staffordshire Medical Institute (NMSI) grant awarded
to Adam P. Sharples (PI), Daniel Turner, Mark Kitchen and Ian Dos-Remedios (Co-I's). This
work was also supported by the UK's Engineering and Physical Sciences Research Council
(EPSRC) and the UK Medical Research Council's (MRC) centre for doctoral training, via a
studentship awarded to the joint first author Piotr Gorski in the group of Adam P. Sharples
(PI). Funds from Keele University and Liverpool John Moores University, UK and The
Norwegian School of Sport Sciences, Oslo, Norway also supported the PhD work by Daniel
Turner & Piotr Gorski in the group of Adam P. Sharples. Philipp Baumert received a fully-
funded Liverpool John Moores PhD scholarship. Mohd Firdaus Maasar received a PhD
studentship via the Malasyisan government agency: Majlis Amanah Rakyat (MARA) via
Barry Drust, Adam P. Sharples and Dr. Andrew Hulten. The physical activity and DNA
methylation study was supported in part by grant from the Russian Science Foundation
(Grant No. 17-15-01436: "Comprehensive analysis of the contribution of genetic, epigenetic
and environmental factors in the individual variability of the composition of human muscle
fibers"; DNA sample collection, genotyping, epigenetic analysis and muscle fibre typing of
Russian subjects).

963
964
965

References

1. Sharples AP, Hughes DC, Deane CS, Saini A, Selman C, and Stewart CE. Longevity and
skeletal muscle mass: the role of IGF signalling, the sirtuins, dietary restriction and
protein intake. Aging Cell. 2015; 14(4):511-23.
2. Morley JE, Anker SD, and von Haehling S. Prevalence, incidence, and clinical impact of
sarcopenia: facts, numbers, and epidemiology-update 2014. Journal of cachexia,
sarcopenia and muscle. 2014; 5(4):253-259.
3. Hughes VA, Frontera WR, Wood M, Evans WJ, Dallal GE, Roubenoff R, et al.
Longitudinal muscle strength changes in older adults: influence of muscle mass,
physical activity, and health. Journals of Gerontology Series A-Biological Sciences &
Medical Sciences. 2001; 56(5):B209-17.
4. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al.
Sarcopenia: European consensus on definition and diagnosis: Report of the European
Working Group on Sarcopenia in Older People. Age Ageing. 2010; 39(4):412-23.
5. Hairi NN, Cumming RG, Naganathan V, Handelsman DJ, Le Couteur DG, Creasey H, et
al. Loss of muscle strength, mass (sarcopenia), and quality (specific force) and its
relationship with functional limitation and physical disability: the Concord Health and
Ageing in Men Project. J Am Geriatr Soc. 2010; 58(11):2055-62.
6. Cederholm T, Cruz-Jentoft AJ, and Maggi S. Sarcopenia and fragility fractures. Eur J
Phys Rehabil Med. 2013; 49(1):111-7.
7. Cooper C, Fielding R, Visser M, van Loon LJ, Rolland Y, Orwoll E, et al. Tools in the
assessment of sarcopenia. Calcif Tissue Int. 2013; 93(3):201-10.
8. Porter MM, Vandervoort AA, and Lexell J. Aging of human muscle: structure, function
and adaptability. Scandinavian journal of medicine & science in sports. 1995;
5(3):129-142.
9. Cruz-Jentoft AJ, Landi F, Topinkova E, and Michel JP. Understanding sarcopenia as a
geriatric syndrome. Curr Opin Clin Nutr Metab Care. 2010; 13(1):1-7.

10.   Beaudart C, Zaaria M, Pasleau F, Reginster J-Y, and Bruyère O. Health Outcomes of
Sarcopenia: A Systematic Review and Meta-Analysis. PLOS ONE. 2017;
12(1):e0169548.
11.   Fuggle N, Shaw S, Dennison E, and Cooper C. Sarcopenia. Best practice & research.
Clinical rheumatology. 2017; 31(2):218-242.
12.   Sharples AP, Stewart CE, and Seaborne RA. Does skeletal muscle have an 'epi'-
memory? The role of epigenetics in nutritional programming, metabolic disease,
aging and exercise. Aging Cell. 2016; 15(4):603-16.
13.   Sharples AP, Seaborne RA, and Stewart CE, *Epigenetics of Skeletal Muscle Aging*, in
Epigenetics of Aging and Longevity, A.M. Vaiserman, Editor. 2018, Academic Press:
Boston. p. 389-416.
14.   Day K, Waite LL, Thalacker-Mercer A, West A, Bamman MM, Brooks JD, et al.
Differential DNA methylation with age displays both common and dynamic features
across human tissues that are influenced by CpG landscape. Genome Biology. 2013;
14(9):R102.
15.   Voisin S, Harvey N, Haupt L, Griffiths L, Ashton K, Coffey V, et al. An epigenetic clock
for human skeletal muscle. bioRxiv. 2019:821009.
16.   Horvath S. DNA methylation age of human tissues and cell types. Genome biology.
2013; 14(10):3156.
17.   Zykovich A, Hubbard A, Flynn JM, Tarnopolsky M, Fraga MF, Kerksick C, et al.
Genome-wide DNA methylation changes with age in disease-free human skeletal
muscle. Aging Cell. 2014; 13(2):360-6.
18.   Sharples AP, Polydorou I, Hughes DC, Owens DJ, Hughes TM, and Stewart CE. Skeletal
muscle cells possess a 'memory' of acute early life TNF-alpha exposure: role of
epigenetic adaptation. Biogerontology. 2016; 17(3):603-17.
19.   Sailani MR, Halling JF, Møller HD, Lee H, Plomgaard P, Pilegaard H, et al. Lifelong
physical activity is associated with promoter hypomethylation of genes involved in
metabolism, myogenesis, contractile properties and oxidative stress resistance in
aged human skeletal muscle. Scientific Reports. 2019; 9(1):3272.
20.   Seaborne RA, Strauss J, Cocks M, Shepherd S, O'Brien TD, van Someren KA, et al.
Human Skeletal Muscle Possesses an Epigenetic Memory of Hypertrophy. Scientific
Reports (Nature) 2018; 8(1):1898.
21.   Turner DC, Seaborne RA, and Sharples AP. Comparative Transcriptome and
Methylome Analysis in Human Skeletal Muscle Anabolism, Hypertrophy and
Epigenetic Memory. Scientific Reports. 2019; 9(1):4251.
22.   Tsumagari K, Baribault C, Terragni J, Varley KE, Gertz J, Pradhan S, et al. Early de novo
DNA methylation and prolonged demethylation in the muscle lineage. Epigenetics.
2013; 8(3):317-32.
23.   Brunk BP, Goldhamer DJ, and Emerson JCP. Regulated Demethylation of
themyoDDistal Enhancer during Skeletal Myogenesis. Developmental Biology. 1996;
177(2):490-503.
24.   Fuso A, Ferraguti G, Grandoni F, Ruggeri R, Scarpa S, Strom R, et al. Early
demethylation of non-CpG, CpC-rich, elements in the myogenin 5'-flanking region: a
priming effect on the spreading of active demethylation. Cell Cycle. 2010; 9(19):3965-
76.

25.   Wu W, Ren Z, Wang Y, Chao Z, Xu D, and Xiong Y. Molecular characterization,
1039   expression patterns and polymorphism analysis of porcine Six1 gene. Molecular
1040   Biology Reports. 2011; 38(4):2619-2632.

1041   26.   Bigot A, Duddy WJ, Ouandaogo ZG, Negroni E, Mariot V, Ghimbovschi S, et al. Age-
1042   Associated Methylation Suppresses SPRY1, Leading to a Failure of Re-quiescence and
1043   Loss of the Reserve Stem Cell Pool in Elderly Muscle. Cell Rep. 2015; 13(6):1172-82.

1044   27.   Barberi L, Scicchitano BM, De Rossi M, Bigot A, Duguez S, Wielgosik A, et al. Age-
1045   dependent alteration in muscle regeneration: the critical role of tissue niche.
1046   Biogerontology. 2013; 14(3):273-292.

1047   28.   Hidestrand M, Richards-Malcolm S, Gurley CM, Nolen G, Grimes B, Waterstrat A, et
1048   al. Sca-1-expressing nonmyogenic cells contribute to fibrosis in aged skeletal muscle.
1049   J Gerontol A Biol Sci Med Sci. 2008; 63(6):566-79.

1050   29.   Pietrangelo T, Puglielli C, Mancinelli R, Beccafico S, Fano G, and Fulle S. Molecular
1051   basis of the myogenic profile of aged human skeletal muscle satellite cells during
1052   differentiation. Exp Gerontol. 2009; 44(8):523-31.

1053   30.   Beccafico S, Riuzzi F, Puglielli C, Mancinelli R, Fulle S, Sorci G, et al. Human muscle
1054   satellite cells show age-related differential expression of S100B protein and RAGE.
1055   Age (Dordr). 2010; 33(4):523-41.

1056   31.   Sharples AP, Al-Shanti N, and Stewart CE. C2 and C2C12 murine skeletal myoblast
1057   models of atrophic and hypertrophic potential: relevance to disease and ageing? J
1058   Cell Physiol. 2010; 225(1):240-50.

1059   32.   Sharples AP, Al-Shanti N, Lewis MP, and Stewart CE. Reduction of myoblast
1060   differentiation following multiple population doublings in mouse C2 C12 cells: a
1061   model to investigate ageing? J Cell Biochem. 2011; 112(12):3773-85.

1062   33.   Sharples AP, Player DJ, Martin NR, Mudera V, Stewart CE, and Lewis MP. Modelling in
1063   vivo skeletal muscle ageing in vitro using three-dimensional bioengineered
1064   constructs. Aging Cell. 2012; 11(6):986-95.

1065   34.   Merritt EK, Stec MJ, Thalacker-Mercer A, Windham ST, Cross JM, Shelley DP, et al.
1066   Heightened muscle inflammation susceptibility may impair regenerative capacity in
1067   aging humans. J Appl Physiol. 2013; 115(6):937-48.

1068   35.   Zwetsloot KA, Childs TE, Gilpin LT, and Booth FW. Non-passaged muscle precursor
1069   cells from 32-month old rat skeletal muscle have delayed proliferation and
1070   differentiation. Cell Prolif. 2013; 46(1):45-57.

1071   36.   Bigot A, Jacquemin V, Debacq-Chainiaux F, Butler-Browne GS, Toussaint O, Furling D,
1072   et al. Replicative aging down-regulates the myogenic regulatory factors in human
1073   myoblasts. Biol Cell. 2008; 100(3):189-99.

1074   37.   Carlson ME and Conboy IM. Loss of stem cell regenerative capacity within aged
1075   niches. Aging Cell. 2007; 6(3):371-82.

1076   38.   Lancioni H, Lucentini L, Palomba A, Fulle S, Micheli MR, and Panara F. Muscle actin
1077   isoforms are differentially expressed in human satellite cells isolated from donors of
1078   different ages. Cell Biol Int. 2007; 31(2):180-5.

1079   39.   Lees SJ, Rathbone CR, and Booth FW. Age-associated decrease in muscle precursor
1080   cell differentiation. Am J Physiol Cell Physiol. 2006; 290(2):C609-15.

1081   40.   Lorenzon P, Bandi E, de Guarrini F, Pietrangelo T, Schafer R, Zweyer M, et al. Ageing
1082   affects the differentiation potential of human myoblasts. Exp Gerontol. 2004;
1083   39(10):1545-54.

41.   Charge SB, Brack AS, and Hughes SM. Aging-related satellite cell differentiation
defect occurs prematurely after Ski-induced muscle hypertrophy. Am J Physiol Cell
Physiol. 2002; 283(4):C1228-41.
42.   Allen RE, McAllister PK, Masak KC, and Anderson GR. Influence of age on
accumulation of alpha-actin in satellite-cell-derived myotubes in vitro. Mech Ageing
Dev. 1982; 18(1):89-95.
43.   Kandalla PK, Goldspink G, Butler-Browne G, and Mouly V. Mechano Growth Factor E
peptide (MGF-E), derived from an isoform of IGF-1, activates human muscle
progenitor cells and induces an increase in their fusion potential at different ages.
Mech Ageing Dev. 2011; 132(4):154-62.
44.   Corbu A, Scaramozza A, Badiali-DeGiorgi L, Tarantino L, Papa V, Rinaldi R, et al.
Satellite cell characterization from aging human muscle. Neurol Res. 2010; 32(1):63-
72.
45.   Shefer G, Van de Mark DP, Richardson JB, and Yablonka-Reuveni Z. Satellite-cell pool
size does matter: defining the myogenic potency of aging skeletal muscle.
Developmental Biology. 2006; 294(1):55-66.
46.   Alsharidah M, Lazarus NR, George TE, Agley CC, Velloso CP, and Harridge SD. Primary
human muscle precursor cells obtained from young and old donors produce similar
proliferative, differentiation and senescent profiles in culture. Aging Cell. 2013;
12(3):333-44.
47.   Seaborne RA, Hughes DC, Turner DC, Owens DJ, Baehr LM, Gorski P, et al. UBR5 is a
novel E3 ubiquitin ligase involved in skeletal muscle hypertrophy and recovery from
atrophy. J Physiol. 2019; 597(14):3727-3749.
48.   Seaborne RA, Strauss J, Cocks M, Shepherd S, O'Brien TD, Someren KAV, et al.
Methylome of human skeletal muscle after acute & chronic resistance exercise
training, detraining & retraining. Scientific Data (Nature). 2018; 5:180213.
49.   Turner DC, Kasper AM, Seaborne RA, Brown AD, Close GL, Murphy M, et al, *Exercising*
Bioengineered Skeletal Muscle In Vitro: Biopsy to Bioreactor, in *Methods in Molecular*
Biology. 2019. p. 1889; 55-79 (Doi: 10.1007/978-1-4939-8897-6_5).
50.   Maksimovic J, Phipson B, and Oshlack A. A cross-package Bioconductor workflow for
analysing methylation array data. F1000Research. 2016; 5(1281).
51.   Pidsley R, Zotenko E, Peters TJ, Lawrence MG, Risbridger GP, Molloy P, et al. Critical
evaluation of the Illumina MethylationEPIC BeadChip microarray for whole-genome
DNA methylation profiling. Genome Biology. 2016; 17(1):208.
52.   Maksimovic J, Gordon L, and Oshlack A. SWAN: Subset-quantile within array
normalization for illumina infinium HumanMethylation450 BeadChips. Genome Biol.
2012; 13(6):R44.
53.   Du P, Zhang X, Huang C-C, Jafari N, Kibbe WA, Hou L, et al. Comparison of Beta-value
and M-value methods for quantifying methylation levels by microarray analysis. BMC
Bioinformatics. 2010; 11(1):587.
54.   Kanehisa M and Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic
Acids Res. 2000; 28(1):27-30.
55.   Kanehisa M, Furumichi M, Tanabe M, Sato Y, and Morishima K. KEGG: new
perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res. 2017;
45(D1):D353-d361.
56.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, and Tanabe M. KEGG as a reference
resource for gene and protein annotation. Nucleic Acids Res. 2016; 44(D1):D457-62.

57. Schmittgen TD and Livak KJ. Analyzing real-time PCR data by the comparative C(T)
method. Nat Protoc. 2008; 3(6):1101-8.
58. McCartney DL, Walker RM, Morris SW, McIntosh AM, Porteous DJ, and Evans KL.
Identification of polymorphic and off-target probe binding sites on the Illumina
Infinium MethylationEPIC BeadChip. Genomics Data. 2016; 9:22-24.
59. Shah N and Sukumar S. The Hox genes and their roles in oncogenesis. Nat Rev
Cancer. 2010; 10(5):361-71.
60. DeLaurier A, Schweitzer R, and Logan M. Pitx1 determines the morphology of muscle,
tendon, and bones of the hindlimb. Dev Biol. 2006; 299(1):22-34.
61. Carpenter EM, Goddard JM, Davis AP, Nguyen TP, and Capecchi MR. Targeted
disruption of Hoxd-10 affects mouse hindlimb development. Development. 1997;
124(22):4505-14.
62. Logan M and Tabin CJ. Role of Pitx1 upstream of Tbx4 in specification of hindlimb
identity. Science. 1999; 283(5408):1736-9.
63. Manley NR and Capecchi MR. Hox group 3 paralogous genes act synergistically in the
formation of somitic and neural crest-derived structures. Dev Biol. 1997; 192(2):274-
88.
64. Choi HY, Jeon H, Yang JW, Ahn JH, and Jung JH. Thyroid-Stimulating Hormone
Receptor Expression on Primary Cultured Human Extraocular Muscle Myoblasts. Curr
Eye Res. 2018; 43(12):1484-1488.
65. Seale P, Ishibashi J, Holterman C, and Rudnicki MA. Muscle satellite cell-specific
genes identified by genetic profiling of MyoD-deficient myogenic cell. Dev Biol. 2004;
275(2):287-300.
66. Tamboli RA, Hajri T, Jiang A, Marks-Shulman PA, Williams DB, Clements RH, et al.
Reduction in inflammatory gene expression in skeletal muscle from Roux-en-Y gastric
bypass patients randomized to omentectomy. PLoS One. 2011; 6(12):e28577.
67. Kato H, Shibahara T, Rahman N, Takakura H, Ohira Y, and Izawa T. Effect of a 9-week
exercise training regimen on expression of developmental genes related to growth-
dependent fat expansion in juvenile rats. Physiol Rep. 2018; 6(19):e13880.
68. Zhang E, He X, Zhang C, Su J, Lu X, Si X, et al. A novel long noncoding RNA HOXC-AS3
mediates tumorigenesis of gastric cancer by binding to YBX1. Genome Biol. 2018;
19(1):154.
69. Li B, Han H, Song S, Fan G, Xu H, Zhou W, et al. HOXC10 Regulates Osteogenesis of
Mesenchymal Stromal Cells Through Interaction with Its Natural Antisense Transcript
lncHOXC-AS3. Stem Cells. 2019; 37(2):247-256.
70. Wang X, Sun Y, Xu T, Qian K, Huang B, Zhang K, et al. HOXB13 promotes proliferation,
migration, and invasion of glioblastoma through transcriptional upregulation of
lncRNA HOXC-AS3. J Cell Biochem. 2019; 120(9):15527-15537.
71. Wang M, Xie H, Shrestha S, Sredni S, Morgan GA, and Pachman LM. Methylation
alterations of WT1 and homeobox genes in inflamed muscle biopsy samples from
patients with untreated juvenile dermatomyositis suggest self-renewal capacity.
Arthritis and rheumatism. 2012; 64(10):3478-3485.
72. Greiwe JS, Cheng B, Rubin DC, Yarasheski KE, and Semenkovich CF. Resistance
exercise decreases skeletal muscle tumor necrosis factor alpha in frail elderly
humans. Faseb J. 2001; 15(2):475-82.

73.   Leger B, Derave W, De Bock K, Hespel P, and Russell AP. Human sarcopenia reveals an
1177   increase in SOCS-3 and myostatin and a reduced efficiency of Akt phosphorylation.
1178   Rejuvenation Res. 2008; 11(1):163-175B.
1179
1180
1181
1182

1a

Attribute ■ Aged Adult SkM Tissue ■ Young Adult SkM Tissue

-3.22   0.00   3.22

1b 'Biological process'

cellular process
biological regulation
metabolic process
developmental process
cellular component organization or biogenesis
response to stimulus
localization
multicellular organismal process
immune system process
reproductive process
locomotion
multi-organism process
biological adhesion
behavior
signaling
cell population proliferation
growth
rhythmic process
biomineralization
detoxification
cell killing
cell aggregation
pigmentation
nitrogen utilization

3998 2665 1332 0 1332 2665 3998

1c 'Cellular component'

cell part
organelle
organelle part
membrane
membrane part
protein-containing complex
extracellular region part
cell junction
extracellular region
synapse part
synapse
membrane-enclosed lumen
supramolecular complex
host cellular component
cell
nucleoid

4511 3007 1503 0 1503 3007 4511

1d 'Molecular function'

binding
catalytic activity
transcription regulator activity
molecular function regulator
molecular transducer activity
transporter activity
structural molecule activity
translation regulator activity
cargo receptor activity
antioxidant activity
small molecule sensor activity
protein tag
protein folding chaperone
cargo adaptor activity

4228 2818 1409 0 1409 2818 4228

1e Regulation of muscle system process

regulation of muscle contraction

regulation of muscle adaptation

regulation of muscle hypertrophy

regulation of relaxation of muscle

66 44 22 0 22 44 66

Figure 1. DNA methylation across the genome in aged human skeletal muscle tissue compared with young adults. 1a. Hierarchical clustering heatmap of the significantly differentially methylated CpG sites, suggesting that aged skeletal muscle tissue demonstrates predominantly hypermethylation (RED) compared with young adult's skeletal muscle (GREEN). **1b,c,d,e**. Significantly enriched CpG sites in gene ontology (GO) pathways for overarching GO terms: 'biological process' **(b),** 'cellular component' **(c)** and 'molecular function' **(d)** ,and most significantly enriched GO term containing the search word 'muscle'; 'regulation of muscle system process' **(e).** All GO terms demonstrating a predominance of hypermethylation in aged compared with young human skeletal muscle. RED hypermethylated and GREEN hypomethylated CpG's.

PATHWAYS IN CANCER

Figure 2. **The top significantly enriched KEGG pathway 'Pathways in Cancer' in aged compared with young adult human skeletal muscle tissue**. This pathway includes well known pathways in skeletal muscle regulation such as: Focal adhesion, MAPK signaling, PI3K-Akt-mTOR signaling, p53 signalling, Jak-STAT signaling, TGF-beta and Notch signalling. In this pathway 266 out of 277 CpG's were hypermethylated (RED) and only the remaining 11 hypomethylated (GREEN) in aged compared young adult muscle tissue. Note, because some genes have multiple CpGs per gene, Partek Pathway (Partek Genomics Suite) used to create this figure, selects the CpG with the lowest p-value (most significant) and uses the beta-value for that particular CpG to colour the pathway diagram. This is accurate therefore in situations where CpG's on the same gene have the same methylation pattern (e.g. all hyper or hypomethylated). However, where multiple CpG's on the same gene have the different methylation pattern (e.g. some hypomethylated and others hypermethylated), only the most significant methylated CpG is chosen and represented in the image. However, in this instance the majority of CpG's (even when there was more than one per gene) were hypermethylated, so in this case depicts an accurate representation of the data. Full CpG lists including the sites that are hypo and hypermethylated in this pathway are included in Suppl. File 1h.

Figure 3. Aged and young adult primary muscle derived stem cells differentiated over 0, 72 h, 7d and 10 d. With corresponding myogenic gene expression and the gene ontology pathways that were enriched across the time-course of differentiation in aged cells. 3a. Light microscope images of aged versus young muscle derived stem cells morphologically depict fewer myotubes in aged versus young cells, particularly at 7 and 10 days. **3b**. MyoD and myogenin gene expression in aged versus young cells over the differentiation time course. Where, in particular, reductions in myogenin were observed at 72 h in aged cells as well as a delayed increase in the upregulation of myogenin compared with young cells. **1c,d,e.** Aged cells also demonstrated significantly altered DNA methylation as they differentiated, whereas young cells did not demonstrate a significant change in methylation as they differentiated. This altered regulation of DNA methylation over the time course of differentiation was most prominent at 7 days and was enriched in GO terms 'regulation of localisation' (**c**), 'regulation of cell communication' (**d**) and 'regulation of signalling' (**e**). RED hypermethylated and GREEN hypomethylated CpG's.

4a

Hierarchical Clustering

Age ■ Aged Cells ■ Young Cells

-4.03   0.00   4.03

4b 'Cytoskeletal protein binding'
0 h aged vs. 0 h young adult cells

actin binding
tubulin binding
actinin binding
tau protein binding
vinculin binding
kinesin binding
troponin C binding
ankyrin binding
spectrin binding
filamin binding
troponin I binding
telethonin binding
tropomyosin binding

24   16   8   0   8   16   24

4c 'Axon guidance'
0 h aged vs. 0 h young adult cells

4d 'Cytoskeletal protein binding'
72 h aged vs. 72 h young adult cells

actin binding
tubulin binding
kinesin binding
spectrin binding
actinin binding
vinculin binding
troponin C binding
ankyrin binding
filamin binding
troponin I binding
telethonin binding
tau protein binding
tropomyosin binding
cytoskeletal adaptor activity

19   12   6   0   6   12   19

4e 'cAMP signaling'
72 h aged vs. 72 h young adult cells

Figure 4. Comparison of DNA methylation in aged versus young muscle stem cells across differentiation. 4a. Hierarchical clustering heatmap of the significantly differentially methylated CpG sites between aged and young muscle stem cells across the entire time-course of differentiation (all time points of 0, 72 h, 7 d and 10 d). **4b.** 0 h aged vs. 0 h young adult muscle stem cells, top significantly enriched GO term 'cytoskeletal protein binding'. **4c.** 0 h aged vs. 0 h young adult, top KEGG pathway 'axon guidance'. **4d.** 72 h aged vs. 72 h young adult top GO term, 'cytoskeletal protein binding', **4e.** 72 h aged vs. 72 h young adult, a top 3 KEGG pathway 'cAMP signaling'. All demonstrating a predominance of hypermethylation in aged compared with young muscle stem cells in these GO terms and KEGG pathways. RED hypermethylated and GREEN hypomethylated CpG's. Note, because some genes have multiple CpGs per gene, Partek Pathway (Partek Genomics Suite) used to create this figure, selects the CpG with the lowest p-value (most significant) and uses the beta-value for that particular CpG to colour the pathway diagram. This is accurate therefore in situations where CpG's on the same gene have the same methylation pattern (e.g. all hyper or hypomethylated). However, where multiple CpG's on the same gene have the different methylation pattern (e.g. some hypomethylated and others hypermethylated), only the most significant methylated CpG is chosen and represented in the image. Full and accurate CpG lists including the sites that are hypo and hypermethylated in these pathways are included in Suppl. File **3n** (for figure 4c) and Suppl. File **4k** (for figure 4e).

5a 'Developmental process' (incl. anatomical structure development') 7 d aged vs. 7 d young adult cells

5b 'Focal adhesion' 7 d aged vs. 7 d young adult cells

5c 'GDP binding' 10 d aged vs. 10 d young adult cells

5d 'Regulation of the actin cytoskeleton' 10 d aged vs. 10 d young adult cells

Figure 5. Comparison of DNA methylation in aged versus young muscle stem cells at 7 d and 10 d of differentiation. 5a. 7 d aged vs. 7 d young adult skeletal muscle stem cells, top significantly enriched GO term, 'anatomical structure morphogenesis'. **5b**. 7 d aged vs. 7 d young adult, top KEGG pathway, 'focal adhesion'. **5c**. 10 d aged vs 10 d young adult, top GO term, 'GDP binding'. **5d.** 10 d aged vs. 10 d young adult, top KEGG pathway, 'regulation of the actin cytoskeleton'. All demonstrating a predominance of hypermethylation in aged compared with young muscle stem cells in these GO terms and KEGG pathways. RED hypermethylated and GREEN hypomethylated CpG's. Note, because some genes have multiple CpGs per gene, Partek Pathway (Partek Genomics Suite) used to create this figure, selects the CpG with the lowest p-value (most significant) and uses the beta-value for that particular CpG to colour the pathway diagram. This is accurate therefore in situations where CpG's on the same gene have the same methylation pattern (e.g. all hyper or hypomethylated). However, where multiple CpG's on the same gene have the different methylation pattern (e.g. some hypomethylated and others hypermethylated), only the most significant methylated CpG is chosen and represented in the image. Full and accurate CpG lists including the sites that are hypo and hypermethylated in these pathways are included in Suppl. File **5i** (for figure 5b) and Suppl. File **6i** (for figure 5d).

6a

6b

6c

Figure 6. SOM profiling of DNA methylation over the time-course of differentiation in aged muscle stem cells. 6a. Demonstrates that there was a larger number of of hypomethylated and hypermethylated CpG sites in aged muscle stem cells, particularly at 7 days of differentiation, compared with young adult muscle stem cells. Confirming the earlier analysis in figure 3, where aged cells seem to have more varied methylation at 7 days compared with young adult cells. **6b.** Venn Diagram depicting the 24 common CpG sites that were altered at every time point of differentiation between aged and young adult muscle stem cells at each time point of differentiation (0, 72 h, 7d and 10 d). **6c**. As with the above analysis in 6a, SOM profiling identified that 16 out of the 24 CpG's also demonstrated the most varied methylation at 7 days of differentiation in aged cells.

7   Aged versus Young Tissue

(6828)

6819

MGC87042
(cg00394316),
C2orf70 (cg23482427),
cg27209395

KIF15 (2 CpG's
cg00702638, cg24888989),
DYRK2 (cg09516963),
FHL2 (cg22454769),
MRPS33 (cg26792755)

1
ABCA17P
(cg02331561)

4996

2189

(5524)

(2719)

Aged Stem Cells 7 d versus
Young Adult Cells 7 d

Aged Stem Cells versus
Young Adult Cells (all time)

Figure 7. Venn diagram of common overlapping CpG sites significantly differentially methylated between aged skeletal muscle tissue and muscle stem cells. Venn diagram depicts commonly varied CpG sites in aged skeletal muscle tissue that were also varied in aged muscle stem cell analysis. Overlap of the tissue and aged cells included common 6 CpG sites on genes KIF15 (2 CpG's Cg00702638 & cg24888989), DYRK2 (cg09516963), FHL2 (cg22454769), MRPS33 (cg26792755) and ABCA17P (cg02331561). Given that aged cells demonstrated the most varied methylation at 7 days versus young cells. When overlapping the 7 d most significantly differentially methylated CpG lists, 4 CpG's (out of the 6 CpG sites identified above) were also identified, including: MGC87042 (cg00394316), C2orf70 (cg23482427), ABCA17P (cg02331561) and cg27209395 (not on an annotated gene). Once more, all of these CpG's (with the exception of C2orf70, cg23482427) were hypermethylated in the tissue analysis as well as the stem cells. With ABCA17P (cg02331561) highlighted across all CpG lists.

HOX Family Genes

8a

Aged Tissue vs. Young Adult Tissue
(6,828 CpG list)

Aged Cells vs. Young
Adult Cells (2,719 CpG list)

Aged Cells 7d vs. Young adults
Cells 7d (5,524 CpG list)

8b

8c

HOXC10
(cg19898108**)**

HOXD8 (cg18448949, cg00035316), **HOXB1** (cg04904318, cg02497558, cg22660933, cg10558129), **HOXC9** (cg02227188), **HOXB3** (cg09952002)

HOXC10 (cg20402783, cg20403938, cg16898193, cg16791250), **HOXB3** (cg23014425, cg04800503), **HOXC-AS2** (cg09605287)

HOXA3
(cg20688917**)**

Difference in Standardised M-Value

| 0 h | 72 h | 7 d | 10 d |

| 0 h | 72 h | 7 d | 10 d |

Figure 8. HOX family of genes and their DNA methylation in aged tissue and muscle stem cells. 8a. Venn diagram identifying 9 commonly differentially methylated HOX genes: including HOXD10, HOXD9, HOXD8, HOXA3, HOXC9, HOXB1, HOXB3, HOXC-AS2 and HOXC10 (note this Venn diagram analysis is by 'gene symbol' not 'probe cg' (CpG site), as some HOX genes also had more than 1 CpG per gene symbol, full Cg lists are located in Suppl. Figure 9 a,b,c,d). **8b.** All HOX family genes by CpG site (cg probe) differentially methylated in aged compared with young skeletal muscle tissue, predominantly all demonstrating hypermethylation. **8c.** SOM profiling depicting the temporal regulation of DNA methylation in aged muscle stem cells as they differentiate in CpG's located amongst the HOX family of genes (depicting the 9 HOX genes identified to be changed, by gene symbol, in both the tissue and cells in Figure 8a). The majority of these HOX CpG sites were differentially methylated at 7 days of differentiation in the aged cells.

Figure 9. Gene expression of the HOX family of genes in aged compared to young adult skeletal muscle tissue (9a) and muscle stem cells at 7 days of differentiation (9b).