1
2
3
4 **Diet-induced loss of adipose Hexokinase 2 triggers hyperglycemia**
5

6 Mitsugu Shimobayashi[1*], Sunil Shetty[1†], Irina C. Frei[1†], Bettina K. Wölnerhanssen[2,3], Diana
7 Weissenberger[1], Nikolaus Dietz[1], Amandine Thomas[1], Danilo Ritz[1], Anne Christin Meyer-
8 Gerspach[2,3], Timm Maier[1], Nissim Hay[4], Ralph Peterli[5], Nicolas Rohner[6,7], Michael N. Hall[1*]
9

[1] Biozentrum, University of Basel, Basel 4056, Switzerland.

[2] St. Clara Research Ltd, St. Claraspital, Basel 4016, Switzerland.

[3] University of Basel, Basel 4056, Switzerland.

[4] Department of Biochemistry and Molecular Genetics, College of Medicine, University of

Illinois at Chicago, Chicago, IL 60607, USA.

[5] Clarunis, Department of Visceral Surgery, University Centre for Gastrointestinal and Liver

Diseases, Basel 4016, Switzerland.

[6] Stowers Institute for Medical Research, Kansas City, Missouri 64110, USA.

[7] Department of Molecular and Integrative Physiology, KU Medical Center, Kansas City,

Kansas 66160, USA

[†] These authors contributed equally to this work.

[*] Corresponding authors:

Klingelbergstrasse 70, Basel 4056, Switzerland

TEL: +41 61 207 2171 (MS)

+41 61 207 2150 (MNH)

e-mail: mitsugu.shimobayashi@unibas.ch (MS)

m.hall@unibas.ch (MNH)

30
31

Summary

Chronically high blood glucose (hyperglycemia) leads to type 2 diabetes, fatty liver disease, and cardiovascular disease[1]. Obesity is a major risk factor for hyperglycemia, but the underlying mechanism is unknown. Here we show that a high fat diet (HFD) in mice causes early loss of expression of the glycolytic enzyme Hexokinase 2 (HK2) specifically in adipose tissue. Adipose-specific knockout of *Hk2* caused enhanced gluconeogenesis and lipogenesis in liver, a condition known as selective insulin resistance[2], leading to glucose intolerance. Furthermore, we observed reduced hexokinase activity in adipose tissue of obese patients, and identified a loss-of-function mutation in the *hk2* gene of naturally hyperglycemic Mexican cavefish. Mechanistically, HFD in mice led to loss of HK2 by inhibiting elongation of *Hk2* mRNA translation. Our findings identify adipose HK2 as a critical mediator of systemic glucose homeostasis, and suggest that obesity-induced loss of adipose HK2 is an evolutionarily conserved, non-cell-autonomous mechanism for hyperglycemia.

Main text

Vertebrates mediate glucose homeostasis by regulating glucose production and uptake in specific organs[3]. High blood glucose stimulates pancreatic beta cells to secrete the hormone insulin which in turn promotes glucose uptake in skeletal muscle and adipose tissue and inhibits glucose production in liver. Obesity is highly associated with insulin insensitivity in these peripheral tissues, leading to hyperglycemia. Although the contribution of adipose tissue to glucose clearance is minor[4,5], glucose metabolism in adipocytes is key for systemic glucose homeostasis[6,7]. Insulin activates the protein kinases mammalian target of rapamycin complex 2 (mTORC2) and AKT in adipose tissue to promote glucose uptake[8,9]. Furthermore, adipose-specific knockout of these insulin signaling components results in local and systemic insulin insensitivity[10-13], highlighting the physiological importance of the adipose insulin-mTORC2-AKT signaling pathway in glucose homeostasis. However, we and others recently reported that diet-induced obesity in mice causes hyperglycemia despite normal insulin signaling in adipose tissue[14,15]. These observations raise the question, how does obesity cause insulin insensitivity and hyperglycemia without affecting the insulin signaling pathway?

HK2 is down-regulated in obese mice and humans

To determine the molecular basis of diet-induced insulin insensitivity in adipocytes and hyperglycemia in mice, we performed an unbiased proteomic analysis on visceral white adipose tissue (vWAT) isolated from C57BL/6JRj wild-type mice fed an HFD or normal diet (ND) for 4 weeks (**Extended Data Fig. 1a-1c**). We detected and quantified 6294 proteins of which 52 and 67 were up- and down-regulated, respectively, in vWAT of HFD-fed mice (**Extended Data Table 1**). The glycolytic enzyme Hexokinase 2 (HK2), expressed in adipose tissue and muscle, was among the proteins significantly down-regulated in vWAT (**Fig. 1a**). HK2 phosphorylates glucose to generate glucose-6-phosphate (G6P), the rate-limiting step in glycolysis, in an insulin-stimulated manner. HK2 caught our attention due to efflux of non-phosphorylated glucose and thus a link of HK2 loss to hyperglycemia. HK2 is the most abundant (~80%) of the three hexokinase isoforms expressed in vWAT but the only one down-regulated upon HFD (**Fig. 1a**). Consistent with a previous study[13], expression of the glucose transporter GLUT4 was not affected in vWAT of 4-week HFD mice. Loss of HK2 was confirmed in vWAT and demonstrated in subcutaneous WAT (sWAT) and brown adipose tissue (BAT) of 4-week HFD mice by immunoblotting (**Fig. 1b**). HK2 expression was unchanged in skeletal muscle (**Fig. 1b**), suggesting that an HFD decreases HK2 expression specifically in adipose tissue. Consistent with reduced HK2 expression, hexokinase activity was decreased in vWAT and sWAT (**Extended Data Fig. 2**). A longitudinal study of HFD-fed mice revealed that HK2 down-regulation in sWAT and vWAT

occurred already at one week of HFD and correlated with hyperglycemia (**Extended Data**
Fig. 3a-3e). GLUT4 was down-regulated only in sWAT and only after $\geq$2 weeks of HFD.
Confirming earlier observations[14,15], insulin signaling was normal at 4 weeks of HFD.
Shifting the diet from HFD to ND for 2 weeks restored HK2 expression and normal blood
glucose levels (**Fig. 1c** and **Extended Data Fig. 3d**), suggesting that loss of HK2 upon HFD
is a transient physiological response. Furthermore, the finding that HK2 expression
inversely correlated with hyperglycemia suggests that loss of adipose HK2 may be causal
for loss of glucose homeostasis. Omental WAT (human vWAT) biopsies from obese non-
diabetic and obese diabetic patients displayed reduced HK activity which inversely
correlated with body mass index (BMI) (**Fig. 1d and 1e** and **Extended Data Table 2**),
consistent with the previous observation that HK2 expression is decreased in adipose tissue
of diabetic patients[16]. These findings suggest that HK2 down-regulation is a key event,
possibly causal, in obesity-induced insulin insensitivity and hyperglycemia in mouse and
human.
98
A loss-of-function mutation in cavefish *hk2*
Mexican cavefish (*Astyanax mexicanus*), also known as blind fish, are hyperglycemic
compared to surface fish from which they are descended[17](**Fig. 1f** and **1g**). Insulin
resistance and hyperglycemia, although a pathological condition in mouse and human, is a
selected trait that allows cave-dwelling fish to survive in nutrient-limited conditions. Among
three independently evolved cavefish isolates, Pachón and Tinaja cavefish (names refer to
the caves from which the fish were isolated) contain a loss-of-function mutation in the insulin
receptor, causing insulin resistance and hyperglycemia[17] (**Fig. 1g**). The third isolate, Molino
cavefish, is the most hyperglycemic but contains a wild-type insulin receptor and displays
normal insulin signaling[17]. Since the phenotype of the Molino fish is similar to that of HFD-
fed mice, we hypothesized that this cavefish may be hyperglycemic due to a loss-of-function
mutation in the *hk2* gene. DNA sequencing of the *hk2* gene of surface fish and the three
cavefish variants revealed a mutation in the *hk2* gene uniquely in Molino. The homozygous,
missense mutation in the coding region of the Molino *hk2* gene changed highly conserved
arginine 42 (R42) to histidine (R42H) (**Fig. 1h** and **Extended Data Fig. 4b**). Based on the
published structure of HK2 [18], R42 forms a salt bridge with aspartic acid 272 (D272) to
stabilize the conformation of HK2 (**Extended Data Fig. 4b**), predicting that R42H
destabilizes HK2 and is thus a loss-of-function mutation. To test this prediction, we over-
expressed wild-type HK2 (HK-WT) and HK2 harboring the Molino mutation (HK2-R42H) in
HEK293T cells which have low intrinsic hexokinase activity. HK2-R42H displayed ~50%
lower hexokinase activity compared to HK2-WT, despite similar expression levels,
suggesting that R42H is indeed a loss-of-function mutation (**Fig. 1i** and **Extended Data Fig.**

4c). To test further whether the HK2-R42H allele is loss-of-function, we generated a knock-

in mouse in which the Molino missense mutation (R42H) is introduced into the mouse *Hk2*

gene by CRISPR-Cas9 gene editing (**Extended Data Fig 5a** and **5b**). Upon crossing

heterozygous $Hk2^{R42H/+}$ mice, we obtained offspring at the non-mendelian ratio of 18:91:49

(R42H homozygous:R42H heterozygous:wild-type) (**Extended Data Table 3**). We note that

our $Hk2^{R42H}$ allele is a full body knock-in and that a full body homozygous *Hk2* knockout is

embryonic lethal[19]. Together, the above findings in cells and mice suggest that R42H is a

partial loss-of-function mutation and may account for the hyperglycemia in Molino cavefish.

In other words, it appears that R42H in Molino is a genetically fixed version of what we

observe in mice as a physiological down-regulation of HK2 in response to HFD. The

findings in Molino provide orthogonal evidence that the down-regulation of HK2 in mice is

physiologically relevant. Further experiments with R42H mice were inconclusive due to

small littermate cohorts and high variation, although in some cases hyperglycemia was

observed.

Adipose-specific *Hk2* knockout causes hyperglycemia

The above findings altogether suggest that adipose-specific loss of HK2 may be a cause of

insulin insensitivity and hyperglycemia. To test this hypothesis, we first generated a stable

HK2 knockdown pre-adipocyte 3T3-L1 cell line (**Extended Data Fig. S6a** and **S6b**). HK2

knockdown pre-adipocytes differentiated normally to produce mature adipocytes (**Extended**

Data Fig. 6c and **6d**). The glycolytic rate was lower in HK2 knockdown adipocytes

compared to controls, as measured by extracellular acidification rate (ECAR) (**Fig. 2a**) and

lactate production (**Fig. 2b**). Furthermore, although basal glucose uptake did not differ,

insulin-stimulated glucose uptake was 50% lower in HK2 knockdown adipocytes (**Fig. 2c**).

Thus, loss of HK2 causes insulin insensitivity in adipocytes *in vitro*.

To examine further the role of adipose HK2 in glucose homeostasis, and in particular

the causality of HK2 loss in hyperglycemia, we generated an adipose-specific *Hk2* knockout

(AdHk2KO) mouse. In AdHk2KO mice, HK2 expression was decreased in vWAT, sWAT,

and BAT but unchanged in skeletal muscle (**Fig. 2d, Extended Data Fig. 7a** and **7b**), similar

to HFD-fed mice (**Fig. 1b**). AdHk2KO mice displayed slightly less fat mass and slightly more

lean mass than controls, but no difference in overall body weight (**Extended Data Fig. 7c**

and **7d**). No significant difference was observed in organ weight, except for a small

decrease in vWAT and a small increase in liver weight (**Extended Data Fig. 7e** and **7f).**

Importantly, AdHk2KO mice were insulin insensitive and severely glucose intolerant (**Fig. 2e**

and **2f**). These findings suggest that loss of HK2 specifically in adipose tissue causes

insulin insensitivity and hyperglycemia.

Regulation of liver metabolism by adipose HK2

Adipose tissue accounts for $\leq$5% of glucose disposal[4,5]. Thus, the severe glucose intolerance observed in AdHk2KO mice (**Fig. 2f**) cannot be explained solely by impaired glucose uptake into adipose tissue. One possible explanation for the severity of the glucose intolerance could be impaired insulin secretion. However, plasma insulin levels were similar in AdHk2KO and control mice (**Extended Data Fig. 7g**). Moreover, insulin signaling (AKT-pS473) was not affected in adipose tissue, skeletal muscle, and liver of AdHk2KO mice (**Fig. 2d** and **Extended Data Fig. 7b**), suggesting that hyperglycemia in AdHk2KO mice is not due to a defect in insulin signaling in peripheral tissues.

Another potential explanation for the severity of glucose intolerance in AdHk2KO mice is derepressed hepatic glucose production. Adipose tissue is known to impinge negatively on glucose production in liver [6,11,13,20], the main glucose producing organ. To examine whether AdHk2KO mice display enhanced hepatic glucose production, we first performed a pyruvate tolerance test (PTT). AdHk2KO mice displayed significantly higher glucose production compared to controls (**Fig. 3a**). We next examined expression of gluconeogenic genes (*G6pc* and *Pepck*) in liver. Consistent with the observed increase in glucose production, gluconeogenic genes were upregulated in AdHk2KO liver (**Fig. 3b**). These findings suggest that enhanced hepatic gluconeogenesis accounts for the severity of glucose intolerance in AdHk2KO mice. Thus, loss of adipose HK2 non-cell-autonomously promotes hepatic gluconeogenesis and thereby hyperglycemia.

We observed enhanced hepatic gluconeogenesis in AdHk2KO mice despite normal systemic insulin signaling, including in liver. To investigate this apparent contradiction, we examined lipogenic gene expression, another readout for hepatic insulin action. Sterol regulatory element-binding protein 1c (SREBP1c) is a transcription factor that activates lipogenic genes and thereby promotes *de novo* lipogenesis in the liver[21]. Expression of SREBP1c and its target fatty acid synthesis genes (*Acly*, *Scd1*, *Fasn*, and *Acc*) were increased in liver of AdHk2KO mice (**Fig. 3c**), suggesting that loss of adipose HK2 also promotes hepatic lipogenesis. The increased hepatic gluconeogenesis and lipogenesis in AdHk2KO mice mimics a condition in diabetic patients known as selective insulin resistance[2] (see below). Thus, loss of adipose HK2 may cause selective insulin resistance and thereby contribute to the pathogenesis of type 2 diabetes.

We note that hepatic triglyceride (TG) levels were unchanged in AdHk2KO mice despite increased expression of lipogenic genes (**Extended Data Fig. 7h**). Most likely, *de novo*-synthesized TG in AdHk2KO liver is delivered to adipose tissue for storage. This is consistent with our observations that adipose tissue mass is largely unaffected in AdHk2KO mice.

HK2 activates ChREBP in adipose tissue

How does adipose HK2 control hepatic gluconeogenesis? As shown previously, adipose-specific knockout of carbohydrate-responsive element binding protein (ChREBP) decreases adipose lipogenesis and increases hepatic gluconeogenesis[20,22]. Furthermore, HFD inhibits adipose ChREBP and lipogenesis via an unknown mechanism proposed to involve glucose or a glucose metabolite[23]. G6P, a glucose metabolite and product of HK2, promotes ChREBP activity in a rat insulinoma cell line[24]. Thus, we hypothesized that HFD-induced loss of HK2 and a concomitant decrease in G6P leads to loss of both ChREBP activity and lipogenesis in adipose tissue. To test this hypothesis, we examined *Chrebp* expression in adipose tissue of AdHk2KO and control mice. ChREBP has two isoforms. Constitutively expressed ChREBPα promotes transcription of ChREBPβ which is then dominant for activation of fatty acid synthesis genes[23]. Expression of *Chrebpβ,* but not *Chrebpα*, was significantly decreased in vWAT, sWAT, and BAT of AdHk2KO mice (**Fig. 3d-3e** and **Extended Data Fig. 7i**). Consistent with reduced *Chrebpβ* expression, fatty acid synthesis genes were down-regulated in vWAT, sWAT, and BAT of AdHk2KO mice (**Fig. 3d-3e** and **Extended Data Fig. 7i**). Thus, HK2 promotes ChREBP activity and lipogenesis in adipose tissue, possibly via G6P. Combined with previous observations[20], our findings suggest that HK2 controls hepatic gluconeogenesis via adipose ChREBP (see below).

Mechanism of HFD-induced HK2 down-regulation

Our findings suggest that HFD causes loss of adipose HK2 and thereby triggers hyperglycemia. How does HFD down-regulate adipose HK2? To answer this question, we first examined *Hk2* mRNA expression in HFD- and ND-fed mice. HFD-fed mice exhibited decreased *Hk2* mRNA in sWAT but not in vWAT and BAT (**Fig. 4a**). This suggests that HK2 synthesis is down-regulated at the transcriptional level in sWAT, and at a post-transcriptional level in vWAT and BAT. To investigate the post-transcriptional mechanism, we performed polysome profiling on vWAT from HFD- and ND-fed mice. In the presence of cycloheximide to maintain ribosome-mRNA complexes, vWAT lysates from HFD-fed mice contained more polysomes and fewer monosomes (80S ribosome) relative to ND-fed mice (**Fig. 4b**). An increase in polysomes can be due to increased translation initiation or decreased elongation. To distinguish these two possibilities, we performed polysome profiling in the absence of cycloheximide, a condition which allows actively translating ribosomes to run off mRNA. Although there were fewer overall polysomes in the absence cycloheximide, we consistently observed more polysomes in vWAT from HFD-fed mice (**Fig. 4c**), suggesting that HFD inhibits or slows translation elongation of a subset of mRNAs. Shifting the diet from HFD to ND for 2 weeks restored elongation (**Fig. 4d**), as described above for glycemia (**Extended

Data Fig. 3d). To determine whether HFD affects elongation of *Hk2* mRNA translation, we measured *Hk2* mRNA in monosome, light polysome, and heavy polysome fractions. *Hk2* mRNA from HFD vWAT was enriched in heavy polysomes, whereas it was enriched in light polysomes in ND vWAT (**Fig. 4e** and **Extended Data Fig. 8a**). Thus, HFD down-regulates HK2 in vWAT by inhibiting elongation of *Hk2* mRNA translation.

It was recently reported that the forkhead transcription factors FOXK1 and FOXK2 promote *Hk2* transcription in adipose tissue[25]. We investigated whether the transcriptional down-regulation of *Hk2* mRNA in sWAT could be due to loss of FOXK1 and FOXK2. *Foxk1* and *Foxk2* expression was significantly decreased in sWAT, but not in vWAT or BAT, of HFD-fed mice (**Fig. 4f, Extended Data Fig. 8b** and **8c**). Thus, HFD appears to down-regulate HK2 in sWAT by inhibiting FOXK1/2 and, thereby, expression of the *Hk2* gene.

Discussion

How obesity causes hyperglycemia is a long-standing question. In this study, we show that a high fat diet causes hyperglycemia by inducing loss of HK2 specifically in adipose tissue. Loss of adipose HK2 leads to hyperglycemia in two ways (**Fig. 4g**). First, HK2 loss results in reduced glucose uptake by adipose tissue due to the inability of adipocytes to trap non-phosphorylated glucose. Second, loss of HK2 in adipocytes de-represses glucose production in liver.

How does loss of HK2 in adipose tissue de-repress hepatic gluconeogenesis? The Kahn group showed that adipose ChREBP normally promotes synthesis of fatty acids including palmitic acid hydroxy stearic acid (PAHSA) which is secreted and suppresses hepatic gluconeogenesis[20]. We demonstrate that loss of HK2 impairs ChREBP activation and prevents fatty acid synthesis. Thus, loss of adipose HK2 may de-represses hepatic gluconeogenesis via down-regulation of circulating PAHSA.

In liver, insulin normally down-regulates gluconeogenesis to decrease blood glucose and up-regulates lipogenesis to increase energy storage. Paradoxically, in type 2 diabetes, insulin fails to inhibit gluconeogenesis but stimulates lipogenesis, hence the liver is selectively insulin resistant[2]. Furthermore, previous findings suggest that selective resistance occurs despite intact hepatic insulin signaling[26]. Consistent with these previous findings, we observed that loss of adipose HK2 causes selective insulin resistance without affecting systemic insulin signaling. Thus, we propose that loss of adipose HK2 is a mechanism for selective insulin resistance and ultimately diabetes.

We found an HK2-R42H mutation in naturally hyperglycemic Mexican cavefish. To date, no human monogenic disease has been reported associated with mutations in the *HK2* gene. However, 12 R42W and 4 R42Q alleles are reported in the genomAD database[27].

267 Further studies are required to determine whether these alleles are loss-of-function and
268 associated with diabetes.
269
Methods
Mouse
Wild-type C57BL/6JRj mice were purchased from JANVIER LABS. Mice carrying *Hk2* with
exons 4-10 flanked by loxP sites (*Hk2$^{fl/fl}$*) were as described[28]. *Adipoq-CreER^{T2}* mice were
provided by Prof. Stefan Offermanns (MPI-HLR, Germany)[29]. *Hk2$^{fl/fl}$* mice were crossed with
Adipoq-CreER^{T2} mice, and resulting Cre positive *Hk2$^{fl/+}$* mice were crossed with *Hk2$^{fl/fl}$* mice
to generate adipocyte-specific *Hk2* knockout (*Adipoq-CreER^{T2}* positive *Hk2$^{fl/fl}$*) mice
(AdHk2KO). *Hk2* knockout was induced by i.p. injection of 1 mg/mouse tamoxifen (Sigma-
Aldrich) in corn oil for 7 days. Littermate *Cre* negative animals were used as a control.
Control mice were also treated with tamoxifen.
The (R42H) modification of the *Hk2* locus using Cas9/CRISPR was carried out by
electroporation of zygotes. C57BL/6J female mice underwent ovulation induction by i.p.
injection of 5 IU equine chorionic gonadotrophin (PMSG; Folligon–InterVet), followed by i.p.
injection of 5 IU human chorionic gonadotropin (Pregnyl–Essex Chemie) 48 hours later. For
recovery of zygotes, C57BL/6J males and females were mated immediately after
administration of human chorionic gonadotropin. Zygotes were collected from oviducts 24
hours after the human chorionic gonadotropin injection, and were then freed from any
remaining cumulus cells by a 1–2 min treatment of 0.1% hyaluronidase (Sigma-Aldrich)
dissolved in M2 medium (Sigma). Prior to electroporation, the zona pellucida was partially
removed by brief treatment with acid Tyrode's solution and the zygotes were washed and
briefly cultured in M16 (Sigma) medium at 37 °C and 5% CO_2. Electroporation with a
mixture of 16 uM cr:trcrRNA hybrid targeting *Hk2* (5'cccttctggagatttctagg<u>cgg</u> 3' - PAM
underlined), 16 µM Cas9 protein and 200 nt ssDNA oligonucleotide carrying the desired
mutations
(5'gcaaggctgggttatatttaataacatagaacctattcttgtcccctgcccccttcaggttgaccagtatctctaccacatgcgtct
ctcagatgagacccttctggagatttctagacatttccggaaggagatggagaaggggctaggagctaccacacaccctacag
cagctgtgaagatgctgcccacctttgtgt 3') (all reagents from IDT) was carried out using 1 mm gap
electroporation cuvette and the ECM830 electroporator (BTX Harvard Apparatus). Two 3
millisecond pulses of 30 V with 100 millisecond intervals were applied as previously
described[30]. Surviving zygotes were washed with M16 medium and transferred immediately
into the oviducts of 8–16 week old pseudopregnant Crl:CD1(ICR) females that had been
mated with sterile genetically vasectomized males[31] the day before zygote transfer (0.5 dpc).
Pregnant females were allowed to deliver and raise their pups until weaning age. In total,
475 zygotes were electroporated and 470 surviving zygotes were transferred into 20 foster

mothers. Possible lethality of NHEJ-mediated disruption of the *Hk2* locus resulted in

substantial reduction of offspring. Only 4 of 20 foster mothers produced live litters with a

total of seven F_0 pups. Two F_0 pups carrying the desired modification (R42H) of the *Hk2*

gene were crossed with C57BL/6JRj mice. Mice of the F_4 generation was used to determine

the mendelian ratio.

Mice were housed at 22 ºC in a conventional facility with a 12 hour light/dark cycle

with unlimited access to water, and normal diet (ND) or high fat diet (HFD: 60% kcal % fat

NAFAG 2127, KLIBA). Only male mice between 6 and 17 weeks of age were used for

experiments.

Plasmids

The R42H mutation was introduced into the pLenti-CMV-ratHK2 by PCR using the oligos

5'atttctaggcACttccggaaggagatggagaaag3' and 5'cttccggaaGTgcctagaaatctccagaagggtc3'.

The desired sequence change was confirmed.

Cell culture

HEK293T cells were obtained from ACTT and cultured in M1 (DMEM supplemented with 4

mM glutamine, 1mM sodium pyruvate, 1x penicillin and streptomycin, and 10% FBS). 3T3-

L1 cells were obtained from ATCC and cultured and differentiated as previously described[32].

In brief, 3T3-L1 preadipocyte cells were maintained in M1 medium at 37 ºC incubator with

5% CO_2. For differentiation, cells were maintained in M1 medium for 2 days after reaching

confluence. The cells were the transferred to M2 medium (M1 medium supplemented with

1.5 µg/mL insulin, 0.5 mM IBMX, 1 µM dexamethasone, and 2 µM rosiglitazone), defined as

day 0 post-differentiation. After 2 days, the cells were transferred to M3 medium (M1 with

1.5 µg/mL insulin). At day 4 post-differentiation, cells were transferred back to M2 medium.

From day 6 post-differentiation, cells were maintained in M3 with medium change every two

days.

For Hk2 knockdown, MISSION shRNA (TRCN0000280118) or control pLKO plasmid

were purchased from Merck and co-transfected with psPAX2 (a gift from Didier Trono:

Addgene plasmid # 12260) and pCMV-VSV-G[33] (a gift from Robert Weinberg: Addgene

plasmid # 8454) into HEK293T cells. Supernatants containing lentivirus were collected one

day after transfection, and used to infect undifferentiated 3T3-L1 cells. Transduced cells

were selected by puromycin. For all experiments, 8-14 days post-differentiated cells were

used.

Human biopsies

Omental white adipose tissue (oWAT) biopsies were obtained from lean subjects with

normal fasting glucose level and body mass index (BMI) < 27 kg/m^2, from obese non-

diabetic subjects with BMI > 35 kg/m^2 HbA1c < 6.0%, and from obese diabetic sujects with

BMI > 35 kg/m^2 HbA1c > 6.1% (**Table S2**). All subjects gave informed consent before the

surgical procedure. Patients were fasted overnight and underwent general anesthesia. All

oWAT specimens were obtained between 8:30 and 12:00am, snap-frozen in liquid nitrogen,

and stored at -80 ºC for subsequent use.

Proteomics

Tissues were pulverized and homogenized in lysis buffer containing 100 mM Tris-HCl pH7.5,

2 mM EDTA, 2 mM EGTA, 150 mM NaCl, 1% Triton X-100, cOmplete inhibitor cocktail

(Roche) and PhosSTOP (Roche). Proteins were precipitated by trichloroacetic acid (Sigma)

and the resulting protein pellets were washed with cold acetone. 25 µg of peptides were

labeled with isobaric tandem mass tags (TMT 10-plex, Thermo Fisher Scientific) as

described previously [34]. Shortly, peptides were resuspended in 20 µl labeling buffer (2 M

urea, 0.2 M HEPES, pH 8.3) and 5 µL of each TMT reagent were added to the individual

peptide samples followed by a 1 h incubation at 25 °C. To control for ratio distortion during

quantification, a peptide calibration mixture consisting of six digested standard proteins

mixed in different amounts was added to each sample before TMT labeling. To quench the

labelling reaction, 1.5 µL aqueous 1.5 M hydroxylamine solution was added and samples

were incubated for another 10 min at 25 °C followed by pooling of all samples. The pH of

the sample pool was increased to 11.9 by adding 1 M phosphate buffer (pH 12) and

incubated for 20 min at 25 °C to remove TMT labels linked to peptide hydroxyl groups.

Subsequently, the reaction was stopped by adding 2 M hydrochloric acid and until a pH < 2

was reached. Finally, peptide samples were further acidified using 5% TFA, desalted using

Sep-Pak Vac 1cc (50 mg) C18 cartridges (Waters) according to the manufacturer's

instructions and dried under vacuum.

TMT-labeled peptides were fractionated by high-pH reversed phase separation using

a XBridge Peptide BEH C18 column (3,5 µm, 130 Å, 1 mm x 150 mm, Waters) on an Agilent

1260 Infinity HPLC system. Peptides were loaded on column in buffer A (20 mM ammonium

formate in water, pH 10) and eluted using a two-step linear gradient from 2% to 10% in 5

minutes and then to 50% buffer B (20 mM ammonium formate in 90% acetonitrile, pH 10)

over 55 min at a flow rate of 42 µl/min. Elution of peptides was monitored with a UV detector

(215 nm, 254 nm) and a total of 36 fractions were collected, pooled into 12 fractions using a

post-concatenation strategy as previously described[35] and dried under vacuum.

Dried peptides were resuspended in 20 µl of 0.1% aqueous formic acid and

subjected to LC–MS/MS analysis using a Q Exactive HF Mass Spectrometer fitted with an

EASY-nLC 1000 (both Thermo Fisher Scientific) and a custom-made column heater set to 60 °C. Peptides were resolved using a RP-HPLC column (75 μm × 30 cm) packed in-house with C18 resin (ReproSil-Pur C18–AQ, 1.9 μm resin; Dr. Maisch GmbH) at a flow rate of 0.2 μL/min. The following gradient was used for peptide separation: from 5% B to 15% B over 10 min to 30% B over 60 min to 45 % B over 20 min to 95% B over 2 min followed by 18 min at 95% B. Buffer A was 0.1% formic acid in water and buffer B was 80% acetonitrile, 0.1% formic acid in water.

The mass spectrometer was operated in DDA mode with a total cycle time of approximately 1 sec. Each MS1 scan was followed by high-collision-dissociation (HCD) of the 10 most abundant precursor ions with dynamic exclusion set to 30 sec. For MS1, 3e6 ions were accumulated in the Orbitrap over a maximum time of 100 ms and scanned at a resolution of 120,000 FWHM (at 200 m/z). MS2 scans were acquired at a target setting of 1e5 ions, accumulation time of 100 ms and a resolution of 30,000 FWHM (at 200 m/z). Singly charged ions and ions with unassigned charge state were excluded from triggering MS2 events. The normalized collision energy was set to 35%, the mass isolation window was set to 1.1 m/z and one microscan was acquired for each spectrum.

The acquired raw-files were converted to the mascot generic file (mgf) format using the msconvert tool (part of ProteoWizard, version 3.0.4624 (2013-6-3)). Using the MASCOT algorithm (Matrix Science, Version 2.4.1), the mgf files were searched against a decoy database containing normal and reverse sequences of the predicted SwissProt entries of Mus musculus (www.ebi.ac.uk, release date 2014/11/24), the six calibration mix proteins[34] and commonly observed contaminants (in total 50214 sequences for Mus musculus) generated using the SequenceReverser tool from the MaxQuant software (Version 1.0.13.13). The precursor ion tolerance was set to 10 ppm and fragment ion tolerance was set to 0.02 Da. The search criteria were set as follows: full tryptic specificity was required (cleavage after lysine or arginine residues unless followed by proline), 3 missed cleavages were allowed, carbamidomethylation (C) and TMT6plex (K and peptide N-terminus) were set as fixed modification and oxidation (M) as a variable modification. Next, the database search results were imported into the Scaffold Q+ software (version 4.3.2, Proteome Software Inc., Portland, OR) and the protein false identification rate was set to 1% based on the number of decoy hits. Proteins that contained similar peptides and could not be differentiated based on MS/MS analysis alone were grouped to satisfy the principles of parsimony. Proteins sharing significant peptide evidence were grouped into clusters. Acquired reporter ion intensities in the experiments were employed for automated quantification and statically analysis using a modified version of our in-house developed SafeQuant R script[34]. This analysis included adjustment of reporter ion intensities, global data normalization by equalizing the total reporter ion intensity across all channels, summation of reporter ion intensities per protein

and channel, calculation of protein abundance ratios and testing for differential abundance

using empirical Bayes moderated t-statistics. Finally, the calculated p-values were corrected

for multiple testing using the Benjamini−Hochberg method (q-value). Deregulated proteins

were selected by log2(fold change) > 0.6 or log2(fold change) < -0.6, q-value < 0.01.

Immunoblots

Adipose tissue, liver, and skeletal muscle were homogenized in lysis buffer containing 100

mM Tris-HCl pH7.5, 2 mM EDTA, 2 mM EGTA, 150 mM NaCl, 1% Triton X-100, cOmplete

inhibitor cocktail (Roche) and PhosSTOP (Roche). Protein concentration was determined by

the Bradford assay, and equal amounts of protein were separated by SDS-PAGE, and

transferred onto nitrocellulose membranes (GE Healthcare). Antibodies used in this study

were as follows: AKT (Cat#4685), AKT-pS473 (Cat#4060), HK2 (Cat#2867), HK1

(Cat#2024), HA tag (Cat#3724) from Cell Signaling, GLUT4 (Cat# NBP2-22214, NOVUS),

and ACTIN (Cat#MAB1501, Millipore).

RNA isolation and quantitative RT-PCR

Total RNA was isolated with TRIzol reagent (Sigma-Aldrich) and RNeasy kit (Qiagen). RNA

was reverse-transcribed to cDNA using iScript cDNA synthesis kit (BIO-RAD).

Semiquantitative real-time PCR analysis was performed using fast SYBR green (Applied

Biosystems). Relative expression levels were determined by normalizing each CT values to

Tbp using the ∆∆CT method. The sequence for the primers used in this study was as

follows. Fasn-fw: 5'GCTGCGGAAACTTCAGGAAAT3', Fasn-rv:

5'AGAGACGTGTCACTCCTGGACTT3', Acc-fw: 5'AAGGCTATGTGAAGGATG3', Acc-rv:

5'CTGTCTGAAGAGGTTAGG3', Cd36-fw: 5' TGGCCTTACTTGGGATTGG3', Cd36-rv: 5'

CCAGTGTATATGTAGGCTCATCCA 3', G6pc-fw: 5'CCATGCAAAGGACTAGGAACAA3',

G6pc-rv: 5' TACCAGGGCCGATGTCAAC 3', Pepck-fw: 5' CCACAGCTGGTGCAGAACA3',

Pepck-rv: 5' GAAGGGTCGATGGCAAA 3', Chrebp-fw:

5'CACTCAGGGAATACACGCCTAC3', Chrebp-rv: 5'ATCTTGGTCTTAGGGTCTTCAGG3',

Chrebp-alpha-fw: 5'CGACACTCACCCACCTCTTC3', Chrebp-alpha-rv:

5'TTGTTCAGCCGGATCTTGTC3', Chrebp-beta-fw: 5'TCTGCAGATCGCGTGGAG3',

Chrebp-beta-rv: 5'CTTGTCCCGGCATAGCAAC3', Hk2-fw: 5'

ACGGAGCTCAACCAAAACCA3', Hk2-rv: 5'TCCGGAACCGCCTAGAAATC3', Foxk1-fw: 5'

GGCTGTCACTCAGAATGGAA3', Foxk1-rv: 5'GAGGCAGATGTGGTAGTGGAG3', Foxk2-

fw: 5' CCACGGGAACTATCAGTGCT3' , Foxk2-rv: 5' GTCATCCTTTGGGCTGTCTC3,

Adipoq-fw: 5'TGACGACACCAAAAGGGCTC3', Adipoq-rv: 5'

ACGTCATCTTCGGCATGACT3', aP2-fw: 5' TCGGTTCCTGAGGATACAAGAT3', aP2-rv:

5'TTTGATGACTGTGGGATTGAAG3', Tbp-fw: 5'TGCTGTTGGTGATTGT3', Tbp-rv:5'

CTTGTGTGGGAAAGAT3'.

Hexokinase assay

Measurements were performed with a hexokinase assay kit (abcam) following

manufacturer's instructions.

Seahorse analyses

Measurements were performed with an XF96 Extracellular Flux Analyzer (Seahorse

Bioscience of Agilent) following manufacturer's instructions.

Lactate measurement

Differentiated adipocytes were starved serum overnight and stimulated with 100 nM insulin

for 2 hours. Extracellular lactate was measured with the Lactate Pro 2 analyzer (Axonlab).

^{3}H-2DG uptake assay

Differentiated adipocytes were starved for serum for 5 hours and then incubated in Krebs

Ringer Phosphate Hepes (KRPH) buffer with or without 100 nM insulin for 20 min. Cells

were incubated with 50 μM 2DG containing 0.25 μCi ^{3}H-2-deoxyglucose (2DG, Perkin

Elmer) for 5 min and washed three times with cold PBS. Cells were lysed in lysis buffer and

cleared by centrifugation at 14,000 g for 10 min. Incorporated ^{3}H-2DG was measured with a

scintillation counter.

Insulin tolerance test, glucose tolerance test, pyruvate tolerance test

For the insulin and glucose tolerance tests, mice were fasted for 6 hours and insulin

Humalog (Lilly, i.p. 0.75 or 0.5 U/kg body weight) or glucose (2 g/kg body weight) was given,

respectively. For the pyruvate tolerance test, mice were fasted for 15 hours and pyruvate (2

g/kg body weight) was administered. Blood glucose was measured with a blood glucose

meter (Accu-Check).

Polysome profile analyses

Tissues were homogenized in polysome lysis buffer (10 mM Tris-HCl, pH7.5, 140 mM KCl, 5

mM MgCl$_2$, 1mM DTT, 0.5% Triton x-100, 0.05% sodium deoxycholate, 40 u/mL RNase

inhibitor) in the presence or absence of 100 μg/mL cycloheximide and the lysates were

centrifuged twice at 14,000 g for 10 min. The cleared lysates were loaded onto 5-50%

sucrose gradients (10 mM Tris-HCl, pH7.5, 140 mM KCl, 5 mM MgCl$_2$, 5% or 50% sucrose)

and centrifuged with TH-641 (Thermo Scientific) at 36,000 rpm for 2 hours at 4 ºC. Fractions

were collected by a Gradient Station (BIOCOMP INSTRUMENTS) with continuous recording

of optical density (OD) at 254 nm. RNA from each fraction was isolated with TRIzol and

qRT-PCR were performed as described[36].

Insulin ELISA

Plasma insulin levels were measured by ultrasensitive mouse insulin ELISA kit (Crystal

Chem) according to the manufacturer's instructions.

Hepatic triglyceride measurement

Hepatic triglyceride levels were measured using a triglyceride assay kit (Abcam) according

to the manufacturer's instructions.

Body composition measurement

Body composition was measured by nuclear magnetic resonance imaging (Echo Medical

Systems).

Study Approval

All animal experiments were performed in accordance with federal guidelines for animal

experimentation and were approved by the Kantonales Veterinäramt of the Kanton Basel-

Stadt. For human biopsies, the study protocol was approved by the Ethikkommission

Nordwest- und Zentralschweiz (EKNZ).

Statistics

Sample size was chosen according to our previous studies and published reports in which

similar experimental procedures were described. The investigators were not blinded to the

treatment groups. All data are shown as the mean ± SEM. Sample numbers are indicated

in each figure legend. For mouse experiments, n represents the number of animals, and for

cell culture experiments, N indicates the number of independent experiments. To determine

the statistical significance between 2 groups, an unpaired two-tailed Student's t test was

performed. For more than 3 groups, one-way ANOVA was performed. For ITT, GTT, PTT,

weigh curve data, two-way ANOVA was performed. All statistical analysis was performed

using GraphPad Prism 8 (GraphPad Software). A p value of less than 0.05 was considered

statistically significant.

Acknowledgements

We thank Stefan Offermanns (MPI-HLR, Germany), Didier Trono (EPFL, Switzerland),

Robert Weinberg (MIT, USA), Christine Riggenbach (St. Claraspital), Christoph Handschin

(Biozentrum), the Imaging Core Facility (Biozentrum), and the Proteomics Core Facility

(Biozentrum) for providing reagents and technical support. We acknowledge support from

the Swiss National Science Foundation (project 179569 and NCCR 182880 to MNH and

161510 to MS), The Louis Jeantet Foundation (MNH), and the Canton of Basel (MNH). The

authors declare no competing financial interests.

Contributions

MS and MNH conceived the project and designed research with additional contributions

from TM and NR. MS, SS, ICF, DW, ND, AT, DR, NR performed experiments and analyzed

data. NH provided reagents and mice. BKW, ACM, and RP contributed to the collection of

human adipose tissue biopsies. MS and MNH wrote the manuscript.

References

1   Petersen, M. C. & Shulman, G. I. Mechanisms of Insulin Action and Insulin
Resistance. *Physiol Rev* **98**, 2133-2223, doi:10.1152/physrev.00063.2017 (2018).
2   Brown, M. S. & Goldstein, J. L. Selective versus total insulin resistance: a pathogenic
paradox. *Cell Metab* **7**, 95-96, doi:10.1016/j.cmet.2007.12.009 (2008).
3   Wasserman, D. H. Four grams of glucose. *Am J Physiol Endocrinol Metab* **296**, E11-
21, doi:10.1152/ajpendo.90563.2008 (2009).
4   Jackson, R. A., Roshania, R. D., Hawa, M. I., Sim, B. M. & DiSilvio, L. Impact of
glucose ingestion on hepatic and peripheral glucose metabolism in man: an analysis
based on simultaneous use of the forearm and double isotope techniques. *J Clin
Endocrinol Metab* **63**, 541-549, doi:10.1210/jcem-63-3-541 (1986).
5   Kowalski, G. M. & Bruce, C. R. The regulation of glucose metabolism: implications
and considerations for the assessment of glucose homeostasis in rodents. *Am J
Physiol Endocrinol Metab* **307**, E859-871, doi:10.1152/ajpendo.00165.2014 (2014).
6   Abel, E. D. *et al.* Adipose-selective targeting of the GLUT4 gene impairs insulin
action in muscle and liver. *Nature* **409**, 729-733, doi:10.1038/35055575 (2001).
7   Shepherd, P. R. *et al.* Adipose cell hyperplasia and enhanced glucose disposal in
transgenic mice overexpressing GLUT4 selectively in adipose tissue. *J Biol Chem*
268, 22243-22246 (1993).
8   Jiang, Z. Y. *et al.* Insulin signaling through Akt/protein kinase B analyzed by small
interfering RNA-mediated gene silencing. *Proc Natl Acad Sci U S A* **100**, 7569-7574,
doi:10.1073/pnas.1332633100 (2003).
9   Beg, M., Abdullah, N., Thowfeik, F. S., Altorki, N. K. & McGraw, T. E. Distinct Akt
phosphorylation states are required for insulin regulated Glut4 and Glut1-mediated
glucose uptake. *Elife* **6**, doi:10.7554/eLife.26896 (2017).
10 Cybulski, N., Polak, P., Auwerx, J., Ruegg, M. A. & Hall, M. N. mTOR complex 2 in
adipose tissue negatively controls whole-body growth. *Proc Natl Acad Sci U S A* **106**,
9902-9907, doi:10.1073/pnas.0811321106 (2009).
11 Kumar, A. *et al.* Fat cell-specific ablation of rictor in mice impairs insulin-regulated fat
cell and whole-body glucose and lipid metabolism. *Diabetes* **59**, 1397-1406,
doi:10.2337/db09-1061 (2010).
12 Shearin, A. L., Monks, B. R., Seale, P. & Birnbaum, M. J. Lack of AKT in adipocytes
causes severe lipodystrophy. *Mol Metab* **5**, 472-479,
doi:10.1016/j.molmet.2016.05.006 (2016).
13 Tang, Y. *et al.* Adipose tissue mTORC2 regulates ChREBP-driven de novo
lipogenesis and hepatic glucose metabolism. *Nat Commun* **7**, 11365,
doi:10.1038/ncomms11365 (2016).
14 Tan, S. X. *et al.* Selective insulin resistance in adipocytes. *J Biol Chem* **290**, 11337-
11348, doi:10.1074/jbc.M114.623686 (2015).
15 Shimobayashi, M. *et al.* Insulin resistance causes inflammation in adipose tissue. *J
Clin Invest* **128**, 1538-1550, doi:10.1172/JCI96139 (2018).
16 Ducluzeau, P. H. *et al.* Regulation by insulin of gene expression in human skeletal
muscle and adipose tissue. Evidence for specific defects in type 2 diabetes. *Diabetes*
50, 1134-1142, doi:10.2337/diabetes.50.5.1134 (2001).
17 Riddle, M. R. *et al.* Insulin resistance in cavefish as an adaptation to a nutrient-limited
environment. *Nature* **555**, 647-651, doi:10.1038/nature26136 (2018).
18 Nawaz, M. H. *et al.* The catalytic inactivation of the N-half of human hexokinase 2
and structural and biochemical characterization of its mitochondrial conformation.
Biosci Rep **38**, doi:10.1042/BSR20171666 (2018).
19 Heikkinen, S. *et al.* Hexokinase II-deficient mice. Prenatal death of homozygotes
without disturbances in glucose tolerance in heterozygotes. *J Biol Chem* **274**, 22517-
22523, doi:10.1074/jbc.274.32.22517 (1999).

20 Vijayakumar, A. *et al.* Absence of Carbohydrate Response Element Binding Protein
in Adipocytes Causes Systemic Insulin Resistance and Impairs Glucose Transport.
Cell Rep **21**, 1021-1035, doi:10.1016/j.celrep.2017.09.091 (2017).
21 Horton, J. D., Goldstein, J. L. & Brown, M. S. SREBPs: activators of the complete
program of cholesterol and fatty acid synthesis in the liver. *J Clin Invest* **109**, 1125-
1131, doi:10.1172/JCI15593 (2002).
22 Ortega-Prieto, P. & Postic, C. Carbohydrate Sensing Through the Transcription
Factor ChREBP. *Front Genet* **10**, 472, doi:10.3389/fgene.2019.00472 (2019).
23 Herman, M. A. *et al.* A novel ChREBP isoform in adipose tissue regulates systemic
glucose metabolism. *Nature* **484**, 333-338, doi:10.1038/nature10986 (2012).
24 Li, M. V. *et al.* Glucose-6-phosphate mediates activation of the carbohydrate
responsive binding protein (ChREBP). *Biochem Biophys Res Commun* **395**, 395-
400, doi:10.1016/j.bbrc.2010.04.028 (2010).
25 Sukonina, V. *et al.* FOXK1 and FOXK2 regulate aerobic glycolysis. *Nature* **566**, 279-
283, doi:10.1038/s41586-019-0900-5 (2019).
26 Titchenell, P. M. *et al.* Direct Hepatocyte Insulin Signaling Is Required for
Lipogenesis but Is Dispensable for the Suppression of Glucose Production. *Cell*
Metab **23**, 1154-1166, doi:10.1016/j.cmet.2016.04.022 (2016).
27 Karczewski, K. J. *et al.* Variation across 141,456 human exomes and genomes
reveals the spectrum of loss-of-function intolerance across human protein-coding
genes. *bioRxiv*, 531210, doi:10.1101/531210 (2019).
28 Patra, K. C. *et al.* Hexokinase 2 is required for tumor initiation and maintenance and
its systemic deletion is therapeutic in mouse models of cancer. *Cancer Cell* **24**, 213-
228, doi:10.1016/j.ccr.2013.06.014 (2013).
29 Sassmann, A., Offermanns, S. & Wettschureck, N. Tamoxifen-inducible Cre-
mediated recombination in adipocytes. *Genesis* **48**, 618-625, doi:10.1002/dvg.20665
(2010).
30 Chen, S., Lee, B., Lee, A. Y., Modzelewski, A. J. & He, L. Highly Efficient Mouse
Genome Editing by CRISPR Ribonucleoprotein Electroporation of Zygotes. *J Biol*
Chem **291**, 14457-14467, doi:10.1074/jbc.M116.733154 (2016).
31 Haueter, S. *et al.* Genetic vasectomy-overexpression of Prm1-EGFP fusion protein in
elongating spermatids causes dominant male sterility in mice. *Genesis* **48**, 151-160,
doi:10.1002/dvg.20598 (2010).
32 Zebisch, K., Voigt, V., Wabitsch, M. & Brandsch, M. Protocol for effective
differentiation of 3T3-L1 cells to adipocytes. *Anal Biochem* **425**, 88-90,
doi:10.1016/j.ab.2012.03.005 (2012).
33 Stewart, S. A. *et al.* Lentivirus-delivered stable gene silencing by RNAi in primary
cells. *RNA* **9**, 493-501 (2003).
34 Ahrne, E. *et al.* Evaluation and Improvement of Quantification Accuracy in Isobaric
Mass Tag-Based Protein Quantification Experiments. *J Proteome Res* **15**, 2537-
2547, doi:10.1021/acs.jproteome.6b00066 (2016).
35 Wang, Y. *et al.* Reversed-phase chromatography with multiple fraction concatenation
strategy for proteome profiling of human MCF10A cells. *Proteomics* **11**, 2019-2026,
doi:10.1002/pmic.201000722 (2011).
36 Panda, A. C., Martindale, J. L. & Gorospe, M. Polysome Fractionation to Analyze
mRNA Distribution Profiles. *Bio Protoc* **7**, doi:10.21769/BioProtoc.2126 (2017).

Fig. 1

a vWAT

b vWAT　sWAT　BAT　muscle

c vWAT　sWAT

d human oWAT

e

f surface fish / Molino

g

	hyperglycemia	insulin receptor	AKT-pS473
surface fish	-	WT	normal
cavefish Tinaja	+	P211L	impaired
cavefish Pachon	+	P211L	impaired
cavefish Molino	++	WT	normal

h

```
LSDETLLEIS  KRFRKEMEKG  human
LSDETLLEIS  RRFRKEMEKG  rat
LSDETLLEIS  RRFRKEMEKG  mouse
LSEDTLQEVS  ERFRKEMEKG  chicken
LSDETLLEVS  SRFEKQMEKG  frog
LSNENLMDVS  KRFRKEMDKG  zebrafish
LSEENLMDVS  RRFRREMDKG  surface fish
LSEENLMDVS  RHFRREMDKG  cave fish molino
31                  50
```

i HEK293T

HK2-WT	-	+	-
HK2-R42H	-	-	+

Fig. 1. Loss of HK2 in obese mouse, human, and cavefish Molino.

a. The protein levels of Hexokinases and GLUT4 in visceral white adipose tissue (vWAT) of normal diet (ND)- and 4-week high fat diet (HFD)-fed wild-type C57BL/6JRj mice. Multiple t test, ****p<0.0001. n=5 (ND) and 5 (HFD).

b. Immunoblots of vWAT, subcutaneous WAT (sWAT), brown adipose tissue (BAT), and skeletal muscle from ND- and 4-week HFD-fed mice. AKT serves as a loading control. n=5 (ND) and 5 (HFD).

c. Immunoblots of vWAT and sWAT from ND-, 2-week HFD-, and 2-week HFD + 2-week ND-fed mice. n=5 (ND), 5 (HFD), and 6 (HFD+ND).

d. HK activity of omental WAT (oWAT) from lean, obese, and obese diabetic patients. Two-way ANOVA, *p<0.05. n=27 (lean), 30 (obese), and 14 (obese diabetic).

e. HK activity positively correlated with BMI. Pearson's correlation analysis.

f. Surface fish and Mexican cavefish Molino.

g. Comparisons of phenotypes in surface fish, Pachón, Tinaja, and Molino[17].

h. Amino acid sequence alignment of the HK2-R42H mutation within vertebrates.

i. HK activity for lysates of HEK293T cells expressing control, HK2-WT, or HK2-R42H. Student's t test, ****p<0.0001. N=4.

Fig. 2

Fig. 2. Loss of adipose HK2 causes glucose intolerance.
a. Extracellular acidification rate of control and HK2 knockdown 3T3-L1 adipocytes in response to glucose (10 mM) and 2-deoxyglucose (2DG, 50 mM). N=8.
b. Lactate secreted into media by control and HK2 knockdown 3T3-L1 adipocytes treated with or without 100 nM insulin. N=3.
c. 2DG uptake in control and HK2 knockdown 3T3-L1 adipocytes treated with or without 100 nM insulin. N=3.
d. Immunoblots of vWAT, sWAT, and BAT of control and adipocyte-specific Hk2 knockout (AdHk2KO) mice. n=10.
e. Insulin tolerance test (ITT) on control and AdHk2KO mice. The mice were fasted for 6 hours and injected with insulin (0.5 U/kg body weight). Two-way ANOVA. *p<0.05, ***p<0.001. n=6 (control) and 10 (AdHk2KO).
f. Gucose tolerance test (GTT) on control and AdHk2KO mice. Mice were fasted for 6 hours and injected with glucose (2 g/kg body weight). Two-way ANOVA for glucose curves and Student's t test for AUC. *p<0.05, **p<0.01. n=7 (control) and 12 (AdHk2KO).

Fig. 3

Fig. 3. HK2 promotes ChREBP-mediated lipogenesis in adipose tissue and suppresses hepatic gluconeogenesis.

a. Pyruvate tolerance test (PTT) on control and AdHk2KO mice. Mice were fasted for 15 hours and injected with pyruvate (2 g/kg body weight). Two-way ANOVA, *p<0.05, **p<0.01. n=5 (control) and 6 (AdHk2KO).

b. mRNA levels of gluconeogenic genes in liver of control and AdHk2KO mice. Multiple t test, *p<0.05, **p<0.01. n=9 (control) and 12 (AdHk2KO).

c. mRNA levels of fatty acid synthesis genes in liver from control and AdHk2KO mice. Multiple t test, *p<0.05, **p<0.01. n=9 (control) and 12 (AdHk2KO).

d-e. mRNA levels of fatty acid synthesis genes in vWAT (d) and sWAT (e) from control and AdHk2KO mice. Multiple t test, *p<0.05, **p<0.01. n=9 (control) and 12 (AdHk2KO).

Fig. 4

Fig. 4. Mechanism of HFD-induced HK2 down-regulation in adipose tissue.
a. *Hk2* mRNA levels in sWAT, vWAT, and BAT from ND- and 4-week HFD-fed mice. Student's t test, **$p < 0.01$. n=10 (sWAT from ND and HFD), 10 (vWAT from ND and HFD), 8 (BAT from ND), 10 (BAT from HFD).
b. Representative polysome profiles of vWAT from ND- and 4-week HFD-fed mice. Tissue lysates were prepared in the presence of cycloheximide (CHX) to prevent ribosome run-off. n=6 (ND) and 6 (HFD).
c. Representative run-off polysome profiles of vWAT from ND- and 4-week HFD-fed mice. Tissue lysates were prepared in the absence of CHX to allow run-off. n=7 (ND) and 7 (HFD).
d. Representative run-off polysome profiles of vWAT from ND-, 2-week HFD-, 2-week HFD + 2-week ND-fed mice. Tissue lysates were prepared in the absence of CHX. n=5 (ND) and 5 (HFD).
e. *Hk2* mRNA abundance in monosome (mono), light polysome (LP), and heavy polysome (HP) fractions obtained in c. Two-way ANOVA, ***$p < 0.001$, ****$p < 0.0001$. n=4.
f. *Foxk1* and *Foxk2* mRNA levels in sWAT from ND- or 4-week HFD-fed mice. Multiple t test, **$p < 0.01$, ***$p < 0.001$. n=10 (ND) and 10 (HFD).
g. Diet-induced loss of adipose HK2 triggers hyperglycemia via reduced glucose clearance in adipocytes (left) and de-repressed hepatic gluconeogenesis (right).

Extended Data Fig. 1

Extended Data Fig. 1. HFD-fed mice are obese, insulin resistant, and glucose intolerant.
a. Body weight of normal diet (ND)- and 4-week high fat diet (HFD)-fed wild-type
C57BL/6JRj mice. Student's t test, *p<0.05. n=5 (ND) and 5 (HFD).
b. Insulin tolerance test (ITT) on ND- and 4-week HFD-fed wild-type C57BL/6JRj mice. The
mice were fasted for 6 hours and injected with insulin (0.5 U/kg body weight). Two-way
ANOVA for glucose curves and Student's t test for AUC. *p<0.05, ****p<0.0001.
n=5 (ND) and 5 (HFD).
c. Glucose tolerance test (GTT) on 4-week HFD- and ND-fed wild-type C57BL/6JRj mice.
Mice were fasted for 6 hours and i.p. injected with glucose (2 g/kg body weight). Two-way
ANOVA for glucose curves and Student's t test for AUC. *p<0.05, **p<0.01, ***p<0.001.
n=5 (ND) and 8 (HFD).

Extended Data Fig. 2

Extended Data Fig. 2.
Hexokinase (HK) activity of vWAT and sWAT from ND- and 4-week HFD-fed mice.
Student's t test, *p<0.05, **p<0.01. n=5 (ND) and 5 (HFD).

Extended Data Fig. 3

Extended Data Fig.3. Adipose HK2 expression correlates hyperglycemia.
a, **b**. Body weight (a) and fasting glucose levels (b) of mice fed a HFD for 0, 1, 2, or 4 weeks.
One-way ANOVA compared to 0-week HFD-fed mice, **p<0.01, ***p<0.001. n=10
c. Fasting insulin levels of mice fed a HFD for 0, 1, 2, or 4 weeks. One-way ANOVA
compared to 0-week HFD-fed mice, *p<0.05. n=5
d. ITT on mice fed a HFD for 0, 1, 2, 4 weeks, and mice fed 2-week HFD and 2week ND.
The mice were fasted for 6 hours and injected with insulin (0.75 U/kg body weight). Oneway
ANOVA, *p<0.05. n=10 (0-week ND), 10(1-week HFD), 10 (2-week HFD), 10 (2-week
HFD), and 6 (2-week HFD + 2-week ND).
e. Immunoblots of vWAT and sWAT from mice fed a HFD for 0, 1, 2, or 4 weeks. Mice were
fasted for 6 hours and injected with insulin (0.75 U/kg body weight). n=5.

Extended Data Fig. 4

a

```
5'-AGG CGC TTC-3' Surface
5'-AGG CGC TTC-3' Pachón
5'-AGG CGC TTC-3' Tinaja
5'-AGG CAC TTC-3' Molino
```

b

c

	HEK293T		
HK2-WT	-	+	-
HK2-R42H	-	-	+
HK2			
HA tag			
ACTIN			

Extended Data Fig. 4. The biological significance of Arginine 42 in HK2.
a. DNA sequence of the Molino mutation.
b. Structural analyses revealed the presence of a salt bridge between Arginine 42 (R42) and Aspartic acid 272 (D272) in the human HK2 (PDB:2NTZ).
c. Immunoblots for lysates of HEK293T cells expressing control, HK2-WT, or HK2-R42H. N=4.

Extended Data Fig. 5

a

b

Extended Data Fig. 5. Generation of HK2-R42H knock-in mice.
a. Sequences of genomic DNA encoding R42H, CRISPR RNA, and template ultramer surrounding of the R42.
b. Sequence chromatograms of wild-type HK2 and R42H knock-in alleles.

Extended Data Fig. 6

Extended Data Fig. 6. Generation and characterization of HK2 knockdown 3T3-L1 adipocytes.
a. Immunoblots of control and HK2 knockdown 3T3-L1 adipocytes. Cells were stimulated
with 100 nM insulin for 25 min. N=4.
b. Hexokinase (HK) activity of control and HK2 knockdown 3T3-L1 adipocytes. Student's t test,
***p<0.001. N=3.
c. Bright filed images of control and HK2 knockdown 3T3-L1 adipocytes.
d. mRNA levels of mature adipocyte markers in control and HK2 knockdown 3T3-L1 adipocytes.
No significant difference in multiple t test. N=3.

Extended Data Fig. 7

Extended Data Fig. 7. Characterization of AdHk2KO mice.
a. Experimental design of tamoxifen treatment, body composition measurement, ITT, and GTT in control and AdHk2KO mice for Fig. 2d-f. Both control and AdHk2KO mice were treated with tamoxifen.
b. Immunoblots of muscle and liver of control and AdHk2KO mice. n=5 (control) and 5 (AdHk2KO). n=10
c-d. Body weight curve (c), lean and fat mass (d) of control and AdHk2KO mice. Student's t test, *p<0.05. n=15 (control) and 17 (AdHk2KO).
e-f. Organ weight for fat tissues (e) and non-fat tissues (f) of control and AdHk2KO mice. Student's t test, *p<0.05. n=12 (control) and 15 (AdHk2KO).
g. Plasma insulin levels of ad libitum fed control and AdHk2KO mice. n=16 (control) and 16 (AdHk2KO).
h. Hepatic triglyceride levels of control and AdHk2KO mice. n=10 (control) and 10 (AdHk2KO).
i. mRNA levels of fatty acid synthesis genes in BAT from control and AdHk2KO mice. Multiple t test, **p<0.01, ***p<0.001, ****p<0.0001. n=10 (control) and 10 AdHk2KO).

Extended Data Fig. 8

Extended Data Fig. 8. Mechanism of HFD-induced HK2 down-regulation in adipose tissue.
a. *Tbp* mRNA abundance in monosome (mono), light polysome (LP), and heavy polysome
(HP) fractions in Fig. 4c. No significant difference in two-way ANOVA. n=4.
b. *Foxk1* and *Foxk2* mRNA levels in vWAT of ND- or 4-week HFD-fed wild-type C57BL6JRj
mice. No significant difference in multiple t test. n=10 (ND) and 10 (HFD).
c. *Foxk1* and *Foxk2* mRNA levels in BAT of ND- or 4-week HFD-fed wild-type C57BL6JRj
mice. No significant difference in multiple t test. n=8 (ND) and 10 (HFD).