Running Title: Deep, Soft, and Dark Sounds Induce ASMR

Deep, soft, and dark sounds induce autonomous sensory meridian response

Takuya Koumura[1,*], Masashi Nakatani[1,2], Hsin-I Liao[1], and Hirohito M. Kondo[1,3,*,†]

[1]Human Information Science Laboratory, NTT Communication Science Laboratories, NTT

Corporation, Atsugi, Kanagawa 243-0198, Japan

[2]Faculty of Environment and Information Studies, Keio University, Fujisawa, Kanagawa 252-0882

Japan

[3]School of Psychology, Chukyo University, Nagoya, Aichi 466-8666, Japan

[*]T.K. and H.M.K. contributed equally to this work.

[†]Corresponding author:

Hirohito M. Kondo, Ph.D.

School of Psychology

Chukyo University

101-2 Yagoto Honmachi, Showa, Nagoya, Aichi 466-8666, Japan

E-mail: kondo@lets.chukyo-u.ac.jp

Tel: +81 52 835 7160

Number of pages: 20 pages with 6 figures and 2 tables

Number of words for abstract, introduction, and discussion: 196, 519, and 1456

Author contributions: T.K., M.N. and H.M.K. designed research; H.M.K. performed research; T.K.

and H.M.K. analyzed data; T.K., H.I.L., H.M.K. wrote the paper.

Acknowledgements: This study was funded by JSPS KAKENHI grant (no. 17K04494 to H.M.K). We

thank the Institute of Advanced Collaborative Research at Chukyo University for generous support.

Conflict of interest statement: The authors declare no competing interests.

1   **Abstract**

There has been a growing interest in the autonomous sensory meridian response (ASMR). The

ASMR is characterized by a tingling sensation around the scalp and neck and often induces a feeling

of relaxation and a reduction of a negative mood. However, it is still unknown what factors affect

the ASMR. The present study focused on stimulus characteristics and individuals' mood states and

personality traits. Participants filled out self-reported questionnaires (the Profile of Mood States,

Beck Depression Inventory, and Big Five Inventory) and reported ASMR estimates throughout a

17-min experiment while listening to binaural tapping and brushing sounds. Cross-correlation

results showed that the ASMR estimates were strongly associated with the acoustic features of

auditory stimuli, such as their amplitude, spectral centroid, and spectral bandwidth. This indicates

that low-pitched sounds with dark timbre trigger the ASMR. The maximum ASMR was observed

around 2 s after the acoustic features changed, suggesting that the sluggishness of multisensory

integration may lead to the ASMR experience. In addition, individual differences in the ASMR

experience were closely linked to participants' mood states, such as anxiety, but not to their

personality traits. Our results provide important clues to understand the mechanisms of

auditory-somatosensory interactions.

Keywords: ASMR, hearing, somatosensory, interaural difference, proximal space

Significant Statements

The autonomous sensory meridian response (ASMR) is characterized by a tingling, electrostatic-like

sensation across the scalp and back of the neck. This phenomenon can be triggered by a variety of

audiovisual stimuli, and many people seek out the ASMR via the internet to receive a feeling of

relaxation and reduce a negative mood. We show that the ASMR is induced about 2 s after acoustic

features, such as the amplitude, spectral centroid, and spectral bandwidth are changed. This

suggests that low-pitched sounds with dark timbre lead to the ASMR experience. The

stimulus-driven ASMR effect is found regardless of the personality traits or mood states of

participants. Our findings provide a critical clue to understand the mechanisms of

auditory–somatosensory interactions.

1   **Introduction**

Many regular-to-heavy users of the internet are familiar with the autonomous sensory

meridian response (ASMR), a phenomenon often characterized by a tingling, electrostatic-like

sensation across the scalp and back of the neck (Barratt and Davis, 2015). As the ASMR can be

triggered by a variety of audiovisual stimuli, content creators have produced hundreds of

ASMR-inducing videos for video-sharing platforms such as YouTube. Many individuals experience

ASMR to receive a feeling of relaxation, reduce a negative mood, and aid sleep (Barratt and Davis,

2015).

In contrast to the general population's interest in ASMR, few scientific studies sought to

clarify the types of stimuli that most effectively trigger the ASMR and how the ASMR is related to

an individual's mental states and personality traits. Recent studies have used web-based surveys of

the Big Five Inventory (BFI) and demonstrated that individuals with higher Openness and lower

Conscientiousness scores reported more frequent ASMR experiences (McErlean and Banissy, 2017;

Fredborg et al., 2018). However, such studies include an inherent sample selection bias in that

participants recruited through the internet may have the tendency to disclose their ASMR

experiences. In addition, how the stimulus characteristics were controlled is unclear, which may

inherently affect the ASMR.

From the perspective of cross-modal interactions, the ASMR is an intriguing phenomenon

because sounds induce a shiver or thrill that corresponds to a tickling sensation on the skin (Grewe

et al., 2009). There is no doubt that the integration of vision, touch, and proprioception plays a

critical role in shaping the body schema (Maravita et al., 2003; de Vignemont et al., 2005; Haggard

et al., 2007). However, few studies have looked at the contribution of audition to the

somatosensory system. Effects of auditory distractors on tactile discrimination are larger when the

distractors are presented close to the head (20 cm) than when they are presented far from the

head (70 cm) (Kitagawa et al., 2005). Self-produced action sounds bias the body schema and its

spatial boundaries (Tajadura-Jimenez et al., 2012). In addition, auditory feedback can modify the

tactile sensation of a listener experiences when rubbing his/her hands together (Jousmäki and Hari,

1998; Guest et al., 2002). When high-frequency components of the auditory feedback increase, the

perceived roughness/moisture of the palmar skin decreases and its smoothness/dryness increases.

Actually, listeners reported a tickling sensation when listening to sounds while the ear of a dummy

head was stroked with a brush (Kitagawa and Igarashi, 2005). However, it is unclear what factors

transform auditory inputs into the tickling sensation.

The present study examined the degree to which the ASMR is affected by the acoustic

features of auditory stimuli, as well as by mood states and personality traits of individuals.

Participants, all of whom had never experienced the ASMR, were instructed to listen to binaural

tapping and brushing sounds from headphones. We hypothesized that there should be no

confounding factor between acoustic features and an individual's states/traits because these

parameters are essentially independent of each other. The ASMR may be similar to aesthetic chills

in terms of a sound-induced emotional experience. After the experiment, we interviewed listeners

to investigate the pleasantness of the ASMR experience.

16   **Materials and Methods**

Listeners

Thirty college students (13 males and 17 females; mean age 20.6 years, range 18–23 years)

participated in this study. Listeners were right-handed with normal hearing. None had ever viewed

any video created to induce the ASMR experience. All listeners gave written informed consent after

the procedures had been fully explained to them. The procedures reported in this study were

approved by the Ethics Committee of School of Psychology at Chukyo University (protocol no.

RS17-020).

Behavioral tasks

Auditory stimuli were made from a file with Creative Commons licenses that was produced

to induce the ASMR and uploaded to the video-sharing site YouTube

(https://www.youtube.com/watch?v=7MZtaAgqoTY). The total duration of each auditory stimulus

was 1,020 s, and visual information was removed from it. Listeners were instructed to dichotically

listen to the stimulus through headphones (Sennheiser HD 599) and continuously indicate their

sensation using a 3-point Likert scale: "no ASMR", "weak ASMR", or "strong ASMR". In this study,

ASMR was defined as a tingling sensation with "zoku-zoku", which is a Japanese onomatopoeia

word for shivering with a feeling of being excited and frightened (Osaka et al., 2003; Osaka et al.,

2004). Their responses were collected via three keys on a computer keyboard with the sampling

rate of 1 kHz. A key press indicating a response was held until a subsequent key press. The stimulus

presentation and data collection were controlled by a PC with Presentation software

(Neurobehavioral Systems, Berkeley, CA, USA). We interviewed listeners separately after the

experiment. In the interviews, they reported where their tingling sensation originated and whether

the sensation moved to other areas of their body. In addition, we checked the degree of their

ticking sensation and pleasant feeling using a visual analogue scale.

Questionnaires of mood states and personality traits

Before the experiment, listeners filled out questionnaires in a quiet room. We obtained

self-reported measures of listener's mood states and personality traits using the POMS, BDI-II, and

the BFI. The POMS measures transient, distinct mood states for the past one week and consists of

six subscales: Tension-Anxiety (9 items), Depression (15 items), Anger-Hostility (12 items), Vigor (8

items), Fatigue (7 items), and Confusion (7 items) (McNair et al., 1971). Listeners rated their mood

by 65 adjectives (e.g., "Furious", "Hopeless" and "Carefree") on a 5-point Likert scale from "not at

all" to "extremely". The BDI is widely used as an indicator of the severity of depression in

adolescents and adults (Beck et al., 1996). Listeners responded to 21 items based on their mental

states over the past two weeks. The BFI is used to measure the individual's basic personality traits:

Extraversion (8 items), Agreeableness (9 items), Conscientiousness (9 items), Neurotisism (8 items),

and Openness (10 items) (John and Srivastava, 2001; Kondo et al., 2017). Listeners rated these 44

items on a 5-point Likert scale ranging from "strongly disagree" to "strongly agree". One listener

was excluded from the subsequent analysis on mood states and personality traits because his/her

data on the questionnaires were missing.

Data analyses

ASMR data analysis

The time-series data of the ASMR estimates were converted to numerical values of 0, 1, and

2 (Fig. 1). Based on the interviews with participants, the ASMR-induced tingling sensation was

visualized on areas of the body (Fig. 2). We performed a correlation analysis to check the

relationships between averaged ASMR estimates, mood states, and personality traits, and

calculated a significance level. Statistical analyses were carried out with Python, R (version 3.1.2),

and IBM SPSS Statistics (version 23).

Acoustic features of auditory inputs

The cochlear spectrogram was calculated from the auditory stimuli presented to each ear.

The sounds were recorded with an artificial ear (TYPE2015, ACO, Tokyo) with the sampling rate of

44.1 kHz. The recorded waveforms were decomposed into frequency components by a gammatone

filter bank (Patterson et al., 1992). The center frequencies of the filters were spaced one equivalent

rectangular bandwidth (ERB) apart (Glasberg and Moore, 1990), and the lowest center frequency

was set to 20 Hz, resulting in 42 filters with the highest center frequency of 20.562 kHz. To simulate

non-linear compression in the auditory periphery, amplitude envelopes of the

frequency-decomposed signals were calculated with Hilbert transformation, powered by 0.3

(McDermott and Simoncelli, 2011), and down-sampled to 2 kHz.

The auditory features obtained for both the left and right channels were the amplitude,

spectral centroid, spectral bandwidth, and instantaneous roughness. The amplitude is the mean of

the cochlear spectrogram over frequencies:

$$Amplitude(t, c) = \text{mean}_f\big(S(f, t, c)\big),$$

where f, t, and c denote a frequency bin in the ERB scale, time, and a channel indicating left or right,

respectively. $S(f, t, c)$ represents a cochlear spectrogram. The spectral centroid indicates the gravity

center in the frequency domain:

$$Centroid(t,c) = \frac{\sum_f f S(f,t,c)}{\sum_f S(f,t,c)}.$$

The spectral bandwidth is the width of the mass distribution in the frequency domain:

$$Bandwidth(t,c) = \sqrt{\frac{\sum_f S(f,t,c)(f-Centroid(t,c))^2}{\sum_f S(f,t,c)}}.$$

Roughness is often calculated from a sound segment with a certain duration. To evaluate temporal

variation of the stimulus, we calculated roughness at every time point, that is, an instantaneous

roughness. For the instantaneous roughness, the cochlear spectrogram at each center frequency

was filtered with a bandpass filter with a passband frequency between 30 and 150 Hz (Arnal et al.,

2015). Instantaneous roughness was defined as

$$Roughness(t,c) = \text{mean}_f\left(\log\left(S_{bp}(f,t,c)^2\right)\right),$$

where S_{bp} denotes the bandpass-filtered spectrogram.

The four features described above were down-sampled to 10 Hz. We computed the average

and the absolute difference of the features in the left and the right ears.

$$AveF(t) = \frac{F(t,left) + F(t,right)}{2},$$

$$DiffF(t) = |F(t,left) - F(t,right)|,$$

where F indicates amplitude, centroid, bandwidth, or roughness. This resulted in totally 8 features

= 4 (amplitude, spectral centroid, spectral bandwidth, and instantaneous roughness) × 2 (average

and absolute difference). We refer to them as Ave- and Diff-amplitude, Ave- and Diff-centroid, Ave-

and Diff-bandwidth, and Ave- and Diff-roughness.

Correlations between ASMR and acoustic features

We calculated cross-correlations between the time course of the ASMR estimates and

acoustic features. The ASMR estimates were z-scored for each listener and averaged over listeners.

The correlations were assessed by Spearman's rank-order correlations. Statistical significance of

the correlation coefficient at the peak of the cross-correlogram was tested by comparing the peak

correlation coefficient with 16,000 cross-correlogram calculated from randomly shifted time

courses. We also calculated a cross-correlogram for each listener. The cross-correlations did not

reach statistical significance for one listener on the amplitude; for two listeners on the spectral

centroid; and for five listeners on the spectral bandwidth. Consequently, we excluded the data of

six listeners from a repeated-measures ANOVA on the time lags of cross-correlations when

comparing between the acoustic features ($N = 24$).

Relationship between ASMR and individual's states and traits

For the factor analysis, we checked the Kaiser–Meyer–Olkin (KMO) statistic using zero-order

correlations and partial correlations to test whether the variables in our dataset were adequate to

correlate. The KMO statistic (0.67) indicated that some common factor underlay our dataset,

because its estimate was higher than 0.50 for proceeding with a satisfactory factor analysis.

Bartlett's test of sphericity showed that correlations between the variables were greater than

those expected by chance: $\chi^2 = 255.04$, $p < 0.001$. The factors were extracted by least squares

estimation and then subjected to an oblique promax rotation.

17   **Results**

ASMR experience

Time-series data of auditory stimuli and ASMR estimates are shown in Figure 1. Tapping and

brushing sounds were used as auditory stimuli to induce the ASMR. The results show not only

interindividual variations of the ASMR estimates but also similar temporal changes across

participants. After the experiment, ninety percent of the participants reported an ASMR-induced

tingling sensation on different parts of their body (Fig. 2a). Of those parts, it mainly originated on

the ears and their vicinities (59%), the neck (44%), the shoulders (44%), and the spine and back

(30%). Our results are consistent with previous findings on the ASMR experience (Barratt and Davis,

2015). A previous study on music-induced chills found that most participants experienced

goosebumps on their arms, whereas less than half felt anything in their spine (Craig, 2005). In this

study, however, small numbers of participants reported the tingling sensation on their arms (15%)

and legs (0%). Using a visual analogue scale (range 0 to 100), we conducted an experiment to check

the subjective intensity of participants' tingling and pleasure (Fig. 2b). The intensity (mean ± SE)

was greater for tickling (51.2 ± 5.3) than for pleasure (35.7 ± 3.7): $t = 2.43$, Cohen's $d = 0.63$, $p =$

0.022. There was no significant correlation between the intensity of the two measures: $r = 0.03$, $p >$

0.87. In addition, the ASMR estimates were positively correlated with the intensity of the tingling

sensation ($r = 0.74$, $p < 0.001$), but not with the intensity of pleasantness ($r = 0.00$, $p > 0.98$). Thus,

the results suggest that the ASMR is not necessary linked with pleasant experiences, at least not for

the stimuli used in the present study.

Correlations between ASMR and acoustic features

We performed a cross-correlation analysis to examine whether and how each acoustic

feature contributed to the ASMR as a function of time (Fig. 3). The acoustic features we examined

were the amplitude, spectral centroid, spectral bandwidth, and instantaneous roughness. We

computed an average (Ave) of the features presented to the left and right ears and the difference

(Diff) in the features between the two ears. For the Ave-features, the ASMR estimates were

positively correlated with the Ave-amplitude ($r_s = 0.37$, $p < 0.001$), indicating that auditory stimuli

with larger sound pressure levels induce greater ASMR estimates (Fig. 3). The ASMR estimates were

negatively correlated with the Ave-centroid ($r_s = -0.48$, $p < 0.001$) and Ave-bandwidth ($r_s = -0.34$, $p <$

0.001). This suggests that low-pitched sounds with dark timbre (a centroid frequency of less than

1.5 kHz) lead to an ASMR experience. In contrast, the correlation between the ASMR estimates and

Ave-roughness did not reach statistical significance ($r_s = 0.24$, $p = 0.088$).

We also found significant cross-correlations between the ASMR estimates and all the

Diff-features except for instantaneous roughness: $r_s = 0.28$, $p < 0.001$ for the Diff-amplitude; $r_s =$

0.25, $p < 0.001$ for the Diff-centroid; $r_s = 0.24$, $p < 0.001$ for the Diff-bandwidth; cf., $r_s = 0.20$, $p =$

0.053 for the Diff-roughness (Fig. 4). This means that as the larger the interaural differences in the

four acoustic features were, the larger the ASMR estimates became. Intriguingly, the peak of

cross-correlation coefficients had a time lag of around 2 s. These results indicate that changes in

the acoustic features of auditory stimuli trigger the ASMR.

We further examined the time lag of cross-correlations using the individual differences

approach. The Ave- and Diff-roughness features were excluded from the subsequent analyses

because their cross-correlations did not reach statistical significance. We performed a 2 (Ave and

Diff) × 3 (amplitude, centroid, and bandwidth) analysis of variance (ANOVA) on the time lags to the

peak of cross-correlations (Fig. 5). The time lag of Diff-features (1.78 ± 0.17 s) was shorter than that

of Ave-features (2.22 ± 0.19 s): $F_{(1, 23)}$ = 25.38, η_p^2 = 0.53, $p < 0.001$. The main effect ofthe acoustic

features did not reach statistical significance—1.93 ± 0.15 s for amplitude; 1.81 ± 0.14 s for

centroid; 2.17 ± 0.26 s for bandwidth; $F_{(2, 46)}$ = 2.45, η_p^2 = 0.10, p = 0.097—and neither did the

interaction: $F_{(2, 46)}$ = 2.51, η_p^2 = 0.10, p = 0.092. Therefore, our results suggest that the ASMR is

caused by interaural differences in the acoustic features and consequently affected by the average

of those features.

Effects of states and traits on ASMR

So far, we have demonstrated effects of acoustic features on the ASMR. Next, we

investigated whether the ASMR is associated with the mood states and personality traits of the

listeners, which were assessed using the Profile of Mood States (POMS), Beck Depression Inventory

(BDI-II), and the BFI (Table 1). Cronbach's alpha of the measures reached a satisfactory level (range

of 0.65 to 0.93), suggesting that these measures have a high level of internal consistency. The

ASMR estimates were significantly correlated with the POMS Anxiety score: r = 0.52, p = 0.004

(Table 2). There were mild positive correlations (0.42 to 0.46) between the ASMR estimates and

POMS Fatigue, Depression, and Confusion scores. In contrast, the correlations with the BFI scores

were generally low (less than 0.32). The overall results indicate that individual's mood states,

relative to his/her personality traits, are involved in the ASMR experience. We used a multiple

regression analysis to specify the degree to which the twelve states/traits scores contribute to the

ASMR estimates. The results of a stepwise analysis showed that the POMS Anxiety score accounted

for 24.5% of the variance of the ASMR estimates: adjusted R^2 = 0.208, $F_{(1, 20)}$ = 6.50, p = 0.02.

However, the contribution of the other measures was not significant. Thus, it is likely that

individuals with anxiety or depressive states are sensitive to the ASMR.

We performed a factor analysis to further examine the relationships between the ASMR

estimates and states/traits scores. The BFI Agreeableness score was excluded from the subsequent

analyses because it was closely related to the BFI Conscientiousness score (r = 0.51, p = 0.005) and

its reliability was relatively low (0.65). Two factors, whose eigenvalues were larger than 1, were

extracted from the 12 subscales (Fig. 6). The first factor, with an eigenvalue of 5.78 before the

promax rotation, was heavily loaded on the ASMR estimates, all the POMS subscores except the

Vigor score, the BDI score, and the BFI Neuroticism score (factor loadings, 0.48 to 0.93). The second

factor, with an eigenvalue of 1.68, was loaded on the BFI Extraversion and Openness scores and the

POMS Vigor score (0.49 to 0.97). The first and second factors were termed the "anxiety" and

"activity", respectively. The contribution of the "anxiety" factor (0.51) to the ASMR estimates was

much greater than that of the "activity" factor (0.05).

17   **Discussion**

The present results demonstrate that the intensity of ASMR estimates varied across listeners,

although its sensation mainly originated on their ears, neck, and shoulders. The ASMR was closely

linked to the tickling sensation but not to the pleasant-unpleasant dimension. The ASMR was

affected by the acoustic features of auditory stimuli. Specifically, the cross-correlation results

showed that the ASMR was induced by interaural differences in the acoustic features, such as the

amplitude, spectral centroid, and spectral bandwidth. The maximum ASMR was observed around 2

s after the acoustic features changed. In addition, it was associated with listener's mood states,

including anxiety. In what follows, we mainly discuss effects of stimulus characteristics on the

ASMR.

Many internet users expect the ASMR to provide a feeling of relaxation (Barratt and Davis,

2015). Our preliminary study using a test-retest paradigm indicated that the ASMR estimates have

high reliability within ASMR-inexperienced participants: $r = 0.87$, $p = 0.001$, $N = 10$. It seems that

the ASMR differs from music-induced chills in terms of physiological, emotional, and

phenomenological aspects. Piloerection during chills is innervated by the sympathetic nervous

system (Lindsley and Sassaman, 1938; Craig, 2005). There is no consensus that the parasympathetic

nervous system is involved in the ASMR (Poerio et al., 2018). A profound emotional experience,

such as weeping, sometimes arises in response to music-induced chills (Sloboda, 1991; Panksepp,

1995; Mori and Iwanaga, 2017), but not in response to the ASMR. In contrast to aesthetic chills, the

ASMR is frequently triggered by a variety of stimuli, such as whispering, close personal attention,

and crisp sounds (Barratt and Davis, 2015). The present study investigated common factors (i.e.,

acoustic features) affecting various ASMR stimuli.

We have demonstrated that the Diff-features of auditory stimuli play an important role in

triggering the ASMR. Specifically, a larger Diff-amplitude parameter resulted in larger ASMR

estimates. The Diff-amplitude used in the present study corresponds to an interaural level

difference (ILD). Thus, the results can be interpreted as indicating that the ASMR is elicited when

listeners perceive sound sources near their ears. This is consistent with the findings that the ASMR

is longer during binaural listening than during diotic listening (Liao et al., 2017) and stronger for

sounds moving around the head than for static sounds (Honda et al., 2019). In a previous study,

listeners reported a stronger tickling sensation when the sound of an ear being tickled was

presented with headphones than when the sound was presented via a loudspeaker placed 80 cm

from their ears (Kitagawa and Igarashi, 2005). Therefore, it is likely that auditory–somatosensory

interactions observed as the ASMR are related to sound localization ability in the space close to the

head.

Most sound localization studies have focused on auditory events in the distal region (i.e.,

distances of more than 1 m from listeners) (Brungart and Rabinowitz, 1999). This may be because

the detection of distant events in an environment is believed to be necessary for the survival of

humans and mammals (Heffner and Heffner, 1992; Kitagawa and Spence, 2006). In general, the ILD

and interaural time difference (ITD) are used as localization cues, respectively, when high- and

low-frequency sounds reach the ears (Rayleigh, 1907; Moore, 2013). ILDs can be accurately

obtained at high frequencies (more than 1.5 kHz) because diffraction due to the head hardly occurs,

whereas ITDs can be easily detected at low frequencies because phase ambiguity hardly occurs.

Intriguingly, a previous study demonstrated that ILDs, even at low frequencies, increase when a

sound source is located less than 50 cm from the head (Brungart and Rabinowitz, 1999).Thus, we

believe that listeners in our experiment used ILDs, as well as ITDs, to estimate the distance and

direction of a sound source, which results in the enhancement of sensitivity to the ASMR.

Our results showed that the Ave-centroid feature was negatively correlated with the ASMR

estimate. The spectral centroid has been shown to be correlated with the perceptual brightness of

musical sounds with clear pitch (Almeida et al., 2017). Although our stimuli were non-musical

natural sounds, a previous study revealed that both pitched and unpitched sounds induce timbre in

a common perceptual space, which includes the spectral centroid as one of its major dimensions

(Lakatos, 2000). Thus, it is likely that sounds with a dark impression are closely linked to the ASMR

experience.

In contrast, the Ave- and Diff-roughness had a limited effect on the ASMR. It is known that

roughness in sounds such as screams induces unpleasant perception (Arnal et al., 2015). The

perceptual attribute can be differentiated from other communication signals such as speech. In this

study, the cross-correlation between ASMR estimates and Ave-roughness was not significant but

was positive. This is consistent with the behavioral results indicating that the ASMR is not

necessarily a pleasant experience for some listeners. Alternatively, listeners might experience an

unpleasant feeling because the auditory stimuli were not accompanied with visual information.

When visual cues are not available, interactions between auditory and somatosensory information

occur easily in the space behind the back (Kóbor et al., 2006; Zampini et al., 2007). It has been

believed that many people expect to receive a feeling of relaxation and/or pleasure from the ASMR.

Thus, a future study should investigate the relationship between the ASMR and unpleasant

feelings.

It may be hypothesized that a sluggish integration of multisensory information, in particular

between audition and tactile sensation, leads to the ASMR. Whether the tickling sensation occurs

or not appears to depend on comparing an internal prediction with actual sensory feedback,

resulting in the phenomenon where a self-produced tactile sensation differs from an externally

produced one (Blakemore et al., 1998). In the present study, the maximum ASMR occurred around

2 s the Diff-features had been changed. Previous studies using auditory streaming showed that

effects of head movement on perceptual formation are maximized 1.8 s after motion onsets

(Kondo et al., 2012; Kondo et al., 2014). Intriguingly, the time scale of auditory–somatosensory

interactions is similar to that of auditory–motor interactions. Thus, there is the possibility that the

tickling sensation accompanying the ASMR is derived from the sluggishness of multisensory

integration processes.

The auditory and somatosensory systems have similar temporal frequency channels (Yau et

al., 2009; Occelli et al., 2011). Many neuropsychological studies have investigated somatosensory

responses in the auditory cortex. Neuronal responses in the caudomedial area of the auditory

cortex have been elicited by cutaneous stimulation on the neck and hands of monkeys (Schroeder

et al., 2001; Fu et al., 2003). Tactile stimulation has been shown to enhance the activity of posterior

auditory areas (Foxe et al., 2002; Kayser et al., 2005; Schürmann et al., 2006). In contrast, little is

known about how neurons in the cerebral cortex encode information on auditory distance. A

substantial study demonstrated that tactile-sensitive neurons in the ventral premotor area respond

to auditory stimuli presented within 30 cm from the head (Graziano et al., 1999) and that neural

responses decrease when they are presented at distances of 50 cm from the head. The traditional

view on sensory systems is that different sensory information is separately analyzed in unisensory

areas and subsequently projected into multisensory areas (Felleman and Van Essen, 1991).

However, recent fMRI studies have revealed that auditory stimulation influences the

somatosensory area (Liang et al., 2013; Pérez-Bellido et al., 2018). Thus, the present findings on the

ASMR could contribute to clarifying neural mechanisms of auditory–somatosensory integration.

The results of a factor analysis showed that listeners with a higher anxiety state were more

sensitive to the ASMR. An early study also reported that ASMR media improve participant's mood,

with the positive effect lasting several hours (Barratt and Davis, 2015). In contrast to previous

findings (McErlean and Banissy, 2017; Fredborg et al., 2018), the ASMR estimates were not

correlated with listener's personality traits. The discrepancy in these results may be due to our

using ASMR-inexperienced listeners in this study. Participants' characteristics and learning effects

probably shape the preference for ASMR media.

Previous studies have focused on the relationship between the ASMR experience and an

individual's personality traits, but have not paid attention to effects of sensory input itself on the

ASMR. An essential aspect of the ASMR is that auditory stimuli produce different forms of

perception at the surface of the body or inside it. The present study demonstrated that the ASMR is

induced by changes in interaural differences in the acoustic features, namely the amplitude,

spectral centroid, and spectral bandwidth. We pointed out that the sluggishness of multisensory

integration may lead to ASMR. We also found that the ASMR is associated with individual's mood

states rather than with his/her personality traits and that people with an anxious state tend to

experience a strong ASMR. Our results provide important clues to understand the mechanisms of

auditory–somatosensory interactions.

References

Almeida A, Schubert E, Smith J, Wolfe J (2017) Brightness scaling of periodic tones. Atten Percept
3   Psychophys 79:1892-1896. https://doi.org/10.3758/s13414-017-1394-6.

4   Arnal LH, Flinker A, Kleinschmidt A, Giraud AL, Poeppel D (2015) Human screams occupy a privileged niche in
5   the communication soundscape. Curr Biol 25:2051-2056.
6   https://doi.org/10.1016/j.cub.2015.06.043.

7   Barratt EL, Davis NJ (2015) Autonomous Sensory Meridian Response (ASMR): a flow-like mental state. PeerJ
8   3:e851. https://doi.org/10.7717/peerj.851.

9   Beck AT, Steer RA, Brown G (1996) Manual for the Beck Depression Inventory-II. San Antonio, TX:
10   Psychological Corporation.

11   Blakemore S-J, Wolpert DM, Frith CD (1998) Central cancellation of self-produced tickle sensation. Nat
12   Neurosci 1:635-640. https://doi.org/10.1038/2870.

13   Brungart DS, Rabinowitz WM (1999) Auditory localization of nearby sources. Head-related transfer functions.
14   J Acoust Soc Am 106:1465-1479. https://doi.org/10.1121/1.427180.

15   Craig DG (2005) An exploratory study of physiological changes during "chills" induced by music Music Sci
16   9:273-287. https://doi.org/10.1177/102986490500900207.

17   de Vignemont F, Ehrsson HH, Haggard P (2005) Bodily illusions modulate tactile perception. Curr Biol
18   15:1286-1290. https://doi.org/10.1016/j.cub.2005.06.067.

19   Felleman DJ, Van Essen DC (1991) Distributed hierarchical processing in the primate cerebral cortex. Cereb
20   Cortex 1:1-47. https://doi.org/10.1093/cercor/1.1.1.

21   Foxe JJ, Wylie GR, Martinez A, Schroeder CE, Javitt DC, Guilfoyle D, Ritter W, Murray MM (2002)
22   Auditory-somatosensory multisensory processing in auditory association cortex: an fMRI study. J
23   Neurophysiol 88:540-543. https://doi.org/10.1152/jn.2002.88.1.540.

24   Fredborg BK, Clark JM, Smith SD (2018) Mindfulness and autonomous sensory meridian response (ASMR).
25   PeerJ 6:e5414. https://doi.org/10.7717/peerj.5414.

26   Fu KMG, Johnston TA, Shah AS, Arnold L, Smiley J, Hackett TA, Garraghty PE, Schroeder CE (2003) Auditory
27   cortical neurons respond to somatosensory stimulation. J Neurosci 23:7510-7515.
28   https://doi.org/10.1523/JNEUROSCI.23-20-07510.2003.

29   Glasberg BR, Moore BCJ (1990) Derivation of auditory filter shapes from notched-noise data. Hear Res
30   47:103-138. https://doi.org/10.1016/0378-5955(90)90170-T.

31   Graziano MS, Reiss LA, Gross CG (1999) A neuronal representation of the location of nearby sounds. Nature
32   397:428-430. https://doi.org/10.1038/17115.

33   Grewe O, Kopiez R, Altenmüller E (2009) Chills as an indicator of individual emotional peaks. Ann N Y Acad

Sci 1169:351-354. https://doi.org/10.1111/j.1749-6632.2009.04783.x.

Guest S, Catmur C, Lloyd D, Spence C (2002) Audiotactile interactions in roughness perception. Exp Brain Res
146:161-171. https://doi.org/10.1007/s00221-002-1164-z.

Haggard P, Christakou A, Serino A (2007) Viewing the body modulates tactile receptive fields. Exp Brain Res
180:187-193. https://doi.org/10.1007/s00221-007-0971-7.

Heffner RS, Heffner HE (1992) Visual factors in sound localization in mammals. J Comp Neurol 317:219-232.
https://doi.org/10.1002/cne.903170302.

Honda S, Ishikawa Y, Konno R, Imai E, Nomiyama N, Sakurada K, Koumura T, Kondo H, Furukawa S, Fujii S,
Nakatani M (2019) Proximal binaural sound can induce subjective frisson. arXiv:190406851.
https://doi.org/arxiv.org/abs/1904.06851.

John OP, Srivastava S (2001) The big-five trait taxonomy: history, measurement, and theoretical perspectives.
In: Handbook of Personality: Theory and Research, 2nd Edition (Pervin LA, John OP, eds), pp 102-138.
New York: Guilford Press.

Jousmäki V, Hari R (1998) Parchment-skin illusion: sound-biased touch. Curr Biol 8:R190.
https://doi.org/10.1016/S0960-9822(98)70120-4.

Kayser C, Petkov CI, Augath M, Logothetis NK (2005) Integration of touch and sound in auditory cortex.
Neuron 48:373-384. https://doi.org/10.1016/j.neuron.2005.09.018.

Kitagawa N, Igarashi Y (2005) Tickle sensation induced by hearing a sound. Jpn J Psychonomic Sci
24:121-122.

Kitagawa N, Spence C (2006) Audiotactile multisensory interactions in human information processing. Jpn
Psychol Res 48:158-173. https://doi.org/10.1111./j.1468-5884.2006.00317.x.

Kitagawa N, Zampini M, Spence C (2005) Audiotactile interactions in near and far space. Exp Brain Res
166:528-537. https://doi.org/10.1007/s00221-005-2393-8.

Kóbor I, Füredi L, Kovács G, Spence C, Vidnyánszky Z (2006) Back-to-front: improved tactile discrimination
performance in the space you cannot see. Neurosci Lett 400:163-167.
https://doi.org/10.1016/j.neulet.2006.02.037.

Kondo HM, Pressnitzer D, Toshima I, Kashino M (2012) Effects of self-motion on auditory scene analysis.
Proc Natl Acad Sci U S A 109:6775-6780. https://doi.org/10.1073/pnas.1112852109.

Kondo HM, Toshima I, Pressnitzer D, Kashino M (2014) Probing the time course of head-motion cues
integration during auditory scene analysis. Front Neurosci 8:170.
https://doi.org/10.3389/fnins.2014.00170.

Kondo HM, Farkas D, Denham SL, Asai T, Winkler I (2017) Auditory multistability and neurotransmitter
concentrations in the human brain. Philos Trans R Soc B 372:20160110.
https://doi.org/10.1098/rstb.2016.0110.

Lakatos S (2000) A common perceptual space for harmonic and percussive timbres. Percept Psychophys 62:1426-1439. https://doi.org/10.3758/bf03212144.

Liang M, Mouraux A, Hu L, Iannetti GD (2013) Primary sensory cortices contain distinguishable spatial patterns of activity for each sense. Nat Commun 4:1979. https://doi.org/10.1038/ncomms2979.

Liao H-I, Nakatani M, Miyazaki H, Furukawa S (2017) Sensing frisson by material sounds. In: 21st Annual Meeting of the Association for the Scientific Study of Consciousness. Beijing, China.

Lindsley DB, Sassaman WH (1938) Autonomic activity and brain potentials associated with "voluntary" control of the pilomotors (mm. Arrectores pilorum). J Neurophysiol 1:342-349. https://doi.org/10.1152/jn.1938.1.4.342.

Maravita A, Spence C, Driver J (2003) Multisensory integration and the body schema: close to hand and within reach. Curr Biol 13:R531-R539. https://doi.org/10.1016/s0960-9822(03)00449-4.

McDermott JH, Simoncelli EP (2011) Sound texture perception via statistics of the auditory periphery: evidence from sound synthesis. Neuron 71:926-940. https://doi.org/10.1016/j.neuron.2011.06.032.

McErlean ABJ, Banissy MJ (2017) Assessing individual variation in personality and empathy traits in self-reported autonomous sensory meridian response. Multisensory Res 30:601-613. https://doi.org/10.1163/22134808-00002571.

McNair DM, Lorr M, Droppleman L (1971) Manual for the Profile of Mood States. San Diego, CA: Educational and Industrial Testing Service.

Moore BCJ (2013) An Introduction to the Psychology of Hearing, 6th Edition. London: Brill.

Mori K, Iwanaga M (2017) Two types of peak emotional responses to music: The psychophysiology of chills and tears. Sci Rep 7:46063. https://doi.org/10.1038/srep46063.

Occelli V, Spence C, Zampini M (2011) Audiotactile interactions in temporal perception. Psychon Bull Rev 18:429-454. https://doi.org/10.3758/s13423-011-0070-4.

Osaka N, Osaka M, Morishita M, Kondo H, Fukuyama H (2004) A word expressing affective pain activates the anterior cingulate cortex in the human brain: an fMRI study. Behav Brain Res 153:123-127. https://doi.org/10.1016/j.bbr.2003.11.013.

Osaka N, Osaka M, Kondo H, Morishita M, Fukuyama H, Shibasaki H (2003) An emotion-based facial expression word activates laughter module in the human brain: a functional magnetic resonance imaging study. Neurosci Lett 340:127-130. https://doi.org/10.1016/S0304-3940(03)00093-4.

Panksepp J (1995) The emotional sources of "chills" induced by music. Music Percept 13:171-207. https://doi.org/10.2307/40285693.

Patterson RD, Robinson K, Holdsworth J, McKeown D, Zhang C, Allerhand M (1992) Complex sounds and auditory images. In: Auditory Physiology and Perception, Proc. 9th Intl. Symp. Hear. (Cazals Y, Demany L, Horner K, eds), pp 429-446. Oxford: Pergamon.

Pérez-Bellido A, Anne Barnes K, Crommett LE, Yau JM (2018) Auditory Frequency Representations in Human
Somatosensory Cortex. Cereb Cortex 28:3908-3921. https://doi.org/10.1093/cercor/bhx255.

Poerio GL, Blakey E, Hostler TJ, Veltri T (2018) More than a feeling: Autonomous sensory meridian response
(ASMR) is characterized by reliable changes in affect and physiology. PLoS ONE 13:e0196645.
https://doi.org/10.1371/journal.pone.0196645.

Rayleigh OM (1907) On our perception of sound direction. Philosophical Magazine 13:214-232.
https://doi.org/10.1080/14786440709463595.

Schürmann M, Caetano G, Hlushchuk Y, Jousmaki V, Hari R (2006) Touch activates human auditory cortex.
Neuroimage 30:1325-1331. https://doi.org/10.1016/j.neuroimage.2005.11.020.

Schroeder CE, Lindsley RW, Specht C, Marcovici A, Smiley JF, Javitt DC (2001) Somatosensory input to
auditory association cortex in the macaque monkey. J Neurophysiol 85:1322-1327.
https://doi.org/10.1152/jn.2001.85.3.1322.

Sloboda JA (1991) Music structure and emotional response: some empirical findings. Psychology of Music
19:110-120. https://doi.org/10.1177/0305735691192002.

Tajadura-Jimenez A, Valjamae A, Toshima I, Kimura T, Tsakiris M, Kitagawa N (2012) Action sounds
recalibrate perceived tactile distance. Curr Biol 22:R516-517.
https://doi.org/10.1016/j.cub.2012.04.028.

Yau JM, Olenczak JB, Dammann JF, Bensmaia SJ (2009) Temporal frequency channels are linked across
audition and touch. Curr Biol 19:561-566. https://doi.org/10.1016/j.cub.2009.02.013.

Zampini M, Torresan D, Spence C, Murray MM (2007) Auditory-somatosensory multisensory interactions in
front and rear space. Neuropsychologia 45:1869-1877.
https://doi.org/10.1016/j.neuropsychologia.2006.12.004.

Figure 1

Figure 1. Auditory stimuli and ASMR estimates. (A) Cochlea spectrograms of the stimuli recorded

by the artificial ear. (B) Results of the ASMR estimates. Top and bottom panels represent

time-series data for each listener and averaged z-scores across the listeners, respectively. The right

panel represents temporal average in each listener.

1   **Figure 2**

Figure 2.   ASMR-induced tingling sensation and pleasant feeling. (a) An illustration of tickling

sensation. Red spots indicate an origin of the sensation, whereas areas colored in pale blue

represent the sensation spreading to other parts of the body. (b) The relationship between tickling

sensation and pleasant feeling. Bars represent the means of subjective rating, whereas circles

indicate individual data. Error bars represent the standard errors of means. The dashed line

represents a linear regression. *$p < 0.05$.

Figure 3

Figure 3.   Results of correlation analysis on the ASMR estimates. Plots of ASMR estimates

(*z*-scores) and each acoustic feature. ASMR estimates are shifted by the peak lags of

cross-correlations. When the Ave-amplitude was larger, the ASMR estimate was greater. In

contrast, when Ave-centroid and Ave-bandwidth were smaller, it was greater. Average values of

corresponding physical dimensions are shown below the panels, although they do not have a

one-to-one correspondence to acoustic feature values. AU: arbitrary unit. ERB: equivalent

rectangular bandwidth scale.

                                                    Figure 4

5   **Figure 4.**   Cross-correlations between the ASMR estimates and acoustic features. Areas shaded in

6   grey and dashed lines represent statistical significance after Bonferroni correction over the eight

7   features. The right panels show magnified plots around peaks of cross-correlations. The positive

8   time lag indicates that the changes in the acoustic features are followed by the ASMR.

1   **Figure 5**

3   **Figure 5.** Time lags of peak cross-correlations for the acoustic features. Box-and-whisker plots

4   indicate the median, 25th and 75th percentiles, and outliers for the distribution across listeners. *p

5   < 0.05.

Figure 6

Figure 6.   Results of the factor analysis. The filled and open circles indicate the subscales of

questionnaires on individual's mood states and personality traits, respectively. Factors 1 and 2 can

be considered the "anxiety" and ''activity'' factors. A-H: Anger-Hostility. T-A: Tension-Anxiety.

1   **Table 1   Descriptive statistics for ASMR estimates and individual's states/traits**

Measure	Mean	SD	Min	Max	Skewness	Kurtosis	Reliability
ASMR	0.84	0.36	0.19	1.66	-0.01	-0.33	N/A
POMS							
Tension-Anxiety	13.5	6.9	1	31	0.43	0.45	0.82
Depression	16.4	12.3	0	46	0.65	0.02	0.93
Anger-Hostility	11.3	9.6	0	34	0.75	-0.25	0.93
Vigor	10.6	6.7	0	30	0.88	1.20	0.91
Fatigue	11.9	6.0	2	27	0.45	0.26	0.86
Confusion	11.4	6.1	0	25	0.01	-0.03	0.84
BDI-II	12.7	8.1	2	31	0.67	-0.12	0.81
BFI							
Extraversion	21.2	6.0	10	34	0.24	-0.63	0.80
Agreeableness	28.0	6.2	10	38	-0.78	1.23	0.65
Conscientiousness	26.2	5.9	13	36	-0.32	-0.42	0.76
Neuroticism	26.7	6.0	9	35	-1.29	1.82	0.80
Openness	29.5	7.0	16	41	-0.13	-0.95	0.76

4   Reliability was calculated using Cronbach's alpha. ASMR, autonomous sensory meridian response;

5   BDI, Beck Depression Inventory; BFI, Big Five Inventory; POMS, Profile of Mood States.

1   **Table 2   Correlations between ASMR estimates and individual's states/traits**

	Correlation	
Measure	*r*	*p*
POMS		
Tension-Anxiety	**.521**	**.004**
Depression	.420	.023
Anger-Hostility	.267	.162
Vigor	-.141	.465
Fatigue	.455	.013
Confusion	.416	.025
BDI-II	.281	.140
BFI		
Extraversion	-.101	.595
Agreeableness	.093	.623
Conscientiousness	.135	.476
Neuroticism	.318	.087
Openness	-.266	.155

4   A value in bold font is significant after a false discovery rate (FDR) correction (*N* = 29).