Quantifying the dynamics of pig movements improves targeted disease surveillance and

control plans

Running title: Time-varying networks improve surveillance in pigs

Gustavo Machado[1*], Jason Ardila Galvis[1], Francisco Paulo Nunes Lopes [2], Joana Voges[2],

Antônio Augusto Rosa Medeiros [2], Nicolas Cespedes Cárdenas[3]

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, Raleigh,

North Carolina.

[2] Secretary of Agriculture, Livestock and Agribusiness of State of Rio Grande do Sul (SEAPI-

RS), Porto Alegre, Brazil.

[3]Department of Preventive Veterinary Medicine and Animal Health, School of Veterinary

Medicine and Animal Science, University of São Paulo, São Paulo, Brazil.

Corresponding author: Dr. Gustavo Machado, gmachad@ncsu.edu

Summary

Tracking animal movements over time can fundamentally determine the success of disease control

interventions throughout targeting farms that are tightly connected. In commercial pig production,

animals are transported between farms based on growth stages, thus it generates time-varying

contact networks that will influence the dynamics of disease spread. Still, risk-based surveillance

strategies are mostly based on a static network. In this study, we reconstructed the static and

temporal pig networks of one Brazilian state from 2017 to 2018, comprising 351,519 movements

and 48 million transported pigs. The static networks failed to capture time-respecting movement

pathways. Therefore, we propose a time-dependent network susceptible-infected (SI) model to

simulate the temporal spread of an epidemic over the pig network globally through the temporal

movement of animals among farms, and locally with a stochastic compartmental model in each

farm, configured to calculate the minimum number of target farms needed to achieve effective

disease control. In addition, we propagated disease on the pig temporal network to calculate the

cumulative contacts as a proxy of epidemic sizes and evaluated the impact of network-based

disease control strategies. The results show that targeting the first 1,000 farms ranked by degree

would be sufficient and feasible to diminish disease spread considerably. Our finding also

suggested that assuming a worst-case scenario in which every movement transmit disease,

pursuing farms by degree would limit the transmission to up to 29 farms over the two years period,

which is lower than the number of infected farms for random surveillance, with epidemic sizes of

2,593 farms. The top 1,000 farms could benefit from enhanced biosecurity plans and improved

surveillance, which constitute important next steps in strategizing targeted disease control

interventions. Overall, the proposed modeling framework provides a parsimonious solution for

targeted disease surveillance when temporal movement data is available.

KEYWORDS: swine movement data, dynamic contact networks, disease surveillance, SI

models.

Introduction

Between-farm animal movement has historically been associated with disease spread (Ortiz-

Pelaez et al., 2006; R. P. Smith et al., 2013). Animal movement data has allowed for the initial

development of network-based approaches, which are dominated by the use of static network

metrics to identify vulnerable farms. In a contact network, each farm (node) can play a role in the

dynamics of disease spread, and some can be classified as vulnerable because they periodically

receive infected animals (a high "in" degree); therefore, the identification of these farms is more

important in scenarios of the absence or low prevalence of disease (Qi et al., 2019). Such

locations may serve as sentinels. On the other hand, farms that are more likely to spread disease

throughout their contact network could be targeted, by which restricting or monitoring outgoing

movements could effectively reduce disease spread (Cárdenas et al., 2018; Payen et al., 2019).

Ultimately, the use of network-based metrics to select farms for isolation or surveillance schemes

has shown great value (Mohr et al., 2018; Firestone et al., 2019; Payen et al., 2019). This is a

direct consequence of the availability of animal movement data (Büttner et al., 2016; Firestone et

al., 2019); however, methodologies have not yet evolved as expected. Most published

approaches make use of static networks to describe animal movement and subsequently make

inferences about outbreaks and epidemics (Büttner et al., 2013, 2018; Marquetoux et al., 2016;

Motta et al., 2017; Alarcón et al., 2019; Kinsley et al., 2019). However, the temporal anomalies

and seasonality of movements can only be identified if movement paths are aggregated to

smaller time windows (Cárdenas et al., 2018; Augusta et al., 2019; Sterchi et al., 2019), which

may not be sufficient to make inferences about disease dynamics within the contact networks.

Disease transmission is not restricted to the historical record of movement (Firestone et al., 2019;

Qi et al., 2019; Sterchi et al., 2019); namely, when disease transmission actually occurs by the

movement of infected animals is likely to be an event in time that is dependent on when animals

are sent and, more importantly, when they are received at the final destination, and this rationale

has been largely overlooked (Chaters et al., 2019; Sterchi et al., 2019). Nevertheless, static

networks have been used to measure the effect of targeted farm removal, on the network

fragmentation by monitoring giant strongly or weakly connected components, which would

create small islands restricting the overall connectivity among farms (Motta et al., 2017;

Mekonnen et al., 2019). Other studies have proposed the use of the basic reproduction number

(R_0) and have measured the contribution of node removal in disease transmission (Marquetoux et

al., 2016; Alarcón et al., 2019; Passafaro et al., 2019), still based on static views of full networks.

Therefore, it is recognized that the current neglected temporality of real-world animal

movements has been a constant roadblock to the advancement of network epidemiology,

especially in the development of methodologies (Pellis et al., 2015).

Stochastic simulations have been used extensively to measure the impact of disease control

strategies in future disease spread scenarios (Smith et al., 2017; Funk et al., 2018; Kim et al.,

2019; Lanzas et al., 2020). Mathematical models have also been used to experimentally spread

disease over contact networks and reveal important dynamic characteristics of epidemics (Miller,

2017; P. Kim and Lee, 2018; Darbon et al., 2019). For example, compartmental models can be

used to describe the epidemic spreading in temporal contact networks, which provides an

opportunity for tracing outbreaks and contact among nodes over time (C. Guinat et al., 2016;

Colman et al., 2019; Ferdousi et al., 2019; Sterchi et al., 2019). Here, we used two years of pig

movement data from one Brazilian state to estimate the optimal number of farms for target

surveillance by tracking between-farm contact pathways and inferring epidemic sizes throughout

disease propagation. We first described the complete pig movements and then challenged the use

of static networks against dynamic networks. We proposed a stochastic compartmental model to

simulate disease spread in a two-year empirical network while accounting for the temporal order

of the movements. Our model formulation allowed the calculation of the minimum number of

farms needed to reduce disease spread and approximate the impact of network-based disease

control strategies (e.g., farm isolation, movement restriction, enhanced biosecurity) on the

expected epidemic sizes. Finally, we applied the proposed model to the time-varying pig

network, ultimately to provide a practical background for the identification of targeted farms,

which were then mapped and described in regard to their biosecurity and infrastructure.

Material and methods

Data collection and entry

The record of pig shipments among all registered farms of one Brazilian state was used to

reconstruct contact networks from 2017 until 2018 (SEAPI-RS, 2018). Information about each

swine farm included farm identification, production types (breeding, certified-swine-breeder,

sow, nursery, wean-to-finisher, finisher, subsistence and others), farms infrastructure (e.g.,

presence of a fence, presence of a cleaning and disinfection station, see table 2 for the full list),

geolocation and farms population capacity. Each pig shipment included date(s), farm of origin

and destination, number of animals transported and purpose of movement (weaning, finishing,

slaughter or other). Movement data with missing information of farm location, production type,

farm of origin or destination were excluded prior to the analysis. Additionally, farms declared

inactive (no pigs raised in the past two years or out of business) were not considered in the

analysis. Movements from or to other states were also excluded from any analysis. Based in

information reported to the local authorities in June 2018 all farms were classified as i)

"commercial" farm with active contracts with integrated swine companies, ii) "independent"

farm that declared to be commercial but did not have active contract with integrated swine

companies and small farm holders that, and iii) "not reported" farms that failed to report to local

authorities their pig operation type. The identification of 1,911 (21.4%) not reported farms,

resulted in regional policy in which the state authorities issued a request to all farms report pig

operation status, within one year, farms that fail to report will have all pig movements blocked

until information is shared. Finally, from the total number of active pig farms 11,849, a subset of

farms 9,500 had registered geographic coordinates which included commercial, independent and

not reported business type, therefore were used for the simulation modeling subsections.

Static and monthly networks reconstruction and analysis

Two directed networks were reconstructed. First, a static network with directed "edges"

represented by pig shipments between two "nodes", where nodes corresponded to pig farms. The

static network movements from January 2017 to December 2018 were analyzed by the

traditional methodology in which movements are aggregated into one snapshot prior to the

reconstruction of the networks (Cárdenas et al., 2018). Second, directed networks were

constructed based on a temporal resolution of 30 day windows, in which directed monthly-

networks were generated. (Lentz et al., 2016). For both complete and monthly networks,

centrality metrics were calculated (Table 1) (Cárdenas et al., 2018). As slaughterhouses

constitute sinks and we were interested in potential disease spread among living animals,

transports to slaughter were excluded from the network analysis, but are kept for descriptive

purposes (see figure 2 and supplement table S1).

Temporal variability of the network structure

Static networks are considered an approximation of all movement paths for a defined period,

e.g., animal movement between pig farms for one month (Büttner et al., 2013; Lentz et al.,

2016). This can artificially create accessibility paths between nodes that are not always

accessible by all "nodes" in a network, which can lead to an overestimation of the connectivity

among the nodes (Holme and Saramäki, 2012; Lentz et al., 2016). To quantify the amounts of

such static error, the number of paths in the static view can be compared with the number of

paths in the time-series network (Lentz et al., 2013, 2016). This ratio is known as the causal

fidelity "c", where:

$$c = \frac{p(P_n)}{\mathrm{p}(\mathrm{P_n})} = \frac{Number\ of\ paths\ in\ the\ temporal\ network}{Number\ of\ paths\ in\ the\ static\ network}. \tag{1}$$

The number of paths is the number of nonzero elements in p or P. A causal fidelity of one means

that the temporal network is well represented by its static network, while networks with lower

causal fidelity values would not represent the temporal paths well because most of the paths are

not causal. Here, we calculated the causal fidelity on timescales that are typically used to

investigate animal movement data: monthly, 6 months and 1 year and for the full two years of

data that were used to reconstruct directed networks (Lentz et al., 2016). We also calculated the

edge loyalty (θ), which measures the fraction of preserved edges (θ) of a given premise between

two consecutive years, $t-1$ and t. To quantify $\theta_i^{t-1,t}$, we define Y_i^{t-1} as the set of edges from

the premise i in the time $t-1$ and Y_i^t as the set of edges from the premise i at time t. Then,

$\theta_i^{t-1,t}$ is given by the Jaccard Index in equation 2 (Valdano et al., 2015; Schulz et al., 2017).

$$\theta_i^{t-1,t} = \frac{|Y_i^{t-1} \cap Y_i^t|}{|Y_i^{t-1} \cup Y_i^t|} \tag{2}$$

156　For the node loyalty, we use the same Jaccard Index,

$$node\ loyalty\ _i^{t-1,t} = \frac{|V^{t-1} \cap V^t|}{|V^{t-1} \cup V^t|} \tag{3}$$

where V_i^{t-1} is the set of farms that are active in period $t-1$, and V_i^t is the set of farms that moved

at least one animal at time t (Valdano et al., 2015; Schulz et al., 2017).

The minimum number of farms needed for effective control disease spreading process over

time-varying network

We developed a stochastic susceptible-infectious (SI) epidemic model to calculate the minimum

number of farms needed to effectively reduce the number of new cases expected to spread

through the contact network. The number of infections after the removal of farms raked by

network metrics betweenness, closeness, page rank and degree, random removals (similar to the

current state of farm surveillance), and without farm removal were the main model outputs.

Ultimately, this approximates the impact of network-based target surveillance, which often

include farm visits, sample collection, etc., to a feasible number of farms. This model framework

integrates infection dynamics in each subpopulation as continuous-time Markov chains using the

Gillespie stochastic simulation algorithm (Widgren et al., 2019). Transitions from $S \rightarrow I$ was in a

two-stage model with a one-way transition depending on $\frac{\beta S_i I_i}{S_i + I_i}$ with β as the transmission rate.

The completely stochastic simulation can be summarized as follows: the empirical movement

data from 2017 until 2018 was used to simulate disease spread over a time-varying network;

here, we assume an initial herd prevalence of 10% and 0.1% of infected pigs in each farm, in

which disease was allowed to be transmitted between and within farms. For the within-farm

transmission rate, we use a set of β values of 0.01, 0.05, 0.1 and 0.2, which can also be

considered the product of the contact rates and the transmission probability between animals into

the farms. Another assumption was that the contact rates among pigs were homogenous. For the

between-farm transmission, we considered the observed number of pigs that moved between

farms and the population size at each farm. As an initial condition, model simulation started with

950 infected farms (10% prevalence) at time "0". Here, we proposed two realistic strategies to

seed initial infected nodes. First, pig farms were selected at random from all farms with at least

one movement from 2017 to 2018. For the second strategy, infected farms were chosen

proportionally to the number of farms in commercial, independent, and not reported. This

allowed the initial infection to be seeded within all types of pig operations. The comparison of

both disease introduction approaches allowed us to test a more realistic scenario since disease

can emerge at any location of the study area and in any pig operation (see Figure 1). The

simulated control intervention was based only on the sequential removal of farms from the time

"0" based on the static network metrics described above. While farms were removed, the number

of new cases per 100 susceptible farms was calculated. We chose an arbitrary number of 2,000

herds to be removed, and at the end of each simulation scenario the incidence of new cases I was

calculated as follows:

$$I = \frac{Number\ of\ new\ infected\ herds\ in\ 2017 - 2018}{(Number\ of\ herds\ that\ traded\ with\ other\ herds\ in\ 2017 - 2018) - (Initial\ number\ of\ infected\ herds\ in\ 2017 - 2018\ that\ traded\ with\ other\ herds\ )}$$

Infection was seeded into all candidate farms randomly 100 times, from which simulations were

run and the median incidence was derived for each run. For a sensitivity analysis, we ran

simulations with a diverse set of β parameters (see supplementary figure S3).

Cumulative contacts as a proxy of epidemic sizes

The modeling setup described by (Payen et al., 2019) was adapted to identify the network-based

metric that can produce the smallest epidemic size distribution at the end of the two years of pig

movements, and consequently allow the generation of a farm hot-list to use to target surveillance

activities. This approach is based on the classic deterministic SI spreading model that follows

movements temporality (Dubé et al., 2008; Nöremark et al., 2011; Konschake et al., 2013).

Thereafter, we will refer to this model as "spread cascade", which is a temporal representation of

the cumulative and consecutive contacts among farms, and therefore can be interpreted as the

size of an epidemic (Payen et al., 2019). The hypothetical disease spread followed the

chronological order of movements over the two years. In short, when there was a directed

interaction, including the three dimensions of time t, infected farm i and susceptible farms j,

disease has successfully been transmitted. These events were counted in favor of each farm i

cascade and the triplet t, i, j and added to the temporal links of the stream; see (Payen et al.,

2019) for more details of the methodology. Overall, we recorded the daily spread cascades for

the two years of data, which are represented by weekly averages and $\pm$ 95% CI (see

supplementary figure S4), and the accumulated spread cascades sizes for each seed node (9,500

farms) in the network at the end of the two years as a proxy for the worst-case scenario in

epidemic propagation (Payen et al., 2019). We used the static network to initially rank all pig

farms in descending order of degree, betweenness, PageRank, and closeness and then removed

the top 10, 250, 500, 750 and 1,000 farms before starting the model simulation while recording

the spread cascade sizes. For comparison with usual surveillance practices, we selected the same

number of farms randomly. The contribution on containing the epidemic spread by each

network-based target surveillance strategy was evaluated by comparison of the final spread

cascade distributions at December 2018 using a nonparametric Kruskal-Wallis test followed by

Dunn's post hoc test, where the p-values below 5% were significant. Once the most efficient

network-based surveillance strategy was defined, the biosecurity, infrastructure and population

of the selected farms were described.

Software

The software used for statistical analysis and graphics was R statistical software (versions 3.4.1,

5.1.3) (R Core Team) with R Studio editor using the igraph 1.2.4 (Csardi and Nepusz, 2006),

tidyverse 1.2.1(Wickham, 2017), SimInf 6.3.0 (Widgren et al., 2019), sf 0.5-3 (Pebesma, 2018),

and brazilmaps 0.1.0 (Siqueira, 2019) packages. Spyder 3.3.3 software for Python version 3.7

and Adobe Illustrator CC 2018 software were also used.

Results

Description of pig movements

The total number of active pig farms in the complete database included 11,849 farms, 351,519

movements and more than 48 million pigs transported. The majority, 40.4%, were identified as

commercial farms, 38.2% of independent pig producers, and 21.4% failed to report pig operation

type (see preliminary table S1 to the description of the population in both pig operation classes).

In the commercial farms there were 17 pig production systems with the number of participant

farms varying from 10 to 465. While we have described movements into the 102 registered

slaughterhouses, they were not accounted for in any network analysis thereafter (see Figure 2).

Shipments from commercial to slaughterhouse totaled about 34% of the total movements, while

18% of the movements were from independent farms into slaughterhouses and 14% were

between commercial farms. Importantly movements from independent pig producers and from

farms that did not report type pig operation types into commercial farms were, 3.35% and 1.97%,

respectively. This interaction between non-commercial and commercial farms raises concerns

about the introduction and spread of known and unknown diseases circulation into highly

connected commercial farms (see supplementary table S1). Among all farms with information

about production types 6,555 (55.32%) finisher farms, 1,816 (15.33%) nurseries, 470 (3.97%)

wean-to-finisher farms, 433 (3.65%) breeding farms, 122 (1.03%) certified swine breeder farm

and 250 (2.11%) other farms, which included insemination stations and isolation units, 2,101

(17.73%) as small pig holders (backyard farms or for subsistence).

Figure 1.

More than 51% of movements were from finisher farms to slaughterhouses, with a monthly

median of 7,657 shipments. The second largest number of movements from nurseries to finisher

farms was 13.51%, with a monthly median of 1,992 transfers. Breeding farms made up for

4.65% of the movements to finisher farms and 4.3% of movements to the nurseries, with 673

monthly median movements. Finally, wean-to-finisher farms sent a small but significant amount

of pigs to slaughter (3.5%), while certified swine breeder farms accounted for 2.8% of

movements to slaughter (see Figure 2).

Figure 2.

Monthly network analysis

From the two years of movement data, we reconstructed the monthly static networks. The

number of farms, movements and animals varied substantially with a steady increase from July

2017 until October of the same year (see Figure 3). The network diameter revealed a positive

increasing trend of 33% from April 2018 through December 2018, December 2018, which

decreased thereafter; this pattern was different from the monthly values from the previous year.

The monthly analysis showed a positive trend of the GSCC in 2017 and a negative trend in 2018,

and the GWCC had a more stable pattern with two consecutive peaks in July and October of

2018. In May 2017 GSCC peaked at 40 farms inflated with mostly commercial farms, and in

November 2018 the GWCC had almost 600 farms in which most were also commercial (see

supplementary Figure S1 and Table S2 for the two years GSCC and GWCC).

Figure 3.

Loyalty and causal fidelity

The node loyalty of 2018 compared with 2017 was 0.65, while edge loyalty was 0.19. This

means that only 65% of the farms that traded pigs in 2017 were preserved in 2018, and 19% of

the edges were preserved between 2017 and 2018. On the other hand, the monthly causal fidelity

varied from 0.33 in July 2018 and reached the highest value of 0.55 in July 2017 (see

supplementary Figure S2). For the network that evaluated time periods of six months, the causal

fidelity was 0.22 for the first half of 2017 and 0.47 for the second half of 2018, and the annual

evaluation was in 2017, 0.18 in 2018, and 0.33 for the entire static network. The causal fidelity

demonstrated a discrepancy between static and temporal network views. The concept of time

does not per se exist in these static networks; thus, the use of traversal network models here

would reduce the accuracy toward overestimating disease outbreaks, and likely mislead the

selection of farms to undergo network-based surveillance (Lentz et al., 2016). In addition, we

calculated the causal error from the causal fidelity, which can be approximated by the inverse of

"c" and represents the overall overestimation of disease outbreaks if static networks are used.

The first six months of 2017 had the worst results, with overestimations at 4.54 and 1.81 for the

July 2018 network.

Number of farms for targeted surveillance and epidemic sizes

The stochastic runs showed that the removal of 2,000 farms did not completely control the

disease from spreading, but it was reduced markedly to a maximum of 1.95 new cases per 100

farms when β of 0.2 was used (the worst-case scenario). Since this study was designed to guide

decisions in the field, we consulted with the local stakeholders before deciding on the final

number of farms to be on the surveillance hot-list (personal communications with Dr. Lopes,

Chair of the Animal Movement and Surveillance Department of the state of Rio Grande do Sul).

Based on local capacities and the average number of new cases was reduced to as low as 7.88

cases per 100 farms with a β of 0.2 when 1,000 farms were removed from the network, the

former was set to be the maximum number of feasible farms to be targeted. Furthermore, we

compared two infectious seed scenarios: a) infections started at random farms and b) stratified

random infections forced to start in proportion to the number of commercial, independent, and

not reported. Both have very similar results which reinforced the importance of independent

farms in the role of spreading diseases into commercial pig farms and vice versa (Figure 4, also

see supplementary figure S3)

Figure 4.

After the removal of the first 1,000 farms with the highest degree, the deterministic model

estimated that the maximum outbreak size would involve 29 farms (Figure 5). As expected,

random removals resulted in a much larger epidemic size, reaching up to 2,593 farms (Figure 5,

also see supplementary table S4 and figure S5).

Figure 5.

Descriptive analysis of biosecurity, infrastructure and population profile of the targeted

farms.

Most of the farms within the 1,000 target farms were commercial (55.6%), see Figure 6, with one

production system contribution with 138 farms, and 29.7% were independent and 14.7% did not

report business operation From the comprehensive list of biosecurity and infrastructure

characteristics of the commercial farms, farms had a median of 2 barns (IQR-1-11), capacities

for housing were approximately 1,100 pigs (IQR=162-9,925), the median number of piglets was

1,122 (IQR-16-7,229), breeding farms had 4 (IQR=1-18) boars, and the gilts population was 544

(IQR-11-3,044). More details about the farms' biosecurity and infrastructures can be found in

Table 2.

Figure 6. The geographic locations of the top 1,000 farms ranked by logarithm of degree.

Discussion

The majority of the movements were from commercial and independent pig farms to

slaughterhouses, but we also demonstrated the presence of a contact network between

independent and farms that did not report pig operation types with commercial enterprises. The

later finding is of concern and a key issue in the face of the introduction and spread of foreign

animal diseases (FADs) into highly connected commercial pig systems. We also demonstrated

that the use of static networks to identify tightly connected farms to be selected for surveillance

activities and to estimate disease spread dynamics would likely mislead disease control efforts

and overestimated epidemic sizes. Thus, our model formulation accounted for the temporal order

of the between-farm movements without overestimating the epidemic sizes while accounting for

impact on disease spread as network-based interventions were implemented, our model also

predicted the impact of network-based target disease control strategies (e.g., farm isolation,

movement restriction, enhancement of biosecurity) on the accumulated number of cases over the

next two years (2017 and 2018). These clearly demonstrate the advantages of temporal networks

to identify key farms to compose disease control contingency plans at national, local and at

production system level for vertical integrated animal production (i.e., swine, poultry), in order

to quickly mitigate an introduction and subsequent spread of infectious diseases. The generation

of a hot-list with farms to undergo complete lockdown or to which additional biosecurity can be

rapidly applied in the face of foreign animal disease and other biological threats to national

agriculture and food supply is a critical contribution of this study.

The frequency of shipments from independent to commercial pig farms reveal the importance of

analyzing the interaction between pig populations of diverse biosecurity levels.

These interactions pose significant risk for the introduction and spread of infectious diseases into

highly connected integrated commercial pig farms. Many studies showed that ASF spread in

China and parts of the EU have been related to pig movements between non-commercial and

commercial pig operations (Claire Guinat et al., 2016; Chenais et al., 2019; Miteva et al., 2020).

The description of interaction between non-commercial and commercial farms remains rare in

the literature (Porphyre et al., 2014). However, a study in Scotland described as much as 8.6% of

movement going from small producers into commercial pig farms (Porphyre et al., 2014),

meanwhile studies in North America have mostly lack the description of such movements

(Kinsley et al., 2019; Sterchi et al., 2019), but for a recent a study in Iowa that described the

presence of such pig flows but did not considered it their network analysis (Passafaro et al.,

2019). Indeed, there is a need to capture movements between non-commercial and commercial

pig farms and vice-versa, which also implies including those into the subsequent network and

modeling analysis. One important limitation of this study is the fact that 21.4 % of the farms did

provided information of the type of pig operation. Briefly, we evaluated the population in this

category, the median number of pig was 415 (IQR 1-1441) with median number of breeding

female of 72 (IQR 1- 483), and median of 1 boar (IQR 0-4), see supplementary table S3 for more

details. By the number of total pigs and breeding females these farms are likely smaller operation

which may include farms with limited biosecurity. We highlight the need for a follow-up study

to collect more information about these farms, including the type of pig operation, farm's

infrastructure and information about biosecurity. Indeed, gathering and analyzing non-

commercial and commercial interactions becomes more important as the pig industry around the

world prepares for the introduction of ASF, which will require tracking animal movements

among all pig farms (Brown and Bevins, 2018; Jurado et al., 2018).

In this study population the static networks failed to capture time-respecting movement

pathways, and consequently deceive disease control efforts. Until now a large amount of studies

have considered static network views to make inferences about disease spread or to calculate

outbreak sizes (Büttner et al., 2013; Lee et al., 2017; Salines et al., 2017). Mostly those studies

ignored the causal fidelity of the contacts, which can be used to quantify the error of the static

representation of a temporal network (Lentz et al., 2013). Thus, without such verification, it is

difficult to decide if and at which time window inferences or metrics from static networks can be

interpreted or used to summarize networks (Lentz et al., 2013; Büttner et al., 2016). It is

dangerous to assume that aggregating contact movements into one snapshot would not

overestimate or introduce edge biases (Vidondo and Voelkl, 2018). It has been suggested that to

some extent, static networks provide useful information about disease related risks (Mohr et al.,

2018). However, most static network studies lacked basic statistical support for their decisions

for data aggregation (Büttner et al., 2013; Lee et al., 2017; Salines et al., 2017), some studies

have rather mention that other studies also used similar time windows, which would support their

decisions to data aggregation (Büttner et al., 2013; Lee et al., 2017; Salines et al., 2017). Finally,

contact network aggregation has been a practice, even when the temporal sequences of the

movements were available (Amaku et al., 2015; Cárdenas et al., 2018). Thus, the use of static

networks tends to artificially connect a larger number of farms, which will consequently bias

surveillance strategies to less relevant farms and compromise resource allocation, therefore when

temporal movement data is available, it is strongly recommended consider such dynamic while

the interest is to study the spread of diseases over time-varying networks.

The number of active pig farms was seasonal and skewed with variation of plus or minus 300

farms over the two-year period, a similar trend was observed in the volume of pigs traded. In

contrast to our observations, the number of active pig farms varied largely compared with other

pig populations in Germany (Büttner et al., 2016), France (Rautureau et al., 2012) and Canada

(Thakur et al., 2016), even though it is difficult to compare distinct networks the comparison

shown diversity especially in the volume of movements (Miele et al., 2019). Some factors that

have contributed to the variations in our study may be associated with hog prices, which tends to

be more volatile than in other countries. In addition, the monthly network diameter almost

doubled in the middle of 2018, which seems to be a highly abnormal pattern in the network. A

possible explanation for this may be related to the global configuration of the network because

the nodes were connected in a way that increased the length of the shortest paths between them,

which was accompanied by an increase in the GWCC. Consequently, during this period, we

observed an increase in the number of nodes and connections; therefore, a given node would

require more steps to reach another farm, within the network which could alone reduce the speed

of disease transmission (Vidondo and Voelkl, 2018). While the GWCC increased over the years,

the GSCC suffered from relevant variations throughout the months, especially in the second half

of 2018, where new paths connected the GSCC with other minor components and consequently

increased the size of the GWCC, which could have implications for disease transmission and

persistence (Schulz et al., 2017; Marquetoux et al., 2016). The sizes of GSCC and GWCC

variation were mostly attributed to variation in the number of commercial farms activities. Farms

that did not report operation types represented the minority of the farms in both components,

especially in April 2018 (see figure S1).

Our results show that different network-based intervention schemes successfully reduced the

spread of disease independently from the local β transmission probability rates at the farm. We

show that targeting farms by degree was the most effective intervention in reducing the number

of new cases, followed by targeting farms by betweenness, which required ~ 1,300 farms to be

quarantined. Finally, as anticipated, random removal had little impact in controlling disease. A

recent study explored the contribution of existing trade duration on disease transmission

probabilities and the impact on epidemic sizes, suggesting that epidemic sizes were sensitive to

network activity (Farine and Whitehead, 2015; Lebl et al., 2016). Indeed, we explicitly

considered node activity while disease spread was simulated. Two other studies attempted to

account for time-dependent disease spread while aiming to reduce R_0. One study used a non-

compartmental model approach and found that the number of farms that needed to be removed

by degree approached 50% of all farms (Kinsley et al., 2019). The second study used a

mechanistic approach in which ASF spread was simulated and found that degree ranking was the

best strategy to contain the disease (Ferdousi et al., 2019). In summary, degree was shown to be

the best strategy in reducing disease burden, while betweenness had the second best

performance. It is worth noting that betweenness is a useful metric when studying disease

diffusion (Farine and Whitehead, 2015), mainly because it is a representation of the shortest path

between nodes. Here independent farms with high betweenness may bridge several commercial

firms; therefore, more details about these unexplored pathways should be investigated in the

future.

Based on the identified minimum number of farms of 1,000 that needed to be removed from the

contact network, we simulated outbreak propagation with a deterministic SI model and identified

the most efficient network-based metric capable of reducing the expected size of spread

cascades. We showed that degree was the most significant intervention, with no more than 29 pig

farms involved per outbreak (Figure 6). In addition, our results demonstrated that targeting more

than 500 farms by degree was efficient in reducing epidemic sizes. Random surveillance was

unexpectedly better than clossenes and PageRank for most of the two years of propagation. One

possible explanation may be that this network is not well connected, and farms that remain to be

removed may be too similar with respect to their clossennes and PageRank values; therefore, the

impact on the propagation is expected to be the one resulting from random removals.

Our results have shown an uneven proportion of commercial and independent pig farms within

the first 1,000 farms with the highest degree. This illustrates the interplay between the two

populations, where infection can arrive at either and propagate in both directions. Here, most

farms of interest lacked important biosecurity features, such as cleaning and disinfection stations

(Dee et al., 2006). A study in Argentina also highlighted the lack of biosecurity on highly

connected farms (Alarcón et al., 2019). On the other hand, in our study, most farms did not

utilize rendering, but preferred to compost dead animals. The former was previously described as

an important risk factor for the spread of porcine reproductive and respiratory syndrome (PRRS)

in the US (Velasova et al., 2012; Silva et al., 2019). In the same study, composting was found to

be the less risky practice when removing cull or dead pigs (Silva et al., 2019). Another relevant

biosecurity point that has been described as a risk for disease introduction is the number of farm

entries; here, we noticed that the majority of the farms had only one main road, which potentially

represents less risk for local disease transmission (Silva et al., 2019). A disproportionate number

of these highly risky farms were nurseries and breeding farms, which has often been proposed as

a desired group to be targeted for surveillance (Dorjee et al., 2013). In addition to the

identification of priority farms, the identification of relevant biosecurity gaps at those farms is

invaluable, especially under the current preparedness for the introduction of foreign animal

diseases such as ASF. One of the best options is the implementation of enhanced biosecurity

plans, such the Secure Pork Supply (SPS) Plan, designed to provide business continuity in the

event of a foreign animal disease outbreak as well as help protect operations from endemic

diseases (Pudenz et al., 2019).

Finally, more comprehensive studies coupling animal movements with local and environmental

routes of transmission while accounting for farms biosecurity and infrastructure are needed to

further explore the mechanisms of pig disease propagation, for which a complete multiscale

mechanistic model is necessary (Qi et al., 2019).

488
Considerations

The use of static networks to target key farms to control the disease from spreading should be

interpreted with caution due to several limitations associated with animal movement aggregation,

as shown in this study and elsewhere (Lentz et al., 2016). An obvious issue here is to assume that

the same pair of farms would trade with each other throughout the entire study periods.

Therefore, we argue that the causal fidelity of the temporal network should be presented before

considering the use of the static networks in the development of disease surveillance or control

strategies (Lentz et al., 2016; Payen et al., 2019). Most of recent studies did not account for this

issue (Büttner et al., 2013, 2018; Marquetoux et al., 2016; Motta et al., 2017; Alarcón et al.,

2019), or considered the temporal network pathway accessibility of each farm.

In this study, we did not examine feed-truck or other vehicle movements or sharing. Such

information is absent from the majority of the current literature (Lentz et al., 2016), but it has

been included in some studies (Augusta et al., 2019; Sterchi et al., 2019). It is critical to

acknowledge that the movement of feed trucks and short-distance transportation of breeding

replacement animals or culled sows contributed to disease transmission (Silva et al., 2019).

Indeed, before these additional contacts are not taken into consideration, our understanding of

disease spread pathways will likely continue to be limited.

Finally, the modeling framework proposed here can be broadly used to calculate the minimum

number of farms to be targeted for the control of disease spread when temporal contact data are

available and to determine the network metrics that should be preferred for selection of target

farms according to the approximation of epidemic sizes. In addition, the proposed framework

could be used as a decision tool for both local official veterinary services and the private sector

for evaluating animal movements and developing a more risk-oriented surveillance strategies. At

production system level, it allows for the creation of a hot-list with farms to which the

reinforcement of biosecurity would be key in the face of an outbreak. The model also provides

an estimation of the amount of personnel and material resources that would need to be allocated

to control the epidemic, which is key for preparedness for future events (Casal et al., 2019).

Future studies are also needed to explore the contribution of movements to slaughter houses,

especially when pig are redirected to other slaughter houses (i.e., slaughter not able to slaughter

all animals) or send back to the farm of origin, this is very common in the US, however little is

known relevance in spreading disease back to the farm, however there is some early discussion

about it (Russell et al., 2020).

Conclusion

The contact among farms of distinct biosecurity by pig movements identify a potential gap that

could result in the introduction and dissemination of FAD into commercial and highly connected

pig populations. One-third of all pig farms traded with the same partners in the course of two

years; consequently, the static networks did not offer reasonable information about the realized

temporal movement pathways. In this scenario, the use of static networks to target vulnerable or

high-risk farms by network metrics would likely overestimate the role of those farms in disease

transmission and limit the accuracy needed to optimize surveillance. Our proposed modeling

framework provides a parsimonious solution to this problem; it handles the temporal order of

movements, calculates the minimum number of farms needed to be targeted and approximates

the expected epidemic size in a worst-case scenario where each movement can effectively

transmit disease. The results from the application of this approach identified priority pig farms

that had remarkably limited biosecurity. This fact imposes important vulnerability to pathogen

introduction and spread; therefore, those farms could benefit from the implementation of

enhanced biosecurity plans, such as what has been promoted and implemented in the US by SPS

plan. Finally, the introduction of risk-based approaches may optimize the costs of routine

surveillance and can be used in preparation for the introduction of FAD into the local pig

industry.

Acknowledgements

544 This work was supported by the Fundo de Desenvolvimento e Defesa Sanitária Animal

545 (FUNDESA-RS) and by the NC State University, College of Veterinary Medicine, Global Health

546 funds.

Authors' contributions

GM, FPNL, JV and AARM coordinated movement data collection. GM conceived paper ideas.

GM, JAG and NCC participated in the design of the study. NCC and GM conducted data

processing and cleaning. GM, JAG and NCC wrote and modified computer algorithms and

conducted analysis. GM, JAG and NCC developed simulation models. GM and NCC wrote the

manuscript and JA edited the manuscript. All authors discussed results and commented on the

manuscript.

Conflict of interest

557 All authors confirm that there are no conflicts of interest to declare

Ethical statement

560 The authors confirm that the ethical policies of the journal, as noted on the journal's author

561 guidelines page. Since this work did not involve animal sampling neither questioner data

562 collection there was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are available from the regional veterinary office.

Restrictions apply to the availability of these data, which were used under confidentiality

agreements.

568
References

Alarcón, L. V, P. CIPRIOTTI, M. Monterubbianessi, C. Perfumo, E. Mateu, and A. Allepuz,

2019: Network analysis of pig movements in Argentina: identification of key farms in the

spread of diseases and relationship with their biosecurity level. *Front. Vet. Sci.*DOI:

10.3389/conf.fvets.2019.05.00006.

Augusta, C., G.W. Taylor, and R. Deardon, 2019: Dynamic contact networks of swine

movement in Manitoba, Canada: Characterization and implications for infectious disease

spread. *Transbound. Emerg. Dis.*DOI: 10.1111/tbed.13220.

Brin, S., and L. Page, 1998: The anatomy of a large scale hypertextual Web search engine

published version. *Comput. Networks ISDN Syst.*DOI: 10.1.1.109.4049.

Brown, V.R., and S.N. Bevins, 2018: A review of African swine fever and the potential for

introduction into the United States and the possibility of subsequent establishment in feral

swine and native ticks. *Front. Vet. Sci.*DOI: 10.3389/fvets.2018.00011.

Büttner, K., J. Krieter, A. Traulsen, and I. Traulsen, 2013: Static network analysis of a pork

supply chain in Northern Germany-Characterisation of the potential spread of infectious

diseases via animal movements. *Prev. Vet. Med.*DOI: 10.1016/j.prevetmed.2013.01.008.

Büttner, K., J. Salau, and J. Krieter, 2016: Quality assessment of static aggregation compared to

the temporal approach based on a pig trade network in Northern Germany. *Prev. Vet.*

*Med.*DOI: 10.1016/j.prevetmed.2016.05.005.

Büttner, K., J. Salau, and J. Krieter, 2018: Effects of data quality in an animal trade network and

their impact on centrality parameters. *Soc. Networks*DOI: 10.1016/j.socnet.2018.01.001.

Cárdenas, N.C., J.O.A. Galvis, A.A. Farinati, J.H.H. Grisi-Filho, G.N. Diehl, and G. Machado,

2018: Burkholderia mallei: the dynamics of networks and disease transmission.

Transbound. Emerg. Dis. **0**, DOI: 10.1111/tbed.13071.

Casal, J., C. Saegerman, S. Bertagnoli, G. Meyer, J.P. Ganière, P. Caufour, K. De Clercq, P.

Jacquiet, C. Hautefeuille, F. Etore, and S. Napp, 2019: A simple method to estimate the

number of doses to include in a bank of vaccines. The case of Lumpy Skin Disease in

France. *PLoS One*DOI: 10.1371/journal.pone.0210317.

Chaters, G.L., P.C.D. Johnson, S. Cleaveland, J. Crispell, W.A. De Glanville, T. Doherty, L.

Matthews, S. Mohr, O.M. Nyasebwa, G. Rossi, L.C.M. Salvador, E. Swai, and R.R. Kao,

2019: Analysing livestock network data for infectious disease control: An argument for

routine data collection in emerging economies. *Philos. Trans. R. Soc. B Biol. Sci.*DOI:

10.1098/rstb.2018.0264.

Chenais, E., K. Depner, V. Guberti, K. Dietze, A. Viltrop, and K. Ståhl, 2019: Epidemiological

considerations on African swine fever in Europe 2014-2018. *Porc. Heal. Manag.*DOI:

10.1186/s40813-018-0109-2.

Colman, E., P. Holme, H. Sayama, and C. Gershenson, 2019: Efficient sentinel surveillance

strategies for preventing epidemics on networks. (Daniela Paolotti, Ed.)*PLOS Comput. Biol.*

15, e1007517, DOI: 10.1371/journal.pcbi.1007517.

Csardi, G., and T. Nepusz, 2006: The igraph software package for complex network research.

InterJournal Complex Syst.

Darbon, A., D. Colombi, E. Valdano, L. Savini, A. Giovannini, and V. Colizza, 2019: Disease

persistence on temporal contact networks accounting for heterogeneous infectious periods.

*R. Soc. Open Sci.*DOI: 10.1098/rsos.181404.

Dee, S.A., J. Deen, and C. Pijoan, 2006: Evaluation of an industry-based sanitation protocol for

full-size transport vehicles contaminated with porcine reproductive and respiratory

syndrome virus. *J. Swine Heal. Prod.*

Dorjee, S., C.W. Revie, Z. Poljak, W.B. McNab, and J. Sanchez, 2013: Network analysis of

swine shipments in Ontario, Canada, to support disease spread modelling and risk-based

disease management. *Prev. Vet. Med.*DOI: 10.1016/j.prevetmed.2013.06.008.

Dubé, C., C. Ribble, D. Kelton, and B. McNab, 2008: Comparing network analysis measures to

determine potential epidemic size of highly contagious exotic diseases in fragmented

monthly networks of dairy cattle movements in Ontario, Canada. *Transbound. Emerg.*

*Dis.*DOI: 10.1111/j.1865-1682.2008.01053.x.

Farine, D.R., and H. Whitehead, 2015: Constructing, conducting and interpreting animal social

network analysis. *J. Anim. Ecol.*DOI: 10.1111/1365-2656.12418.

Ferdousi, T., S.A. Moon, A. Self, and C. Scoglio, 2019: Generation of swine movement network

and analysis of efficient mitigation strategies for African swine fever virus. DOI:

10.1371/journal.pone.0225785.

Firestone, S.M., Y. Hayama, R. Bradhurst, T. Yamamoto, T. Tsutsui, and M.A. Stevenson, 2019:

Reconstructing foot-and-mouth disease outbreaks: a methods comparison of transmission

network models. *Sci. Rep.*DOI: 10.1038/s41598-019-41103-6.

Freeman, L.C., 1978: Centrality in social networks conceptual clarification. *Soc. Networks*DOI:

10.1016/0378-8733(78)90021-7.

Funk, S., A. Camacho, A.J. Kucharski, R.M. Eggo, and W.J. Edmunds, 2018: Real-time

forecasting of infectious disease dynamics with a stochastic semi-mechanistic model.

*Epidemics*DOI: 10.1016/j.epidem.2016.11.003.

Guinat, C., A. Relun, B. Wall, A. Morris, L. Dixon, and D.U. Pfeiffer, 2016: Exploring pig trade

patterns to inform the design of risk-based disease surveillance and control strategies. *Sci.*

*Rep.*DOI: 10.1038/srep28429.

Guinat, Claire, A. Gogin, S. Blome, G. Keil, R. Pollin, D.U. Pfeiffer, and L. Dixon, 2016:

Transmission routes of African swine fever virus to domestic pigs: Current knowledge and

future research directions. *Vet. Rec.*DOI: 10.1136/vr.103593.

Holme, P., and J. Saramäki, 2012: Temporal networks. *Phys. Rep.*DOI:

10.1016/j.physrep.2012.03.001.

Jurado, C., M. Martínez-Avilés, A.D. La Torre, M. Štukelj, H.C. de C. Ferreira, M. Cerioli, J.M.

Sánchez-Vizcaíno, and S. Bellini, 2018: Relevant measures to prevent the spread of African

Swine Fever in the European Union Domestic Pig Sector. *Front. Vet. Sci.*DOI:

10.3389/fvets.2018.00077.

Kim, M., D. Paini, and R. Jurdak, 2019: Modeling stochastic processes in disease spread across a

heterogeneous social system. *Proc. Natl. Acad. Sci. U. S. A.*DOI:

10.1073/pnas.1801429116.

Kim, P., and C.H. Lee, 2018: Epidemic Spreading in Complex Networks with Resilient Nodes:

Applications to FMD. *Complexity*DOI: 10.1155/2018/5024327.

Kinsley, A.C., A.M. Perez, M.E. Craft, and K.L. Vanderwaal, 2019: Characterization of swine

movements in the United States and implications for disease control. *Prev. Vet. Med.*DOI:

10.1016/j.prevetmed.2019.01.001.

Konschake, M., H.H.K. Lentz, F.J. Conraths, P. Hövel, and T. Selhorst, 2013: On the Robustness

of In- and Out-Components in a Temporal Network. *PLoS One*DOI:

10.1371/journal.pone.0055223.

Lanzas, C., K. Davies, S. Erwin, and D. Dawson, 2020: On modelling environmentally

transmitted pathogens. *Interface Focus* **10**, 20190056, DOI: 10.1098/rsfs.2019.0056.

Lebl, K., H.H.K. Lentz, B. Pinior, and T. Selhorst, 2016: Impact of network activity on the

spread of infectious diseases through the german pig trade network. *Front. Vet. Sci.*DOI:

10.3389/fvets.2016.00048.

Lentz, H.H.K., A. Koher, P. Hövel, J. Gethmann, C. Sauter-Louis, T. Selhorst, and F.J. Conraths,

2016: Disease spread through animal movements: A static and temporal network analysis of

pig trade in Germany. *PLoS One*DOI: 10.1371/journal.pone.0155196.

Lentz, H.H.K., T. Selhorst, and I.M. Sokolov, 2013: Unfolding Accessibility Provides a

Macroscopic Approach to Temporal Networks. *Phys. Rev. Lett.*DOI:

10.1103/PhysRevLett.110.118701.

Marquetoux, N., M.A. Stevenson, P. Wilson, A. Ridler, and C. Heuer, 2016: Using social

network analysis to inform disease control interventions. *Prev. Vet. Med.*DOI:

10.1016/j.prevetmed.2016.01.022.

Mekonnen, G.A., G. Ameni, J.L.N. Wood, S. Berg, and A.J.K. Conlan, 2019: Network analysis

of dairy cattle movement and associations with bovine tuberculosis spread and control in

emerging dairy belts of Ethiopia. *BMC Vet. Res.*DOI: 10.1186/s12917-019-1962-1.

Miele, V., C. Matias, S. Robin, and S. Dray, 2019: Nine quick tips for analyzing network data.

(Francis Ouellette, Ed.)*PLOS Comput. Biol.* **15**, e1007434, DOI:

10.1371/journal.pcbi.1007434.

Miller, J.C., 2017: Mathematical models of SIR disease spread with sexual-network and other

transmission mechanisms. *Infect. Dis. Model.*DOI: 10.1016/j.idm.2016.12.003.

Miteva, A., A. Papanikolaou, A. Gogin, A. Boklund, A. Bøtner, A. Linden, A. Viltrop, C.G.

Schmidt, C. Ivanciu, D. Desmecht, D. Korytarova, E. Olsevskis, G. Helyes, G.

Wozniakowski, H.H. Thulke, H. Roberts, J.C. Abrahantes, K. Ståhl, K. Depner, L.C.

González Villeta, M. Spiridon, S. Ostojic, S. More, T.C. Vasile, V. Grigaliuniene, V.

Guberti, and R. Wallo, 2020: Epidemiological analyses of African swine fever in the

European Union (November 2018 to October 2019). *EFSA J.*DOI:

10.2903/j.efsa.2020.5996.

Mohr, S., M. Deason, M. Churakov, T. Doherty, and R.R. Kao, 2018: Manipulation of contact

network structure and the impact on foot-and-mouth disease transmission. *Prev. Vet.*

*Med.*DOI: 10.1016/j.prevetmed.2018.05.006.

Motta, P., T. Porphyre, I. Handel, S.M. Hamman, V. Ngu Ngwa, V. Tanya, K. Morgan, R.

Christley, and B.M.D. Bronsvoort, 2017: Implications of the cattle trade network in

Cameroon for regional disease prevention and control. *Sci. Rep.*DOI: 10.1038/srep43932.

Nöremark, M., N. Håkansson, S.S. Lewerin, A. Lindberg, and A. Jonsson, 2011: Network

analysis of cattle and pig movements in Sweden: Measures relevant for disease control and

risk based surveillance. *Prev. Vet. Med.*DOI: 10.1016/j.prevetmed.2010.12.009.

Ortiz-Pelaez, A., D.U. Pfeiffer, R.J. Soares-Magalhães, and F.J. Guitian, 2006: Use of social

network analysis to characterize the pattern of animal movements in the initial phases of the

2001 foot and mouth disease (FMD) epidemic in the UK. *Prev. Vet. Med.*DOI:

10.1016/j.prevetmed.2006.04.007.

Passafaro, T.L., A.F.A. Fernandes, B.D. Valente, N.H. Williams, and G.J.M. Rosa, 2019:

Network analysis of swine movements in a multi-site pig production system in Iowa, USA.

*Prev. Vet. Med.*DOI: 10.1016/J.PREVETMED.2019.104856.

Payen, A., L. Tabourier, and M. Latapy, 2019: Spreading dynamics in a cattle trade network:

Size, speed, typical profile and consequences on epidemic control strategies. *PLoS One*DOI:

10.1371/journal.pone.0217972.

Pebesma, E., 2018: Sf: Simple Features for R. R package version 0.5-3. .

Pellis, L., F. Ball, S. Bansal, K. Eames, T. House, V. Isham, and P. Trapman, 2015: Eight

challenges for network epidemic models. *Epidemics*DOI: 10.1016/j.epidem.2014.07.003.

Porphyre, T., L.A. Boden, C. Correia-Gomes, H.K. Auty, G.J. Gunn, and M.E.J. Woolhouse,

2014: How commercial and non-commercial swine producers move pigs in Scotland: A

detailed descriptive analysis. *BMC Vet. Res.*DOI: 10.1186/1746-6148-10-140.

Pudenz, C.C., L.L. Schulz, and G.T. Tonsor, 2019: Adoption of secure pork supply plan

biosecurity by U.S. Swine producers. *Front. Vet. Sci.*DOI: 10.3389/fvets.2019.00146.

Qi, L., G. Beaunée, S. Arnoux, B.L. Dutta, A. Joly, E. Vergu, and P. Ezanno, 2019:

Neighbourhood contacts and trade movements drive the regional spread of bovine viral

diarrhoea virus (BVDV). *Vet. Res.*DOI: 10.1186/s13567-019-0647-x.

Rautureau, S., B. Dufour, and B. Durand, 2012: Structural vulnerability of the French swine

industry trade network to the spread of infectious diseases. *Animal*DOI:

10.1017/S1751731111002631.

Russell, L.E., J. Polo, and D. Meeker, 2020 (16. February): The Canadian 2014 porcine epidemic

diarrhoea virus outbreak: Important risk factors that were not considered in the

epidemiological investigation could change the conclusions. *Transbound. Emerg.*

*Dis.*tbed.13496, DOI: 10.1111/tbed.13496. Blackwell Publishing Ltd.

Schulz, J., A. Boklund, T.H.B. Halasa, N. Toft, and H.H.K. Lentz, 2017: Network analysis of pig

movements: Loyalty patterns and contact chains of different holding types in Denmark.

*PLoS One*DOI: 10.1371/journal.pone.0179915.

SEAPI-RS, 2018: SEAPI, Secretaria da Agricultura, Pecuária e arrigação [Online] Available at

http://www.agricultura.rs.gov.br/inicial.

Silva, G.S., G. Machado, K.L. Baker, D.J. Holtkamp, and D.C.L. Linhares, 2019: Machine-

learning algorithms to identify key biosecurity practices and factors associated with

breeding herds reporting PRRS outbreak. *Prev. Vet. Med.*DOI:

10.1016/j.prevetmed.2019.104749.

Siqueira, P., 2019: brazilmaps: Brazilian Maps from Different Geographic Levels. .

Smith, R.P., A.J.C. Cook, and R.M. Christley, 2013: Descriptive and social network analysis of

pig transport data recorded by quality assured pig farms in the UK. *Prev. Vet. Med.* **108**,

167–177, DOI: 10.1016/j.prevetmed.2012.08.011.

Smith, T.A., N. Chitnis, M. Penny, and M. Tanner, 2017: Malaria Modeling in the Era of

Eradication. *Cold Spring Harb. Perspect. Med.* **7**, a025460, DOI:

10.1101/cshperspect.a025460.

Sterchi, M., C. Faverjon, C. Sarasua, M.E. Vargas, J. Berezowski, A. Bernstein, R. Grütter, and

H. Nathues, 2019: The pig transport network in Switzerland: Structure, patterns, and

implications for the transmission of infectious diseases between animal holdings. *PLoS*

*One*DOI: 10.1371/journal.pone.0217974.

Thakur, K.K., C.W. Revie, D. Hurnik, Z. Poljak, and J. Sanchez, 2016: Analysis of Swine

Movement in Four Canadian Regions: Network Structure and Implications for Disease

Spread. *Transbound. Emerg. Dis.*DOI: 10.1111/tbed.12225.

Valdano, E., C. Poletto, A. Giovannini, D. Palma, L. Savini, and V. Colizza, 2015: Predicting

Epidemic Risk from Past Temporal Contact Data. *PLoS Comput. Biol.*DOI:

10.1371/journal.pcbi.1004152.

Velasova, M., P. Alarcon, S. Williamson, and B. Wieland, 2012: Risk factors for porcine

reproductive and respiratory syndrome virus infection and resulting challenges for effective

disease surveillance. *BMC Vet. Res.*DOI: 10.1186/1746-6148-8-184.

Vidondo, B., and B. Voelkl, 2018: Dynamic network measures reveal the impact of cattle

markets and alpine summering on the risk of epidemic outbreaks in the Swiss cattle

population. *BMC Vet. Res.*DOI: 10.1186/s12917-018-1406-3.

Wasserman, S., and K. Faust, 1994: Social Network Analysis: Methods and Applications :

Cambridge University Press.

Wickham, H., 2017: Tidyverse: Easily install and load "tidyverse" packages. .

Widgren, S., P. Bauer, R. Eriksson, and S. Engblom, 2019: Siminf: An R package for data-driven

stochastic disease spread simulations. *J. Stat. Softw.*DOI: 10.18637/jss.v091.i12.

Table 1. Description of network terminology and definition applied to pig movements.

Parameter	Definition	References
Nodes	A pig farms.	(Wasserman and Faust, 1994)
Edge	A link between two nodes.	Wasserman and Faust (1994)
Degree (k)	The number of unique contacts to and from a farm. When direction was taken into account, the ingoing and outgoing contacts were separated: the "out" degree is the number of contacts originating from a specific farm, and the "in" degree is the number of contacts directed toward a specific premise.	(Wasserman and Faust, 1994)
PageRank	Measure of the Google PageRank, a link analysis algorithm that produces a ranking importance of all the farms from 0 to 1.	(Brin and Page, 1998)
Closeness centrality	Closeness centrality measures how many steps are required to access every other vertex from a given node. This measure can be directed as incoming steps or outgoing steps.	(Freeman, 1978)
Betweenness	The frequency at which a node is located on the shortest path between any two pairs of nodes in the network.	(Freeman, 1978)

Giant weakly connected component (GWCC)	The proportion of nodes that are connected in the largest component when directionality of movement is ignored.	Wasserman and Faust (1994)
Giant strongly connected component (GSCC)	The proportion of nodes that are connected in the largest component when directionality of movement is considered.	(Wasserman and Faust, 1994)

769 **Table 2.** Description of biosecurity and infrastructure of the targeted farms.

Variable	Overall proportion
Presence of isolation barn(s)	No=77.6% Yes=22.4%
Fence around the farm	No=54.7% Yes=45.3%
Cleaning and disinfection station(s)	No=72.7% Yes=27.3%
Vegetation barrier around the farm	No=86.7% Yes=13.3%
Single entry point	No=72.4% Yes=27.6%
Bird nets	No=59.6% Yes=40.4%
Pig loading and unloading outside the farm's designated clean area	No=61.3% Yes=38.7%
Special clothing dedicated for farm use (shower in)	No=77.7% Yes=22.3%
Feed bins inside the designated clean area while loading is external	No=60.3% Yes=39.7%
*Vulnerability score ** Low Medium High Not informed	 0.21% 47.4% 17.8% 34.8%
Veterinary care From commercial system Independent veterinary Absent Not informed	 63.5% 4.5% 1.1% 30.9%
Farm type Nursery Breeding farm Finisher farm Certified swine breeder farm Wean-to-finisher Not informed	 32.23% 20.2% 15.1% 5.65% 1.49% 25.33%
Water sources Pond Natural water fountain Public source Lake Water cistern	 42.8% 22.2% 9.9% 1.5% 0.3%

Not informed	23.3%
Feed source	
From production system	68.7%
Own feed mill	7.8%
Not informed	23.5%
Destination of dead animals	
Compost	76.0%
Incinerator	0.23%
Bury	0.23%
Septic pit	0.24%
Not informed	23.3%
Manure destination	
Lagoon	28.9%
Septic pit	2.48%
Not informed	68.62%
Barn infrastructure	
Concrete or bricks	51.9%
Wood	1.29%
Mixed of wood and concrete	23.4%
Not informed	23.5%

770   ***This measures the level of vulnerability the introduction of pathogen into certified-swine-

771   breeder-farms, briefly is a combination of risks measurements form distance to farms to presence

772   of biosecurity and infrastructure of the farm.

 Figure Legend

Figure 1. Geographical location of farm density by type of pig operation, commercial,

independent, and not reported.

Figure 2. Description of pig movements between all farms types from January 2017 to

December 2018. The arrows size shows the percentage of movements and the color is given by

the origin of the movements. The size of the circles represented the total number of farms.

Figure 3. Description of the monthly variation of the network metrics of the pig movements

from January 2017 until December 2018.

Figure 4. Simulated disease spread dynamics within the swine contact network. Simulations

assumed two seed infection scenarios a) random, where 950 pig farms (10% prevalence) were

infected at day "0", and b) stratified, in which the proportion of the 950 pig farms (10%

prevalence) was equally distributed by commercial, independent, and not reported farms

operation types. The simulations assumed within farm prevalence of 0.1%.

Figure 5. The two years spread cascades sizes for each target-network surveillance strategy. In

light blue, the estimated spread cascade, which had a maximum value of 29 farms while not

targeting farms, resulted in a cascade of 2,624 farms (black).

Figure 6. The geographic locations of the top 1,000 farms ranked by degree.