Title: Tracking niche change through time: simultaneous inference of

environmental niche evolution and estimation of contemporary niches

Authors: Xia Hua[1,2*], Marcel Cardillo[1], Lindell Bromham[1]

[1]Macroevolution and Macroecology, Research School of Biology, Australian National

University, Canberra ACT 0200 Australia

[2]Mathematical Sciences Institute, Australian National University, Canberra ACT 0200

Australia

[*]Corresponding author:

Xia Hua

Mathematical Sciences Institute

Australian National University

Canberra ACT 0200

Australia

Tel: +61 2 6125 9773

E-mail: xia.hua@anu.edu.au

Abstract:

The environmental niche of a species is often studied from two different perspectives. Niche

modelling correlates a species' niche to known observations from contemporary species

distribution. Niche evolution applies phylogenetic methods on summary statistics of species

distribution to infer the patterns of changing niche over time. Even though the two areas of

research are fundamentally linked and are typically both based on the same species

distribution data, modelling the contemporary niche and inferring niche history are never

combined in the same analytical framework. Here we provide a new way to combine both

niche modelling and niche evolution, using distribution data and phylogenies. Doing so

allows us to simultaneously infer the history of niche evolution and estimate the

environmental niche of each tip species on the phylogeny. It also avoids some limitations of

the separate methods, such as the failure of environmental niche models to account for the

role of history in shaping current species distribution, and the use of unrealistic models of

change in phylogenetic comparative methods. The novelty of our method is that it explicitly

models three fundamental processes of niche evolution – adaptation, speciation, and dispersal

– applying a reversible jump algorithm to infer the occurrences of these processes on

phylogeny. Simulations demonstrate high accuracy of our method for estimating the

environmental niche of tip species and reasonable power to identify the occurrences of the

three processes on phylogeny. We also prove the efficacy of our approach using real-world

data. We show that our method of combining niche modelling with niche evolution is far

more effective at predicting salt tolerance in Australian *Acacia* species than using

environmental niche modelling alone. We also show that we can correctly infer periods of

evolutionary change in response to changing climate: our method identified two peak times in

drought adaptation events in the evolutionary history of Australian *Acacia*, which correspond

to the two drying periods in Australia during the Cenozoic. Therefore, we believe our method

provides a promising approach to integrate different research areas in understanding species

distribution.

Key words: Ecological niche modelling, Phylogenetic comparative methods, species

distribution, niche evolution, phylogeny, *Acacia*, salt tolerance, drought tolerance.

Running head: simultaneous inference of niche & niche evolution

A species' environmental niche describes the environmental conditions under which it

is able to maintain positive population growth (the 'fundamental niche' of Hutchinson 1957;

Holt 2009). Ecologists often have strong motivations for describing a species' environmental

niche, given observations made about its current distribution. One is to define the range of

conditions under which the species can persist: this information is useful for current

conservation planning (e.g. Weber et al. 2017) and for predicting future distributions given a

change in environmental conditions (e.g. Catullo et al. 2015; González-Orozco et al. 2016).

Prediction of the present-day environmental niche from data on species occurrence is referred

to as environmental niche modelling (ENM) or species distribution modelling (SDM). In

addition, we may be interested in reconstructing the history of change in a species niche over

time, in order to describe the way lineages respond to environmental change (e.g. Cang et al.

2016) and diversify to occupy a range of conditions (e.g.Castro-Insua et al. 2018). Typically,

these two areas of inference – modelling species present-day niches and reconstructing niche

evolution over time – have been conducted separately. But these two areas are tightly linked,

because the current niches of species are the product of evolutionary history, and

reconstructions of niche evolution usually proceed from data on current species niches.

Therefore, it is likely that a powerful approach for estimating current niches and

reconstructing their underlying history is to do both simultaneously, in a single analytical

framework (Smith et al. 2018).

While there have been advances in incorporating evolutionary information to improve

current niche estimation (Smith et al. 2018), such as estimating a species' niche by

aggregating its subspecies' niches (Maguire et al. 2018) or modelling niches of multiple

species while accounting for their phylogenetic non-independence (Morales-Castilla et al.

2017), we still lack a method that can simultaneously infer the history of niche evolution and

estimate current niches. In this study, we present a method that combines, in a single

statistical framework, an environmental niche model (ENM) and a phylogenetic comparative

method (PCM) to reconstruct niche evolution.

ENM estimates a species environmental niche as a mathematical representation of the

environmental conditions in the places where the species is known to occur or not occur

(Peterson 2001). ENM methods make the strong assumption that a species current

distribution is determined by its current environmental niche. But the observed distribution of

a species may not be an exact reflection of its environmental niche. Firstly, the available

occurrence data are typically an incomplete sample of the species true occurrences. A sparse

sampling of occurrence records is used to estimate a continuous range (extent of occurrence),

and those records may not be randomly sampled from the species distribution. ENM methods

have been developed to model and account for sampling biases, for example most species are

more likely to be recorded from areas close to population centres or roads (Fithian et al.

2015). In addition, while many studies reliably record the presence of a species in a given

sample location, explicit information on the absence of the species from sampling locations is

lacking, so we must model the probability that the lack of occurrence data represents true

absence or failure to sample. Secondly, the current distribution reflects not only the

environmental tolerances of the species, but also reflects other ecological processes such as

interspecific competition, as well as evolutionary history. The species may not fully occupy

its potential range if it has not had the time or opportunity to disperse to all parts of the range,

or if competitive exclusion has confined it to a suboptimal part of the environmental niche.

ENM methods typically use observed species occurrences without accounting for the

potential role of history and chance in determining current distributions of species.

Combining ENM with niche evolution models allows us to explicitly model how history has

shaped a species niche, leading to its current distribution.

The evolution of niches has previously been modelled in the framework of

phylogenetic comparative methods (PCM), which model the evolution of biological traits in a

group of species based on their current trait values and phylogenetic relationships

(Felsenstein 1985). PCMs have been applied to analyses of niche evolution by considering

the environmental niche to be a trait (Kozak and Wiens 2010). However, modelling niches in

this way raises two important problems. First, a species niche is often quantified as a

distribution of values along one or more environmental axes, but PCMs were designed to

work with point estimates rather than distributions (Hardy 2006). Second, niche evolution

involves complex (and still incompletely understood) biological processes, which may not be

adequately characterized by the kinds of models (usually Brownian Motion or one of its

variants) used to characterize trait evolution in PCMs. Models for trait evolution used in

PCMs have been criticized as lacking biological realism (Mu□nkemu□ller et al. 2015) and

being hard to interpret, because a range of different biological processes can lead to the

identical outcomes as expected by these evolutionary models (Revell et al. 2008; Losos

2011).

We can overcome many of the limitations of these two approaches by combining the

strengths of both. To combine ENM with niche evolution models and to address limitations

in using PCM to study niche evolution, we introduce a novel process-based model for niche

evolution. Our approach involves modelling explicitly the occurrence on phylogenies of three

biological processes that are expected to drive niche evolution and influence species

distribution: *in situ* adaptation, gradient speciation, and dispersal. By *in situ* adaptation, we

refer to a process where the environmental niche of the species shifts to encompass a new

range of conditions, for example, through the evolution of greater tolerance to drought.

Gradient speciation is a process where different populations of an ancestral species diverged

along an environmental axis and eventually evolve into distinct species. Dispersal is a

process where a species extends its current distribution to areas where conditions would

allow it to persist.

Where these processes occur on the phylogeny, and how they influence the

environmental niche of a species, define the species' history of niche evolution. We take a

Bayesian approach to simultaneously infer the history of niche evolution and the

environmental niche of each tip species on the phylogeny. First, we propose a history of

niche evolution. We then let the environmental niche of each tip species to evolve from the

root to the tip following the history of niche evolution. The inferred environmental niche of

each tip species allows us to assess the fit of the history of niche evolution by calculating the

likelihood of the observed distribution of each species given its inferred niche using ENM.

Last, we decide whether to accept the proposed history of niche evolution based on its fit to

the distribution data. These steps repeat until we get a good approximation of the posterior of

the history of niche evolution. The logic of our approach is similar to PCMs, but it extends

PCM application to traits that are not point estimates and allows more biologically

informative model of niche evolution.

We use simulations to demonstrate the power of our method to correctly infer the

history of niche evolution, and the accuracy of our method to estimate the environmental

niche of each tip species. We also demonstrate how the method can be used to test

hypotheses of niche evolution in real datasets, using the example of drought tolerance and

salt tolerance in Australian members of the plant genus *Acacia*. Australian *Acacia* is a good

case study for our method because they are widely distributed across Australia, often a

dominant part of the flora, and include specialists that are adapted to many different

environmental conditions, from desert taxa to alpine specialists to coastal dune species.

Australian *Acacia* is also a data-rich case study, because it is a large (~800 species) and

widely-distributed species group, with geographic distribution data for nearly all species and

a well-sampled phylogeny (Mishler et al. 2014). We focus on the evolution of drought

tolerance in Australian *Acacia* to test the power of our method to identify processes of niche

evolution. This allows us to test our inferred patterns of niche evolution against independent

data on changing climate. Specifically, there have been three significant drying periods in

Australia during the Cenozoic Era: around 28-23 Million years ago (Ma), 14-5 Ma, and

shortly after 5 Ma (Owen et al. 2017). Our method uses only contemporary species

distribution data, plus a phylogeny, so comparison of our results to this climate change

history provides an independent test of the usefulness of our method. If our method can

correctly infer the history of niche evolution, we would expect to see rapid evolution of

climatic niches during these periods as *Acacia* species developed adaptations to drier

climates. In addition, we use adaptation to high salinity in some *Acacia* species to evaluate

the performance of our method in estimating the environmental niche of each *Acacia* species.

Salt tolerance for several *Acacia* species has been well studied, so we can assemble a list of

known salt-tolerant species. We can then test if our method identifies these salt-tolerant

Acacia species as having significantly higher fitness under high salinity than the other *Acacia*

species, according to their estimated niche along a salinity axis.

METHODS

Basic model structure

Figure 1 summarizes the basic model structure of our approach. There are three major

components of our method: the phylogenetic comparative method (PCM), the model of niche

evolution, and the environmental niche model (ENM). We can express these three

components in terms of conditional probability as follows. PCM estimates

p(model|phylogeny), which is the probability of a certain history of niche evolution along a

given time-scaled phylogeny. The model of niche evolution estimates

p(niche|model,phylogeny), which is the probability of evolving the inferred contemporary

niche of each tip species given the history of niche evolution and the phylogeny. ENM

estimates p(presence and absence|niche), which is the probability of observing the presence

and/or absence of a species given our inferred niche of the species – this information allows

us to assess the fit of the inferred niche to the presence and/or absence locations of the

species. The product of these probabilities gives the probability of observing the presence

and/or absence of a species (the data) given its contemporary niche inferred from a history of

niche evolution along a phylogeny (the inference). Given the prior for each parameter in the

three components, we can make Bayesian inferences about both contemporary (tip) niches

and the history of niche evolution, as:

p(niche,model|presence and absence,phylogeny)

$= p$(niche|presence and absence)$*p$(model|niche,phylogeny) $\propto p$(presence and

absence|niche)$*p$(niche|model,phylogeny)$*p$(model|phylogeny)$*p$(parameters)

This gives the general framework of our method. The inputs are (presence and

absence,phylogeny): distribution data with environmental conditions, such as the degree of

aridity or salinity, at each presence location, and in some cases, reports of definite absence of

the species from other locations, plus a time-scaled phylogeny of the taxa. The outputs are

(niche,model): the posterior distribution of the environmental niche of each tip species and

the posterior distribution of the history of niche evolution. Figure 1 summarizes the

parameters to estimate in the Bayesian inference. We describe these parameters in each of the

three components of our approach in the next three sections.

The model of niche evolution: process-based model

For the model of niche evolution, we developed a process-based model, in which

three kinds of biological processes are involved in niche evolution and are modelled as events

that occur on the phylogeny. The three processes are (1) *in situ* adaptation, (2) gradient

speciation, and (3) dispersal. Each event has distinct properties modelled by local and global

parameters. Local parameters are those that are estimated for each event, such as the location

of the event on the phylogeny (see the PCM section) and the way that a species niche changes

during the event (see details below). Global parameters are those that are shared among

events, such as adaptation rate and dispersal rate (Fig. 1). To make the model of niche

evolution tractable, we assume that a species environmental niche evolves in a deterministic

way according to the order and the type of events that occur along the path from the root to

the tip, so p(niche|model,phylogeny)=1. Before we describe how a niche evolves in each type

of event, it is important to note that in these processes, fitness is expressed as probability of a

species persisting under particular environmental conditions. While described at the species

level, it is assumed to result from differential survival and reproduction of individuals. For

example, if the environment shifts, only those individuals of a species with tolerance to the

new environments may be able to reproduce, and in this way the range of environmental

conditions a species can tolerate can shift over time in response to environmental change.

We define three aspects of a species niche (Fig. 1) that change during the processes of

niche evolution. One is the fundamental niche of the species, the range of conditions under

which the species can persist. One is the available niche of the species, the range of

conditions in the potential areas that the species could expand into, if it had the ability to

persist under those conditions. Another is the realised niche of the species, the actual

distribution of the species or the environmental conditions under which it is currently found.

All these niches are specified with respect to a particular environmental factor (e.g. drought

or salt), but can be extended to include multiple environmental factors. In our current model,

we focus on the evolution of tolerance to extreme conditions, so environmental conditions are

assumed to follow a continuous gradient on an environmental axis, becoming more extreme

towards the right of the axis (Fig. 2).

Our model of niche evolution starts from the root of the phylogeny, so we need to

parameterize the three aspects of the ancestral species niche at the root. The fundamental

niche of the root species is assumed to follow $f(x) = e^{-(x/\alpha)^{\beta}}$ on the environmental axis, so

that under the least extreme condition, fitness $f(0) = 1$. Parameter α and β control the scale

and the shape of the niche: the larger the α, the more spread out the niche; the smaller the β,

the slower the fitness declines towards extreme conditions (Fig. 2). The available niche of the

root species is assumed to follow a normal distribution on the environmental axis. The

realized niche of the root species equals the product of its fundamental niche and its available

niche: that is, the distribution of the root species is at equilibrium, including all the areas of

suitable habitat that the species is able to disperse to and occupy. Without niche evolution,

the three niches are inherited by all the descendent lineages with no changes.

Now we can describe how the three aspects of niches change during *in situ*

adaptation, gradient speciation, or dispersal events. An *in situ* adaptation event (Fig. 2a) is

triggered by a change in the environmental conditions in the species distribution range. With

the environmental change, the species no longer has high fitness in its distribution area. Some

individuals die, some survive. This differential survival of individuals leads to shift in the

species realized niche and fundamental niche along the environmental axis. To model this

process, we first model the environmental change as a shift by n towards extreme conditions

in both the species' realized niche and available niche (Fig. 2a). In other words, the

environmental change influences both the species' distribution area and the areas it is able to

disperse to. Then we model the species new realized niche as the product of the species

fundamental niche and the shifted realized niche, because the species fundamental niche

describes differential survival of individuals under different environmental conditions and the

shifted realized niche describes the new environmental conditions in the species distribution

area before the species disperses to other areas (dispersal is modelled as a separate event). As

a result, the final amount of shift from the old realized niche to the new realized niche is not

n and we denote it as n' in Figure 2a. Along with the change in the realized niche, the species

fundamental niche also evolves into higher fitness in extreme conditions. According to our

parameterization of the fundamental niche, higher α leads to higher fitness in extreme

conditions and lower β leads to less abrupt decline in fitness towards extreme conditions

(Fig. 2a), so increasing fitness in extreme conditions is reflected in increasing α values and

decreasing β values. So, we model changes in the fundamental niche as the amount of change

in α and β, which is $\gamma n'$ and $-\gamma n'$ on a logarithmic scale, where γ is adaptation rate. A

species with higher adaptation rate evolves its fundamental niche faster in response to

environmental deterioration, by evolving higher α and lower β.

When a gradient speciation event occurs (Fig. 2b), two populations of an ancestral

species become distinct species, each occupying different conditions along an environmental

axis. In Figure 2b, where the ancestral species splits on the axis is determined by parameter p,

such that a proportion p of individuals on the right of the ancestral species realized niche

become a species distinct from the remaining 1-p individuals on the left of the realized niche.

One new species consists of individuals with higher fitness under the extreme end of the

conditions that the ancestral species could tolerate. The other new species will have, on

average, lower fitness under these extreme conditions. For both species, the amount of

change in α and β during gradient speciation is $\gamma n'$ and $-\gamma n'$ on the logarithmic scale, where

n' is difference in the mean of realized niche between the new species and the ancestral

species. A gradient speciation event can occur at any node in the phylogeny. When this

happens, the two daughter species are randomly assigned to the two descendant lineages in

the phylogeny, based on the sign of p. With a non-zero extinction rate, a gradient speciation

event can also occur along a branch in the phylogeny with one of the two daughter species

going extinct. When this happens, the gradient speciation effectively leads to a punctuated

change in the realized niche and the fundamental niche of the ancestral species. The sign of

parameter p decides which species remains and so the other species is regarded as extinct.

Between any two events, either *in situ* adaptation or gradient speciation, dispersal

occurs (Fig. 2c), which is a continuous process of a species to extend its realized niche along

the environmental axis until it reaches equilibrium. The species is at equilibrium when its

realized niche equals the product of its fundamental niche and its available niche. Dispersal

rate (d) defines how fast a species reaches the equilibrium. We consider dispersal after each

in situ adaptation event or each gradient speciation event, because these events cause a lag

between the species realized niche and the available niche during which the species

distribution is not at equilibrium. Since we parameterize the available niche as a continuous

distribution, which does not allow any discrete barrier for dispersal, we introduce another

parameter x to model a dispersal barrier by truncating the available niche under non-extreme

conditions (truncating the side left to x in Fig. 2c). We only model dispersal barriers under

non-extreme conditions, because a species distribution under extreme conditions is

constrained by the fundamental niche and so little signal for the presence of dispersal barrier

under extreme conditions can be detected. The mechanism for a barrier is not important for

our model, instead we treat x as a nuisance parameter. For example, we model x and p as two

independent parameters, but this does not exclude the possibility that the location of the

barrier (x) may coincide with the location where species splits on the environmental axis (p)

during a gradient speciation event, which may suggest competitive exclusion during the

speciation event.

In summary, our process-based model includes three types of processes: dispersal, *in*

situ adaptation, and gradient speciation. Each type of process is described by two parameters:

dispersal rate (d) and the location of barriers on the environmental axis (x) for a dispersal

event; adaptation rate (γ) and the amount of environmental change (n) for an *in situ*

adaptation event; adaptation rate (γ) and the location of species split on the environmental

axis (p) for a gradient speciation event (Fig. 1 and Fig. 2). To reduce the risk of over-

parameterization, dispersal rate and adaptation rate are assumed to be constant over time and

across lineages, so they are global parameters (Fig. 1). Each *in situ* adaptation event or

gradient speciation event is assumed to be a discrete event, occurring at a specific time point

on the phylogeny. Dispersal event is a continuous process, happening in-between two

discrete events. So, for each *in situ* adaptation event and gradient speciation event, we have

an additional parameter (t) to describe the location on the phylogeny when the event occurs

(Fig. 1). Last, we have two parameters (mean and variance of a normal distribution) to define

the available niche and two parameters (α and β) to define the fundamental niche of the

ancestral species at the root of the phylogeny (Fig. 1).

ENM: inhomogeneous Poisson point process

The model described above shows how we model the evolution of the three aspects of

niche forwards in time along the phylogeny, starting from the niches at the root, evolving

through a series of events of the three different biological processes, resulting in the niches at

the tips. The three niches of each tip are then used to calculate *p*(presence and absence|niche)

using ENM. In principle, any niche modelling method (ENM) can be used to calculate

p(presence and absence|niche). Here we use an inhomogeneous Poisson point process

(Renner et al. 2015), because this process can account for sampling bias, and so is applicable

to both presence/absence data or presence-only data without generating pseudo-absence

locations (Fithian et al. 2015). The process models the number of recorded presences of a

species per unit area as a Poisson random variable. The inferred rate of recording a presence

of the species in the area depends on the abundance of the species in that area and the

sampling intensity in that area (see details in Fithian et al. 2015). Below, we describe how the

three aspects of niches are linked to parameters in the ENM, assuming equal sampling

intensity across the area (unequal sampling intensity is accounted for later in the Case study

section).

We can start by considering a dataset where we have presence-only data on

distribution: that is, where our knowledge of species niches are informed by records of that

species at particular sampling locations for which we can derive relevant environmental data.

In brief, the log-likelihood of the presence-only data of each tip species is

$$\ell_P = \sum_{i \in I_{PO}} \log(\lambda_{r_i}) - \mathcal{D} \int \lambda_f \lambda_a ds$$

, where I_{PO} is the set of all the species presence-only locations. In the first term, λ_{r_i} is the

realized niche, or the rate of recording a presence of the species under the environmental

condition at the i^{th} presence location. The second term is the integral of the product between

the fundamental niche (λ_f) and the available niche (λ_a), and $\mathcal{D}$ is the geographic area of the

available niche. This term describes the overall probability of recording a presence of the

species in the species available niche. Often, $\mathcal{D}$ cannot be measured accurately without

knowing the dispersal ability of a species. However, it can be used to adjust how informative

presence-only locations are for inferring the species fundamental niche. For example, a

species with widely scattered presence records has a higher $\mathcal{D}$ value than a species with more

spatially concentrated records, so in the species fundamental niche, the same amount of

decrease in fitness in conditions not in the presence locations will increase the model

likelihood more for the species with higher $\mathcal{D}$ value.

Now we can consider the case where in addition to presence data (indicating points

where the species is known to occur), we also have absence data (locations where the species

was actively sought but not found). Absence of a species is often inferred from surveys

within a local area, such as quadrat sampling. The log-likelihood of the presence-absence

data of each tip species is

$$\ell_A = \sum_{i \in I_{PA}} \left[-y_i \log\left(1 - e^{-\lambda_{r_i}}\right) + (1 - y_i)\lambda_{r_i} \right]$$

, where I_{PA} is the set of all the presence and absence locations of the species, with $y_i = 1$ if

the species is present at the i^{th} location and $y_i = 0$ if the species is absent at the i^{th} location.

We don't need to account for the available niche here, because the survey is complete and so

wherever the species does not occur, it is assumed to be absent.

To make our model analytically tractable, we converted the continuous environmental

axis into bins. The minimum and maximum bins are determined by the minimum and

maximum values of environmental conditions in the study area. The three aspects of niche

were converted into discrete distributions using the probability that values from these niche

distributions fall in the range of each of the bins. For the bin containing the minimum or the

maximum value, it is the probability that values from these niche distributions are lower or

higher than the minimum or the maximum value.

PCM: reversible-jump model

Our model of niche evolution includes three biological processes: *in situ* adaptation,

gradient speciation, and dispersal. A phylogenetic comparative method (PCM) is needed to

estimate the number and the location of each type of event on the phylogeny. Here we

assume events occur across a phylogeny under an inhomogeneous Poisson process and use

reversible jump Markov Chain Monte Carlo (rjMCMC; Green 1995) to implement the

process in a Bayesian framework. Several PCM studies have used rjMCMC to study jumps in

evolutionary processes of phenotypic evolution (e.g., Eastman et al. 2011; Uyeda and

Harmon 2014) and diversification (Rabosky et al. 2014). In our implementation, each jump is

an occurrence of a discrete event: an *in situ* adaptation event or a gradient speciation event.

As in these previous studies, our implementation involves several steps: 1) adding a new

event to the model, 2) removing an existing event from the model, 3) updating the location of

an existing event on the phylogeny, 4) updating parameters specific to that event, 5) updating

parameters shared among events, 6) updating the occurrence rate of an event. Details on these

steps are given in the Appendices.

To calculate p(model|phylogeny), we assume *in situ* adaptation events occur

randomly according to a homogeneous Poisson process with rate r_A, and gradient speciation

events occur randomly according to an inhomogeneous Poisson process with overall rate r_S

(Fig. 1). The occurrence of gradient speciation is inhomogeneous because a speciation event

occurring along a branch must be accompanied by an extinction event and the occurrence

probability of the extinction event depends on the time (t) when the speciation event occurs,

which is $p_E(t) = \frac{\mu(1-e^{-(\lambda-\mu)t})}{\lambda-\mu e^{-(\lambda-\mu)t}}$ (Kendall 1948), where λ is speciation rate and μ is extinction

rate. Given that a gradient speciation event occurs, the probability of the event occurring at a

node relative to the probability of the event occurring along a branch is the number of nodes

in the phylogeny divided by the integral of $\lambda p_E(t)$ over all the branch lengths of the

phylogeny. When a gradient speciation event occurs at a node, it has equal probability to

occur at any node in the phylogeny. When a gradient speciation event occurs on a branch, the

relative probability that the event occurs on a specific branch depends on the integral of

$p_E(t)$ from the starting and the ending times of that branch, and the relative probability of the

event to occur at time t in that branch is $p_E(t)$.

Simulation study

We performed a simulation study to assess the performance of our method given the

assumptions that species niches evolve according to our process-based model, and processes

occur across phylogenies under the inhomogeneous Poisson process. First, we generated

simulated datasets by evolving niches along phylogenies, then sampling locations from the

resulting realized niche of each tip species to give species presence and absence data.

Phylogenies were simulated under a birth-death process using the R package diversitree

(FitzJohn 2012) with 100 tips, a birth rate of 0.1, and a death rate of 0.03. All the simulated

phylogenies had the same available niche along an arbitrary environmental axis at the root,

with mean of 1 and standard deviation of 2, and the same fundamental niche at the root, with

α of 3 and β of 2. On each simulated phylogeny, we randomly placed 20 *in situ* adaptation

events with equal probability of placement at any location on the phylogeny, and we placed

20 gradient speciation events with different probabilities of different locations, as described

above for p(model|phylogeny). Parameters specific to each event (x, n, p) were randomly

drawn from their corresponding prior distributions (see Appendices). Species niches were

then evolved through the randomly placed events along the path from the root to each tip

according to our process-based model. This produced the available niche, the realized niche,

and the fundamental niche of each tip species. During this process, the environmental axis

was discretised into 10 bins and dispersal rate was fixed to 1 across the phylogeny. Two

simulation schemes were applied to adaption rate (γ). In one scheme, γ was fixed to 0.03

across the phylogeny. In the other scheme, γ on each branch was randomly drawn from an

exponential distribution with mean of 0.03, so that different lineages may have different

adaptation rates. We generate two sets of sample data for the analysis. One dataset is

presence-only data, in which presence-only locations were randomly drawn for each tip

species from the species realized niche (λ_r). The number of presence-only locations is drawn

from a Poisson distribution with mean equal to the overall probability of recording a presence

of the species in the species available niche. The other dataset is presence and absence data,

in which 1000 presence-absence locations were randomly drawn for each tip species with

equal probability from each bin of the environmental axis, of which the i^{th} location has $e^{-\lambda_{r_i}}$

probability to be an absence location.

For each simulated phylogeny and the sampled presence-only and presence-absence

data of the tip species on the phylogeny, we ran two independent MCMCs for 10^6

generations, with a thinning interval of 1000 and the first 600,000 generations discarded as

burnin. We tested two aspects of the performance of our method: (1) accuracy of estimating

the fundamental niche of each tip species, and (2) power to correctly identify the occurrences

of *in situ* adaptation events and gradient speciation events on the phylogeny. To assess the

accuracy of our method to estimate the fundamental niche of each tip species, we first

estimated, for each simulation, the fundamental niche of each tip species as the average of the

values of parameters α and β of each tip species in the posterior samples. We then compared

these estimated values of parameters α and β of each tip species to their corresponding

values in the simulation, using linear regression. Last, we summarized the regression results

over all the simulations by the slope estimate and the R squared of the regression model.

Slope values close to 1 and R squared values close to 1 indicate a high level of accuracy in

our estimation of the fundamental niche of tip species.

To assess the power of our method to identify *in situ* adaptation events, we first

calculated for each branch the posterior probability of at least one *in situ* adaptation event

occurring on each branch, by counting the proportion of posterior samples with *in situ*

adaptation events on that branch. We then identified branches that had significantly higher

posterior probability than expected if all the branches had equal probability of having

adaptation events. Following Shi and Raboksy (2015), a branch had significantly higher

posterior probability than expected, if its marginal odds ratio was larger than a threshold

arbitrarily set to 5. This is essentially hypothesis testing based on Bayes factor for a model

with adaptation events on the branch (against a model without adaptation events on the

branch), and Bayes factor larger than 5 is generally considered as substantial evidence for the

model. Next, we calculated the power of our method as the proportion of branches with true

adaptation events that passed the test: in other words, how many branches with true

adaptation events we have identified. We also calculated the false positive rate of our method

as the proportion of branches that passed the test but with no true adaptation events: in other

words, how many inferred adaptation events were not true adaptation events.

Last, we wished to find out what kind of branch is more likely to pass the test. We

expected that events that occurred more recently, or events with larger influence on niche

evolution, are more likely to be correctly identified by our method, so we measured two

properties of each branch. For a branch with true adaptation events, one property is the

average of the ages of all the *in situ* adaptation events placed on the branch during simulation.

The other property is the sum of the value of parameter *n* over all the *in situ* adaptation events

placed on the branch during simulation, with a larger value suggesting more niche evolution

due to *in situ* adaptation on that branch. For a branch with no true adaptation events, the same

two properties were measured, but they were measured by the events included in the posterior

samples instead of the events placed on the branch during simulation. We divided all the

branches into groups according to the two properties. For each group, we calculated the ratio

of the sum of the posterior probabilities of all the branches with true adaptation events against

the sum of the posterior probabilities of all the branches with no true adaptation events. A

ratio larger than 1 suggests that, in the group, a branch that passed the test is more likely to

contain true adaptation events. We applied the same steps to assess the power of our method

to identify gradient speciation events, except that the second property was measured by the

average of the absolute value of parameter *p* over all the gradient speciation events placed on

the branch during simulation.

Case study: drought and salt tolerance in Acacia

For the *Acacia* case study, we used a published dataset that consists of 132,295

presence locations of 508 *Acacia* species (~80% described *Acacia* species in Australia) as

well as a published phylogenetic tree that includes the same 508 *Acacia* species (see Mishler

et al. 2014). We dated the phylogenetic tree by the relaxed clock model using R package ape

(Paradis and Schliep 2018) with the root date constrained between 21 and 23.9 million years

ago, a range of crown divergence dates previously estimated for *Acacia* (Miller et al. 2013).

We assembled species presence-absence survey data from New South Wales

(www.environment.nsw.gov.au), Victoria (www.data.vic.gov.au), and West Australia

(naturemap.dpaw.wa.gov.au) government databases. We only included data from complete

floristic surveys that were conducted later than 2005 to ensure consistency in taxonomy, and

under government custodianship to ensure survey quality. As a result, our presence-absence

survey data included a total of 27,850 survey sites across three states of Australia. These

presence-absence data were analysed together with the presence locations. We extracted the

degree of aridity and salinity at each presence location of each species and each survey site

from the geographic layer of maximum monthly aridity index (Williams et al. 2012) for the

drought tolerance analysis, and from the geographic layer of root zone soil electrical

conductivity (Bui et al. 2017) for the salt tolerance analysis.

For both analyses, we modelled sampling bias in the presence locations. This bias

results from the fact that the majority of presence locations for most species are in the form of

presence-only records that are typically collected opportunistically, so their spatial

distributions tend to be biased towards regions that are more easily accessible to researchers

(Fithian et al.2015). To correct for this bias, we included two factors that may influence

sampling effort at a location: distance to the nearest road (using data from

www.ga.gov.au/geoscience/national) and whether it is in a protected area, such as a

conservation reserve or national park (from

www.environment.gov.au/land/nrs/science/capad).

Following Fithian et al. (2015), the log-likelihood of the presence-only data of each

tip species is modified as,

$$\ell_P = \sum_{i \in I_{PO}} \left[\log(\lambda_{r_i}) + \delta z_i \right] - \frac{\mathcal{D}}{n} \sum_{i \in I_{BG}} \lambda_{f_i} \lambda_{a_i} e^{\delta z_i}$$

, where z_i are values of the factors that influence sampling effort at the i^{th} location and δ are

the coefficients of these factors, which are assumed to be constant across tip species and were

updated in the same way as the other global parameters in each MCMC (see Fig. 1 and

Appendices). Note that the second term is no longer a direct integral of $\lambda_f \lambda_a$, because we also

need to account for sampling bias. We can only approximate the integral by randomly

sampling n locations from the surroundings of the species presence locations to construct a

set of background locations I_{BG} and averaging the term to integrate over all the background

locations. For each *Acacia* species, background locations were sampled randomly from areas

within an arbitrary large radius (20 *km*) of each presence location of the species. The total

number of background locations was ten times the number of the species presence locations,

with a minimum of 1000 for species with few presence locations. The area ($\mathcal{D}$) is the total

area that falls within the 20 *km* radius of each species presence location. Note that although

the radius is arbitrarily defined and background locations were sampled randomly from the

area, whether the species is potentially present at i^{th} background location depends on both its

fundamental niche λ_{f_i} and its available niche λ_{a_i}. Based on the environmental conditions

over all the background locations, we used Sturges' formula (Sturges 1926) to decide the bin

size to discretise the aridity index and the electrical conductivity.

The starting values for the available niche and the fundamental niche at the root, and

the coefficients for the two factors that may influence sampling effort, were set to their

maximum likelihood estimates given the likelihood of each tip species' presence-only data

and presence-absence data, assuming that no events of niche evolution occurred on the

phylogeny. The starting value for dispersal rate (d) was set to 1 and for adaptation rate (γ) to

0.03. Dispersal rate and the coefficients for factors on sampling effort had rather flat priors,

an exponential distribution with rate of 0.1. The parameters for the niches at the root and the

adaptation rate (γ) had relatively narrow priors, a lognormal distribution with logarithm mean

of their initial values and logarithm standard deviation of 0.5. The narrow priors for the root

niches were used to ensure that the root niches do not exceed the maximum and minimum

observed values for aridity and salinity in Australia too often during MCMCs. The narrow

prior for γ was used to ensure a finite likelihood value, as parameters of the fundamental

niche change proportionally to γ on a log scale and so there is a narrow range of γ that gives

finite likelihood values. The starting value for the occurrence rate of *in situ* adaptation events

and the occurrence rate of gradient speciation events was set to 0.1. Because the simulation

study suggests a tendency to overestimate the number of *in situ* adaptation events and

gradient speciation events (see Results), we used a narrow prior for the occurrence rates of

both events (r_A and r_S), an exponential distribution with rate of 10, in order to reduce the

degree of overestimation. For both studies, we ran 20 independent MCMCs for 1×10^6

generations, with a thinning interval of 1000 and the first 600,000 generations discarded as

burnin.

We used the climatic history of Australia to generate *a priori* hypotheses which we

then tested against the data. Miller et al (2013) estimated the age of the common ancestor of

extant *Acacia* as 23Ma at the earliest, and since then there have been two key periods of

declining temperatures and increasing aridity: a long period from 14-5Ma, followed by a

brief warmer and wetter period at the start of the Pliocene, then a second period of

aridification through the late Pliocene and through the Pleistocene (Owen et al. 2017). So we

would expect our method to infer more environmental change for the *in situ* adaptation

events, along the axis of drought tolerance, between 14-5 Ma and shortly after 5 Ma,

compared to other periods. To test this, we plotted the histogram of the dates of *in situ*

adaptation events in the posterior samples, with the date of each event weighted by the

amount of environmental change (*n*) for that event. We then corrected the relative frequency

in each bin or date interval of the histogram by the number of lineages on the phylogeny

during that interval, so that the frequency gives the relative amount of environmental change

that each lineage had experienced during the date interval in history, as inferred by our

method. Using the same method, we calculated the frequency for the salt tolerance analysis

as a comparison to the draught tolerance analysis.

In order to test the accuracy of our method for predicting current niches, we used

independent information on the salinity tolerance of *Acacia* species from the literature to

compile an *a priori* list of known salt-tolerant species. The list of known salt-tolerators

included 12 *Acacia* species in the phylogeny (see Table S1 in Appendices), allowing us to

ask whether our method correctly identifies these species as salt-tolerators. Note that this is

an incomplete list as we only included studies that did physiological experiments on the

species over a range of salinity conditions, so that we were confident that each species on the

list was salt tolerant. We then calculated the mean of parameter α and β of the fundamental

niche of each tip species from the posterior samples. If our method gives reasonable estimates

for the fundamental niche of each tip species, then the estimated α of the 12 known salt-

tolerant *Acacia* species should be at the high end of the distribution of values of other *Acacia*

species and the estimated β of these salt-tolerant species should be at the low end of the

distribution of values of other *Acacia* species. A Welch's *t*-test was used to test for the

significance in these differences. As a comparison to standard ENM that do not incorporate

niche evolution, we also estimated the environmental niche along the salinity axis of each

Acacia species from only ENM, using the MultispeciesPP R package (Fithian et al. 2015)

which implements the same ENM model as used by our method and corrects sampling bias

across species. This ENM analysis was applied to the same dataset as our method, i.e., the

same species distribution data, the same data on soil electrical conductivity, and the same

variables to correct for sampling bias. We used three customised natural spline bases of the

soil electrical conductivity to predict a species' environmental niche along salinity axis. We

then tested if the 12 known salt-tolerance *Acacia* species have higher predicted fitness, i.e.,

larger area under their predicted niche curve, under high salinity than the other *Acacia*

species, using Welch's *t*-test. We used different threshold to define high salinity and reported

the lowest *p* value of the Welch's *t*-test.

RESULTS

Simulation study

Overall, simulations suggested that our method performed well under the assumptions

of the method, with high accuracy in estimating the fundamental niches of tip species and

reasonably high power to identify the occurrence of processes of niche evolution on the

phylogeny. Over all the simulations, the average R squared is above 0.8 and the average

slope is around 1, suggesting that the estimates of both parameters α and β of the

fundamental niches of tip species are highly correlated with their true values in the

simulations (Fig. 3). The power of our method to correctly identify branches with *in situ*

adaptation events (the true positive adaptation) is on average 0.8 over simulations, and the

power to correctly identify branches with gradient speciation events (the true positive

speciation) is on average 0.5 over all simulations (Fig. 3). However, our method has a high

false positive rate for both *in situ* adaptation events and gradient speciation events, which is

on average 0.6 over all simulations (Fig. 3). For *in situ* adaptation events, these false positives

tend to be events with low amount of environmental change (or low influence on niche

evolution; Fig. 4b). In other words, inferred *in situ* adaptation events with lower amount of

environmental change are more likely to be 'false' events. For gradient speciation events, the

false positives tend to be old events closer to the root (Fig. 4a). This suggests that our method

does not reliably identify gradient speciation events that occurred in the early diversification

history of a species group, at least within this parameter space. This is not surprising given

that these gradient speciation events do not influence the available niche and so their

influence on species distribution can be quickly erased by dispersal.

Note that the sum of true positive rate and false positive rate for both *in situ*

adaptation and gradient speciation is over one (Fig. 3), suggesting that our method

overestimates the number of both types of events. However, our method still accurately

estimates the fundamental niche of each tip species (Fig. 3). So falsely inferred events overfit

the history of niche evolution to species presence-only and presence-absence data, but they

have minimal influence on the accuracy of the inferred amount of niche evolution from the

root to each tip species. This is also suggested by the result that inferred gradient speciation

events with older dates and inferred *in situ* adaptation events with smaller amounts of

environmental change are more likely to be false events (Fig. 4). In other words, species

distribution data are less informative for events with less influence on niche evolution, so the

number of falsely inferred events in the posterior samples is more sensitive to the prior than

the number of true events. Therefore, we suggest using narrow priors on the event occurrence

rates (r_A and r_S) in any empirical application of our method, in order to reduce the number

false events.

An unrealistic assumption of our method is constant adaptation rates across lineages,

which is unlikely to be true in the real world, as species' responses to natural selection

depend on many factors. To assess the performance of our method when this assumption is

violated, we simulated the adaptation rate of each branch on each simulated phylogeny as a

random variable. Note that the actual amount of niche evolution is on the exponential scale of

the adaptation rate, so little variation in the adaptation rate will cause large differences in the

fundamental niche. As expected, our method has reduced accuracy in estimating the

fundamental niche of each tip species under the violation, with the average R squared

reduced to 0.5 and the average slope reduced to 0.5 (Fig. 3). However, the violation does not

affect the power of our method to identify the occurrence of niche evolution on the

phylogeny. The true positive and false positive rates of both *in situ* adaptation events and

gradient speciation events do not differ significantly from those under the simulation scenario

of constant adaptation rate (Fig. 3).

Drought and salt tolerance in Acacia

Applying our method to drought tolerance and salt tolerance in *Acacia* species gives

results that are consistent with expectations, suggesting that our method provides useful

insights into niche evolution in the real world. First, for both drought tolerance and salt

tolerance, *in situ* adaptation and gradient speciation events tend to have high occurrence in

clades that are distributed predominantly in semiarid and arid areas of Australia (mostly

Central and Western Australia, Fig. 5 and Fig. 6), such as the *victoriae*/*pyrifolia* clade (Miller

et al. 2011), while few events occur in the large clade of a recent radiation in relatively mesic

areas (mostly Northern and Eastern Australia, Fig. 5 and Fig. 6).

Second, for drought tolerance, the inferred amount of niche evolution per lineage due

to *in situ* adaptation events peaks at 13 Ma and near the present, corresponding to the two

drying periods of Australia at 14-5 Ma and after 5 Ma (Fig. 7). For salt tolerance, the inferred

amount of niche evolution per lineage due to *in situ* adaptation events peaks near the present,

with two additional lower peaks around 10 Ma and 17 Ma (Fig. 7). As a result, the inferred

amount of niche evolution due to *in situ* adaptation events does not correlate between drought

tolerance and salt tolerance (Kendall's rank correlation coefficient=0.04, p=0.13; compare

Fig. 5a and Fig. 6a). However, the inferred amount of niche evolution due to gradient

speciation events significantly correlates between drought tolerance and salt tolerance

(compare Fig. 5b and Fig. 6b): the average value of parameter p of the gradient speciation

events inferred on a branch is correlated between the analysis on drought tolerance and the

analysis on salt tolerance (Kendall's rank correlation coefficient=0.12, p<0.001).

Third, for salt tolerance, the estimated values of parameter α and β of known salt-

tolerant species are significantly higher than the estimated values of the other *Acacia* species

(Welch's t=2.48, p=0.01). Since species with higher α have higher fitness under extreme

condition and species with lower β have less abrupt decline in fitness towards extreme

condition, these results suggest that our methods successfully predict higher fitness under

high salinity of the known salt-tolerant species compared to remaining species. In contrast,

using only ENM without accounting for shared history of niche evolution among species, we

cannot identify the known salt-tolerant species. Their estimated niches do not have higher

fitness under high salinity compared to the other *Acacia* species (Welch's t=0.85, p=0.21).

DISCUSSION

Environmental niche models (ENMs) range from correlative models that infer species

fundamental niches or potential geographic distributions from observed occurrence-

environment correlations, to mechanistic models that incorporate processes based on the

physiological or functional traits of species (Kearney and Porter 2009; Dormann et al. 2012).

However, no current approach accounts for the historical role of speciation and adaptation in

determining species present-day niches. While there have been several recent developments

of ENMs that incorporate a phylogenetic dimension (Smith et al. 2018; Morales-Castilla et al.

2017; Maguire et al. 2018), these have mostly aimed to account for the statistical problem of

non-independence due to relatedness among contemporary species, rather than to model the

historical development of niches through evolutionary time. By combining ENMs and niche

evolution models in the same statistical framework, the approach we present in this paper

allows for species present-day niches to be shaped by evolutionary history and allows us to

test historical hypotheses of niche evolution. Adding historical processes to ENMs results in

better predictions of present-day niches from contemporary data. Simulations show that the

method accurately estimates the fundamental niche of species and has reasonable power to

identify the occurrences of *in situ* adaptation events and gradient speciation events on the

phylogeny. Using the method, we were able to infer temporal patterns in drought tolerance

and salt tolerance in *Acacia* species that are consistent with predictions based on major trends

in Australia's climatic history over the past 23Myr. We were also able to identify known salt-

tolerant *Acacia* species with this method, while the standard ENM could not.

Drought and salt tolerance in Acacia

The climatic history of Australia through the second half of the Cenozoic has been

characterized by increasing aridification punctuated by several brief periods of more mesic

climates (Byrne et al. 2008). The contraction of humid forests to coastal areas, and their

replacement by the more open, sclerophyll vegetation that now covers much of the continent,

began around 33Ma following the establishment of the Antarctic Circumpolar Current and a

global drop in temperatures at the Eocene-Oligocene boundary. During this period

Australia's flora became dominated by arid-adapted lineages, especially in the large families

of Myrtaceae, Fabaceae and Proteaceae (Crisp and Cook 2013). To survive drought, species

in these families have evolved a range of strategies to cope with water stress. These strategies

have been broadly classified into escape, avoidance, and tolerance (Delzon 2015). In *Acacia*,

all three strategies are seen, for example phyllode shedding (escape), scleromorphic

structures such as sunken stomata or needle-shaped phyllodes (avoidance), and physiological

adaptations such as low leaf water potential (Moore 2013). The wide range of mechanisms

adopted by *Acacia* species to cope with water stress are summarized in their geographic

distributions, and hence their climatic niches.

Because aridification of Australia has occurred at varying rates and was more rapid in

the periods 28-23Ma, 14-5Ma and after 5Ma and because *Acacia* originated around 23 Ma at

the earliest, we hypothesized that bursts of niche evolution along the drought axis (aridity

index) would correspond with 14-5Ma and after 5Ma periods. Our method was able to

reconstruct both temporal and spatial variation in the degree of niche evolution consistent

with these expectations. We inferred peak times in adaptation to drought that correspond to

these two major drying periods, and we reconstructed a concentration of niche evolution

events in both drought and salinity tolerance in clades distributed predominantly in arid and

semiarid regions.

The amount of niche evolution in the aridity index and root zone electrical

conductivity attributable to gradient speciation events is highly correlated across branches of

the *Acacia* phylogeny. This result is consistent with the observation that saline soils are

frequently associated with aridity, so many species are adapted to both high salinity and high

aridity. However, we found no correlation in the occurrence of adaptation events between

drought tolerance and salt tolerance: while adaptation events in drought tolerance occurred

primarily in the major drying periods, adaptation events in salt tolerance occurred primarily

near the tips (Fig. 6). This kind of "tippy" macroevolutionary pattern is consistent with the

finding in grasses that salt tolerance evolves frequently but does not give rise to a large clade

of salt-tolerant descendants (Bennett et al. 2013; Bromham et al. 2016), suggesting that the

evolution of salt tolerance may be a macroevolutionary "dead-end" (Bromham et al 2016).

Consequently, even if salt tolerance has evolved many times through old adaptation events,

these events rarely gave rise to persistently salt tolerant lineages of large clades of related salt

tolerant taxa, leaving less phylogenetic signal for us to reconstruct adaptation events in the

deep part of the phylogeny than adaptation events near the tip of the phylogeny.

We also found that, in *Acacia* salt tolerance, branches with high occurrence of

gradient speciation events tend to also have high occurrence of *in situ* adaptation (Kendall's

rank correlation coefficient=0.35, p<0.001). This result suggests an important role of gradient

speciation induced by environmental change in the evolution of tolerance to high salinity in

Acacia. The prevalence of gradient speciation in *Acacia* species along the salinity axis may

provide a mechanism for the high lability of salt tolerance (Bromham et al. 2016).

Another contribution of our study to understanding the macroevolutionary pattern of

tolerance is that it provides a continuous estimate of the degree of tolerance for tip species.

Previous studies on macroevolutionary patterns of tolerance have required species to be

categorized as either tolerant or sensitive to an environmental stress, based on existing

evidence or expert opinions (e.g., Bennett et al. 2013; Moray et al. 2015; Moray et al. 2016;

Bromham et al. 2016). These studies had a common methodological issue that the cutoff for

distinguishing tolerant from sensitive species is arbitrary, and many species cannot be

assigned definitively to one category or the other. Our method was able to identify known

salt-tolerant *Acacia* species by their values of parameters α and β, suggesting that these

parameters can give informative estimates of the degree of salt tolerance from real data. Our

method therefore provides a more flexible approach to analyzing macroevolutionary patterns

of tolerance by estimating the degree of tolerance from current species distribution data,

which are available in some form for large numbers of species.

Process-based model of niche evolution

Unlike previous methods that incorporate evolutionary history into niche models

(Smith et al. 2018), our method does not use phylogenetic distance as a measure of how

much information on a species' environmental niche can be derived from the distribution data

of other species, an approach that assumes strong phylogenetic niche conservatism

(Mu☐nkemu☐ller et al. 2015). Instead, we explicitly inferred events of niche evolution on

the phylogeny from species distribution data and let the data tell us the amount of niche

evolution occurred on each branch of a phylogeny. As a result, the amount of inferred niche

evolution along the ancestral lineage of any two species describes the relatedness in their

niches. But as with all methods, our method has its own assumptions. One major assumption

is that the environmental niche of a species has evolved according to our process-based

model. Our model includes three fundamental processes of niche evolution – *in situ*

adaptation, gradient speciation, and dispersal - but the modelling of each process is highly

simplified in order to reduce the number of parameters and limit calculation time. For

example, we use one parameter (p) to model the process of gradient speciation by splitting

the realized niche at a particular point, irrespective of the cause of the speciation event, but

different causes may result in different ways of splitting the realized niche. For example,

incipient species under divergent selection may actively move apart from each other to

occupy their optimal environments rather than simply cut their realized niche at a particular

point. However, even with highly simplified processes of niche evolution, our model is

already much more biologically informative than a pure mathematical model, such as the

Brownian-motion related models used by most previous PCMs (e.g. Revell et al. 2008; Losos

2011).

The other major assumption is that the occurrence of events of niche evolution

follows a compound Poisson process. Like other applications of the compound Poisson

process on molecular evolution and trait evolution, this assumption makes our model of niche

evolution weakly identifiable (Moore et al. 2016); that is, the model can explain variation in

the data equally well either by including lots of events each with a small influence on niche

evolution, or by inferring few events with a large influence on niche evolution. This caused

the large number of falsely inferred events from the simulation data using our method, where

gradient speciation events deep in the phylogeny, and *in situ* adaptation events with small

amounts of environmental change, were inferred on branches along which no such events

occurred. However, simulations also suggested that inference of true events with large

influence on niche evolution were less sensitive to the assumption, and so our method still

gives accurate estimates on the environmental niche of tip species and reliable inferences on

the occurrence of major events of niche evolution. Despite possible violations of these

assumptions in the real world, the *Acacia* case study demonstrated that our model was able to

capture occurrences of important events in niche evolution and give sensible estimation of the

environmental niche of tip species.

Two additional assumptions of our method are a constant dispersal rate and a constant

adaptation rate across the phylogeny. Simulations showed that variation in adaptation rate

does not influence the power of our method to infer the history of niche evolution, but it

reduces the accuracy of our method to estimate the fundamental niche of tip species. These

assumptions can be relaxed, for example, by sampling these rates as random variables in

MCMCs. However, whether treating adaptation rate as a random variable can improve the

performance of our method requires further investigation. In particular, when both adaptation

rate and the number of events of niche evolution are free to change on each branch, the model

becomes unidentified because the amount of niche evolution depends on the product of

adaptation rate and the number of events. This problem is similar to allowing the molecular

evolution rate to change freely in molecular dating (Rannala 2002). In molecular dating the

solution is to add calibrations on divergence dates (Ronquist et al. 2012). Similarly, we can

add other types of input data in addition to species distribution data to our combined analysis

of niche modelling and niche evolution.

Four types of data may be used to 'calibrate' the analysis. First, information on

paleoclimate is useful for adjusting priors for dates of *in situ* adaptation events. For example,

we know there were three major drying periods in Australia during the Cenozoic – in this

paper, we used this as an independent source of historical data against which to test the

inferences from our model. However, for studies aiming to make inferences about the way

species adapt over time, this kind of *a priori* information on changing climate could be used

to inform the analysis, for example by increasing the prior probability of the occurrence of *in*

situ adaptation events during the evolution of drought tolerance in these periods. Second,

evidence on the tolerance range of tip species can be used as another source of data to inform

the fundamental niche of the species. For example, if we know the absolute tolerance limits

of a species, we can add another term in the likelihood function to calculate the probability

that the species cannot survive under conditions that are more extreme than the known

tolerance limits. Third, species presence/absence not only records the environmental

conditions where the species can or cannot live, it also gives co-distribution patterns between

different species and the existence of potential barriers to dispersal, for example if a species

is present only on one side of a mountain range (Morales-Castilla et al. 2017). This

information can be used to inform the occurrence of gradient speciation and parameter x in

the dispersal event. It is also possible to combine our method with existing methods in

historical biogeography to reconstruct ancestral geographic distributions. For example, in

each MCMC, we sample a history of niche evolution and reconstruct the ancestral

fundamental niches. These ancestral niches can be used with a method in historical

biogeography (e.g., Quintero et al. 2015) to reconstruct ancestral realized niches under

reconstructed paleoclimates. A fourth type of data, fossil occurrence records, can then

calibrate both ancestral fundamental niches and ancestral realized niches. Methods are

available to calibrate ancestral geographic distribution with fossil records and use the

reconstructed distribution and past climate data to reconstruct the climatic niche of ancestral

lineages (Meseguer et al. 2015).

CONCLUSION

Paleoclimate, species tolerance range, species distributions, fossil records and

phylogeny are a group of tightly linked data, where each type of data carries information on

the other types of data. Our method makes it possible to use all these types of data to study

niche evolution and estimate species niche. We have shown that combining niche evolution

models with environmental niche modelling led to greater predictive power of the

environmental tolerances in contemporary species from distribution data. We have also

demonstrated that this method could reconstruct past history of niche evolution. The logical

next step is to combine our method with methods in historical biogeography towards an

integrated approach to understanding species' geographic distribution through the lens of

both niche evolution and geographic range evolution. With the exponential growth of data

and the fast development of methods in these highly linked areas of macroecology and

macroevolution, integrating data and methods in these areas is very much needed to relax

biologically unrealistic assumptions in studying complex biological problems. A critical step

towards an integrated approach is to identify common biological processes in these areas.

Speciation, adaptation, and dispersal are the three fundamental processes underlying both

niche evolution and geographic range evolution. By explicitly modelling the three processes,

our method paves a promising path to a holistic understanding of species distribution.

SOFTWARE AVAILABILITY

Source code for the method and for the simulation is available at

https://github.com/huaxia1985/Phyloniches

FUNDING

This work was supported by an Australian Research Council Discovery Project

(DP160103915) and a Centre for Biodiversity Analysis Ignition Grant.

ACKNOWLEDGEMENTS

This paper is dedicated to the memory of John le Salle: we are grateful for his support and

encouragement in this and other projects, and we will miss his infectious enthusiasm for

biodiversity data and research. We thank Joe Miller, Elisabeth Bui, John le Salle, Rebecca

Pirzl, and Lee Belbin for useful discussions. We thank Joe Miller for providing the *Acacia*

distribution data. We thank Caroline Chong for her contribution to some of the preliminary

groundwork for this project.

REFERENCES

Bennett T.H., Flowers T.J., Bromham L. 2013. Repeated evolution of salt-tolerance in

grasses. Biol. Lett. 9:20130029.

Bromham L., Hua X., Cardillo M. 2016. Detecting macroevolutionary self-destruction from

phylogenies. Syst. Biol. 65:109–127.

Bui E., Wilford J., de Caritat P. 2017. Predictive spatial modelling for mapping soil salinity at

continental scale, EGU General Assembly, Vienna, Austria. doi:

10.13140/RG.2.2.25552.20483.

Byrne M., Yeates D.K., Joseph L., Kearney M., Bowler J., Williams M.A., Cooper S.,

Donnellan S.C., Keogh J.S., Leys R., Melville J., Murphy D.J., Porch N., Wyrwoll K.H.

2008. Birth of a biome: insights into the assembly and maintenance of the Australian arid

zone biota. Mol. Ecol. 17:4398–4417.

Cang F.A., Wilson A.A., Wiens J.J. 2016. Climate change is projected to outpace rates of

niche change in grasses. Biol. Lett. 12:20160368.

Castro-Insua A., Gómez-Rodríguez C., Wiens J.J., Baselga A. 2018. Climatic niche

divergence drives patterns of diversification and richness among mammal families. *Sci.*

Rep. 8:8781.

Catullo R.A., Ferrier S., Hoffmann A.A. 2015. Extending spatial modelling of climate change

responses beyond the realized niche: estimating, and accommodating, physiological limits

and adaptive evolution. Glob. Ecol. Biogeogr. 24:1192–1202.

Crisp M.D., Cook L.G. 2013. How was the Australian flora assembled over the last 65

million years? A molecular phylogenetic perspective. Annu. Rev. Ecol. Evol. Syst.

44:303–324.

Delzon S. 2015. New insight into leaf drought tolerance. Funct. Ecol. 29:1247–1249.

Dormann C.F., Schymanski S.J., Cabral J., Chuine I., Graham C., Hartig F., Kearney M.,

Morin X., Römermann C., Schröder B., Singer A. 2012. Correlation and process in species

distribution models: bridging a dichotomy. J. Biogeogr. 39:2119–2131.

Eastman J.M., Alfaro M.E., Joyce P., Hipp A.L., Harmon L.J. 2011. A novel comparative

method for identifying shifts in the rate of character evolution on trees. Evolution

65:3578–3589.

Felsenstein J. 1985. Phylogenies and the comparative method. Am. Nat. 12:1.

Fithian W., Elith J., Hastie T., Keith D.A. 2015. Bias correction in species distribution

models: pooling survey and collection data for multiple species. Methods Ecol. Evol.

6:424–438.

FitzJohn R.G. 2012. Diversitree: comparative phylogenetic analyses of diversification in R.

Methods Ecol. Evol. 3:1084–1092.

González-Orozco C.E., Pollock L.J., Thornhill A.H., Mishler B.D., Knerr N., Laffan S.W.,

Miller J.T., Rosauer D.F., Faith P.D., Nipperess D.A., Kujala H., Linke S., Butt N.,

Külheim C., Crisp M.D., Bruber B. 2016. Phylogenetic approaches reveal biodiversity

threats under climate change. Nat. Clim. Change 6:1110–1115.

Green P.J. 1995. Reversible jump Markov Chain Monte Carlo computation and Bayesian

model determination. Biometrika 82:711–732.

Hardy C.R. 2006. Reconstructing ancestral ecologies: challenges and possible solutions.

Diversity Distrib. 12:7–19.

Hutchinson G.E. 1957. Concluding remarks. Cold Spring Harb. Symp. 22:415–427.

Holt R.D. 2009. Bringing the Hutchinsonian niche into the 21st century: ecological and

evolutionary perspectives. Proc. Natl Acad. Sci. USA 106:19659–19665.

Kearney M., Porter W. 2009. Mechanistic niche modelling: combining physiological and

spatial data to predict species' ranges. Ecol. Lett. 12:334–350.

Kendall D.G. 1948. On the generalized ''Birth-and-Death'' process. Ann. Math. Statist.

19:1–15.

Kozak K., Wiens J. 2010. Accelerated rates of climatic-niche evolution underlie rapid

species diversification. Ecol. Lett. 13:1378–1389.

Losos J.B. 2001. Seeing the forest for the trees: the limitations of phylogenies in comparative

biology. Am. Nat. 177:709–727.

Maguire K.C., Shinneman D.J., Potter K.M., Hipkins V.D. 2018. Intraspecific niche models

for ponderosa pine (*Pinus ponderosa*) suggest potential variability in population-level

response to climate change. Syst. Biol. 67:965–978.

Meseguer A.S., Lobo J.M., Ree R., Beerling D.J., Sanmartin I. 2015. Integrating fossils,

phylogenies, and niche models into biogeography to reveal ancient evolutionary history:

the case of *Hypericum* (Hypericaceae). Syst. Biol. 64:215–232.

Miller J.T., Murphy, D.J., Brown G.K., Richardson D.M., González-Orozco C.E. 2011. The

evolution and phylogenetic placement of invasive Australian *Acacia* species. Diversity

Distrib. 17:848–860.

Miller J.T., Murphy D.J., Ho S.Y.W., Cantrill D.J., Seigler D. 2013. Comparative dating of

Acacia: combining fossils and multiple phylogenies to infer ages of clades with poor fossil

records. Aust. J. Bot. 61:436–445.

Mishler B.D., Knerr N., González-Orozco C.E., Thornhill A.H., Laffan S.W., Miller J.T.

2014. Phylogenetic measures of biodiversity and neo- and paleo-endemism in Australian

Acacia. Nat. Comm. 5:4473.

Moore G.M. 2013. Adaptations of Australian tree species relevant to water scarcity in the

urban forest. Arboric. Urban. For. 39:109–115.

Moore B.R., Ho□hna S., May M.R., Rannala B., Huelsenbeck J.P. 2016. Critically

evaluating the theory and performance of Bayesian analysis of macroevolutionary

mixtures. Proc. Natl. Acad. Sci. USA 113:9569–9574.

Morales-Castilla I., Davies T.J., Pearse W.D., Peres-Neto P. 2017. Combining phylogeny and

co-occurrence to improve single species distribution models. Glob. Ecol. Biogeogr.

26:740–752.

Moray C., Hua X., Bromham L. 2015. Salt tolerance is evolutionarily labile in a diverse set

of angiosperm families. BMC Evol. Biol. 15:90.

Moray C., Goolsby E., Bromham L. 2016. The phylogenetic association between salt

tolerance and heavy metal hyperaccumulation in angiosperms. Evol. Biol. 43:119–130.

Mu□nkemu□ller T., Boucher F.C., Thuiller W., Lavergne S. 2015. Phylogenetic niche

conservatism – common pitfalls and ways forward. Funct. Ecol. 29:627–639.

Owen C.L., Marshall D.C., Hill K.B.R., Simon C. 2017. How the aridification of Australia

structured the biogeography and influenced the diversification of a large lineage of

Australian cicadas. Syst. Biol. 66:569–589.

Paradis E., Schliep K. 2018. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics, bty633.

Peterson A.T. 2001. Predicting species' geographic distributions based on ecological niche

modeling. Condor 103:599–605.

Quintero I., Keil R., Jetz W., Crawford F.W. 2015. Historical biogeography using species

geographic ranges. Syst. Biol. 64:1059–1073.

Rabosky D. L. 2014. Automatic detection of key innovations, rate shifts, and density-

dependence on phylogenetic trees. PLOS ONE 9:e89543.

Rannala B. 2002. Identifiability of parameters in MCMC Bayesian inference of phylogeny.

Syst. Biol. 51:754–760.

Renner I.W., Elith J., Baddeley A., Fithian W., Hastie T., Phillips S.J., Popovic G., Warton

D.I. 2015. Point process models for presence-only analysis. Methods Ecol. Evol. 6:366–

379.

Revell L.J., Harmon L.J., Collar D.C. 2008. Phylogenetic signal, evolutionary process, and

rate. Syst. Biol. 57:591–601.

Ronquist F., Klopfstein S., Vilhelmsen L., Schulmeister S., Murray D.L., Rasnitsyn A.P.

2012. A total-evidence approach to dating with fossils, applied to the early radiation of the

Hymenoptera. Syst. Biol. 61:973–999.

Shi J.J., Rabosky D.L. 2015. Speciation dynamics during the global radiation of extant bats.

Evolution 69:1528–1545.

Smith A.B., Godsoe W., Rodríguez-Sánchez F., Wang H., Warren D. 2018. Niche estimation

above and below the species level. Trends Ecol. Evol. 34:260–273.

Sturges H.A. 1926. The choice of a class interval. J. Am. Stat. Assoc. 21:65–66.

Uyeda J.C., Harmon L.J. 2014. A novel Bayesian method for inferring and interpreting the

dynamics of adaptive landscapes from phylogenetic comparative data. Syst. Biol. 63:902–

918.

Weber M.M., Stevens R.D., Diniz-Filho J.A., Grelle C.E. 2017. Is there a correlation between

abundance and environmental suitability derived from ecological niche modelling? A

meta-analysis. Ecography 40: 817–828.

Williams K.J., Belbin L., Austin M.P., Stein J.L., Ferrier S. 2012. Which environmental

variables should I use in my biodiversity model? Int. J. Geogr. Inf. Sci. 26:2009–2047.

1015

Figure 1. Illustration of the basic model structure. Our method uses a Bayesian framework

that includes three major components: the phylogenetic comparative method (PCM), the

model of niche evolution, and the environmental niche model (ENM). The PCM component

estimates the location (*t*) of each *in situ* adaptation event and gradient speciation event on the

phylogeny, by assuming that these events occur across a phylogeny under an inhomogeneous

Poisson process with rate r_A and r_S. The distribution of these events on the phylogeny gives a

history of niche evolution that we want to infer. The niche evolution component defines how

three aspects of species niches (fundamental niche, realized niche, available niche) evolve

through all the events (*in situ* adaptation, gradient speciation, dispersal) from the root to each

tip along the phylogeny. This is done by estimating both local parameters (*n*, *p*, *x*) and global

parameters (, *d*, root niches) for all the events that describe how niches evolve. The model

infers the three niches of each tip species. The ENM component assesses the fit of the

inferred niches to the presence and/or absence locations of the tip species (species

distribution data). It also accounts for sampling bias of these locations by estimating how

included factors affect sampling bias ( ). The fit of the inferred niches to the data then

feedbacks to our inferences of the history of niche evolution and the niches of tip species.

1032

1033

1034

Figure 2. Illustration of the process-based model for niche evolution. There are three types of

events: *in situ* adaptation (a), gradient speciation (b), and dispersal (c). Each type of event

may change three aspects of niches along an environmental axis (with more extreme

environmental conditions towards the right of the axis). In each plot, the solid curve is the

niche before the event and the dashed curve is the niche after the event if any change happens

(realized niche after the event is also shaded in grey). Bold letters are parameters to be

estimated for the type of event. Each type of event is described by two parameters: dispersal

rate (*d*) and the location of barrier on the environmental axis (*x*) for a dispersal event;

adaptation rate ( ) and the number of steps of environmental change (*n*) for an *in situ*

adaptation event; adaptation rate ( ) and the location of species split on the environmental

axis (*p*) for a gradient speciation event. *n'* is the difference in the mean of the realized niche

before and after an event. Together with adaptation rate ( ), they determine change in

fundamental niche through   and , where increase in   leads to higher fitness under

extreme condition and decrease in   leads to less abrupt decline in fitness towards extreme

condition.

1050

1051

Figure 3. Performance of our method on simulated data. Eight statistics were used to assess

the performance. To assess the accuracy of our method for predicting tip species niches, we

used the R^2 and the slope of regressions of estimated α and β values of each tip species

against their true (simulated) values. To assess the power of our method to correctly identify

events of niche evolution, we used the proportion of branches with true speciation events

(True positive speciation) or true adaptation event (True positive adaptation) that were

identified as having significantly higher posterior probability of having speciation or

adaptation events than expected. To assess the false positive rate of our method, we used the

proportion of identified branches (with significantly higher posterior probability of having

speciation or adaptation events than expected) that did not have a true speciation event (False

positive speciation) or true adaptation event (False positive adaptation). The top row shows

the performance of our method under the exact assumptions of the method. The bottom row

shows the performance of our method when the assumption of constant adaptation rate is

violated.

1066

1067

Figure 4. The confidence that a branch is correctly identified to contain true speciation events

(a) and true adaptation events (b). The confidence is measured by the logarithm of the ratio in

the posterior probability of an identified branch with true events to an identified branch with

no true events. A positive value (red) suggests that the branch is more likely to be correctly

identified, that is the identified branch contains true events. A negative value (blue) suggests

that the branch is more likely to be falsely identified, that is the identified branch does not

contain true events. The confidence depends on two properties of the branch: how much

niche evolution occurred on that branch (x-axis) and how old the events are on the branch (y-

axis).

1077

1078

1079

Figure 5. Phylogenetic distribution of inferred adaptation events (a) and speciation events (b)

in the evolutionary history of *Acacia* drought tolerance. (a) The grey scale gives the average

total amount of environmental change (parameter n) in the inferred adaptation events on each

branch over the posterior samples. Each tip species is color coded according to its current

distribution in Australia. The map shows the spatial variation in the aridity index in Australia,

with lower values in more arid area. (b) The color scale gives the average amount of niche

evolution (parameter p) of inferred speciation events on each branch over the posterior

samples. Negative values suggest the lineage has lower fitness under extreme conditions than

its ancestral lineage. Positive value suggests the lineage has higher fitness under extreme

conditions than its ancestral lineage. Short branch lengths are elongated to help view the

speciation events near the tips.

1091

1092

1093

Figure 6. Phylogenetic distribution of inferred adaptation events (a) and speciation events (b)

in the evolutionary history of *Acacia* salt tolerance. (a) The grey scale gives the average total

amount of environmental change (parameter *n*) in the inferred adaptation events on each

branch over the posterior samples. Each tip species is color coded according to its current

distribution in Australia. The map shows the spatial variation in the soil electrical

conductivity index in Australia, with higher values in more salty area. The 12 known salt-

tolerant species are marked by asterisks. (b) The color scale gives the average amount of

niche evolution (parameter *p*) of inferred speciation events on each branch over the posterior

samples. Negative value suggests the lineage has lower fitness under extreme conditions than

its ancestral lineage. Positive value suggests the lineage has higher fitness under extreme

conditions than its ancestral lineage. Short branch length is elongated to help view the

speciation events near the tips.

1106

1107

1108

1109   Figure 7. Histogram of the inferred dates of *in situ* adaptation events in *Acacia* drought

1110   tolerance (solid) and salt tolerance (dashed). The relative frequency of the histogram is

1111   weighted by the amount environmental change in each event and is corrected by the number

1112   of lineages in the phylogeny during each time interval.

1113

1114

1115

APPENDICES:

Reversible-jump model:

The implementation involves six steps: 1) adding a new event to the model, 2)

removing an existing event from the model, 3) updating the location of an existing event on

the phylogeny, 4) updating parameters specific to that event, 5) updating parameters shared

among events, 6) updating the occurrence rate of an event.

1122

In steps 1-2, the acceptance probability of the addition or the removal of an event, or

of a proposed model that has one more or one fewer events than the current model, is *min*(1,

LikelihoodRatio× PriorRatio×ProposalRatio×Jacobian). *LikelihoodRatio* is the ratio of the

likelihood of the proposed model to the current model. *PriorRatio* is the ratio of the

probability, p(model|phylogeny), of the proposed model to the current model. *ProposalRatio*

is the relative probability of adding or deleting a move to the current model and we followed

arguments in Rabosky (2014, equation 6-7) to set *ProposalRatio*. *Jacobian* is included to

account for parameter transformation. We sampled values of each parameter for each event

from the same prior distribution as all the other events, so there is no parameter

transformation in our implementation and so *Jacobian*=1. For an *in situ* adaptation event,

parameter n (the number of steps of environmental change) has a discrete prior of equal

probability for any integer between one and the number of bins on the environmental axis.

For a gradient speciation event, parameter p (the location on the environmental axis where

species split) has a discrete prior with equal probability for a sequence from -0.9 to 0.9 by 0.2

increment. For the dispersal process following an *in situ* adaptation event or a gradient

speciation event, parameter x (the location of dispersal barrier on the environmental axis) has

a discrete prior with equal probability for any integer between 0 and the number of bins on

the environmental axis.

1141

1142    In steps 3-4, an existing event was randomly chosen to update. The location of the

1143 chosen event on the phylogeny was updated by either shifting the location by a small random

1144 quantity that was sampled from a uniform distribution or by moving the event up and down

1145 one branch. The frequency of the two ways of proposing a new location is arbitrarily set to

1146 10:1. The proposed location is accepted with probability

1147 $min⬚(1, LikelihoodRatio \times PriorRatio)$, where *PriorRatio* is the ratio of the prior probability of

1148 the proposed location to the current location of the chosen event. Parameter *x* and *n* or *p* for

1149 the chosen event is updated with equal probability of moving the current value to its adjacent

1150 lower or higher value in its discrete prior. For example, if the current value for *x* is 1, the

1151 proposed value can be either 0 or 2. The proposed value is accepted with probability $min⬚(1,$

1152 $LikelihoodRatio \times ProposalRatio)$, where *ProposalRatio* is included because the proposal

1153 probability is not symmetric: the proposed value can only be higher or lower than the current

1154 value, if the current value is at the minimum or maximum bound of the discrete prior.

1155

1156    In steps 5-6, each global parameter was updated, including the mean and the variance

1157 of the root available niche, parameter α and β of the root fundamental niche, dispersal rate

1158 (*d*), adaptation rate (γ), the occurrence rate of *in situ* adaptation events (r_A), and the overall

1159 occurrence rate of gradient speciation events (r_S). The mean of the root available niche is

1160 updated by adding its current value with a uniform random variable between -ε and ε, where

1161 ε is a tuning parameter, and the accepted probability is $min⬚(1, LikelihoodRatio)$. All the

1162 other parameters are strictly positive, so they are updated by multiplying the current value

1163 with $e^{\eta(U-0.5)}$, where *U* is a uniform random variable between 0 and 1, and η is a tuning

1164 parameter. The proposed value is accepted with probability $min⬚(1,$

1165 $LikelihoodRatio \times e^{\eta(U-0.5)}))$.

1166

1167   *Table S1. Known salt-tolerant Acacia species.*

1168   We identified salt-tolerant *Acacia* species from studies that did experiments on *Acacia*

1169   species over a range of salinity conditions. The reported salinity conditions are the highest

1170   salinity that a species was treated with in the studies and was able to tolerate.

Species	Salinity Conditions	Reference
A. ampliceps	600mM	1,7
A. auriculiformis	12-31 dS/m	2-3
A. cyclops	10-48 dS/m	4
A. harpophylla	30 dS/m	5
A. ligulata	> 4 dS/m	6
A. patagiata	10-48 dS/m	4
A. redolens	10-48 dS/m	4
A. salicina	> 4 dS/m	6
A. saligna	> 4 dS/m	6
A. sibilans	> 4 dS/m	6
A. stenophylla	> 4 dS/m	6
A. victoriae	> 4 dS/m	6

1171

Reference:

1. Theerawitaya C., Tisarum R., Samphumphuang T., Singh H.P., Cha-Um S., Kirdmanee

C., Takabe T. 2015. Physio-biochemical and morphological characters of halophyte

legume shrub, *Acacia ampliceps* seedlings in response to salt stress under greenhouse.

Front. Plant Sci. 6:630.

2. Miah M.A. 2013. Salt tolerances of some mainland tree species select as through nursery

screening. Pak. J. Biol. Sci. 16:945–949.

3. Patel A.D., Jadeja H., Pandey A.N. 2010. Effect of salinization of soil on growth, water

status and nutrient accumulation in seedlings of *Acacia auriculiformis* (Fabaceae). J. Plant

Nutrition 33: 914–932.

4. Craig G.F., Bell D.T., Atkins C.A., 1990. Response to salt and waterlogging stress of ten

taxa of Acacia selected from naturally saline areas of Western Australia. Aust. J. Bot.

38:619–630.

5. Reichman S.M., Bellairs S.M., Mulligan D.R. 2006. The effects of temperature and

salinity on *Acacia harpophylla* (brigalow) (Mimosaceae) germination. Rangeland J.

28:175–178.

6. Aswathappa N., Marcar N.E., Thompson L.A., 1987. Salt tolerance of Australian tropical

and subtropical Acacias. In: Turnbull J. (Ed.), Australian Acacias in Developing

Countries. ACIAR Proceedings, No. 16, pp. 70–73.