Effect of high frequency and low intensity pulsed electric field on protecting SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced cell injury via ERK pathway

Wei-Ting Chen[1,2,¶], Guan-Bo Lin[1,2,¶], Yu-Yi Kuo[1,2,¶], Chih-Hsiung Hsieh[1,2], Chueh-Hsuan Lu[1,2], Yi-Kun Sun[2], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author

E-mail: cychao@phys.ntu.edu.tw (CYC)

¶ These authors contributed equally to this work.

Abstract

Among various neurodegenerative diseases (NDDs), Alzheimer's disease (AD) is the most common form. Excessive aggregation of β-amyloid protein (Aβ) is thought to be a major cause in the pathogenesis of AD. A growing number of studies also have suggested that reactive oxygen species (ROS) play a critical role in the onset and progression of AD. The present study was aimed to investigate the neuroprotective effect of the high frequency and low intensity pulsed electric field (H-LIPEF) against hydrogen peroxide (H_2O_2) and Aβ-induced cytotoxicity in SH-SY5Y cells. The systematic study on frequency- and amplitude-dependent neuroprotective effect of pulsed electric field (PEF) was performed, and the result showed that H-LIPEF at 200 Hz conferred the best protective effect to the SH-SY5Y cells. The underlying mechanisms were confirmed to be due to the activation of extracellular signal-regulated kinase (ERK) pathway and the downstream prosurvival and antioxidant proteins. Given the fact that electric field can be modified to focus on specific area and applicable in a non-contact manner, we propose that H-LIPEF holds great potential for neuroprotection and curing the patients with NDDs. Furthermore, it is anticipated that the combination of such physical stimulation with drugs or natural compounds will be even more effective in treating NDDs.

Introduction

Neurodegenerative diseases (NDDs) are characterized by progressive deterioration of brain function, resulting from the gradual degeneration or death of neurons. Due to increased human lifespan, deaths due to NDDs have been increasing over the years, and caused significant impact on the quality of life of patients and their families. The World Health Organization (WHO) predicts that by 2040, NDDs will overtake cancer to become the second leading cause of death after cardiovascular diseases. Among various notorious NDDs, Alzheimer's disease (AD) is the most common neurodegenerative disorder. The typical symptoms of AD include the deposition of cortical plaques formed by aggregation of β-amyloid protein (Aβ) or hyperphosphorylation of Tau protein which finally formed the neurofibrillary tangles [1]. Scientists have long believed that these abnormal structures are prime suspects in damaging and killing nerve cells and cause progressive brain degeneration and deterioration of cognitive function among elderly people. However, there is currently no cure for AD or any other common type of NDDs. Existing medications can only help to control or delay the symptoms. One limitation for AD drug development is the presence of the blood-brain barrier (BBB), since more than 98% of small molecule drugs, and ~100% of large molecule drugs can not pass the BBB [2]. Therefore, effective treatment of AD remains one of the greatest challenges in modern medicine.

Our previous study used the non-invasive low intensity pulsed electric field

(LIPEF) in combination with fucoidan to prevent oxidative stress-induced mouse motor

neuron death [3]. It was found that LIPEF suppressed the hydrogen peroxide (H_2O_2)-

enhanced expression of ROCK protein and increased the phosphorylation of Akt in the

H_2O_2-treated NSC-34 cell line. In this work, the non-contact LIPEF stimulation was

employed to study human neural cell line SH-SY5Y, which was widely used as the

neurodegenerative damage model, and investigate the neuroprotective effect of LIPEF

against oxidative damage induced by H_2O_2. Besides, we also extended the

neuroprotective study to the AD disease model by Aβ protein administration.

Interestingly, it was observed that the LIPEF parameters used in the mouse motor

neuron are not effective in human neural cell line. The systematic study on frequency-

and amplitude-dependent neuroprotective effect of LIPEF was performed, and the

result found that the high frequency LIPEF (H-LIPEF) at 200 Hz conferred the best

protective effect to the SH-SY5Y cells. Since the alternating current (AC) signal has

only one frequency and intensity, it is time-consuming and painstaking to find the

specific AC electric signal to achieve certain biological effect. Besides, the specific AC

signal is definitely not enough to target complex signal transductions in a biological

process or disease treatment. Therefore, such method will be confined to a small

number of diseases which will discount its applicability. Unlike the AC signal, the

pulsed electric field (PEF) can be viewed as the composition of multiple sinusoidal

subcomponents with different frequencies and intensities simultaneously [3]. We

further propose that the multi-frequency component nature of PEF greatly facilitates

the development of new disease therapeutics to fulfill different demands.

In this study, we applied the H-LIPEF stimulation to human neural cell line SH-

SY5Y, and examined the prosurvival effect of such physical stimulation against

oxidative damage induced by H_2O_2 and Aβ-induced cytotoxicity. The results showed

that exposing the cells to H-LIPEF at 200 Hz significantly ameliorated the H_2O_2- and

Aβ-induced cytotoxicity in SH-SY5Y cells. Examination of the underlying mechanism

also showed that 200 Hz H-LIPEF could activate extracellular signal-regulated kinase

(ERK) pathway and downstream neuroprotective proteins. These findings implicate

that H-LIPEF is a promising electrotherapy, which sheds light on novel treatments for

AD or other NDDs.

Materials and methods

Experimental setup for exposure of the cells to non-contact

H-LIPEF

The LIPEF device described previously [3] was used for exposure of electric field with

various strengths and frequencies. The cells were placed between two copper flat and

parallel electrodes. Consecutive pulses with pulse duration 2 ms, different electric field

intensities and different frequencies were applied across the electrodes. Cells treated

with continuous LIPEF were kept in a humidified incubator of 5% CO_2 and 95% air at

37℃. After optimization of the pulse parameters, we used the H-LIPEF stimulation to

produce the electric field in a non-contact manner, and examined its neuroprotection

effect against the oxidative insult caused by H_2O_2 to SH-SY5Y human neural cells.

Cell culture and treatment

The SH-SY5Y cell was purchased from Bioresource Collection and Research Center

(Hsinchu, Taiwan) and maintained in MEM/F-12 mixture containing 10% fetal bovine

serum (HyClone; GE Healthcare Life Sciences) and 1% penicillin-streptomycin,

supplemented with 1 mM sodium pyruvate and 0.1 mM non-essential amino acids. Cells

were maintained in a humidified incubator composed of 5% CO_2 and 95% air at 37℃.

Cells were harvested with 0.05% trypsin−0.5 mM EDTA solution (Gibco; Thermo Fisher

Scientific, Inc.) and prepared for in vitro experiments. Cells were pretreated with H-

LIPEF for 4 h and then challenged with 500 μM H_2O_2 in the continuous administration

of H-LIPEF for another 24 h. For the AD disease model, the cells were pretreated with

H-LIPEF for 4 h and then challenged with 25 or 50 μM $A\beta_{25-35}$ protein solution in the

continuous administration of H-LIPEF for another 4 days. The Aβ stock solution was

prepared by solubilizing the Aβ$_{25-35}$ peptide (Sigma-Aldrich; Merck KGaA) in sterile

deionized water to a concentration of 1 mM and then incubated at 37℃ for 3 days to

allow self-aggregation before treatment. After the treatment, the cell viability was

determined.

Cell viability assay

Viability of SH-SY5Y cells after treatments with or without H-LIPEF was assessed by

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma-Aldrich;

Merck KGaA) assay. Briefly, the medium was replaced with MTT solution (0.5 mg/mL

in SH-SY5Y culture medium) and incubated at 37℃ for 4 h. During this period, the

ability of mitochondrial dehydrogenases to reduce the MTT to formazan crystal is an

indicator of cellular viability. Following MTT reaction, formazan dissolution was

performed by using 10% sodium dodecyl sulfate (SDS) and 0.01 M HCl. The optical

density of each well was then determined at 570 nm subtracting the background at 690

nm using an ELISA microplate reader. The cell viability was calculated based on the

intensity of formazan, and expressed as percentage of the non-treated control.

Western blot analysis

After the treatments, cells were washed with phosphate buffered saline (PBS) and then

harvested and lysed in RIPA lysis buffer (EMD Millipore) containing fresh protease

and phosphatase inhibitor cocktail (EMD Millipore). The lysates were then centrifuged,

and the supernatant was collected and used for the determination of protein

concentration. Equal amounts of protein extracts were loaded in the 10% SDS-PAGE

wells and transferred to polyvinylidene fluoride membranes. After drying for 1 h at

room temperature (RT), the membranes were probed with ROCK (Abcam, plc.), p-ERK,

Nrf2, p-CREB, Bcl-2, Bax (Cell Signaling Technology, Inc.), and GAPDH (GeneTex,

Inc.) antibodies overnight at 4°C. Subsequently, the washed membranes were incubated

with horseradish peroxidase-coupled secondary antibodies (Jackson ImmunoResearch

Laboratories, Inc.) for 1 h at RT. All the antibodies were diluted at the optimal

concentration according to the manufacturer's instructions. Finally, protein bands were

visualized with an enhanced chemiluminescence substrate (Advansta, Inc.) and

detected with the Amersham Imager 600 imaging system (GE Healthcare Life

Sciences). The images were analyzed with Image Lab software (Bio-Rad Laboratories,

Inc.). For normalization of proteins, GAPDH was used as an internal control.

Statistical analysis

161   Each data point represents the average from larger or equal to three independent

experiments and is expressed as the mean ± standard deviation. Statistical analyses were

performed using OriginPro 2015 software (OriginLab). Differences of statistical

significance were determined by a one-way analysis of variance (ANOVA), followed

by Tukey's post-hoc test. P-value < 0.05 was considered to indicate a statistically

significant difference.

Results

In vitro-applied H-LIPEF

The H-LIPEF was applied in a non-contact manner with the apparatus consisting of two

thin parallel conductive electrodes mounted on the insulating acrylic frame (Fig 1A).

The culture dishes were placed between the parallel electrodes so the cells were not in

contact with the electrodes directly. Therefore, the H-LIPEF stimulation device can

provide the electric field in a non-invasive manner, avoiding the harmful effects caused

by conduction current and Joule effect via contact electrodes. The pulsed electric signal

was from a function generator and amplified by a voltage amplifier. In the study, the

applied electric signal was characterized by a pulse train waveform (Fig 1B) with

various field intensities (1-60 V/cm) and frequencies (2-499 Hz) for optimization of the

treatment.

Fig 1. The experimental setup of H-LIPEF. (A) Image of the H-LIPEF stimulation device. The culture dishes were placed in the stimulation device constructed of two parallel electrodes. The blue arrows shown here indicate the schematic representation of non-invasive H-LIPEF exposure on cells. (B) Schematic representation of the pulse train waveform applied on the electrodes.

Effect of H-LIPEF exposure on H_2O_2- and Aβ−induced cytotoxicity in SH-SY5Y cells

The oxidative stress plays a critical role in the etiology and exacerbates the NDDs progression. We applied H_2O_2 to SH-SY5Y cells as an oxidative stress and looked into the neuroprotective effect of H-LIPEF technique. As shown in Fig 2A, administration of H_2O_2 on SH-SY5Y cells for 24 h decreased the cell viability in a dose-dependent manner. Treatment of SH-SY5Y cells with 500 µM H_2O_2 for 24 h reduced the cell viability to 52% compared to the control cells, which was used in the subsequent experiments. To examine whether H-LIPEF could produce protection and increase the cell viability against the oxidative insult, cells were pretreated with H-LIPEF for 4 h and then exposed to H_2O_2 in the continuous administration of H-LIPEF for another 24 h. In our previous study, the low frequency LIPEF with parameters of 2 Hz, 60V/cm,

and 2 ms pulse duration was used in the mouse motor neurons NSC-34 [3], but the

result showed that this parameter combination is ineffective in the SH-SY5Y cells. In

this study, we used H-LIPEF in our experiments, and found that the H-LIPEF

stimulation using the parameters of 200 Hz, 10 V/cm, and 2 ms pulse duration produced

significant protection against the oxidative insult caused by H_2O_2 to SH-SY5Y human

neural cell line (Fig 2B). To examine whether the frequency and intensity are both

critical parameters in the neuroprotective effect, the systematic study on frequency- and

amplitude-dependent effects was performed. The intensity of H-LIPEF was firstly fixed

at 10 V/cm and the frequency was varied from 2 Hz to 499 Hz. The upper limit was set

to 499 Hz since for 2 ms pulse duration, two pulses will overlap for frequency greater

or equal to 500 Hz. The results showed that for frequency less than or equal to 100 Hz,

their survival rates were not significantly different from the H_2O_2-treated group. At high

frequency region, the H-LIPEF conferred protective effect to the H_2O_2-treated SH-

SY5Y cells, and the maximum protective effect was at near 200 Hz (Fig 2C). For the

intensity-dependent effect, the frequency was fixed at 200 Hz, and the intensity varied

from 1~60 V/cm was examined. As shown in Fig 2D, it was obvious that there was a

maximum at 10 V/cm, indicating that the best protective effect was at 10 V/cm.

Therefore, we conclude that both the frequency and intensity are both critical

parameters in the neuroprotective effect, and the optimum protective parameters (200

Hz, 10 V/cm, and 2 ms pulse duration) were used in the following experiments. For AD

disease model, it was found that Aβ treatment reduced the viability of SH-SY5Y cells

due to the cytotoxicity of the aggregated Aβ (Fig 3A). As shown in Fig 3B and 3C, the

H-LIPEF treatment greatly improved the viability and restored the cell morphology of

Aβ-treated SH-SY5Y cells, indicating the curative effect of H-LIPEF in AD disease

model in vitro.

Fig 2. Effect of H-LIPEF on H_2O_2-induced cytotoxicity in SH-SY5Y cells. (A) Dose-

dependent curve of SH-SY5Y cells treated with different concentrations of H_2O_2 and

the cell viability was measured by MTT assay at 24 h after the H_2O_2 treatment. (B) SH-

SY5Y cells were pretreated using 200 Hz H-LIPEF with field intensity of 10 V/cm and

challenged with or without 500 μM H_2O_2. The cell viability was also measured by MTT

assay at 24 h after the H_2O_2 treatment. (C) SH-SY5Y cells were pretreated with 10 V/cm

PEF at different pulse frequencies and challenged with 500 μM H_2O_2 for 24 h and then

the cell viability was measured by MTT assay. (D) SH-SY5Y cells were pretreated

using 200 Hz PEF with different field intensities and challenged with 500 μM H_2O_2 for

24 h and then the cell viability was measured by MTT assay. Data represent the mean

± standard deviation (n=3). ***P < 0.001.

Fig 3. Effect of H-LIPEF on Aβ-induced cytotoxicity in SH-SY5Y cells. (A) The cell

viability of SH-SY5Y cells treated with 25 or 50 μM Aβ for 4 days. (B) The SH-SY5Y

cells were pretreated with H-LIPEF for 4 h and then exposed to 50 μM Aβ in the

continuous administration of H-LIPEF for another 4 days, and the cell viability was

measured by MTT assay. (C) Representative light microscopy images of SH-SY5Y

cells after Aβ and/or H-LIPEF treatment. The integrity of the cells was destructed by

Aβ, and the H-LIPEF treatment caused the protective effect and retained the cell

morphology. Scale bar = 100 μm. Data represent the mean ± standard deviation (n=3).

[***]P < 0.001.

Effect of H-LIPEF treatment on ROCK protein expression in SH-SY5Y cells

In our previous study, the neuroprotective effect of LIPEF was found to be associated

with the inhibition of the H_2O_2-induced activation of ROCK pathway in the motor

neuron cell line NSC-34 [3]. In this work, we examined the ROCK protein expressions

in order to investigate whether the protective effect of high frequency LIPEF on human

neuroblastoma SH-SY5Y cells was related to the ROCK pathway. As shown in Fig 4,

the expression of ROCK protein is slightly higher after H_2O_2 treatment but almost

unaffected after the treatment of both H-LIPEF and H$_2$O$_2$. This phenomenon points out

that although the H-LIPEF treatment can protect SH-SY5Y cells, the main mechanism

is not associated with the ROCK pathway. Therefore, other prosurvival protein

pathways should be considered in the following experiments.

Fig 4. Effect of H-LIPEF on expressions of ROCK protein in SH-SY5Y cells. (A)

Western blot analysis of ROCK protein expressions. (B) Quantification of ROCK

expression level after H-LIPEF, H$_2$O$_2$, or H-LIPEF+H$_2$O$_2$ treatment. The expression

levels were normalized to GAPDH and each relative expression level was compared

with control. Data represent the mean ± standard deviation (n=3).

Effect of H-LIPEF exposure on ERK-Nrf2/CREB signalling

pathways in H$_2$O$_2$-treated SH-SY5Y cells

We then studied the signalling pathways that could participate in the neuroprotective

mechanism of H-LIPEF stimulation. ERK is one of the major cascades of mitogen-

activated protein kinases (MAPK) family and has been known to be widely associated

with cell survival [4]. Upon phosphorylation, the active ERK proteins enter into the

nucleus where they can phosphorylate certain transcription factors, thus regulating the

expression of specific genes that contribute to cell survival [5]. Therefore, in our study,

we examined the phosphorylation of ERK and the downstream transcription factors

including cAMP response element binding protein (CREB) and nuclear factor erythroid

2-related factor 2 (Nrf2) by western blot analysis.

As shown in Fig 5A, in comparison to the untreated control cells, the level of

phosphorylated ERK (p-ERK) was increased in the cells exposed to H-LIPEF for 24 h.

In contrast, the level of p-ERK in the cells was decreased when exposed to 500 μM

H_2O_2 for 24 h without the treatment of H-LIPEF. In addition, the result showed that the

H-LIPEF treatment induced a significant recovery of the p-ERK level in the cells

exposed to 500 μM H_2O_2. This result indicates that the neuroprotection provided by H-

LIPEF could be related to the recovery of the H_2O_2-induced decline of p-ERK. In the

following, we further examined the expression levels of Nrf2 and p-CREB proteins

using western blot analysis. Nrf2 is a key transcription factor that regulates the

expression of antioxidant proteins participating in the protection of oxidative damage

triggered by injury and inflammation. As shown in Fig 5B, the level of Nrf2 was also

increased in H-LIPEF treated SH-SY5Y cells compared to the untreated cells. In

contrast, the level of Nrf2 in the cells exposed to 500 μM H_2O_2 for 24 h but without the

treatment of H-LIPEF was decreased. Meanwhile, the H-LIPEF treatment was found to

recover the expression of Nrf2 in the cells exposed to 500 μM H_2O_2 obviously. This

result showed that the neuroprotective effect was due in part to the H-LIPEF induced

Nrf2 expression and the antioxidant effect. Next, CREB is also an important

transcription factor triggering the expression of prosurvival proteins. As shown in Fig

5C, in comparison to the untreated control cells, the level of p-CREB was dramatically

increased when cells were under the H-LIPEF treatment alone. On the other hand, the

level of p-CREB in cells was decreased when exposed to 500 µM H_2O_2 for 24 h without

the treatment of H-LIPEF. Moreover, the result found that the H-LIPEF treatment

obviously recovered the expression of p-CREB in the cells exposed to 500 µM H_2O_2.

Furthermore, it is widely known that CREB phosphorylation regulates Bcl-2 family

proteins including anti-apoptotic Bcl-2 and pro-apoptotic Bax proteins, and the ratio of

Bcl-2/Bax determines whether neurons undergo survival or death after an apoptotic

stimulus. Therefore, the expressions of Bcl-2 and Bax protein levels were also

examined in the study. As shown in Fig 5D, for SH-SY5Y cells under the H-LIPEF

treatment alone, the expression levels of Bcl-2 and Bax were increased and decreased

respectively, leading to a higher elevation of the Bcl-2/Bax ratio compared to the

untreated control cells. Meanwhile, in comparison to the untreated control cells, the

level of Bcl-2/Bax ratio of SH-SY5Y cells was found to decrease when exposed to 500

µM H_2O_2, indicating that the cells had undergone apoptosis. Moreover, it was observed

that the H-LIPEF treatment notably restored the expression of Bcl-2/Bax ratio in the

cells exposed to 500 µM H_2O_2. To sum up, the results showed that the H-LIPEF

treatment induced the anti-apoptotic effect by regulating the Bcl-2 family proteins via

modulating the ERK/CREB pathways and thus promoted the cell survival against

oxidative stress.

Fig 5. Effect of H-LIPEF on ERK/Nrf2 and ERK/CREB signalling pathways and

related protein expressions. Western blot analysis of p-ERK (A), Nrf2 (B), p-CREB

(C), Bcl-2 and Bax (D) protein expressions. The expression levels were normalized to

GAPDH and each relative expression level was compared with control. Data represent

the mean $\pm$ standard deviation (n=3). ***P < 0.001 and **P < 0.01.

Discussion

The focus of this study was to investigate the neuroprotective effect of H-LIPEF

on SH-SY5Y cells. The results demonstrated that treating the cells with high frequency

(~200 Hz) and low intensity (~10 V/cm) pulsed electric field produced the most

significant protection against H_2O_2- and Aβ-induced apoptosis in SH-SY5Y human

neural cells. The western blot results showed that H-LIPEF triggered the ERK

signalling pathway and mediated the activation of CREB and Nrf2 transcription factors.

The electric stimulation had long been used in medical applications, such as wound

healing [6], muscle relaxation [7], and increasing the blood circulation [8]. However,

these treatments are often involving the electric current passing through the injury site

caused by direct contact between body and implanted electrodes. The invasive or

contact manner greatly prohibits the feasibility of the treatment or even causes harmful

effects to the cells or body such as electrolysis, burns and muscle spasms [9]. Therefore,

the electrotherapy with the non-invasive or non-contact manner will be a preferred

modality. In our previous study, we had demonstrated a non-contact electric stimulation

device [10]. The device operates as a parallel capacitor which generates electric field

between two parallel electrodes (Fig 1A). Therefore, the electric field with such

capacitive coupling mode can pass through biological samples to achieve specific

biological effect without the actual current flowing through the cells or media via

contact electrodes. It is worthy to note that the field strength passing through the culture

dishes or media is the same as the provided voltage across the electrodes. Besides, the

electric field waveform used here is a PEF, which is characterized by its multi-

frequency component nature. A pulse train signal is composed of multiple sinusoidal

subcomponents with different frequencies and intensities [3]. When the PEF enters the

biological samples, it encounters different proteins, cells and tissues with different

dielectric constants and is therefore dispersed to multiple frequencies electrically. The

multi-frequency sinusoidal waves may interact with molecules, proteins, or organelles

simultaneously to induce specific bioeffect during the PEF exposure. In our previous

studies, the non-contact LIPEF had shown promises in enhancing the activity of

anticancer agent for pancreatic cancer treatment [11] and in the neuroprotection of

mouse motor neurons [3]. In this work, we applied the LIPEF with different parameters

to the human neural cells and examined its neuroprotective effect. Unexpectedly, the

LIPEF parameter used in the mouse motor neurons NSC-34 is not effective in human

SH-SY5Y cells. Actually, scientists had found that the mouse model is quite different

from the human individuals [12]. The differences provide explanation for why some

drugs or treatments that successfully treat mice do not work in people, which have been

hampering the discovery and development of potential drugs or treatments of many

diseases. Therefore, finding treatments for animal models or cell types closer to humans

will be more important and meaningful. Here, our results showed that for human SH-

SY5Y cells, the neuroprotective effect of LIPEF was more pronounced in the high

frequency region especially near 200 Hz which was quite different from the parameter

at 2 Hz for the NSC-34 mouse cell line. For higher frequency LIPEF, it consists more

high frequency components in its composition. Therefore, the result points out that it

could be due to such high frequency components participating in the neuroprotective

effect of H-LIPEF in human SH-SY5Y neural cells. The results also showed that the

therapeutic window of the H-LIPEF intensity for SH-SY5Y cells was near 10 V/cm

which was smaller than the most effective parameter (60 V/cm) for NSC-34 cells.

Therefore, one can see that the pulse frequency and the field intensity are critical

parameters for therapeutic applications. The suitable parameters (frequency, intensity,

and pulse duration) of LIPEF could vary with different types of cells or diseases and

need to be fine-tuned and optimized. Furthermore, we propose that one may also

combine multiple pulses with different parameters to simultaneously treat the target

region.

As for the neuroprotective mechanism, it was found previously that the LIPEF

treatment obviously suppressed the H_2O_2-enhanced expression of ROCK protein in

NSC-34 cells. However, for SH-SY5Y cells, the H-LIPEF simulation did not have

significant impact on ROCK protein expression. It is thus known that the

neuroprotective mechanism is not through the ROCK pathway. In the western blot

results, the study proved that such neuroprotective effect induced by H-LIPEF was

through activation of ERK pathway. The ERK pathway plays an important role in

regulating of proliferation and differentiation as well as survival of various cell types

[13]. It transmits external cues, including growth factors into signalling cascade via the

phosphorylation of ERK. Several studies have suggested that the stimulation of ERK

pathway may be a potential way for neuroprotection [4, 5]. Here, the study showed that

the physical stimulation by H-LIPEF alone significantly enhanced the expression level

of p-ERK, indicating H-LIPEF could activate the ERK signalling pathway. Since the

ERK pathway is initiated by the binding of epidermal growth factor (EGF) to the

epidermal growth factor receptor (EGFR), we speculate that H-LIPEF could have an

interaction with EGFR and cause the conformation change of EGFR, thus triggering

the ERK pathway. Previous study had shown that magnetic field could alter the

conformation of TRPV1 ion channel, making it more favorable for capsaicin binding

[14]. The results implicate that the PEF or magnetic field with different parameters

could activate different signalling pathways. In the study, H-LIPEF with 200 Hz

frequency and 10 V/cm field intensity was found to significantly trigger the ERK

pathway and produce the neuroprotective effect. Subsequently, the phosphorylation of

ERK protein could result in the activation of its kinase activity and lead to

phosphorylation of many downstream targets including certain transcription factors

containing CREB and Nrf2 required for neuroprotection.

CREB is an important transcription factor activated by phosphorylation from ERK

or other kinases. It triggers expression of neurotrophins and antiapoptotic proteins such

as brain-derived neurotrophic factor (BDNF) and B-cell lymphoma 2 (Bcl-2) [15].

Besides, CREB downregulation is related to the pathology of AD [16], and thus

increasing the CREB expression has been considered as a potential therapeutic strategy

for AD. In the study, it was observed that H-LIPEF could increase the level of p-CREB

dramatically. Moreover, the result found that the H-LIPEF treatment notably recovered

the expression of p-CREB in the cells exposed to 500 µM H_2O_2. On one hand, it is

known that CREB phosphorylation regulates Bcl-2 family proteins including anti-

apoptotic Bcl-2 and pro-apoptotic Bax proteins. The antiapoptotic protein, Bcl-2, has

been proposed to be a pivotal component for neuroprotection [17]. It suppresses

apoptosis by heterodimerizing with Bax protein and neutralizes the effects of Bax.

When Bcl-2 is present in excess, cells are protected against apoptosis. It should be also

pointed out that when Bax is excessive and the homodimers of Bax dominate, cells are

susceptible to programed cell death. Therefore, it is the ratio of Bcl-2 to Bax which

determines the fate of a cell [18]. In the results, we found that the H-LIPEF treatment

obviously restored the expression of Bcl-2/Bax ratio in the cells exposed to 500 µM

H_2O_2, indicating the prosurvival effect of H-LIPEF in SH-SY5Y cells.

Another possible mechanism for neuroprotection of the H-LIPEF stimulation may

be associated with the Nrf2 protein. Nrf2 is also an important transcription factor

activated by ERK pathway. It regulates the expression of various antioxidant proteins

and maintains their redox states in mammalian cells [19]. The results showed that the

H-LIPEF treatment obviously recovered the expression of Nrf2 in the cells exposed to

500 µM H_2O_2, indicating that the neuroprotective effect was due in part to the H-LIPEF

induced Nrf2 expression and the antioxidant effect.

In the study, the results confirmed the neuroprotective effect of H-LIPEF on SH-

SY5Y cells. Given the fact that the electric field can be modified to focus on a targeted

region, it is proposed that H-LIPEF treatment could be applied to different organs of

the human body. In particular, the non-contact manner and the multi-frequency nature

of the H-LIPEF technique will greatly reduce the risk and consuming time of the

treatment, respectively. Besides, our results also demonstrated the ability of H-LIPEF

to rescue the Aβ-induced cytotoxicity. Since the accumulation of misfolded proteins in

brain is also the typical symptom of several NDDs, the study could become an inspiring

paradigm for future exploration for optimal LIPEF conditions for the treatment of other

NDDs.

In summary, the study demonstrates for the first time that the physical stimulation

by H-LIPEF can protect SH-SY5Y human neural cells from H_2O_2- and Aβ-induced

cytotoxicity. The underlying molecular mechanism for the neuroprotective effect is

partially due to the activation of ERK pathway and the function of some downstream

proteins, such as CREB, Nrf2, and Bcl-2. Further studies are necessary to confirm the

optimal LIPEF conditions for other NDDs and the involvement of other proteins in the

neuroprotective effect. Finally, it is also advisable for combining H-LIPEF with NDD

drugs or natural compounds to attain better curative effect for NDD patients.

Funding

This work was supported by grants from Ministry of Science and Technology (MOST 108-2112-M-002-016 and 105-2112-M-002-006-MY3 to CYC) and Ministry of Education (MOE 106R880708 to CYC) of the Republic of China. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors have declared that no competing interests exist.

Acknowledgments

The authors would like to acknowledge the service provided by the Research Core Facilities 3 Laboratory of the Department of Medical Research at National Taiwan University hospital for use of flow cytometry system.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo, Chih-Yu Chao.

Formal analysis: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo, Chih-Hsiung Hsieh,

Chueh-Hsuan Lu, Yi-Kun Sun, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo.

Writing – original draft: Wei-Ting Chen, Guan-Bo Lin, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References:

1.   Ross CA, Poirier MA. Protein aggregation and neurodegenerative disease. Nat
Med. 2004; 10:S10-S17.

2.   Yang Y, Zhang X, Ye D, Laforest R, Williamson J, Liu Y, et al. Cavitation dose
painting for focused ultrasound-induced blood-brain barrier disruption. Sci Rep. 2019;
9:2840.

3.   Hsieh CH, Lu CH, Kuo YY, Lin GB, Chao CY. The protective effect of non-
invasive low intensity pulsed electric field and fucoidan in preventing oxidative stress-
induced motor neuron death via ROCK/Akt pathway. PLoS One. 2019; 14:e0214100.

4.   Mohamed SK, Ahmed AAE, Elmorsy EM, Nofal S. ERK activation by zeranol
has neuroprotective effect in cerebral ischemia reperfusion. Life Sci. 2019; 227:137-
144.

5.   Almeida RD, Manadas BJ, Melo CV, Gomes JR, Mendes CS, Grãos MM, et al.
Neuroprotection by BDNF against glutamate-induced apoptotic cell death is mediated
by ERK and PI3-kinase pathways. Cell Death Differ. 2005; 12:1329-1343.

6. Gentzkow GD, Miller KH. Electrical stimulation for dermal wound healing. Clin
Podiatr Med Surg. 1991; 8:827-841.

7. Windsor RE, Lester JP, Herring SA. Electrical stimulation in clinical practice.
Physician Sportsmed. 1993; 21:85-93.

8. Park RJ, Son H, Kim K, Kim S, Oh T. The effect of microcurrent electrical
stimulation on the foot blood circulation and pain of diabetic neuropathy. J Phys Ther
Sci. 2011; 23:515-518.

9. Antov Y, Barbul A, Mantsur H, Korenstein R. Electroendocytosis: exposure of
cells to pulsed low electric fields enhances adsorption and uptake of macromolecules.
Biophys J. 2005; 88:2206-2223.

10. Hsieh CH, Lu CH, Chen WT, Ma BL, Chao CY. Application of non-invasive low
strength pulsed electric field to EGCG treatment synergistically enhanced the inhibition
effect on PANC-1 cells. PLoS One. 2017; 12:e0188885.

11. Lu CH, Lin SH, Hsieh CH, Chen WT, Chao CY. Enhanced anticancer effects of
low-dose curcumin with non-invasive pulsed electric field on PANC-1 cells.
OncoTargets Ther. 2018; 11:4723-4732.

12. Uhl EW, Warner NJ. Mouse models as predictors of human responses:
evolutionary medicine. Curr Pathobiol Rep. 2015; 3:219-223.

13. Sun Y, Liu WZ, Liu T, Feng X, Yang N, Zhou HF. Signaling pathway of
MAPK/ERK in cell proliferation, differentiation, migration, senescence and apoptosis.
J Recept Signal Transduct. 2015; 35:600-604.

14. Chen WT, Lin GB, Lin SH, Lu CH, Hsieh CH, Ma BL, et al. Static magnetic field
enhances the anticancer efficacy of capsaicin on HepG2 cells via capsaicin receptor
TRPV1. PLoS One. 2018; 13:e0191078.

15. Finkbeiner S. CREB couples neurotrophin signals to survival messages. Neuron.
2000; 25:11-14.

16. Rosa E, Fahnestock M. CREB expression mediates amyloid β-induced basal
BDNF downregulation. Neurobiol Aging. 2015; 36:2406-2413.

17. Howard S, Bottino C, Brooke S, Cheng E, Giffard RG, Sapolsky R.
Neuroprotective effects of bcl-2 overexpression in hippocampal cultures: interactions
with pathways of oxidative damage. J Neurochem. 2002; 83:914-923.

18. Pepper C, Hoy T, Bentley DP. Bcl-2/Bax ratios in chronic lymphocytic leukaemia
and their correlation with in vitro apoptosis and clinical resistance. Br J Cancer. 1997;
76:935-938.

19. Kubben N, Zhang W, Wang L, Voss TC, Yang J, Qu J, et al. Repression of the
antioxidant NRF2 pathway in premature aging. Cell. 2016; 165:1361-1374.

Figure 1

A

parallel electrode

acrylic frame

parallel electrode

B

Pulse duration

Field strength (V/cm)

0   one cycle or period

Time

Figure 2

Figure 3

Figure 4

A

B

Figure 5