Title

A new model for the HPA axis explains dysregulation of stress hormones on the timescale

of weeks

Omer Karin, Moriya Raz, Avichai Tendler, Alon Bar, Yael Korem Kohanim, Tomer Milo, Uri

Alon*

Dept. Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel 76100

*Corresponding author, email: uri.alon@weizmann.ac.il

Abstract

Stress activates a complex network of hormones known as the Hypothalamic-Pituitary-Adrenal

(HPA) axis. The HPA axis is dysregulated in chronic stress and psychiatric disorders, but the

origin of this dysregulation is unclear and cannot be explained by current HPA models. To

address this, we developed a new mathematical model for the HPA axis that incorporates

changes in the total functional mass of the HPA hormone-secreting glands. The mass changes

are caused by the HPA hormones which act as growth factors for the glands in the axis. We

find that the HPA axis shows the property of dynamical compensation, where gland masses

adjust over weeks to buffer variation in physiological parameters. These mass changes explain

the experimental findings on dysregulation of cortisol and ACTH dynamics in alcoholism,

anorexia and postpartum. Dysregulation occurs for a wide range of parameters, and is

exacerbated by impaired glucocorticoid receptor (GR) feedback, providing an explanation for

the implication of GR in mood disorders. These findings suggest that gland-mass dynamics

may play an important role in the pathophysiology of stress-related disorders.

<u>Author Summary</u>

The HPA axis is a neuroendocrine axis that is activated in response to stressors. The classical

description of this axis includes three hormones that act in a cascade, with the final hormone

cortisol inhibiting the two upstream hormones, ACTH and CRH. This classical picture has

timescales of hours due to hormone half-lives, and cannot explain phenomena on the scale of

weeks to months associated with this axis, such as the dysregulation observed in depression,

alcohol addiction, postpartum, and other conditions. Here, we use a minimal-model approach

to add to the classical model two known interactions in which CRH and ACTH not only

regulate downstream hormones, but also act as growth factors for the cells that secrete these

hormones. This creates a physiological circuit that can maintain total cell mass and buffer

parameter changes. It has a fragility in which after prolonged stress, the total cell functional

masses grow and take weeks to return to baseline. This is sufficient to explain the specific

dynamics of hormone dysregulation found in several contexts. It also quantifies the effect of

the cortisol (glucocorticoid) receptor on resilience to prolonged stress. Our findings suggest

that interactions between hormones and cell functional mass may play an important role in

HPA axis regulation on the timescale of weeks to months.

Synopsis

The HPA axis helps the body adapt to stress, but becomes dysregulated after prolonged activation, with clinical consequences. The origin of this dysregulation is unclear. We provide a mechanism for dysregulation based on the effect of the HPA hormones as growth factors for their downstream glands.

- A mathematical model that includes gland functional mass dynamics, introduces a new slow timescale of weeks to the HPA axis; previous models had only fast timescales of hours.

- The gland masses grow during prolonged activation, providing dynamical compensation, and recover with overshoots over weeks after withdrawal of activation.

- These overshoots are sufficient to explain the observed HPA dysregulation in pathological conditions, and clarify the role of glucocorticoid receptors in resilience to prolonged stress.

Introduction

A major hormone system that responds to stress is the HPA axis (Tsigos & Chrousos, 2002; Zavala *et al*, 2019; Melmed *et al*, 2015; Hosseinichimeh *et al*, 2015). Activation of the HPA axis results in the secretion of cortisol, which has receptors in almost all cell types, and exerts widespread effects on metabolism, immunity and behavior, to help the organism cope with stress. The HPA axis is organized in a cascade of hormones (Figure 1A): physiological and psychological stresses cause secretion of CRH from the hypothalamus (H in Figure 1A). CRH causes the pituitary corticotroph cells to secrete ACTH, which in turn causes the adrenal cortex to secrete cortisol. Cortisol negatively feeds back on the secretion of the two upstream hormones.

The HPA axis is dysregulated in a wide range of physiological and pathological conditions. HPA dysregulation is measured by an assay of HPA function called the CRH test. In the CRH test, CRH is administered and cortisol and ACTH dynamics are measured for a few hours (Figure 1B). Major depression is associated with a blunted (reduced) ACTH response (Gold *et al*, 1986b; Bardeleben & Holsboer, 1989; Holsboer *et al*, 1984) in the CRH test (Figure 1C), as well as with elevated baseline cortisol (Murphy, 1991). Elevated cortisol and blunted ACTH responses are also observed in other conditions that involve prolonged HPA axis activation, including anorexia nervosa (Gold *et al*, 1986a), alcohol abuse disorder (von Bardeleben *et al*, 1989) (Figure 1D), and pregnancy (Magiakou *et al*, 1996).

In these conditions, one can see three stages of HPA axis dysregulation after HPA over-activation stops (e.g. by weight normalization after anorexia, cessation of alcohol consumption, and childbirth respectively). The stages are defined in Figure 1D. The first stage occurs right after the over-activation stops. In this early withdrawal stage, ACTH is blunted and cortisol is high. In the second stage of intermediate withdrawal, which occurs 2-6 weeks after resolution of HPA over-activation, cortisol returns to baseline levels but ACTH remains blunted. This ACTH blunting may be causal for some of the clinical aspects of these conditions, since ACTH is co-regulated with other peptides that modulate pain and mood (Adinoff *et al*, 2005; Guillemin *et al*, 1977; Vale *et al*, 1981; Racz *et al*, 2008; Peciña *et al*, 2019; Marrazzi & Luby, 1986). In the third stage, months after withdrawal, both ACTH and cortisol normalize (for alcohol abuse disorder this time-point was not measured in (von Bardeleben *et al*, 1989)) (Figure 1D).

76
Figure 1. After prolonged stress, ACTH response is blunted for weeks after cortisol response normalizes. (A) Schema of the classic
HPA axis. CRH causes the secretion of ACTH and cortisol. (B) In the CRH test, the secretion of these hormones is measured after CRH
administration. (C) Patients suffering from major depressive disorder (MDD) show a blunted response to CRH (black line), compared with
control (gray line) - data from (von Bardeleben *et al*, 1988). (D) Patients suffering from anorexia and admitted to treatment show a blunted
ACTH response and hypercortisolemia, which resolves within 6-24 months after weight normalization- data from (Gold *et al*, 1986a).
However, 3-4 weeks after weight normalization, cortisol dynamics are normal whereas ACTH dynamics are blunted. Pregnancy is associated
with elevated cortisol levels due to CRH secretion by the placenta. 3 weeks after delivery, cortisol levels and dynamics return to normal,
whereas ACTH dynamics are blunted- data from (Magiakou *et al*, 1996). After 12 weeks, ACTH dynamics normalize as well. Individuals
recovering from alcohol abuse show hypercortisolemia and blunted ACTH response after admission – data from (von Bardeleben *et al*, 1989).
After 2-6 weeks, these individuals show normal cortisol dynamics, but blunted ACTH responses persist. In all panels, patient data are denoted
by thin gray line, and controls by a thicker black line.

The dysregulation of the HPA axis is not understood by the existing mechanistic models
of the HPA axis. The models cannot show the middle withdrawal phase with persistent blunting
of ACTH responses despite the resolution of hyper-cortisolemia. This is because in current
models, the dynamics of ACTH are strongly associated with the dynamics of cortisol, and so
once cortisol normalizes, so should ACTH, within minutes to hours. Likewise, current models
cannot explain how a deficient ACTH response produces a normal cortisol response, given that
ACTH is the main regulator of cortisol secretion. Explaining this dysregulation requires a

process on the scale of weeks that decouples the dynamics of ACTH and cortisol. This
timescale cannot be readily explained by existing models of the HPA axis, where the relevant
timescale is the lifetime of hormones, which is minutes to hours (Bingzheng *et al*, 1990). One
important process with potentially a timescale of weeks is epigenetic regulation of the
sensitivity of the cortisol receptor GR (glucocorticoid receptor) (McGowan *et al*, 2009;
Anacker *et al*, 2011; Watkeys *et al*, 2018; Cohen *et al*, 2012, 20; Turner *et al*, 2010; Schaaf &
Cidlowski, 2002). This process, however, does not break the association between ACTH and
cortisol, and cannot explain, on its own, the observed dysregulation.

Here we provide a mechanism for HPA axis dysregulation, and more generally for HPA
dynamics on the scale of weeks. To do so, we add to the classic model two additional
interactions which are experimentally characterized but have not been considered on the
systems level. These are the interactions in which the HPA hormones act as the primary growth
factors for the cells in their downstream glands. This causes the functional mass of the HPA
glands to change over time, where by 'functional mass of a gland' we mean the total capacity
of the cells for the secretion of a hormone. A large body of research, beginning with Hans Selye
in the 1930s, showed that the mass and number of adrenal cortisol-secreting cells increases
under stress. Subsequent studies established the role of ACTH as the principle regulator of the
functional mass of the adrenal cortex (Swann, 1940; Lotfi & de Mendonca, 2016). Imaging
and post-mortem studies also showed that adrenal mass increases in humans suffering from
major depression (Nemeroff *et al*, 1992; Amsterdam *et al*, 1987; Dorovini-Zis & Zis, 1987;
Dumser *et al*, 1998; Szigethy *et al*, 1994; Rubin *et al*, 1996; Ludescher *et al*, 2008), and returns
to its original size after remission (Rubin *et al*, 1995).

Similarly, CRH causes the growth of pituitary corticotrophs that secrete ACTH.
Prolonged administration of CRH, or a CRH-secreting tumor, leads to increases in corticotroph
cell mass (Gertz *et al*, 1987; Westlund *et al*, 1985; Carey *et al*, 1984; Horvath, 1988;
Schteingart *et al*, 1986; O'Brien *et al*, 1992; Asa *et al*, 1992) as well as ACTH output (Bruhn
et al, 1984; Young & Akil, 1985). Adrenalectomy, which removes negative feedback inhibition
from the HPA axis, shows similar effects in rodents (Bruhn *et al*, 1984; Westlund *et al*, 1985;
Gulyas *et al*, 1991; McNicol *et al*, 1988), and leads to increased proliferation of corticotrophs
(Gulyas *et al*, 1991) which is potentiated by CRH treatment.

Changes in functional masses can occur by hypertrophy (enlarged cells) and/or
hyperplasia (more cells); the exact mechanism does not matter for the present analysis. The
changes in functional mass take weeks, due to the slow turnover time of cell mass. Such

changes in functional mass have been shown in other hormonal axes (the insulin-glucose

system) to provide important functions, including dynamical compensation, in which gland

mass changes buffer variations in physiological parameters (Karin *et al*, 2016).

We therefore asked whether the interplay of interactions between hormones and gland

mass in the HPA axis can explain the observed dysregulation of the HPA axis on the timescale

of weeks in the pathological and physiological situations mentioned above. We also tested

other putative slow processes such as epigenetic regulation of GR (and more generally, GR

resistance), slow changes in the input signal, or changes in the removal rate of cortisol.

Our model incorporates both the hormonal interactions and the gland mass dynamics

in the HPA axis. The model incorporates cortisol feedback through the high-affinity cortisol

receptor MR (mineralocorticoid receptor) and the low-affinity receptor GR. We find that

prolonged HPA activation enlarges the functional masses of the pituitary corticotrophs and

adrenal cortex, and that the recovery of these functional masses takes weeks after stress is

removed. The dynamics of this recovery explains the observed HPA dysregulation: ACTH

responses remain blunted for weeks after cortisol has normalized. Other putative slow

processes that we tested cannot explain this dysregulation because they do not break the strong

association between ACTH and cortisol. We further show that the GR protects the HPA axis

against this dysregulation after high levels of stress, providing an explanation for the

association between deficient GR feedback and depression. Finally, we demonstrate the

physiological advantages conferred by the control of functional mass. Thus, functional mass

changes provide an integrated explanation of HPA dysregulation and dynamics on the scale of

weeks to months.

Results

<u>A new model for HPA axis dynamics that includes functional mass changes</u>

We begin by showing that the classic HPA model cannot produce the observed

dysregulation. We then add new equations for the gland masses and show that they are

sufficient to explain the dysregulation.

The classical understanding of the HPA model is described by several minimal models

(Andersen *et al*, 2013; Gupta *et al*, 2007; Sriram *et al*, 2012a; Bangsgaard & Ottesen, 2017).

These models are designed to address the timescale of hours to days, and capture HPA

dynamics on this timescale including circadian and ultradian rhythms. The input to these HPA

models is the combined effects of physical and psychological stresses, including low blood

glucose, low blood pressure, inflammation signals, psychological stressors, or effects of drugs

such as alcohol. All inputs acting at a given time-point are considered as a combined input

signal which we denote as u. The concentration of the three hormones CRH, ACTH and cortisol

are x_1, x_2, x_3. The three-hormone cascade, with feedback by x_3, is described by

165
$$\frac{dx_1}{dt} = k_1 g_1(x_3)u - w_1 x_1 \quad [1]$$

166
$$\frac{dx_2}{dt} = k_2 g_2(x_3)x_1 - w_2 x_2 \quad [2]$$

167
$$\frac{dx_3}{dt} = k_3 x_2 - w_3 x_3 \quad [3]$$

where the hormone secretion parameters are k_1, k_2, k_3, and hormone removal rates are

w_1, w_2, w_3. Hormone half-lives, given by $\log 2 / w_i$, are minutes for x_1 and x_2, and 90 minutes

for x_3 (Bingzheng $et\ al$, 1990) (Table 1). The feedback functions $g_1(x_3)$ and $g_2(x_3)$ are Hill

equations which describe the negative effect of cortisol on secretion of x_1 and x_2 (see Methods).

These equations have a single stable steady-state solution (Andersen $et\ al$, 2013). Their

response to prolonged stress, namely a pulse of input $u(t)$ that lasts for a few weeks, shows

elevated hormones during the stress, and a return to baseline within hours after the stress is

over. Figure 2A shows a simulated CRH test, in which external CRH is added at a given time-

point. ACTH shows no blunted ACTH responses in either early, intermediate or late

withdrawal phases (Figure 2A). This behavior can also be shown analytically (Methods).

To describe HPA dynamics over weeks, we add to the classic model two interactions

between the hormones and the total *functional mass* of the cells that secrete these hormones.

We introduce two new variables, the functional mass of the corticotrophs, C(t), and the

functional mass of the adrenal cells that secrete cortisol, A(t). To focus on the role of the mass,

we separate the secretion parameters into a product of secretion per cell times the total cell.

The secretion parameter of ACTH is thus $k_2 = b_2 C$, where C is the corticotroph mass and b_2

is the rate of ACTH secretion per unit corticotroph mass. The parameter b_2 includes the

metabolic capacity of the corticotrophs, the number of CRH receptors and the total blood

volume which dilutes out ACTH. A similar equation describes the secretion parameter of

cortisol, $k_3 = b_3 A$. To isolate the effects of mass changes, we assume for simplicity that the

per-unit-biomass secretion rates b_2 and b_3 are constant, whereas A(t) and C(t) can vary with

time.

This introduces two new equations for the functional masses, which have a slow timescale

of weeks. Corticotrophs proliferate under control of x_1, and adrenal cortex cells under control

of x_2, and thus

193
$$\frac{dC}{dt} = C(k_C x_1 - w_C) \quad [4]$$

194
$$\frac{dA}{dt} = A(k_A x_2 - w_A) \quad [5]$$

Where the cell-mass production rates are $k_C x_1$ and $k_A x_2$, and the cell-mass removal rates

are w_C and w_A. The removal rates are taken from experimental data that indicate cell half–lives

of days-weeks (Swann, 1940; Westlund *et al*, 1985; Gulyas *et al*, 1991). We use half-life of 6

days for C and 12 days for A (parameters given in Table 1), but the results do not depend

sensitively on these parameters, as shown below.

The new model thus has five equations, three on the fast timescale of hours, and two on the

slow timescale of weeks. One can prove that they have a single stable steady state. We non-

dimensionalized the equations to provide hormone and cell-mass steady state levels of 1

(Methods Eq. 6-10).

Model shows HPA axis dysregulation after prolonged activation

The HPA model with gland mass changes captures the experimentally observed

dysregulation (Figure 2AB). In response to a prolonged stress input *u* of a few weeks or longer,

the adrenal gland mass A and corticotroph mass C both grow. When the pulse is over, the

glands are large (Figure 2B - early withdrawal). They gradually return to baseline, but the

corticotroph mass returns with an undershoot, dropping below baseline mass and then returning

over several weeks (Figure 2B). These qualitative properties of the dynamics can be shown

analytically (Methods), and do not depend on model parameters. These transients of gland

masses are at the core of the hormone dysregulation.

Figure 2. Model with functional mass dynamics shows ACTH blunting for weeks even after cortisol normalizes. (A) The classic model of HPA axis dynamics without gland mass dynamics produces elevated levels of stress hormones during prolonged stress. However, it does not produce blunted ACTH responses in the CRH test, and, after cessation of the stressor, all hormones return to baseline within hours. (B) To account for the control of corticotroph growth by CRH and adrenal cortex growth by ACTH, we added to classic HPA model two equations that represent the dynamics of the functional mass of corticotrophs (C) and the adrenal cortex (A). *(Inset)* Such dynamics explain, for example, the enlarged adrenals of stressed rats (inset, right) compared with control (inset, left), adapted from (Selye, 1952). The model shows the three distinct phases of HPA axis dysregulation observed in experiments. At the end of a prolonged stress period, the adrenal mass is enlarged and the corticotroph mass is slightly enlarged, which results in hypercortisolemia and blunted ACTH responses in the CRH test. After a few weeks, corticotroph mass drops below baseline, while adrenal mass is slightly enlarged, causing normal cortisol dynamics with blunted ACTH responses to the CRH test. Finally, after a few months both tissue masses return to normal, leading to normalization of both cortisol and ACTH dynamics. In all panels, simulations are of a CRH test (Methods), where 'case' is after stress and 'control' is after basal HPA axis activation, as described in Figure 3.

The changed masses of the glands affect the response of the hormones to a CRH test. In early withdrawal, cortisol is high and ACTH responses are blunted (Figure 2B). Then, for a period of several weeks in intermediate withdrawal, cortisol has returned to its original baseline but ACTH remains blunted. Finally, after several months, both cortisol and ACTH return to their original baselines. Thus, the model recapitulates the experimentally observed dynamics of Figure 1.

To understand these dynamics in detail, we plot in Figure 3 the full behavior of the functional masses and hormones during and after the pulse of input. Importantly, the qualitative

conclusions are insensitive to the precise values of the model parameters. Figure 3 shows the dynamics of corticotroph and adrenal masses (Figure 3B), as well as the hormone levels (Figure 3C) during and after a prolonged pulse of stress input. To compare to CRH tests, we also simulated a CRH test at each time-point (Figure 3D). To summarize the result of the CRH tests, we plot the ratio of the peak hormone level after CRH administration relative to a control CRH test without the prolonged stress input (Figure 3E). Blunted responses correspond to values less than 1.

Figure 3. Model dynamics of HPA axis after prolonged stress. Numerical solution of the HPA model after a prolonged pulse of input (u=4) lasting 3 months, followed by recovery to baseline input u=1(A). During the pulse, gland masses (B) increase over weeks, leading to exact adaptation of ACTH and CRH levels after a few weeks (C) despite the increased input level. After stress ends, gland masses adjust back over weeks. During this adjustment period, the HPA axis is dysregulated. (D) To model a CRH test, we add CRH to the simulation and follow the hormones over several hours. The response is defined as the maximum response to a CRH test (case) relative to the maximum response to a CRH test in steady-state with basal input conditions (control). In E, the response to a CRH test given at time *t* is shown as a function of *t*. thus, this is the predicted response to CRH tests done at different days during and after the stressor. The model (black lines) shows blunted (reduced) responses to CRH tests after the stress similar to those observed in Figure 1, as well as a mismatch between cortisol and ACTH dynamics that develops a few weeks after cessation of stress.

One can see two phases during the stress pulse. The initial phase occurs after the onset of the stressor and before adaptation to the stressor (Marked ONSET in Figure 3). It lasts several weeks. In this phase, the increase in input *u* causes elevated levels and responses of CRH, ACTH and cortisol (Figure 3C). However, over weeks of stress input, the corticotroph and adrenal masses grow (Figure 3B). These gland masses thus effectively adjust to the stressor, as example of the more general phenomenon of dynamical compensation in physiological systems (Karin *et al*, 2016). The mass growth causes a return to baseline of hypothalamic CRH and ACTH, due to negative feedback by cortisol. More precisely, a larger adrenal functional mass means that less ACTH is needed to produce a concentration of cortisol that drives ACTH down to baseline.

Such a return to baseline is called *exact adaptation*. Exact adaptation is a robust feature

of this circuit due to a mathematical principle in the functional mass equations, Eq. 4,5, called

integral feedback (Karin *et al*, 2016) (Methods). The only steady-state solution of Eq. 4,5 is

that the hormones x_1 and x_2 balance proliferation and growth parameters. Exact adaptation does

not occur in models without the effects of functional mass changes - the hormones do not adapt

to the stressor (Figure S1, Supplementary Information 1).

The enlarged functional masses results in elevated cortisol levels during the stress

period, but in adapted (that is, baseline) levels of CRH, ACTH, and blunted responses of CRH

and ACTH to inputs (Figure 3E). During prolonged stress there is thus a transition from an

elevated to a blunted response of ACTH that occurs due to changes in functional masses, and

results from cortisol negative feedback.

At the end of the prolonged stress pulse, the Early Withdrawal (or EW) phase, the

functional masses are abnormal, and take weeks to months to recover. This fundamental

process is the reason for the hormonal dysregulation that is the subject of this study. In the first

weeks after the stressor is removed, the adrenal and corticotroph functional masses shrink,

accompanied by dropping cortisol levels. ACTH responses are blunted, and blunting may even

worsen over time.

Then, cortisol and CRH levels and responses simultaneously normalize. This marks the

beginning of the next phase, Intermediate Withdrawal (IW, Figure 3). In this phase, ACTH

responses remain blunted, despite the fact that cortisol is back to baseline, because adrenal

functional mass is enlarged, and corticotroph functional mass is deficient. Finally, over time,

the entire dynamics of the HPA axis normalize (Late Withdrawal, LW in Figure 3), and the

system has fully recovered.

These recovery phases are robust features of the HPA model with mass dynamics. After

withdrawal of the stressor, cortisol and CRH levels and dynamics recover *together*, before the

recovery of ACTH. This order occurs regardless of parameter values such as turnover times of

the tissues (proof in Methods). The intuitive reason for this is that before CRH returns to

baseline, the growth rate of the pituitary corticotrophs is negative, preventing ACTH from

returning to baseline.

We conclude that the gland-mass model is sufficient to explain the dynamics of

recovery from chronic HPA activation in the conditions mentioned in the introduction –

anorexia, alcohol addiction, and pregnancy (Figure 3). In order to explain the timescales of

recovery, the only model parameters that matter are the tissue turnover times. Good agreement

is found with turnover times on the scale of 1-3 weeks for the corticotrophs and adrenal cortex

cells (see Figure S2 for comparison of different turnover times, Supplementary Information 2).

The model therefore explains how ACTH responses remain blunted despite normalization of

cortisol baseline and dynamics.

We also tested several alternative mechanisms with a slow timescale of weeks. We

tested models with constant gland masses and the following processes to which we assigned

time constants on the order of one month: GR resistance following HPA activation (Figure S1,

purple lines), slow changes in input signal (Figure S1, blue lines), and slow changes in cortisol

removal rate (Figure S1, gray lines). None of these models shows the dysregulation that we

consider. The reason is that these slow processes do not cause a mismatch between ACTH and

cortisol needed to capture the blunting of ACTH despite normal cortisol responses during

intermediate withdrawal.

<u>Deficient GR feedback exacerbates HPA dysregulation following prolonged stress</u>

We next considered the role of glucocorticoid receptor (GR) in recovery from

prolonged HPA activation. GR mediates the negative feedback of cortisol on CRH and ACTH

secretion. Impaired feedback by GR is observed in many cases of depression. For example,

administration of dexamethasone, which binds the GR in the pituitary, fails to suppress cortisol

secretion in the majority of individuals suffering from depression (Coppen *et al*, 1983).

Reduced expression of GR and impaired GR function in people with depression was also

demonstrated in *post mortem* brains (López *et al*, 1998; Pandey *et al*, 2013; Webster *et al*,

2002; McGowan *et al*, 2009) and in peripheral tissues (Pariante, 2004). The feedback strength

of the GR is regulated epigenetically, and is affected by early-life adversity (Weaver *et al*,

2004; McGowan *et al*, 2009).

The relation between depression and impaired GR function seems paradoxical, since

GR signaling mediates many of the detrimental effects associated with high cortisol levels such

as hippocampal atrophy (Sapolsky *et al*, 1985). One explanation for the association between

impaired GR feedback and depression is that impaired GR feedback leads to failure of the HPA

axis to terminate the stress response on the timescale of hours, leading to excessive cortisol

levels (Anacker *et al*, 2011; De Kloet *et al*, 2005). GR feedback also plays a role in ultradian

and circadian rhythms in the HPA axis (Sriram *et al*, 2012b; Walker *et al*, 2010; Stavreva *et*

al, 2009).

Here we present an additional explanation for the association between GR feedback

and depression, based on the functional mass dynamics model. Model simulations show that

negative feedback by GR protects the HPA axis against large changes in hormone levels and

responses on the timescale of weeks after a prolonged stress input (Figure 4A). The impaired

dynamics of cortisol and ACTH is reduced when GR affinity is high. In other words, dysregulation is more severe the weaker the feedback from GR.

Figure 4. GR provides resilience to the HPA axis against prolonged stressors. Here we show the dynamics of the HPA axis during and after a prolonged pulse of stress, described in Figure 3, for K_{GR}=2 (strong feedback), K_{GR}=4 (moderate feedback), and K_{GR}=8 (weak feedback). Stronger feedback from the GR attenuates the dysregulation of all HPA axis hormones.

The reason for this effect is as follows. After an increase in stress levels ($u_1 \rightarrow u_2$), the adrenal gland mass increases (Figure 4B). When GR feedback is weak ($K_{GR} \gg u_2$), the adrenal increases by about a factor of u_2/u_1. However, if GR feedback is strong ($K_{GR} \ll u_2$), the adrenal increases to a smaller extent, because less cortisol is required to inhibit ACTH to the level required for precise adaptation. The smaller adrenal mass means a smaller dysregulation of cortisol and ACTH after stress. Strong GR feedback therefore provides resilience to the HPA axis against stress on the slow timescale.

We also tested the effects of variation in all other model parameters. We find that the hormone production and removal rates affect baseline levels but not the dynamics on the timescale of weeks. The parameters that do affect the timescale of weeks are the cell turnover times. We find that the present results are found for wide variations in these parameters (Figure S2, Supplementary Information 2).

Mass changes provide robustness and dynamic compensation to the HPA axis

Finally, we discuss potential advantages provided by the functional mass changes in

the HPA axis. The first advantage is organ size control for the corticotrophs and adrenal cortex.

The circuit provides a way for the cell populations to maintain their steady-state mass, by

balancing growth and removal. Growth and removal must be precisely balanced in order to

avoid aberrant growth or atrophy of the tissue. Thus, although the cells turn over, the feedback

through the hormones couples with the hormone trophic effects to set the masses at a functional

level.

The second beneficial feature is the robustness of the steady-state hormone levels with

respect to physiological parameters. The simple form of the equations for the masses C(t) and

A(t) (Eq. 4-5) locks the hormones CRH and ACTH into a unique steady-state determined only

by the growth and removal parameters of the tissues. Thus $CRH_{st} = \frac{w_C}{k_C}$, and $ACTH_{st} = \frac{w_A}{k_A}$.

This is remarkable because all other model parameters do not affect these steady-states.

Similarly, the cortisol baseline level, $CORT_{st} \approx \frac{k_1 k_C}{w_1 w_C} u$, depends on very few parameters. As

appropriate for a stress-response hormone, cortisol baseline depends on the input u to the

hypothalamus (averaged over weeks), which corresponds to physiological and psychological

stresses. It is independent on almost all other parameters, including the production rate per cell

(b_2, b_3) or the removal rates (w_2, w_3) of CRH and ACTH, as well as the rates of the

proliferation and removal of the adrenal cortex cells, k_A and w_A. This robustness of baseline

hormone levels is due to the ability of the functional masses to grow or shrink to buffer changes

in these parameters.

This robustness also makes the steady-state level of cortisol and ACTH *independent of*

total blood volume. This is because blood volume only enters through the production

parameters b_2, b_3. These parameters describe the secretion rate per cell, and since hormone

concentration is distributed throughout the circulation, these parameters are inversely

proportional to blood volume (k_1 relates to the portal vein system that directly connects the

hypothalamus and pituitary, and not to total blood volume (Owens & Nemeroff, 1991)). The

functional masses therefore adjust to stay proportional to blood volume.

The third feature of the present model is dynamical compensation, in which masses C

and A change to make the fast-timescale (minute-hour) response of the system to acute stress

inputs u invariant to changes in the production rates b_2, b_3 as well as several other model

parameters. A full analysis of dynamical compensation in the HPA axis is provided in the SI

(Supplementary Information 3).

<u>Discussion</u>

In this study we provide a mechanism to explain HPA axis dysregulation, based on the
effects of the hormones on the gland masses. These effects are well characterized but have not
been considered before on the systems level. We develop a minimal mathematical model that
includes gland mass changes, and show that it is sufficient to explain the observed
dysregulation following prolonged stress, alcohol abuse, anorexia and postpartum.

The blunted ACTH response which we study may have clinical implications that last
for weeks after stress is removed. One set of psychopathological implications is due to the fact
that ACTH secretion is tightly linked with the secretion of endogenous opioids such as β-
endorphin. β-endorphin is cleaved from the same polypeptide as ACTH, the protein POMC
(Guillemin *et al*, 1977; Rivier *et al*, 1982; Hargreaves *et al*, 1987). β-endorphin is secreted in
response to CRH from the anterior pituitary and from POMC-neurons in the hypothalamus. Its
secretion, like that of ACTH, is suppressed by cortisol (Lim *et al*, 1982). β-endorphin is the
endogenous ligand of the mu-opioid receptor (MOR), the primary target of addictive drugs
such as morphine and heroin. It mediates euphoria and analgesia in humans and animals
(Hawkes, 1992; Berridge & Kringelbach, 2008; Peciña *et al*, 2006; Drewnowski *et al*, 1992;
Chelnokova *et al*, 2014; Buchel *et al*, 2018). Dysregulation of β-endorphin occurs in depression
(Peciña *et al*, 2019), anorexia (Marrazzi & Luby, 1986; Marrazzi *et al*, 1990), and substance
abuse disorders (Roth-Deri *et al*, 2008; Racz *et al*, 2008; Kiefer *et al*, 2002). Our results suggest
that the dysregulation of β-endorphin, which we assume is similar to that of ACTH, persists
for many weeks after cessation of stress. It can therefore contribute to pain and anhedonia in
depression and addiction withdrawal.

The present mass model also provides insight into a physiological hallmark of several
psychiatric disorders: deficient GR feedback. Impaired GR feedback is implicated in
depression (De Kloet *et al*, 2005), and several genetic and environmental factors have been
associated with this impaired feedback (Spijker & Van Rossum, 2009; Bet *et al*, 2009).
Deficient GR feedback has been suggested to affect ultradian (Walker *et al*, 2010) and circadian
(Sriram *et al*, 2012b) HPA rhythms. We find that strong GR feedback (high GR affinity to
cortisol) can also work via functional mass dynamics on the timescale of weeks: it protects the
HPA axis from dysregulation after prolonged activation. Reduced GR feedback causes larger
dysregulation after prolonged stress. This effect is directly mediated by changes in adrenal and
corticotroph masses: strong feedback allows smaller changes of adrenal mass during the stress
period, and hence to a smaller dysregulation after stress is removed.

Since chronic stress can by itself lead to reduced GR sensitivity ("glucocorticoid
resistance") (Cohen *et al*, 2012), the present findings suggest that prolonged stress makes the
HPA less resilient to the next prolonged stress. Perhaps this reduced resilience is relevant to
the progression of depressive episodes (Kendler *et al*, 2001).

The interactions between functional masses and hormones which underlie the present
HPA axis dysregulation also provide exact adaptation to CRH and ACTH (Figure 3) during
stress. Exact adaption has been extensively studied in the context of biochemical circuits
(Ferrell Jr, 2016), but less is known about the role of exact adaptation in physiological circuits,
and, in particular, about its relevance to psychopathologies. One exception is mass changes in
the insulin-glucose system, in which exact adaptation can provide control of glucose levels and
dynamics (Karin *et al*, 2016) despite changes in insulin resistance, a form of dynamical
compensation. It would be fascinating to study what functional roles exact adaptation has in
the HPA axis, in order to understand the trade-offs that impact its dysregulation. The present
modelling approach may be able to address such questions.

From a medical point of view, if this model is correct, functional masses are potential
targets to address HPA dysregulation. Measuring these masses and their dynamics using
imaging may provide clinically relevant information. Interventions that seek to normalize
functional masses can potentially reduce the extent of dysregulation during and after prolonged
stresses. One class of interventions may use control-engineering approaches to control the
masses. Such a feedback controller can work by periodically measuring masses or hormones
and administering HPA agonists or antagonist at computed doses in order to guide the HPA
axis to desired functional masses and hormone level goals (Ben-Zvi *et al*, 2009).

Methods.

HPA axis model.

We model HPA axis using the following non-dimensionalized equations, in which hormone and cell mass steady-states are equal to 1:

$$\frac{dx_1}{dt} = w_1(g_1(x_3)u - x_1) \quad [6]$$

$$\frac{dx_2}{dt} = w_2(Cg_2(x_3)x_1 - x_2) \quad [7]$$

$$\frac{dx_3}{dt} = w_3(Ax_2 - x_3) \quad [8]$$

$$\frac{dC}{dt} = w_C C(x_1 - 1) \quad [9]$$

$$\frac{dA}{dt} = w_A A(x_2 - 1) \quad [10]$$

The cortisol feedback on CRH is through the GR and MR receptors, and on ACTH only through the GR receptor. We model the effect of the high-affinity MR receptor, $M(x_3)$, as a Michaeleis-Menten function in the saturated regime, because cortisol is thought to saturate MR at physiological levels (Andersen *et al*, 2013). Thus $M(x_3) = 1/x_3$, where the Michaelis-Menten constant is absorbed in the production term parameters. We model the cooperative, low-affinity GR receptor using a Hill function $G(x_3) = 1/(1 + \left(\frac{x_3}{K_{GR}}\right)^n)$ with n=3, where K_{GR} is GR receptor halfway-effect parameter. We model the combined receptor effects multiplicatively so that $g_1(x_3) = M(x_3)G(x_3)$, and $g_2(x_3) = G(x_3)$. All parameters are provided in Table 1. Code to simulate the equations and generate the figures is provided in https://github.com/omerka-weizmann/hpa_dynamics. The quasi-steady state of Eqs. [8-10] can be solved by setting time derivatives equal to zero, yielding at basal input u=1:

$$x_2 \approx C^{1/2}A^{-1/2} \quad [11]$$

$$x_3 \approx (CA)^{1/2} \quad [12]$$

$$x_1 \approx (CA)^{-1/2} \quad [13]$$

Alternative models.

The model without functional mass dynamics is provided by Eq. [1-3]. To this model, we added, instead of gland-mass dynamics, several alternative biological processes that have a slow timescale, potentially on the order of a month. The first process is glucocorticoid resistance, where chronically elevated cortisol levels cause weaker feedback from the GR (Cohen *et al*, 2012; Schaaf & Cidlowski, 2002). One possible mechanism involves epigenetic

470   effects such as DNA methylation (Watkeys *et al*, 2018). To model GR resistance, we added a

471   variable R that modifies the effective binding coefficient, yielding:

$$G(x_3) = \frac{1}{1 + \left(R \frac{x_3}{K_{GR}}\right)^n} \quad [14]$$

473   For R we use the following equation, based on a model of leptin resistance by Jacquier et al.

474   (Jacquier *et al*, 2015). It describes the decline of R with CORT levels:

$$\frac{dR}{dt} = w_R(f(x_3) - h(x_3)R) \quad [15]$$

For high cortisol levels to induce strong resistance, we use the simple forms: *f(x)=1, h(x)=* $1 +$

λx^2.

The second alternative slow process is a slow decrease in input, *u*. We considered an

exponentially decreasing input signal (Figure S1). The third alternative process is a putative

slow decrease in cortisol clearance rate. To model this, we added a term C_R to the removal term

in equation [8]:

$$\frac{dx_3}{dt} = w_3(A \cdot x_2 - x_3 C_R^{-1}) \quad [16]$$

483   Because there is no well-characterized biological process that governs removal on the scale of

484   weeks, we use a putative description in which cortisol reduces its own removal rate by

485   increasing C_R:

$$\frac{dC_R}{dt} = w_{C_R}(x_3 - C_R) \quad [17]$$

487   All putative processes were provided with a month timescale, by setting $w_{C_R} = w_R =$

488    $\frac{\log(2)}{30} day^{-1}$. Simulation results are shown in the SI.

489   **CRH test.**

490   We modeled the CRH test by adding the following equation for the concentration of externally

491   administrated CRH, denoted x_{1E}:

$$\frac{dx_{1E}}{dt} = w_{CRH_E}(\delta(t) - x_{1E}) \quad [18]$$

493   Where $\delta(t)$ describes a dose D of external CRH injected at T_{inj} and lasting a time W:

$$\delta(t) = \begin{cases} D & T_{inj} < t \le T_{inj} + W \\ 0 & otherwise \end{cases}$$

495   and the removal rate is w_{CRH_E} as described in (Saphier *et al*, 1992). Extrinsically administrated

496   CRH causes the pituitary to secrete ACTH, as described by adding CRH_E to the intrinsic CRH

497   (units of CRH_E are set to have equal biological effect to CRH). Thus, Eq. 7 was modified to:

$$498 \qquad \frac{dx_2}{dt} = w_2(Cg_2(x_3)(x_1 + x_{1E}) - x_2) \quad [19]$$

499 The dose D and pulse width W were calibrated to provide the observed mean CRH test results

500 in non-stressed control subjects, providing D=20 and W=30 min.

501 Proof for dysregulation of ACTH after cortisol normalization.

We briefly show how the model Eq. 6-10 provides ACTH dysregulation even after cortisol

normalizes. Consider a prolonged stressor, like the one presented in Figure 3A (that is, a pulse

increase in the input u). Since ACTH and CRH are adapt to the stressor, within hours after

withdrawal of the stressor, CRH and ACTH levels drop to below their pre-stressor baseline,

whereas cortisol remains above its pre-stressor baseline. CRH and cortisol recover

simultaneously over a few weeks because CRH dynamics depend only on cortisol and the input

u. It therefore remains to be shown that ACTH does not recover before cortisol. Since ACTH

recovers from below baseline $x_2 < 1$, it recovers with a *positive temporal derivative* $\frac{dx_2}{dt} > 0$.

If (by negation) this happens before CRH and cortisol recover, we will come to a contradiction:

since CRH is below its baseline $x_1 < 1$, C has a negative derivative due to Eq. 9; the derivative

of A is zero due to Eq 8. Because ACTH levels are approximately proportional to $C^{\frac{1}{2}}A^{-\frac{1}{2}}$ (Eq.

15) we find that ACTH has a *negative temporal derivative* when it crosses its baseline $\frac{dx_2}{dt} < 0$.

We come to a contradiction. We conclude that CRH and cortisol return to baseline before

ACTH does.

516 Cortisol can show normal fast-timescale response despite ACTH blunting.

We now use the HPA model Eq. 6-10 to show that when CRH and cortisol levels return to

baseline after stress (point IW in Figure 3), their entire dynamics in response to any fast-

timescale input (such as a CRH test) normalize, even if ACTH responses have not yet

normalize. The mass of the adrenal cortex and pituitary corticotrophs at baseline is

denoted A_0, C_0, and their size at the timepoint IW where cortisol and CRH first normalize ($x_3 =$

$x_1 = 1$) is $\lambda_A A_0, \lambda_P C_0$. Because cortisol and CRH are at baseline, and both are proportional to

the product of gland masses AC (Eq. 11-13), as noted above, one obtains $\lambda_A \lambda_C = 1$. Replacing

$\widetilde{x_2} = \lambda_C x_2$ in Eq. 6-8 yields revised equations for the fast timescale dynamics:

$$525 \qquad \frac{dx_1}{dt} = w_1(u(t) \cdot g_1(x_3) - x_1) \quad [20]$$

$$526 \qquad \frac{d\widetilde{x_2}}{dt} = \frac{dx_2}{dt} \cdot \frac{1}{\lambda_C} = \frac{1}{\lambda_C} w_2(\lambda_C C_0 x_1 g_2(x_3) - \lambda_C \widetilde{x_2}) = w_2(C_0 x_1 g_2(x_3) - \widetilde{x_2}) \quad [21]$$

$$527 \qquad \frac{dx_3}{dt} = w_3(\lambda_A A_0 x_2 - x_3) = w_3(A_0 \widetilde{x_2} - x_3) \quad [22]$$

Note that Eq. [20-21] are the same as Eq. [6-8] when $A = A_0$ and $C = C_0$, that is, they are

independent of λ_A, λ_C. In addition, the initial conditions for the hormones are also independent

of λ_A, λ_C, because $x_3 = x_1 = 1$ are at baseline, while x_2, which is proportional to $C^{\frac{1}{2}}A^{-\frac{n_1}{1+n_1}} =$

$C_0^{\frac{1}{2}}A_0^{-\frac{1}{2}} \cdot \lambda_C$, scales with λ_C, so that $\widetilde{x_2}_{ST}$ is also independent of λ_A, λ_C. We conclude that after

cortisol and CRH return to baseline, the fast timescale dynamics of cortisol and CRH to any

input (such as a CRH test) are equal to the dynamics before the stressor. In the simulations,

after CRH and cortisol have normalized for the first time, they only deviate from baseline

slightly (at most 9%), and so this consideration holds approximately for further times in the

scenario of Figure 3.

Table 1. Parameter values.

Parameter	Value
w_1	0.17/min (Andersen *et al*, 2013)
w_2	0.035/min (Andersen *et al*, 2013)
w_3	0.0091/min (Andersen *et al*, 2013)
w_C	0.099/day
w_A	0.049/day
K_{GR}	4
w_{CRH_E}	0.016/min (Saphier *et al*, 1992)
W	30 min
D	20
w_{C_E}	0.023/day
λ	1
w_R	0.023/day
n	3 (Andersen *et al*, 2013)

539
540

541   **References**

542   Adinoff B, Junghanns K, Kiefer F & Krishnan-Sarin S (2005) Suppression of the HPA axis
543   stress-response: implications for relapse. *Alcohol. Clin. Exp. Res.* **29:** 1351–1355

544   Amsterdam JD, Marinelli DL, Arger P & Winokur A (1987) Assessment of adrenal gland
545   volume by computed tomography in depressed patients and healthy volunteers: a
546   pilot study. *Psychiatry Res.* **21:** 189–197

547   Anacker C, Zunszain PA, Carvalho LA & Pariante CM (2011) The glucocorticoid receptor:
548   pivot of depression and of antidepressant treatment? *Psychoneuroendocrinology* **36:**
549   415–425

550   Andersen M, Vinther F & Ottesen JT (2013) Mathematical modeling of the hypothalamic-
551   pituitary-adrenal gland (HPA) axis, including hippocampal mechanisms. *Math. Biosci.*
552   **246:** 122–138

553   Asa SL, Kovacs K, Hammer GD, Liu B, Roos BA & Low MJ (1992) Pituitary corticotroph
554   hyperplasia in rats implanted with a medullary thyroid carcinoma cell line
555   transfected with a corticotropin-releasing hormone complementary
556   deoxyribonucleic acid expression vector. *Endocrinology* **131:** 715–720

557   Bangsgaard EO & Ottesen JT (2017) Patient specific modeling of the HPA axis related to
558   clinical diagnosis of depression. *Math. Biosci.* **287:** 24–35

559   von Bardeleben U, Heuser I & Holsboer F (1989) Human CRH stimulation response during
560   acute withdrawal and after medium-term abstention from alcohol abuse.
561   *Psychoneuroendocrinology* **14:** 441–449

562   Bardeleben U von & Holsboer F (1989) Cortisol Response to a Combined Dexamethasone-
563   Human Corticotrophin-Releasing Hormone Challenge in Patients with Depression. *J.*
564   *Neuroendocrinol.* **1:** 485–488

565   von Bardeleben U, Stalla GK, Müller OA & Holsboer F (1988) Blunting of ACTH response to
566   human CRH in depressed patients is avoided by metyrapone pretreatment. *Biol.*
567   *Psychiatry* **24:** 782–786

568   Ben-Zvi A, Vernon SD & Broderick G (2009) Model-Based Therapeutic Correction of
569   Hypothalamic-Pituitary-Adrenal Axis Dysfunction. *PLOS Comput. Biol.* **5:** e1000273

570   Berridge KC & Kringelbach ML (2008) Affective neuroscience of pleasure: reward in humans
571   and animals. *Psychopharmacology (Berl.)* **199:** 457–480

572   Bet PM, Penninx BWJH, Bochdanovits Z, Uitterlinden AG, Beekman ATF, Schoor NM van,
573   Deeg DJH & Hoogendijk WJG (2009) Glucocorticoid receptor gene polymorphisms
574   and childhood adversity are associated with depression: New evidence for a gene–
575   environment interaction. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **150B:** 660–
576   669

Bingzheng L, Zhenye Z & Liansong C (1990) A mathematical model of the regulation system
of the secretion of glucocorticoids. *J. Biol. Phys.* **17:** 221–233

Bruhn TO, Sutton RE, Rivier CL & Vale WW (1984) Corticotropin-Releasing Factor Regulates
Proopiomelanocortin Messenger Ribonucleic Acid Levels in vivo. *Neuroendocrinology*
39: 170–175

Buchel C, Miedl S & Sprenger C (2018) Hedonic processing in humans is mediated by an
opioidergic mechanism in a mesocorticolimbic system. *eLife* **7:** e39648

Carey RM, Varma SK, Drake Jr CR, Thorner MO, Kovacs K, Rivier J & Vale W (1984) Ectopic
secretion of corticotropin-releasing factor as a cause of Cushing's syndrome: a
clinical, morphologic, and biochemical study. *N. Engl. J. Med.* **311:** 13–20

Chelnokova O, Laeng B, Eikemo M, Riegels J, Løseth G, Maurud H, Willoch F & Leknes S
(2014) Rewards of beauty: the opioid system mediates social motivation in humans.
Mol. Psychiatry **19:** 746–747

Cohen S, Janicki-Deverts D, Doyle WJ, Miller GE, Frank E, Rabin BS & Turner RB (2012)
Chronic stress, glucocorticoid receptor resistance, inflammation, and disease risk.
Proc. Natl. Acad. Sci. **109:** 5995–5999

Coppen A, Abou-Saleh M, Milln P, Metcalfe M, Harwood J & Bailey J (1983) Dexamethasone
Suppression Test in Depression and other Psychiatric Illness. *Br. J. Psychiatry* **142:**
498–504

De Kloet ER, Joëls M & Holsboer F (2005) Stress and the brain: from adaptation to disease.
Nat. Rev. Neurosci. **6:** 463

Dorovini-Zis K & Zis AP (1987) Increased adrenal weight in victims of violent suicide. *Am. J.*
Psychiatry

Drewnowski A, Krahn DD, Demitrack MA, Nairn K & Gosnell BA (1992) Taste responses and
preferences for sweet high-fat foods: evidence for opioid involvement. *Physiol.*
Behav. **51:** 371–379

Dumser T, Barocka A & Schubert E (1998) Weight of adrenal glands may be increased in
persons who commit suicide. *Am. J. Forensic Med. Pathol.* **19:** 72–76

Ferrell Jr JE (2016) Perfect and near-perfect adaptation in cell signaling. *Cell Syst.* **2:** 62–67

Gertz BJ, Contreras LN, McComb DJ, Kovacs K, Tyrrell JB & Dallman MF (1987) Chronic
administration of corticotropin-releasing factor increases pituitary corticotroph
number. *Endocrinology* **120:** 381–388

Gold PW, Gwirtsman H, Avgerinos PC, Nieman LK, Gallucci WT, Kaye W, Jimerson D, Ebert
M, Rittmaster R & Loriaux DL (1986a) Abnormal hypothalamic–pituitary–adrenal
function in anorexia nervosa. *N. Engl. J. Med.* **314:** 1335–1342

Gold PW, Loriaux DL, Roy A, Kling MA, Calabrese JR, Kellner CH, Nieman LK, Post RM, Pickar
D, Gallucci W, Avgerinos P, Paul S, Oldfield EH, Cutler GB & Chrousos GP (1986b)
Responses to Corticotropin-Releasing Hormone in the Hypercortisolism of
Depression and Cushing's Disease. *N. Engl. J. Med.* **314:** 1329–1335

Guillemin R, Vargo T, Rossier J, Minick S, Ling N, Rivier C, Vale W & Bloom F (1977) beta-
Endorphin and adrenocorticotropin are selected concomitantly by the pituitary
gland. *Science* **197:** 1367–1369

Gulyas M, Pusztai L, Rappay G & Makara GB (1991) Pituitary corticotrophs proliferate
temporarily after adrenalectomy. *Histochemistry* **96:** 185–189

Gupta S, Aslakson E, Gurbaxani BM & Vernon SD (2007) Inclusion of the glucocorticoid
receptor in a hypothalamic pituitary adrenal axis model reveals bistability. *Theor.*
Biol. Med. Model. **4:** 8

Hargreaves KM, Schmidt EA, Mueller GP & Dionne RA (1987) Dexamethasone alters plasma
levels of beta-endorphin and postoperative pain. *Clin. Pharmacol. Ther.* **42:** 601–607

Hawkes CH (1992) Endorphins: the basis of pleasure? *J. Neurol. Neurosurg. Psychiatry* **55:**
247–250

Holsboer F, Von Bardeleben U, Gerken A, Stalla GK & Müller OA (1984) Blunted
corticotropin and normal cortisol response to human corticotropin-releasing factor
in depression. *N. Engl. J. Med.* **311:** 1127

Horvath E (1988) Pituitary hyperplasia. *Pathol. - Res. Pract.* **183:** 623–625

Hosseinichimeh N, Rahmandad H & Wittenborn AK (2015) Modeling the hypothalamus–
pituitary–adrenal axis: A review and extension. *Math. Biosci.* **268:** 52–65

Jacquier M, Soula HA & Crauste F (2015) A mathematical model of leptin resistance. *Math.*
Biosci. **267:** 10–23

Karin O, Swisa A, Glaser B, Dor Y & Alon U (2016) Dynamical compensation in physiological
circuits. *Mol. Syst. Biol.* **12:**

Kendler KS, Thornton LM & Gardner CO (2001) Genetic Risk, Number of Previous Depressive
Episodes, and Stressful Life Events in Predicting Onset of Major Depression. *Am. J.*
Psychiatry **158:** 582–586

Kiefer F, Horntrich M, Jahn H & Wiedemann K (2002) Is withdrawal-induced anxiety in
alcoholism based on β-endorphin deficiency? *Psychopharmacology (Berl.)* **162:** 433–
437

Lim AT, Khalid B a. K, Clements J & Funder JW (1982) Glucocorticoid and Mineralocorticoid
Effects on Adrenocorticotropin and β-Endorphin in the Adrenalectomized Rat. *J. Clin.*
Invest. **69:** 1191–1198

López JF, Chalmers DT, Little KY & Watson SJ (1998) Regulation of Serotonin1A,
Glucocorticoid, and Mineralocorticoid Receptor in Rat and Human Hippocampus:
Implications for the Neurobiology of Depression. *Biol. Psychiatry* **43:** 547–573

Lotfi CFP & de Mendonca PO (2016) Comparative effect of ACTH and related peptides on
proliferation and growth of rat adrenal gland. *Front. Endocrinol.* **7:** 39

Ludescher B, Najib A, Baar S, MacHann J, Schick F, Buchkremer G, Claussen CD & Eschweiler
GW (2008) Increase of Visceral Fat and Adrenal Gland Volume in Women with
Depression: Preliminary Results of a Morphometric Mri Study. *Int. J. Psychiatry Med.*
38: 229–240

Magiakou MA, Mastorakos G, Rabin D, Dubbert B, Gold PW & Chrousos GP (1996)
Hypothalamic corticotropin-releasing hormone suppression during the postpartum
period: implications for the increase in psychiatric manifestations at this time. *J. Clin.*
Endocrinol. Metab. **81:** 1912–1917

Marrazzi MA & Luby ED (1986) An auto-addiction opioid model of chronic anorexia nervosa.
Int. J. Eat. Disord. **5:** 191–208

Marrazzi MA, Mullings-Britton J, Stack L, Powers RJ, Lawhorn J, Graham V, Eccles T & Gunter
S (1990) Atypical endogenous opioid systems in mice in relation to an auto-addiction
opioid model of anorexia nervosa. *Life Sci.* **47:** 1427–1435

McGowan PO, Sasaki A, D'Alessio AC, Dymov S, Labonté B, Szyf M, Turecki G & Meaney MJ
(2009) Epigenetic regulation of the glucocorticoid receptor in human brain associates
with childhood abuse. *Nat. Neurosci.* **12:** 342–348

McNicol AM, Kubba MAG & McTeague E (1988) The mitogenic effects of corticotrophin-
releasing factor on the anterior pituitary gland of the rat. *J. Endocrinol.* **118:** 237-NP

Melmed S, Polonsky KS, Larsen PR & Kronenberg HM (2015) Williams textbook of
endocrinology Elsevier Health Sciences

Murphy BE (1991) Steroids and depression. *J. Steroid Biochem. Mol. Biol.* **38:** 537–559

Nemeroff CB, Krishnan KRR, Reed D, Leder R, Beam C & Dunnick NR (1992) Adrenal gland
enlargement in major depression: a computed tomographic study. *Arch. Gen.*
Psychiatry **49:** 384–387

O'Brien T, Young Jr WF, Davlla DG, Schelthauer BW, Kovacs K, Horvath E, Vale W & van
Heerden JA (1992) Cushing's syndrome associated with ectopic production of
corticotrophin-releasing hormone, corticotrophin and vasopressin by a
phaeochromocytoma. *Clin. Endocrinol. (Oxf.)* **37:** 460–467

Owens MJ & Nemeroff CB (1991) Physiology and pharmacology of corticotropin-releasing
factor. *Pharmacol. Rev.* **43:** 425–73

Pandey GN, Rizavi HS, Ren X, Dwivedi Y & Palkovits M (2013) Region-specific alterations in glucocorticoid receptor expression in the postmortem brain of teenage suicide victims. *Psychoneuroendocrinology* **38:** 2628–2639

Pariante CM (2004) Glucocorticoid Receptor Function In Vitro in Patients with Major Depression. *Stress* **7:** 209–219

Peciña M, Karp JF, Mathew S, Todtenkopf MS, Ehrich EW & Zubieta J-K (2019) Endogenous opioid system dysregulation in depression: implications for new therapeutic approaches. *Mol. Psychiatry* **24:** 576–587

Peciña S, Smith KS & Berridge KC (2006) Hedonic hot spots in the brain. *Neurosci. Rev. J. Bringing Neurobiol. Neurol. Psychiatry* **12:** 500–511

Racz I, Schürmann B, Karpushova A, Reuter M, Cichon S, Montag C, Fürst R, Schütz C, Franke PE, Strohmaier J, Wienker TF, Terenius L, Ösby U, Gunnar A, Maier W, Bilkei-Gorzó A, Nöthen M & Zimmer A (2008) The opioid peptides enkephalin and β-endorphin in alcohol dependence. *Biol. Psychiatry* **64:** 989–997

Rivier C, Brownstein M, Spiess J, Rivier J & Vale W (1982) In vivo corticotropin-releasing factor-induced secretion of adrenocorticotropin, beta-endorphin, and corticosterone. *Endocrinology* **110:** 272–278

Roth-Deri I, Green-Sadan T & Yadid G (2008) β-Endorphin and drug-induced reward and reinforcement. *Prog. Neurobiol.* **86:** 1–21

Rubin RT, Phillips JJ, McCracken JT & Sadow TF (1996) Adrenal gland volume in major depression: relationship to basal and stimulated pituitary-adrenal cortical axis function. *Biol. Psychiatry* **40:** 89–97

Rubin RT, Phillips JJ, Sadow TF & McCracken JT (1995) Adrenal gland volume in major depression. Increase during the depressive episode and decrease with successful treatment. *Arch. Gen. Psychiatry* **52:** 213–218

Saphier PW, Faria M, Grossman A, Coy DH, Besser GM, Hodson B, Parkes M, Linton EA & Lowry PJ (1992) A comparison of the clearance of ovine and human corticotrophin-releasing hormone (CRH) in man and sheep: a possible role for CRH-binding protein. *J. Endocrinol.* **133:** 487–495

Sapolsky RM, Krey LC & McEwen BS (1985) Prolonged glucocorticoid exposure reduces hippocampal neuron number: implications for aging. *J. Neurosci.* **5:** 1222–1227

Schaaf MJM & Cidlowski JA (2002) Molecular mechanisms of glucocorticoid action and resistance. *J. Steroid Biochem. Mol. Biol.* **83:** 37–48

Schteingart DE, Lloyd RV, Akil H, Chandler WF, Ibarra-Perez G, Rosen SG & Ogletree R (1986) Cushing's Syndrome Secondary to Ectopic Corticotropin-Releasing Hormone-Adrenocorticotropin Secretion∗*. *J. Clin. Endocrinol. Metab.* **63:** 770–775

Selye H (1952) The story of the adaptation syndrome.(Told in the form of informal,
illustrated lectures.). *Story Adapt. Syndr. Form Informal Illus. Lect.*

Spijker AT & Van Rossum EF (2009) Glucocorticoid receptor polymorphisms in major
depression. Focus on glucocorticoid sensitivity and neurocognitive functioning. *Ann*
NY Acad Sci **1179:** 199–215

Sriram K, Rodriguez-Fernandez M & Doyle FJ (2012a) Modeling Cortisol Dynamics in the
Neuro-endocrine Axis Distinguishes Normal, Depression, and Post-traumatic Stress
Disorder (PTSD) in Humans. *PLoS Comput. Biol.* **8:** e1002379

Sriram K, Rodriguez-Fernandez M & Iii FJD (2012b) Modeling Cortisol Dynamics in the
Neuro-endocrine Axis Distinguishes Normal, Depression, and Post-traumatic Stress
Disorder (PTSD) in Humans. *PLOS Comput. Biol.* **8:** e1002379

Stavreva DA, Wiench M, John S, Conway-Campbell BL, McKenna MA, Pooley JR, Johnson TA,
Voss TC, Lightman SL & Hager GL (2009) Ultradian hormone stimulation induces
glucocorticoid receptor-mediated pulses of gene transcription. *Nat. Cell Biol.* **11:**
1093–1102

Swann HG (1940) THE PITUITARY-ADRENOCORTICAL RELATIONSHIP. *Physiol. Rev.* **20:** 493–
521

Szigethy E, Conwell Y, Forbes NT, Cox C & Caine ED (1994) Adrenal weight and morphology
in victims of completed suicide. *Biol. Psychiatry* **36:** 374–380

Tsigos C & Chrousos GP (2002) Hypothalamic–pituitary–adrenal axis, neuroendocrine factors
and stress. *J. Psychosom. Res.* **53:** 865–871

Turner JD, Alt SR, Cao L, Vernocchi S, Trifonova S, Battello N & Muller CP (2010)
Transcriptional control of the glucocorticoid receptor: CpG islands, epigenetics and
more. *Biochem. Pharmacol.* **80:** 1860–1868

Vale W, Spiess J, Rivier C & Rivier J (1981) Characterization of a 41-residue ovine
hypothalamic peptide that stimulates secretion of corticotropin and beta-endorphin.
Science **213:** 1394–1397

Walker JJ, Terry JR & Lightman SL (2010) Origin of ultradian pulsatility in the hypothalamic–
pituitary–adrenal axis. *Proc. R. Soc. B Biol. Sci.* **277:** 1627–1633

Watkeys OJ, Kremerskothen K, Quidé Y, Fullerton JM & Green MJ (2018) Glucocorticoid
receptor gene (NR3C1) DNA methylation in association with trauma,
psychopathology, transcript expression, or genotypic variation: A systematic review.
Neurosci. Biobehav. Rev. **95:** 85–122

Weaver ICG, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S, Szyf M &
Meaney MJ (2004) Epigenetic programming by maternal behavior. *Nat. Neurosci.* **7:**
847–854

Webster MJ, Knable MB, O'grady J, Orthmann J & Weickert CS (2002) Regional specificity of
brain glucocorticoid receptor mRNA alterations in subjects with schizophrenia and
mood disorders. *Mol. Psychiatry* **7:** 985

Westlund KN, Aguilera G & Childs GV (1985) Quantification of Morphological Changes in
Pituitary Corticotropes Produced by in Vivo Corticotropin-Releasing Factor
Stimulation and Adrenalectomy. *Endocrinology* **116:** 439–445

Young E & Akil H (1985) Changes in releasability of ACTH and beta-endorphin with chronic
stress. *Neuropeptides* **5:** 545–548

Zavala E, Wedgwood KC, Voliotis M, Tabak J, Spiga F, Lightman SL & Tsaneva-Atanasova K
(2019) Mathematical modelling of endocrine systems. *Trends Endocrinol. Metab.*

765