Small non-coding RNA CjNC110 influences motility, autoagglutination, AI-2 localization, and chicken colonization in *Campylobacter jejuni*

Amanda J. Kreuder[1,2#], Brandon Ruddell[2], Kathy Mou[2*], Alan Hassall[1,2], Qijing Zhang[2], Paul J. Plummer[1,2]

[1] Department of Veterinary Diagnostic and Production Animal Medicine, College of Veterinary Medicine, Iowa State University, Ames, IA, United States

[2] Department of Veterinary Microbiology and Preventive Medicine, College of Veterinary Medicine, Iowa State University, Ames, IA, United States

#Address correspondence to Amanda J. Kreuder, akreuder@iastate.edu

* Current Address: United States Department of Agriculture, Agricultural Research Service, National Animal Disease Center, Ames, IA United States; and Oak Ridge Institute for Science and Education, Oak Ridge, TN, United States

Amanda J. Kreuder and Brandon Ruddell contributed equally to this work. Author order was determined on the basis of seniority.

Abstract

Small non-coding RNAs are involved in many important physiological functions in pathogenic microorganisms. Previous studies have identified the presence of non-coding RNAs in the major zoonotic pathogen *Campylobacter jejuni*, however, few have been functionally characterized to date. CjNC110 is a conserved ncRNA in *C. jejuni,* located downstream of the *luxS* gene which is responsible for the production of the quorum-sensing molecule autoinducer-2 (AI-2)*.* In this study, we utilized strand specific high-throughput RNAseq to identify potential targets or interactive partners of CjNC110 in a sheep abortion clone of *C. jejuni*. This data was utilized to focus further phenotypic evaluation of the role of CjNC110 in growth, motility, autoagglutination, quorum sensing, and chicken colonization in *C. jejuni*. Inactivation of the CjNC110 ncRNA led to a statistically significant decrease in autoagglutination ability as well as increased motility when compared to wild-type. Extracellular AI-2 detection was decreased in ΔCjNC110, however, intracellular AI-2 accumulation was significantly increased with a concurrent increase in LuxS expression, suggesting a key role of CjNC110 in modulating the transport of AI-2. Notably, ΔCjNC110 also showed a significant defect in the ability to colonize chickens. Complementation of CjNC110 restored all phenotypic changes back to wild-type levels. The collective results of the phenotypic and transcriptomic changes observed in our data provide valuable insights into the pathobiology of *C. jejuni* sheep abortion clone and strongly suggest that CjNC110 plays an important role in regulation of energy taxis, flagellar glycosylation, cellular communication via quorum sensing, and chicken colonization in this important zoonotic pathogen.

Introduction

Campylobacter jejuni is a major foodborne and zoonotic pathogen that causes enteritis in humans [1]. A sheep abortion (SA) clone, with IA3902 as the prototype strain, has recently emerged as the predominant cause of ovine abortion [2] and as an important pathogen in foodborne outbreaks of human gastroenteritis [3] in the United States. *C. jejuni* clone SA is also widely distributed in the cattle population in the U.S. [4]. The genome of IA3902 is quite syntenic with that of *C. jejuni* type-strain NCTC 11168 [5] although it is much more virulent than NCTC 11168 in inducing systemic infection and abortion in animals. Despite its hypervirulence, *C. jejuni* IA3902 does not harbor any virulence factors known to be associated with abortion in *C. fetus* subsp. *fetus* [6, 7]. Previously, the *luxS* gene, which mediates autoinducer-2 (AI-2) production, and the genes coding capsular polysaccharide have been identified as critical for intestinal colonization and/or translocation across gut epithelium into the bloodstream [8] [9]. Point mutations in the major outer membrane protein encoded by *porA* have also been shown to be sufficient to cause the abortion phenotype when compared to the syntenic non-abortive strain NCTC 11168 [10]. The fact that relatively mild changes in genomic sequences have led to significantly enhanced ability to cause disease by *C. jejuni* IA3902, as described above, suggests that even slight differences in sequence variations or gene regulation may have a major impact on virulence variation among differing strains of *C. jejuni*.

C. jejuni has only three known sigma factors that regulate gene transcription: σ^{70} (encoded by *rpoD*), σ^{54} (encoded by *rpoN*) and σ^{28} (encoded by *fliA*) [11]. Besides control at the transcriptional level, regulation of gene expression can also occur by post-transcriptional

modulation of mRNA translation, stability and processing, for which small non-coding RNAs are
the primary players [12-14]. Non-coding RNAs (ncRNAs) can be rapidly produced and serve to
regulate multiple different targets within a cell in a variety of ways to coordinate rapid responses
to changing environments. Thus, regulation of cellular processes by ncRNAs can provide
several advantages to the bacteria when compared to the traditional model of protein-mediated
regulation [15].
Prior to completion of the transcriptional start site map via high throughput RNA
sequencing (RNAseq) of *H. pylori* [16], the ε-proteobacteria were thought not to be capable of
using small and antisense RNA as a regulatory mechanism, partly due to a lack of the small RNA
chaperone protein Hfq [17]. Indeed, attempts using computational approaches to identify
ncRNAs in *Campylobacter* failed to yield any potential candidates, with only 3 potential loci
being identified in *Helicobacter* [18]. Recently, clear evidence that *C. jejuni* also harbors these
important regulators has been published, revealing the existence and expression of a wealth of
ncRNAs present in strains such as NCTC 11168, 81-176, 81116, RM1221 and IA3902 [19-24].
Dugar *et al.* [21], when comparing the transcriptomes of 4 different *C. jejuni* isolates, observed a
large variation in transcriptional start sites as well as expression patterns of both mRNA and
ncRNA between strains. This suggests that variations in the existence and expression of ncRNAs
even among closely related strains may play a role in differentiating virulence. Despite these
advances, identification of the functions of ncRNAs in *Campylobacter* has been slow to follow,
with most lacking functionally characterization.
The first report attempting to elucidate the role of two previously identified non-coding
RNAs in *C. jejuni* suggests these ncRNAs may play a role in flagellar biosynthesis; however, the
authors were unable to demonstrate phenotypic changes following inactivation of either non-
coding RNA [25]. A later paper, however, did establish the role of an RNA antitoxin (*cjrA*) as
the first noncoding antisense small RNA functionally characterized in *Campylobacter* to date
[26]. Beyond simply establishing the existence of non-coding RNA transcripts in
Campylobacter, there is critical need to continue to determine the physiological functions of
these potential regulators in this important zoonotic pathogen.
Our previous work has demonstrated the *in vivo* and *in vitro* expression of numerous
ncRNAs in *C. jejuni* IA3902 [24], including several that are conserved in other strains of
Campylobacter [21]. One of the expressed ncRNAs identified in our study, the conserved small
RNA CjNC110, is located in the intergenic region immediately downstream of the *luxS* gene.
This ncRNA was of particular interest to our group as previous work in our lab has already
highlighted the importance of the *luxS* gene in the virulence of *C. jejuni* [8]. In *Campylobacter*,
the *luxS* gene is known to serve two important functions, production of the quorum sensing
molecule autoinducer-2 (AI-2) and conversion of *S*-ribosylhomocysteine to homocysteine in the
active methyl cycle (AMC) [27]. Of particular note, following the identification of CjNC110, it
has also been suggested that different methods of generation of the *luxS* mutation may have led
to polar effects on this downstream ncRNA that may help explain observed differences in
phenotypes and gene expression between studies of *luxS* mutants in *Campylobacter* [28].
Based on these observations, in this study we chose to focus on elucidating the role of
CjNC110 in the pathobiology of *C. jejuni* IA3902. Specifically, we investigated the effect of
mutagenesis of either CjNC110 or *luxS* as well as combined mutation of both genes on the
transcriptional landscape during exponential and stationary stages of growth of *C. jejuni* IA3902
using high-throughput RNA sequencing (RNAseq). Using RNAseq, we successfully
demonstrated differences in the transcriptional landscape between our mutant constructs and

IA3902 wild-type. Based on the RNAseq data, we then utilized several phenotypic models to
further demonstrate that inactivation of CjNC110 affects motility, autoagglutination activity,
autoinducer-2 (AI-2) activity and chicken colonization. The work presented here provides
compelling evidence that expression of the ncRNA CjNC110 is important for the pathobiology
of *C. jejuni* IA3902.

Materials and methods

Ethics Statement

All studies involving animals were approved by the Iowa State University Institutional Animal
Care and Use Committee (IACUC) prior to initiation (1-18-8675-G) and followed all appropriate
animal care guidelines as set forth in the "Public Health Service Policy on Humane Care and Use
of Laboratory Animals" developed by the U.S. Department of Health and Human Services,
National Institutes of Health Office of Laboratory Animal Welfare. Humane euthanasia was
performed using carbon dioxide asphyxiation as recommended by the American Veterinary
Medical Association "Guidelines for the Human Euthanasia of Animals."

Bacterial strains, plasmids, primers and culture conditions

C. jejuni IA3902 was initially isolated from an outbreak of sheep abortion in Iowa during
2006 and has been utilized by our laboratory as the prototypical isolate of clone SA [2]. W7 is a
highly motile variant of the commonly utilized laboratory strain *C. jejuni* NCTC 11168 [8]. *C.*
jejuni strains and their isogenic mutants were routinely grown in Mueller-Hinton (MH) broth or
agar plates (Becton-Dickinson, Franklin Lakes, NJ) at 42°C under microaerophilic conditions
(5% O_2, 10% CO_2, 85% N_2). For the mutant strains containing a chloramphenicol resistance
cassette, 5 μg/mL chloramphenicol was added to either the broth or agar plates when appropriate.
For strains containing a kanamycin resistance cassette, 30 μg/mL kanamycin was added to either
the broth or agar plates when appropriate.
For genetic manipulations, *Escherichia coli* competent cells were grown at 37°C on
Luria-Bertani (LB) agar plates or broth (Becton-Dickinson, Franklin Lakes, NJ) with shaking at
125 rpm. When appropriate, 50 μg/mL kanamycin, 20 μg/mL chloramphenicol or 100 μg/mL
ampicillin was added to the broth or agar plates for selection of colonies. *Vibrio harveyi* strains
were grown in autoinducer broth (AB) at 30°C with shaking at 175 rpm as described previously
[29].
All strains used in this study are described in **Table 1**, with all relevant plasmids listed in
Table A in S1 Appendix and primer sequences and LNA probe sequence in **Table B in S1**
Appendix. All strains were maintained in 20% glycerol stocks at -80°C and passaged from those
stocks as needed for experimental procedures.

156   **Table 1.** Bacterial strains utilized in this study.

Strain	Description	Source or Reference
Campylobacter jejuni		
W7	Wild-type motile variant of NCTC 11168	[8]
W7 ΔCjNC110	W7 ΔCjNC110::Cm^R	This study
Sheep Abortion (SA) IA3902	Wild-type *C. jejuni*	[2]
IA3902 ΔCjNC110	IA3902 ΔCjNC110::Cm^R	This study
IA3902 ΔluxS	IA3902 ΔluxS::Kan^R	[30]
IA3902 ΔCjNC110ΔluxS	IA3902 ΔCjNC110::Cm^R ΔluxS::Kan^R	This study
IA3902 ΔCjNC110c	IA3902 ΔCjNC110::Cm^R CjNC110::Kan^R	This study
Escherichia coli		
DH5α	*fhuA2* Δ(*argF-lacZ*)U169 *phoA gln*V44 Φ80 Δ(*lacZ*)M15 *gyr*A96 *recA1 relA1 endA1 thi-1 hsdR*17	New England Biolabs, Ipswich, MA
Vibrio harveyi		
BB152	AI-1-/AI-2+; luxM::Tn5	[29]
BB170	AI-2 reporter strain; luxN::Tn5	[31]

Kan^R = kanamycin resistance cassette

Cm^R = chloramphenicol resistance cassette

Creation of *C. jejuni* ΔCjNC110 and ΔCjNC110ΔluxS mutants in IA3902

An isogenic CjNC110 mutant of *C. jejuni* IA3902 was constructed via deletional
mutagenesis utilizing a combination of synthetic double-stranded DNA (dsDNA) fragments and
traditional cloning methods. Based on previously published data depicting the proposed
transcriptional start site for CjNC110 [21], the coding region of CjNC110 in IA3902 and the
prototypical *C. jejuni* strain 11168 were first confirmed to be identical. Then, a 200 bp section of
the IA3902 genome starting 20 bp upstream and including the entire 137 nt transcript of the
CjNC110 sequence as predicted in Dugar *et al.* [21] was replaced with 820 bp of the promoter
and coding sequence of the chloramphenicol acetyltransferase (*cat*) gene of *Campylobacter coli*
plasmid C-589 [32]. Synthetic dsDNA including approximately 500 bp upstream and 500 bp
downstream of the region replaced with the *cat* cassette was then synthesized in 4 fragments of
500 bp each with overlapping homologous ends (Integrated DNA Technologies, Coralville, IA).
The Gibson Assembly method was then utilized to assemble the synthetic dsDNA fragments [33]
using the Gibson Assembly Master Mix (New England Biolabs, Ipswich, MA). Following
assembly, primers (CJnc110F2 and CJnc110R2) were designed to amplify a 1785 bp product of
the assembled dsDNA; PCR amplification was achieved using TaKaRa Ex Taq DNA
Polymerase (ClonTech, Mountain View, CA). This amplified PCR product was then cloned into
the pGEM-T Easy Vector using T4 ligase (Promega, Madison, WI), resulting in the construction
of pCjNC110::cat which was then transformed into chemically competent *E. coli* DH5α (New
England Biolabs, Ipswich, MA). Transformants were then selected on LB agar plates containing

chloramphenicol (20 μg/ml), ampicillin (100 μg/ml), and ChromoMax IPTG/X-Gal Solution
(Fisher Scientific, Pittsburgh, PA). pCjNC110::cat was purified from the transformed *E. coli*
using the QIAprep Miniprep kit (QIAGEN, Germantown, MD) and confirmed by PCR to contain
the construct again using the CJNC110F2 and CJNC110R2 primers.
The pCjNC110::cat plasmid DNA was then introduced to *C. jejuni* W7 as a suicide
vector and the deletion transferred into the genome of *C. jejuni* W7 via homologous
recombination. Transformants were selected on MH agar plates containing chloramphenicol (5
μg/ml) and deletional mutagenesis was again confirmed via PCR analysis and Sanger sequencing
to create *C. jejuni* W7ΔCjNC110. Following confirmation, natural transformation was used to
move the gene deletion into *C. jejuni* IA3902 as previously described [34] to create *C. jejuni*
IA3902ΔCjNC110. Natural transformation was again used to move the CjNC110 gene deletion
into the previously created *luxS* insertional mutant *C. jejuni* IA3902ΔluxS [8] to create the
double knockout mutant *C. jejuni* IA3902ΔCjNC110ΔluxS. Transformants were selected on MH
agar plates containing chloramphenicol (5 μg/mL) and kanamycin (30 μg/mL) and confirmed via
PCR analysis and Sanger sequencing of the CjNC110 region along with the entire upstream
(*luxS*) and downstream (CjSA_1137) genes using primers Cj1198F1 and Cj1199R3. All colonies
were screened for presence/absence of motility as described below, and only colonies with
verified motility were used for future studies. Expression of CjNC110 in wild-type and
elimination of expression in the ΔCjNC110 mutant was confirmed via northern blot analysis
using an LNA custom designed probe (QIAGEN) and 15μg of total RNA as previously described
[35].
200

Creation of *C. jejuni* ΔCjNC110 complement in IA3902

Complementation of ΔCjNC110 was achieved via insertion of the coding region into the
intergenic region of the 16S and 23S rRNA operon (*rrs-rrl*) of IA3902ΔCjNC110 via
homologous recombination using plasmid pRRK. A copy of the CjNC110 gene including 190bp
upstream of the transcription initiation location (total of 570bp) was cloned into an *XbaI* site of
pRRK located upstream of the kanamycin resistance determinant to create pCjNC110c as
previously described in our lab [36] with some modifications of the original protocol [37].   Briefly,
the CjNC110 gene and flanking regions of IA3902 were amplified using forward and reverse
primers designed with an *XBaI* restriction enzyme site (CjNC110cF1 and CjNC110cR1).
Following PCR amplification with TaKaRa Ex Taq DNA Polymerase (ClonTech, Mountain
View, CA), the PCR product and pRRK plasmid were digested with the *XBaI* restriction
endonuclease (New England Biolabs, Ipswich, MA). The digested plasmid was then incubated
twice with Antarctic Phosphatase (New England Biolabs, Ipswich, MA) to prevent self-ligation
of cut plasmid. Following ligation and transformation, transformants were selected on LB agar
plates containing chloramphenicol (20 μg/ml) and kanamycin (50 μg/ml kanamycin) yielding a
plasmid containing the construct pRRK::CjNC110 in *E. coli* DH5α. Following PCR
confirmation and Sanger sequencing to confirm preservation of correct gene sequence, the
plasmid was then utilized as a suicide vector and the gene was inserted into the intergenic region
of the 23S rRNA operon (*rrs-rrl*) of IA3902ΔCjNC110 via homologous recombination to
generate IA3902ΔCjNC110c transformants. The transformants were selected on MH agar plates
containing chloramphenicol (5 μg/ml) and kanamycin (30 μg/ml) and confirmed via PCR
analysis and Sanger sequencing to contain both the expected ΔCjNC110 deletional mutation and
ΔCjNC110c insertion, validating successful genetic complementation. All positive colonies

were screened for presence/absence of motility as described below and a motile complement was
selected to create IA3902ΔCjNC110c.
226

Growth curves

Two separate growth curves were completed in triplicate utilizing the same method as
described below with the only variation being the addition of the complement strain ΔCjNC110c
to the second growth curve. To begin the growth curve, the A_{600} of overnight cultures were
adjusted to 0.5 using sterile MH broth on a Genesys 10S VIS spectrophotometer (Thermo
Scientific™, Waltham, MA). Cultures were then diluted 1:10 for a final targeted starting A_{600} of
0.05 in 90 mL sterile MH broth and placed in a sterile 250 mL Erlenmeyer glass flask. Cultures
were incubated at 42°C under microaerophilic conditions with shaking at 125 rpm for 30 hours
with removal of samples from the flasks at designated time points [3, 6, 9, 12, 24, 27 (2^{nd} growth
curve only) and 30 hours]. For the first growth curve (which included the following strains:
IA3902 wild-type, ΔCjNC110, ΔluxS and ΔCjNC110ΔluxS), samples were processed as
described below for RNA isolation as well as assessed for A_{600} and actual colony counts using
the drop-plate method as previously described [38]. For the second growth curve (which
included ΔCjNC110c in addition to the four previous strains), samples were collected and
processed for A_{600} and as described below for assessment of AI-2 levels via the bioluminescence
assay. The A_{600} over time were statistically analyzed for the two independent growth curves,
using a two-way ANOVA with repeated measures and Dunnett's multiple comparison test
(GraphPad Prism).
245

Total RNA extraction

Broth culture samples processed for total RNA extraction were either centrifuged at 8000
x g for 2 minutes (for sequencing and qRT-PCR) or 10,000 x g for 4 minutes (Northern blot) at
4°C immediately following collection. Following pelleting of the cells, the supernatant was
decanted and 1 mL QIAzol Lysis Reagent (QIAGEN) was added to resuspend the cell pellet and
protect the RNA. QIAzol-protected cultures were then stored at -80°C for up to two months prior
to proceeding with total RNA isolation. Total RNA isolation was performed using the miRNeasy
Mini Kit (QIAGEN) followed by purification using the RNeasy MinElute Cleanup kit
(QIAGEN) as previously described [24]. RNA quality was measured using the Agilent 2100
Bioanalyzer RNA 6000 Nano kit (Agilent Technologies, Santa Clara, CA) and all RNA samples
utilized for downstream RNAseq library preparation had an RNA integrity number (RIN) of
>9.0, indicating high quality RNA. Verification of complete removal of any contaminating DNA
was performed via PCR amplification of a portion of the CjSA_1356 gene, which is unique to *C.
jejuni* IA3902, using primers SA1356F and SA1356R [39].
260

RNAseq library preparation and sequencing

The 3 hour (exponential phase) and 12 hour (stationary phase) timepoints were selected
for RNAseq analysis based on assessment of $\log_{10}$ CFU/mL (**Fig A in S3 Appendix**). 2.5 µg of
confirmed DNA-free total RNA was treated with Ribo-Zero rRNA Removal Kit for Bacteria
according to the manufacturer's instructions (Illumina, San Diego, CA). Following rRNA
removal, the ribosomal depleted total RNA was purified using the RNeasy MinElute Cleanup kit
(QIAGEN) using the same modifications as described previously [24]. Following clean-up, the
RNA quality, quantity, and rRNA removal efficiency was then analyzed via the Agilent 2100

Bioanalyzer RNA 6000 Pico kit (Agilent Technologies). Library preparation for sequencing on
the Illumina HiSeq platform was completed using the TruSeq stranded mRNA HT library
preparation kit (Illumina) with some modifications as described previously [24]. The pooled
library was then submitted to the Iowa State University DNA Facility and sequenced on an
Illumina HiSeq 2500 machine in high-output single read mode with 100 cycles.
274

Differential gene expression analysis of RNAseq data
Rockhopper (http://cs.wellesley.edu/~btjaden/Rockhopper/) was used to analyze the
differences in gene expression between strains and timepoints; the standard settings of the
program were utilized for analysis [40]. Following computational analysis via Rockhopper, a
change in gene expression was deemed significant when the Q-value (false discovery rate) was
below 5% and a >1.5 fold change in expression was observed. Visual assessment of read count
data was performed using the Integrated Genome Viewer (IGV)
(https://www.broadinstitute.org/igv/) [41, 42]. Assessment of function of the differentially
expressed genes was performed using the Clusters of Orthologous Groups (COG) [43] as
previously described in IA3902 [5]. Venn diagrams were used to depict overlap of genes
differentially regulated in multiple mutant strains and were generated using BioVenn [44]
(http://www.cmbi.ru.nl/cdd/biovenn/index.php). Kegg Pathways was used to perform metabolic
pathway analysis [45] (http://www.genome.jp/kegg/pathway.html).
288

Northern Blot
For northern blot detection of CjNC110, bacterial cells from WT, ΔCjNC110, and
ΔCjNC110c were grown to exponential phase of growth in MH broth and total RNA isolation
performed as described above. The total RNA input was normalized to 15 μg using Qubit™ BR
RNA Assay Kit (Invitrogen™, USA) for each respective strain. For northern blot detection, total
RNA from bacterial cells (15 μg) and a pre-stained DynaMarker® RNA High Ladder
(Diagnocine LLC, Hackensack, NJ) were loaded onto 1.3% denaturing agarose gel (Bio-rad,
Hercules, CA), consisting of 1x MOPS buffer (Fisher, USA) and 5% solution of 37%
formaldehyde (Fisher, USA), electrophoresed at 100V for 1h and transferred to positively-
charged nylon membrane (Roche, Indianapolis, IN) using a Vacuum Blot System (Bio-Rad)
transferring at 60mbar for 2 hours. To make RNA cross-linking solution, 245 μl of 12.5 M 1-
methylimidazole (Sigma, St. Louis, MO) was suspended in 9 mL RNase free water, maintaining
a pH of 8.0. Then, a total of 0.753 g 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC)
was added and the total volume adjusted to 24 mL with RNase-free water. RNA was then cross-
linked to the membrane by placing the nylon membrane, side devoid of RNA, on saturated 3mm
Whatman chromatography paper and incubating it in the freshly prepared EDC solution at 60°C
for 2 hours. After cross-linking the membrane was washed with RNase-free water. Next,
CjNC110-LNA probe (/5'DigN/ GCACATCAGTTTCAT/3'Dig_N/) (QIAGEN) was added to
15 mL DIG EasyHyb™ Buffer (Roche) to reach a final concentration of 25 ng/mL.
Hybridization was performed using a hybridization oven and bottle, rotating at 42°C for 12
hours. After overnight incubation, subsequent washes and incubations were performed again
using the hybridization chamber. The blot was first washed twice with low stringency buffer (2x
SSC, 0.1% SDS) for 15 minutes at 50°C, and three times with high stringency buffer (0.1% SSC
and 0.1% SDS) for 10 minutes at 50°C. DIG Washing and DIG Blocking Buffers were prepared
using the DIG Wash and Block Set (Roche, USA). The blot was washed twice with Washing
Buffer for 10 minutes at 37°C. Next, the blot was incubated in 1x Blocking Buffer for 3 hours at

room temperature. DIG antibody solution was added by mixing Blocking Buffer with DIG
antibody at a ratio of 1:5,000. The blot was then incubated for 30 minutes at room temperature.
Next, excess DIG antibody was removed by rinsing with the DIG Washing Buffer a total of four
times for 15 minutes each. For development and detection, the membrane was placed in
Development/Detection Buffer for 10 minutes, and 2 mL ready-to-use CDP-*Star*® (Roche,
USA) was added to cover the blot completely. To image the blot, Chemidoc Imager (Bio-Rad)
was used, exposing the blot for 5 minutes on the chemiluminescence setting. Image layout and
annotations were performed using GraphPad Prism. ImageLab software (Bio-Rad) was used to
generate a standard curve using RF and nt size of the ladder to predict the average band size of
the target RNA.
325

TargetRNA2 and RNAFold
To attempt to computationally determine potential targets regulated by CjNC110 and to
compare results to our RNAseq data, the freely available TargetRNA2 program
(http://cs.wellesley.edu/~btjaden/TargetRNA2/) was utilized to predict potential targets of
interest *a priori* [46]. The standard program settings were utilized, which included searching the
region 80 nt upstream and 20 nt downstream of the predicted translational start site of all genes
in the IA3902 genome. To attempt to predict the secondary structure of the CjNC110 non-
coding RNA, the standard settings of the RNAFold webserver was used for prediction analysis
(http://rna.tbi.univie.ac.at/cgi-bin/RNAfold.cgi) [47, 48]. Both the 137 nt read length predicted
from Dugar *et al.* [21] and the 152 nt length predicted from our northern blot results were used
for analysis via these methods.
337

Enumeration of *luxS* Transcriptional Levels Utilizing qRT-PCR
To validate the RNAseq results and compare *luxS* expression between mutants,
quantitative reverse transcriptase PCR (qRT-PCR) was performed using total RNA isolated from
IA3902 wild-type, ΔCjNC110, and ΔCjNC110c purified in three separate biological replicates of
broth cultures grown to 12 hours from the second growth curve described above. Total RNA
was extracted and qRT-PCR was performed using the iScript cDNA Synthesis Kit (Bio-Rad,
Hercules, CA) and 1000 ng of total RNA according to the manufacturer's instructions as
previously described [49]. Each sample was normalized to the same starting amount of cDNA
using Qubit™ BR DNA Assay Kit (Invitrogen). cDNA purity was measured using the NanoDrop
ND-1000 spectrophotometer. qPCR assays were run using the SsoAdvanced™ Universal
SYBR® Green Supermix (Bio-Rad, Hercules, CA) and the CFX Maestro™ Real-Time PCR
detection system (Bio-Rad). Dilutions of cDNA template for both standards and all unknowns
were run in triplicate with reaction volumes of 10 μl. Amplification of converted cDNA
occurred with 35 cycles of denaturation at 95°C for 10 seconds and then annealing for each
primer pair (**Table 3**) at 58°C for 30 seconds. Prior to analysis, both standards curves (16S and
luxS) were experimentally validated to have high efficiency 90%> of amplification and precision
R^2=0.98 or greater. Relative fold change in *luxS* mRNA expression between IA3902 wild-type,
ΔCjNC110, and ΔCjNC110c was calculated using the ISU Gallup Method Equation [50].
Statistical analyses was performed using one-way ANOVA (GraphPad Prism) to determine
significance in gene expression levels. A p-value of <0.05 was considered significant.
358

359 ## *Vibrio harveyi* bioluminescence assay

360   Culture samples collected from time points 3, 6, 9 and 12 hours (2 mL each timepoint)
361 from the second growth curve described above were centrifuged at 20,000 x g for 5 minutes at
362 4ºC. The supernatant was then removed from the cell pellet and filter-sterilized using a 0.2 µm
363 syringe filter to create cell-free supernatant representative of the extracellular environment (E-
364 CFS); both the E-CFS and the remaining cell pellet were then frozen at -80ºC. For processing of
365 the cell pellet to create intracellular cell-free supernatant (I-CFS), samples were thawed on ice
366 and then washed twice with PBS; following each wash, the cells were spun again at 8000 x g and
367 the resulting supernatant was completely removed from the cell pellet. Next, the cell pellet was
368 resuspended in 600 µl of ice-cold MH broth. The resuspended cell pellets were then lysed using a
369 Bullet Blender® (NextAdvance, Troy, NY). To lyse the cells, 1.5 mL microcentrifuge tubes
370 (Corning, Corning, NY) were loaded with sterilized 0.9-2.0 mm diameter stainless steel beads
371 (NextAdvance). The Bullet Blender was used for 5 minutes at max speed and stored at 4°C.
372 Lysed cellular debris were spun at 8000 x g for 2 minutes using a microcentrifuge, and the
373 resulting supernatant filtered using a 0.20 µm sterile syringe filter, resulting in intracellular cell-
374 free supernatant (I-CFS). Both E-CFS and I-CFS were frozen at -80 ºC until proceeding with the
375 bioluminescence assay.
376   Autoinducer-2 levels within the collected supernatant for both intracellular (I-CFS) and
377 extracellular (E-CFS) samples were then enumerated using the *Vibrio harveyi* bioluminescence
378 assay as previously described [51]. Briefly, 10 µL of each CFS sample was added to 90 µL AB
379 media containing a 1:5000 dilution of the reporter strain, *V. harveyi* strain BB170 [31] in
380 triplicate. Relative light units (RLU) were measured every 15 minutes over 8 hours using the
381 FLUOstar Omega (BMG Labtech, Ortenburg, Germany). MH broth and AB media were used as
382 negative controls, while E-CFS collected from *V. harveyi* strain BB152 [29] was used as a
383 positive control. The timepoints utilized for analysis were those occurring during the nadir of
384 values for the negative control wells and at a standardized duration of time (approximately 3
385 hours) following initiation of increasing values for the positive wells. Results reported are the
386 average of three independent growth curves with three technical replicates used at each
387 timepoint. Statistical analysis was conducted using a two-way ANOVA with repeated measures
388 and Sidak's multiple comparisons test (GraphPad Prism). A p-value of <0.05 was considered
389 significant.
390

391 Motility assay

392   Motility was determined via inoculation of plates consisting of MH broth with 0.4% agar
393 as previously described in our laboratory [8]. Briefly, the A_{600} of overnight cultures was adjusted
394 to 0.3 using sterile MH broth on a Genesys 10S VIS spectrophotometer (Thermo Scientific™). A
395 1 µL volume inoculation stick was then dipped into a set volume of the standardized culture
396 contained in the bottom of a 15 mL conical tube which was then used to make a stab inoculation
397 into the center of the freshly made motility agar (MH broth with 0.4% Bacto agar) with a new
398 inoculation stick for each plate. Plates were incubated at 42°C under microaerophilic conditions
399 as described above with the exception that the plates were incubated right-side up and in a single
400 layer. Measurement of the outermost reach of the halo was performed at 30 hours following
401 inoculation for the first set of experiments and at 24 hours for the second set of experiments. All
402 strains were assessed for average motility using six biological replicates consisting of six
403 technical replicates during each independent study. The six experiments were statistically
404 analyzed using a one-way ANOVA and differences between each strain assessed via Tukey's
405 multiple comparisons test (GraphPad Prism). A p-value of <0.05 was considered significant.

406
Autoagglutination assay

Autoagglutination was assessed according to the method described previously [52] with some modifications. Briefly, the A_{600} of overnight cultures was adjusted to 1.0 in sterile Dulbecco's phosphate buffered saline (PBS) (Corning cellgro, Manassas, VA) using a Genesys 10S VIS spectrophotometer (Thermo Scientific™). The suspension was then aliquoted (2 mL each) into standard glass culture tubes. One subset of cultures were kept at controlled room temperature (25°C) under microaerophilic conditions; the others were incubated at 37°C microaerophilic. At 24 hours, 1 mL of the upper aqueous phase was carefully removed and A_{600} measured to determine autoagglutination activity. All strains were assessed in quadruplicate at each temperature in three independent experiments. The three experiments were statistically analyzed using a one-way ANOVA and differences between each strain assessed via Tukey's multiple comparisons test (GraphPad Prism). A p-value of <0.05 was considered significant.

Liquid Chromatography with Tandem Mass Spectrometry-Based Proteomic Analysis

For extraction of whole cell protein, bacterial cells from IA3902 wild-type, ΔCjNC110 and ΔCjNC110c were grown to exponential phase of growth at 16 hours on MH agar. Lawn cultures were then harvested from the plates using 1 mL of MH broth and resuspended in MH broth. Collected lawns were normalized to an A_{600} of 0.2 and aliquoted into 1 mL technical replicates. This was repeated for three biological replicates. To wash the cells, each aliquot was re-suspended in 1 mL of PBS, cells were spun down at 8000 x g for 3 minutes, and the supernatant was decanted. Next, 50 µl of lysis buffer (filtered sterilized water + 1% Triton X-100) was added to the protein pellets for resuspension (adapted from [53]). The cells were then mechanically sheared by boiling at 96°C for 5 minutes followed by rapid pipetting. After 3 cycles of boiling-shearing, total cell extracts were centrifuged at $10,000 \times g$ for 5 min to remove cell debris. The supernatants were collected, and protein concentration was determined using the Qubit™ Protein Assay and Fluorometer (Invitrogen™). Crude protein was then submitted to the Iowa State Protein Facility. Crude protein extracts were digested overnight in solution with trypsin/Lys-C (25-50 µg). After digestion, Pierce™ Peptide Retention Time Calibration Mixture (PRTC) (ThermoFisher Scientific) was spiked into the samples to serve as an internal control (25 fmol/µL). The PRTC areas were used to normalize collision energies, retention times, and peak intensities to allow for quantitative analysis between samples.

Proteomic analysis was performed via liquid chromatography with tandem mass spectrometry (LC-MS/MS) using a Q Exactive™ Hybrid Quadrupole-Orbitrap Mass Spectrometer (Thermo Scientific™) system. The system was coupled with EASY-nLC 1200 nanopump with integrated auto-sampler (Thermo Scientific™). Liquid chromatography was used to separate the peptides followed by fragmentation and MS/MS analysis. The resulting intact fragmentation pattern was compared to a theoretical fragmentation pattern (MASCOT) to find peptides that can be used to identify high-confidence proteins. Minora Feature Detector was used for label-free quantification to detect and quantify discovered isotopic clusters.

Abundances of proteins were compared by grouping by strain and averaging the abundances of each protein identified. For data filtering and data imputation, PANDA-view, a freely available proteomic software (https://sourceforge.net/projects/panda-view/), was used [54]. Briefly, only proteins identified in two out of three biological replicates were included in

analysis. Nearest neighbor (kNN) imputation was used to fill any missing protein values [55]. All
data was log transformed using $\log_2$. For statistical analysis, two-way ANOVA with false
discovery rate correction set at 0.05 was performed (Graphpad Prism). Graphical Hierarchical
clustering heat maps were generated using freely available MetaboAnalyst software [56]. The
top 20 proteins were group and analyzed using the following heat map settings: distance
measurement of Euclidean and the average clustering algorithm [56, 57].
457

Chicken colonization

Chicken colonization studies were performed as previously described in our laboratory
utilizing 1 day-old broiler chicks obtained from a commercial hatchery [8]. A subset of chicks
from each group were screened via cloacal swabs plated on MH agar containing both selective
supplement and growth supplement (MH+sel+sup) (Oxoid, ThermoFisher Scientific) and found
to be negative for *Campylobacter* carriage prior to inoculation. At 3 days of age, chicks were
inoculated by oral gavage with 200 µl of bacterial suspension containing approximately 1×10^7
of one of the tested strains. Groups of chicks inoculated with specific strains were housed in
separate brooders with no contact between the groups. At pre-determined timepoints post-
inoculation (5 days, 12 days, and 19 days), a minimum of 5 chicks from each group were
randomly selected for humane euthanasia which was immediately followed by necropsy. Cecal
contents were harvested aseptically and stored on ice until further processing could be
completed. Cecal samples were then weighed and subjected to a 10-fold serial dilution series,
plated on MH+sel+sup agar, and grown under routine culture conditions as described above.   To
confirm that the isolates recovered were in fact the mutant strain and no cross contamination
occurred, representative colonies were selected from each strain and timepoint, grown in
MH+sel+sup broth, and plated on the respective antimicrobial selective agar of the mutant strain
as described above. In addition, a minimum of two CFU from each group was collected for PCR
analysis and confirmation of no cross-contamination between biological groups. The detection
limit of the assay was determined to be 100 CFU/g; for isolates with no detectable colonies
within the optimal target range of 30-300 CFU on the initial dilution plate, a count of 3×10^3 was
used to enable statistical analysis. For statistical analysis, one-way ANOVA and Tukey's
multiple comparison test was used to determine significant differences in colonization between
biological groups (GraphPad Prism), using the null hypothesis that colonization rates between
groups are the same. A p-value of <0.05 was considered significant.
483

Data availability

485
486 The RNAseq dataset generated in this publication has been deposited in the NCBI Database
487 under BioProject ID Number PRJNA590513
488 (http://www.ncbi.nlm.nih.gov/bioproject/590513).
489

Results

Northern blotting and RNAseq validate expression of small RNA
CjNC110 and mutant constructs

To validate expression of CjNC110 in IA3902, northern blots comparing IA3902 wild-type,
ΔCjNC110 and ΔCjNC110c using 15 μg of total RNA extracted at log phase of growth from
cultures collected at the same OD_{600} were performed. Northern blot analysis clearly
demonstrates that small RNA CjNC110 is expressed by IA3902 wild-type and ΔCjNC110c but
not ΔCjNC110 (**Fig 1**). Using ImageLab, a standard curve was generated using RF and nt size of
the ladder to predict the IA3902 CjNC110 average band size, which was 152bp. This is slightly
larger than the size that was initially predicted in Dugar *et al.* which was 137bp [21].

Fig 1. Northern blotting demonstrates expression of CjNC110 in wild-type IA3902 and
ΔCjNC110c but not in ΔCjNC110. Cultures for RNA extraction were collected at log phase of
growth and northern blot analysis was conducted using 15 μg of total RNA in each of three
separate lanes per strain tested.

The RNAseq data that was compiled for each of the mutants allowed further visualization
of the unique read signature related to the mutated regions via the Integrated Genome Viewer as
presented in **Fig 2A** and as compared to the anticipated mutated gene structure (**Fig 2B**). In all
timepoints and replicates sequenced for the ΔCjNC110 and ΔCjNC110ΔluxS mutants, no reads
mapped to the reported location of the CjNC110 small RNA; this was anticipated based on the
method of mutant construction which removed that entire region of the genome en bloc and
replaced it with an antibiotic resistance marker. Rockhopper did fail to identify CjNC110 as a
small RNA candidate in this dataset which was somewhat unexpected. Further analysis of the
data revealed that expression levels of CjNC110 transcripts were relatively low in the IA3902
wild-type at both stages of growth, which may explain why it was not recognized by Rockhopper
as a unique ncRNA.
515
516 **Fig 2. Graphic view of mutations**. (A) A screen capture depicting the average RNAseq
517 expression results from IGV of the *luxS* and CjNC110 regions of the genome, corresponding to
518 the genome structure as depicted in (B) of all of the strains sequencing using RNAseq.
519
For the ΔluxS and ΔCjNC110ΔluxS mutants, the location of the insertion of the
kanamycin resistance cassette into the coding sequence of the *luxS* gene can be directly observed
as a sudden decrease in the amount of reads downstream of the location of the insert. In the
ΔluxS mutant, reads mapping to CjNC110 can also be seen which indicates that the mutation
does not interfere with ability of the downstream small RNA to be transcribed. In the
ΔCjNC110ΔluxS mutant, no reads are present that map to the CjNC110 coding region,
confirming that both genes are inactivated in that strain.
527

Differential gene expression analysis of RNAseq data indicates upregulation of *luxS* in ΔCjNC110 mutant and significant gene expression changes in ΔCjNC110Δ*luxS* associated with flagellar biosynthesis

Overall, 24 barcoded libraries were sequenced yielding over 109 million reads, with close
to 100 million high quality reads aligning to either the genome or pVir plasmid of *C. jejuni*
IA3902 and averaging 4,553,847 reads per library (**Tables C and D in S1 Appendix**). The vast
majority of reads (average of 83% of total reads) mapped to protein coding genes of the

chromosome, with only 7% of reads mapping to ribosomal RNA on average following rRNA
depletion with Ribo-Zero (median of 3%). Over half of the libraries contained less than or equal
to 3% ribosomal RNA reads, which is consistent with the manufacturer's predicted rRNA
removal efficiency. One third of the libraries did not exhibit efficient rRNA removal (>10%
rRNA reads); the reason for this is unclear. Despite this difference in efficiency, all samples
provided a more than adequate number of non-rRNA reads and thus were successfully utilized in
the analysis. Less than 1% of reads mapped to antisense regions of the annotated genome.
A summary of the differences in numbers of genes with increased and decreased
expression in the various strains and timepoints is given in **Table 2**, with a listing of specific
genes given in **Table 3**. In the ΔCjNC110 mutant strain, six genes were found to be
downregulated and four genes were found to be upregulated when compared to the IA3902 wild-
type strain during exponential growth (complete information **Table E in S1 Appendix**). In
addition, a previously described non-coding RNA, CjNC140 [21] was found to be upregulated in
the mutant condition. During the stationary phase, 16 genes were found to be downregulated and
seven genes were found to be upregulated in the mutant strain when compared to wild-type. Of
the regulated genes, three genes (*neuB2, hisF, ptmA*) were downregulated in the ΔCjNC110
mutant during both the exponential and stationary phases, and one hypothetical protein coding
gene (CjSA_1261) was found to be upregulated in both conditions. Of the differentially
expressed genes, five separate operons predicted by Rockhopper demonstrated multiple genes
within the operon affected by the mutant condition. Analysis of functionality via the COG
database revealed that multiple upregulated genes (*luxS, cetA*, *cetB*) were present in the "Signal
transduction mechanisms" category; multiple downregulated genes were also included in the
"Cell wall/membrane biogenesis" (*neuB2, ptmB,* CjSA_1352) category and "Posttranslational
modification, protein turnover, chaperones" (*tpx,* CjSA_0687) functional category.
560
561 **Table 2**. Summary of differential gene expression results between mutant strains.

	Condition					
	ΔCjNC110		ΔluxS		ΔCjNC110ΔluxS	
	3 h	12 h	3 h	12 h	3 h	12 h
Protein-coding genes						
Number downregulated	6	16	1	6	61	32
Number upregulated	4	7	14	6	47	29
Non-coding RNA genes						
Number downregulated	0	0	0	0	25[b]	2
Number upregulated	1[a]	0	1[a]	0	1[a]	0

[a] = previously described ncRNA, CjNC140 [21]

[b] = 17 tRNA genes, 3 known RNA genes (tmRNA, SRP, 6S) and 4 newly predicted ncRNA

562
563 **Table 3.** Summary of differential gene expression (>1.5 fold change, q <0.05) in the IA3902
564 ΔCjNC110, ΔluxS and ΔCjNC110ΔluxS mutants as determined by RNAseq.

	Exponential (3 h)						Stationary (12 h)			
	ΔCjNC110*	**ΔluxS***	**ΔCjNC110ΔluxS***				**ΔCjNC110***	**ΔluxS***	**ΔCjNC110ΔluxS***	
Genes downregulated	hisF	CjSA_1350	CjSA_0014	flgG	flgI	rplF	panC	CjSA_0560	rnhB	aroB
	pseA		CjSA_0041	mogA	CjSA_1388	rpsH	CjSA_0559	CjSA_0620	CjSA_0158	CjSA_1137
	neuB2		flgD	aspB	CjSA_1389	rpsN	CjSA_0687	CjSA_1349	CjSA_0160	CjSA_1164
	ptmA		flgE	hydD	flgK	rplE	Tpx	CjSA_1350	CjSA_0284	pyrH
	CjSA_1352		CjSA_0067	pseB	moaE	rplX	CjSA_0785	Acs	motB	CjSA_1349
	rpsN		accD	pseC	nuoD	rplN	ciaB	leuC	trpD	CjSA_1350
			trxA	CjSA_1233	CjSA_1562	rpmC	CjSA_1102		CjSA_0344	CjSA_1414
			panB	neuC2	CjSA_1568	rplP	petC		CjSA_0372	Acs
			motB	CjSA_1350	CjSA_1577	CjSA_CjSRP1	CjSA_1137		CjSA_0389	CjSA_1461
			fliN	CjSA_1351	CjSA_t0035	CjSA_t0002	CjSA_1244		CjSA_0396	rnhA
			rpsU	CjSA_0788	CjSA_t0036	CjSA_t0005	hisF		sdhA	CjSA_1560
			frdB	flgL	CjSA_t0037	CjSA_t0007	neuB2		sdhB	leuC
			flag	rpmH	CjSA_t0038	CjSA_t0012	CjSA_1266		uxaA	leuB
			fliS	CjSA_0920	CjSA_t0039	CjSA_t0013	ptmB		CjSA_0560	leuA
			CjSA_0521	CjSA_1093	CjSA_t0043	CjSA_t0015	ptmA		CjSA_0616	CjSA_pVir0030
			CjSA_0569	CjSA_1102	CjNC130	CjSA_t0017	CjSAt0002		trmD	
			pstS	dctA	secY	CjSA_t0018				
			hslV	luxS	rplO	CjSA_t0020				
			flgH	CjSA_1182	rpsE	CjSA_t0021				
			flgG2	porA	rplR	CjSA_t0033				
Genes upregulated	CjSA_0008	dnaN	dnaN	trpF	cetB	CjSA_1449	hcrA	flgE	dsbI	CjSA_1107
	cetA	CjSA_0008	CjSA_0008	trpB	cetA	rloH	CjSA_0716	flgG2	CjSA_0040	omp50
	cetB	trpF	gltD	trpA	CjSA_1129	nuoM	CjSA_0717	flgG	CjSA_0041	CjSA_1180
	CjSA_1261	trpB	folk	CjSA_0337	CjSA_1131	CjSA_1549	CjSA_1107	hcrA	flgD	pseB
	CjNC140	CjSA_0337	CjSA_0076	CjSA_0370	CjSA_1145	euC	CjSA_1108	CjSA_1107	flgE	CjSA_1262
		CjSA_0370	CjSA_0105	CjSA_0372	CjSA_1164	leuB,	omp50	nrfA	flgC	neuB2
		CjSA_0732	CjSA_0110	sdhA	cbpA	leuA	luxS		flgB	neuC2
		CjSA_1017	dgkA	sdhB	CjSA_1259	CjNC140			flgH	flgI
		CjSA_1131	pyrC	CjSA_0490	CjSA_1260	CjSA_pVir0042			flgG2	CjSA_1387
		CjSA_1301	Tal	CjSA_0491	CjSA_1301	CjSA_pVir0044			flgG	flgK
		CjSA_1352	CjSA_0305	CjSA_0836	CjSA_1343	CjSA_pVir0025			hcrA	CjSA_1562
		CjSA_1449	trpD	Cfa	tagF				CjSA_0716	p19
		CjSA_1549							flgS	CjSA_pVir0042
		CjSA_pVir004							flgL	CjSA_pVir0013
		CjNC140							CjSA_0969	

* = additional small RNAs predicted by Rockhopper present with differential expression; see S4, S5, and S6 Tables for additional information

In the ΔluxS mutant strain, one gene was found to be downregulated and 14 genes were
found to be upregulated when compared to the IA3902 wild-type strain during exponential
growth (summary **Table 3**, complete information **Table F in S1 Appendix**). Similar to the
ΔCjNC110 mutant, the previously described non-coding RNA CjNC140 was also found to be
upregulated in the mutant condition during exponential growth only. At 12 hours, six genes were
found to be downregulated and six upregulated in the ΔluxS mutant strain when compared to
wild-type. Of the differentially expressed genes, only one gene, CjSA_1350, a putative
methyltransferase, was found to be downregulated in the ΔluxS mutant during both exponential
and stationary phases. Three separate operons predicted by Rockhopper demonstrated multiple
genes within the operon affected by the mutant condition (two operons upregulated, one operon
downregulated). Analysis of functionality via the COG database revealed that multiple both up-
(*trpF, trpB*) and down- (CjSA_0620, *leuC*) regulated genes were present in the "Amino acid
transport and metabolism" functional category; multiple upregulated genes were also included in
the "Cell motility" (*flgE, flgG2, flgG*) functional category. Only two genes were differentially
expressed in both the ΔCjNC110 and ΔluxS mutants, CjSA_0008 (upregulated during
exponential growth hours) and CjSA_1107 (upregulated during stationary phase); both genes are
annotated as hypothetical proteins at this time.
In the ΔCjNC110ΔluxS double knockout mutant, a large increase in both downregulated
and upregulated genes when compared to wild-type was observed at both timepoints (summary
Table 3, complete information **Table G in S1 Appendix**). Sixty-one protein coding genes,
seventeen tRNA genes, three known ncRNA genes (tmRNA, SRP, 6S) and seven newly
predicted ncRNAs were downregulated during exponential phase compared to IA3902 wild-type.
In addition, 47 protein coding genes were upregulated in exponential phase, along with the
previously described ncRNA CjNC140, which was also observed to be upregulated in the
ΔCjNC110 and ΔluxS single knockout mutants. During stationary phase, 32 genes and two
newly predicted non-coding RNAs were found to be downregulated, while 29 genes were
upregulated in the mutant strain when compared to wild-type. Of the observed genes with
differential expression, many were observed to show altered expression during both exponential
and stationary phase when compared to wild-type; however, most were differentially expressed
in the opposite direction between the two timepoints. Only two genes (*motB,* CjSA_1350) were
downregulated in both conditions, while no genes were observed to be upregulated at both
timepoints. Many of the differentially upregulated genes demonstrated in both background
mutants, ΔCjNC110 and ΔluxS, were observed in the double knockout (**Figs 3A** and **3B**), while
fewer of the downregulated genes were shared (**Figs 4A** and **4B**).
599
600   **Fig 3. Venn diagram depicting the overlap of shared upregulated genes during the**
601   **exponential and stationary growth phases.** BioVenn was used to compare the lists of known
602   genes upregulated in all 3 mutant strains during both exponential and stationary growth.
603
604   **Fig 4. Venn diagram depicting the overlap of shared downregulated genes during the**
605   **exponential and stationary growth phases.** BioVenn was used to compare the lists of known
606   genes downregulated in all 3 mutant strains during both exponential and stationary growth.
607
Of the differentially expressed genes affected by the ΔCjNC110ΔluxS double knockout
mutation, 20 separate operons predicted by Rockhopper (predicted operon structure listed in
Table J in S2 Appendix) demonstrated multiple genes within the operon affected by the mutant

condition (eight operons upregulated, four operons downregulated, and eight operons that
showed opposing regulation at the different timepoints). **Figure A in S3 Appendix** shows the
number of genes in each COG category that were differentially expressed in the double mutant
condition during either exponential growth. The categories most affected by these mutations
were "Energy production and conversion," "Amino acid transport and metabolism,"
"Translation," "Cell wall/membrane biogenesis," "Cell motility," and "Signal transduction
mechanisms."
KEGG Pathways was used to determine the effect of the mutations on important
pathways in *C. jejuni* pathobiology. **Figs 5A** and **5B** compare the genes observed to be
differentially expressed in the double knockout mutant during both exponential and stationary
phases for flagellar assembly. A large number of genes associated with flagellar assembly were
noted to be differentially regulated in the double knockout mutant as compared to the single
knockout mutants alone (**Fig C in S3 Appendix**).
624
625 **Fig 5. KEGG Pathway analysis demonstrates alteration of expression of multiple genes**
626 **involved in flagellar assembly in *C. jejuni* in the ΔCjNC110ΔluxS mutant during both**
627 **exponential and stationary growth phases.** Blue color indicates downregulation of gene
628 expression, red color indicates up regulation of gene expression, green indicates that the gene is
629 present in *C. jejuni* IA3902, and white indicates that the gene is not present in IA3902.
630

Identification of additional non-coding RNAs in IA3902

In addition to determining differential gene expression between the mutant and wild-type
strains of IA3902, Rockhopper has the capability to predict non-coding RNAs present within the
data. Prior to further manual analysis, Rockhopper predicted a total of 59 ncRNAs present in the
data (57 chromosomal, 2 pVir). Of those 59, five align with previously identified ncRNAs
(CjNC20, CjNC60, CjNC120, CjNC140, CJpv2), all of which were predicted to be conserved in
IA3902 by Dugar *et al.* [21]. Manual curation of the list decreased the number of predicted
ncRNA candidates to 40 (**Table H in S1 Appendix**). Candidate non-coding RNAs were
discarded if they were related to the 16S or 23S genes as these were thought to be spuriously
identified due to Ribo-Zero depletion differences between replicates; in addition, a single
antisense RNA was discarded due to incorrect annotation of the *rnpB* gene to the wrong strand in
IA3902. Of those remaining, 6 were demonstrated transcripts in the region of genes annotated as
pseudogenes in IA3902 (CjSA1052, CjSA1323, CjSA1444, CjSA1543 and CjSA1630 – 2
separate ncRNA predictions). Two of the 40 remaining ncRNA candidates could potentially be
classified as *cis* as they are transcribed opposite of transcribed regions of other genes; 25 would
likely be considered *trans* ncRNAs as they are primarily located within intergenic regions; and
seven are classified as anti-sense RNAs.
648

TargetRNA2 and RNAFold predictions suggest potential targets and secondary structure of CjNC110

The TargetRNA2 program predicted either 21 or 16 regulatory targets of CjNC110 in
IA3902 based on the standard input parameters of the program and a 137 nt or 152 nt length,
respectively (**Table I in S1 Appendix**). Of the predicted targets, only a single gene (CjSA_1137,
annotated as an 2OG-Fe(II) oxygenase) was found to show significantly altered transcription in
the CjNC110 mutant as compared to wild-type; this significant decrease was observed during

both exponential and stationary growth. This gene is located immediately downstream of
CjNC110. No genes were noted to overlap with the protein expression data.
To assess secondary structure of the CjNC110, RNAFold was used to calculate the
structure of CjNC110 using both the 137nt and 152nt predicted sizes, both of which yielded a
prediction of a Y-shaped stem-loop structure (**Fig 6**). When comparing the Target2RNA data to
the RNAFold predictions, all of the predicted interaction areas of the non-coding RNA were
grouped into the 3 unpaired loop regions of the RNA structure.
663
664 **Fig 6. Structural prediction using RNAfold of the 137nt and 152 nt length options for**
665 **CjNC110 RNA.** For both the 137nt and 152nt predicted sizes, the structural prediction for
666 CjNC110 contains 3 stem loops, with the unpaired loops being the region most likely to interact
667 with mRNAs as predicted by TargetRNA2.
668

qRT-PCR confirms RNAseq results that *luxS* is transcribed at higher levels in ΔCjNC110 when compared to IA3902 wild-type

To further validate the results of the RNAseq and confirm that the increased expression
of *luxS* in the ΔCjNC110 mutant was repeatable, qRT-PCR was performed on samples collected
at 12 hours from the second growth curve using targets specific for the 3' region of the *luxS*
gene. **Fig 7** demonstrates the expression levels of *luxS* in the ΔCjNC110 mutant compared to
wild-type and confirms that increased *luxS* mRNA levels are present during stationary phase in
the mutant, resulting in a fold change increase of 3.20 when compared to IA3902 wild-type.
Additionally, ΔCjNC110c was tested to determine if complementation of the mutation returned
luxS to wild-type levels; interestingly, ΔCjNC110c also demonstrated a fold change increase of
3.77 when compared to IA3902 wild-type. One-way ANOVA with Tukey's multiple
comparison testing demonstrated significant fold change difference ($p < 0.05$) in *luxS*
transcription between both ΔCjNC110 and ΔCjNC110c when compared to IA3902 wild-type.
682
683 **Fig 7. qRT-PCR confirms *luxS* is transcribed at higher levels in the ΔCjNC110 and**
684 **ΔCjNC110c backgrounds when compared to IA3902 wild-type (WT) (mean ± SEM).** *C.*
685 *jejuni* cultures were collected at timepoint 12 hours for RNA conversion cDNA for qRT-PCR to
686 assesses expression of mRNA *luxS*. Each bar represents the average fold change expression from
687 three biological replicates. Significance is denoted by "*" above each timepoint.
688

Growth of mutant strains during exponential growth is consistent with wild-type, but significant effects are seen in motility and autoagglutination

The growth of IA3902 wild-type, ΔCjNC110, ΔluxS, and ΔCjNC110ΔluxS was evaluated
over a period of 30 hours and differences tested for statistical significance ($p < 0.05$) using two-
way ANOVA with Dunnet's repeated measures of the A_{600} readings at each time point. ANOVA
did identify a statistically significant difference between strains in both growth curves ($p < 0.05$),
however, multiple comparison analysis of individual time points and strains when compared to
wild-type growth revealed that the only significant difference was a decrease in the A_{600} of
ΔCjNC110 after 27 and 30 hours in both growth curves performed (**Fig A in S3 Appendix** and
Fig 8). In both experiments, after 27 hours represents the point where a decline in the A_{600} begins
to occur in all strains tested, and further examination of the actual CFU/mL over the course of

the first growth curve revealed that by 30 hours cell death was beginning to occur and colony
counts were decreasing (**Fig A in S3 Appendix**). As further growth is assumed to be ceased at
this time, there is likely minimal biological significance to the finding of differences after 27
hours in the assays performed using samples collected during exponential growth prior to 24
hours, however, there is possibly biological significance to the difference noted during the
stationary phase which may warrant further investigation.
707
708 **Fig 8. Shaking growth curve of wild-type IA3902 (WT) and isogenic mutants (mean ±**
709 **SEM) confirm normal exponential growth.** Results of four replicates (A_{600}) of a shaking
710 growth curve performed in 250 mL Erlenmeyer flasks under microaerophilic conditions in MH
711 broth. Samples were pulled from this growth curve and used for qRT-PCR and AI-2
712 bioluminescence assays. Significance is denoted by "*" above each timepoint.
713
Motility of the mutant strains in semi-solid agar was compared to wild-type IA3902 at 30
hours post-inoculation for all mutant isolates, and all isolates were confirmed to be highly motile
(**Fig 9**). Statistical analysis via one-way ANOVA indicated that there was a significant difference
between strains (p <0.0001). Further post-hoc analysis via Tukey's multiple comparisons test
did not reach statistical significance (p>0.05) when comparing ΔCjNC110 to wild-type despite
motility for the ΔCjNC110 strain observed to be consistently increased above the wild-type
phenotype in all biological replicates performed. Results did reach statistical significance
(p<0.05), however, when comparing wild-type to both the ΔCjNC110ΔluxS and ΔluxS mutants
which displayed statistically significant decreased motility; this result is consistent with previous
reports of decreased motility of *luxS* mutants in the IA3902 background [8]. As many of the
strains had migrated to the edge of the plate by 30 hours, we hypothesized that by reading the
plates sooner we may observe a larger difference in the motility of the ΔCjNC110 strain,
therefore, a second set of experiments was performed with measurements taken at 24 hours post-
inoculation. When measured at 24 hours, motility for the ΔCjNC110 strain did demonstrate a
statistically significant increase (p<0.05) when compared to all other strains tested, and motility
of both the ΔCjNC110ΔluxS and ΔluxS mutants was again noted to be significantly decreased.
Following complementation of CjNC110, no significant difference was observed when
compared to IA3902 wild-type in either experiment, confirming that complementation of
CjNC110 quantitatively restored the phenotypic change back to wild-type levels.
733
734 **Fig 9. Motility assays confirm isogenic mutants remain motile, with ΔluxS motility**
735 **decreased and ΔCjNC110 motility increased over IA3902 wild-type (mean± SEM).** Each
736 bar represents the average motility of six biological replicates consisting of six technical
737 replicates during each independent study. Significance is denoted by "*" above each bar when
738 comparing wild-type to other respective strains at each independent timepoint.
739
Autoagglutination activity was measured at 25ºC and 37ºC following 24 hours of
incubation (**Fig 10**) with an increase in optical density correlating to decreased autoagglutination
ability, and a decrease in optical density correlating to increased autoagglutination ability. A
statistically significant difference (p <0.0001) between strains at both 25ºC and 37ºC was noted
based on initial analysis via one-way ANOVA. When compared to IA3902 wild-type,
autoagglutination activity for ΔluxS and ΔCjNC110ΔluxS was noted to be increased at a
statistically significant level (p<0.05) at both temperatures, while ΔCjNC110 exhibited

statistically significant decreased autoagglutination activity, again at both temperatures.
Complementation of CjNC110 restored the autoagglutination ability of ΔCjNC110 to wild-type
levels. These results indicate that *luxS* and CjNC110 function to influence autoagglutination
activity in opposing directions; interestingly, when mutation of both genes was combined, the
results favored the ΔluxS mutation but did return closer to wild-type levels.
752
Fig 10. ΔluxS bolsters autoagglutination and ΔCjNC110 hinders autoagglutination of
IA3902 (mean± SEM). Autoagglutination as determined by optical density (A_{600}) was measured
following a 24 hour incubation at either 25°C or 37°C. Each bar represents the average
autoagglutination of 4 biological replicates consisting of 3 technical replicates during each
independent study. Significance is denoted by "*" above each bar when comparing wild-type to
other respective strains at each independent timepoint.
759

Vibrio harveyi bioluminescence assay demonstrates decreased extracellular AI-2 levels with a concomitant increase in intracellular AI-2 in ΔCjNC110

Bioluminescence activity was measured using the *Vibrio harveyi* assay as an
approximation for extracellular (using E-CFS) AI-2 levels at various time points during the
growth of IA3902 wild-type and mutant strains (**Fig 11A**). As expected, both the IA3902ΔluxS
and ΔCjNC110ΔluxS mutant strains demonstrated no bioluminescence activity at any point
during growth, indicating a complete lack of AI-2 production which is consistent with the
absence of a functional LuxS protein. For the ΔCjNC110 mutant, bioluminescence was
determined to be statistically significantly decreased when compared to wild-type at time points
6, 9 and 12 hours ($p <0.05$) of the growth curve, occurring during mid to late exponential phase
and early stationary phase. Complementation of ΔCjNC110 completely restored extracellular
AI-2 to wild-type levels.
773
774 **Fig 11. Bioluminescence activity as measured via the *Vibrio harveyi* bioassay over the**
775 **course of growth (mean ± SEM) demonstrates (A) decreased extracellular AI-2 levels with**
776 **(B) a concomitant increase in intracellular AI-2 in ΔCjNC110** Each bar represents the
777 average RLU (AI-2 activity level) from three biological replicates consisting of three technical
778 replicates each. Significance is denoted by "*" above each timepoint.
779
To determine if AI-2 production or transport across the cell membrane was impaired
leading to this observed decrease in extracellular AI-2, the intracellular bioluminescence activity
was also assessed utilizing I-CFS obtained from the cell pellet of the same culture as the E-CFS.
Results of this assay demonstrated that intracellular AI-2 levels increased to statistically
significant ($p <0.05$) levels in the ΔCjNC110 mutant when compared to IA3902 wild-type at
both the 9 hour and 12 hour timepoints. (**Fig 11B).** Again, complementation of ΔCjNC110
corrected the phenotypic alteration of increased intracellular AI-2 presence. These results
indicate that the observed decrease in extracellular AI-2 activity is not due to impaired
production, and instead suggests that transport of AI-2 across the cell membrane is altered in the
ΔCjNC110 mutant.
790

Liquid Chromatography with Tandem Mass Spectrometry confirms increased LuxS protein abundance in both ΔCjNC110 and ΔCjNC110c

Proteomic analysis was conducted to reveal the most highly abundant proteins and changes in their translational level at exponential phase of growth to compare to transcriptomic data. Whole protein extract LC-MS/MS analysis resulted in a total of 262 identified proteins that were discovered in each biological group. Two-way ANOVA with FDR correction identified statistically significant differences in protein abundance; these proteins are illustrated in **Table 4**. When comparing wild-type to ΔCjNC110, proteins ThiJ and FrdA had a statistically significant fold change difference of 19.7 and -11.3, respectively. When comparing wild-type to ΔCjNC110c, proteins CjSA_0166 (lipoprotein) and RplN both had statistically significant fold change of -9.8, and RplU had a fold change of -4.9. In addition, proteins RecJ and AtpF demonstrated a statistically significant fold change of 8.0 and 9.2, respectively. There were several proteins that were identified as significantly different when comparing wild-type to both ΔCjNC110c and ΔCjNC110, including: LuxS, which exhibited increased expression, as well as CjSA_0067 (iron-sulfur protein) and CjSA_0489 (rhodanese-like domain-containing protein), both of which exhibited decreased expression in both backgrounds.

Table 4: Significant differentially expressed proteins identified from proteomic analysis ΔCjNC110 and ΔCjNC110c versus IA3902 wild-type

Protein ID	Geometric Mean (Abundance)		Fold Change vs WT	P-value*
	WT	Mutant		
ΔCjNC110				
CjSA_0067 - iron-sulfur protein	23726566.41	851708.3688	-27.9	<0.0001
CjSA_0489 - rhodanese-like domain-containing	35962749.77	2965820.801	-12.1	0.0001
ThiJ - thiazole monophosphate synthesis protein	1589344.014	31307392.07	19.7	<0.0001
FrdA - fumarate reductase flavoprotein subunit	3178688.029	280958.9826	-11.3	0.0002
LuxS - S-ribosylhomocysteine lyase	3913424.009	50859008.46	13.0	<0.0001
ΔCjNC110c				
CjSA_0067 - iron-sulfur protein	23726566.41	1123835.93	-21.1	<0.0001
CjSA_0489 - rhodanese-like domain-containing	35962749.77	2767208.654	-13.0	0.0001
CjSA_0166 – lipoprotein	25429504.23	2581896.969	-9.8	0.0006
RplN - 50S ribosomal protein L14	10327587.87	1048576	-9.8	0.0007
RplU - 50S ribosomal protein L21	15653696.03	3178688.029	-4.9	0.0007
RecJ - single-stranded-DNA-specific exonuclease	2408995.053	19271960.42	8.0	0.002
AtpF - ATP synthase subunit B	2965820.801	27254667.8	9.2	0.001
LuxS - S-ribosylhomocysteine lyase	3913424.009	54509335.6	13.9	<0.0001
CjSA_0926 - lipoprotein	1703416.738	23726566.41	13.9	0.0001

Geometric mean calculated by backtransforming log2 transformation value

Hierarchical heat maps also reveal contrasting protein expression patterns when
comparing IA3902 wild-type to ΔCjNC110 and ΔCjNC110c. The top 20 hierarchical clustered
proteins are illustrated for both ΔCjNC110 and ΔCjNC110c compared to wild-type IA3902 in
Fig 12. Heat maps can provide meaningful clusters of data that can drive biomarker discovery
that may not be picked up by a stricter statistical-based test, and can also shrink the data to find
the most meaningful changes in the dataset. We note overlap between FDR significant proteins
and proteins identified after heat map analysis, including ThiJ, LuxS, CJSA_0489, and
CJSA_0166. Several additional proteins were identified via this approach that did not reach the
FDR threshold for significance but exhibited clear differences in translational patterns when
compared to IA3902 wild-type. In the ΔCjNC110 background, protein PycB demonstrated
increased expression when compared to IA3902 wild-type, while proteins with decreased
expression included CJSA_0387, CJSA_1389, NifU, FlaA, FlgD, and RpoZ. In addition, various
proteins classified as periplasmic (CjSA_0190, CjSA_0854, CjSA_1107, CjSA_1178,
CjSA1419, and CjSA1577) and lipoprotein (CjSA_0969) also exhibited decreased expression
compared to wild-type. Complementation of the CjNC110 mutation resulted in reversal of these
expression changes for all of the proteins described with the notable exception of LuxS which
exhibited increased expression in both backgrounds; this finding was consistent with the results
of the qPCR which demonstrated increased gene expression of *luxS* in both strains.

Fig 12. **Heatmaps reveal differential protein clustering patterns induced in both ΔCjNC110**
and ΔCjNC110c when compared to IA3902 wild-type (WT). All raw abundances were
normalized, imputed, transformed to $\log_2$ scale values, and clustered based on resulting
hierarchical patterns. The top 20 proteins were grouped and analyzed using heat map settings:
distance measurement of Elucidean and average clustering algorithm.

Colonization of chickens is impaired in the ΔCjNC110 and absent in
ΔluxS and ΔCjNC110ΔluxS mutants, but is restored to wild-type
colonization levels in ΔCjNC110c

To determine if mutation of ΔCjNC110, ΔluxS and ΔCjNC110ΔluxS affected the ability
of IA3902 to colonize chickens, cecal colonization levels were monitored for 3 weeks following
oral inoculation of approximately 200 µl of 1×10^7 CFU per strain. On DPI 5, a statistically
significant decrease in colonization was noted for both the ΔCjNC110 and ΔluxS strains
compared to wild-type; complementation of the ΔCjNC110 mutant restored colonization to wild-
type levels (**Fig 13**). On DPI 12 and 19, a statistically significant decrease in colonization was
noted for the ΔCjNC110, ΔluxS, and ΔCjNC110ΔluxS strains compared to wild-type with no
colonization of ΔluxS and ΔCjNC110ΔluxS apparent at these time points; complementation of
the ΔCjNC110 mutant again restored colonization levels to wild-type levels for both DPI 12 and
19. These results indicate that the presence of CjNC110 is necessary for optimal colonization,
while *luxS* is required for colonization of IA3902 in chickens.

Fig 13. Chicken colonization is impaired in ΔCjNC110 and absent in ΔluxS and
ΔCjNC110ΔluxS mutants compared to wild-type (mean ± SEM). Results are reported as $\log_{10}$
CFU/mL of ceca content for each sampling day [DPI 5 (Week 1), DPI 12 (Week 2), and DPI 19

(Week 3)]. Each bar represents the average colonization levels for each biological group with
weekly technical replicates using a minimum of 6 birds. Significance is denoted by "*" above
each timepoint.
858

Discussion
859

Over the past several years, RNAseq has become the method of choice to analyze the
transcriptome of bacteria under various conditions as it allows evaluation of the entire
transcriptome rather than only previously annotated regions as with older technology such as
microarrays [58-60]. Recent work in *Campylobacter* has demonstrated that RNAseq can be
useful for analyzing transcriptomic differences between wild-type and mutant strains following
inactivation of protein-coding genes [61, 62]. In theory, RNAseq should also be able to be
utilized for the same purpose of discovering the global effects of inactivation of ncRNAs. Small
RNAs have been shown to be able to stabilize mRNA transcripts by multiple mechanisms [63],
which should lead to increased levels of transcript availability and identification using RNAseq.
In contrast, some interactions with ncRNAs have also been shown to increase transcript turnover
by targeting transcripts for RNase degradation or exposing RNase cleavage sites [63], which
should lead to decreased levels of transcripts available for identification via RNAseq. Thus,
assessment of differential gene expression via RNAseq should be useful for uncovering
interactions of ncRNAs with their targets that directly leads to altered levels of mRNA
transcripts in the cell. One limitation of the use of RNAseq data for discovery of targets of small
RNAs is that some target interactions may not lead to direct changes in mRNA transcript levels
but instead affect the translation efficiency of the target mRNA; these types of interactions
cannot be determined via this approach. Therefore, use of multiple additional methods such as
computational approaches and proteomics to determine additional potential targets of ncRNAs is
also warranted. The results of our work clearly indicate that RNAseq can be a useful tool for
investigating the role of ncRNAs in gene expression in bacteria and can successfully be used to
inform the design of phenotypic studies to begin to elucidate the function of ncRNAs in
pathogenic bacteria.
During stationary phase in the ΔCjNC110 mutant, expression of the *luxS* mRNA was
determined to be statistically significantly upregulated when compared to wild-type using
RNAseq; during exponential growth, increased expression was also present but did not reach
statistical significance. This initially suggested that CjNC110 may normally interact with the
luxS transcript to impact AI-2 expression and quorum sensing. qRT-PCR confirmed the
observed increase in *luxS* transcription seen via RNAseq, and increased expression of the LuxS
protein was also observed via LC-MS/MS, clearly indicating that the response to the loss of
CjNC110 expression was to increase LuxS protein production. However, further investigation
revealed that despite the increased *luxS* expression and LuxS protein abundance in the CjNC110
mutant, AI-2 extracellular concentrations were decreased. This defect was corrected via
complementation of CjNC110 despite the fact that *luxS* expression and LuxS protein abundance
remained increased in the complement. The reason for the increased *luxS* expression and LuxS
protein abundance in CjNC110c is currently unknown but is most likely related to additional
unknown effects of CjNC110 on the AMC due to constitutive expression of CjNC110 in the
complement. The increased *luxS* expression and LuxS protein abundance in the CjNC110c,
combined with complementation back to wild-type of extracellular AI-2 activity, does clearly

demonstrate, however, that a change in LuxS protein abundance is not the cause of the observed
difference in extracellular AI-2 concentrations.
Having ruled-out decreased LuxS abundance as a cause for the decreased extracellular
AI-2, potential hypotheses for the observed difference included either decreased production of
AI-2 by LuxS, increased degradation of AI-2, or altered kinetics of AI-2 movement into or out of
the extracellular environment. To assess for altered kinetics of AI-2 movement, intracellular AI-
2 levels were measured in the ΔCjNC110 mutant compared to wild-type and revealed an
accumulation of AI-2 intracellularly in ΔCjNC110 that was corrected to wild-type levels in
ΔCjNC110c. This indicates that impairment of AI-2 transport out of or increased import into the
cell was the reason for the observed decrease in extracellular AI-2 in the CjNC110 mutant, and
not decreased production or increased degradation. In *E. coli,* the YdgG (TqsA) protein, part of
the AI-2 exporter superfamily, has been described to export AI-2 molecules [64]. To date,
however, no similar homologue has been found in *C.jejuni*. Previous BlastP analysis has shown
additional potential homologues of the YdgG protein exist within the *Campylobacterales* family,
but none directly within the *C. jejuni* species [65]. Within our proteomics data, several
periplasmic and lipoproteins which could be located in the cytoplasmic membrane were
identified to exhibit decreased expression in ΔCjNC110 that was corrected to wild-type levels in
ΔCjNC110c. The function of many of these proteins has yet to be elucidated in *Campylobacter,*
therefore, future work assessing whether these proteins could serve as an AI-2 exporter in *C.*
jejuni is warranted. In other species of bacteria such as *E. coli*, uptake of AI-2 from the
extracellular environment has also been shown to be mediated by an active LSR system [66].
Comparative genomics has again failed to identify a homologous AI-2 uptake system in *C.*
jejuni, however, several ABC-type transporters whose function remains to be elucidated do exist
in *Campylobacter* sp. [27]. Several studies evaluating the concentration of AI-2 over time
following addition of exogenous AI-2 to cell culture have attempted to determine whether or not
AI-2 is able to be internalized in *C. jejuni* with mixed results [67, 68]. Thus, further research
into an AI-2 uptake system in *C. jejuni* may also be warranted.
Interestingly, analysis of transcriptome changes in the ΔluxS mutant were not as
enlightening as was expected. The list of genes identified does not overlap with the list of
differentially expressed genes generated via microarray in *C. jejuni* strain 11168 which utilized
the exact same mutant construct that was moved via homologous recombination into IA3902
[69]; however, differences in the strain of *C. jejuni* utilized in this study (IA3902) and culture
conditions may explain why there was no overlap of differentially expressed genes. No genes
associated with the AMC pathway were identified in our RNAseq data as have been previously
reported via microarray in other strains of *C. jejuni* [70], however, there was very minimal
expression of many of the genes associated with the AMC pathway in our data (*metE, metF, pfs*)
which may have led to an inability to detect a difference. Decreased transcription of *flaA* has
previously been associated with *luxS* mutation in strain 11168 [71]; expression of *flaA* during
exponential growth was decreased 1.4-fold compared to wild-type, however, this change was not
found to be statistically significant; no difference was present in *flaA* levels during stationary
growth. A single gene, CjSA_1350, which is annotated as a putative methyltransferase, was
found to be down-regulated during both exponential and stationary growth in the ΔluxS mutant
of IA3902. The LuxS enzyme is critical for the formation of *S*-adenosylmethionine (SAM),
which is a major methyl donor necessary for methylation of DNA, proteins, carbohydrates and
other molecules important to the physiology of prokaryotes [72]. It is plausible that expression of
this methyltransferase is sensitive to decreased availability of products from the activated methyl

cycle due to *luxS* mutation; however, recent data generated within our lab group suggests that
this gene does not encode an active methyltransferase in IA3902 [73]. No change was detected in
the expression of CjNC110 in the ΔluxS construct, however, an additional ncRNA, CjNC140,
was noted to exhibit increased expression during exponential growth phase.
In distinct contrast to the lack of similar findings to previous studies exhibited by the
IA3902ΔluxS mutant, when the ΔCjNC110 and ΔluxS mutations were combined into a double
knockout mutant (ΔCjNC110ΔluxS), many of the previously noted transcriptional changes
attributed to mutation of the *luxS* gene alone became apparent [70]. The He *et al.* [70] study was
performed in a different strain of *C. jejuni*, 81-176, which also encodes and expresses CjNC110
[21]. After attempting to compare various strains and methods of mutation of the *luxS* gene and
finding wide variation in phenotypical and transcriptional changes, Adler *et al.* [28] suggested
that some of the differences reported between *luxS* mutants in various studies may be due to
unknown polar affects caused by alteration of expression of CjNC110 based on the mutation
strategy used for the *luxS* mutation. It seems highly likely based on the data generated in our
study that some of the changes attributed to *luxS* mutation alone in the He *et al.* [70] study may
in fact be due to inactivation of the CjNC110 transcript as well. Of the 57 genes listed as
differentially expressed by the *luxS* mutant in He *et al.*[70], 21 were also found to be
differentially expressed in our ΔCjNC110ΔluxS RNAseq data; again, none were found to be
differentially expressed only in the ΔluxS mutant, while in contrast, three genes (*tpx, ptmB,*
ptmA) were differentially expressed only in the ΔCjNC110 mutant. Of the 21 overlapping genes,
only the flagellar genes *flgE, flgG2,* and *flgG* were differentially expressed in both the ΔluxS and
ΔCjNC110ΔluxS mutants. These findings strongly suggest that in the previous study the main
driver for the transcriptional changes seen was an inadvertent inactivation of both CjNC110 and
luxS, and not *luxS* alone. Of the genes identified in both our study and He *et al.* [70], a large
number of the hook-basal body associated proteins (FlgD, FlgE, FlgG, FlgG2, FlgH, FlgI, and
FlgK) that are under the control of σ^{54} (RpoN) promotors [74, 75] were identified. Expression
from σ^{54} promotors has been shown to require activation of the FlgRS two-component regulatory
system [74, 76]. Many of these showed opposite differences based on growth phase (decreased
exponential, increased stationary), and it has been previously demonstrated that regulation by
these sigma-factors is growth cycle dependent, with σ^{54}-regulated genes typically expressed
between σ^{70}- and σ^{28}-associated genes [74], which may help explain the differences seen here. Of
particular note is the fact that *flgS* was observed to be upregulated in the ΔCjNC110ΔluxS
mutant. It has been previously suggested that an additional unknown factor may control the
temporal regulation of σ^{54} dependent flagellar genes [77]; therefore, it is reasonable to consider
based on our results that *luxS* or CjNC110 may play a role in this regulation.
Several additional phenotypes of interest were identified in our research, including
alteration of the ability to colonize chickens by ΔCjNC110, ΔluxS and ΔCjNC110ΔluxS. The
abolishment of the ability of ΔluxS to colonize chickens has previously been described in
IA3902 but not in NCTC 11168; the reason for this difference is currently unknown, but has
been speculated to be related to strain-specific differences in the metabolism of methionine or *S-*
adenosylmethionine (SAM) recycling which are also affected by the *luxS* mutation [27].
Additional work in our lab has shown that the *luxS* mutation in IA3902 does disrupt the AMC,
which functions to recycle SAM and produce methionine [49], however, DNA methylation in
IA3902 was not affected by the *luxS* mutation [73]. Comparative genomics reveals that NCTC
11168 does harbor an additional genomic system for methionine biosynthesis in the form of
metAB which is not present in IA3902 and may represent an alternative pathway for synthesis of

this required amino acid and an explanation for the observed difference in colonization ability.
Mutation of ΔCjNC110 in our work also led to a significantly decreased colonization ability in
IA3902 compared to wild-type, indicating that this ncRNA has the potential to play a key role in
modulating colonization by *C. jejuni*. Further work is ongoing in our lab to determine if mutation
of CjNC110 in NCTC 11168 also displays a colonization defect, or, if similar to *luxS,* the
phenotype differs from IA3902.
997   Motility is considered critical for the *in vivo* virulence of *C. jejuni* and requires a
functional flagellar apparatus [78, 79]; however, this alone is not sufficient for normal motility to
be present [80]. In our study, a statistically significant increase in motility was found between the
ΔCjNC110 mutant and wild-type when motility was assessed at 24 hours rather than 30 hours.
Interestingly, when assessing the transcriptomics data, the *cetAB* operon (*Campylobacter* energy
taxis proteins A and B), which is known to mediate energy taxis response in *Campylobacter*
[81]*,* was statistically significantly upregulated in the ΔCjNC110 mutant when compared to wild-
type during the exponential phase of growth. Expression of *cetAB* was increased during the
stationary phase as well, however, did not reach the level of statistical significance. Defects in
both *cetA* and *cetB* have been shown to lead to altered motility phenotypes, particularly in
response to migration towards critical factors in *Campylobacter* metabolism such as sodium
pyruvate and fumarate [80]. Investigation of expression of *cetA* and *cetB* has shown that levels of
the gene products are unaffected by mutation of sigma factors σ^{54} or σ^{28}, indicating that
transcription of the *cetAB* operon is likely controlled by σ^{70} or another yet unknown transcription
factor [81]. Based on this information, it seems plausible that CjNC110 might normally act as a
repressor of the CetAB energy taxis system. Attempts to determine the location of the interaction
between the *cetAB* operon and CjNC110 using computational prediction programs such as
TargetRNA2 failed to identify an interaction region, however, computational methods of target
identification are particularly unrewarding in non-model organisms such as *Campylobacter* due
to a lack of conservation of small RNAs in these species [18]. Our proteomics data also failed to
detect the CetA and CetB proteins, therefore, we could not validate that increased expression of
cetA and *cetB* resulted in increased protein expression levels. Many of the proteins identified in
our study exhibited extremely high expression levels, which, without fractioning, prevented
proteins of lesser abundance from being detected. Whole cell proteomic analysis with fractioning
and time-course analysis would be required to reveal the entire proteomic network regulated by
CjNC110 and validate expression patterns of proteins of lesser abundance than the 262 proteins
that were identified in our study. Further work to determine if an interaction between CjNC110
and the *cetAB* operon exists and leads to increased expression of the CetA and CetB proteins is
warranted.
Minimal changes in genes associated with the flagellar apparatus were noted in the
transcriptome of ΔCjNC110 which may also help to explain why an increase in energy taxis
would allow for an increase in observed motility with the assumption that normally functioning
flagella are present. In contrast to the ΔCjNC110 strain, however, both the ΔluxS and
ΔCjNC110ΔluxS mutants did display altered expression of genes associated with the flagellar
apparatus during at least one stage of growth. For the ΔluxS mutant, a small number of flagellar
associated genes were observed to be upregulated during stationary growth. No additional genes
known to be related to motility in *C. jejuni* appeared to be affected by the *luxS* mutation;
therefore, a reasonable explanation for the decreased motility cannot be ascertained from this
data alone. For the ΔCjNC110ΔluxS mutant, however, there were a large number of genes
involved in assembly of the flagellar apparatus that demonstrated dysregulation when both the

luxS and CjNC110 mutations were combined. Many genes related to flagellar assembly were
decreased in expression during exponential growth; however, the same genes were increased in
expression during stationary growth. Our data in this study demonstrated a statistically
significant decrease in motility of the ΔCjNC110ΔluxS mutant when compared to both wild-type
and the ΔCjNC110 mutant. This suggests that the decrease in motility is unrelated to the
CjNC110 mutation when present by itself, but when both the ΔluxS and ΔCjNC110 mutations
are combined, the effect of decreased motility observed in the *luxS* mutation is maintained.
The presence of normal flagella has also been associated with autoagglutination ability
and is considered necessary but not sufficient for this important trait [80]. Previous studies in *C.*
jejuni have demonstrated that interactions between modifications on adjacent flagellar filaments,
particularly those provided by protein glycosylation, are required for normal autoagglutination
ability [82]. In addition, mutants defective in autoagglutination ability have also been shown to
display a decrease in adherence and invasion ability *in vitro* as well as attenuation in disease
models [82]. In the present study, the ΔCjNC110 mutant exhibited decreased autoagglutination
when compared to wild-type at both 25ºC and 37ºC. Interestingly, two genes that have been
associated with flagellar glycosylation, *ptmA* and *neuB2*, were downregulated in our RNAseq
data in the ΔCjNC110 mutant when compared to wild-type during both exponential and
stationary growth phases. The *ptmA* and *neuB2* genes have been previously shown to be involved
in production of Leg5Am7Ac and PseAm, two important structural glycans involved in flagellar
glycosylation in *C. jejuni* [83, 84]. A decrease in the amount of flagellar glycosylation related to
decreased production of these genes could in theory lead to decreased autoagglutination ability.
The fact that we observed a decrease in autoagglutination activity in the ΔCjNC110 mutant,
combined with RNAseq data indicating a decrease in the presence of the mRNA transcripts of
the genes *ptmA* and *neuB2* that affect flagellar glycosylation, suggests that these findings
represent a true phenotypic change in the CjNC110 mutant and an additional potential area of
regulation for the CjNC110 non-coding RNA. In this case, and in contrast to the *cetAB* operon
where removal of CjNC110 led to an increase in expression, inactivation of CjNC110 led to a
decrease in the expression of these two genes which were predicted to belong to the same operon
in our Rockhopper analysis. In other bacterial species, small ncRNAs have been shown to lead to
stabilization of the target mRNA [85]. This suggests that CjNC110 may normally serve in *trans*
to prevent degradation of the *ptmA* and *neuB2* mRNA transcripts after they have been produced,
leading to increased longevity of the mRNA and thus increasing levels of the translated protein.
By removing CjNC110, stabilization of the mRNA message of *ptmA* and *neuB2* may not occur
as normal, leading to increased turnover of the message and decreased protein production. Again,
attempts to determine the location of the interaction between *ptmA*, *neuB2* and CjNC110 using
computational prediction programs such as TargetRNA2 failed to identify an interaction region,
and our proteomics data failed to identify either protein.
In contrast to ΔCjNC110, the ΔluxS mutant exhibited statistically significant increased
autoagglutination at both 25°C and 37°C. This finding is the opposite of what was observed in a
different strain of *C. jejuni,* 81-176*,* where the *luxS* mutation led to a decrease in
autoagglutination ability [71]. Mutation of the *luxS* gene by Jeon *et al.* [71] was accomplished
using the same method as He *et al.* [70], thus it is plausible that the results obtained may also be
a result of the *luxS* gene inactivation in combination with polar affects to CjNC110. However,
there are also known differences between strains of *C. jejuni* in the type of flagellar glycosylation
present, particularly between NCTC 11168 (considered sytenic to IA3902) and 81-176, which
may help to further explain the differences observed between autoagglutination studies

performed using different *C. jejuni* strains [86]. It is unclear at this time what mechanism
allowed for increased autoagglutination ability in the ΔluxS mutant in this particular strain of *C.*
jejuni. In addition, while these changes in autoagglutination are statistically significant, the
biological relevance at this time is unknown, particularly as decreased to absent colonization of
chickens was again confirmed in this mutant. When measured in terms of percent of normal
autoagglutination of the IA3902 wild-type strain, for the ΔluxS mutant, autoagglutination ability
was observed to be 172% and 168% of the wild-type at 25°C and 37°C, respectively. For the
ΔCjNC110 mutant, autoagglutination was observed to be 53% at 25°C and 56% at 37°C of the
wild-type. Following complementation of CjNC110, on a percentage basis autoagglutination
levels were much closer to wild-type demonstrating autoagglutination activity of 101% at 25ºC
and 107% at 37ºC. However, when compared to the percentage of autoagglutination possible,
which would be considered 100% at an A_{600} of 0.00 and 0% at an A_{600} of 1.0, both the wild-type
and mutant strains still exhibited a relatively high ability to autoagglutinate**.** For comparison,
mutations in the *flaA/flaB* and *flbA* genes of *C. jejuni* strain 81-176 led to a decrease from A_{600}
0.031 (96.9%) to 0.731 (24.9%) and 0.654 (35.6%), respectively [52]. While not reflected in the
transcriptomic data, our proteomic heat map data did show a decrease in FlaA translation in
ΔCjNC110, thus it is possible that the decreased autoagglutination observed in this mutant may
also be related to a decrease in FlaA protein abundance [52]. Differences in virulence displayed
in both *in vitro* and *in vivo* models are likely to be expected with such dramatic changes in AAG
activity as displayed in the *flaA/flab* and *flbA* mutations. As neither the *luxS* gene nor the
CjNC110 are known to be directly involved in flagellar glycosylation, it seems reasonable that
inactivation of these genes could lead to a modification of autoagglutination ability without
complete loss or gain of function. Further work including assessment for changes in the flagellar
glycosylation of these mutants when compared to the IA3902 wild-type and additional *in vitro*
and *in vivo* analysis of flagellar glycosylation is warranted.
Several additional putative ncRNAs were also identified via the Rockhopper program
which allowed for confirmation of expression of five non-coding RNAs conserved among
Campylobacter species that were previously predicted to be present in IA3902 [21]. The majority
of the other identified putative non-coding RNAs that were predicted by the program have not
been previously reported in *C. jejuni.* Based on the data presented in Dugar *et al.* [21] that
suggests that many small RNAs in *Campylobacter* are strain-specific, the fact that the predicted
RNAs were not previously identified does not mean that they are not real. That being said, there
are limitations to the identification of non-coding RNAs by Rockhopper. One-third of the
initially predicted ncRNAs were manually discarded as they mapped to the region of the 16S or
23S genes and were likely identified due to differences between replicates in Ribo-Zero rRNA
depletion efficiency. Another subset of the identified non-coding RNAs were recognized by the
program because the genome file utilized by Rockhopper does not include pseudogenes,
therefore, reads that mapped to areas of putative pseudogenes were classified as non-coding
RNA. While it is possible that these transcripts no longer serve as functional mRNA transcripts
and now instead act as functional non-coding RNAs, it is probably more likely that these
transcripts either do form functional proteins or are simply a non-functional remnant of what
once was a functional protein-coding message.
While not explored further in our phenotypic studies, Rockhopper did identify two
additional non-coding RNAs, CjNC140 and CjNC130/6S, as being differential expressed in our
mutants; these ncRNAs have been previously identified and confirmed to be transcribed in
multiple other *Campylobacter* species [21]. Of particular interest is the CjNC140 non-coding

RNA. Predicted to be transcribed in the intergenic region upstream of *porA*, this ncRNA was upregulated during exponential growth in all three of the mutant strains when compared to wild-type (**Fig D in S3 Appendix**). As increased expression was not observed during stationary growth, this suggests that CjNC140 may serve as a regulator involved in mediating changes during different stages of bacterial growth. The fact that its expression is similarly altered under all 3 mutant conditions also suggests that the three genes, *luxS*, CjNC110 and CjNC140, may normally interact within the cell in some way to facilitate these changes. Point mutations in the nearby *porA* have also been recently determined to be sufficient to cause the abortion phenotype [10], making this ncRNA particularly interesting for further study. Futher work is ongoing in our lab to determine the role of CjCN140 in the pathophysiology of *C. jejuni* as well.

Conclusion

The results of our research clearly demonstrate a significant role for CjNC110 in the pathobiology of *C. jejuni* IA3902. The collective results of the phenotypic and transcriptomic changes observed in our data complement each other and suggest that CjNC110 may be involved in regulation of energy taxis, flagellar glycosylation, cellular communication via quorum sensing, and chicken colonization in *C. jejuni* IA3902. This work provides for the first time valuable insights into the potential regulatory targets of the CjNC110 small RNA in the zoonotic pathogen *C. jejuni* and suggests a wide range of research avenues into the role of non-coding RNAs in the pathobiology of this important zoonotic pathogen.

Acknowledgements

The authors would like to thank Dr. Lei Dai and Dr. Zuowei Wu for technical assistance in mutant construction and northern blot technique.

References

1.   World Health Organization. WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007-2015. 2015 [cited 2015 December 3]. Available from: http://apps.who.int/iris/bitstream/10665/199350/1/9789241565165_eng.pdf.

2.   Sahin O, Plummer PJ, Jordan DM, Sulaj K, Pereira S, Robbe-Austerman S, et al. Emergence of a tetracycline-resistant Campylobacter jejuni clone associated with outbreaks of ovine abortion in the United States. J Clin Microbiol. 2008;46(5):1663-71. Epub 2008/03/07. doi: JCM.00031-08 [pii]10.1128/JCM.00031-08. PubMed PMID: 18322054; PubMed Central PMCID: PMC2395063.

3.   Sahin O, Fitzgerald C, Stroika S, Zhao S, Sippy RJ, Kwan P, et al. Molecular evidence for zoonotic transmission of an emergent, highly pathogenic Campylobacter jejuni clone in the United States. J Clin Microbiol. 2012;50(3):680-7. Epub 2011/12/23. doi: JCM.06167-11 [pii]

1168   10.1128/JCM.06167-11. PubMed PMID: 22189122; PubMed Central PMCID: PMC3295108.
1169

1170   4.   Tang Y, Meinersmann RJ, Sahin O, Wu Z, Dai L, Carlson J, et al. Wide but Variable
1171   Distribution of a Hypervirulent Campylobacter jejuni Clone in Beef and Dairy Cattle in the
1172   United States. Appl Environ Microbiol. 2017;83(24). doi: 10.1128/AEM.01425-17. PubMed
1173   PMID: 28970227; PubMed Central PMCID: PMCPMC5717212.
1174

1175   5.   Wu Z, Sahin O, Shen Z, Liu P, Miller WG, Zhang Q. Multi-omics approaches to
1176   deciphering a hypervirulent strain of Campylobacter jejuni. Genome biology and evolution.
1177   2013;5(11):2217-30. doi: 10.1093/gbe/evt172. PubMed PMID: 24201373; PubMed Central
1178   PMCID: PMC3845652.
1179

1180   6.   Grogono-Thomas R, Blaser MJ, Ahmadi M, Newell DG. Role of S-layer protein
1181   antigenic diversity in the immune responses of sheep experimentally challenged with
1182   Campylobacter fetus subsp. fetus. Infect Immun. 2003;71(1):147-54. PubMed PMID: 12496160;
1183   PubMed Central PMCID: PMC143156.
1184

1185   7.   van Putten JP, van Alphen LB, Wosten MM, de Zoete MR. Molecular mechanisms of
1186   campylobacter infection. Current topics in microbiology and immunology. 2009;337:197-229.
1187   doi: 10.1007/978-3-642-01846-6_7. PubMed PMID: 19812984.
1188

8.   Plummer P, Sahin O, Burrough E, Sippy R, Mou K, Rabenold J, et al. Critical role of
LuxS in the virulence of Campylobacter jejuni in a guinea pig model of abortion. Infect Immun.
2012;80(2):585-93. Epub 2011/12/07. doi: IAI.05766-11 [pii]10.1128/IAI.05766-11. PubMed
PMID: 22144479; PubMed Central PMCID: PMC3264297.

9.   Sahin O, Terhorst SA, Burrough ER, Shen Z, Wu Z, Dai L, et al. Key Role of Capsular
Polysaccharide in the Induction of Systemic Infection and Abortion by Hypervirulent
Campylobacter jejuni. Infect Immun. 2017;85(6). doi: 10.1128/IAI.00001-17. PubMed PMID:
28373351; PubMed Central PMCID: PMCPMC5442617.
1197

1198   10.   Wu Z, Periaswamy B, Sahin O, Yaeger M, Plummer P, Zhai W, et al. Point mutations in
1199   the major outer membrane protein drive hypervirulence of a rapidly expanding clone of
1200   Campylobacter jejuni. Proc Natl Acad Sci U S A. 2016;113(38):10690-5. doi:
1201   10.1073/pnas.1605869113. PubMed PMID: 27601641; PubMed Central PMCID:
1202   PMCPMC5035848.
1203

1204   11.   Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, et al. The genome
1205   sequence of the food-borne pathogen Campylobacter jejuni reveals hypervariable sequences.
1206   Nature. 2000;403(6770):665-8. doi: 10.1038/35001088. PubMed PMID: 10688204.
1207

1208   12.   Papenfort K, Vogel J. Regulatory RNA in bacterial pathogens. Cell host & microbe.
1209   2010;8(1):116-27. doi: 10.1016/j.chom.2010.06.008. PubMed PMID: 20638647.
1210

13.   Storz G, Vogel J, Wassarman KM. Regulation by small RNAs in bacteria: expanding
frontiers. Molecular cell. 2011;43(6):880-91. doi: 10.1016/j.molcel.2011.08.022. PubMed PMID:
21925377; PubMed Central PMCID: PMC3176440.

14.   Caldelari I, Chao Y, Romby P, Vogel J. RNA-mediated regulation in pathogenic bacteria.
Cold Spring Harbor perspectives in medicine. 2013;3(9):a010298. doi:
10.1101/cshperspect.a010298. PubMed PMID: 24003243.
1217
1218   15.   Beisel CL, Storz G. Base pairing small RNAs and their roles in global regulatory
1219   networks. FEMS microbiology reviews. 2010;34(5):866-82. doi: 10.1111/j.1574-
1220   6976.2010.00241.x. PubMed PMID: 20662934; PubMed Central PMCID: PMC2920360.
1221
16.   Sharma CM, Hoffmann S, Darfeuille F, Reignier J, Findeiss S, Sittka A, et al. The
primary transcriptome of the major human pathogen Helicobacter pylori. Nature.
2010;464(7286):250-5. Epub 2010/02/19. doi: nature08756 [pii]10.1038/nature08756. PubMed
PMID: 20164839.
17.   Valentin-Hansen P, Eriksen M, Udesen C. The bacterial Sm-like protein Hfq: a key
player in RNA transactions. Mol Microbiol. 2004;51(6):1525-33. PubMed PMID: 15009882.
1228
1229   18.   Livny J, Teonadi H, Livny M, Waldor MK. High-throughput, kingdom-wide prediction
1230   and annotation of bacterial non-coding RNAs. PLoS One. 2008;3(9):e3197. doi:
1231   10.1371/journal.pone.0003197. PubMed PMID: 18787707; PubMed Central PMCID:
1232   PMC2527527.
1233
19.   Chaudhuri RR, Yu L, Kanji A, Perkins TT, Gardner PP, Choudhary J, et al. Quantitative
RNA-seq analysis of the Campylobacter jejuni transcriptome. Microbiology. 2011;157(Pt
10):2922-32. Epub 2011/08/06. doi: mic.0.050278-0 [pii]10.1099/mic.0.050278-0. PubMed
PMID: 21816880; PubMed Central PMCID: PMC3353397.
20.   Butcher J, Stintzi A. The transcriptional landscape of Campylobacter jejuni under iron
replete and iron limited growth conditions. PLoS One. 2013;8(11):e79475. doi:
10.1371/journal.pone.0079475. PubMed PMID: 24223952; PubMed Central PMCID:
PMC3815106.
1242
21.   Dugar G, Herbig A, Forstner KU, Heidrich N, Reinhardt R, Nieselt K, et al. High-
resolution transcriptome maps reveal strain-specific regulatory features of multiple
Campylobacter jejuni isolates. PLoS Genet. 2013;9(5):e1003495. Epub 2013/05/23. doi:
10.1371/journal.pgen.1003495PGENETICS-D-13-00603 [pii]. PubMed PMID: 23696746;
PubMed Central PMCID: PMC3656092.
22.   Porcelli I, Reuter M, Pearson BM, Wilhelm T, van Vliet AH. Parallel evolution of
genome structure and transcriptional landscape in the Epsilonproteobacteria. BMC Genomics.
2013;14(1):616. Epub 2013/09/14. doi: 1471-2164-14-616 [pii]10.1186/1471-2164-14-616.
PubMed PMID: 24028687.
23.   Taveirne ME, Theriot CM, Livny J, Dirita VJ. The Complete Campylobacter jejuni
Transcriptome during Colonization of a Natural Host Determined by RNAseq. PLoS One.
2013;8(8):e73586. Epub 2013/08/31. doi: 10.1371/journal.pone.0073586PONE-D-13-20915
[pii]. PubMed PMID: 23991199; PubMed Central PMCID: PMC3749233.

24.   Kreuder AJ, Schleining JA, Yaeger M, Zhang Q, Plummer PJ. RNAseq Reveals Complex
Response of Campylobacter jejuni to Ovine Bile and In vivo Gallbladder Environment. Front
Microbiol. 2017;8:940. doi: 10.3389/fmicb.2017.00940. PubMed PMID: 28611744; PubMed
Central PMCID: PMCPMC5447181.
1260
1261   25.   Le MT, van Veldhuizen M, Porcelli I, Bongaerts RJ, Gaskin DJ, Pearson BM, et al.
1262   Conservation of sigma28-Dependent Non-Coding RNA Paralogs and Predicted sigma54-
1263   Dependent Targets in Thermophilic Campylobacter Species. PLoS One. 2015;10(10):e0141627.
1264   doi: 10.1371/journal.pone.0141627. PubMed PMID: 26512728; PubMed Central PMCID:
1265   PMC4626219.
1266
1267   26.   Shen Z, Patil RD, Sahin O, Wu Z, Pu XY, Dai L, et al. Identification and functional
1268   analysis of two toxin-antitoxin systems in Campylobacter jejuni. Mol Microbiol.
1269   2016;101(6):909-23. doi: 10.1111/mmi.13431. PubMed PMID: 27291507.
1270
1271   27.   Plummer PJ. LuxS and quorum-sensing in Campylobacter. Front Cell Infect Microbiol.
1272   2012;2:22. doi: 10.3389/fcimb.2012.00022. PubMed PMID: 22919614; PubMed Central
1273   PMCID: PMCPMC3417632.
1274
1275   28.   Adler L, Alter T, Sharbati S, Golz G. Phenotypes of Campylobacter jejuni luxS mutants
1276   are depending on strain background, kind of mutation and experimental conditions. PLoS One.
1277   2014;9(8):e104399. doi: 10.1371/journal.pone.0104399. PubMed PMID: 25093839; PubMed
1278   Central PMCID: PMC4122453.
1279
1280   29.   Bassler BL, Wright M, Showalter RE, Silverman MR. Intercellular signalling in Vibrio
1281   harveyi: sequence and function of genes regulating expression of luminescence. Mol Microbiol.
1282   1993;9(4):773-86. PubMed PMID: 8231809.
1283
1284   30.   Plummer P, Zhu J, Akiba M, Pei D, Zhang Q. Identification of a key amino acid of LuxS
1285   involved in AI-2 production in Campylobacter jejuni. PLoS One. 2011;6(1):e15876. Epub
1286   2011/01/26. doi: 10.1371/journal.pone.0015876. PubMed PMID: 21264316; PubMed Central
1287   PMCID: PMC3019222.
1288
1289   31.   Bassler BL, Wright M, Silverman MR. Multiple signalling systems controlling
1290   expression of luminescence in Vibrio harveyi: sequence and function of genes encoding a second
1291   sensory pathway. Mol Microbiol. 1994;13(2):273-86. PubMed PMID: 7984107.
1292
1293   32.   Wang Y, Taylor DE. Chloramphenicol resistance in Campylobacter coli: nucleotide
1294   sequence, expression, and cloning vector construction. Gene. 1990;94(1):23-8. PubMed PMID:
1295   2227449.
1296
1297   33.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. Enzymatic
1298   assembly of DNA molecules up to several hundred kilobases. Nature methods. 2009;6(5):343-5.
1299   doi: 10.1038/nmeth.1318. PubMed PMID: 19363495.
1300

1301   34.   Wang Y, Taylor DE. Natural transformation in Campylobacter species. J Bacteriol.
1302   1990;172(2):949-55. PubMed PMID: 2404960; PubMed Central PMCID: PMC208523.
1303
1304   35.   Kim SW, Li Z, Moore PS, Monaghan AP, Chang Y, Nichols M, et al. A sensitive non-
1305   radioactive northern blot method to detect small RNAs. Nucleic Acids Res. 2010;38(7):e98. doi:
1306   10.1093/nar/gkp1235. PubMed PMID: 20081203; PubMed Central PMCID: PMCPMC2853138.
1307
36.   Muraoka WT, Zhang Q. Phenotypic and genotypic evidence for L-fucose utilization by
Campylobacter jejuni. J Bacteriol. 2011;193(5):1065-75. Epub 2011/01/05. doi: JB.01252-10
[pii]10.1128/JB.01252-10. PubMed PMID: 21193610; PubMed Central PMCID: PMC3067607.
37.   Karlyshev AV, Wren BW. Development and application of an insertional system for gene
delivery and expression in Campylobacter jejuni. Appl Environ Microbiol. 2005;71(7):4004-13.
Epub 2005/07/08. doi: 71/7/4004 [pii]10.1128/AEM.71.7.4004-4013.2005. PubMed PMID:
16000815; PubMed Central PMCID: PMC1169003.
38.   Chen CY, Nace GW, Irwin PL. A 6 x 6 drop plate method for simultaneous colony
counting and MPN enumeration of Campylobacter jejuni, Listeria monocytogenes, and
Escherichia coli. Journal of microbiological methods. 2003;55(2):475-9. PubMed PMID:
14529971.
1319
1320   39.   Luo Y, Sahin O, Dai L, Sippy R, Wu Z, Zhang Q. Development of a loop-mediated
1321   isothermal amplification assay for rapid, sensitive and specific detection of a Campylobacter
1322   jejuni clone. The Journal of veterinary medical science / the Japanese Society of Veterinary
1323   Science. 2012;74(5):591-6. PubMed PMID: 22188995.
1324
1325   40.   McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA, et al.
1326   Computational analysis of bacterial RNA-Seq data. Nucleic Acids Res. 2013;41(14):e140. doi:
1327   10.1093/nar/gkt444. PubMed PMID: 23716638; PubMed Central PMCID: PMC3737546.
1328
1329   41.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
1330   Integrative genomics viewer. Nature biotechnology. 2011;29(1):24-6. doi: 10.1038/nbt.1754.
1331   PubMed PMID: 21221095; PubMed Central PMCID: PMC3346182.
1332
1333   42.   Thorvaldsdottir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-
1334   performance genomics data visualization and exploration. Briefings in bioinformatics.
1335   2013;14(2):178-92. doi: 10.1093/bib/bbs017. PubMed PMID: 22517427; PubMed Central
1336   PMCID: PMC3603213.
1337
1338   43.   Galperin MY, Makarova KS, Wolf YI, Koonin EV. Expanded microbial genome
1339   coverage and improved protein family annotation in the COG database. Nucleic Acids Res.
1340   2015;43(Database issue):D261-9. doi: 10.1093/nar/gku1223. PubMed PMID: 25428365;
1341   PubMed Central PMCID: PMC4383993.
1342
1343   44.   Hulsen T, de Vlieg J, Alkema W. BioVenn - a web application for the comparison and
1344   visualization of biological lists using area-proportional Venn diagrams. BMC Genomics.

1345   2008;9:488. doi: 10.1186/1471-2164-9-488. PubMed PMID: 18925949; PubMed Central
1346   PMCID: PMC2584113.

1347

1348   45.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference
1349   resource for gene and protein annotation. Nucleic Acids Res. 2015. doi: 10.1093/nar/gkv1070.
1350   PubMed PMID: 26476454.

1351

1352   46.   Kery MB, Feldman M, Livny J, Tjaden B. TargetRNA2: identifying targets of small
1353   regulatory RNAs in bacteria. Nucleic Acids Res. 2014;42(Web Server issue):W124-9. doi:
1354   10.1093/nar/gku317. PubMed PMID: 24753424.

1355

1356   47.   Gruber AR, Lorenz R, Bernhart SH, Neubock R, Hofacker IL. The Vienna RNA
1357   websuite. Nucleic Acids Res. 2008;36(Web Server issue):W70-4. doi: 10.1093/nar/gkn188.
1358   PubMed PMID: 18424795; PubMed Central PMCID: PMC2447809.

1359

1360   48.   Lorenz R, Bernhart SH, Honer Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et al.
1361   ViennaRNA Package 2.0. Algorithms for molecular biology : AMB. 2011;6:26. doi:
1362   10.1186/1748-7188-6-26. PubMed PMID: 22115189; PubMed Central PMCID: PMC3319429.

1363

1364   49.   Mou KT, Plummer PJ. The impact of the LuxS mutation on phenotypic expression of
1365   factors critical for Campylobacter jejuni colonization. Vet Microbiol. 2016;192:43-51. doi:
1366   10.1016/j.vetmic.2016.06.011. PubMed PMID: 27527763.

1367

1368   50.   Gallup JM, Ackermann MR. Addressing fluorogenic real-time qPCR inhibition using the
1369   novel custom Excel file system 'FocusField2-6GallupqPCRSet-upTool-001' to attain consistently
1370   high fidelity qPCR reactions. Biol Proced Online. 2006;8:87-152. doi: 10.1251/bpo122. PubMed
1371   PMID: 17033699; PubMed Central PMCID: PMCPMC1592462.

1372

1373   51.   Surette MG, Bassler BL. Quorum sensing in Escherichia coli and Salmonella
1374   typhimurium. Proc Natl Acad Sci U S A. 1998;95(12):7046-50. PubMed PMID: 9618536;
1375   PubMed Central PMCID: PMC22733.

1376

1377   52.   Misawa N, Blaser MJ. Detection and characterization of autoagglutination activity by
1378   Campylobacter jejuni. Infect Immun. 2000;68(11):6168-75. PubMed PMID: 11035721; PubMed
1379   Central PMCID: PMC97695.

1380

1381   53.   Hobb RI, Fields JA, Burns CM, Thompson SA. Evaluation of procedures for outer
1382   membrane isolation from Campylobacter jejuni. Microbiology. 2009;155(Pt 3):979-88. doi:
1383   10.1099/mic.0.024539-0. PubMed PMID: 19246768; PubMed Central PMCID:
1384   PMCPMC2763183.

1385

54.   Chang C, Xu K, Guo C, Wang J, Yan Q, Zhang J, et al. PANDA-view: an easy-to-use
tool for statistical analysis and visualization of quantitative proteomics data. Bioinformatics.
2018;34(20):3594-6. doi: 10.1093/bioinformatics/bty408. PubMed PMID: 29790911; PubMed
Central PMCID: PMCPMC6184437.

55. Beretta L, Santaniello A. Nearest neighbor imputation algorithms: a critical evaluation.
BMC Med Inform Decis Mak. 2016;16 Suppl 3:74. doi: 10.1186/s12911-016-0318-z. PubMed
PMID: 27454392; PubMed Central PMCID: PMCPMC4959387.

1393
1394 56. Chong J, Yamamoto M, Xia J. MetaboAnalystR 2.0: From Raw Spectra to Biological
1395 Insights. Metabolites. 2019;9(3). doi: 10.3390/metabo9030057. PubMed PMID: 30909447;
1396 PubMed Central PMCID: PMCPMC6468840.

1397
1398 57. Key M. A tutorial in displaying mass spectrometry-based proteomic data using heat
1399 maps. BMC bioinformatics. 2012;13 Suppl 16:S10. doi: 10.1186/1471-2105-13-S16-S10.
1400 PubMed PMID: 23176119; PubMed Central PMCID: PMCPMC3489527.

1401
58. Croucher NJ, Thomson NR. Studying bacterial transcriptomes using RNA-seq. Curr Opin
Microbiol. 2010;13(5):619-24. Epub 2010/10/05. doi: S1369-5274(10)00136-0
[pii]10.1016/j.mib.2010.09.009. PubMed PMID: 20888288; PubMed Central PMCID:
PMC3025319.

59. van Vliet AH. Next generation sequencing of microbial transcriptomes: challenges and
opportunities. FEMS Microbiol Lett. 2010;302(1):1-7. Epub 2009/09/09. doi: FML1767
[pii]10.1111/j.1574-6968.2009.01767.x. PubMed PMID: 19735299.

60. van Opijnen T, Camilli A. Transposon insertion sequencing: a new tool for systems-level
analysis of microorganisms. Nat Rev Microbiol. 2013;11(7):435-42. doi: 10.1038/nrmicro3033.
PubMed PMID: 23712350; PubMed Central PMCID: PMC3842022.

1412
1413 61. Butcher J, Handley RA, van Vliet AH, Stintzi A. Refined analysis of the Campylobacter
1414 jejuni iron-dependent/independent Fur- and PerR-transcriptomes. BMC Genomics. 2015;16:498.
1415 doi: 10.1186/s12864-015-1661-7. PubMed PMID: 26141822; PubMed Central PMCID:
1416 PMC4491227.

1417
1418 62. Chandrashekhar K, Kassem, II, Nislow C, Gangaiah D, Candelero-Rueda RA,
1419 Rajashekara G. Transcriptome analysis of Campylobacter jejuni polyphosphate kinase (ppk1 and
1420 ppk2) mutants. Virulence. 2015;6(8):814-8. doi: 10.1080/21505594.2015.1104449. PubMed
1421 PMID: 26537695.

1422
1423 63. Waters LS, Storz G. Regulatory RNAs in bacteria. Cell. 2009;136(4):615-28. doi:
1424 10.1016/j.cell.2009.01.043. PubMed PMID: 19239884; PubMed Central PMCID: PMC3132550.

1425
1426 64. Herzberg M, Kaye IK, Peti W, Wood TK. YdgG (TqsA) controls biofilm formation in
1427 Escherichia coli K-12 through autoinducer 2 transport. J Bacteriol. 2006;188(2):587-98. doi:
1428 10.1128/JB.188.2.587-598.2006. PubMed PMID: 16385049; PubMed Central PMCID:
1429 PMCPMC1347309.

1430
65. Golz G, Sharbati S, Backert S, Alter T. Quorum sensing dependent phenotypes and their
molecular mechanisms in Campylobacterales. Eur J Microbiol Immunol (Bp). 2012;2(1):50-60.
doi: 10.1556/EuJMI.2.2012.1.8. PubMed PMID: 24611121; PubMed Central PMCID:
PMCPMC3933990.

66.   Xavier KB, Bassler BL. Regulation of uptake and processing of the quorum-sensing
autoinducer AI-2 in Escherichia coli. J Bacteriol. 2005;187(1):238-48. doi:
10.1128/JB.187.1.238-248.2005. PubMed PMID: 15601708; PubMed Central PMCID:
PMCPMC538819.

67.   Holmes K, Tavender TJ, Winzer K, Wells JM, Hardie KR. AI-2 does not function as a
quorum sensing molecule in Campylobacter jejuni during exponential growth in vitro. BMC
Microbiol. 2009;9:214. doi: 10.1186/1471-2180-9-214. PubMed PMID: 19814796; PubMed
Central PMCID: PMC2772989.

68.   Adler L, Alter T, Sharbati S, Golz G. The signalling molecule Autoinducer-2 is not
internalised in Campylobacter jejuni. Berl Munch Tierarztl Wochenschr. 2015;128(3-4):111-6.
PubMed PMID: 25876270.

69.   Plummer PJ. The mechanisms and roles of quorum sensing in the pathobiology of
Campylobacter jejuni. Graduate Theses and Disseratations: Iowa State University; 2009.

70.   He Y, Frye JG, Strobaugh TP, Chen CY. Analysis of AI-2/LuxS-dependent transcription
in Campylobacter jejuni strain 81-176. Foodborne pathogens and disease. 2008;5(4):399-415.
doi: 10.1089/fpd.2008.0106. PubMed PMID: 18713060.

71.   Jeon B, Itoh K, Misawa N, Ryu S. Effects of quorum sensing on flaA transcription and
autoagglutination in Campylobacter jejuni. Microbiology and immunology. 2003;47(11):833-9.
PubMed PMID: 14638994.

72.   Parveen N, Cornell KA. Methylthioadenosine/S-adenosylhomocysteine nucleosidase, a
critical enzyme for bacterial metabolism. Mol Microbiol. 2011;79(1):7-20. doi: 10.1111/j.1365-
2958.2010.07455.x. PubMed PMID: 21166890; PubMed Central PMCID: PMC3057356.

73.   Mou KT. Mechanisms and roles of the LuxS system, methyl recycling, and DNA
methylation on the physiology of Campylobacter jejuni. Graduate Theses and Disserations: Iowa
State University; 2015.

74.   Wosten MM, Wagenaar JA, van Putten JP. The FlgS/FlgR two-component signal
transduction system regulates the fla regulon in Campylobacter jejuni. The Journal of biological
chemistry. 2004;279(16):16214-22. doi: 10.1074/jbc.M400357200. PubMed PMID: 14960570.

75.   Malik-Kale P, Parker CT, Konkel ME. Culture of Campylobacter jejuni with sodium
deoxycholate induces virulence gene expression. J Bacteriol. 2008;190(7):2286-97. Epub
2008/01/29. doi: JB.01736-07 [pii]10.1128/JB.01736-07. PubMed PMID: 18223090; PubMed
Central PMCID: PMC2293209.

76.   Joslin SN, Hendrixson DR. Activation of the Campylobacter jejuni FlgSR two-
component system is linked to the flagellar export apparatus. J Bacteriol. 2009;191(8):2656-67.
doi: 10.1128/JB.01689-08. PubMed PMID: 19201799; PubMed Central PMCID: PMC2668382.

77.   Hendrixson DR, DiRita VJ. Transcription of sigma54-dependent but not sigma28-
dependent flagellar genes in Campylobacter jejuni is associated with formation of the flagellar
secretory apparatus. Mol Microbiol. 2003;50(2):687-702. PubMed PMID: 14617189.
1482
1483   78.   Wassenaar TM, Bleumink-Pluym NM, van der Zeijst BA. Inactivation of Campylobacter
1484   jejuni flagellin genes by homologous recombination demonstrates that flaA but not flaB is
1485   required for invasion. The EMBO journal. 1991;10(8):2055-61. PubMed PMID: 2065653;
1486   PubMed Central PMCID: PMC452888.
1487
1488   79.   Yao R, Burr DH, Doig P, Trust TJ, Niu H, Guerry P. Isolation of motile and non-motile
1489   insertional mutants of Campylobacter jejuni: the role of motility in adherence and invasion of
1490   eukaryotic cells. Mol Microbiol. 1994;14(5):883-93. PubMed PMID: 7715450.
1491
1492   80.   Golden NJ, Acheson DW. Identification of motility and autoagglutination Campylobacter
1493   jejuni mutants by random transposon mutagenesis. Infect Immun. 2002;70(4):1761-71. PubMed
1494   PMID: 11895937; PubMed Central PMCID: PMC127829.
1495
1496   81.   Hendrixson DR, Akerley BJ, DiRita VJ. Transposon mutagenesis of Campylobacter
1497   jejuni identifies a bipartite energy taxis system required for motility. Mol Microbiol.
1498   2001;40(1):214-24. PubMed PMID: 11298288.
1499
1500   82.   Guerry P, Ewing CP, Schirm M, Lorenzo M, Kelly J, Pattarini D, et al. Changes in
1501   flagellin glycosylation affect Campylobacter autoagglutination and virulence. Mol Microbiol.
1502   2006;60(2):299-311. doi: 10.1111/j.1365-2958.2006.05100.x. PubMed PMID: 16573682;
1503   PubMed Central PMCID: PMC1424674.
1504
1505   83.   Logan SM, Kelly JF, Thibault P, Ewing CP, Guerry P. Structural heterogeneity of
1506   carbohydrate modifications affects serospecificity of Campylobacter flagellins. Mol Microbiol.
1507   2002;46(2):587-97. PubMed PMID: 12406231.
1508
1509   84.   McNally DJ, Aubry AJ, Hui JP, Khieu NH, Whitfield D, Ewing CP, et al. Targeted
1510   metabolomics analysis of Campylobacter coli VC167 reveals legionaminic acid derivatives as
1511   novel flagellar glycans. The Journal of biological chemistry. 2007;282(19):14463-75. doi:
1512   10.1074/jbc.M611027200. PubMed PMID: 17371878.
1513
1514   85.   Papenfort K, Vanderpool CK. Target activation by regulatory RNAs in bacteria. FEMS
1515   microbiology reviews. 2015;39(3):362-78. doi: 10.1093/femsre/fuv016. PubMed PMID:
1516   25934124; PubMed Central PMCID: PMC4542691.
1517
1518   86.   Merino S, Tomas JM. Gram-negative flagella glycosylation. International journal of
1519   molecular sciences. 2014;15(2):2840-57. doi: 10.3390/ijms15022840. PubMed PMID:
1520   24557579; PubMed Central PMCID: PMC3958885.
1521
1522
1523

Supporting information

S1 Appendix: Supporting Tables A through I

Table A. Plasmids used in this study.

Table B. Primers used in this study.

Table C. Summary of RNAseq results following rRNA depletion and strand specific library preparation on the Illumina HiSeq 2500 – successfully aligned reads.

Table D. Summary of RNAseq results following rRNA depletion and strand specific library preparation on the Illumina HiSeq 2500 – read mapping results.

Table E. Differential gene expression in the IA3902 ΔCjNC110 mutant as determined by RNAseq.

Table F. Differential gene expression in the IA3902 ΔluxS mutant as determined by RNAseq.

Table G. Differential gene expression in the IA3902 ΔCjNC110ΔluxS mutant as determined by RNAseq.

Table H. List of non-coding RNAs predicted by Rockhopper to exist using RNAseq transcriptomics.

Table I. Targets predicted for CjNC110 in *C. jejuni* IA3902, 137nt and 152nt predicted length, by TargetRNA2.

S2 Appendix: Suuporting Table J. List of operons in *C. jejuni* IA3902 predicted by Rockhopper to exist using RNAseq transcriptomics.

S3 Appendix: Supporting Figures A through D

Fig A. Shaking growth curve utilized for RNA extraction for RNAseq of wild-type IA3902 and isogenic mutants (mean ± SEM). A_{600} (A) and log10 CFU/mL (B) results of four replicates of a shaking growth curve performed in 250 mL Erlenmeyer flasks under microaerophilic conditions in MH broth. Analysis via two-way ANOVA revealed a statistically significant difference between strains ($p < 0.05$), however, multiple comparison analysis of individual time points and strains when compared to wild type growth revealed that the only time point where the differences was considered to be statistically significant was at 30 hours for all strains (denoted by *).

Fig B. COG functional categories of differentially expressed genes in the ΔCjNC110ΔluxS mutant. Clusters of Orthologous Groups (COG) categories are indicated on the x-axis, with the number of genes enriched shown on the y-axis; blue bars indicate increased expression, red bars indicate decreased expression. C - Energy

production and conversion; D - Cell cycle control, mitosis and meiosis; E - Amino acid
transport and metabolism; F - Nucleotide transport and metabolism; G - Carbohydrate
transport and metabolism; H - Coenzyme transport and metabolism; I - Lipid transport
and metabolism; J – Translation; K – Transcription; L - Replication, recombination and
repair; M - Cell wall/membrane biogenesis; N - Cell motility; O - Posttranslational
modification, protein turnover, chaperones; P - Inorganic ion transport and metabolism;
Q - Secondary metabolites biosynthesis, transport and catabolism; R - General function
prediction only; S - Function unknown; T - Signal transduction mechanisms; U -
Intracellular trafficking and secretion; V - Defense mechanisms; W - Extracellular
structures

1580

1581   **Fig C. KEGG Pathway for flagellar assembly in *C. jejuni:* genes affected by the**
1582   **single mutation of either ΔCjNC110 (A) or ΔluxS (B) mutation during stationary**
1583   **growth phase.** Blue color indicates downregulation of gene expression, red color
1584   indicates up regulation of gene expression, green indicates that the gene is present in *C.*
1585   *jejuni* IA3902, and white indicates that the gene is not present in IA3902.

1586

1587   **Fig D. Graphic view of CjNC140 expression from RNAseq**. (A) A screen capture from
1588   IGV of the *porA* and CjNC140 regions of the genome, corresponding to the genome
1589   structure as depicted in (B), of all of the strains sequenced using RNAseq.

1590

1591

1592

Ladder	A.	B.	C.
Strain	IA3902 WT	ΔCjNC110c	ΔCjNC110
Total RNA	15 μg/well	15 μg/well	15 μg/well

Shared upregulated genes

Exponential

Stationary

Shared downregulated genes

ΔCjNC110

10   1

ΔluxS →   1

ΔCjNC110ΔluxS

Exponential

ΔCjNC110

1   ΔluxS

ΔCjNC110ΔluxS

Stationary

152 nt length

137 nt length

0 ────── 1