Small non-coding RNA CjNC110 influences motility, autoagglutination, AI-2 localization,

hydrogen peroxide sensitivity and chicken colonization in *Campylobacter jejuni*

Amanda J. Kreuder[a,b#], Brandon Ruddell[b], Kathy Mou[b*], Alan Hassall[a,b], Qijing Zhang[b], Paul J.

Plummer[a,b]

[a] Department of Veterinary Diagnostic and Production Animal Medicine, College of Veterinary

Medicine, Iowa State University, Ames, IA, USA

[b] Department of Veterinary Microbiology and Preventive Medicine, College of Veterinary

Medicine, Iowa State University, Ames, IA, USA

Running Head: CjNC110 small RNA in *Campylobacter jejuni*

#Address correspondence to Amanda J. Kreuder, akreuder@iastate.edu

* Present Address: United States Department of Agriculture, Agricultural Research Service,

National Animal Disease Center, Ames, IA United States; and Oak Ridge Institute for Science

and Education, Oak Ridge, TN, United States

Amanda J. Kreuder and Brandon Ruddell contributed equally to this work. Author order was

determined on the basis of seniority.

ABSTRACT

Small non-coding RNAs are involved in many important physiological functions in pathogenic microorganisms. Previous studies have identified the presence of non-coding RNAs in the major zoonotic pathogen *Campylobacter jejuni*, however, few have been functionally characterized to date. CjNC110 is a conserved ncRNA in *C. jejuni,* located downstream of the *luxS* gene which is responsible for the production of the quorum-sensing molecule autoinducer-2 (AI-2). In this study, we utilized strand specific high-throughput RNAseq to identify potential targets or interactive partners of CjNC110 in a sheep abortion clone of *C. jejuni*. This data was then utilized to focus further phenotypic evaluation of the role of CjNC110 in motility, autoagglutination, quorum sensing, hydrogen peroxide sensitivity and chicken colonization in *C. jejuni*. Inactivation of the CjNC110 ncRNA led to a statistically significant decrease in autoagglutination ability as well as increased motility and hydrogen peroxide sensitivity when compared to wild-type. Extracellular AI-2 detection was decreased in ΔCjNC110, however, intracellular AI-2 accumulation was significantly increased, suggesting a key role of CjNC110 in modulating the transport of AI-2. Notably, ΔCjNC110 also showed a decreased ability to colonize chickens. Complementation of CjNC110 restored all phenotypic changes back to wild-type levels. The collective results of the phenotypic and transcriptomic changes observed in our data provide valuable insights into the pathobiology of *C. jejuni* sheep abortion clone and strongly suggest that CjNC110 plays an important role in regulation of energy taxis, flagellar glycosylation, cellular communication via quorum sensing, oxidative stress tolerance and chicken colonization in this important zoonotic pathogen.

INTRODUCTION

Campylobacter jejuni is a major foodborne and zoonotic pathogen that causes enteritis in

humans (1). A sheep abortion (SA) clone, with IA3902 as the prototype strain, has recently

emerged as the predominant cause of ovine abortion and as an important pathogen in foodborne

outbreaks of human gastroenteritis in the United States (2, 3). *C. jejuni* clone SA is also widely

distributed in the cattle population in the U.S. (4). The genome of IA3902 is quite syntenic with

that of *C. jejuni* type-strain NCTC 11168 although it is much more virulent than NCTC 11168 in

inducing systemic infection and abortion in animals (5). Despite its hypervirulence, *C. jejuni*

IA3902 does not harbor any virulence factors known to be associated with abortion in *C. fetus*

subsp. *fetus* (6, 7). Previously, the *luxS* gene, which mediates autoinducer-2 (AI-2) production,

and the genes encoding the capsular polysaccharide have been identified as critical in IA3902 for

intestinal colonization and/or translocation across gut epithelium into the bloodstream (8, 9).

Point mutations in the major outer membrane protein encoded by *porA* of IA3902 have also been

shown to be sufficient to cause the abortion phenotype when compared to the syntenic non-

abortive strain NCTC 11168 (10). The fact that relatively mild changes in genomic sequences

have led to significantly enhanced ability to cause disease by *C. jejuni* IA3902, as described

above, suggests that even slight differences in sequence variations or gene regulation may have a

major impact on virulence variation among differing strains of *C. jejuni*.

C. jejuni has only three known sigma factors that regulate gene transcription: σ^{70}

(encoded by *rpoD*), σ^{54} (encoded by *rpoN*) and σ^{28} (encoded by *fliA*) (11). Besides control at the

transcriptional level, regulation of gene expression can also occur by post-transcriptional

modulation of mRNA translation, stability and processing, for which small non-coding RNAs are

the primary players (12-14). Non-coding RNAs (ncRNAs) can be rapidly produced and serve to

regulate multiple different targets within a cell in a variety of ways to coordinate rapid responses

to changing environments. Thus, regulation of cellular processes by ncRNAs can provide

several advantages to the bacteria when compared to the traditional model of protein-mediated

regulation (15).

Prior to completion of the transcriptional start site map via high throughput RNA

sequencing (RNAseq) of *H. pylori* (16), the ε-proteobacteria were thought not to be capable of

using small and antisense non-coding RNA as a regulatory mechanism, partly due to a lack of

the RNA chaperone protein Hfq (17). Indeed, initial attempts using computational approaches to

identify ncRNAs in *Campylobacter* failed to yield any potential candidates, with only 3 potential

loci being identified in *Helicobacter* (18). Recently, however, clear evidence that *C. jejuni* also

harbors these important regulators has been published, revealing the existence and expression of

a wealth of ncRNAs present in strains such as NCTC 11168, 81-176, 81116, RM1221 and

IA3902 (19-24). Dugar *et al.* (21), when comparing the transcriptomes of 4 different *C. jejuni*

isolates, observed a large variation in transcriptional start sites as well as expression patterns of

both mRNA and ncRNA between strains. This suggests that variations in the existence and

expression of ncRNAs even among closely related strains may play a role in differentiating

virulence. Despite these advances, identification of the functions of ncRNAs in *Campylobacter*

has been slow to follow, with most lacking both functional characterization as well as

mechanistic investigation.

The first report attempting to elucidate the function of two previously identified non-

coding RNAs in *C. jejuni* suggests these ncRNAs may play a role in flagellar biosynthesis;

however, the authors were unable to demonstrate phenotypic changes following inactivation of

either non-coding RNA (25). A later paper, however, did establish the role of an RNA antitoxin

(*cjrA*) as the first non-coding antisense small RNA functionally characterized in *Campylobacter*

to date (26). Just recently, the small RNA pair, CjNC180/CjNC190, has been identified to alter

colonization in a newly developed 3D *in vitro* model of disease in strain NCTC 11168 (27); the

mechanism by which this occurs remains to be fully determined. Thus, beyond simply

establishing the existence of non-coding RNA transcripts in *Campylobacter*, there is critical need

to continue to determine the physiological functions of these potential regulators in this

important zoonotic pathogen.

Our previous work has demonstrated the *in vivo* and *in vitro* expression of numerous

ncRNAs in IA3902 (24), including several that are conserved in other strains of *C. jejuni* (21).

One of the expressed ncRNAs identified in our study, the conserved small RNA CjNC110, is

located in the intergenic region immediately downstream of the *luxS* gene. This ncRNA was of

particular interest to our group as previous work in our lab has already highlighted the

importance of the *luxS* gene in the virulence of *C. jejuni* IA3902 (8). In *Campylobacter*, the *luxS*

gene is known to serve two important functions, production of the quorum sensing molecule

autoinducer-2 (AI-2) and conversion of *S*-ribosylhomocysteine to homocysteine in the activated

methyl cycle (AMC) (28). Of particular note, following the identification of CjNC110, it has also

been suggested that different methods of generation of the *luxS* mutation may have led to polar

effects on this downstream ncRNA that may help explain observed differences in phenotypes

and gene expression between various studies of *luxS* mutants in *Campylobacter* (29).

Based on these observations, in this study we chose to focus on beginning to elucidate the

functional role of CjNC110 in the pathobiology of *C. jejuni* IA3902, as well as investigate the

transcriptomic and phenotypic differences between single mutation of CjNC110 and *luxS* versus

co-mutation of both genomic regions. Using high-throughput RNA sequencing (RNAseq), we

successfully demonstrated differences in the transcriptional landscape following mutagenesis of

CjNC110 and identified a number of potential mRNA targets for CjNC110 regulation. From

these potential targets, we then demonstrated that inactivation of CjNC110 affects several

important phenotypes in IA3902, including motility, autoagglutination activity, AI-2

localization, hydrogen peroxide sensitivity, and chicken colonization; our results also

demonstrate that these phenotypes consistently differ from those of seen with inactivation of

luxS. The work presented here provides compelling evidence that expression of the ncRNA

CjNC110 is important for the pathobiology of *C. jejuni* IA3902, and lays the foundation for

future work investigating the mechanism of regulation of post-transcriptional gene expression in

IA3902 by CjNC110.

RESULTS

Northern blot analysis validates expression of small ncRNA CjNC110 in IA3902

wild-type and mutant constructs ΔluxS and ΔCjNC110c, and absence in ΔCjNC110.

Expression of CjNC110 had been previously validated via northern blot analysis in several

strains of *C. jejuni*, including NCTC 11168, RM1221, 81-176, and 81116, and further genetic

analysis suggested CjNC110 to be present in IA3902 as well (21). Therefore, to begin to study

the role of CjNC110 in IA3902, insertional deletion of the entire predicted coding sequence was

utilized to construct ΔCjNC110 in IA3902, and complementation was achieved via re-insertion

of CjNC110 into the 16S and 23S rRNA operon (*rrs-rrl*) of ΔCjNC110 via homologous

recombination using plasmid pRRK, creating ΔCjNC110c (**Table 1** lists all strains used in this

study). To validate expression of CjNC110 in IA3902, an initial northern blot using the same

probe sequence as Dugar *et al*. (21) and 15 μg of total RNA extracted at early stationary phase of

growth from cultures collected at the same A_{600} was performed to compare IA3902 wild-type,

ΔCjNC110 and ΔCjNC110c. The results of this northern blot analysis clearly demonstrate that

small ncRNA CjNC110 is expressed by IA3902 wild-type and ΔCjNC110c but not ΔCjNC110

(**Fig. 1**), validating expression of CjNC110 in IA3902 and confirming successful mutant and

complement constructs. Visual inspection of the northern blot data from both Dugar *et al.* and

the work presented here also demonstrates the presence of multiple bands of differing sizes in

both studies (21). All bands were eliminated in the ΔCjNC110 mutant and restored in the

ΔCjNC110c complement, which suggests processing of the CjNC110 transcript may be

occurring; further work is needed to investigate this observation.

As previous studies have questioned the method of mutant generation of ΔluxS on

CjNC110 expression (29), a second northern blot comparing wild-type IA3902 to a previously

constructed ΔluxS mutant utilized by our lab (8) was performed and confirmed that CjNC110

expression in the IA3902ΔluxS mutant is also equivalent to wild-type levels (**Fig. S1**).

Homologous recombination was utilized to transfer the CjNC110 mutation into this ΔluxS

mutant to create the double-knockout mutant, ΔCjNC110ΔluxS, with the goal of comparing the

transcriptomic and phenotypic differences between mutation of ΔluxS or ΔCjNC110 alone

versus in combination (ΔCjNC110ΔluxS).

Differential gene expression analysis of RNAseq data suggests several potential

regulatory targets for CjNC110. To begin to identify potential regulatory targets for CjNC110,

RNAseq was utilized to compare gene expression changes between ΔCjNC110, ΔluxS and

ΔCjNC110ΔluxS mutants. Total RNA was isolated from an *in vitro* growth curve performed in

triplicate and colony counts (CFU/mL) over time were used to select samples representative of

exponential (3 hours) and early stationary (12 hours) phases of growth (**Fig. S2A**). ANOVA did

identify a statistically significant difference in growth between strains (P <0.05), however,

multiple comparison analysis of individual time points and strains when compared to wild-type

growth revealed that the only significant difference was a decrease in the A_{600} of ΔCjNC110 at

30 hours (**Fig S2B)** which represents the point where a decline in the A_{600} was noted to occur in

all strains tested. Further examination of the actual CFU/mL over the course of the growth curve

revealed that by 30 hours cell death was beginning to occur and colony counts were decreasing.

As further growth is assumed to be ceased at this time, there is likely minimal effect of this

difference on samples collected during growth prior to 24 hours, however, there is possibly

biological significance to the difference noted during the late stationary phase which may

warrant further investigation. It should be also be noted that the use of CFU/mL more clearly

allowed for delineation of actual growth stage of each strain as the A_{600} results were noted to

significantly lag the results of actual colony counts; therefore, the CFU/mL data was utilized to

determine selection of timepoints for RNAseq.

Sequencing of the total RNA libraries prepared from the growth curves was performed on

an Illumina HiSeq 2500 in high-output single read mode with 100 cycles, and Rockhopper was

utilized to analyze the resulting data for differential gene expression between the mutants (30).

Overall, 24 barcoded libraries were sequenced yielding over 109 million reads, with close to 100

million high quality reads aligning to either the genome or pVir plasmid of *C. jejuni* IA3902 and

averaging 4,553,847 reads per library (**Tables S1 and S2**). The vast majority of reads (average of

83% of total reads) mapped to protein coding genes of the chromosome, with only 7% of reads

mapping to ribosomal RNA on average following rRNA depletion with Ribo-Zero (median of

3%). Over half of the libraries contained less than or equal to 3% ribosomal RNA reads, which is

consistent with the manufacturer's predicted rRNA removal efficiency. One third of the libraries

did not exhibit efficient rRNA removal (>10% rRNA reads); the reason for this is unclear.

Despite this difference in efficiency, all samples provided a more than adequate number of non-

rRNA reads and thus were successfully utilized in the analysis. Less than 1% of reads mapped to

antisense regions of the annotated genome. Rockhopper also identified a number of putative

ncRNAs in the dataset; **Table S3** includes the curated list of identified ncRNA candidates.

Following computational analysis via Rockhopper, a change in gene expression was

deemed significant when the Q-value (false discovery rate) was <0.05 and a >1.5 fold change in

expression was observed. A summary of the differences in numbers of genes with significantly

increased and decreased expression in the various strains and timepoints is given in **Table 2**,

with a listing of specific genes given in **Table 3**. In the ΔCjNC110 mutant strain, six genes were

found to be downregulated and four genes were found to be upregulated when compared to the

IA3902 wild-type strain during exponential growth (complete information **Table S4)**. In

addition, a previously described ncRNA, CjNC140 (21) was found to be upregulated in the

mutant condition. During early stationary phase, 16 genes were found to be downregulated and

seven genes were found to be upregulated in the mutant strain when compared to the wild-type

strain. Of the differentially expressed genes, three genes [*neuB2, hisF, ptmA*] were

downregulated in the ΔCjNC110 mutant during both the exponential and stationary phases, and

one, CJSA_1261 was found to be upregulated in both conditions. In addition, five separate

operons predicted by Rockhopper demonstrated multiple genes within the operon affected by the

mutant condition. Analysis of functionality via the COG database revealed that multiple

upregulated genes (*luxS*, *cetA*, *cetB*) were present in the "Signal transduction mechanisms"

category; multiple downregulated genes were included in the "Cell wall/membrane biogenesis"

(*neuB2, ptmB,* CJSA_1352) and "Post-translational modification, protein turnover, chaperones"

(*tpx,* CJSA_0687) functional categories. These differentially expressed genes represent potential

targets of CjNC110 regulation in IA3902, and thus were used to inform further phenotypic study

of the function of this ncRNA. Genes highlighted in bold in **Table 3** indicate potential mRNA

targets of CjNC110 that were further investigated using phenotypic assays.

ΔluxS displays minimal alterations in gene expression via RNAseq, with more

significant gene expression changes observed in ΔCjNC110ΔluxS. Differential gene

expression analysis via RNAseq and Rockhopper was also performed for ΔluxS and

ΔCjNC110ΔluxS and compared to wild-type IA3902 to assess the difference between single

mutation of *luxS* and concurrent mutation of both CjNC110 and *luxS*. In the ΔluxS mutant, a

surprisingly small number of genes were noted to be differentially expressed. One gene was

found to be downregulated and 14 genes were found to be upregulated when compared to

IA3902 wild-type during exponential growth (summary **Table 3**, complete information **Table**

S5). Similar to the ΔCjNC110 mutant, the previously described ncRNA CjNC140 was also found

to be upregulated in the mutant condition during exponential growth only. During early

stationary phase, six genes were found to be downregulated and six upregulated in the ΔluxS

mutant strain when compared to wild-type. Of the differentially expressed genes, only one gene,

CJSA_1350, a putative methyltransferase, was found to be downregulated in the ΔluxS mutant

during both exponential and stationary phases. Three separate operons predicted by Rockhopper

demonstrated multiple genes within the operon affected by the mutant condition (two operons

upregulated, one operon downregulated). Analysis of functionality via the COG database

revealed that multiple up- (*trpF, trpB*) and down- (CJSA_0620, *leuC*) regulated genes were

present in the "Amino acid transport and metabolism" functional category; multiple upregulated

genes were also included in the "Cell motility" (*flgE, flgG2, flgG*) functional category. Only two

genes were differentially expressed in both the ΔCjNC110 and ΔluxS mutants, CJSA_0008

(upregulated during exponential phase) and CJSA_1107 (upregulated during stationary phase);

both genes are annotated as hypothetical proteins at this time. Despite the critical role that LuxS

plays in the AMC pathway, significant effects on the expression of other genes within the AMC

pathway were not observed.

In the ΔCjNC110ΔluxS double knockout mutant, however, a large increase in both

downregulated and upregulated genes when compared to wild-type was observed at both

timepoints (summary **Table 3**, complete information **Table S6**). Sixty-one protein coding genes,

seventeen tRNA genes, three known ncRNA genes (tmRNA, SRP, 6S) and seven newly

predicted ncRNAs were downregulated during exponential phase compared to IA3902 wild-type.

In addition, 47 protein coding genes were upregulated in exponential phase, along with the

previously described ncRNA CjNC140, which was also observed to be upregulated in the

ΔCjNC110 and ΔluxS single knockout mutants. During early stationary phase, 32 genes and two

newly predicted ncRNAs were found to be downregulated, while 29 genes were upregulated in

the mutant strain when compared to wild-type. Of the observed genes with differential

expression, many were observed to show altered expression during both exponential and

stationary phase when compared to wild-type; however, most were differentially expressed in the

opposite direction between the two timepoints. Only two genes (*motB,* CJSA_1350) were

downregulated in both conditions, while no genes were observed to be upregulated at both

timepoints. Many of the differentially upregulated genes demonstrated in both background

mutants, ΔCjNC110 and ΔluxS, were observed in the double knockout (**Figs. 2A** and **2B**), while

fewer of the downregulated genes were shared (**Figs. 2C** and **2D**).

Of the differentially expressed genes affected by the ΔCjNC110ΔluxS double knockout

mutation, 20 separate operons predicted by Rockhopper demonstrated multiple genes within the

operon affected by the mutant condition (eight operons upregulated, four operons

downregulated, and eight operons that showed opposing regulation at the different timepoints).

Figure S3 shows the number of genes in each COG category that were differentially expressed

in the double mutant condition during either exponential growth. The categories most affected by

these mutations were "Energy production and conversion," "Amino acid transport and

metabolism," "Translation," "Cell wall/membrane biogenesis," "Cell motility," and "Signal

transduction mechanisms." Using KEGG Pathways (31), a significant effect was observed in the

flagellar assembly pathway in the double knockout mutant as compared to either mutant alone

(**Figs. S4 and S5**). Of particular interest, this affect was also observed to be altered based on

growth phase, with decreased expression of σ^{54}-associated genes noted during exponential phase,

and increased expression noted during early stationary phase.

Phenotypic evaluation of increased *luxS* expression unexpectedly reveals alteration

of AI-2 transport in ΔCjNC110. Utilizing the potential targets identified in the RNAseq data

for ΔCjNC110, phenotypic testing was next performed to begin to determine the physiologic role

of CjNC110 in IA3902. Further investigation of a potential interaction with *luxS,* which was

identified as demonstrating significantly increased expression in ΔCjNC110 during early

stationary phase, was initially pursued due to the proximity of the two genes. LuxS activity is

traditionally measured using the *Vibrio harveyi* bioluminescence activity assay as an

approximation of AI-2 production (32). Using this assay as previously described, AI-2 levels

were initially evaluated within the extracellular environment by collecting cell-free supernatant

(E-CFS) at various time points during the growth of IA3902 wild-type and mutant strains. To

collect these samples, a second set of growth curves independent of those utilized for RNAseq

was performed in triplicate to include the ΔCjNC110c (**Fig. 3**). Similar to the first growth curves

utilized for RNAseq, ANOVA again identified a statistically significant difference between the

growth of the strains (P <0.05), however, the only significant difference was a decrease in the

A_{600} of ΔCjNC110 after 27 hours, similar to the difference seen in the first set of growth curves.

This defect was corrected in the ΔCjNC110c complement in the second set of growth curves,

indicating that while the biological significance is unknown, this observation is repeatable and

related to the loss of CjNC110.

As expected based on previous studies (8, 33), the results of the *V. harveyi*

bioluminescence assay demonstrated that both the ΔluxS and ΔCjNC110ΔluxS mutant strains

displayed no bioluminescence activity at any point during growth, indicating a complete lack of

AI-2 production which is consistent with the absence of a functional LuxS protein (**Fig. 4A**). For

the ΔCjNC110 mutant, however, bioluminescence was determined to be statistically significantly

decreased when compared to wild-type at time points 6, 9 and 12 hours (P <0.05) of the growth

curves, occurring during mid to late exponential phase and early stationary phase. This is in

distinct contrast to the expected result based on RNAseq analysis which demonstrated increased

luxS mRNA expression. Complementation of ΔCjNC110 completely restored extracellular AI-2

to wild-type levels, strongly suggesting that the observed defect in extracellular AI-2 was a true

phenotype warranting further exploration.

As the RNAseq results (i.e. – increased *luxS* gene expression) did not match the results of

the AI-2 assay (i.e.- decreased extracellular AI-2), qRT-PCR was performed on samples

collected at 12 hours from the same growth curve as the AI-2 assay using targets specific for the

3' region of the *luxS* gene. **Fig. 5** demonstrates the expression levels of *luxS* in the ΔCjNC110

mutant compared to wild-type and confirms that increased *luxS* mRNA levels are indeed present

during early stationary phase in the mutant, resulting in a fold change increase of 3.20 when

compared to IA3902 wild-type ($P <0.05$). Additionally, ΔCjNC110c was tested to determine if

complementation of the mutation returned *luxS* to wild-type levels; interestingly, ΔCjNC110c

also demonstrated a fold change increase of 3.77 when compared to IA3902 wild-type ($P <0.05$).

Proteomic analysis using liquid chromatography with tandem mass spectrometry (LC-

MS/MS) was then conducted to determine if the relative abundance of the LuxS protein was also

increased in ΔCjNC110. Whole protein extract LC-MS/MS analysis resulted in a total of 262

identified proteins that were discovered in each biological group. Two-way ANOVA with FDR

correction was used to identify statistically significant differences in protein abundance; these

statistically significant proteins are listed in **Table 4**. In both ΔCjNC110 and ΔCjNC110c, the

abundance of LuxS was noted to be statistically significantly increased ($P <0.0001$). Hierarchical

heat maps were also constructed to further investigate protein expression patterns when

comparing IA3902 wild-type to ΔCjNC110 and ΔCjNC110c. The top 20 hierarchical clustered

proteins are illustrated for both ΔCjNC110 and ΔCjNC110c compared to wild-type IA3902 in

Fig S6; LuxS was also identified via this approach as being differentially expressed in both

ΔCjNC110 and ΔCjNC110c. Therefore, the proteomics data confirmed the RNAseq and qRT-

PCR observations of significantly increased expression of LuxS when comparing wild-type to

both ΔCjNC110 and ΔCjNC110c.

As decreased LuxS abundance was not the cause of the observed decrease in extracellular

AI-2, several additional hypotheses were proposed, including: 1) decreased activity of the LuxS

protein leading to decreased AI-2 production, 2) increased degradation of AI-2, or 3) altered AI-

2 transport into or out of the extracellular environment. To assess for altered AI-2 transport, the

traditional *V. harveyi* bioluminescence activity assay was modified as described in the Materials

and Methods to assess for intracellular AI-2 levels using intracellular cell free supernatant (I-

CFS) obtained from the cell pellet of the same growth curve sample as the E-CFS. Results of

this assay are shown in **Fig. 4B** and clearly demonstrate that intracellular AI-2 levels increased to

statistically significant (P <0.05) levels in the ΔCjNC110 mutant when compared to IA3902

wild-type at both timepoints tested. Again, complementation of ΔCjNC110 corrected the

phenotypic alteration of increased intracellular AI-2 presence to wild-type. When taken together,

these results indicate that the observed decrease in extracellular AI-2 activity is not due to

impaired activity of LuxS, but instead strongly suggests that transport of AI-2 across the cell

membrane is altered in the ΔCjNC110 mutant leading to decreased extracellular and increased

intracellular AI-2 accumulation.

Mutation of CjNC110 significantly increases motility and hydrogen peroxide

sensitivity while decreasing autoagglutination in IA3902, opposite of the effects seen with

mutation of *luxS*. To further investigate the role of CjNC110 in IA3902, several additional

phenotypes suggested by the RNAseq results were assessed, including motility, hydrogen

peroxide sensitivity, and autoagglutination. Loss of c*etA* and *cetB* has been previously shown to

affect energy taxis and motility (34), therefore, the motility of the mutant strains in semi-solid

agar was initially compared to wild-type IA3902 at 30 hours post-inoculation. The results of this

assay confirmed that all isolates were highly motile, and statistical analysis via one-way

ANOVA indicated that there was a significant difference between strains (P <0.0001) (**Fig. 6**).

Further post-hoc analysis via Tukey's multiple comparisons test did not reach statistical

significance (P >0.05) when comparing ΔCjNC110 to wild-type despite motility for the

ΔCjNC110 strain observed to be consistently increased above the wild-type phenotype in all

biological replicates performed. Results did reach statistical significance (P <0.05), however,

when comparing wild-type to both the ΔCjNC110ΔluxS and ΔluxS mutants which displayed

statistically significant decreased motility; this result is consistent with previous reports of

decreased motility of *luxS* mutants in the IA3902 background (8). As many of the strains had

migrated to the edge of the plate by 30 hours, a second set of experiments was performed with

measurements taken at 24 hours post-inoculation. When measured at 24 hours, motility for the

ΔCjNC110 strain did demonstrate a statistically significant increase (P <0.05) when compared to

all other strains tested, and motility of both the ΔCjNC110ΔluxS and ΔluxS mutants was again

noted to be significantly decreased (**Fig. 6**). Following complementation of CjNC110, no

significant difference was observed when compared to IA3902 wild-type in either experiment,

confirming that complementation of CjNC110 quantitatively restored the phenotypic change

back to wild-type levels. These results indicate an increased motility of ΔCjNC110, which

differs from ΔluxS which displays decreased motility in IA3902.

Based on the potential regulatory interaction identified in our RNAseq data of CjNC110

with *tpx,* which has been previously demonstrated to encode a dedicated hydrogen peroxide

reductase in *Campylobacter* sp., a hydrogen peroxide (H_2O_2) disk inhibition assay was performed

as previously described (35). **Fig. 7** demonstrates the results of this assay which indicates that the

lack of CjNC110 significantly impacts H_2O_2 sensitivity when compared to wild-type as increased

sensitivity was observed in both CjNC110 mutant backgrounds (ΔCjNC110 and

ΔCjNC110ΔluxS) (P <0.05). Complementation of CjNC110 led to a significantly decreased

sensitivity to H_2O_2 compared to wild-type, indicating that the potential overexpression of

CjNC110 in ΔCjNC110c may lead to overcorrection of the phenotypic change (P <0.05). There

was no significant difference (P >0.05) noted between wild-type and ΔluxS, which is consistent

with previous studies of *luxS* mutation in other strains of *C. jejuni* such as NCTC 11168 (36).

These results indicate that CjNC110 may play a key role in sensing and responding to oxidative

stress in the environment for IA3902, and again represent a differing phenotype between

ΔCjNC110 and ΔluxS.

Differential expression of several genes associated with flagellar modification [*pseA,*

ptmA, ptmB, neuB2, and CJSA_1261] was also observed in ΔCjNC110; these genes have been

previously shown to affect the autoagglutination ability of *C. jejuni* (37-39). Therefore,

autoagglutination was also assessed as previously described (40) to determine if the observed

gene expression changes in ΔCjNC110 led to a phenotypically observable alteration.

Autoagglutination activity was measured at both 25ºC and 37ºC following 24 hours of incubation

with an increase in optical density correlating to decreased autoagglutination ability, and a

decrease in optical density correlating to increased autoagglutination ability. A statistically

significant difference ($p <0.0001$) between strains at both 25ºC and 37ºC was noted based on

initial analysis via one-way ANOVA (**Fig. 8**). When compared to IA3902 wild-type, ΔCjNC110

exhibited statistically significant decreased autoagglutination activity ($P <0.05$) at both

temperatures, while autoagglutination activity for ΔluxS and ΔCjNC110ΔluxS was noted to be

increased at a statistically significant level ($P <0.05$), again at both temperatures.

Complementation of CjNC110 restored the autoagglutination ability of ΔCjNC110 to wild-type

levels. These results indicate that *luxS* and CjNC110 function to influence autoagglutination

activity in opposing directions; interestingly, when mutation of both genes was combined, the

results favored the ΔluxS mutation but did return closer to wild-type levels.

Sustained colonization of chickens is impaired in ΔCjNC110 and absent in mutants

ΔCjNC110ΔluxS and ΔluxS, but is restored to wild-type colonization levels in ΔCjNC110c.

The ability to colonize the host is the critical first step in establishment of infection by

Campylobacter sp. Previous studies have demonstrated a loss of colonization ability of ΔluxS in

IA3902 (8), therefore, chicken colonization was utilized to determine if a similar phenotype was

present in either ΔCjNC110 or ΔCjNC110ΔluxS. Three-day old chicks were orally inoculated

with approximately 200 µl of 10^7 CFU per strain (wild-type, ΔCjNC110, ΔluxS,

ΔCjNC110ΔluxS and ΔCjNC110c). Once weekly for 3 weeks following inoculation, 6 chicks

were humanely euthanized and cecel contents collected for analysis of colonization levels via

plating of serial dilutions and colony counts. For ΔCjNC110, while colonization was still present

at all timepoints, a statistically significant decrease in colonization was noted compared to wild-

type on days post-inoculation (DPI) 5, 12, and 19 (P <0.05); complementation of the ΔCjNC110

mutant restored colonization to wild-type levels at all timepoints analyzed (**Fig. 9**). Similar to

previous studies, while some colonization, albeit significantly decreased, was noted for ΔluxS at

DPI 5, no colonization was observed by DPI 12 and 19 (P <0.05). For the ΔCjNC110ΔluxS

mutant, normal colonization levels compared to wild-type were observed on DPI 5, however, by

DPI 12 and 19, no colonization of ΔCjNC110ΔluxS was detectable (P <0.05). These results

indicate that the presence of CjNC110 is necessary for optimal colonization, while *luxS* is

required for establishing sustained colonization of IA3902 in chickens.

DISCUSSION

In other species of bacteria that harbor known RNA chaperones such as Hfq, methods of

identification of cognate mRNA binding partners of ncRNAs are available via several

straightforward approaches [recently reviewed in (41)]. As *Campylobacter* sp. lack all known

RNA chaperones, identification of targets via these methods is not feasible in these pathogens;

previous attempts at computational predictions have also failed (18). However, recent work in

Campylobacter has demonstrated that RNAseq can be useful for analyzing transcriptomic

differences between wild-type and mutant strains following inactivation of protein-coding genes

(42, 43). Indeed, over the past several years, RNAseq has become the method of choice to

analyze the transcriptomes of bacteria under various conditions as it allows evaluation of the

entire transcriptome rather than only previously annotated regions as with older technology such

as microarrays (44-46). In theory, RNAseq should also be able to be utilized for the same

purpose of discovering the global effects of inactivation of ncRNAs. Small ncRNAs have been

shown to be able to stabilize mRNA transcripts by multiple mechanisms (47), which should lead

to increased levels of transcript availability and identification of significant differences using

RNAseq. Some interactions with ncRNAs have also been shown to increase transcript turnover

by targeting transcripts for RNase degradation or exposing RNase cleavage sites (47), which

should lead to decreased levels of transcripts available for identification via RNAseq. Thus,

assessment of differential gene expression via RNAseq should be useful for uncovering potential

interactions of ncRNAs with mRNA targets when that interaction directly leads to altered levels

of mRNA transcripts in the cell; target interactions that do not lead to direct changes in mRNA

transcript levels but instead affect the translation efficiency of the target mRNA or are the result

of direct protein-ncRNA interactions cannot be determined via this approach. While our

differential gene expression results do not guarantee a direct mRNA-ncRNA interaction or

provide a mechanism for that interaction, the results of the work presented here clearly indicate

that RNAseq can be a useful tool for investigating the role of ncRNAs in post-transcriptional

regulation of gene expression in bacteria, and can successfully be used to inform the design of

phenotypic studies to begin to elucidate the function of ncRNAs in pathogenic bacteria who lack

RNA chaperones.

One particular area of interest from the onset of our work was the potential for an

interaction between CjNC110 and *luxS.* Our results related to a potential interaction between

luxS and CjNC110 were surprising in that despite increased LuxS expression, extracellular AI-2

levels were decreased while intracellular AI-2 increased; this ultimately suggested that CjNC110

plays a key role in the transport of AI-2 between the intra- and extracellular environment of

IA3902. In *E. coli,* the YdgG (TqsA) protein, part of the AI-2 exporter superfamily, has been

described to export AI-2 molecules (48); additional *in vitro* studies assessing the membrane

permeability of AI-2 also demonstrate that it is hydrophilic with low affinity towards lipids,

which strongly suggests the need for a transport system (49). Previous BlastP analysis has shown

potential homologues of the YdgG protein exist within the *Campylobacterales* family, but none

directly within the *C. jejuni* species (50). Within our proteomics data, several periplasmic and

lipoproteins which could be located in the cytoplasmic membrane were identified to exhibit

decreased expression in ΔCjNC110 that was corrected to wild-type levels in ΔCjNC110c. The

function of many of these proteins has yet to be elucidated in *Campylobacter,* therefore, future

work assessing whether these proteins may be regulated by CjNC110 and could serve as an AI-2

exporter in *C. jejuni* is warranted. In other species of bacteria such as *E. coli*, uptake of AI-2

from the extracellular environment has also been shown to occur and is mediated by an active

LSR system (51). Comparative genomics has again failed to identify a homologous AI-2 uptake

system in *C. jejuni*, however, a number of ABC-type transporters whose function remains to be

elucidated do exist in *Campylobacter* sp. (28). Several studies evaluating the concentration of

AI-2 over time following addition of exogenous AI-2 to cell culture have attempted to determine

whether or not AI-2 is able to be internalized in *C. jejuni* with mixed results (52, 53). Thus,

further research into an AI-2 uptake system in *C. jejuni* and the role that CjNC110 may play in

regulation of that uptake may also be warranted.

Based on the AI-2 results, the increased expression of LuxS in ΔCjNC110 and failure of

complementation of *luxS* gene expression and LuxS protein abundance in ΔCjNC110c was

unexpected, particularly as all other phenotypes tested, including AI-2 detection intra- and

extracellularly, fully complemented to wild-type. The reason for the increased *luxS* expression

and LuxS protein abundance in ΔCjNC110 and ΔCjNC110c is currently unknown but one

reasonable hypothesis is that this increased expression may be related to additional unknown

effects of CjNC110 on the AMC leading to feedback regulation of *luxS* expression. While

regulation of the AMC in other Proteobacteria has been studied in detail, the regulatory pathways

of the AMC are currently unknown in *Campylobacter* as this genus lacks all of the known

transcription factors and riboswitches typically involved regulation (54). As CjNC110 is located

immediately downstream of *luxS,* en bloc removal of the CjNC110 coding sequence could also

have led to disruption of normal termination of the *luxS* transcript, thus altering the natural

mRNA stability; this effect would have been maintained in the complement as well. Further

investigation to determine the location of a terminator via the WebGesTer database failed to

identify a predicted terminator sequence for *luxS* (http://pallab.serc.iisc.ernet.in/gester/) (55).

Therefore, additional work is needed to determine which hypothesis might explain the increase

of LuxS in both ΔCjNC110 and ΔCjNC110c. Regardless of the cause, our data clearly

demonstrates that the increased *luxS* expression and LuxS protein abundance in ΔCjNC110 is

unrelated to the observed changes in extracellular and intracellular AI-2 concentrations which

fully complemented back to wild-type levels in ΔCjNC110c.

Several other important phenotypes were also affected by mutation of CjNC110 based on

the RNAseq data, including motility, autoagglutination, hydrogen peroxide sensitivity, and

chicken colonization. Motility is considered critical for the *in vivo* virulence of *C. jejuni* and

requires a functional flagellar apparatus (56, 57); however, this alone is not sufficient for normal

motility to be present (58). In our RNAseq data, the *cetAB* operon (*Campylobacter* energy taxis

proteins A and B), which is known to mediate the energy taxis response in *Campylobacter* (34)*,*

was statistically significantly upregulated in the ΔCjNC110 mutant when compared to wild-type

during the exponential phase of growth; expression of *cetAB* was increased during stationary

phase as well, however, did not reach the level of statistical significance.   Defects in both *cetA*

and *cetB* have been shown to lead to altered motility phenotypes, particularly in response to

migration towards critical factors in *Campylobacter* metabolism such as sodium pyruvate and

fumarate (58). Thus, the increase in *cetAB* correlates well to the statistically significant increase

in motility found between the ΔCjNC110 mutant and wild-type when motility was assessed at 24

hours. Investigation of expression of *cetA* and *cetB* has shown that levels of the gene products

are unaffected by mutation of sigma factors σ^{54} or σ^{28}, indicating that transcription of the *cetAB*

operon is likely controlled by σ^{70} or another yet unknown transcription factor (34). Based on this

information, it seems plausible that CjNC110 might normally act as a repressor of the CetAB

energy taxis system. Our proteomics data failed to detect the CetA and CetB proteins, therefore,

we could not validate that increased transcriptional expression of *cetA* and *cetB* resulted in

increased protein expression levels. Many of the proteins identified in our study exhibited

extremely high expression levels, which, without fractioning, prevented proteins of lesser

abundance from being detected. Whole cell proteomic analysis with fractioning and time-course

analysis would be required to reveal the entire proteomic network regulated by CjNC110 and

validate expression patterns of proteins of lesser abundance than the 262 proteins that were

identified in our study. Further work to determine if a direct interaction between CjNC110 and

the *cetAB* operon exists and leads to increased expression of the CetA and CetB proteins is

warranted.

No statistically significant differences in the expression of genes associated with the

flagellar apparatus were noted in the transcriptome of ΔCjNC110; this may help to explain why

an increase in energy taxis would allow for an increase in observed motility with the assumption

that normally functioning flagella are present. The presence of normal flagella has also been

associated with autoagglutination ability and is considered necessary but not sufficient for this

important trait (58). Previous studies in *C. jejuni* have demonstrated that in addition to the

presence of normal flagella, interactions between modifications on adjacent flagellar filaments,

particularly those provided by O-linked protein glycosylation, are required for normal

autoagglutination ability (59). In the present study, the ΔCjNC110 mutant exhibited decreased

autoagglutination when compared to wild-type at both 25ºC and 37ºC. Several genes that have

been associated with flagellar glycosylation in other strains of *C. jejuni* were downregulated in

our RNAseq data in the ΔCjNC110 mutant when compared to wild-type during both exponential

and stationary growth phases. The O-linked protein glycosylation system which is responsible

for flagellar glycosylation has been extensively studied in only one strain of *C. jejuni* (81-176)

and has not been studied to date in IA3902 (38); this system is one of the most diverse genomic

loci in *Campylobacter* spp. and also represents one of the key areas of variation between IA3902

and NCTC 11168 (5).   Further genetic analysis of the flagellar modification region of the

IA3902 genome reveals the presence of genes associated with both of the previously identified

pseudaminic acid and legionaminic acid flagellar glycosylation pathways known to exist in

Campylobacter sp.; this analysis and the subsequent gene expression data comparison of

ΔCjNC110 to wild-type is presented in **Table S7**. The pseudaminic acid pathways (encoded by

the *pseA-I* genes which are fully annotated in IA3902) lead to flagellar glycosylation with either

pseudaminic acid (PseAc) (Pse5Ac7Ac), which is the major glycan found on *Campylobacter*

flagella, or derivatives such as PseAm (Pse5Am7Ac) which has an acetamidino substitution (60).

In our RNAseq dataset, *pseA* was significantly decreased in expression during exponential

growth, and also decreased without reaching significance during stationary phase. Mutation of

pseA has previously been shown to lead to flagellar modification with only PseAc, and not

PseAm (38, 60); *pseA* has also been shown to be required for optimal adherence and invasion of

human epithelial cells and virulence in a ferret model (59).

The legionaminic acid (LegAm) flagellar glycosylation pathway, encoded by the *ptmA-H*

genes, has been less extensively studied in *C. jejuni*, in part because it is not present in 81-176; in

fact, much of the work that has been done in *Campylobacter* spp. related to this pathway has

been in *C. coli* (37, 39). NCTC 11168 does contain homologs to all of the *ptm* genes (Cj1324-

Cj1332) albeit with different gene nomenclature (*leg* or *neu*) often used to describe the genes,

and mutation of these genes in NCTC 11168 has been shown to decrease autoagglutination

ability as well as reduce but not eliminate the ability to colonize chickens (61).   While not

currently annotated as such, our further analysis has identified that in IA3902, gene locus

CJSA_1261-1268 represent homologs to all of the *ptm* genes present in NCTC 11168 (**Table**

S7). Based on this data, it is reasonable to suggest that IA3902 also harbors an active LegAm

flagellar glycosylation pathway. Several genes related to legionaminic acid flagellar

glycosylation were identified in our RNAseq dataset as being differentially expressed in the

CjNC110 mutant during both stages of growth, which may help explain altered autoagglutination

levels of ΔCjNC110. In particular, *ptmA* and *neuB2* (homologous to *ptmC)* were both observed

to be statistically significantly decreased at both exponential and stationary phase, while *ptmB*

was statistically significantly decreased at stationary phase and decreased but not reaching

statistical significance during exponential phase. Mutation of the *ptmA-G* genes in *C. coli* results

in the loss of all LegAm derivatives and a conversion to glycosylation with PseAc only; these

mutants still retain a normal flagellar apparatus and motility (37, 39, 62). In our RNAseq data,

CJSA_1261 (homologous to *ptmG*) was the sole flagellar modification gene that demonstrated a

statistically significant increase in expression during both growth phases; *ptmG* has been shown

to be involved in conversion of LegAm to various derivatives (Leg5AM7Ac and

Leg5AmNMe7Ac) (37). Alterations in *ptmG* levels in the CjNC110 mutant are of particular

interest as it has been recently shown to be also regulated by another small non-coding RNA pair

in *C. jejuni*, CjNC180/CjNC190 in NCTC 11168 (27). While CjNC180/CjNC190 were not

detected in this dataset by Rockhopper, these ncRNAs are present in IA3902 and CjNC180 has

previously been demonstrated to be significantly upregulated in the *in vivo* host gallbladder

environment in IA3902 (24). When taken together, this data suggests that CjNC110 may play a

key role in modulation of flagellar glycosylation in IA3902, thus directly affecting

autoagglutination ability. Other potential causes for decreased autoagglutination ability include

alteration of the flagellar filament. While not identified via any other approach, the FlaA protein

was demonstrated to be decreased via hierarchal heat mapping in our proteomics data in the

CjNC110 mutant with a return to wild-type levels in the complement (**Fig. S6**); if this change

corresponds to an alteration of the flagellar structure, it could in theory also explain the observed

decrease in autoagglutination ability. Thus, further work including determination of the normal

flagellar glycosylation of IA3902 and assessment for changes in both the flagellar filament

structure and glycosylation of the ΔCjNC110 mutant when compared to the IA3902 wild-type is

warranted.

The RNAseq data also demonstrated that in the ΔCjNC110 background *tpx* was

significantly downregulated during the exponential phase of growth, prompting phenotypic

assessment of hydrogen peroxide sensitivity. Thiol peroxidase (Tpx) has been previously shown

to function as a hydrogen peroxide reductase in NCTC 11168, however, in that strain, Tpx-

defective mutants were not shown to be sufficient to cause alteration of hydrogen peroxide

sensitivity via disc diffusion and required a second mutation of the Bcp gene to observe

increased sensitivity (35). IA3902 also harbors a Bcp homolog that is 96% homologous to

NCTC 11168; this gene did not demonstrate altered expression via RNAseq analysis, and

functional evaluation of this gene in IA3902 has not been performed to date. While our results

clearly indicate that the loss of CjNC110 increases hydrogen peroxide sensitivity, and that the

expression of *tpx* in the mutant decreased, it is not possible to definitively state that the observed

phenotype is due solely to an interaction of CjNC110 with *tpx* alone. Further analysis of

potential alterations of CjNC110 with additional known oxidative stress genes in the RNAseq

data did indicate that *sodB,* which has been previously identified to contribute to hydrogen

peroxide sensitivity in *C. jejuni* (63), demonstrated a Q-value that reached the threshold for

significance ($Q = 0.004$), however, the fold change just missed the cutoff for significance at -1.4.

Thus, further investigation for the potential interaction of CjNC110 with both *tpx* and additional

genes that control the oxidative stress response in *C. jejuni* such as *sodB* is also warranted.

When taken together, the results of the *in vitro* phenotypic studies, which identified

several key phenotypes affected by the loss of CjNC110, strongly suggested that mutation of

CjNC110 may interfere with normal colonization of IA3902. Indeed, our results show that

mutation of ΔCjNC110 led to a significantly decreased although still present colonization ability

in IA3902 compared to wild-type, indicating that this ncRNA has the potential to play a key role

in modulating colonization by *C. jejuni*. The exact mechanism for the decreased colonization

ability remains unknown, but could be due to any combination of factors identified in the

phenotypic screening, including increased sensitivity to ROS, decreased autoagglutination

ability, or altered quorum sensing ability; further work to elucidate how these factors may

interact to affect *in vivo* colonization is thus warranted. The loss of the ability of ΔluxS to

sustain colonization in chickens, as demonstrated in this study, has previously been described in

IA3902 but the same phenotype was not observed in the closely related NCTC 11168 (8); the

reason for this difference is currently unknown, but has been speculated to be related to strain-

specific differences in the metabolism of methionine or *S*-adenosylmethionine (SAM) recycling

which are also affected by the *luxS* mutation (28). Additional work in our lab has shown that the

luxS mutation in IA3902 does disrupt the AMC, which functions to recycle SAM and produce

methionine (64), however, DNA methylation in IA3902 was not affected by the *luxS* mutation

(65). Comparative genomics reveals that NCTC 11168 does harbor an additional genomic

system for methionine biosynthesis in the form of *metAB* which is not present in IA3902 and

may represent an alternative pathway for synthesis of this required amino acid and an

explanation for the observed difference in colonization ability. Further work is ongoing in our

lab to determine if differing methionine metabolism may explain the difference in colonization

ability of *luxS* mutants between strains, as well as to assess if mutation of CjNC110 in NCTC

11168 also leads to a defect in colonization, or, if similar to *luxS,* the phenotype differs from

IA3902.

After attempting to compare various strains and methods of mutation of the *luxS* gene and

finding wide variation in phenotypical and transcriptional changes, Adler *et al.* (29) suggested

that some of the differences reported between *luxS* mutants of *C. jejuni* in various studies may be

due to unknown polar affects caused by alteration of expression of CjNC110 based on the

mutation strategy used for the *luxS* mutation. Our northern blot analysis confirmed that CjNC110

was still expressed at the same level as the wild-type strain in our *luxS* mutant construct. In

addition, our phenotypic data clearly demonstrates very different phenotypes between

ΔCjNC110 and ΔluxS in IA3902 for all phenotypes tested, with variation in the phenotype of

ΔCjNC110ΔluxS based on the dominant phenotype of the two mutants. Thus, based on the work

presented here, for our *luxS* mutant constructs, we can definitively state that polar effects on

CjNC110 are not present.

Initial analysis of transcriptome changes in the ΔluxS mutant in our study revealed very

few genes identified as differentially expressed, and the list of genes identified did not overlap

with the list of differentially expressed genes generated primarily via microarray in various other

strains (66-68). One explanation for this discrepancy may be differences between strains of *C.*

jejuni or culture conditions when compared to these previous studies. However, in distinct

contrast to the lack of similar findings to previous gene expression studies exhibited by the

ΔluxS mutant, when the ΔCjNC110 and ΔluxS mutations were combined into ΔCjNC110ΔluxS,

many of the previously noted transcriptional changes attributed to mutation of the *luxS* gene

alone in a different strain of *C. jejuni*, 81-176, which also encodes and expresses CjNC110,

became apparent (67). Of the 57 genes listed as differentially expressed by the *luxS* mutant under

various conditions in He *et al.*(67), 23 were also found to be differentially expressed in our

RNAseq data (any mutant construct). Of the 23 overlapping genes, no genes were found only in

the ΔluxS mutant, and only the flagellar genes *flgE, flgG2,* and *flgG* and a putative helix-turn-

helix protein CJSA_1449 were differentially expressed in both the ΔluxS and ΔCjNC110ΔluxS

mutants. In contrast, three genes (*tpx, ptmB, ptmA*) were differentially expressed only in the

ΔCjNC110 mutant. The He *et al.* study (67) also reported increased sensitivity to H_2O_2 in their

luxS mutant construct, which more closely aligns with our ΔCjNC110 and ΔCjNC110ΔluxS

mutant data and not ΔluxS. These findings strongly suggest that as hypothesized by Adler *et al.*

(29), in some previous studies of ΔluxS, the main driver for the transcriptional and phenotypic

changes seen was likely an inadvertent inactivation of both CjNC110 and *luxS,* and not *luxS*

alone. Thus, further work is needed to re-evaluate the expression of CjNC110 in various *luxS*

mutant constructs to determine if alteration of expression of CjNC110 is present and may

account for some of the differences seen in phenotypes associated with *luxS* mutation in *C.*

jejuni.

Of the genes identified in both our study and He *et al.* (67), a large number of the

flagellar assembly hook-basal body associated proteins (FlgD, FlgE, FlgG, FlgG2, FlgH, FlgI,

and FlgK) that are under the control of σ^{54} (RpoN) promotors were identified as differentially

expressed (69, 70). Expression from σ^{54} promotors has been shown to require activation of the

FlgRS two-component regulatory system (69, 71). Many of these genes showed opposite

expression based on growth phase (decreased exponential, increased stationary), and it has been

previously demonstrated that regulation by these sigma-factors is growth cycle dependent, with

σ^{54}-regulated genes typically expressed between σ^{70}- and σ^{28}-associated genes (69). It has been

suggested that an additional unknown factor may control the temporal regulation of σ^{54}-

dependent flagellar genes (72), and the intergenic region between *luxS* and CJSA_1137 where

CjNC110 is located has previously been identified to demonstrate differential expression in an

rpoN (σ^{54}) mutant in NCTC 11168 (19). Therefore, it is reasonable to consider based on

previous studies and our results that the genomic region encompassing *luxS* and CjNC110 may

play a role in the growth-phase dependent regulation of flagellar assembly.

666   While not explored further in our phenotypic studies, Rockhopper did identify two

additional ncRNAs, CjNC140 and CjNC130/6S, as being differential expressed in our mutants;

these ncRNAs have been previously identified and confirmed to be transcribed in multiple other

C. jejuni strains (21). Predicted to be transcribed in the intergenic region upstream of *porA*, the

CjNC140 ncRNA was upregulated during exponential growth in all three of the mutant strains

when compared to wild-type (**Fig. S7**). As increased expression was not observed during

stationary growth, this suggests that CjNC140 may serve as a regulator involved in mediating

changes during different stages of bacterial growth. The fact that its expression is similarly

altered under all 3 mutant conditions also suggests that the three genes, *luxS*, CjNC110 and

CjNC140, may normally interact within the cell in some way to facilitate these changes. Point

mutations in the nearby *porA* have also been recently determined to be sufficient to cause the

abortion phenotype (10), making this ncRNA particularly interesting for further study.

678   Finally, while the specific mechanisms by which ncRNA CjNC110 interacts with each

individual target were not elucidated in this study, the research presented herein clearly identified

several specific roles of CjNC110 in the pathophysiology of *C. jejuni* IA3902. Indeed, even for

well-studied ncRNAs such as MicA in *Enterobacteriacae,* new publications are still being

produced which describe new roles and mechanisms of action even after a decade of research

[recently reviewed in (73)]. Future work is necessary to investigate the hypotheses generated

herein for genes and proteins that may be regulated by CjNC110 and to determine the

mechanism by which CjNC110 interacts with each of the potential mRNA targets identified in

this study.

In summary, the results of our research clearly demonstrate a significant role for

CjNC110 in the pathobiology of *C. jejuni* IA3902. The collective results of the phenotypic and

transcriptomic changes observed in our data complement each other and suggest that CjNC110

may be involved in regulation of energy taxis, flagellar glycosylation, cellular communication

via quorum sensing, oxidative stress and chicken colonization in *C. jejuni* IA3902. This work

provides for the first time valuable insights into the potential regulatory targets of the CjNC110

small ncRNA in the zoonotic pathogen *C. jejuni* and suggests a wide range of research avenues

into the role of ncRNAs in the pathobiology of this important zoonotic pathogen.

MATERIALS AND METHODS

Bacterial strains, plasmids, primers and culture conditions. *C. jejuni* IA3902 was

initially isolated from an outbreak of sheep abortion in Iowa during 2006 and has been utilized

by our laboratory as the prototypical isolate of clone SA (3). W7 is a highly motile variant of the

commonly utilized laboratory strain *C. jejuni* NCTC 11168 (8). *C. jejuni* strains and their

isogenic mutants were routinely grown in Mueller-Hinton (MH) broth or agar plates (Becton-

Dickinson, Franklin Lakes, NJ) at 42°C under microaerophilic conditions (5% O_2, 10% CO_2,

85% N_2). For the mutant strains containing a chloramphenicol resistance cassette, 5 μg/mL

chloramphenicol was added to either the broth or agar plates when appropriate. For strains

containing a kanamycin resistance cassette, 30 μg/mL kanamycin was added to either the broth

or agar plates when appropriate.

For genetic manipulations, *Escherichia coli* competent cells were grown at 37°C on

Luria-Bertani (LB) agar plates or broth (Becton-Dickinson, Franklin Lakes, NJ) with shaking at

125 rpm. When appropriate, 50 μg/mL kanamycin, 20 μg/mL chloramphenicol or 100 μg/mL

ampicillin was added to the broth or agar plates for selection of colonies. *Vibrio harveyi* strains

were grown in autoinducer broth (AB) at 30°C with shaking at 175 rpm as described previously

(74).

All strains used in this study are described in **Table 1**, with all relevant plasmids listed in

Table S8 and primer sequences and LNA DIG-labeled probe sequence listed in **Table S9**. **Fig.**

S8 illustrates the genetic modification strategy described below to create the mutants utilized in

this study. All strains were maintained in 20% glycerol stocks at -80°C and passaged from those

stocks as needed for experimental procedures.

Creation of *C. jejuni* ΔCjNC110 and ΔCjNC110ΔluxS mutants in IA3902. An

isogenic CjNC110 mutant of *C. jejuni* IA3902 was constructed via deletional mutagenesis

utilizing a combination of synthetic double-stranded DNA (dsDNA) fragments and traditional

cloning methods. Based on previously published data depicting the proposed transcriptional start

site for CjNC110 (21), the coding region of CjNC110 in IA3902 and the prototypical *C. jejuni*

strain NCTC 11168 were first confirmed to be identical. Then, a 200 bp section of the IA3902

genome starting 20 bp upstream and including the entire 137 nt transcript of the CjNC110

sequence as predicted in Dugar *et al.* (21) was replaced with 820 bp of the promoter and coding

sequence of the chloramphenicol acetyltransferase (*cat*) gene of *Campylobacter coli* plasmid C-

589 (75). Synthetic dsDNA including approximately 500 bp upstream and 500 bp downstream of

the region replaced with the *cat* cassette was then synthesized in 4 fragments of 500 bp each with

overlapping homologous ends (Integrated DNA Technologies, Coralville, IA). The Gibson

Assembly method was then utilized to assemble the synthetic dsDNA fragments using the

Gibson Assembly Master Mix (New England Biolabs, Ipswich, MA) (76). Following assembly,

primers (CjNC110F2 and CjNC110R2) were designed to amplify a 1785 bp product of the

assembled dsDNA; PCR amplification was achieved using TaKaRa Ex Taq DNA Polymerase

(ClonTech, Mountain View, CA). This amplified PCR product was then cloned into the pGEM-T

Easy Vector using T4 ligase (Promega, Madison, WI), resulting in the construction of

pCjNC110::cat which was then transformed into chemically competent *E. coli* DH5α (New

England Biolabs, Ipswich, MA). Transformants were then selected on LB agar plates containing

chloramphenicol (20 μg/ml), ampicillin (100 μg/ml), and ChromoMax IPTG/X-Gal Solution

(Fisher Scientific, Pittsburgh, PA). pCjNC110::cat was purified from the transformed *E. coli*

using the QIAprep Miniprep kit (QIAGEN, Germantown, MD) and confirmed by PCR to contain

the construct again using the CjNC110F2 and CjNC110R2 primers.

The pCjNC110::cat plasmid DNA was then introduced to *C. jejuni* W7, a highly motile

variant of NCTC 11168, as a suicide vector and the deletion transferred into the genome of *C.*

jejuni W7 via homologous recombination. Transformants were selected on MH agar plates

containing chloramphenicol (5 μg/ml) and deletional mutagenesis was again confirmed via PCR

analysis and Sanger sequencing to create *C. jejuni* W7ΔCjNC110. Following confirmation,

natural transformation was used to move the gene deletion into *C. jejuni* IA3902 as previously

described to create *C. jejuni* IA3902ΔCjNC110 (77). Natural transformation was again used to

move the CjNC110 gene deletion into the previously created *luxS* insertional mutant *C. jejuni*

IA3902ΔluxS (8) to create the double knockout mutant *C. jejuni* IA3902ΔCjNC110ΔluxS.

Transformants were selected on MH agar plates containing chloramphenicol (5 μg/mL) and

kanamycin (30 μg/mL) and confirmed via PCR analysis and Sanger sequencing of the CjNC110

region along with the entire upstream (*luxS*) and downstream (CJSA_1137) genes using primers

Cj1198F1 and Cj1199R3. All colonies were screened for presence/absence of motility as

described below, and only colonies with verified motility were used for future studies.

Expression of CjNC110 in wild-type and elimination of expression in the ΔCjNC110 mutant was

confirmed via northern blot analysis using an LNA custom designed probe and 15 μg of total

RNA as described below.

Creation of *C. jejuni* ΔCjNC110 complement in IA3902. Complementation of

ΔCjNC110 was achieved via insertion of the coding region into the intergenic region of the 16S

and 23S rRNA operon (*rrs-rrl*) of IA3902ΔCjNC110 via homologous recombination using

plasmid pRRK. A copy of the CjNC110 gene including 190bp upstream of the transcription

initiation location (total of 570bp) was cloned into an *XbaI* site of pRRK located upstream of the

kanamycin resistance determinant to create pRRK-CjNC110 as previously described in our lab

(78) with some modifications of the original protocol (79).   Briefly, the CjNC110 gene and

flanking regions of IA3902 were amplified using forward and reverse primers designed with an

XBaI restriction enzyme site (CjNC110cF1 and CjNC110cR1). Following PCR amplification

with TaKaRa Ex Taq DNA Polymerase (ClonTech, Mountain View, CA), the PCR product and

pRRK plasmid were digested with the *XBaI* restriction endonuclease (New England Biolabs,

Ipswich, MA). The digested plasmid was then incubated twice with Antarctic Phosphatase (New

England Biolabs, Ipswich, MA) to prevent self-ligation of cut plasmid. Following ligation and

transformation, transformants were selected on LB agar plates containing chloramphenicol (20

μg/ml) and kanamycin (50 μg/ml kanamycin) yielding a plasmid containing the construct

pRRK::CjNC110 in *E. coli* DH5α. Following PCR confirmation and Sanger sequencing to

confirm preservation of correct gene sequence, the plasmid was then utilized as a suicide vector

and the gene was inserted into the intergenic region of the 23S rRNA operon (*rrs-rrl*) of

IA3902ΔCjNC110 via homologous recombination to generate IA3902ΔCjNC110c

transformants. The transformants were selected on MH agar plates containing chloramphenicol

(5 μg/ml) and kanamycin (30 μg/ml) and confirmed via PCR analysis and Sanger sequencing to

contain both the expected ΔCjNC110 deletional mutation and ΔCjNC110c insertion, validating

successful genetic complementation. All positive colonies were screened for presence/absence

of motility as described below and a motile complement was selected to create

IA3902ΔCjNC110c. Expression of CjNC110 in ΔCjNC110c was confirmed via northern blot

analysis using an LNA custom designed probe and 15 μg of total RNA as described below.

Growth curves. Two separate growth curves were completed in triplicate utilizing the

same method as described below with the only variation being the addition of the complement

strain ΔCjNC110c to the second growth curve. To begin the growth curve, the A_{600} of overnight

cultures were adjusted to 0.5 using sterile MH broth on a Genesys 10S VIS spectrophotometer

(Thermo Scientific™, Waltham, MA). Cultures were then diluted 1:10 for a final targeted

starting A_{600} of 0.05 in 90 mL MH broth and placed in a sterile 250 mL Erlenmeyer glass flask.

Cultures were incubated at 42°C under microaerophilic conditions with shaking at 125 rpm for

30 hours with removal of samples from the flasks at designated time points [3, 6, 9, 12, 24, 27

(2^{nd} growth curve only) and 30 hours]. For the first growth curve (which included the following

strains: IA3902 wild-type, ΔCjNC110, ΔluxS and ΔCjNC110ΔluxS), samples were processed as

described below for RNA isolation as well as assessed for A_{600} and actual colony counts using

the drop-plate method as previously described (80). For the second growth curve (which

included ΔCjNC110c in addition to the four previous strains), samples were collected and

processed for A_{600} and as described below for qRT-PCR and assessment of AI-2 levels via the

bioluminescence assay. The A_{600} over time were statistically analyzed for the two independent

growth curves, using a two-way ANOVA with repeated measures and Dunnett's multiple

comparison test (GraphPad Prism).

Total RNA extraction. Broth culture samples processed for total RNA extraction were

either centrifuged at 8000 x *g* for 2 minutes (for sequencing and qRT-PCR) or 10,000 x *g* for 4

minutes (northern blot) at 4°C immediately following collection. Following pelleting of the cells,

the supernatant was decanted and 1 mL QIAzol Lysis Reagent (QIAGEN) was added to

resuspend the cell pellet and protect the RNA. QIAzol-protected cultures were then stored at -

80°C for up to two months prior to proceeding with total RNA isolation. Total RNA isolation

was performed using the miRNeasy Mini Kit (QIAGEN) followed by purification using the

RNeasy MinElute Cleanup kit (QIAGEN) as previously described (24). RNA quality was

measured using the Agilent 2100 Bioanalyzer RNA 6000 Nano kit (Agilent Technologies, Santa

Clara, CA) and all RNA samples utilized for downstream RNAseq library preparation had an

RNA integrity number (RIN) of >9.0, indicating high quality RNA. Verification of complete

removal of any contaminating DNA was performed via PCR amplification of a portion of the

CJSA_1356 gene, which is unique to *C. jejuni* IA3902, using primers SA1356F and SA1356R

(81).

RNAseq library preparation and sequencing. The 3 hour (exponential phase) and 12

hour (stationary phase) timepoints were selected for RNAseq analysis based on assessment of

$\log_{10}$ CFU/mL (**Fig. S2A**). 2.5 µg of confirmed DNA-free total RNA was treated with Ribo-Zero

rRNA Removal Kit for Bacteria according to the manufacturer's instructions (Illumina, San

Diego, CA). Following rRNA removal, the ribosomal depleted total RNA was purified using the

RNeasy MinElute Cleanup kit (QIAGEN) using the same modifications as described previously

(24). Following clean-up, the RNA quality, quantity, and rRNA removal efficiency was then

analyzed via the Agilent 2100 Bioanalyzer RNA 6000 Pico kit (Agilent Technologies). Library

preparation for sequencing on the Illumina HiSeq platform was completed using the TruSeq

stranded mRNA HT library preparation kit (Illumina) with some modifications as described

previously (24). The pooled library was then submitted to the Iowa State University DNA

Facility and sequenced on an Illumina HiSeq 2500 machine in high-output single read mode with

100 cycles.

Differential gene expression analysis of RNAseq data. Rockhopper

(http://cs.wellesley.edu/~btjaden/Rockhopper/) was used to analyze the differences in gene

expression between strains and timepoints; the standard settings of the program were utilized for

analysis (30). Following computational analysis via Rockhopper, a change in gene expression

was deemed significant when the *Q*-value (false discovery rate) was <0.05 and a >1.5 fold

change in expression was observed. Visual assessment of read count data was performed using

the Integrated Genome Viewer (IGV) (https://www.broadinstitute.org/igv/) (82, 83). Assessment

of function of the differentially expressed genes was performed using the Clusters of

Orthologous Groups (COG) (84) as previously described in IA3902 (5). Venn diagrams were

used to depict overlap of genes differentially regulated in multiple mutant strains and were

generated using BioVenn (http://www.cmbi.ru.nl/cdd/biovenn/index.php) (85). KEGG Pathways

was used to perform metabolic pathway analysis (http://www.genome.jp/kegg/pathway.html)

(31).

In addition to determining differential gene expression between the mutant and wild-type

strains of IA3902, Rockhopper has the capability to predict non-coding RNAs present within the

data. Prior to further manual analysis, Rockhopper predicted a total of 59 ncRNAs present in

IA3902 (57 chromosomal, 2 pVir). Manual curation was performed to remove candidate non-

coding RNAs if they were related to the 16S or 23S genes as these were thought to be spuriously

identified due to Ribo-Zero depletion differences between replicates. In addition, a single

antisense RNA was discarded due to incorrect annotation of the *rnpB* gene to the wrong strand in

IA3902.

Northern blot detection of CjNC110. For northern blot detection of CjNC110, two

separate experiments utilizing identical conditions were performed as previously described with

some modifications (86). Experiment 1 included wild-type IA3902, ΔCjNC110, and

ΔCjNC110c, while the second included wild-type and ΔluxS. For both, bacterial cells were

grown to early stationary phase of growth in MH broth and total RNA isolation performed as

described above. The total RNA input was normalized to 15 μg using Qubit™ BR RNA Assay

Kit (Invitrogen™, USA) for each respective strain. For northern blot detection**,** total RNA from

bacterial cells (15 μg) and a pre-stained DynaMarker® RNA High Ladder (Diagnocine LLC**,**

Hackensack, NJ) were loaded onto 1.3% denaturing agarose gel (Bio-rad, Hercules, CA),

consisting of 1x MOPS buffer (Fisher, USA) and 5% solution of 37% formaldehyde (Fisher,

USA), electrophoresed at 100V for 1h and transferred to positively-charged nylon membrane

(Roche, Indianapolis, IN) using a Vacuum Blot System (Bio-Rad) transferring at 60mbar for 2

hours. To make RNA cross-linking solution, 245 μl of 12.5 M 1-methylimidazole (Sigma, St.

Louis, MO) was suspended in 9 mL RNase free water, maintaining a pH of 8.0. Then, a total of

0.753 g 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) was added and the total volume

adjusted to 24 mL with RNase-free water. RNA was then cross-linked to the membrane by

placing the nylon membrane, side devoid of RNA, on saturated 3mm Whatman chromatography

paper and incubating it in the freshly prepared EDC solution at 60°C for 2 hours. After cross-

linking the membrane was washed with RNase-free water. Next, CjNC110-LNA DIG-labeled

probe (/5'DigN/ GCACATCAGTTTCAT/3'Dig_N/) (QIAGEN) was added to 15 mL DIG

EasyHyb™ Buffer (Roche) to reach a final concentration of 25 ng/mL. Hybridization was

performed using a hybridization oven and bottle, rotating at 54°C for 12 hours. After overnight

incubation, subsequent washes and incubations were performed again using the hybridization

chamber. The blot was first washed twice with low stringency buffer (2x SSC, 0.1% SDS) for 15

minutes at 60°C, and three times with high stringency buffer (0.1% SSC and 0.1% SDS) for 10

minutes at 60°C. DIG Washing and DIG Blocking Buffers were prepared using the DIG Wash

and Block Set (Roche, USA). The blot was washed twice with Washing Buffer for 10 minutes at

37°C. Next, the blot was incubated in 1x Blocking Buffer for 3 hours at room temperature. DIG

antibody solution was added by mixing Blocking Buffer with DIG antibody at a ratio of 1:5,000.

The blot was then incubated for 30 minutes at room temperature. Next, excess DIG antibody was

removed by rinsing with the DIG Washing Buffer at 25°C a total of four times for 15 minutes

each. For development and detection, the membrane was placed in Development/Detection

Buffer for 10 minutes, and 2 mL ready-to-use CDP-*Star*® (Roche, USA) was added to cover the

blot completely. To image the blot, Chemidoc Imager (Bio-Rad) was used, exposing the blot for

5 minutes on the chemiluminescence setting. Image layout and annotations were performed

using GraphPad Prism. ImageLab software (Bio-Rad) was used to generate a standard curve

using RF and nt size of the ladder to predict the average band size of the target RNA.

887   **Enumeration of *luxS* transcriptional levels utilizing qRT-PCR.** To validate the

RNAseq results and compare *luxS* expression between mutants, quantitative reverse transcriptase

PCR (qRT-PCR) was performed using total RNA isolated from IA3902 wild-type, ΔCjNC110,

and ΔCjNC110c purified in three separate biological replicates of broth cultures grown to 12

hours from the second growth curve described above. Total RNA was extracted and qRT-PCR

was performed using the iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA) and 1000 ng of

total RNA according to the manufacturer's instructions as previously described (64). Each

sample was normalized to the same starting amount of cDNA using Qubit™ BR DNA Assay Kit

(Invitrogen). cDNA purity was measured using the NanoDrop ND-1000 spectrophotometer.

qPCR assays were run using the SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad,

Hercules, CA) and the CFX Maestro™ Real-Time PCR detection system (Bio-Rad). Dilutions of

cDNA template for both standards and all unknowns were run in triplicate with reaction volumes

of 10 µl. Amplification of converted cDNA occurred with 35 cycles of denaturation at 95°C for

10 seconds and then annealing for each primer pair (**Table S9**) at 58°C for 30 seconds. Prior to

analysis, both standards curves (16S and *luxS*) were experimentally validated to have high

efficiency 90%> of amplification and precision R^2=0.98 or greater. Relative fold change in *luxS*

mRNA expression between IA3902 wild-type, ΔCjNC110, and ΔCjNC110c was calculated using

the ISU Gallup Method Equation (87). Statistical analyses was performed using one-way

ANOVA (GraphPad Prism) to determine significance in gene expression levels. A *P*-value of

<0.05 was considered significant.

Liquid chromatography with tandem mass spectrometry-based proteomic analysis.

For extraction of whole cell protein, bacterial cells from IA3902 wild-type, ΔCjNC110 and

ΔCjNC110c were grown to stationary phase of growth at 16 hours on MH agar. Lawn cultures

were then harvested from the plates using 1 mL of MH broth. Collected lawns were normalized

to an A_{600} of 0.2 and aliquoted into 1 mL technical replicates. This was repeated for three

biological replicates. To wash the cells, each aliquot was re-suspended in 1 mL of PBS, cells

were spun down at 8000 x *g* for 3 minutes, and the supernatant was decanted. Next, 50 µl of lysis

buffer (filtered sterilized water + 1% Triton X-100) was added to the protein pellets for

resuspension [adapted from (88)]. The cells were then mechanically sheared by boiling at 96°C

for 5 minutes followed by rapid pipetting. After 3 cycles of boiling-shearing, total cell extracts

were centrifuged at $10,000 \times g$ for 5 min to remove cell debris. The supernatants were collected,

and protein concentration was determined using the Qubit™ Protein Assay and Fluorometer

(Invitrogen™). Crude protein was then submitted to the Iowa State Protein Facility.

Crude protein extracts were digested overnight in solution with trypsin/Lys-C (25-50 μg). After

digestion, Pierce™ Peptide Retention Time Calibration Mixture (PRTC) (ThermoFisher

Scientific) was spiked into the samples to serve as an internal control (25 fmol/μL). The PRTC

areas were used to normalize collision energies, retention times, and peak intensities to allow for

quantitative analysis between samples.

Proteomic analysis was performed via liquid chromatography with tandem mass

spectrometry (LC-MS/MS) using a Q Exactive™ Hybrid Quadrupole-Orbitrap Mass

Spectrometer (Thermo Scientific™) system. The system was coupled with EASY-nLC 1200

nanopump with integrated auto-sampler (Thermo Scientific™). Liquid chromatography was

used to separate the peptides followed by fragmentation and MS/MS analysis. The resulting

intact fragmentation pattern was compared to a theoretical fragmentation pattern (MASCOT) to

find peptides that can be used to identify high-confidence proteins. Minora Feature Detector was

used for label-free quantification to detect and quantify discovered isotopic clusters.

Abundances of proteins were compared by grouping by strain and averaging the

abundances of each protein identified. For data filtering and data imputation, PANDA-view, a

freely available proteomic software (https://sourceforge.net/projects/panda-view/), was used

(89). Briefly, only proteins identified in two out of three biological replicates were included in

analysis. Nearest neighbor (kNN) imputation was used to fill any missing protein values (90). All

data was log transformed using $\log_2$. For statistical analysis, two-way ANOVA with false

discovery rate (*Q*-value) correction set at 0.05 was performed (Graphpad Prism). Graphical

hierarchical clustering heat maps were generated using freely available MetaboAnalyst software

(91). The top 20 proteins were group and analyzed using the following heat map settings:

distance measurement of Euclidean and the average clustering algorithm (91, 92).

***Vibrio harveyi* bioluminescence assay.** Culture samples collected from time points 3, 6,

9 and 12 hours (2 mL each timepoint) from the second growth curve described above were

centrifuged at 20,000 x *g* for 5 minutes at 4ºC. The supernatant was then removed from the cell

pellet and filter-sterilized using a 0.2 μm syringe filter to create cell-free supernatant

representative of the extracellular environment (E-CFS). The cell pellet was then washed twice

with PBS; following each wash, the cells were spun again at 8000 x *g* and the resulting

supernatant was completely removed from the cell pellet. Both the E-CFS and the cell pellet

were then frozen at -80ºC. For processing of the cell pellet to create intracellular cell-free

supernatant (I-CFS), samples were thawed on ice and then resuspended in 600 μl of ice-cold MH

broth. The resuspended cell pellets were then lysed using a Bullet Blender® (NextAdvance,

Troy, NY). To lyse the cells, 1.5 mL microcentrifuge tubes (Corning, Corning, NY) were loaded

with sterilized 0.9-2.0 mm diameter stainless steel beads (NextAdvance). The Bullet Blender was

used for 5 minutes at max speed and stored at 4°C. Lysed cellular debris were spun at 8000 x *g*

for 2 minutes using a microcentrifuge, and the resulting supernatant filtered using a 0.20 μm

sterile syringe filter, resulting in intracellular cell-free supernatant (I-CFS). Both E-CFS and I-

CFS were frozen at -80 ºC until proceeding with the bioluminescence assay.

Autoinducer-2 levels within the collected supernatant for both intracellular (I-CFS) and

extracellular (E-CFS) samples were then enumerated using the *Vibrio harveyi* bioluminescence

assay as previously described (32). Briefly, 10 μL of each CFS sample was added to 90 μL AB

media containing a 1:5000 dilution of the reporter strain, *V. harveyi* strain BB170 (93), in

triplicate. Relative light units (RLU) were measured every 15 minutes over 8 hours using the

FLUOstar Omega (BMG Labtech, Ortenburg, Germany). MH broth and AB media were used as

negative controls, while E-CFS collected from *V. harveyi* strain BB152 (74) was used as a

positive control. The timepoints utilized for analysis were those occurring during the nadir of

values for the negative control wells and at a standardized duration of time (approximately 3

hours) following initiation of increasing values for the positive wells. Results reported are the

average of three independent growth curves with three technical replicates used at each

timepoint. Statistical analysis was conducted using a two-way ANOVA with repeated measures

and Sidak's multiple comparisons test (GraphPad Prism). A *P*-value of <0.05 was considered

significant.

Motility assay. Motility was determined via inoculation of plates consisting of MH broth

with 0.4% agar as previously described in our laboratory (8). Briefly, the A_{600} of overnight

cultures was adjusted to 0.3 using sterile MH broth on a Genesys 10S VIS spectrophotometer

(Thermo Scientific™). A 1 μL volume inoculation stick was then dipped into a set volume of the

standardized culture contained in the bottom of a 15 mL conical tube which was then used to

make a stab inoculation into the center of the freshly made motility agar (MH broth with 0.4%

Bacto agar) with a new inoculation stick for each plate. Plates were incubated at 42°C under

microaerophilic conditions as described above with the exception that the plates were incubated

right-side up and in a single layer. Measurement of the outermost reach of the halo was

performed at 30 hours following inoculation for the first set of experiments and at 24 hours for

the second set of experiments. All strains were assessed for average motility using six biological

replicates consisting of six technical replicates during each independent study. The six

experiments were statistically analyzed using a one-way ANOVA and differences between each

strain assessed via Tukey's multiple comparisons test (GraphPad Prism). A P-value of <0.05 was

considered significant.

988   **Autoagglutination assay.** Autoagglutination was assessed according to the method

described previously (40) with some modifications. Briefly, the A_{600} of overnight cultures was

adjusted to 1.0 in sterile Dulbecco's phosphate buffered saline (PBS) (Corning cellgro,

Manassas, VA) using a Genesys 10S VIS spectrophotometer (Thermo Scientific™). The

suspension was then aliquoted (2 mL each) into standard glass culture tubes. One subset of

cultures were kept at controlled room temperature (25°C) under microaerophilic conditions; the

others were incubated at 37°C microaerophilic. At 24 hours, 1 mL of the upper aqueous phase

was carefully removed and A_{600} measured to determine autoagglutination activity. All strains

were assessed in quadruplicate at each temperature in three independent experiments. The three

experiments were statistically analyzed using one-way ANOVA and differences between each

strain assessed via Tukey's multiple comparisons test (GraphPad Prism). A P-value of <0.05 was

considered significant.

Hydrogen peroxide disk inhibition assay. The hydrogen peroxide (H_2O_2) disk inhibition

assay was performed as previously described with some modifications (63). Briefly, *C. jejuni*

IA3902 wild-type, and mutant strains were grown to log phase in MH broth. The bacteria were

harvested by centrifugation at 12,000 x *g* for 2 minutes and subsequently normalized by

resuspension into MH broth to A_{600} of 1.0. Next, 24 mL of melted MH agar was combined with

1 mL of the normalized bacterial suspension on petri plates, mixed well, and allowed to solidify.

20 μL of aqueous 3% H_2O_2 solution (Sigma-Aldrich, St. Louis, MO) was then pipetted onto 6-

mm diameter disk, which was placed at the center of the plate containing the bacterial agar

suspension. Plates were incubated for 24 hours at 42°C under microaerophilic conditions. The

zones of H_2O_2 sensitivity were measured in millimeters (mm), by measuring the diameter of the

clear zones. The H_2O_2 disk inhibition assay was repeated three times using freshly prepared 3%

H_2O_2 with 4 technical replicates per strain. The data was statistically evaluated using one-way

ANOVA and Dunnett's multiple comparison test. A *P*-value of <0.05 was deemed significant.

Chicken colonization. All studies involving animals were approved by the Iowa State

University Institutional Animal Care and Use Committee (IACUC) prior to initiation (1-18-

8675-G) and followed all appropriate animal care guidelines. Chicken colonization studies were

performed as previously described in our laboratory utilizing 1 day-old broiler chicks obtained

from a commercial hatchery (8). A subset of chicks from each group were screened via cloacal

swabs plated on MH agar containing both selective supplement and growth supplement

(MH+sel+sup) (Oxoid, ThermoFisher Scientific) and found to be negative for *Campylobacter*

carriage prior to inoculation. At 3 days of age, chicks were inoculated by oral gavage with 200

μl of bacterial suspension containing approximately 1×10^7 of one of the tested strains. Groups

of chicks inoculated with specific strains were housed in separate brooders with no contact

between the groups. At pre-determined timepoints post-inoculation (5 days, 12 days, and 19

days), 6 chicks from each group were randomly selected for humane euthanasia which was

immediately followed by necropsy. Cecal contents were harvested aseptically and stored on ice

until further processing could be completed. Cecal samples were then weighed and subjected to a

10-fold serial dilution series, plated on MH+sel+sup agar, and grown under routine culture

conditions as described above.   To confirm that the isolates recovered were in fact the mutant

strain and no cross contamination occurred, representative colonies were selected from each

strain and timepoint, grown in MH+sel+sup broth, and plated on the respective antimicrobial

selective agar of the mutant strain as described above. In addition, a minimum of two CFU from

each group was collected for PCR analysis and confirmation of no cross-contamination between

biological groups. The detection limit of the assay was determined to be 100 CFU/g; for samples

that failed to reach the optimal target range of 30-300 CFU on the initial dilution plate, a count of

3×10^3 was used to enable statistical analysis. For statistical analysis, one-way ANOVA and

Tukey's multiple comparison test was used to determine significant differences in colonization

between biological groups (GraphPad Prism), using the null hypothesis that colonization rates

between groups are the same. A *P*-value of <0.05 was considered significant.

Data availability. The RNAseq dataset generated in this publication has been deposited

in the NCBI Database under BioProject ID Number PRJNA590513

(http://www.ncbi.nlm.nih.gov/bioproject/590513).

1042

1043   **ACKNOWLEDGEMENTS**

1044   The authors would like to thank Dr. Lei Dai and Dr. Zuowei Wu for technical assistance

1045   in mutant construction and northern blot technique.

1046

1047

REFERENCES

1048

1049   1.   WHO. 2015. WHO estimates of the global burden of foodborne diseases: foodborne

1050   disease burden epidemiology reference group 2007-2015.

1051   http://apps.who.int/iris/bitstream/10665/199350/1/9789241565165_eng.pdf. Accessed

1052   December 3, 2019.

1053   2.   Sahin O, Fitzgerald C, Stroika S, Zhao S, Sippy RJ, Kwan P, Plummer PJ, Han J, Yaeger

1054   MJ, Zhang Q. 2012. Molecular evidence for zoonotic transmission of an emergent, highly

1055   pathogenic *Campylobacter jejuni* clone in the United States. J Clin Microbiol 50:680-7.

1056   3.   Sahin O, Plummer PJ, Jordan DM, Sulaj K, Pereira S, Robbe-Austerman S, Wang L,

1057   Yaeger MJ, Hoffman LJ, Zhang Q. 2008. Emergence of a tetracycline-resistant

1058   *Campylobacter jejuni* clone associated with outbreaks of ovine abortion in the United

1059   States. J Clin Microbiol 46:1663-71.

1060   4.   Tang Y, Meinersmann RJ, Sahin O, Wu Z, Dai L, Carlson J, Plumblee Lawrence J,

1061   Genzlinger L, LeJeune JT, Zhang Q. 2017. Wide but variable distribution of a

1062   hypervirulent *Campylobacter jejuni* clone in beef and dairy cattle in the United States.

1063   Appl Environ Microbiol 83.

1064   5.   Wu Z, Sahin O, Shen Z, Liu P, Miller WG, Zhang Q. 2013. Multi-omics approaches to

1065   deciphering a hypervirulent strain of *Campylobacter jejuni*. Genome Biol Evol 5:2217-

1066   30.

1067   6.   Grogono-Thomas R, Blaser MJ, Ahmadi M, Newell DG. 2003. Role of S-layer protein

1068   antigenic diversity in the immune responses of sheep experimentally challenged with

1069   *Campylobacter fetus* subsp. *fetus*. Infect Immun 71:147-54.

7.   van Putten JP, van Alphen LB, Wosten MM, de Zoete MR. 2009. Molecular mechanisms

of *Campylobacter* infection. Curr Top Microbiol Immunol 337:197-229.

8.   Plummer P, Sahin O, Burrough E, Sippy R, Mou K, Rabenold J, Yaeger M, Zhang Q.

2012. Critical role of LuxS in the virulence of *Campylobacter jejuni* in a guinea pig

model of abortion. Infect Immun 80:585-93.

9.   Sahin O, Terhorst SA, Burrough ER, Shen Z, Wu Z, Dai L, Tang Y, Plummer PJ, Ji J,

Yaeger MJ, Zhang Q. 2017. Key role of capsular polysaccharide in the induction of

systemic infection and abortion by hypervirulent *Campylobacter jejuni*. Infect Immun 85.

10.   Wu Z, Periaswamy B, Sahin O, Yaeger M, Plummer P, Zhai W, Shen Z, Dai L, Chen SL,

Zhang Q. 2016. Point mutations in the major outer membrane protein drive

hypervirulence of a rapidly expanding clone of *Campylobacter jejuni*. Proc Natl Acad Sci

U S A 113:10690-5.

11.   Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, Chillingworth T,

Davies RM, Feltwell T, Holroyd S, Jagels K, Karlyshev AV, Moule S, Pallen MJ, Penn

CW, Quail MA, Rajandream MA, Rutherford KM, van Vliet AH, Whitehead S, Barrell

BG. 2000. The genome sequence of the food-borne pathogen *Campylobacter jejuni*

reveals hypervariable sequences. Nature 403:665-8.

12.   Papenfort K, Vogel J. 2010. Regulatory RNA in bacterial pathogens. Cell Host Microbe

8:116-27.

13.   Storz G, Vogel J, Wassarman KM. 2011. Regulation by small RNAs in bacteria:

expanding frontiers. Mol Cell 43:880-91.

14.   Caldelari I, Chao Y, Romby P, Vogel J. 2013. RNA-mediated regulation in pathogenic

bacteria. Cold Spring Harb Perspect Med 3:a010298.

15.   Beisel CL, Storz G. 2010. Base pairing small RNAs and their roles in global regulatory

networks. FEMS Microbiol Rev 34:866-82.

16.   Sharma CM, Hoffmann S, Darfeuille F, Reignier J, Findeiss S, Sittka A, Chabas S,

Reiche K, Hackermuller J, Reinhardt R, Stadler PF, Vogel J. 2010. The primary

transcriptome of the major human pathogen *Helicobacter pylori*. Nature 464:250-5.

17.   Valentin-Hansen P, Eriksen M, Udesen C. 2004. The bacterial Sm-like protein Hfq: a key

player in RNA transactions. Mol Microbiol 51:1525-33.

18.   Livny J, Teonadi H, Livny M, Waldor MK. 2008. High-throughput, kingdom-wide

prediction and annotation of bacterial non-coding RNAs. PLoS One 3:e3197.

19.   Chaudhuri RR, Yu L, Kanji A, Perkins TT, Gardner PP, Choudhary J, Maskell DJ, Grant

AJ. 2011. Quantitative RNA-seq analysis of the *Campylobacter jejuni* transcriptome.

Microbiology 157:2922-32.

20.   Butcher J, Stintzi A. 2013. The transcriptional landscape of *Campylobacter jejuni* under

iron replete and iron limited growth conditions. PLoS One 8:e79475.

21.   Dugar G, Herbig A, Forstner KU, Heidrich N, Reinhardt R, Nieselt K, Sharma CM. 2013.

High-resolution transcriptome maps reveal strain-specific regulatory features of multiple

Campylobacter jejuni isolates. PLoS Genet 9:e1003495.

22.   Porcelli I, Reuter M, Pearson BM, Wilhelm T, van Vliet AH. 2013. Parallel evolution of

genome structure and transcriptional landscape in the *Epsilonproteobacteria*. BMC

Genomics 14:616.

23.   Taveirne ME, Theriot CM, Livny J, Dirita VJ. 2013. The complete *Campylobacter jejuni*

transcriptome during colonization of a natural host determined by RNAseq. PLoS One

8:e73586.

24. Kreuder AJ, Schleining JA, Yaeger M, Zhang Q, Plummer PJ. 2017. RNAseq reveals

complex response of *Campylobacter jejuni* to ovine bile and *in vivo* gallbladder

environment. Front Microbiol 8:940.

25. Le MT, van Veldhuizen M, Porcelli I, Bongaerts RJ, Gaskin DJ, Pearson BM, van Vliet

AH. 2015. Conservation of sigma28-dependent non-coding RNA paralogs and predicted

sigma54-dependent targets in thermophilic *Campylobacter* Species. PLoS One

10:e0141627.

26. Shen Z, Patil RD, Sahin O, Wu Z, Pu XY, Dai L, Plummer PJ, Yaeger MJ, Zhang Q.

2016. Identification and functional analysis of two toxin-antitoxin systems in

Campylobacter jejuni. Mol Microbiol 101:909-23.

27. Alzheimer M, Svensson SL, Konig F, Schweinlin M, Metzger M, Walles H, Sharma CM.

2020. A three-dimensional intestinal tissue model reveals factors and small regulatory

RNAs important for colonization with *Campylobacter jejuni*. PLoS Pathog 16:e1008304.

28. Plummer PJ. 2012. LuxS and quorum-sensing in *Campylobacter*. Front Cell Infect

Microbiol 2:22.

29. Adler L, Alter T, Sharbati S, Golz G. 2014. Phenotypes of *Campylobacter jejuni luxS*

mutants are depending on strain background, kind of mutation and experimental

conditions. PLoS One 9:e104399.

30. McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA,

Vanderpool CK, Tjaden B. 2013. Computational analysis of bacterial RNA-Seq data.

Nucleic Acids Res 41:e140.

31. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2015. KEGG as a

reference resource for gene and protein annotation. Nucleic Acids Res

doi:10.1093/nar/gkv1070.

32. Surette MG, Bassler BL. 1998. Quorum sensing in *Escherichia coli* and *Salmonella*

typhimurium. Proc Natl Acad Sci U S A 95:7046-50.

33. Plummer P, Zhu J, Akiba M, Pei D, Zhang Q. 2011. Identification of a key amino acid of

LuxS involved in AI-2 production in *Campylobacter jejuni*. PLoS One 6:e15876.

34. Hendrixson DR, Akerley BJ, DiRita VJ. 2001. Transposon mutagenesis of

Campylobacter jejuni identifies a bipartite energy taxis system required for motility. Mol

Microbiol 40:214-24.

35. Atack JM, Harvey P, Jones MA, Kelly DJ. 2008. The *Campylobacter jejuni* thiol

peroxidases Tpx and Bcp both contribute to aerotolerance and peroxide-mediated stress

resistance but have distinct substrate specificities. J Bacteriol 190:5279-90.

36. Elvers KT, Park SF. 2002. Quorum sensing in *Campylobacter jejuni*: detection of a *luxS*

encoded signalling molecule. Microbiology 148:1475-81.

37. McNally DJ, Aubry AJ, Hui JP, Khieu NH, Whitfield D, Ewing CP, Guerry P, Brisson

JR, Logan SM, Soo EC. 2007. Targeted metabolomics analysis of *Campylobacter coli*

VC167 reveals legionaminic acid derivatives as novel flagellar glycans. J Biol Chem

282:14463-75.

38. McNally DJ, Hui JP, Aubry AJ, Mui KK, Guerry P, Brisson JR, Logan SM, Soo EC.

2006. Functional characterization of the flagellar glycosylation locus in *Campylobacter*

jejuni 81-176 using a focused metabolomics approach. J Biol Chem 281:18489-98.

39.   Logan SM, Kelly JF, Thibault P, Ewing CP, Guerry P. 2002. Structural heterogeneity of

carbohydrate modifications affects serospecificity of *Campylobacter* flagellins. Mol

Microbiol 46:587-97.

40.   Misawa N, Blaser MJ. 2000. Detection and characterization of autoagglutination activity

by *Campylobacter jejuni*. Infect Immun 68:6168-75.

41.   Saliba AE, S CS, Vogel J. 2017. New RNA-seq approaches for the study of bacterial

pathogens. Curr Opin Microbiol 35:78-87.

42.   Butcher J, Handley RA, van Vliet AH, Stintzi A. 2015. Refined analysis of the

Campylobacter jejuni iron-dependent/independent Fur- and PerR-transcriptomes. BMC

Genomics 16:498.

43.   Chandrashekhar K, Kassem, II, Nislow C, Gangaiah D, Candelero-Rueda RA,

Rajashekara G. 2015. Transcriptome analysis of *Campylobacter jejuni* polyphosphate

kinase (ppk1 and ppk2) mutants. Virulence 6:814-8.

44.   Croucher NJ, Thomson NR. 2010. Studying bacterial transcriptomes using RNA-seq.

Curr Opin Microbiol 13:619-24.

45.   van Vliet AH. 2010. Next generation sequencing of microbial transcriptomes: challenges

and opportunities. FEMS Microbiol Lett 302:1-7.

46.   van Opijnen T, Camilli A. 2013. Transposon insertion sequencing: a new tool for

systems-level analysis of microorganisms. Nat Rev Microbiol 11:435-42.

47.   Waters LS, Storz G. 2009. Regulatory RNAs in bacteria. Cell 136:615-28.

48.   Herzberg M, Kaye IK, Peti W, Wood TK. 2006. YdgG (TqsA) controls biofilm formation

in *Escherichia coli* K-12 through autoinducer 2 transport. J Bacteriol 188:587-98.

49.   Kamaraju K, Smith J, Wang J, Roy V, Sintim HO, Bentley WE, Sukharev S. 2011.

Effects on membrane lateral pressure suggest permeation mechanisms for bacterial

quorum signaling molecules. Biochemistry 50:6983-93.

50.   Golz G, Sharbati S, Backert S, Alter T. 2012. Quorum sensing dependent phenotypes and

their molecular mechanisms in *Campylobacterales*. Eur J Microbiol Immunol (Bp) 2:50-

60.

51.   Xavier KB, Bassler BL. 2005. Regulation of uptake and processing of the quorum-

sensing autoinducer AI-2 in *Escherichia coli*. J Bacteriol 187:238-48.

52.   Holmes K, Tavender TJ, Winzer K, Wells JM, Hardie KR. 2009. AI-2 does not function

as a quorum sensing molecule in *Campylobacter jejuni* during exponential growth in

vitro. BMC Microbiol 9:214.

53.   Adler L, Alter T, Sharbati S, Golz G. 2015. The signalling molecule Autoinducer-2 is not

internalised in *Campylobacter jejuni*. Berl Munch Tierarztl Wochenschr 128:111-6.

54.   Leyn SA, Suvorova IA, Kholina TD, Sherstneva SS, Novichkov PS, Gelfand MS,

Rodionov DA. 2014. Comparative genomics of transcriptional regulation of methionine

metabolism in *Proteobacteria*. PLoS One 9:e113714.

55.   Mitra A, Kesarwani AK, Pal D, Nagaraja V. 2011. WebGeSTer DB--a transcription

terminator database. Nucleic Acids Res 39:D129-35.

56.   Wassenaar TM, Bleumink-Pluym NM, van der Zeijst BA. 1991. Inactivation of

Campylobacter jejuni flagellin genes by homologous recombination demonstrates that

flaA but not *flaB* is required for invasion. EMBO J 10:2055-61.

57.   Yao R, Burr DH, Doig P, Trust TJ, Niu H, Guerry P. 1994. Isolation of motile and non-

motile insertional mutants of *Campylobacter jejuni*: the role of motility in adherence and

invasion of eukaryotic cells. Mol Microbiol 14:883-93.

58.   Golden NJ, Acheson DW. 2002. Identification of motility and autoagglutination

Campylobacter jejuni mutants by random transposon mutagenesis. Infect Immun

70:1761-71.

59.   Guerry P, Ewing CP, Schirm M, Lorenzo M, Kelly J, Pattarini D, Majam G, Thibault P,

Logan S. 2006. Changes in flagellin glycosylation affect *Campylobacter*

autoagglutination and virulence. Mol Microbiol 60:299-311.

60.   Thibault P, Logan SM, Kelly JF, Brisson JR, Ewing CP, Trust TJ, Guerry P. 2001.

Identification of the carbohydrate moieties and glycosylation motifs in *Campylobacter*

jejuni flagellin. J Biol Chem 276:34862-70.

61.   Howard SL, Jagannathan A, Soo EC, Hui JP, Aubry AJ, Ahmed I, Karlyshev A, Kelly

JF, Jones MA, Stevens MP, Logan SM, Wren BW. 2009. *Campylobacter jejuni*

glycosylation island important in cell charge, legionaminic acid biosynthesis, and

colonization of chickens. Infect Immun 77:2544-56.

62.   Guerry P, Doig P, Alm RA, Burr DH, Kinsella N, Trust TJ. 1996. Identification and

characterization of genes required for post-translational modification of *Campylobacter*

coli VC167 flagellin. Mol Microbiol 19:369-78.

63.   Palyada K, Sun YQ, Flint A, Butcher J, Naikare H, Stintzi A. 2009. Characterization of

the oxidative stress stimulon and PerR regulon of *Campylobacter jejuni*. BMC Genomics

10:481.

64.   Mou KT, Plummer PJ. 2016. The impact of the LuxS mutation on phenotypic expression

1225   of factors critical for *Campylobacter jejuni* colonization. Vet Microbiol 192:43-51.

1226   65.   Mou KT. 2015. Mechanisms and roles of the LuxS system, methyl recycling, and DNA

1227   methylation on the physiology of *Campylobacter jejuni*. PhD. Iowa State University,

1228   Graduate Theses and Disserations.

1229   66.   Plummer PJ. 2009. The mechanisms and roles of quorum sensing in the pathobiology of

1230   *Campylobacter jejuni*. PhD. Iowa State University, Graduate Theses and Disseratations.

1231   67.   He Y, Frye JG, Strobaugh TP, Chen CY. 2008. Analysis of AI-2/LuxS-dependent

1232   transcription in *Campylobacter jejuni* strain 81-176. Foodborne Pathog Dis 5:399-415.

1233   68.   Jeon B, Itoh K, Misawa N, Ryu S. 2003. Effects of quorum sensing on *flaA* transcription

1234   and autoagglutination in *Campylobacter jejuni*. Microbiol Immunol 47:833-9.

1235   69.   Wosten MM, Wagenaar JA, van Putten JP. 2004. The FlgS/FlgR two-component signal

1236   transduction system regulates the *fla* regulon in *Campylobacter jejuni*. J Biol Chem

1237   279:16214-22.

1238   70.   Malik-Kale P, Parker CT, Konkel ME. 2008. Culture of *Campylobacter jejuni* with

1239   sodium deoxycholate induces virulence gene expression. J Bacteriol 190:2286-97.

1240   71.   Joslin SN, Hendrixson DR. 2009. Activation of the *Campylobacter jejuni* FlgSR two-

1241   component system is linked to the flagellar export apparatus. J Bacteriol 191:2656-67.

1242   72.   Hendrixson DR, DiRita VJ. 2003. Transcription of sigma54-dependent but not sigma28-

1243   dependent flagellar genes in Campylobacter jejuni is associated with formation of the

1244   flagellar secretory apparatus. Mol Microbiol 50:687-702.

73.   Van Puyvelde S, Vanderleyden J, De Keersmaecker SC. 2015. Experimental approaches

to identify small RNAs and their diverse roles in bacteria--what we have learnt in one

decade of MicA research. Microbiologyopen 4:699-711.

74.   Bassler BL, Wright M, Showalter RE, Silverman MR. 1993. Intercellular signalling in

Vibrio harveyi: sequence and function of genes regulating expression of luminescence.

Mol Microbiol 9:773-86.

75.   Wang Y, Taylor DE. 1990. Chloramphenicol resistance in *Campylobacter coli*:

nucleotide sequence, expression, and cloning vector construction. Gene 94:23-8.

76.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. 2009.

Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods

6:343-5.

77.   Wang Y, Taylor DE. 1990. Natural transformation in *Campylobacter* species. J Bacteriol

172:949-55.

78.   Muraoka WT, Zhang Q. 2011. Phenotypic and genotypic evidence for L-fucose

utilization by *Campylobacter jejuni*. J Bacteriol 193:1065-75.

79.   Karlyshev AV, Wren BW. 2005. Development and application of an insertional system

for gene delivery and expression in *Campylobacter jejuni*. Appl Environ Microbiol

71:4004-13.

80.   Chen CY, Nace GW, Irwin PL. 2003. A 6 x 6 drop plate method for simultaneous colony

counting and MPN enumeration of *Campylobacter jejuni*, *Listeria monocytogenes*, and

Escherichia coli. J Microbiol Methods 55:475-9.

81.   Luo Y, Sahin O, Dai L, Sippy R, Wu Z, Zhang Q. 2012. Development of a loop-mediated

isothermal amplification assay for rapid, sensitive and specific detection of a

Campylobacter jejuni clone. J Vet Med Sci 74:591-6.

82.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov

JP. 2011. Integrative genomics viewer. Nat Biotechnol 29:24-6.

83.   Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. Brief Bioinform 14:178-

92.

84.   Galperin MY, Makarova KS, Wolf YI, Koonin EV. 2015. Expanded microbial genome

coverage and improved protein family annotation in the COG database. Nucleic Acids

Res 43:D261-9.

85.   Hulsen T, de Vlieg J, Alkema W. 2008. BioVenn - a web application for the comparison

and visualization of biological lists using area-proportional Venn diagrams. BMC

Genomics 9:488.

86.   Kim SW, Li Z, Moore PS, Monaghan AP, Chang Y, Nichols M, John B. 2010. A

sensitive non-radioactive northern blot method to detect small RNAs. Nucleic Acids Res

38:e98.

87.   Gallup JM, Ackermann MR. 2006. Addressing fluorogenic real-time qPCR inhibition

using the novel custom Excel file system 'FocusField2-6GallupqPCRSet-upTool-001' to

attain consistently high fidelity qPCR reactions. Biol Proced Online 8:87-152.

88.   Hobb RI, Fields JA, Burns CM, Thompson SA. 2009. Evaluation of procedures for outer

membrane isolation from *Campylobacter jejuni*. Microbiology 155:979-88.

89.   Chang C, Xu K, Guo C, Wang J, Yan Q, Zhang J, He F, Zhu Y. 2018. PANDA-view: an

1289   easy-to-use tool for statistical analysis and visualization of quantitative proteomics data.

1290   Bioinformatics 34:3594-3596.

1291   90.   Beretta L, Santaniello A. 2016. Nearest neighbor imputation algorithms: a critical

1292   evaluation. BMC Med Inform Decis Mak 16 Suppl 3:74.

1293   91.   Chong J, Yamamoto M, Xia J. 2019. MetaboAnalystR 2.0: From Raw Spectra to

1294   Biological Insights. Metabolites 9.

1295   92.   Key M. 2012. A tutorial in displaying mass spectrometry-based proteomic data using heat

1296   maps. BMC Bioinformatics 13 Suppl 16:S10.

1297   93.   Bassler BL, Wright M, Silverman MR. 1994. Multiple signalling systems controlling

1298   expression of luminescence in *Vibrio harveyi*: sequence and function of genes encoding a

1299   second sensory pathway. Mol Microbiol 13:273-86.

1300

 FIGURE LEGENDS

 FIG 1. Northern blotting demonstrates expression of CjNC110 in wild-type IA3902 and

 ΔCjNC110c and absence of expression in ΔCjNC110. Cultures for RNA extraction were

 collected at early stationary phase of growth and northern blot analysis was conducted using 15

 μg of total RNA in each of three separate replicates per strain tested.   The arrow indicates the

 most prominent band which corresponds to the previously predicted size of CjNC110 in *C. jejuni*

 (21).

 FIG 2. Venn diagram depicts overlap of shared up- and downregulated genes during the

 exponential and stationary growth phases of all 3 mutant strains. BioVenn was used to

 compare the lists of known genes identified by Rockhopper as upregulated in the RNAseq data

 in all 3 mutant strains during both exponential [A] and stationary [B] growth; downregulated

 genes were also compared for all 3 mutant strains during both exponential [C] and stationary [D]

 growth.

 FIG 3. Shaking growth curve of wild-type IA3902 (WT) and isogenic mutants (mean ±

 SEM) confirms normal exponential growth. Results of four replicates (A_{600}) of a shaking

 growth curve performed in 250 mL Erlenmeyer flasks under microaerophilic conditions in MH

 broth. Samples were pulled from this growth curve and used for qRT-PCR and AI-2

 bioluminescence assays; for the results of the growth curve from which the RNAseq samples

 were taken, please see **Fig. S2A and S2B.** Significance is denoted by "*" above each timepoint.

FIG 4. Bioluminescence activity as measured via the *Vibrio harveyi* bioassay over the course

of growth demonstrates (A) decreased extracellular AI-2 levels with (B) a concomitant

increase in intracellular AI-2 in ΔCjNC110 when compared to wild-type IA3902 (WT)

(mean ± SEM). The *V. harveyi* bioluminescence assay was used initially to detect AI-2 levels in

the extracellular CFS to approximate LuxS activity (A). A second experiment was performed to

investigate AI-2 levels in the intracellular CFS (B). Each bar represents the average RLU (AI-2

activity level) from three biological replicates consisting of three technical replicates each.

Significance is denoted by "*" above each timepoint.

1330

1331 **FIG 5. qRT-PCR confirms *luxS* is transcribed at higher levels in the ΔCjNC110 and**

1332 **ΔCjNC110c backgrounds when compared to IA3902 wild-type (WT) (mean ± SEM).** *C.*

1333 *jejuni* cultures were collected at 12 hours for RNA conversion to cDNA for qRT-PCR to assess

1334 expression of mRNA *luxS*. Each bar represents the average fold change expression from three

1335 biological replicates. Significance is denoted by "*" above each timepoint.

1336

FIG 6. Motility assays confirm isogenic mutants remain motile, with ΔCjNC110 motility

increased and ΔluxS motility decreased over IA3902 wild-type (WT) (mean± SEM).

Swarming motility in semi-solid agar was assessed via measurement in mm of the outermost

reach of growth at both 30 hours and 24 hrs. Each bar represents the average motility of six

biological replicates from six independent studies. Significance is denoted by "*" above each bar

when comparing wild-type to other respective strains at each independent timepoint.

1343

FIG 7. ΔCjNC110 displays increased hydrogen peroxide (H_2O_2) sensitivity *in vitro* when

compared to wild-type IA3902 (WT) (mean ±SEM). Cell cultures normalized to an A_{600} of 1.0

were inoculated into melted MH agar then incubated with a 6 mm disk soaked in 3% H_2O_2 placed

at the center of the solidified MH agar plate; after 24 hours the zone of sensitivity to H_2O_2 was

measured in mm. Each bar represents the average of three biological replicates from four

independent studies. Significance is denoted by "*" above each bar when comparing wild-type to

other respective strains at each independent timepoint.

1351

FIG 8. Autoagglutination ability of IA3902 is decreased in ΔCjNC110 and increased in

ΔluxS when compared to wild-type IA3902 (WT) (mean± SEM). Autoagglutination as

determined by optical density (A_{600}) was measured following a 24 hour incubation at either 25°C

or 37°C. Each bar represents the average autoagglutination of 3 biological replicates consisting

of 4 technical replicates during each independent study. An increase in optical density correlates

to decreased autoagglutination ability, and a decrease in optical density correlates to increased

autoagglutination ability. Significance is denoted by "*" above each bar when comparing wild-

type to other respective strains at each independent timepoint.

1360

FIG 9. Sustained colonization in chickens is impaired in ΔCjNC110 and absent in ΔluxS

and ΔCjNC110ΔluxS mutants compared to wild-type (mean ± SEM). Results of a single

study investigating the colonization ability of mutants strains compared to wild-type (WT)

IA3902 are reported as $\log_{10}$ CFU/mL of ceca content for each sampling day [DPI 5 (Week 1),

DPI 12 (Week 2), and DPI 19 (Week 3)]. Each bar represents the average colonization levels for

each biological group with weekly technical replicates using a minimum of 6 birds. Significance

is denoted by "*" above each timepoint.

1368

1369   **TABLE 1.** Bacterial strains utilized in this study.

Strain	Description	Source or Reference
Campylobacter jejuni		
W7	Wild-type motile variant of NCTC 11168	(8)
W7 ΔCjNC110	W7 ΔCjNC110::Cm[R]	This study
Sheep Abortion (SA) IA3902	Wild-type *C. jejuni*	(3)
IA3902 ΔCjNC110	IA3902 ΔCjNC110::Cm[R]	This study
IA3902 ΔluxS	IA3902 ΔluxS::Kan[R]	(33)
IA3902 ΔCjNC110ΔluxS	IA3902 ΔCjNC110::Cm[R] ΔluxS::Kan[R]	This study
IA3902 ΔCjNC110c	IA3902 ΔCjNC110::Cm[R] CjNC110::Kan[R]	This study
Escherichia coli		
DH5α	*fhuA2* Δ(*argF-lacZ*)U169 *phoA gln*V44 Φ80 Δ(*lacZ*)M15 *gyrA*96 *recA1 relA1 endA1 thi-1 hsdR*17	New England Biolabs, Ipswich, MA
Vibrio harveyi		
BB152	AI-1-/AI-2+; luxM::Tn5	(74)
BB170	AI-2 reporter strain; luxN::Tn5	(93)

Kan[R] = kanamycin resistance cassette

Cm[R] = chloramphenicol resistance cassette

1370

1371

1372   **TABLE 2**. Summary of differential gene expression results between mutant strains.

	Condition					
	ΔCjNC110		ΔluxS		ΔCjNC110ΔluxS	
	3 h	12 h	3 h	12 h	3 h	12 h
Protein-coding genes						
Number downregulated	6	16	1	6	61	32
Number upregulated	4	7	14	6	47	29
Non-coding RNA genes						
Number downregulated	0	0	0	0	25[b]	2
Number upregulated	1[a]	0	1[a]	0	1[a]	0

[a] = previously described ncRNA, CjNC140 (21)

[b] = 17 tRNA genes, 3 known RNA genes (tmRNA, SRP, 6S) and 4 newly predicted ncRNA

1373
1374

1375 **TABLE 3.** Summary of differential gene expression (>1.5 fold change, Q <0.05) in the IA3902 ΔCjNC110, ΔluxS and
1376 ΔCjNC110ΔluxS mutants as determined by RNAseq and analyzed via Rockhopper.

	Exponential (3 h)						Stationary (12 h)			
	ΔCjNC110*	ΔluxS*	ΔCjNC110ΔluxS*				ΔCjNC110*	ΔluxS*	ΔCjNC110ΔluxS*	
Genes downregulated	*hisF*	CJSA_1350	CJSA_0014	*mogA*	CJSA_1389	CJSA_t0002	*panC*	CJSA_0560	*rnhB*	*aroB*
	pseA		CJSA_0041	*aspB*	*flgK*	CJSA_t0005	CJSA_0559	CJSA_0620	CJSA_0158	CJSA_1137
	neuB2		*flgD*	CJSA_0788	*moaE*	CJSA_t0007	CJSA_0687	CJSA_1349	CJSA_0160	CJSA_1164
	ptmA		*flgE*	*flgL*	*nuoD*	CJSA_t0012	***tpx***	CJSA_1350	CJSA_0284	*pyrH*
	CJSA_1352		CJSA_0067	*rpmH*	CJSA_1562	CJSA_t0013	CJSA_0785	*acs*	*motB*	CJSA_1349
	rpsN		*accD*	CJSA_0920	CJSA_1568	CJSA_t0015	*ciaB*	*leuC*	*trpD*	CJSA_1350
			trxA	CJSA_1093	CJSA_1577	CJSA_t0017	CJSA_1102		CJSA_0344	CJSA_1414
			panB	CJSA_1102	*secY*	CJSA_t0018	*petC*		CJSA_0372	*acs*
			motB	*dctA*	*rplO*	CJSA_t0020	CJSA_1137		CJSA_0389	CJSA_1461
			fliN	*luxS*	*rpsE*	CJSA_t0021	CJSA_1244		CJSA_0396	*rnhA*
			rpsU	CJSA_1182	*rplR*	CJSA_t0033	*hisF*		*sdhA*	CJSA_1560
			frdB	*porA*	*rplF*	CJSA_t0035	***neuB2***		*sdhB*	*leuC*
			flaG	*hydD*	*rpsH*	CJSA_t0036	CJSA_1266		*uxaA*	*leuB*
			fliS	*pseB*	*rpsN*	CJSA_t0037	***ptmB***		CJSA_0560	*leuA*
			CJSA_0521	*pseC*	*rplE*	CJSA_t0038	***ptmA***		CJSA_0616	CJSA_pVir0030
			CJSA_0569	CJSA_1233	*rplX*	CJSA_t0039	CJSAt0002		*trmD*	
			pstS	*neuC2*	*rplN*	CJSA_t0043				
			hslV	CJSA_1350	*rpmC*	CjNC130				
			flgH	CJSA_1351	*rplP*					
			flgG2	*flgI*	CJSA_CjSRP1					
			flgG	CJSA_1388	*ssrA*					
Genes upregulated	CJSA_0008	*dnaN*	*dnaN*	*trpF*	*cetB*	CJSA_1449	*hcrA*	*flgE*	*dsbI*	CJSA_1107
	cetA	CJSA_0008	CJSA_0008	*trpB*	*cetA*	*rloH*	CJSA_0716	*flgG2*	CJSA_0040	*omp50*
	cetB	*trpF*	*gltD*	*trpA*	CJSA_1129	*nuoM*	CJSA_0717	*flgG*	CJSA_0041	CJSA_1180
	CJSA_1261	*trpB*	*folk*	CJSA_0337	CJSA_1131	CJSA_1549	CJSA_1107	*hcrA*	*flgD*	*pseB*
	CjNC140	*trpD*	CJSA_0076	CJSA_0370	CJSA_1145	*leuC*	*omp50*	CJSA_1107	*flgE*	CJSA_1262
		CJSA_0337	CJSA_0105	CJSA_0372	CJSA_1164	*leuB*	***luxS***	*nrfA*	*flgC*	*neuB2*
		CJSA_0370	CJSA_0110	*sdhA*	*cbpA*	*leuA*	**CJSA_1261**		*flgB*	*neuC2*
		CJSA_0732	*dgkA*	*sdhB*	CJSA_1259	CjNC140			*flgH*	*flgI*
		CJSA_1017	*pyrC*	CJSA_0490	CJSA_1260	CJSA_pVir0042			*flgG2*	CJSA_1387
		CJSA_1131	*Tal*	CJSA_0491	CJSA_1301	CJSA_pVir0044			*flgG*	*flgK*
		CJSA_1301	CJSA_0305	CJSA_0836	CJSA_1343	CJSA_pVir0025			*hcrA*	CJSA_1562
		CJSA_1352	*trpD*	*Cfa*	*tagF*				CJSA_0716	*p19*
		CJSA_1449							*flgS*	CJSA_pVir0042
		CJSA_1549							*flgL*	CJSA_pVir0013
		CJSA_pVir0042							CJSA_0969	
		CjNC140								

* = additional small RNAs predicted by Rockhopper present with differential expression; see S4, S5, and S6 Tables for additional information

BOLD indicates a gene for which further phenotypic investigation of the functional significance of the observed differential expression was performed [*pseA, neuB2 (ptmC), CJSA_1261 (ptmG), ptmA, ptmB* = autoagglutination; *cetA, cetB* = motility; *Tpx* = hydrogen peroxide sensitivity; and *luxS* = AI-2 assay]

1377

1378

1379

1380   **TABLE 4:** Significant differentially expressed proteins identified from proteomic analysis
1381   ΔCjNC110 and ΔCjNC110c versus IA3902 wild-type

Protein ID	Geometric Mean (Abundance)		Fold Change vs WT	Q-value*
	WT	Mutant		
ΔCjNC110				
CJSA_0067 - iron-sulfur protein	23726566.41	851708.3688	-27.9	<0.0001
CJSA_0489 - rhodanese-like domain-containing	35962749.77	2965820.801	-12.1	0.0001
ThiJ - thiazole monophosphate synthesis protein	1589344.014	31307392.07	19.7	<0.0001
FrdA - fumarate reductase flavoprotein subunit	3178688.029	280958.9826	-11.3	0.0002
LuxS - S-ribosylhomocysteine lyase	**3913424.009**	**50859008.46**	**13.0**	**<0.0001**
ΔCjNC110c				
CJSA_0067 - iron-sulfur protein	23726566.41	1123835.93	-21.1	<0.0001
CJSA_0489 - rhodanese-like domain-containing	35962749.77	2767208.654	-13.0	0.0001
CJSA_0166 – lipoprotein	25429504.23	2581896.969	-9.8	0.0006
RplN - 50S ribosomal protein L14	10327587.87	1048576	-9.8	0.0007
RplU - 50S ribosomal protein L21	15653696.03	3178688.029	-4.9	0.0007
RecJ - single-stranded-DNA-specific exonuclease	2408995.053	19271960.42	8.0	0.002
AtpF - ATP synthase subunit B	2965820.801	27254667.8	9.2	0.001
LuxS - S-ribosylhomocysteine lyase	**3913424.009**	**54509335.6**	**13.9**	**<0.0001**
CJSA_0926 - lipoprotein	1703416.738	23726566.41	13.9	0.0001

Geometric mean calculated by back-transforming log2 transformation value

*Two-way ANOVA FDR set at 0.05

WT = wild-type

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392
1393