Characterization of the Intestinal Absorption of Morroniside from

***Cornus officinalis Sieb. et Zucc* via a Caco-2 Cell Monolayer Model**

Short title: Intestinal Absorption of Morroniside via Caco-2 Cell

Renjie Xu[1, 2¶], Hongdan Zhu [3¶], Lingmin Hu[4], Beimeng Yu[3], Xiaohua

Zhan[5], Yichu Yuan[6*], Ping Zhou[1*]

[1] Department of Clinical pharmacy, Shaoxing Women and Children's Hospital,

Shaoxing, Zhejiang, China.

[2] Department of Pharmacy, Xinhua Hospital Affiliated to Shanghai Jiao Tong

University School of Medicine, Shanghai, China.

[3] Neonatal Intensive Care Unit, Shaoxing Women and Children's Hospital, Shaoxing,

Zhejiang, China.

[4] Department of Laboratory, Shaoxing Seventh People's Hospital, Shaoxing,

Zhejiang, China.

[5] The Third Maternal wards, Shaoxing Women and Children's Hospital, Shaoxing,

Zhejiang, China.

[6] Department of Urology, The Second Affiliated Hospital, School of Medicine,

Zhejiang University, Hangzhou, Zhejiang, China

¶These authors contributed equally to this work.

*Correspondence should be addressed to Yichu Yuan; ycyuan@zju.edu.cn and Ping

Zhou; 290772309@qq.com

Abstract

Morroniside is biologically active polyphenols found in *Cornus officinalis Sieb. et*

Zucc (CO) which exhibits a broad spectrum of pharmacological activities, such as

protecting nerves, preventing diabetic liver damage and renal damage. However, little

data is available regarding its mechanism of intestinal absorption. Here, a human

intestinal epithelial cell model Caco-2 cell *in vitro* cultured was applied to study on

the absorption and transport of morroniside, the effects of time, donor concentration,

pH, temperature and inhibitors, the absorption and transport of morroniside were

investigated. The bidirectional permeability of morroniside from the apical (AP) to

the basolateral (BL) side and in the revese direction was studied.   When

administrated was set at three tested concentrations (5, 25 and 100μM), the P_{app} value

in AP-to-BL direction was ranged from 1.59 to 2.66×10^{-6}cm/s. In the reverse

direction, BL-to-AP, the value was ranged from 2.67 to 4.10×10^{-6}cm/s. The data

indicated that morroniside transport was both pH- and temperature- dependent. The

morroniside's permeability process affected by treatment with various inhibitors, such

as the multidrug resistance protein inhibitors MK571 and benzbromarone, the breast

cancer resistance protein inhibitor apigenin. It can be found that the mechanisms of

intestinal absorption of morroniside may involve multiple transport pathways like

passive diffusion as well as efflux protein-mediated active transport especially the

multidrug resistance protein2 and breast cancer resistance protein. After CO was

added, $P_{app}AB$ increased significantly by about 125.26%, it can therefore be assumed

that some ingredients in the crude material promote morroniside's absorption in the

small intestine.

Keywords: Caco-2; Morroniside; multidrug resistance protein; absorption

mechanism

Introduction

Traditional Chinese medicines (TCM) are natural therapeutic remedies that have been

widely used for thousands of years (1). Morroniside (Fig 1), one of the most

important iridoid glycosides, is the main active ingredient of *Cornus officinalis Sieb.*

et Zucc (CO). It is a rich source of iridoid glycosides and has been used as a

traditional Chinese medicinal herb for centuries (2). Various pharmacological studies

have indicated that morroniside was effective in the treatment of Alzheimer's disease

(3) for protecting nerves (4), preventing diabetic liver damage (5) and renal damage

(6), having beneficial effects on lipid metabolism and inflammation (7) and anti-

anaphylactic activity (8). Since morroniside and its correlative plant extracts have

exhibited these pharmacological effects, it would be hopeful that morroniside to be

developed to some promising preparations of herbal medicinal products.

Several studies have been conducted to determine the concentration of

morroniside in biological matrices (9-11). The absolute oral bioavailability of

morroniside was calculated to be 7.0, 6.1 and 3.6% for low, medium and high dosage,

respectively (9, 12). It seemed like that the plasma levels of morroniside

administrated intravenously were much higher than those after oral administration. It

is well known that oral administration is the main route for applying TCM and they

should be absorbed in the gastrointestinal tract to exhibit pharmacologic effects (13).

The intestinal absorption barrier is a major factor controlling the absorption and oral

bioavailability of drugs(14-16) and it is here that the first steps of pharmacokinetics

occur after oral intake. Therefore, exploration of the intestinal absorption mechanism

of morroniside is necessary not only for the *in vivo* pharmacokinetics study but also to

provide some key information for their effective delivery system.

The aim of present study was to further investigate the intestinal absorptive

characteristics of morroniside using Caco-2 cells. This model is extensively used

because of its morphological and functional similarities to the human small intestinal

epithelium, and it has been recognized by the FDA as a viable model that replicates

human intestinal absorption (17-19). The author aimed at further revealing the reason

of low bioavailability of morroniside and providing a theoretical basis for the

development of formulations.

81   Fig 1 The chemical structure of morroniside

Materials and Methods

Materials and Reagents

Transwell permeable polycarbonate inserts (0.4 μm) and 12-well cell culture plates

were obtained from Corning. (Cambridge, MA, USA). Caco-2 cell line was

generously provided by Cell Bank of the Chinese Academy of Sciences (Shanghai,

China). Dulbecco's Modified Eagle's Medium (DMEM) was obtained from Gibco

Laboratories (Life Technologies Inc., Grand Island, NY, USA). Hanks' balanced salt

solution (HBSS, powder form) was obtained from Sigma Chemical Co. (Deisenhofen,

Germany). Fetal bovine serum was purchased from Hyclone (Logan, UT, USA). 100×

nonessential amino acids, 100× penicillin and streptomycin, 0.25% trypsin with

ethylene- diaminetetraacetic acid (EDTA), 1 M 4-(2-hydroxyethyl)-1-piperazi-

neethanesulfonic acid (HEPES), and BSA were all purchased from Invitrogen Corp.

(Carlsbad,CA). Verapamil, MK 571, indomethacin, benzbromarone, apigenin, sodium

vanadate and cimetidine used in this study were obtained from Aladdin Industrial Inc.

(Shanghai, China). Morroniside (purity > 98.0%) and loganin (internal standard (IS),

purity > 98.0%) were obtained from the National Institute for the Control of

Pharmaceutical and Biological Products (Beijing, China). All other reagents (typically

analytical grade or better) were used as received.

Herb extract preparation

The extract *was* prepared according to reported methods (20). CO (100 g) crude

material was decocted with 1000 mL water for 2 h. Then, the filtrate was collected

and the residue was decocted with 1000 mL water for another 2 h. Finally, the two

batches of filtrates were combined and concentrated to 100 mL. The solution was

centrifuged to pellet any insoluble molecules then filter sterilized with a 0.22 μm

(PEM) syringe filter. The contents of morroniside in preparation turned out about

16.02mg/g in crude material after determined by a HPLC method(9).

Caco-2 Cell Culture and Cytotoxicity Assay

The conditions for Caco-2 cell culture have been described previously(21). In brief,

Cells were cultured in a humidified atmosphere of 5% CO_2 and 95% air at 37 °C. The

culture medium (DMEM) was supplemented with 10% (v/v) fetal bovine serum, 1%

nonessential amino acids, 1% penicillin, and streptomycin. The medium was changed

every 2-3 days. When the cell monolayer reached 80-90% confluence, the cells were

detached with a solution of trypsin (0.5 mg/ml) and EDTA (ethylene diamine

tetraacetic acid, 0.2 mg/ml) (PAA) and reseeded at a density of 5×10^4 cells/cm^2.

For Cytotoxicity Assay, the cells were incubated in 96-well plate for 24 h,

MTT solution was prepared at 1 mg/mL in PBS and was filtered through a 0.2 μm

filter. Then, 20 μL of MTT was added into each well. Cells were incubated for 4 h at

37 °C with 5% CO_2, 95% air and complete humidity. After 4 h, the MTT solution was

removed and replaced with 150 μL of DMSO. The plate was further incubated for 5

min at room temperature, and the optical density (OD) of the wells was determined

using a plate reader at a test wavelength of 490 nm.

Transport of Analytes across the Caco-2 Monolayer

An initial stock solution of morroniside in methanol was prepared. The stock solution

was then diluted with HBSS. The transepithelial electrical resistance (TEER) value

was measured with a Millipore ERS voltameter (Millipore, Massa- chusetts, USA) in

order to evaluate and determine the monolayer integrity. The monolayers were ready

for experiments from 19 to 22 days after seeding. Only monolayers that demonstrated

a TEER value above 400 $\Omega \times cm^2$ were used for the experiment.

During incubation, the culture medium was refreshed every other day. The culture

medium was discarded, 500 μL of HBSS was added to each well and incubated for 20

min at 37 ℃. Afterward, each of the well was rinsed 2 times with HBSS at 37 ℃.

The test compounds were added to either the apical (AP, 0.5 mL) or basolateral side

(BL, 1.0 mL), while the receiving chamber contained the corresponding volume of

blank HBSS medium. Incubation was performed at 37℃ for 180 min, with shaking at

50 rpm. To assess the drugs transport from AP to BL, at the incubation of 5, 15, 30,

45, 60, 90, 120 and 180 min, a 50 μL of the solution from BL or AP side was

collected, and replaced with an equal volume of HBSS. The effect on uptake was

investigated at 4 ℃ and 37 ℃. One plate was placed in a cell incubator at 37 ℃,

with 5% CO_2, and the other was in a 4 ℃ refrigerator. Caco-2 cells were seeded at 5

$\times 10^4$ cells cm^2 in a 12-well plate. After 19 day of incubation, the effect of

temperature on cellular uptake was investigated by 60 min of incubation with

morroniside at 25 μg/mL. To study the effect of pH on the morroniside transport, the

experiments were performed in HBSS buffer, which was adjusted to pH 6.5 or pH 7.4

on the apical side (0.5 mL) depending on the experiment, and to pH 7.4 on the

basolateral side (1.5 mL).

Efflux and influx transporters were investigated for their effects on the

transport flux of morroniside. Several ABC transporter inhibitors, one p-glycoprotein

inhibitor (100μM verapamil) (22), three multidrug resistance- related protein inhibitor

(100μM MK571, 50μM benzbromarone and 200μM indomethacin) (23-25), and one

breast cancer resistance protein inhibitor (25μM apigenin) (26) were used to

determine the transporters involved in the efflux transport of morroniside.; Inhibiting

the influx during morroniside transport across the Caco-2 cell monolayers was

investigated by adding 50μM sodium vanadate and cimetidine to evaluate the

selectivity of Na/K-ATPase (27).

Sample Processing

Samples were treated with a protein precipitation method. After spiked with 4-fold

volume of acetonitrile containing 5% acetic acid and 70 ng/mL IS, the tubes were

vortexed for 1min. The tubes were centrifuged at 14,000rpm for 10min, and the

supernatants were collected and filtered through a 0.22μm hydrophobic membrane.

An aliquot of 2 μL was injected into the LC-MS/MS system for analysis.

UPLC-MS/MS analytical methods

An API 5500 triple-quadrupole mass spectrometer (Applied Biosystems-SCIEX,

Concord, Canada) equipped with an electrospray ionization (ESI) source were used

for chromatographic analyses. The separation was achieved with a ZORBAX Eclipse

Plus C_{18} column (50 mm × 2.1 mm, 1.8 μm). The mobile phase consisted of 0.1%

acetic acid in aqueous solution as solvent A and 100% acetonitrile solution as solvent

B, and the flow rate was 0.4 mL/min. The following gradient elution was used: 0-1.0

min 1% B; 1.0-2.2 min 1→29% B; 2.2-2.3 min 29→95% B; 2.3-3.3 min 95% B; 3.3-

3.4 min 95-1% B, and followed by a 1.0 min re-equilibration to the initial condition.

The column was maintained at 30℃ and the autosampler tray at 30℃.

173   Analytical procedures were evaluated with negative electrospray ionization

174   (ESI) mode. Typical parameters for the Turbo Ionspray source set were: ion spray

175   source temperature at 550 ℃, ionspray voltage (IS) at 5000V; Gas 1 and Gas 2

176 (nitrogen) at 50 psi; CUR at 50; CAD at -2.0. Other parameters were shown in Table

177 1.

178 Table 1 Optimized MRM parameters for the determination of morroniside and IS.

Compounds	Q1(Da)	Q3(Da)	DP(V)	CE(eV)	CXP(V)	Detected ion
Morroniside	451.0	243.0	-100	-30	-15	M⁻
Loganin	435.2	227.0	-100	-23	-15	M⁻

Statistical analysis

UPLC-MS/MS data acquisition was performed using Analyst 1.5.2 and MultiQuant

2.1.1 software (Applied Biosystems). All of the data shown are the mean (standard

deviation, n = 3. The data were analyzed by SPSS 20.0 Software. In the Caco-2 cell

model, rate of transport is obtained from amount transported versus time curve using

linear regression. The permeability of a compound is calculated using the following

equation: $P_{app} = (dQ/dt)/(A \cdot C_0)$

where, dQ/dt is the rate of drug transport, C_0 is the initial concentration of the

compound in the donor chamber and A represents thethe surface area of the cell

monolayer. The efflux ratio P was determined by calculating the ratio of P_{app} in the

secretory (BA) direction divided by that in the absorptive (AB) direction, according to

the following equation:

191              $P = P_{app}BA / P_{app}AB$

Authors had access to information that could identify individual participants during or

after data collection.

Results

Cytotoxicity in Caco-2 Cells

Viability of cells was directly measured using MTT test to evaluate the cytotoxicity of

morroniside toward Caco-2 cells prior to transport experiments. As shown in Fig 2,

An increase in the concentrations of morroniside was accompanied by a slight

decrease in the viability of Caco-2 cells, although the difference was not statistically

significant. Even at the highest concentration (200 μM), the cell viability rates of

Caco-2 cells treated with the three compounds reduced only by 8%, indicating that in

our experimental design, morroniside was nontoxic to the growth of Caco-2 cells, as

was 0.63mg/mL CO (containing 25μM morroniside).

Fig 2 Cytotoxicity of morroniside in Caco-2 cells as evaluated by the MTT assay. Data are the mean

values ± standard variation of five replications. Different letters indicate a significant difference at $p <$

206   0.05.

LC/MS/MS analysis to quantify morroniside

The retention time for morroniside and IS were approximately 1.85 and 2.01 min,

respectively. As shown in Fig 3, any significant peaks interfering with morroniside or

IS were not observed in fresh blank HBSS. Representative chromatograms are

presented in Fig 3, including that of morroniside and IS in fresh blank HBSS and a

sample 5 minutes after transport experiments. Calibration curves, constructed using

linear least- squares regression, showed good linearity within a concentration range of

4-1000 ng/mL morroniside in HBSS. A typical calibration curve equation for

morroniside was y=0.007x+0.0055, where y represents the ratio of the morroniside

peak area to the loganin peak area (x is the concentration of morroniside). The

calibration curves displayed good linearity as indicated by high correlation

coefficients (R^2=0.9997). The lower limit of quantification (LLOQ) in the collected

samples was 4 ng/mL.

Fig 3 Representative multiple reaction monitoring (MRM) chromatograms of morroniside (1) and IS

(2) in fresh blank HBSS (A) and a sample (B) 5 minutes after transport experiments by LC/MS/MS

(7.81ng/mL for morroniside and 70ng/mL for IS)

Uptake of morroniside by Caco-2 Cells

Results of the cellular uptakes of different concentrations (5, 25 and 100μM) of

morroniside into Caco-2 cell monolayers from the AP compartment over 2 h were

shown in Fig 4. As shown in Fig 4, the uptakes of morroniside into Caco-2 cells

occurred quickly. $P_{app}AB$ of morroniside ranged from 1.59 at 100μM to $2.66 \times 10^{-}$

^{6}cm/s at 5μM (AP to BL) and from 2.67 at 100μM to 4.10×10^{-6}cm/s at 5μM (BL to

AP)

Fig 4 Time-course of morroniside (5, 25 and 100μM) transport across the Caco-2 cell monolayers from

apical (AP) to basolateral (BL) and from BL to AP (n=3)

Effect of pH and temperature on the permeation of morroniside

The assay studying the AP to BL transport in HBSS with various pH values was

shown in Table 2. The $P_{app}AB$ values of morroniside at pH 7.4 were significantly

higher than those at pH 6.0 (p<0.05), indicating an easier transport of morroniside at a

higher pH (7.4) than at lower a pH (6.0).

Temperature plays an important role in the transport of drugs, it can not only

affect the activity of the carrier protein, but also the state of drugs in solvent(28). As

presented in Table 2, the morroniside transport across Caco-2 cell monolayers at low

temperature (4℃) significantly reduced because $P_{app}AB$ was decreased from $1.94 \times 10^{-}$

^{6}cm/s to 0.6×10^{-6}cm/s.

Table 2 $P_{app}AB$ of morroniside (25 μM) in human intestinal Caco-2cells previously treated at different

243   PH values and temperatures (n = 3, p < 0.05).

Conditions	$P_{app}AB$
4℃	0.60±0.19*
37℃	1.94±0.64
Ph 6.0	1.85±0.60△
Ph 7.4	0.59±0.14

*Denote results significantly different from those of the 37℃ experiments

△Denote results significantly different from those of the Ph 7.4 experiments (p < 0.05).

Effects of Various Compounds on Morroniside Transport

With the participation of 50μM of sodium vanadate or cimetidine, the uptake of

morroniside did not significantly increase or decrease. These results indicated that the

influx transporters like OATs or Na^+/K^+ pump contributed little to the transport of

morroniside. (Table 3)

Table 3Inhibitory effects of influx transporters on morroniside transport in Caco-2 cell monolayers.

Inhibitors	Transporter	$P_{app}AB_{(\times 10^{-6} cm/s)}$	$P_{app}BA_{(\times 10^{-6} cm/s)}$	P
Control		1.90±0.5	3.06±0.6	1.60
Sodium vanadate	Na^+/K^+pump	1.89±0.4	3.11±0.7	0.78
Cimetidine	OATs	1.82±0.5	3.1±0.7	1.23

MRP and BCRP are present in the intestinal tract of humans, which are responsible

for the transport and efflux of compounds or drugs (29). Fig 5 showed the $P_{app}AB$

values increased, but had no effect on the $P_{app}BA$, the data indicated that morroniside

was effluxed by MRP.

BCRP was found to be located at the apical side in the cell membrane (30).

Apigenin was used to determine the effect of BCRP on the transport of 25μM

morroniside. After addition of apigenin (25μM), the $P_{app}AB$ of morroniside increased

significantly (p < 0.05), which resulted in a decrease in the efflux ratio by 72.05%

(Fig 5). It can therefore be assumed that morroniside is also a substrate of BCRP.

Fig 5 Inhibitory effects of exflux transporters on morroniside transport in human intestinal

265   Caco-2 cell monolayers: AP to BL (a) and from BL to AP (b) (n = 3)

*Denote results significantly different from those of the control experiments (p < 0.05).

**Denote results significantly different from those of the control experiments (p < 0.001)

DISCUSSION

To further study the transport characters of morroniside, we investigated the

relationship of time and concentration to transport. The concentration of morroniside

increased approximately linearly with time from 0 to 60 min. After 60 min, the

concentration gradient between the two sides had greatly decreased, result in the

curves turned to reach a platform (Fig 4). A possible reason is that the concentration

gradient between the two sides had greatly decreased after 60 min, which resulted in

much less transport and led to a platform. From the above results, it is concluded that

passive transfer is the main transport way. In our previous work, loganin (31), a

structural analog of morroniside, showed good intestinal permeability by using the

human intestinal Caco-2 cells model. It seemed like the cell permeability of

morroniside is far lower than loganin.

Our results showed that the $P_{app}BA$ values of the morroniside was higher

than their $P_{app}AB$ values ($p < 0.05$), indicating that some transporters might be

involved in the transport of morroniside in B-A direction.

Accounting for the limit of detection of the method, compounds which are

completely absorbed in the human intestine typically exhibit P_{app} values of $>70\times10^{-6}$

cm/s in the Caco-2 transwell system, whereas compounds with poor absorption

($<20\%$) have P_{app} values of $<10\times10^{-6}$ cm/s(32). The P_{app} values of morroniside

determined was at the level of 10^{-6} which indicated poor intestinal absorption about

morroniside. It is also important to note that, strong metabolism may be regarded as

another major cause of the low oral bioavailability of morroniside *in vivo*. However,

further studies are required to clarify whether metabolism was the main factor in the

in vivo process of morroniside.

The data illustrating how pH and temperature effect permeation of

morroniside, show that morroniside transport is both pH-dependent and temperature-

dependent, these results indicate, that some transporters may be involved in the efflux

of morroniside. In fact, previous research indicated that both the influx (33) and efflux

transport like BCRP (34, 35) were activated at lower pH levels. It can therefore be

assumed that morroniside is also a substrate of BCRP. This could be indeed verified

in the next subsequent experiments.

To determine whether P-gp is involved in the transport of these compounds,

their transports were studied in the presence of verapamil, a known inhibitor for P-gp.

But the $P_{app}AB$ and $P_{app}BA$ of morroniside did not change significantly before and

after the addition of verapamil, which indicated that P-gp might not be involved in the

transport of morroniside. (Fig 5)

MRP has different isoforms, from which MRP2 is located at apical side in

the cell membrane and MRP3 was the main basolaterally localized MRP(36).

Indomethacin and benzbromarone were used as MRP2 and MRP3 inhibitors,

respectively. The results in Fig 5 showed that $P_{app}AB$ values caused a significant

increase in the AP to BL flux ($p < 0.001$) after Indomethacin was added. This in turn

indicated that morroniside is a substrate of MRP2. Co-treatment with 200μM

benzbromarone did not alter morroniside's $P_{app}AB$ or the $P_{app}BA$ values, and the

efflux ratio reduced only by 6.64%.

After 0.63mg/mL CO (containing 25μM morroniside) was added, $P_{app}AB$

increased significantly by about 125.26%, and $P_{app}BA$ value was no significant

difference compared with the control group. Many ingredients, especially iridoid

glycosides are contained in CO. To our knowledge, the most abundant bioactive

iridoid glycosides, which are widely distributed in CO, are loganin, morroniside and

sweroside (37). Loganin, which was found to be a substrate of BCRP, in our previous

study, may reduce efflux of morroniside and lead to a lower $P_{app}AB$ in the crude

material group (31). But the compounds contained in CO are complex, and more

research is needed to clarify the intestinal absorption characteristics of this herb

preparation.

Conclusions

In the present study, morroniside was shown to be poor-absorbed compound in the

Caco-2 cell monolayer model and the mechanisms were mainly by passive diffusion

and with efflux protein-mediated active transport. Transporters, BCRP and MRPs

(especially MRP2) are vital for morroniside transport in the intestine. These new

findings provided useful information for predicting its oral bioavailability,

pharmacokinetics, and the clinical application as well as determination of the

bioactive substance basis of morroniside.

Acknowledgments

This work was partially supported by the Cross-fund of Biomedical Engineering of

Shanghai Jiaotong University (No. XH2398) and by Xinhua hospital scientific

research project (No. XH2053).

References

1.   Zhu M, Tang Y, Duan JA, Guo J, Guo S, Su S, et al. Roles of paeoniflorin and

senkyunolide I in SiWu decoction on antiplatelet and anticoagulation activities. J Sep

Sci 2010;33(21):3335-40.

2.   Czerwińska ME, Melzig MF. *Cornus mas* and *Cornus Officinalis*—Analogies

and Differences of Two Medicinal Plants Traditionally Used. Front Pharmacol. 2018;

9 Article 894.

3.   Bhakta HK, Park CH, Yokozawa T, Min BS, Jung HA, Choi JS. Kinetics and

molecular docking studies of loganin, morroniside and 7-O-galloyl-D-sedoheptulose

derived from Corni fructus as cholinesterase and beta-secretase 1 inhibitors. Arch

Pharm Res. 2016;39(6):794-805.

4.   Jeong EJ, Kim TB, Yang H, Kang SY, Kim SY, Sung SH, et al. Neuroprotective

iridoid glycosides from Cornus officinalis fruits against glutamate-induced toxicity in

HT22 hippocampal cells. Phytomedicine. 2012;19(3-4):317-21.

5.   Park CH, Noh JS, Kim JH, Tanaka T, Zhao Q, Matsumoto K, et al. Evaluation of

Morroniside, Iridoid Glycoside from Corni Fructus, on Diabetes-Induced Alterations

such as Oxidative Stress, Inflammation, and Apoptosis in the Liver of Type 2

Diabetic db/db Mice. Biol Pharm Bull. 2011;34(10):1559-65.

6.   Yokozawa T, Yamabe N, Kim HY, Kang KS, Hur JM, Park CH, et al. Protective

Effects of Morroniside Isolated from Corni Fructus against Renal Damage in

Streptozotocin-Induced Diabetic Rats. Biol Pharm Bull. 2008;31(7):1422-8.

7.   Takeda Y, Tanigawa N, Sunghwa F, Ninomiya M, Hagiwara M, Matsushita K, et

al. Morroniside cinnamic acid conjugate as an anti-inflammatory agent. Bioorg Med

Chem Lett. 2010;20(16):4855-7.

8. Oku H, Ogawa Y, Iwaoka E, Ishiguro K. Allergy-Preventive Effects of

Chlorogenic Acid and Iridoid Derivatives from Flower Buds of Lonicera japonica.

Biol Pharm Bull. 2011;34(8):1330-3

9. Li X, Wang Q, Zhang X, Sheng X, Zhou Y, Li M, et al. HPLC study of

pharmacokinetics and tissue distribution of morroniside in rats. J Pharm Biomed

Anal. 2007;45(2):349-55.

10. Yuan GY, Zhang R, Li X, LiW. Simultaneous HPLC-MS Determination of

Loganin, Morroniside and Paeoniflorin in Rat Plasma; Pharmacokinetics of Liuwei

Dihuang Pills. Indian J Pharm Sci. 2019;81(1):129-37.

11. Zhao M, Tao JH, Qian DW, Liu P, Shang EX, Jiang S et al. Simultaneous

determination of loganin, morroniside, catalpol and acteoside in normal and chronic

kidney disease rat plasma by UPLC-MS for investigating the pharmacokinetics of

Rehmannia glutinosa and *Cornus officinalis* Sieb drug pair extract. J Chromatogr B

2016;1009-1010:122-9.

12.   Li B, Lei S, Xiong S, Chen S, Zhang Z. Pharmacokinetics and

Pharmacodynamics of Morroniside: A Review. Nat Prod Commun. DOI:

10.1177/1934578X19856526.

13.   Liu Y, Yang L. Early metabolism evaluation making traditional Chinese

medicine effective and safe therapeutics. J Zhejiang Univ Sci B. 2006;7(2):99-106.

14.   Medina MA, Pérez-Sánchez A, Borrás-Linares I, Barrajón-Catalán E, Arráez-

Román D, González-Álvarez I, et al. Evaluation of the intestinal permeability of

rosemary (Rosmarinus officinalis L.) extract polyphenols and terpenoids in Caco-2

cell monolayers. Plos One. 12(2): e0172063.

15.   Peng Y, Fan M, Peng CS, Wang MY, Li XB. Alleviating the Intestinal

Absorption of Rhein in Rhubarb through Herb Compatibility in Tiaowei Chengqi

Tang in Caco-2 Cells. Evid-Based Compl Alt. DOI: 10.1155/2018/7835128.

16.   Lu J, Liu L, Zhu X, Wu L, Chen Z, Xu Z, et al. Evaluation of the Absorption

Behavior of Main Component Compounds of Salt-Fried Herb Ingredients in Qing'e

Pills by Using Caco-2 Cell Model. Molecules. DOI: 10.3390/molecules23123321.

17. Xu RJ, Peng Y, Wang MY, Li X. Intestinal Absorption of Isoalantolactone and

Alantolactone, Two Sesquiterpene Lactones from Radix Inulae, Using Caco-2 Cells.

Eur J Drug Metab Ph. 2018;44(2):295-303.

18. Liu LJ, Guo LN, Zhao C, Wu XW, Wang RF, Liu C. Characterization of the

Intestinal Absorption of Seven Flavonoids from the Flowers of Trollius chinensis

Using the Caco-2 Cell Monolayer Model. Plos One. 10(3): e0119263.

19. Prasadani WC, Senanayake CM, Jayathilaka N, Ekanayake S, Seneviratne KN.

Effect of three edible oils on the intestinal absorption of caffeic acid: An *in vivo* and

in vitro study. Plos One. 12(6): e0179292.

20. Tao JH, Zhao M, Jiang S, Zhang W, Xu BH, Duan JA. UPLC-Q-TOF/MS-based

metabolic profiling comparison of four major bioactive components in normal and

CKD rat plasma, urine and feces following oral administration of Cornus officinalis

Sieb and Rehmannia glutinosa Libosch herb couple extract. J Pharm Biomed Anal.

2018;161:254-61.

21. Wang Y, Wang H, Wu S, Li D, Chen S. Effect of Gelsemium elegans and

Mussaenda pubescens, the Components of a Detoxification Herbal Formula, on

Disturbance of the Intestinal Absorptions of Indole Alkaloids in Caco-2 Cells. Evid-

Based Compl Alt. DOI: 10.1155/2017/6947948.

22. Zhang B, Zhu XM, Hu JN, Ye H, Luo T, Liu XR, et al. Absorption Mechanism

of Ginsenoside Compound K and Its Butyl and Octyl Ester Prodrugs in Caco-2 Cells.

J Agr Food Chem. 2012;60(41):10278-84.

23. Prime-Chapman HM, Fearn RA, Cooper AE, Moore V, Hirst BH. Differential

multidrug resistance-associated protein 1 through 6 isoform expression and function

in human intestinal epithelial Caco-2 cells. J Pharmacol Exp Ther. 2004;311(2):476-

84.

24. Zhang L, Lin G, Kovacs B, Jani M, Krajcsi P, Zuo Z. Mechanistic study on the

intestinal absorption and disposition of baicalein. Eur J Pharm Sci. 2007;31(3-4):221-

31.

25. Ursic D, Berginc K, Zakelj S, Kristl A. Influence of luminal monosaccharides on

secretion of glutathione conjugates from rat small intestine *in vitro*. Int J Pharm.

2009;381(2):199-204.

26. Zhang SZ, Yang XN, Morris ME. Combined effects of multiple flavonoids on

breast cancer resistance protein (ABCG2)-mediated transport. Pharm Res.

2004;21(7):1263-73.

27. Nait Chabane M, Al Ahmad A, Peluso J, Muller CD, Ubeaud G. Quercetin and

naringenin transport across human intestinal Caco-2 cells. J Pharm Pharmacol.

2009;61(11):1473-83.

28. Wu AG, Zeng B, Huang MQ, Li SM, Chen JN, Lai XP. The absorption and

transport of magnolol in Caco-2 cell model. Chinese Journal of Integrative Medicine.

2012;19(3):206-11.

29. Lev Bromberga VA. Effects of polyether-modified poly(acrylic acid) microgels

on doxorubicin transport in human intestinal epithelial Caco-2 cell layers. J Control

Release. 2003;88:11-22.

30. Xue CF, Guo JM, Qian DW, Duan JA, Shu Y. Absorption of flavonoids from

Abelmoschus manihot extract by in situ intestinal perfusion. Yao Xue Xue Bao.

2011;46(4):454-9.

31. Xu RJ, Yuan YC, Qi J, Zhou J, Guo X, Zhang J, et al. Elucidation of the

Intestinal Absorption Mechanism of Loganin in the Human Intestinal Caco-2 Cell

Model. Evid-Based Compl Alt. DOI: 10.1155/2018/8340563.

32. Willenberg I, Michael M, Wonik J, Bartel LC, Empl MT, Schebb NH.

Investigation of the absorption of resveratrol oligomers in the Caco-2 cellular model

of intestinal absorption. Food Chem. 2015;167:245-50.

33. Matthew D. Troutman1 DRT. Novel Experimental Parameters to Quantify the

Modulation of Absorptive and Secretory Transport of Compounds by P-Glycoprotein

in Cell Culture Models of Intestinal Epithelium. Pharm Res. 2003;20(8):1210-24.

34. Crowe A, Diep S. pH dependent efflux of methamphetamine derivatives and their

reversal through human Caco-2 cell monolayers. Eur J Pharmacol. 2008;592(1-3):7-

12.

35. Citi S. Proein Kinase Inhibitors Prevent Junction Dissociation Induced by Low

Extracellular Calciumin MDCK Epithelial Cells. *J Cell Biol.* 1992;117:169-78.

36. Duan J, Xie Y, Luo H, Li G, Wu T, Zhang T. Transport characteristics of

isorhamnetin across intestinal Caco-2 cell monolayers and the effects of transporters

on it. Food. Chem. Toxicol. 2014;66:313-20.

37. Cai H, Cao G, Cai B. Rapid simultaneous identification and determination of the

multiple compounds in crudeFructus Corniand its processed products by HPLC–

MS/MS with multiple reaction monitoring mode. Pharm Biol. 2012;51(3):273-8.

Fig 1 The chemical structure of morroniside

Fig 2 Cytotoxicity of morroniside in Caco-2 cells as evaluated by the MTT assay. Data are the mean values ± standard variation of five replications. Different letters indicate a significant difference at p < 0.05.

Fig 3 Representative multiple reaction monitoring (MRM) chromatograms of morroniside (1) and IS (2) in fresh

blank HBSS (A) and a sample (B) 5 minutes after transport experiments by LC/MS/MS (7.81ng/mL for

morroniside and 70ng/mL for IS)

Fig 4 Time-course of morroniside (5, 25 and 100μM) transport across the Caco-2 cell monolayers from apical

(AP) to basolateral (BL) and from BL to AP (n=3)

Fig 5 Inhibitory effects of exflux transporters on morroniside transport in human intestinal Caco-2 cell

monolayers: AP to BL (a) and from BL to AP (b) (n = 3)

*Denote results significantly different from those of the control experiments (p < 0.05).

**Denote results significantly different from those of the control experiments (p < 0.001)