Title: Opportunities for prioritizing and expanding conservation enterprise in India using carnivores as flagships

Authors: Arjun Srivathsa[1,2,3], Iravatee Majgaonkar[4], Sushma Sharma[3], Priya Singh[5], Girish Arjun Punjabi[6], Malaika Mathew Chawla[7], Aditya Banerjee[4]

Author affiliations:

School of Natural Resources and Environment, University of Florida, Gainesville, USA

Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, USA

Wildlife Conservation Society–India, Bengaluru, India

Ashoka Trust for Research in Ecology and the Environment, Bengaluru, India

Researchers for Wildlife Conservation, National Centre for Biological Sciences, Bengaluru, India

Wildlife Conservation Trust, Mumbai, India

College of Science and Engineering, James Cook University, Townsville, Australia

Corresponding author:

Arjun Srivathsa

Email: asrivathsa@wcsindia.org

Abstract

Conservation interventions in developing countries are frequently thwarted by socio-economic agendas, severely limiting the scope and rigor of biodiversity and habitat conservation. Very few ecological assessments incorporate human interests in conservation prioritization, creating asynchrony between planning and implementation. For conservation actions to be logistically feasible, multiple criteria including ecological, social, economic and administrative aspects must be considered. Understanding how these different dimensions interact spatially is also important for gauging the potential for conservation success. Here, we use a guild of select mammalian carnivores in India to (i) generate distribution maps at a spatial scale that is relevant to management, (ii) examine ecological, social and biogeographic factors associated with their distribution, quantify key threats, and identify areas critical for their conservation, (iii) use prioritization tools for balancing habitat conservation, human needs and economic growth, and (iv) evaluate the spatial congruence between areas with high conservation potential, and areas currently in focus for protection efforts, conservation investments, and infrastructure development. We find that the current Protected Area system does not adequately cover or represent diverse habitats, that there is immense potential for States to increase financial investments towards alternative conservation strategies, and, most infrastructure projects may be putting important carnivore habitats at risk. Our framework allowed for identifying locations where conservation investments would lead to the highest dividends for flagship carnivores and associated species across habitats. We make a case for re-evaluating how large-scale prioritization assessments are made, and for broadening the purview of conservation policies in India and other developing countries.

Keywords carnivores; conservation investments; economic development; spatial conservation prioritization; species distribution models

Introduction

Conservation is a politically challenging endeavor, particularly in developing economies where

governments have to frequently negotiate a balance between limited conservation funding and

pressing economic agendas (Waldron et al. 2013; Lindsey et al. 2016). In this context, debates on

what to conserve and how, the use of relevant biodiversity surrogates, and trade-offs in

conservation planning are integral parts of the conservation discourse (Westgate et al. 2014;

Lentini & Wintle 2015). The advent of decision-support tools have enabled scientists to

quantitatively determine priority hotspots, prescribe ideal reserve designs, and better plan land-

use management for conservation (Moilanen et al. 2005; Ball et al. 2009; Gregory et al. 2012).

However, these aspects are rarely in tandem with on-ground implementation due to socio-

political, cultural and economic hurdles (Game et al. 2013), seldom meeting requirements of

representativeness and complementarity in species and habitat conservation (sensu Margules and

Pressey 2000). As a consequence, allocation of research efforts, human resources and

conservation funding in many countries do not agree with global priorities and goals for

conservation (Halpern 2006; Jenkins et al. 2013; Brum et al. 2017). Given this background,

conservation prioritization assessments, more often than not, become limited to an academic

exercise.

Integrating federal research and decision-support tools with conservation interventions have been

successful in select countries and landscapes, mostly in the global north (Wintle et al. 2019). The

global south, on the other hand, exemplifies certain asynchronies in priorities, where balancing

human well-being, economic development, and safeguarding nature at the same time, present

formidable challenges (Sandker et al. 2012; Barlow et al. 2018). Many developing countries in

the tropics have witnessed rapid loss of species and habitats, accounting for some of the highest

extinction rates in the world (Pimm et al. 2014; Ceballos et al. 2017). But these tropical countries

are also investing heavily on infrastructure development, relying on capitalist approaches to

economic growth, often at a cost to natural habitats and threatened species (Barlow et al. 2018).

The expanding and growing economy in these countries, together with a globalizing human

society aspiring for better standards of living, continue to sustain this demand. Such scenarios

call for multi-criteria conservation planning assessments that stretch beyond taxonomic,

phylogenetic or ecological considerations (Brum et al. 2017; Sibarani et al. 2019) to additionally

acknowledge, address and incorporate socio-economic, political and administrative factors.

Globally, Protected Area networks are dominated by forested biomes, with 2–6 % of tropical,

subtropical and temperate savanna grasslands and shrublands afforded strict protection (Jenkins

& Joppa 2009). Most published literature and policy perspectives deal with prioritizing global

species-rich hotspots in tropical forests, or Protected Areas and woodland systems in relatively

wealthy, temperate countries (Martin et al. 2012). Furthermore, management approaches that

decouple human-use areas and wilderness areas in pursuit of "intactness" (Mittermeier et al.

2003), tend to bias protection efforts towards landscapes with very low human densities or

access (Joppa and Pfaff 2009; Venter et al. 2014). Such approaches undervalue many ecosystems

and species that do not fit the parochial view of nature and wilderness. For example, savanna

grassland systems, semi-arid scrublands, ravines, deserts and pasturelands may not have very

low human densities, but they support unique assemblages of species adapted to survive in such

ecosystems (e.g., Brito et al. 2013). Even cultivated agricultural lands support many large

mammalian species that persist in transformed food webs and systems (Magioli et al. 2019).

Such multi-use areas rarely figure in global priority locations for conservation (Jenkins et al.

2013), whilst being lost at an alarming rate or rapidly transitioning into unsustainable use

(Dobrovolski et al. 2013). Basic ecological information on key indicator species that represent

the health of these habitats is severely inadequate or inaccurate (Hurlbert & Jetz 2007; Ghosh-

Harihar et al. 2019). This creates an imperative need for identifying and using apposite flagship

species (Verissimo et al. 2013; Shen et al. 2019) or species assemblages as surrogates to

conserve such diverse yet neglected ecosystems.

Terrestrial mammals that belong to the order Carnivora receive among the highest investments in

terms of conservation funds, human resources and public support (Treves & Karanth 2003;

Smith et al. 2012). That some of the most admired species worldover are large wild felids further

testifies their charismatic appeal (Macdonald et al. 2015). Mammalian carnivores represent a

spectrum of habitat associations (Crooks et al. 2011; Bateman & Fleming 2012), with some

showing extremely high specificity (e.g., Ethiopian wolf *Canis simensis* in the highlands of

Ethiopia) and others representing a range of eclectic land cover types (e.g., golden jackal *Canis*

aureus). Certain predators like the leopard *Panthera pardus*, coyote *Canis latrans* and puma

Puma concolor even thrive in radically human-modified habitats, sometimes sharing space with

high human densities (Gehrt & Riley 2010; Vynne et al. 2011; Athreya et al. 2015). Recent

studies have also highlighted the variety of critical ecosystem services and benefits carnivores

provide, all of which contribute towards human well-being through direct or indirect

mechanisms (O'Bryan et al. 2018). Yet, carnivores are highly threatened across the world, with

populations facing massive range contractions and risks of local extinctions, primarily due to

anthropogenic factors (Ripple et al. 2014). Taken together, their natural capital, threatened status,

and the ecological and trophic niches they occupy, makes carnivores potential flagships in their

respective habitats for targeting conservation efforts (Macdonald & Loveridge 2010).

Here, we use India as a case study and focus on nine species/sub-species (henceforth 'species')

of carnivores– dhole *Cuon alpinus*, golden jackal, Indian wolf *Canis lupus pallipes*, Tibetan wolf

Canis lupus chanco, Indian fox *Vulpes bengalensis*, red fox *Vulpes vulpes*, desert fox *Vulpes*

vulpes pusilla, Tibetan fox *Vulpes ferrilata* and striped hyena *Hyaena hyaena* (Fig. 1). We:

(i) generate countrywide distribution maps of the carnivores, at a spatial scale and resolution that

is relevant to administration

(ii) examine ecological, social and biogeographic factors associated with their distribution,

quantify key threats, and identify landscapes and habitats critical for their conservation

(iii) use prioritization tools to make a case for conserving a consortium of habitats with wild

canids and hyenas as flagships, while balancing human needs and economic development

(iv) evaluate the spatial congruence between areas with high conservation potential, and areas

currently in focus for protection efforts, conservation investment, and infrastructure

development.

In doing so, we provide a roadmap for implementing a decentralised approach to long-term

monitoring of carnivores and the habitats they represent, through periodic assessments of their

distribution patterns, threats, human interests and administrative management potential.

Materials and Methods

Carnivore presence records: We collected carnivore distribution data in three phases. In phase 1, we used citizen-science data from countrywide web-based surveys for three months in 2018 (see SI Appendix, section 1 for detailed survey protocol). In addition to our own survey, we also included carefully vetted records from other citizen-science portals (India Biodiversity Portal: http://www.indiabiodiversity.org; iNaturalist: https://www.inaturalist.org). In phase 2, we extracted data from wildlife, nature and photography pages on social media, online wildlife photo-repositories and reliable blog articles in 2018–19 (SI Appendix, section 1). In phase 3, we extracted information from published studies, unpublished theses, forest department reports and openly accessible project reports submitted to funding agencies in 2019. We thoroughly verified and validated each record, ensuring correct species identification, geographic location, and time (month and year) of record. We considered only reliable records of species presence, corresponding to the period from January 2015 to December 2018 (SI Appendix, section 1).

Distribution models: We treated administrative sub-districts ('*tehsil*' or '*taluk*') in the country as independent spatial units or 'sites'(mainland India has 2342 sub-districts). For each species, we first defined a plausible distribution range based on all available information from field guides, State forest department checklists, published literature and our own field-based knowledge. Every presence record was then assigned to an administrative sub-district within the species' plausible range. We used an occupancy modelling framework that accounted for partial detectability to map distribution patterns (MacKenzie et al. 2018). Since our data pertained to presence-only information, we created detection histories using the following method– for each species, a given site within its plausible range was labelled 'D' (detected) if it had at least one detection in the four-year period; months with detections were assigned '1' and months with no detections were assigned '0'. Sites were labelled 'ND' (not detected) when the focal species was not detected across four years, but there was at least one detection of any of the other eight species; all months in these sites were assigned '0'. Sites that did not have detections of any of the nine species during the four-year period were labelled 'NS' (not surveyed; see Powney et al. 2019). For analyses, we collapsed data from 48 months into four 12-month blocks, resulting in one temporal replicate per year. At the sub-district scale (mean = 1400 sq. km; range = 3 – 51,000 sq. km), we assumed that the distribution status of the focal species remained stable

during the four-year period. For each species, we built a set of candidate models with singular

and additive effects of a set of explanatory variables, based on specific *a priori* predictions (SI

Appendix, section 2). We fit occupancy models to detection/non-detection data using package

'unmarked' in program R v3.4.1 (R Core Team 2018).

Explanatory variables for modeling distributions: We used a combination of remotely sensed

data, government-generated figures, and estimates from published studies to compile information

on explanatory variables. These variables were chosen based on their expected influence on one

or more species. We used 12 land-use land cover categories, synthesized and combined from a

total of 152 categories classified by Roy et al. (2015). Climate and topography data were

extracted from remotely sensed satellite imagery. Data on Protected Areas, linear infrastructure

(railways/roadways) and human population densities were obtained from web-based open data

sources. Population data on livestock (sheep, goat and cattle) and free-ranging dogs were sourced

from government livestock census, and data on large wild prey (for dhole and Indian wolf) were

based on published literature. All the variables were re-processed at the sub-district scale for

analyses. Details of these variables, including category, description, and data source and year are

in Table S1.

Threat assessment: Given the paucity of quantitative information on human-induced threats to

wild canids and hyenas in India, we created a conservation score for each species by combining

their protection status, area of occupancy in India, and threat information from surveys of field

experts. For protection status, we included the IUCN Red List status (scores: 1 = CR; 2 = EN; 3

= VU; 4 = NT; 5 = LC), CITES Appendix category (5 = Appendix 1; 3 = Appendix 2; 1 =

Appendix 3), and India's Wildlife Protection Act Schedule (5 = Schedule l; 3 = Schedule 2; 1 =

Schedule 3). Scores for area of occupancy were based on the predicted proportion of habitat that

a species currently occupies within its plausible range in India (scores: 1 to 10; 1 = species

occupies 0–10% of plausible range; 10 = species occupies 90–100% of plausible range). Through

surveys of field experts, we elicited scores for perceived population trend (1 = decreasing; 3 =

stable; 5 = increasing). We also obtained scores for 'level of threat' attributed to habitat loss,

prey decline, direct persecution, road-related mortality, illegal trade and negative interactions

with free-ranging dogs (1 = fatal; 2 = high; 3 = medium; 4 = low; 5 = not a threat). The total

scores for each category were averages from scores of individual experts. The final conservation

score was obtained by summing across all categories, weighting area of occupancy at 0.5,

protection status at 0.3 and expert responses on threats at 0.2. The unequal weighting is because

area of occupancy is a quantitatively estimated metric, protection status is derived from global

datasets but without the same analytical rigor, and threat information is based on expert opinions.

We re-scaled the sum out of 100; a lower conservation score implied a higher threatened status

(within India).

Spatial prioritization: We used software Zonation 4.0 (Moilanen et al. 2014) to identify regions

of high and low conservation values with respect to the focal carnivore species and their habitats

in India. Zonation's algorithm ranks cells (raster pixels) within the area of interest through an

iterative cell-removal process. This is implemented by first assuming that all cells are to be

retained, and then, removing cells that cause the smallest marginal loss in the overall

conservation value (Moilanen et al. 2014). The process is repeated until no more cells are left;

cells with the least value are removed first and those with the highest values are retained towards

the end. The relative weights assigned to the input features and the cell-specific value of each

feature determines the hierarchy of cell removal. The output raster file thus includes pixels

ranked by relative conservation priority values. We chose the 'Core Area Zonation' cell removal

rule because it prioritizes units with the most important and rare features while minimizing loss

in the overall conservation value. We also modified the algorithm such that the final

conservation values are assigned to clusters of pixels grouped by sub-district boundaries. As

primary input features, we used (i) carnivore diversity indices calculated from estimated

occupancy probabilities, and (ii) extent of important habitats in each sub-district (Table S2). We

chose diversity indices rather than individual species distributions after preliminary exploratory

analyses showed better spatial representation of important areas and greater area of retention

with the former. Diversity indices used as input features are shown in Fig. S1. In addition to

these, we also included human poverty index and projected human population for 2020, both of

which qualified as "costs" in our assessment. Our rationale was that sub-districts with higher

human populations and higher poverty require focus on economic growth and infrastructure

development and should therefore be of lower priority for conservation (see Table S2 for

details). Following exploratory runs with different combinations of settings for Boundary Length

Penalty (BLP) and Warp Factor (WP), we set BLP at 0 and WP at 100 as a trade-off between

computation time and reliability of spatial maps.

Finally, we compared locations of high conservation priority and (i) their spatial congruence with

current Protected Areas, (ii) State-wise conservation likelihood based on current budgetary

spending in forest and wildlife sectors, (iii) their overlap with locations of current/planned

infrastructure projects such as solar farms, large- and small-hydropower projects, large mines,

and irrigation projects– all of which are likely to affect the focal species and their habitats.

Results

We collated a total of 4437 presence records of the target species across three phases (Table 1).

Model-averaged occupancy estimates ranged from 0.21 (SE 0.02) for desert fox to 0.75 (SE

0.002) for golden jackal (Fig. 2). We could not formally analyze or generate estimates for

Tibetan fox because the data were too sparse. Tibetan wolf occupied the least overall area

(152,180 sq. km) and golden jackal was the most widely distributed species (2,259,361 sq. km).

Combining protection status, distribution extent, population status and anthropogenic threats (the

latter two elicited from surveys of field experts in India; n = 45), golden jackal and red fox

scored the highest, while Indian fox and striped hyena scored the least (Table 1).

Carnivore–habitat associations generally agreed with our predictions. Forest cover and large wild

prey were important for dhole and Indian wolf (Table 2). Extent of grasslands, scrublands,

ravines and open (barren) habitats were positively associated with all fox species and striped

hyena. Production agroforests were preferred by dhole and jackal, and striped hyenas preferred

rocky outcrops. Considered together, grasslands, scrublands, open/ravine habitats and forests

were key habitats for wild canids and hyenas. Dhole occupancy was negatively associated with

cattle and human densities, and positively with extent of Protected Areas– underscoring their

importance for dhole source populations. The magnitudes and directions of effects with respect

to human settlements, linear infrastructure and free-ranging dog densities varied across species

(Table 2). Interestingly, high probabilities of golden jackal occurrence were clustered around

large settlements with high densities of humans and free-ranging dogs. This may be attributed to

high availability of provisioned food resources, and their ability to adapt to human-modified
areas. While these results were based on singular covariate effects, the final occupancy estimates
for all species were derived from averaging across models with comparable statistical support
(based on AICc scores and weights; SI Appendix section 2, Table S3). We generated both,
distribution maps at the sub-district level (Fig. 2), and maps depicting occurrence probabilities
within plausible habitats (Fig. S2).

Based on results from prioritization analysis, we chose sub-districts that accounted for the top
30% priority scores (n = 703) as the most important areas for conserving wild canids, hyenas,
and their habitats (Fig. 3). The high priority areas– which we term 'Canid Conservation Units'
(CCUs)– cover 1,475,558 sq. km, of which 545,070 sq. km (37%) constitute focal habitats (Fig.
3). We further classified these areas into tier 1, tier 2 and tier 3, based on level of priority (top 10,
20, or 30% scores). Protected Areas currently cover around 26% of key habitats within CCUs
(Fig. 3). Summed conservation ranks of CCUs showed that 12 States fared relatively better than
India's 17 other States (Fig. 4). Of these, Madhya Pradesh, Karnataka, Odisha and Chhattisgarh
have the most conservation potential and likelihood of success, when the States' budgetary
allocation for forests and wildlife sectors is taken into account. Rajasthan, Gujarat, Andhra
Pradesh, Maharashtra, West Bengal, Bihar, Tamil Nadu and Uttar Pradesh have high
conservation scores, but will need to invest higher monetary resources to be able to work towards
conserving CCUs in their States (Fig. 4).

We extracted information on numbers and spatial locations of infrastructure projects across the
country from multiple online databases of federal and State government ministries. A total of
1143 projects were implemented or commissioned in the last five years (roughly coinciding with
the period considered for carnivore distribution assessment). We could correctly assign spatial
locations to 1073 projects: 413 solar farms, 214 irrigation projects, 20 large mines, 205 large-
and 221 small-hydropower projects. Of these, 41% overlapped with CCUs (Fig. 5), suggestive of
high risks to carnivores and their habitats. Only 18% of the projects were in low conservation
priority areas (sub-districts with bottom 30% priority scores), i.e., locations that ranked high in
terms of projected human populations and poverty index.

Discussion

Given the oft-overlooked caveats with global-scale assessments, such as data inaccuracy or inadequacy and broad generalization of inferences (Ocampo-Peñuela et al. 2016; Norman & White 2019), our study demonstrates the utility of carefully designed large-scale investigations integrated with local knowledge and contexts. We present more nuanced distribution maps depicting species occurrences as probabilities within their Area of Habitat (Brooks et al. 2019; Fig. S2), rather than Minimum Convex Polygons that over- or under-estimate species ranges (Ocampo-Peñuela et al. 2016; Ramesh et al. 2017). Our approach also rectifies errors in current IUCN distribution maps; spatial mismatch between the two ranged from 167,897 sq. km for wolf (*Canis lupus pallipes* and *Canis lupus chanco*, combined) to 2,123,324 sq. km for red fox (*Vulpes vulpes* and *Vulpes vulpes pusilla*, combined). These findings are extremely relevant to past and future studies that use IUCN range maps for large-scale conservation assessments, reviews or syntheses. The focal species together represented multiple important habitats hitherto neglected in terms of conservation and management. Our estimates for Tibetan wolf and desert fox had lower reliability due to gaps in data (SI Appendix section 2), which we hope to redress in future surveys.

About 5% of India's 3.29 million sq. km land area is protected as National Parks and Wildlife Sanctuaries. Most of these are in forest habitats, owing to the country's history of forest-centric protection regimes that stem from a British-colonial legacy of caching forestlands for timber extraction (see Ghosh-Harihar et al. 2019). Consequently, non-forest habitats that were either non-arable or unfit for timber production were termed "wastelands" (Singh et al. 2006). Decades of sidelining these habitats from mainstream conservation discourse has rendered them fragmented, isolated and degraded in most parts of the country (Vanak et al. 2015; Mondal et al. 2019). The systematic marginalization, compounded by other factors, resulted in the near-extinction of grassland-dependent species like the Great Indian Bustard *Ardeotis nigriceps*, Jerdon's courser *Rhinoptilus bitorquatus*, and perhaps many others (Rahmani 2012). We foresee immense potential for expanding the conservation enterprise in India through demarcation of Conservation Reserves and Community Reserves, which currently cover *c.* 5,000 sq. km. Employing mechanisms such as creation of Village Reserves with equitable regulation can protect species and habitats without restricting sustainable use of resources in human-dominated

landscapes. Our study also identifies locations where other logistically, economically, and

socially feasible management options could be explored. If prioritized and carefully protected,

the CCUs could potentially safeguard around 23,830 sq. km of grasslands, 99,950 sq. km of

scrublands, 124,450 sq. km of open habitats or ravines, and 232,430 sq. km of forests– together

with the current Protected Area network. This, of course, should not dissuade practitioners from

prioritizing individual species or habitats in areas beyond CCUs for targeting conservation

efforts.

Supporting the world's second largest human population, India continues to witness rapid land-

use changes, with land conversions being relatively easier for grasslands, scrublands and even

agricultural lands (Baka 2014; Vanak et al. 2015; Kennedy et al. 2019). But ecological

considerations rarely figure in current policies of landscape management and land-use planning.

Instead, flawed notions of 'compensating for damage' through afforestation initiatives– when

forests are cleared for infrastructure projects– are implemented in grasslands and scrublands

(Vanak et al. 2015). Similar afforestation schemes globally promulgated for carbon sequestration

and combating climate change (Bond et al. 2019) and exploiting these beleaguered habitats for

industrial-scale solar farms, further increase their fragility while also bearing negative

consequences for local communities (Yenneti et al. 2016; this study). The glaring knowledge

gaps about these habitats impede our ability to comprehend their importance or plan their

management (Ratnam et al. 2016). For example, our results did not show a significant effect of

agricultural lands on carnivore occupancy. But we posit that it may be a scale-dependent effect;

at finer spatial scales, agricultural areas likely serve as secondary habitats, offer temporary

refuges, and facilitate connectivity for these carnivores (Vanak & Gompper 2010; Athreya et al.

2015; Majgaonkar et al. 2019). Changes in crop type, transitions from seasonal to year-round

cultivation, expansion of permanent irrigation, and intensification of pesticide use (impacting

lower trophic levels) could create altered landscapes (UNCCD 2017), leading to local extinctions

and/or shifts in carnivore communities (e.g., Majgaonkar et al. 2019). Future studies will need to

undertake detailed investigations on these aspects to offer deeper ecological insights and

augment our findings.

Carnivores have long held the fascination of wildlife managers, governments and the civil

society (Treves & Karanth 2003; Smith et al. 2012). But they also present complicated

conservation paradigms because of their negative interactions with people (Ripple et al. 2014;

Montgomery et al. 2018); although, wild canids and hyenas in India are generally not associated

with major threats to human life or property (Srivathsa et al. 2019). At present, wildlife

monitoring in India predominantly involves national estimation of tiger–prey, Asiatic lion, Asian

elephant and one-horned rhinoceros populations. No protocol or framework exists for monitoring

other species groups like wild canids, small carnivores, amphibians, riverine fauna, or ungulate

herbivores in high-altitude or non-woodland habitats– a trend that appears to be commonplace

for many species and habitats in developing countries (see Dornelas et al. 2014). Bridging some

of these gaps, our work adds to a growing body of studies that leverage emerging opportunities

in citizen science, social media and open source data for conservation monitoring (Newman et al.

2012; Ramesh et al. 2017; Toivonen et al. 2019).

Our focal species align themselves mid-way between obscurity and ubiquity, i.e., they are neither

so rare that citizens are unaware of their existence, nor are they so abundant that calling for their

conservation seems futile. And since these carnivores also occupy mid–high trophic levels, we

believe they can serve as ideal flagships across important habitats. Our aim is to build a

monitoring system with real-time documentation of distribution and threat data on these

carnivores, coupled with periodic analyses that would incrementally update the findings

presented here. We anticipate the application of our approach to other poorly-studied taxa for

which intensive resource allocation is not possible. This would complement government-driven

monitoring programs with decentralized local efforts, fostering citizens' involvement in

scientific research while also garnering public support for conservation (Newman et al. 2012).

Conclusion

The window of opportunity for implementing conservation initiatives in developing economies is

becoming narrow. For spatial prioritization results to be relevant for local and federal

governments, decision-support tools should incorporate human needs, politics and economic

interests (Ban et al. 2013; Dickman et al. 2015). While our study does address these aspects, we

recognize greater scope for refining and increasing the rigor of this exercise as we move forward.

For instance, examining habitat connectivity and social thresholds of human–carnivore

coexistence, quantifying region-specific threats and species-wise infrastructure impacts, and

tracking shifts in human populations towards urban centres and away from forest-dependent or

agrarian livelihoods, would all contribute towards adaptively building upon the results presented

here. Nonetheless, we strongly argue for broadening current management notions of wilderness

and natural habitats in India and recommend exploring and implementing alternative

conservation approaches beyond forested Protected Areas. Certainly, executing these approaches

would entail choosing habitat-specific flagships relevant to ecological, regional and social

contexts, and developing monitoring techniques that can be co-opted by conservation managers

and planners. In sum, our study represents an 'ensemble approach', combining a range of data

sources, methods and analytical frameworks, and incorporates human population, poverty,

infrastructure projects, and administrative potential and likelihood in conservation prioritization.

We believe these are crucial for critically evaluating large-scale prioritization assessments, and

for rethinking country-level conservation policies in India and other developing economies.

Acknowledgements

We express gratitude to all the people who proactively participated and contributed data as part

of our citizen-science surveys. We are also grateful to all the field experts in India who

contributed to the threat assessment surveys. We thank Wildlife Conservation Society–India,

IUCN Canid Specialist Group, Nature Conservation Foundation, Nature in Focus, Conservation

India, IUCN Hyaena Specialist Group, R. Sreenivasan, V. Athreya and R. Chakravarty for their

support. A.S was supported by the University of Florida, Wildlife Conservation Society's

Christensen Conservation Leaders Scholarship and Wildlife Conservation Network's Sidney

Byers Fellowship. G.A.P was supported by IDEA Wild. We thank M.A. Agnishikhe, A. Das, T.

Kothawalla, D. Ganguly, K. Chauhan and A. Simon for their enthusiastic assistance in data

compilation and processing. The authors received no funding for this study.

Author contributions

A.S., I.M., S.S., P.S and G.A.P. designed the research; all authors performed the research and

analyzed the data; A.S., I.M., P.S. and G.A.P. wrote the paper.

References

Athreya, V., Srivathsa, A., Puri, M., Karanth, K. K., Kumar, N. S., and Karanth, K. U. (2015).

Spotted in the news: using media reports to examine leopard distribution, depredation, and

management practices outside protected areas in Southern India. *PLoS One*, 10(11), e0142647.

Baka, J. (2014). What wastelands? A critique of biofuel policy discourse in South India.

Geoforum, 54, 315-323.

Ball, I. R., Possingham, H. P., and Watts, M. (2009). Marxan and relatives: software for spatial

conservation prioritisation. *Spatial conservation prioritisation: Quantitative methods and*

*computational tools,*185-195.

Ban, N.C., Mills, M., Tam, J., Hicks, C.C., Klain, S., Stoeckl, N., Bottrill, M.C., Levine, J.,

Pressey, R.L., Satterfield, T. and Chan, K.M. (2013). A social–ecological approach to

conservation planning: embedding social considerations. *Front. Ecol. Environ.*, 11(4), 194-202.

Barlow, J., Franca, F., Gardner, T. A., Hicks, C. C., Lennox, G. D., Berenguer, E., Castello, L.,

Economo, E. P., Ferreira, J., Guenard. B., et al. (2018). The future of hyperdiverse tropical

ecosystems. *Nature*, 559 (7715), 517–526.

Bateman, P. W., and Fleming, P. A. (2012). Big city life: carnivores in urban environments. *J.*

Zool., 287(1), 1-23.

Bond, W. J., Stevens, N., Midgley, G. F., and Lehmann, C. E. (2019). The Trouble with Trees:

Afforestation plans for Africa. Trends in ecology and evolution. doi:

https://doi.org/10.1016/j.tree.2019.08.003

Brito, J.C., Godinho, R., Martínez Freiría, F., Pleguezuelos, J.M., Rebelo, H., Santos, X., Vale,

C.G., Velo Antón, G., Boratyński, Z., Carvalho, S.B., et al. (2013). Unravelling biodiversity,

evolution and threats to conservation in the Sahara Sahel. *Biol. Rev.*, 89(1), 215-231.

Brooks, T.M., Pimm, S.L., Akçakaya, H.R., Buchanan, G.M., Butchart, S.H., Foden, W., Hilton-

Taylor, C., Hoffmann, M., Jenkins, C.N., Joppa, L. and Li, B.V. (2019). Measuring terrestrial

area of habitat (AOH) and its utility for the IUCN red list. *Trends Ecol. Evol*. doi:

https://doi.org/10.1016/j.tree.2019.06.009

Brum, F. T., Graham, C. H., Costa, G. C., Hedges, S. B., Penone, C., Radeloff, V. C., Rondinini.

C., Loyola, R., and Davidson, A. D. (2017). Global priorities for conservation across multiple

dimensions of mammalian diversity. *P. Natl. A. Sci.,* 114(29), 7641-7646.

Ceballos, G., Ehrlich, P. R., and Dirzo, R. (2017). Biological annihilation via the ongoing sixth

mass extinction signalled by vertebrate population losses and declines. *P. Natl. A. Sci.,* 114(30),

E6089-E6096.

Crooks, K. R., Burdett, C. L., Theobald, D. M., Rondinini, C., and Boitani, L. (2011). Global

patterns of fragmentation and connectivity of mammalian carnivore habitat. *Philos. T. R. Soc. B.*,

366(1578), 2642-2651.

Dornelas, M., Gotelli, N. J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., and Magurran, A.

E. (2014). Assemblage time series reveal biodiversity change but not systematic loss. *Science*,

344(6181), 296-299.

Dickman, A. J., Hinks, A. E., Macdonald, E. A., Burnham, D., and Macdonald, D. W. (2015).

Priorities for global felid conservation. *Conserv. Biol.*, 29(3), 854-864.

Dobrovolski, R., Loyola, R. D., Guilhaumon, F., Gouveia, S. F., and Diniz-Filho, J. A. F. (2013).

Global agricultural expansion and carnivore conservation biogeography. *Biol. Conserv.,* 165,

162-170.

Game, E. T., Kareiva, P., and Possingham, H. P. (2013). Six common mistakes in conservation

priority setting. *Conserv. Biol.,* 27(3), 480-485.

Gehrt, S. D., and Riley, S. P. (2010). Coyotes (*Canis latrans*). *Urban carnivores: Ecology,*

*conflict, and conservation,*79-95.

Ghosh-Harihar, M., An, R., Athreya, R., Borthakur, U., Chanchani, P., Chetry, D., Datta, A.,

Harihar, A., Karanth, K.K., Mariyam, D. and Mohan, D. (2019). Protected areas and biodiversity

conservation in India. *Biol. Conserv.,* 237, 114-124.

Gregory, R., Failing, L., Harstone, M., Long, G., McDaniels, T., and Ohlson, D.

(2012). *Structured Decision Making: A Practical Guide to Environmental Management*

Choices. (pp. 1–299). Wiley.

Halpern, B. S., Pyke, C. R., Fox, H. E., Chris Haney, J., Schlaepfer, M. A., and Zaradic, P.

(2006). Gaps and mismatches between global conservation priorities and spending. *Conserv.*

Biol., 20(1), 56-64.

Hurlbert, A. H., and Jetz, W. (2007). Species richness, hotspots, and the scale dependence of

range maps in ecology and conservation. *P. Natl. A. Sci.,* 104(33), 13384-13389.

Jenkins, C. N., and Joppa, L. (2009). Expansion of the global terrestrial protected area system.

Biol. Conserv., 142(10), 2166-2174.

Jenkins, C. N., Pimm, S. L., and Joppa, L. N. (2013). Global patterns of terrestrial vertebrate

diversity and conservation. *P. Natl. A. Sci.,* 110(28), E2602-E2610.

Joppa, L. N., and Pfaff, A. (2009). High and far: biases in the location of protected areas. *PloS*

one, 4(12), e8273.

Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch Mordo, S., and Kiesecker, J. (2019).

Managing the middle: A shift in conservation priorities based on the global human modification

gradient. *Global Change Biology*, 25(3), 811-826.

Lentini, P. E., and Wintle, B. A. (2015). Spatial conservation priorities are highly sensitive to

choice of biodiversity surrogates and species distribution model type. *Ecography*, 38(11), 1101-

1111.

Lindsey, P.A., Balme, G.A., Funston, P.J., Henschel, P.H., and Hunter, L.T.B. (2016). Life after

Cecil: channeling global outrage into funding for conservation in Africa. *Conserv. Lett.*, 9(4),

296-301.

Macdonald, D., and Loveridge, A. (2010). *The biology and conservation of wild felids (Vol. 2)*.

Oxford University Press, U.K.

Macdonald, E. A., Burnham, D., Hinks, A. E., Dickman, A. J., Malhi, Y., and Macdonald, D. W.

(2015). Conservation inequality and the charismatic cat*: Felis felicis*. *Global Ecol. Conserv.,* 3,

851-866.

MacKenzie, D. I., Nichols, J. D., Royle, J. A., Pollock, K. H., Bailey, L., and Hines, J. E. (2018).

Occupancy estimation and modeling: inferring patterns and dynamics of species occurrence.

Elsevier.

Magioli, M., Moreira, M. Z., Fonseca, R. C. B., Ribeiro, M. C., Rodrigues, M. G., and de Barros,

K. M. P. M. (2019). Human-modified landscapes alter mammal resource and habitat use and

trophic structure. *P. Natl. A. Sci.,* 116(37), 18466-18472.

Majgaonkar, I., Vaidyanathan, S., Srivathsa, A., Shivakumar, S., Limaye, S., and Athreya, V.

(2019). Land sharing potential of large carnivores in human modified landscapes of western

India. *Conserv. Sci. Prac.*, 1(5), e34.

Margules, C. R., and Pressey, R. L. (2000). Systematic conservation planning. *Nature*,

405(6783), 243.

Martin, L. J., Blossey, B., and Ellis, E. (2012). Mapping where ecologists work: biases in the

global distribution of terrestrial ecological observations. *Front. Ecol. Environ.*, 10(4), 195-201.

Mittermeier, R. A., Mittermeier, C. G., Brooks, T. M., Pilgrim, J. D., Konstant, W. R., Da

Fonseca, G. A., and Kormos, C. (2003). Wilderness and biodiversity conservation. *P. Natl. A.*

Sci., 100(18), 10309-10313.

Moilanen, A., Franco, A. M., Early, R. I., Fox, R., Wintle, B., and Thomas, C. D. (2005).

Prioritizing multiple-use landscapes for conservation: methods for large multi-species planning

problems. *P. R. Soc. B.,* 272(1575), 1885-1891.

Moilanen, A., Pouzols, F. M., Meller, L., Veach, V., Arponen, A., Leppänen, J., and Kujala, H.

(2014). *Zonation–Spatial Conservation Planning Methods and Software. Version 4. User*

Manual. C-BIG Conservation Biology Informatics Group Department of Biosciences. University

of Helsinki, Finland

Mondal, T., Qureshi, S. Q., Jhala, Y. V., and Basu, P. (2019). An assessment of land use land

cover change in Central Highland of Deccan Peninsula and Semi-Arid tracts of India.

*BioRxiv,*665794.

Montgomery, R.A., Elliott, K.C., Hayward, M.W., Gray, S.M., Millspaugh, J.J., Riley, S.J.,

Kissui, B.M., Kramer, D.B., Moll, R.J., Mudumba, T. and Tans, E.D. (2018). Examining evident

interdisciplinarity among prides of lion researchers. *Front. Ecol. Evol.,* 6, 49.

Newman, G., Wiggins, A., Crall, A., Graham, E., Newman, S., and Crowston, K. (2012). The

future of citizen science: emerging technologies and shifting paradigms. *Front. Ecol. Environ.,*

10(6), 298-304.

Norman, K. E., and White, E. P. (2019). Hotspot prioritizations show sensitivity to data type.
bioRxiv, 685735.

O'Bryan, C. J., Braczkowski, A. R., Beyer, H. L., Carter, N. H., Watson, J. E., and McDonald-
Madden, E. (2018). The contribution of predators and scavengers to human well-being. *Nat.*
Ecol. Evol., 2(2), 229.

Ocampo-peñuela, N., Jenkins, C. N., Vijay, V., Li, B. V, and Pimm, S. L. (2016). Incorporating
explicit geospatial data shows more species at risk of extinction than the current Red List. *Sci.*
Adv., 2(11), 1–10 (e1601367).

Pimm, S.L., Jenkins, C.N., Abell, R., Brooks, T.M., Gittleman, J.L., Joppa, L.N., Raven, P.H.,
Roberts, C.M. and Sexton, J.O. (2014). The biodiversity of species and their rates of extinction,
distribution, and protection. *Science,* 344(6187), p.1246752.

Powney, G. D., Carvell, C., Edwards, M., Morris, R. K., Roy, H. E., Woodcock, B. A., and Isaac,
N. J. (2019). Widespread losses of pollinating insects in Britain. *Nature comm.,* 10(1), 1018.

Rahmani, A. R., Kalra, M., and Khan, N. I. (2012). *Threatened birds of India: their conservation*
requirements. Oxford University Press.

Ramesh, V., Gopalakrishna, T., Barve, S., and Melnick, D. J. (2017). IUCN greatly
underestimates threat levels of endemic birds in the Western Ghats. *Biol. Conserv.*, 210, 205-
221.

Ratnam, J., Tomlinson, K. W., Rasquinha, D. N., and Sankaran, M. (2016). Savannahs of Asia:
antiquity, biogeography, and an uncertain future. *Philos. T. R. Soc. B.*, 371(1703), 20150305.

Ripple, W.J., Estes, J.A., Beschta, R.L., Wilmers, C.C., Ritchie, E.G., Hebblewhite, M., Berger,
J., Elmhagen, B., Letnic, M., Nelson, M.P. and Schmitz, O.J. (2014). Status and ecological
effects of the world's largest carnivores. *Science,* 343(6167), 1241484.

Roy, P.S., Behera, M.D. Murthy, M.S.R. Roy, A. Singh, S., Kushwaha, S.P.S., Jha, C.S.

Sudhakar, S., Joshi, P.K., Reddy, S., et al. (2015). New vegetation type map of India prepared

using satellite remote sensing: Comparison with global vegetation maps and utilities. *Int. J Appl.*

Earth Obs., 39, 142-159.

Sandker, M., Ruiz-perez, M., and Campbell, B. M. (2012). Trade-Offs Between Biodiversity

Conservation and Economic Development in Five Tropical Forest Landscapes. *Environ.*

Manage., 50, 633–644.

Shen, X., Li, S., McShea, W.J., Wang, D., Yu, J., Shi, X., Dong, W., Mi, X. and Ma, K.,

Effectiveness of management zoning designed for flagship species in protecting sympatric

species. (2019). *Conserv. Biol.* https://doi.org/10.1111/cobi.13345

Sibarani, M. C., Di Marco, M., Rondinini, C., and Kark, S. (2019). Measuring the surrogacy

potential of charismatic megafauna species across taxonomic, phylogenetic and functional

diversity on a megadiverse island. *J. Appl. Ecol.*, 56(5), 1220-1231.

Singh, P., Rahmani, A. R., Wangchuk, S., Mishra, C., Singh, K. D., and Narain, P. (2006).

Report of the task force on grasslands and deserts. *Planning Commission, Government of India,*

New Delhi.

Smith, R. J., Verissimo, D., Isaac, N. J. B., and Jones, K. E. (2012). Identifying Cinderella

species: uncovering mammals with conservation flagship appeal. *Conserv. Lett.*, 5, 205–212.

Srivathsa, A., Puri, M., Karanth, K. K., Patel, I., and Kumar, N. S. (2019). Examining human–

carnivore interactions using a socio-ecological framework: sympatric wild canids in India as a

case study. *Roy. Soc. Open Sci.,* 6(5), 182008.

Treves, A., and Karanth, K. U. (2003). Human   carnivore conflict and perspectives on carnivore

management worldwide. *Conserv. Biol.,* 17(6), 1491-1499.

Toivonen, T., Heikinheimo, V., Fink, C., Hausmann, A., Hiippala, T., Järv, O., Tenkanen, H. and

Di Minin, E., (2019). Social media data for conservation science: a methodological overview.

Biol. Conserv., 233, 298-315.

United Nations Convention to Combat Desertification. (2017). The Global Land Outlook, first

edition. Bonn, Germany.

Vanak, A. T., and Gompper, M. E. (2010). Multi scale resource selection and spatial ecology of

the Indian fox in a human dominated dry grassland ecosystem. *J. Zool*., 281(2), 140-148.

Vanak, A. T., Kulkarni, A., Gode, A., Sheth, C., and Krishnaswamy, J. (2015). Extent and Status

of Semiarid Savanna Grasslands in Peninsular India. In *Ecology and Management of Grassland*

Habitats in India. ENVIS Bulletin, 17. ENVIS Centre of Wildlife Institute of India, Dehradun.

Venter, O., Fuller, R.A., Segan, D.B., Carwardine, J., Brooks, T., Butchart, S.H., Di Marco, M.,

Iwamura, T., Joseph, L., O'Grady, D. and Possingham, H.P. (2014). Targeting global protected

area expansion for imperiled biodiversity. *PLoS Biol.*, 12(6), e1001891.

Veríssimo, D., Fraser, I., Girão, W., Campos, A. A., Smith, R. J., and MacMillan, D. C. (2013).

Evaluating conservation flagships and flagship fleets. *Conserv. Lett.*, 7(3), 263-270.

Vynne, C., Keim, J. L., Machado, R. B., Marinho-Filho, J., Silveira, L., Groom, M. J., and

Wasser, S. K. (2011). Resource selection and its implications for wide-ranging mammals of the

Brazilian Cerrado. *PloS one*, 6(12), e28939.

Waldron, A., Mooers, A.O., Miller, D.C., Nibbelink, N., Redding, D., Kuhn, T.S., Roberts, J.T.,

and Gittleman, J.L. 2013. Targeting global conservation funding to limit immediate biodiversity

declines. *P. Natl. A. Sci.* 110(29): 12144-12148.

Westgate, M. J., Barton, P. S., Lane, P. W., and Lindenmayer, D. B. (2014). Global meta-
analysis reveals low consistency of biodiversity congruence relationships. *Nature commun.*, 5,
3899.
680
681   Wintle, B. A., Kujala, H., Whitehead, A., Cameron, A., Veloz, S., Kukkala, A., Moilanen, A.,
682   Gordon, A., Lentini, P.E., Cadenhead, N.C. and Bekessy, S. A. (2019). Global synthesis of
683   conservation studies reveals the importance of small habitat patches for biodiversity. *P. Natl. A.*
684   *Sci.,* 116(3), 909-914.
685
686   Yenneti, K., Day, R., and Golubchikov, O. (2016). Spatial justice and the land politics of
687   renewables: Dispossessing vulnerable communities through solar energy mega-projects.
688   *Geoforum*, 76, 90-99.

Tables

Table 1. Summary of data records collated from three survey phases, estimated occupancy

(standard errors in parentheses), extent of occurrence in India and conservation score for the

focal carnivore species

Table 2. Estimated slope coefficients from singular covariate models for each of the carnivore

species. Abbreviations: fcov– combined forest cover; dfor– tropical dry forests; tfor– temperate

forests; gsor– grasslands, scrublands, open (barren) habitats and ravines; ohab– open (barren)

habitats; agri– agricultural areas; prod– production agroforests; hamt– high altitude mountains;

habt– combined area of plausible habitats (species-specific); rock– rocky outcrops and

escarpments; pptn– annual precipitation; rugg– terrain ruggedness; prey– wild prey index; elev–

elevation; catl– cattle population; shot– sheep and goat population; lstk– livestock (cattle, sheep

and goat) population; hpop– human population; dogs– free-ranging dog population; sett– human

settlements; ifra– density of roads and railways; resv– extent of Protected Areas

Figure captions

Figure 1. The nine focal carnivore species/sub-species in this study. From left to right, top row to bottom row (photo credits in parentheses): dhole (Sushvin Gowda), golden jackal (Santosh Doddagoudar), Indian wolf (Anup Deodhar), Tibetan wolf (Aditya Chavan), Indian fox (Anuroop Krishnan), red fox (Ismail Shariff), desert fox (Manish Vaidya), Tibetan fox (Aditya Chavan) and striped hyena (Kalyan Varma).

Figure 2. Estimated sub-district level occupancy probabilities for the focal carnivore species across India. Range of estimates: dhole (0.03–0.96), golden jackal (0.51–0.99), Indian wolf (0.02–0.55), Tibetan wolf (0.20–0.65), Indian fox (0.11–0.93), red fox (0.04–0.99), desert fox (0.02–0.99) and striped hyena (0.26–0.66). Model averaged estimates and associated standard errors are in Table 1.

Figure 3. Spatial maps of results from prioritization analysis. (a) Sub-district level conservation priority scores for India. (b) Sub-districts with top 30% priority scores, classified as tiers 1– top 10% scores, 2– top 20%, and 3– top 30%, collectively termed Canid Conservation Units 'CCUs'. (c) spatial overlap between Protected Areas of India and CCUs identified in this study.

Figure 4. Summed CCU scores for each State in India, against the State's average budgetary spending in wildlife and forest sectors (expressed as a proportion of annual State budget 2015– 2018). Vertical black lines on both axes are corresponding median values. Dots in black represent States with high CCU scores and higher budget allocation; yellow– high CCU scores, lower budget; grey– low CCU scores, low budget; blue– low CCU scores, higher budget. States with dashed lines do not have any sub-districts in the top 30% priority scores.

Figure 5. Infrastructure projects in key habitats. Left panel– Sub-district-wise numbers of infrastructure projects (irrigation projects, large mines, solar farms, large- and small-hydropower projects) implemented or commissioned in the past five years in India. Right panel– Sub-districts where infrastructure projects overlap with CCUs; dashed areas represent other CCUs with no projects or no available data on the key projects.

Table 1. Summary of data records collated from three survey phases, estimated occupancy (standard errors in parentheses), extent of occurrence in India and conservation score for the focal carnivore species

Species	Phase 1 records	Phase 2 records	Phase 3 records	Estimated occupancy (SE)	Area occupied (sq. km)	Conservation score (scaled to 100)
dhole	290	417	82	0.46 (0.01)	249,606	59
golden jackal	1054	377	86	0.75 (0.002)	2,259,361	69
Indian wolf	331	177	8	0.33 (0.003)	293,947	60
Tibetan wolf	92	60	1	0.36 (0.01)	152,180	66
Indian fox	418	213	1	0.34 (0.001)	890,831	49
red fox	58	54	95	0.50 (0.02)	274,292	69
desert fox	28	115	0	0.21 (0.02)	174,580	55
Tibetan fox	3	10	0	–	–	–
striped hyena	226	185	56	0.41 (0.001)	957,478	49

Table 2. Estimated slope coefficients from singular covariate models for each of the carnivore species. Abbreviations: fcov– combined forest cover; dfor– tropical dry forests; tfor– temperate forests; gsor– grasslands, scrublands, open (barren) habitats and ravines; ohab– open (barren) habitats; agri– agricultural areas; prod– production agroforests; hamt– high altitude mountains; habt– combined area of plausible habitats (species-specific); rock– rocky outcrops and escarpments; pptn– annual precipitation; rugg– terrain ruggedness; prey– wild prey index; elev– elevation; catl– cattle population; shot– sheep and goat population; lstk– livestock (cattle, sheep and goat) population; hpop– human population; dogs– free-ranging dog population; sett– human settlements; ifra– density of roads and railways; resv– extent of Protected Areas

	dhole	golden jackal	Indian wolf	Tibetan wolf	Indian fox	red fox	desert fox	striped hyena
fcov	0.26 (0.18)	0.02 (0.15)	–	–	–	–	–	–
dfor	–	–	0.12 (0.11)	–	–	–	–	0.12 (0.12)
tfor	–	–	–	0.04 (0.61)	–	–	–	–
gsor	–	0.78 (0.62)	0.12 (0.13)	0.20 (0.54)	0.25 (0.12)	1.09 (0.80)	1.14 (0.73)	0.22 (0.13)
ohab	–	–	–	–		–	1.01 (1.27)	–
agri	–	0.16 (0.19)	0.01 (0.13)	–	0.08 (0.13)	1.07 (0.63)	0.35 (17.0)	0.06 (0.13)
prod	0.14 (0.13)	1.54 (0.89)	–	–	–	–	–	0.18 (0.11)
hamt	–	–	–	0.10 (0.92)	–	0.89 (0.54)	–	–
habt	0.34 (0.26)	0.14 (0.08)	0.17 (0.14)	0.33 (0.38)	0.27 (0.12)	0.60 (0.82)	2.65 (1.17)	0.01 (0.09)
rock	–	–	0.01 (0.15)	–	0.08 (0.13)	–	–	0.51 (0.32)
pptn	0.28 (0.20)	0.49 (0.19)	0.01 (0.13)	1.82 (1.14)	0.19 (0.12)	12.92 (9.43)	–	–
rugg	0.21 (0.15)	0.29 (0.12)	0.49 (0.20)	0.98 (1.57)	0.14 (0.12)	1.46 (0.74)	–	0.03 (0.12)
prey	0.70 (0.17)	–	0.43 (0.15)	–	–	–	–	–
elev	–	–	–	0.64 (0.69)	–	1.65 (0.67)	–	–
catl	0.32 (0.18)	–	–	0.12 (0.84)	–	–	–	–
shot	–	–	0.09 (0.12)	0.39 (0.66)	–	–	–	–
lstk	–	–	–	–	–	–	–	0.02 (0.11)
hpop	0.37 (0.19)	4.45 (1.55)	0.34 (0.19)	0.24 (0.95)	0.46 (0.27)	0.16 (0.58)	0.08 (5.68)	0.14 (0.14)
dogs	0.003 (0.13)	0.64 (0.46)	0.48 (0.26)	1.95 (1.81)	0.20 (0.16)	0.90 (0.58)	–	0.24 (0.18)
sett	0.03 (0.22)	0.60 (0.41)	0.21 (0.15)	3.77 (13.61)	0.18 (0.15)	1.21 (1.73)	–	0.10 (0.12)
ifra	0.26 (0.21)	0.79 (0.58)	0.20 (0.15)	10.77 (16.09)	–	0.49 (0.51)	0.14 (11.8)	0.02 (0.13)
resv	0.59 (0.20)	–	–	–	–	–	–	–

a.

Priority ranking

top 10% 20 30 40 50 60 70 80 90 100%

b.

Canid Conservation Units (CCU)

- tier 1
- tier 2
- tier 3

c.

- CCU
- Protected Areas

AP– Andhra Pradesh; AR– Arunachal Pradesh; AS– Assam; BR– Bihar; CG– Chhattisgarh; GA– Goa; GJ– Gujarat; HP– Himachal Pradesh; HR– Haryana; JH– Jharkhand; JK– Jammu Kashmir; KA– Karnataka; KL– Kerala; MH– Maharashtra; ML– Meghalaya; MP– Madhya Pradesh; OR– Odisha; PB– Punjab; RJ–Rajasthan; TN– Tamil Nadu; TS– Telangana; UK– Uttarakhand; UP– Uttar Pradesh; WB– West Bengal

Number of projects

1   29

Canid Conservation
Units (CCU)