Host environmental conditions induce small fungal cell size and alter population heterogeneity in *Cryptococcus neoformans*

Xin Zhou[1], Hanna Zafar[1], Poppy Sephton-Clark[2], Sally H. Mohamed[3], Ambre Chapuis[4], Maria Makarova[1], Donna M. MacCallum[4], Rebecca A. Drummond[3], Ivy M. Dambuza[4*], Elizabeth R. Ballou[1*]

[1]*Institute of Microbiology and Infection, School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK*

[2]*Genome Sequencing and Analysis Program, Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, 02142, Massachusetts, USA*

[3]*Institute of Microbiology and Infection, Institute of Immunology and Immunotherapy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK*

[4]*Institute of Medical Science, University of Aberdeen, Aberdeen, UK*

[5]*Medical Research Council Centre for Medical Mycology, University of Exeter, Exeter, EX4 4QD, UK*

*To whom correspondence should be addressed:

Ivy M. Dambuza: i.m.dambuza@exeter.ac.uk

Elizabeth R. Ballou: e.r.ballou@bham.ac.uk

Funding

XZ is supported by a studentship from the Darwin Trust. HZ is supported by an NC3Rs PhD studentship (NC/R001472/1). PSC was funded by a BBSRC MIBTP PhD studentship. RAD and SHM are supported by the Academy of Medical Sciences (SBF004_1008) and the Medical Research Council (MR S024611_1). MM is funded by the University of Birmingham School of Biosciences. AC and DMM were supported by the NC3Rs (NC/N002482/1). IMD is funded by the Wellcome Trust (102705 and 097377) and the MRC Centre for Medical Mycology (MR/N006364/1). ERB is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (211241/Z/18/Z).

Author summary

The human fungal pathogen *Cryptococcus neoformans* causes 200,000 HIV-associated deaths each year. In the lung, *Cryptococcus* makes an unusual yeast-to-titan morphological switch that contributes to disease development by altering immune polarization and introducing aneuploidy underlying host stress and drug resistance. Specifically, a proportion of 5 um haploid yeast endoreduplicate and swell, converting to large (> 10 um) polyploid titan cells that can then produce genetically distinct daughter cells. We recently developed an *in vitro* protocol for inducing large titan cells and additionally observed a novel small "titanide" cell type. Here we investigate the nature and origin of these small cells, demonstrating that they emerge during both *in vitro* and *in vivo* mouse-passaged titan induction in the well characterised lab strain H99 and are also apparent in a titanizing clinical isolate, Zc8. We show that these titanide cells share features with titan mothers (lipid order) and with spores produced during heterothalic mating. Finally, we show that the capacity of clinical isolates to produce both titan and titanide cells impacts aneuploidy and the emergence of drug resistance *in vivo*.

Abstract

Fungal morphology significantly impacts the host response. Filamentation and tissue penetration by *Candida* and *Aspergillus* species are essential for virulence, while growth as a yeast allows the thermal dimorphic fungi *Coccidiodes, Histoplasma,* and *Talaromyces* to reside inside phagocytes and disseminate. The basidiomycete *Cryptococcus neoformans* exhibits an unusual yeast-to-titan transition thought to enhance pathogenicity by increasing fungal survival in the host lung and dissemination to the central nervous system. In a common laboratory strain (H99), *in vitro* and *in vivo* titan induction yields a heterogenous population including >10 μm titan cells, 5-7 μm yeast cells and 2-4 μm titanides. Previous reports have shown that titan cells are associated with enhanced virulence and the generation of aneuploid cells that facilitate stress adaptation and drug resistance, while small (>10 μm) cells are associated with increased dissemination. However, the relationship between titan cells, small cells, and titanides remains unclear. Here, we characterize titanides and small cells in H99 and three clinical isolates and show that titanides share the lipid membrane order of their titan mothers and the G_0 quiescent-like DNA staining of mating spores. In addition, we show that both titanizing and non-titanizing isolates exhibit altered capsule structure and PAMP exposure over time during *in vitro* culture, and generate aneuploidy *in vivo*.

Introduction

Cryptococcus neoformans is a human fungal pathogen causing life-threatening pneumonia and meningitis in immunocompromised and immunocompetent individuals [1-3]. In the host lung, inhaled *C. neoformans* spores and yeast proliferate and can undergo an unusual morphogenic switch from 5-7μm haploid yeast to large (>10 μm) polyploid titan cells with altered cell wall and capsule organization [4-7]. This is a form of fungal gigantism, with individual cells reaching 50 to 100 μm [4-6, 8-10]. Titan cells also produce aneuploid small (<10 μm) daughter cells, although the importance of these in disease progression is unclear [3, 11-14]. Titan cells enhance the dissemination and virulence of *C. neoformans* in animal and mini-host models of infection [4, 5, 15-18]. Yet, titan cells block phagocytosis, and phagocyte-mediated dissemination, of non-titan cells. Mutants deficient in titan cells exhibit reduced virulence, and mutants that produce fewer small cells disseminate more slowly [15, 19-21].

A pressing clinical question is how titan cells influence patient outcome in a clinical setting. Clinical isolates vary widely in their expression of known virulence factors including capsule, melanin, and urease, a result of the wide range of infecting genotypes even in relatively localized settings and within individual patients [13, 22-26]. Given this diverse virulence portfolio, it is unsurprising that a screen of clinical isolates demonstrated that fungal cell size alone is insufficient to explain *in vivo* outcomes [22]. However, two recent analyses of clinical isolates highlighted a role for small cells in disease progression. Following long term growth in capsule-inducing medium, Fernandes *et al*. observed either giant (>15 μm including capsule) or thick-walled micro (<1 μm) cells in *C. tetragattii* and *C. neoformans* isolates, respectively, and correlated this with CD4 clinical outcomes [13]. Rare *C. neoformans* isolates that produced both giant and micro cells were highly virulent. Mukaremera *et al*. also observed an inverse correlation between virulence in humans and the presence of large cells following culture in DMEM+serum, conditions in which the virulent, titanizing isolate KN99α failed form titan cells [22]. Denham et al showed the emergence of small cells in the lung over the course of infection and showed that small cells predominate in the brain [19]. Together both studies suggest a role for small cells in infection progression, however the link between small and titan cells remains undetermined. By studying H99 titan-induced cultures, we identified a distinct small cell type which we termed titanides [14]. Titanides have distinct morphological properties: oval cell shapes, small cell size (2-3 μm) and thin cell walls [14]. These are morphologically distinct from previously described micro

cells (round, thick cell wall, <1 μm) and drop cells (5μm), both of which have been associated with *in vivo* infection [8, 27, 28].

Genetic and genome-wide analyses in *C. neoformans* H99 have shown that the yeast-to-titan transition is a complex phenotype regulated by multiple interacting pathways[14, 20, 29-31]. Titan cells have been observed in *C. neoformans* VNI, VNII, VNB, *C. deneoformans* and *C. gattii* clinical and environmental isolates during growth in established *in vitro* titan-inducing conditions that recapitulate the host environment [14, 29, 31, 32]. However, in these conditions, we and others also observed a striking variation in size range and morphology regardless of the presence of classic titan cells (>10 μm cell body) [14, 29, 31]. Given evidence of induced population size heterogeneity across and within species, as well as evidence demonstrating conserved genetic regulation of titanization across species, we hypothesized that even in isolates that do not exhibit classic >10 μm "titan" cells, exposure to *in vitro* titan-inducing conditions might trigger important changes in titan-related phenotypes (size distribution, cell wall organization, ploidy, capsule), with implications for fungal virulence.

To investigate this, we examined the impact of *in vitro* titan-inducing conditions on four representative clinical isolates: H99, Zc1, Zc8, and Zc12, selected based on their genetic relatedness and their range of titan phenotypes: titanizing (H99, Zc8) or non-titanizing (Zc1, Zc12). Here, we fully characterized these isolates and find that, like titanizing strains, 'non-titanizing' strains also display population heterogeneity in size distribution, cell wall organization, and DNA content following exposure to inducing conditions. We identify titanide cells both during *in vitro* culture and *in vivo* and demonstrate that titanides are the progeny of polyploid titan cells, sharing their lipid membrane organization. Moreover, we show that titanides share features of quiescent cells and spores. In addition, we find that the 'non-titanizing' Zc12 isolate, despite being defective in producing titan and titanide cells, do produce small cells and can generate aneuploid offspring *in vivo*. Finally, we observe that *in vivo* aneuploidy does not directly correlate with drug resistance. Together, this work confirms the importance of small cells for disease progression, identifies titanides as a specific subset of small cells originating from titan mothers, and raises questions about the role of population heterogeneity in disease progression.

Results

In this study, we examine clinical isolates previously collected from the ACTA Lusaka

Trial in Lusaka, Zambia [26, 33]. We previously showed that Zambian clinical and

environment strains collected by Vanhove *et al.* vary widely in their capacity to form titan

cells, irrespective of originating niche or genotype [14]. Here, we further characterize

titanization in four strains- H99, Zc1, Zc8 and Zc12- selected based on their capacity to form

titan cells *in vitro*. *In vivo* induced titan cells are characterized by four defining features:

enlarged cell sizes, high ploidy, tightly compacted capsule, and the presence of a single, large

vacuole [6, 34]. Using this definition, established by Nielsen and Zaragoza, we previously

classified Zc8 as titanizing, and Zc1 and Zc12 as non-titanizing [6, 14]. We also showed that

Zc1 exhibits reduced virulence compared to H99, and that all four isolates produce melanin

to comparable levels [14].

***In vitro* titan inducing conditions trigger a range of cell size and DNA content changes**

across four clinical isolates

All four strains are of clinical origin, have been fully sequenced and genotyped, and are

genetically closely related (VNI). During standard *in vitro* YPD culture, cells were uniform in

size, generally round with no defects in budding, and cell size was similar across and within

all strains. To more fully characterize the impact of titan induction on all four strains, we first

examined cell size distribution across the four clinical isolates under *in vitro* titanizing

conditions. Consistent with our previous classification, H99 and Zc8 produce highly

heterogeneous populations comprising titans (>10 μm), yeasts (>4μm and <10μm), and

titanides (>1 μm and <3 μm oval cells), while Zc1 and Zc12 populations are more uniform

(Fig 1A). After 24 hours, cells >10 μm account for about 15% of the H99 total population,

but only approximately 2% in Zc8 (Fig 1B). Moreover, induced non-titan Zc8 cells (<10 μm)

are smaller on average than induced non-titan H99 cells (3.80 μm ± 0.10 vs 7.15 μm ± 0.14

SEM, p<0.0001, n>200), and are smaller on average than Zc1 or Zc12 induced cells (4.44 μm

± 0.05, 4.57 μm ± 0.06 SEM, respectively, p<0.0001, n>200).

Given the low frequency of titans detected in induced Zc8 cultures, which can easily be

missed when relying on cell counts with small numbers (n=200), we assessed cell size spread

and DNA content across the population of a large number of cells by flow cytometry (up to

n=10,000). Using this approach, for H99 we observed population structure characteristic of

typical and titan populations, with a minority sub-population of cells distinguishable as a shift

to the right by forward scatter (FSC, a common proxy for cell size) (Fig 1Ci, n>8000) and

DNA content >4C (12.3%, Fig 1D, see Fig S1A for matched YPD controls). However, while

6.4% of induced Zc8 cells had DNA content >4C, this FSC shift was not apparent as H99

(Fig 1Cii). In addition, the median FSC of cells with >4C DNA content was lower than in the

H99 population, suggesting these Zc8 polyploid cells may be smaller than H99 polyploid

cells. Despite this overall reduction in frequency, a similar stepwise population of polyploid

cells with a long tail to the right was apparent for Zc8 (Fig 1D). Unlike H99, the vast

majority of cells in titan-induced Zc1 and Zc12 populations had DAPI staining consistent

with 1C DNA content. However, when we closely examined DNA content, we did identify

rare polyploid 4C cells in Zc1 isolates (2.8%) and Zc12 (<1%). These cells were similar in

FSC to H99 4C cells, suggesting the presence of rare titan cells in the Zc1 and Zc12 cultures,

which likely require acquisition of larger sample size in order to detect (Fig 1C iii). These

data suggest that exposure to titan inducing conditions has a general impact on both clinical

isolate ploidy and cell size.

Induced cells exhibit changes in PAMPs exposure

C. neoformans titan and typical cells recovered from mouse lung have different cell wall

composition, with an increase in chitin and mannose (including mannan, mannoproteins, and

capsular mannans) observed in cells >10 μm [7, 18]. We therefore asked whether similar

differences could be observed in *in vitro* induced cells. Chitin (Calcofluor white, CFW),

chitosan (EosinY), and cell wall mannan (ConA) were measured following exposure to

inducing conditions for 24 hours. Microscopy and flow cytometry revealed an overall

decrease in PAMP exposure following titan induction relative to YNB (Fig 2A, 2Bi, H99

representative plot). Gating on positively staining cells only, chitin, mannan, and chitosan

staining correlated with cell size, with larger cells staining more intensely than smaller cells,

the majority of which were below the threshold of detection (Fig 2Bii). Moreover, distinct

high chitin populations were apparent in H99, Zc1, and Zc8 cultures (Fig S2). Quantitative

analysis by flow cytometry using the same approach to identify titan cells as taken in Fig 1C

again revealed a sub-population of large cells in all four induced cultures (Fig S2). Similar to

in vivo-derived H99 titan cells, *in vitro*-induced H99, Zc1, Zc8, and Zc12 cells displayed a

distinct CFW-high population that correlated with the largest cells (Fig 2B, Fig S2) [7, 18].

The largest cells in all isolates had comparable chitosan and cell wall mannan staining to

typical cells (Fig S2). Moreover, in long-term induced cultures (72 h), overall size

heterogeneity was more apparent, with increased numbers of small cells and distinct high and

low chitin peaks present in both Zc1 and Zc12 populations (Fig 2C). Together, these data

suggest that changes in cell wall composition occur during low density growth in serum

regardless of induction of cells larger than 10um.

Growth in titan inducing conditions can alter virulence relative to yeast-phase growth

Next, we assessed relative virulence of the four isolates using a wax moth larva model [35,

36]. When pre-grown in YPD, the non-titanizing isolate Zc12 was significantly more virulent

than the other three isolates (n=10, Gehan-Breslow-Wilcoxon test, p=0.00070) (Fig 3A),

killing a majority of larva within 5 days, whereas a majority of Zc8 infected larvae survived

to pupation [37]. However, when cells were pre-grown for 24 hr in titan-inducing conditions

prior to infection, H99, Zc1, and Zc8 became as virulent as Zc12 (Fig 3A) [14]. Comparing

the impact of inducing condition on the change in virulence revealed that no change was

observed for Zc12, but induced Zc1, Zc8, and H99 cells increased in virulence relative to un-

induced cells (Fig 3B). The observation that exposure to titan inducing conditions does not

increase the virulence of Zc12 raised the question of what impact titan inducing conditions

might have on Zc1 to alter its virulence.

Cell density is sufficient to change capsule size and structure during capsule induction

To further explore the differences in disease outcome resulting from titan induction, we

examined capsule structure from yeast-phase and titan-induced stages. The *C. neoformans*

cell wall is covered by a protective capsule, which not only impairs phagocytosis by

macrophages, but also masks PAMPs from being engaged by host PRRs[38]. We first

examined growth during incubation in 10% FCS at high cell density (OD_{600} 0.1), conditions

established to induce robust capsule but not titan cells [14]. Capsule was visualized using

india ink and the capsule to cell body ratio measured (Fig 4B, C). There was no relationship

between wax moth virulence and capsule size: H99 and Zc12 showed an overall larger

capsule size compared to Zc1 and Zc8 (Fig 4B) (p<0.0001, n>100). We previously showed

that the anti-capsule IgM antibody Crp127 binds O-mannosylation sites in capsular

polysaccharide. In standard capsule inducing conditions, Crp127 binding was highly

heterogeneous across and within all four strains, with both stained and unstained cells

apparent (Fig 4C), consistent with our previous reports assessing binding during YPD and

DMEM-induced growth [39]. As assessed by flow cytometry, H99 spanned unstained to high

binding, with 51.8% of cells showing intermediate or high binding. For Zc8, the vast majority

stained with intermediate Median Fluorescence Intensity (MFI (67.7%). For Zc1 the majority

of cells are unstained (86.4%), while 48.1% of cells from Zc12 exhibited intermediate

binding (Fig 4C).

Simply reducing the cell density leads to titan induction:10% FCS at $OD_{600}=0.001$.

After induction, all isolates expressed robust capsule and were reactive to Crp127 IgM.

Analysis of capsule to cell body ratio for each of the populations revealed that Zc12 cells

display reduced capsule compared to the other isolates ($p<0.0001$, $n>100$) (Fig 5B). Across

the four isolates, binding by Crp127 remained heterogeneous, but binding patterns changed

compared to high density capsule induction (Fig 5A, C), consistent with our observation that

Crp127 binding is dynamic throughout titan development [39]. Similar to high-density

capsule induction, for H99 and Zc12, more than 50% of the total population was positive for

IgM binding, but H99 staining was higher overall, while Zc12 remained more intermediate.

For Zc1, which was predominately negative for IgM under high-density capsule conditions,

~40% of cells became positively stained. However, for Zc8 the pattern was reverse: only 5%

of Zc8 cells, the largest cells present, were positive for IgM binding, whereas a majority of

Zc8 cells were positive for IgM under high density capsule induction (Fig 4C, D). Together,

these data suggest that differences in cell density during exposure to inducing conditions

($OD600 = 0.1$ vs 0.001) significantly impact capsule structure, with a high potential for

altered capsule expression and organization across genetically related clinical isolates, even

in absence of large-scale cell size changes (Zc1, Zc12).

What are titanides?

Given the lack of correlation between conventional virulence factors and relative virulence in

the *G. mellonella* model, we asked whether there was a correlation with the presence of small

cells during infection. Specifically, we previously identified a third distinct cell type in

induced cultures: oval cells 2-3 µm in diameter with thin cell walls and a clear, well defined

nucleus, which we termed titanides [14]. These cells emerge 24 hrs after serum exposure and

are the predominant cell type in H99 induced cultures after 3 days (Fig 1A, 6A). We

observed titanide cells in Zc8 *in vitro* cultures that shared these features (Fig 1A, 6A) and in

in vivo lung extracts (Fig S3A), where thick walled "micro" cells could also be observed.

Viability analysis of individual cells isolated from *in vitro* cultures by microdissection was

100% within 48 hours of serum induction for both H99 and Zc8 titanides. Titanides are

therefore distinct from previously described micro cells (<1 µm) and drop cells (5 µm),

which have thick cell walls and, for the latter, disorganized nuclei and low viability [8, 27,

28].

We previously speculated that titanide cells are the daughters of titan mothers. To track

their origin, we examined changes in subcellular membrane organization caused by

titanization using the ratiometric dye di-4-ANEPPDHQ (Fig 6B)[40]. We observed that the

plasma membrane in titan cells is significantly more ordered compared to yeast cells (~5 μm),

which had lower lipid packing values (generalized polarization, GP) (Fig 6C). In cells <3 μm,

lipids in the plasma membrane were highly ordered, comparable to lipids in titan mother cells,

further suggesting these cells are the direct progeny of titan mothers (Fig 6C). Finally, live

cell imaging of *in vitro* titan cells revealed that titan daughter cells emerge every 90 minutes

and are on average <3.5 μm in diameter (Fig 6D, S4), identifying these as titanide cells.

Examination of DNA content by flow cytometry for small cell populations revealed that

when grown under inducing conditions, all four clinical isolates contained cells with apparent

"sub-1C" DNA content, represented as stepwise peaks to the left (Fig 1D). These "sub-1C"

populations were not present when cells were grown in YPD (Fig S1 ii). We speculated that

these peaks might represent titanide cells, and in fact they are the smallest cells by FSC (Fig

S1B). Microscopy to compare cell size in high density capsule inducing conditions vs. low

density conditions also showed that these small cells are specifically generated under titan

induction (Fig 6E).

Similar "sub-1C" peaks have been observed during flow cytometry-assisted cell cycle

analysis of *Saccharomyces cerevisiae* and are representative of G0 or quiescent cells [41, 42].

We speculated that this peak might therefore represent G0 quiescent cells, suggesting that

titanides might share similarity to *C. neoformans* spores produced by mating, which are

quiescent prior to germination. To test this hypothesis, we first established that quiescent C.

neoformans spores exhibit sub-1C DNA content similar to S. cerevisiae quiescent cells. H99,

Zc1, Zc8, and Zc12 MATalpha cells were crossed with KN99 MATa cells and incubated for

2 weeks to ensure robust sporulation. Zc1 and Zc12 formed robust hyphae and sporulating

basidia after 7 days, comparable with H99. Zc8, a MATalpha isolate, exhibited a profound

defect in the formation of hyphae, with sporulating basidia rarely visible even after 3 months

(Fig S3B). The mating front, representing hyphae, basidia, spores, and some contaminating

yeast, was excised, and filtered through 40 um filters to exclude large particles. The resulting

population was fixed, stained and analysed by flow cytometry alongside yeast-phase cells to

identify 1C and 2C gates and a "sub-1C" background of <2.5%, set as a threshold for false

positives using actively growing YPD cultures. This analysis revealed a "sub-1C" peak to be

a major population in robust mating reactions (H99: 37%; Zc1:34%; Zc12:53%) and a minor

population (10%) in the Zc8 mating front, in which basidia were rarely observed (Fig 6F,

S3B). Direct comparison to titan-induced H99 cultures revealed a similar "sub-1C" peak (8%)

suggesting that this population represents quiescent G0 cells specifically generated during

titan induction.

Modelling fungal-phagocyte interaction *in vitro* reveals strain specific phenotypes

To better understand the impact of titan and small cells on disease development, we

analyzed the infection process of all four strains following titan induction using an *in vitro*

model. We observed significant variation in the percent of macrophages containing at least

one internalized *C. neoformans* cell across the four isolates after titan induction, while YNB

grown cells were phagocytosed similarly across the four isolates (Fig 7A). H99 showed the

lowest macrophage infection rate and was significantly reduced compared to the other 3

strains (Fig 7A) (compared to H99: Zc1, p=0.0063; Zc8, p=0.0131; Zc12, p=0.0164). As

shown in Fig 7B, for both H99 and Zc8, engulfed cells were typical cells (titanides and yeasts)

rather than titan cells. Furthermore, capsule staining with 18B7 revealed that H99 titan cells

physically associated with macrophages but were not engulfed (Fig 7C). We therefore asked

whether the reduced uptake of H99 cells overall relative to Zc8 (Fig 7A) is a consequence of

the higher proportion of titan cells in this population, similar to in vivo reports. Filtered titan

(>10 μm) cells or typical (<10 μm) H99 cells were used to infect macrophages separately.

When only titan cells were present, the average macrophage infection was 21.6% ± 5.5

(SEM), while for typical H99 cells the rate was 82.2% ± 1.6 (SEM) (Fig 7D, p=0.0002,

n=500), similar to uptake of the total populations for the three Zc strains. Mixed cultures of

titanized cells were co-incubated with macrophages in vitro and observed via time-lapse

imaging for 18h. Intracellular Proliferation Rate (IPR), frequency of vomocytosis (vomo %),

and frequency of macrophage death (Mφ death %) were measured. No significant difference

in IPR was observed for the four strains, indicating that they maintain similar intracellular

growth rates (Fig 7E). Zc8 induced the highest frequency of vomocytosis, although it was not

statistically significantly different (p=0.05 vs. Zc12) (Fig 7F), but did induce significantly

more macrophage death (p<0.001) (Fig 7G; n=400). This suggests that there is no correlation

between titanization capacity and vomocytosis or macrophage death. These results are

consistent with previous studies *in vivo* showing that titan cells are more resistant to

phagocytosis and with the observation that these cells exert a protective effect on the uptake

of smaller cells and further validate the overlap between in vitro induction and in vivo

induction [5, 12].

In vivo titan inducing conditions match *in vitro* predictions

Overall, *in vitro* phenotyping by microscopy and flow cytometry for cell size and DNA

content predicts that: 1) H99 and Zc8 isolates will form titans *in vivo*, with the overall size of

Zc8 populations smaller than that of H99, 2) a minority of Zc1 titan cells might be detected, 3)

Zc12 will fail to produce detectable titans *in vivo*, and 4) small cells will be apparent for all

four isolates and will proliferate to a similar degree within phagocytes. We therefore assessed

all four strains in a murine inhalation model of cryptococcosis, the gold standard for

determining titan capacity. C57BL/6 female mice were infected intra-nasally with $5x10^5$ cells

pre-cultured in sabouroud dextrose medium according to standard infection protocols, and

brain and lung fungal burdens were measured after 10 days. Work using H99-derived

mutants has demonstrated a role for titan cells in survival and proliferation in the pulmonary

environment as early as day 7 [15]. When fungal cells from lung homogenates were analysed

microscopically for titanization *in vivo*, similar size distributions were observed as seen in

vitro (Fig 8A). Specifically, robust titanization was observed for both H99 and Zc8 (65% and

68% of cells recovered from the lung, respectively, >10 um), with Zc8 population covering a

significantly smaller size range than H99 (Kolmogorov-Smirnov test for cumulative

distribution, p=0.0001). A minority (8%) of Zc1 cells were >10 um. No Zc12 cells >10 um

were observed. Similar to in vitro induced Zc1 and Zc12 cells, the *in vivo* populations as a

whole were smaller and more uniform overall. All four clinical strains showed similar fungal

burdens in the lungs on day 10 (Fig 8B, p>0.05) as assessed by CFUs, demonstrating similar

capacities to proliferate early during infection. Moreover, brain CFUs analysed at day 10 post

infection were comparably low across all four isolates, with Zc1 CFUs slightly higher than

the others (Fig 8C; p<0.01). Taken together, these results support our predictions that *in vitro*

titanization reflects *in vivo* titanization.

Clinical isolates show different capacities to produce aneuploid colonies

In vivo polypoid titan cells were reported to be able generate aneuploid progeny with

improved stress resistance compared to typical cells from the same culture [12]. Our *in vitro*

model predicts that high titanizing isolates H99 and Zc8 will yield greater aneuploidy *in vivo*.

It remains unclear whether ploidy variation in CNS cryptococcosis reflects the degree of

ploidy variation observed in the lung. However, because titan daughter cells are smaller on

average and readily phagocytosed, we predict that ploidy variation following dissemination

to the brain will reflect the degree of ploidy variation observed in the lung. To test this, we

determined the impact of titanization on the genetic diversity of offspring and the impact on

different sites. We examined the ploidy of individual colonies originating from lung and

brain CFUs for all four isolates (Fig 9A, Fig S5, n=30 for lung CFUs, n≥10 for brain CFUs,

across 3 mice per isolate). Colonies were assessed within 72 hours of recovery from mice, at

the point when a small colony was just visible on YPD agar (~36 generations), a measure of

stable aneuploidy in the absence of environmental stress. Fig 9A shows the ploidy of

individual colonies relative to the median ploidy of all colonies from that particular lineage.

Interestingly, ploidy variation occurred for all isolates, including Zc12. There was no

significant difference in relative ploidy between colonies from lung vs. brain isolates for any

of the isolates. We observed that ploidy of H99 and Zc8 colonies varied widely, while Zc1

colonies were more uniform (p=0.02 compared to H99). This is consistent with a high degree

of titanization driving aneuploidy in H99 and Zc8 infections, and relatively low titanization

in the Zc1 isolate. However, we also observed widespread instances of aneuploidy in Zc12

CFUs, despite finding no evidence of true titanization in this strain either *in vitro* or *in vivo*.

380   Aneuploidy is associated with increased resistance to fluconazole, which continues to

be used as a frontline antifungal in resource limited settings. Fluconazole resistance in *C.*

neoformans is linked to aneuploidy of chromosomes 1, 4, 6, and 10 due to the presence of

genes on these chromosomes that directly enable fluconazole resistance, however the

contribution of titanization to resistance remains unclear [43]. We therefore assessed the

impact of altered ploidy on drug resistance of individual colonies isolated from the lungs and

brains of infected mice (Fig 9B). Relative to parent isolates, we observed a wide range in

capacity to proliferate in the presence of fluconazole. The distribution of resistance for Zc12-

derived colonies was significantly limited compared to those of H99, Zc8, or Zc1 (p≤0.0002).

For H99, both fluconazole resistant and sensitive colonies were identified. For Zc8, colonies

with increased resistance to fluconazole tended to have increased ploidy, but a colony with

reduced overall ploidy and increased drug resistance was also observed. For Zc1, aneuploid

colonies were more sensitive to the drug. Within individual lineages there was no correlation

between increased or decreased ploidy and degree of fluconazole resistance (R^2<0.3 in all

cases). This is consistent with titanization facilitating ploidy shifts in progeny and with

aneuploidy in these isolates being random rather than specific amplification of chromosomes

involved in fluconazole resistance occurring prior to drug exposure.

Discussion

The first clinical and murine model reports of cells consistent with giant/titan cells also reported significant size and morphological heterogeneity. Examination of cells from fungal abscesses revealed a highly heterogeneous population of cells ranging from 2-100 um[9, 10]. Consistent with this, infection of mice with laboratory strains yields a highly heterogeneous population comprised of large, polyploid titans and typical yeast as well as small, thick walled "micro" cells in lung extracts[5, 8, 20, 44]. However, replicating this heterogeneity in vitro presents a challenge to modeling *in vivo* infection. When cultured on rich medium, clinically heterogenous populations become uniform, comprised of only typical yeast[9, 10]. Moreover, this heterogeneity may be site specific within patients: clinical examination of *C. neoformans* isolates taken directly from CSF failed to identify cells that cross the defining 10 μm cutoff [45]. Whether this is because titan cells cannot escape the host lung or because the CSF environment is not supportive of *de novo* titan cell formation remains unclear [34, 46]. Even during co-culture with immune cells that replicate host conditions and elicit titan cells, heterogeneity is not observed [44].

Some degree of heterogeneity is observed under capsule inducing conditions: Fernandes et al. reported that 5 days exposure to DMEM (37°C, 5%CO2) yielded giant and micro cells and rare isolates that showed both giant and micro cells were significantly more virulent[47]. Interestingly, these authors also reported wide variation in size and fungal morphology within and between isolates and across serotypes [13, 47]. Mukaremera et al. also observed that strains that produce small cells were more virulent, using 5 days growth in DMEM+serum (37°C, 5%CO2), conditions in which the high virulence titanizing laboratory isolate KN99a failed to produce titans. However, neither group report observing the wide variation in cell size observed in established titan inducing protocols, nor do they observe small oval cells consistent with titanides[14, 29, 31]. Despite this, these works, together with the findings of Denham et al. reveal a role for small cells in infection progression [19]. While it is established that the presence of titans can affect drug resistance and host immune responses, contributing to poor disease outcome, the role of small cells is less clear [4, 5, 15, 20]. Therefore, a pressing question is the capacity of clinical and environmental isolates to generate population heterogeneity regardless of classic titanization capacity and the effect of this population heterogeneity on disease outcome.

Investigating population heterogeneity *in vitro*

Here, we expand upon the *in vivo* definition of titan cells to encompass the wide

phenotypic diversity observed during *in vitro* induction of three clinical isolates from the

ACTA Lusaka Trial in Lusaka, Zambia [26, 33]. Our data add to existing knowledge by

specifically studying clinical isolate titanization using an established in vitro titan induction

protocol. After in vitro titanization, *C. neoformans* populations are highly heterogeneous with

multiple sub-populations: titan (>10um), typical/small (>4 μm), and titanide (2-4 μm) cells,

as well as micro (1 μm) cells [5, 8-10, 14, 44]. Building on our previous findings, we used 4

clinical strains from the same genetic clade (VNI) which we previously identified as having

different capacities to produce titan cells [14]. These four strains are closely related based on

genome analysis, however evolved variation still exists between them.

All 4 strains demonstrated significant population heterogeneity after exposure to inducing

conditions, regardless of titan status. Relative to Zc8, H99 is strongly titanizing *in vitro* when

the *in vivo* definition (> 10 μm) is applied, but Zc8-induced cells exhibit overall population

structure consistent with titanization, including polyploidy (>4C), a relative increase in cell

size, and overall changes in PAMP organization. By combining microscopy and flow

cytometry, we propose a model in which predictive outcomes can be made about *in vivo*

strain "titan-signature" status. We additionally note that researchers can readily identify titan

cells through visual inspection of liquid cultures in tissue culture plates using bright-field

microscopy, with titans being large cells with a single vacuole and an intensely dark cell wall

apparent in H99, Zc8, and Zc1 cultures, but DNA content analysis offers important

information about overall population structure (Fig 1A).

Variation in cell size and morphology and impact on host cell interaction

Population heterogeneity appears to be a general feature of titanizing isolates. Here we

observed that both titanizing strains H99 and Zc8 are highly heterogeneous after induction,

with three sub-populations that maintain distinct cell sizes and DNA contents. Moreover,

titan cells from both strains exhibit changes in PAMP exposure and capsule size and

composition, features important for the virulence of *C. neoformans*. Surprisingly, non-

titanizing induced strains Zc1 and Zc12 also show population heterogeneity as measured by

DNA content and cell wall structure, despite a lack of true titan cells. Consistent with

previous findings, we directly showed that the presence of titan cells leads to decreased

phagocytosis rates of small cells [5, 12]. Denham et al. showed that at later infection stages,

in both lungs and brains the predominant population is the smaller cryptococcal cells around

10μm in total diameter[19]. This matches our *in vitro* observation that in later titan-induced

cultures, the proportion of small cells increases, consistent with the increased replication rate

of titan mothers and the impact of cell culture density on the generation of new titan

mothers[14, 29, 31].

We additionally observed changes in capsule and IgM binding upon titan induction relative

to other capsule inducing protocols. IgM mediates complement-dependent phagocytosis and

has been shown to specifically inhibit the yeast-to-titan transition [48, 49].Interestingly, IgM

has been shown to be particularly important in the clearance of small particles (2-5μm) [50].

Our data here suggest differences in epitope exposure on small vs. large cells may exert a

protective effect by reducing overall phagocytosis. Future work will examine the impact of

titan induction on complement-dependent, opsonin-independent uptake and will address the

relative pathogenic potential of titanide vs yeast vs spores in murine models of infection.

How do titanides contribute to disease progression?

There is growing evidence of a high degree of genome plasticity during *C. neoformans in*

vivo growth [51-53]. Semighini *et al.* previously demonstrated, using comparative genome

hybridization arrays in H99-derived strains, that passage through mice in the absence of drug

stress results in large scale genome alterations, including aneuploidy, insertions, and

deletions [51]. Gusa *et al.* recently demonstrated that host temperature is sufficient to induce

transposon-mediated rearrangements in the *C. neoformans* genome [52].However,

titanization adds to and enhances this variation: titan cells produce aneuploid daughter cells

that promote *in vitro* stress resistance relative to yeast in the same population [12]. We add to

these previous reports by showing that high titanizing H99 and Zc8, low-titanizing Zc1 and

non-titanizing Zc12 isolates all yield aneuploid daughters to some degree when exposed to

the host microenvironment. We reveal that titanization can drive the emergence of aneuploid

offspring in the host by comparing the ploidy distribution of cells recovered from infected

mice. We observed large scale changes in genome content relative to parent isolates, and this

weakly correlates with degree of titanization (H99>Zc8>Zc1>Zc12). We hypothesize that, in

addition to strongly inducing titanization, during infection, the host microenvironment itself

is sufficient to induce aneuploidy and exerts a selective pressure, as has been observed for *C.*

albicans [54]. In line with this, we previously reported that daughter cell colonies derived

from *in vitro*-induced titan cells are mostly diploid, in conflict with reports from Gerstein *et*

al. that colonies derived from *in vivo*-derived titan cells are haploid or aneuploid [12, 14].

This suggests that daughter cells derived from the two conditions may differ in their tolerance

to aneuploidy or capacity for genome maintenance. Fluconazole exposure, oxidative stress,

and high temperature drive *in vivo C. albicans* aneuploidy, including whole ploidy reduction

(diploidy to haploidy), individual chromosome aneuploidy (Chr 5), and segmental aneuploidy,

and the same is likely true for *C. neoformans* even in the absence of titanization [54-57].

Future work will more fully probe the molecular mechanisms differentiating these clinical

isolates and their impact on virulence.

What is the environmental significance of "sub-1C" cells?

Ashton et al. recently suggested that the existence of robust quiescent cells such as

spores might contribute to the broad ecological distribution of specific sub-clades, however

the rarity of MATa *C. neoformans* isolates in the environment suggests an alternate origin for

these cells[58, 59]. Here, we demonstrate that titanides observed in induced cultures originate

from titan mothers and that these cells exist in a in G0-like state similar to spores generating

by mating. Titanide cells have unique features including oval shape, smaller size and altered

DNA content compared with typical yeast cells. The ability to induce titan cells in conditions

likely to be encountered by Cryptococcus in the environment make it plausible that titan cells

may form in the environment and are not just limited to patient lungs. In addition, we also we

detected "sub-1C" populations from the Zc12 isolate in titan-inducing conditions, suggesting

that the capacity to produce quiescent cells is not simply a consequence of high ploidy

mothers. This raises the hypothesis that "quiescent" cells proposed to drive population

expansion in environmental isolates may be generated via regulatory pathways driving

titanization [58].

Impact of population heterogeneity on drug resistance

Gerstein *et al.* previously demonstrated that *in vivo*-derived titans provide daughters with

a survival advantage compared to *in vivo*-derived typical cells[12]. Therefore, we predict the

presence of titanides also contributes to disease progression by affecting drug resistance and

stress adaption. Titan-derived colonies are resistant to fluconazole via aneuploidy of

chromosomes 1, 4, 10, 11, and/or 12 when propagated in the presence of fluconazole,

suggesting that titanization drives fluconazole resistance *in vivo* [12, 43]. Our data support

these findings, with overall increases in population diversity in titanizing strains compared to

non-titanizing strains following passage through mice (Fig 7A). However, we observed no

specific correlation between increased ploidy and increased fluconazole resistance (Fig 7B).

This is consistent with *in vivo*-induced aneuploidy being random with regard to particular

chromosomes, rather than a specific increase of chromosomes involved in fluconazole

resistance. Finally, induced but not un-induced cells from all isolates showed high but similar

virulence when using the galleria model and we found no differences in capacity to control

early lung fungal burdens during a mouse model of cryptococcosis. Our data therefore

demonstrate that in response to the host environment signals both non-titanizing and

titanizing isolates generate heterogeneous populations which will have implications for drug

resistance and host cell interactions. Future studies will investigate their impact on long term

immune responses including investigation of T cell polarization mediated by the different

clinical isolates.

Acknowledgements

We thank the doctors, patients, and their families involved in the ATCA Lusaka trial, without

whom this research would not have been possible. We thank the staff of the University of

Birmingham BMSU and the University of Aberdeen Medical Research Facility for their

assistance. 18B7 anticapsule antibody was a generous gift from Arturo Casadevall. Crp127

IgM anticapsule antibody was a generous gift from Robin May. We are grateful to Debbie

Wilkinson and Lucinda Wight in the University of Aberdeen Microscopy and Histology Core

Facility for their expert help with TEM experiments.

Materials and methods

Strains, media and growth conditions: All *C. neoformans* strains used in this study are

listed in Table 1.

Strains	lineages	Clades	Source
H99	VNI	Clinical isolate	North Carolina
Zc1	VNI	Clinical isolate	Zambia
Zc8	VNI	Clinical isolate	Zambia
Zc12	VNI	Clinical isolate	Zambia
CNV111	VN1	H99-derived	[60]

Strains were cultured routinely on YPD (1% yeast extract, 2% bacto-peptone, 2% glucose, 2%

bacto-agar) plates stored at 4°C. For mating reactions, cells were mixed from YPD plates

with KN99a directly on Murashige Skoog agar (M5519 Sigma-Aldrich) and incubated in the

dark at room temperature for at least 2 weeks. Titan cell *in vitro* induction was performed as

described in [14]. Briefly, cells were pre-cultured at 37°C, 150rpm, in 10ml YNB without

amino acids (Sigma-Aldrich) in which 2% of glucose was added according to manufacturer's

protocol. After 24 hrs, cells were normalized to $OD_{600}=0.001$ in $1\times PBS$ (phosphate-buffered

saline) with 10% of Heat inactivated Fetal Bovine Serum (HI-FBS) (Sigma-Aldrich F9665)

and incubated at 37°C, 5%CO_2 for 24 hrs unless otherwise indicated.

***Galleria mellonella* infection model:** *G. mellonella* were purchased from LiveFoods, Ltd

(UK) and used within 3 days of arrival. Worms were maintained at 16°C until use, and only

worms lacking melanization were included in the experiment. Worms were allocated

randomly to each treatment group (10 per group), with the average weight of worms across

treatment groups maintained within 0.002 g. Cryptococcus strains were incubated overnight

in YPD (5% yeast extract, 10% bactopeptone, 2% D-glucose) at 150 rpm, 30 °C, or YNB

(and then exposed to 10% serum in PBS at OD600, 5%CO_2, 37 °C for 16 hrs. Cells were

washed in sterile PBS, and then normalized to 5x 10^6/ml. Strains were randomly allocated to

the different groups. Worms were injected into the back left pro-leg with 20 ul innoculum.

Worms were maintained in the dark at 30 °C and observed for 10 days for survival, with

blinding for the identities of the different treatment groups. Survival was analyzed by

Kaplan-Meyer survival curve and significance was assessed by Log rank (Mantel-Cox) test.

Murine infection model: Female C57BL/6 mice were used at 8-10 weeks of age and were

maintained in individually-ventilated cages under specific pathogen-free conditions at the

Biomedical Research facility at the University of Birmingham (Birmingham, UK). Animals

were provided with food and water *ad libitum.* All experimentation conformed to the terms

and conditions of United Kingdom Home Office license for research on animals

(PPL/PBE275C33) and the University of Birmingham ethical committee.

Cryptococcosis Infection Model: Yeast was grown in SabDex broth (Sigma-Aldrich),

grown at 30°C with shaking for 18-24 hours. Yeast cells were washed in PBS, counted, and

delivered to mice intranasally while under isofluorane anaesthesia. Animals were infected

with 2x10^5 CFU. For analysis of lung and brain fungal burdens, animals were euthanized and

organs weighed, homogenized in PBS, and serially diluted before plating onto YPD agar

supplemented with Penicillin/Streptomycin (Invitrogen). Colonies were counted after

incubation at 37°C for 48 hours.

Preparation of Lungs for FACS: Lungs were removed aseptically from euthanized animals,
finely minced with a scalpel, and digested in digest buffer (RPMI supplemented with 10%
FBS, 1% penicillin/streptomycin [Invitrogen], 1mg/mL collagenase D [Sigma], 1mg/mL
dispase [Sigma], 40 µg/mL DNAse [Roche]) at 37°C for 30 minutes. Digested lungs were
smashed through a 70µM filter, washed and red blood cells lysed on ice using BD
PharmLyse solution. Lung-isolated cells were washed in PBS/2mM EDTA, and then stained
with fluorophore-conjugated antibodies in the presence of anti-CD16/32 and 0.5% BSA for
30 minutes on ice. Samples were washed in PBS/0.5% BSA/0.01% sodium azide and
acquired using the BD Fortessa instrument equipped with BD FACS Diva software (BD
Biosciences). FlowJo (TreeStar) was used for the final analysis. Anti-mouse antibodies used
in this study were: CD45 (30-F11), Ly6G (1A8), Ly6C (HK1.4), SiglecF (S17007L), all from
Biolegend, and CD11b (M1/70), F4/80 (REA126), MHC Class II (REA813), all from
Miltenyi Biotec.

Cell wall staining: Cells from 24 hrs induced titan cultures were collected, washed twice
with 1xPBS and resuspended in 500ul McIIvaine buffer. Cells were stained with EosinY
(5µg/ml) in dark for 10mins and repeated for washes with McIIvaine buffer, then
resuspended in 500ul 1×PBS. CFW (10 ug/ml) and ConA (50 ug/ml) were added into
samples and maintained in the dark for 10 mins. After washing twice with 1×PBS, cells were
imaged with a Zeiss Axio Observer and analysed for fluorescent intensities with Attune NxT
Flow cytometer.

J774.1 cell culture: For most *in vitro* infection assays, we used the J774.1 murine
macrophage cell line. Cells were routinely passaged in Dulbecco's modifies Eagle medium
(DMEM+) culture media with serum (DMEM, low glucose, from Sigma-Aldrich; 10% HI-
FBS; 1% penicillin-streptomycin solution from Sigma-Aldrich; 1% 200mM L-glutamine
from Sigma-Aldrich). All assays were performed with cells in passages 4 to 10.

***In vitro* phagocytosis assays:** The intracellular proliferation assay was performed as
previously described [61]. 24h before infection, 1×10^5 macrophages J774 were seeded into
12-well plastic plates [62] in 1 ml DMEM culture media with serum (DMEM+) and activated
with 10 U/ml IFNγ. Before infection, *C. neoformans* cells were collected from titanization

culture and washed three times with sterile 1×PBS. After normalising to $1×10^6$ cells/ml in

300 μl DMEM+, cells were opsonized with 10μg/ml anticapsular 18B7 antibody by

incubating at 37°C for 30min [63]. Then opsonized *C. neoformans* cells were added into

activated J774.1 macrophages to give a multiplicity of infection (MOI) ratio of 10:1. After 2

hrs (time point 0), infected wells were washed at least three time with pre-warm 1×PBS until

all un-engulfed *C. neoformans* cells were removed from the wells.

Human monocytes were infected with opsonized cells for 4h without IFNγ activation, and at

this point all supernatants were collected for cytokine qualification. 4h post-infection,

monocytes were divided into two groups: with or without IFNγ (10 U/ml) activation and then

infection continued for another 20h. Total 24h later, all supernatants were collected again

with LPS (100ng/ml) as a positive control added at the same time as fungal cells and

uninfected as negative control.

Macrophage infection rates: At time point 0, attached *C. neoformans* cells and

macrophages were stained with Calcoflour white (CFW, 10 μg/ml) and Lyso-Tracker Red

(ThermoFisher, 1 μM/ml) in dark for 10 min and then fixed with 1 ml 4% paraformaldehyde

(PFA) in 1×PBS at 37°C for 5min. After washing with 1×PBS, at least 5 frames from each

single infected well were randomly selected and imaged with Nikon Ti-E microscope,

capturing 500 cells. All images were analysed with FIJI and experiments were repeated 3

times for biological replicates.

Filtered Titans phagocytosis assays: After 24h of *in vitro* induction, H99 titanised total

population was passed through an 11μm filter to separate titan (>11μm) and typical cells

(<11μm). Then both populations were washed, normalized and opsonize independently

before adding to infection assays. Activation and infection of macrophages with opsonized *C.*

neoformans cells followed the same protocol as described above.

Intracellular proliferation and vomocytosis assays. At time point 0 of phagocytosis assays,

extracellular cells were removed by washing with warm fresh DMEM+. All samples were

maintained at 37°C, 5%CO_2 in the Nikon Ti-E microscope chamber and imaged for 18h via

time-lapse. Images were taken every 15 min and compiled into single movie files for analysis

using NIS elements or FIJI software. Movies then were scored visually for intracellular

proliferation rate, macrophage death and vomocytosis percentage. Vomocytosis scoring

followed the guidelines described in Andrew et al., 2017 [64].

Flow cytometry for cell ploidy: Cells were fixed and stained according to the protocol of

[14]. Cells were fixed with 50% methanol for 15 mins, washed 3 times with 1×PBS and

stained with 3 ug/ml DAPI. Cells were analysed for DNA content using an Attune NxT Flow

cytometer with violet laser. All flow cytometry data were analysed with FlowJo. Doublets

and clumps were excluded using recommended gating system of FSC-H vs FSC-A, which

was determined to be more parsimonious than SSC-H vs SSC-W followed by FSC-H vs.

FSC-W. Autofluorescence in the DAPI channel was excluded by comparison to unstained

cells. Gates for 1C and 2C were established with haploid controls cultured in YPD liquid

media as described in the text.

Cell sizes measurement: Cell diameter was measured using FIJI, with frames randomly

selected, all cells in a given frame analyzed, and at least 5 images acquired per sample for

each of two independent runs. Total cell number of each samples was >200. All statistical

analyses were performed using Graphpad Prism.

Measurements of membrane lipid order *in vivo*: Cells from 24 hrs induced titan cultures

were collected, washed twice with 1xPBS. Following resuspension in fresh medium

containing 5 μM di-4-ANEPPDHQ, cells were incubated for 30min at 30°C. Cells were then

transferred into a glass-bottomed microscope dish. Imaging was performed on a Zeiss LSM

780 confocal microscope equipped with a 32 element GaAsP Quasar detector. A 488 nm

laser was selected for fluorescence excitation of di-4-ANEPPDHQ. The detection windows

were set to 510–580 nm and 620–750 nm. The images were analyzed using a plug-in

compatible with FIJI.

685
686
687 **Figure legends**

688
Fig 1. Titan cells formation varies among clinical isolates *in vitro*.
(A) *C. neoformans* cells from all indicated strains after 24hrs titan induction, Scale bar: 10μm. (B)
Cell body sizes of titan induced cells (n>200). (C) Flow cytometry for cells after titanization with
FSC-A as a proxy for cell size. i) Histograms for cell size (FSC-A) are shown for all indicated isolates
with a dotted line indicating the distinct shoulder of size increase. ii) The gray sub-population
indicates the size threshold identified for cells with DNA content >4C. iii) FSC-A vs SSC-A scatter
plots for all indicated isolates with a dotted line indicating the threshold for H99 cell sizes (D) Flow
cytometry analysis for DNA content after staining with DAPI. 1C and 2C populations were identified

relative to YPD grown yeast cells from matched parent strains (Fig S1A).4C peaks were established
according to doubled MFI of DAPI staining relative to 2C. "T" indicates pre-titanized cells. For all
figures, data for a representative biological replicate is presented (n>3).
700
Fig 2. Non-titanising strains also exhibit population heterogeneity. (A) PAMPs staining for all
indicated strains cells from both YNB yeast and 24hrs titan induction, Mannan (ConA, Red), Chitosan
(EosinY, Green), Chitin (CFW, Blue). YNB grown yeast scale bar 5μm; titanized cells scale bar 10
μm. (B) Stained cells from (A) were analysed by flow cytometry to determine MFI indicating
exposure of Chitin (CFW, right); Mannan (ConA, center); and Chitosan (EosinY, left). (C) Flow
cytometry analysis of PAMPs staining for indicated cells from long-term(72hrs) titan inductions with
MFI indicating exposure of Chitin (CFW, right); Mannan (ConA, center); and Chitosan (EosinY, left).
"T" indicates pre-titanized cells.
709
710
711 **Fig 3. Disease outcomes varied from basal to titan-induced conditions on worm model**
712 (A)Survival curves of worms infected by YPD-grown or titan induced cells of four indicated strains. (B)
713 Comparison of survival curves of worms infected with cells from both conditions. "T" indicates pre-
714 titanized cells. For all figures, data for a representative biological replicate is presented (n>3).
715
716
Fig4. Phenotype characterization of clinical isolates at yeast phase.
(A) *C. neoformans* cells from all indicated strains at yeast phase, Scale bar: 20μm. (B) Total capsule
and cell body diameters were measured and a ratio reported for each cell. (n>100). ****: p<0.0001.(C)
i) Representative images of capsule structure from capsule induction condition stained with both India
ink and anti-IgM (Crp127) FITC(Green). ii) Histograms for IgM staining and gates show peaks for
different levels of staining with corresponding frequency and MFI reported. Negative: cells failed to
bind. Scale bar: 10μm. For all figures, data for a representative biological replicate is presented (n>3).
724
Fig5. Capsule structure from titan induction
(A) Representative images of capsule structure of titanized cells stained with India ink and anti-IgM
(Crp127) FITC(Green); bar scale: 10μm. (B) Total capsule and cell body diameters were measured
and a ratio reported for each cell. (n>100). ****: p<0.0001. (C, D) Flow cytometry analysis of IgM
staining with frequency and MFI reported. Ci) Gates show peaks for different levels of staining, with
(D) corresponding MFI indicated. Cii) Size (FSC-A) vs IgM-FITC intensity, with different
populations highlighted as in (Ci). "T" indicates pre-titanized cells. For all figures, data for a
representative biological replicate is presented (n>3).
733
734
Fig 6. Titanides from both *in vivo* and *vitro*
(A) TEM of titanide cells from H99 and Zc8 from *in vitro* titan induction. Scale bar:500nm (B)
Single-plane pseudocolored generalized polarization (GP) images. Color bar designates the range of
GP values where red shows high membrane order and blues show low membrane order. Note plasma
membrane zones were used to quantify mean GP values. Scale bar: 5μm. (C) A plot representing
mean GP values quantified at the plasma membrane region as a function of cell size in H99
cells.(D)Cell body sizes of titan daughters from live imaging of titan cells budding(n=30). (E) Cell
body sizes of all indicate strains after capsule induction(n>100). [49] Flow cytometry for DNA
content of mating fronts from the indicated isolates crossed with KN99**a**.1C and 2C populations were
identified relative to YPD-agar grown yeast cells from matched parent strains and MFI was recorded
for sub-1C, 1C, 2C peaks. "T" indicates pre-titanized cells.
746
Fig 7: *In vitro* titan cells interaction with host immune cells.
(A) Infection rates for the 4 clinical strains, showing the percentage of macrophages infected with at
least one internalized *C. neoformans* cell after 2h co-incubation. Data were assessed as not normal by
Shapiro-Wilk and analysed by Welch's ANOVA with Dunnett's test for multiple comparisons.

*p<0.02; ***p=0.0008. (B) Representative images showing macrophages (Red, Lysotracker) infected
with cryptococcal cells (Blue, CFW). Scale bar: 10 μm. (C) Macrophages infection rates infected with
only titan or typical cells (typical cells <10 um, titan cells > 11 um). Data were assessed for normality
by Shapiro-Wilk and analysed by Student's T-test with Welsh's correction for unequal variance
(determined by F-test). ***p=0.0002. (D) Representative images showing macrophages (Red)infected
with H99 yeast and titan cells. Fungi were co-incubated with anti-capsule antibody 18b7 plus anti-
mouse IgG-FITC (Green) and calcofluor white (CFW) for chitin (blue). Scale bar: 20μm. (**E-G**)
Macrophages and pre-titanized *C. neoformans* were co-incubated for 18 hours and assessed via live
imaging for (**E**) IPR, [49] vomocytosis, and (**G**) macrophage lysis. (**E**) IPR: mean intracellular
proliferation rate. Data were assessed for normality by Shapiro-Wilk and analysed by One-Way
ANOVA. [49] Mf death %: frequency of macrophage death indicated by cell lysis. Data were
assessed for normality by Shapiro-Wilk and analysed by One-Way ANOVA. *p=0.05; ***p<0.001.
(**G**) Vomo %: frequency of vomocytosis. Data were assessed as not normal by Shapiro-Wilk and
analysed by Kruskal-Wallis with Dunn's correction
Data for a representative biological replicate is presented (n>3). "T" indicates pre-titanized cells.
766
Fig 8. Titanizaiton and disease outcomes on mice model
(A) Cell body sizes of fungal cells collected from murine lungs (n>70 cells for each isolate). (B, C)
Infectious burdens for the indicated strains. Murine lungs (B) and brains (C) were assessed for
dissemination 10 days post-infection via colony forming units of total homogenates (n=5 per strain).
Data are presented as Log CFU/gram. (B) Data were assessed as normal by Shapiro-Wilk and
analysed by One-Way ANOVA with Holm-Sidak's multiple comparisons test. Data are presented as
Log CFU/gram. (C) Data were assessed as not normal by Shapiro-Wilk and analysed by Kruskal-
Wallis ANOVA with Dunn's correction for multiple comparisons. (D, E) Immune cell recruitment to
lungs for the indicated strains. Lung homogenates were analysed by FACS for the indicated cell
populations identified using the indicated markers for (D) Neutrophils (CD45+, CD11b+Ly6G+) or
(E) Alveolar Macrophages (CD45+ Ly6C- Ly6G- CD11b- SiglecF+). Data were assessed as normal
by Shapiro-Wilk and analysed by One-Way ANOVA with Holm-Sidak's multiple comparisons test.
779
Fig 9: Clinical isolates produce aneuploid daughter cells with altered drug resistance *in vivo*.
(A) Degree of heterogeneity in ploidy across the individual colony forming units recovered from host
tissue. 1C and 2C populations were identified relative to YPD-grown colonies from matched parent
strains and MFI was recorded for 1C and 2C peaks. Data are presented relative to median 1C MFI for
all colonies for a given isolate. Data were assessed as not normal by Shapiro-Wilk and analysed by
Kruskal-Wallis ANOVA compared to H99 with Dunn's correction for multiple comparisons. (B)
Fluconazole resistance relative to parent for individual colonies aggregated in (A). Data are plotted
relative to parent MIC and as a function of colony ploidy as assessed in (A). Data were assessed as
not normal by Shapiro-Wilk and analysed by Kruskal-Wallis ANOVA with Dunn's correction for
multiple comparisons.
790
791
Supplemental Figures
Fig S1. Supplemental information for DNA content analysis used Fig 3. (A) i) Histograms of
DAPI staining intensity for YPD grown cells, and 1C and 2C peaks are indicated based on the MFI.
ii)FSC-A vs DNA content scatter plots show the correlation of size increase with DNA content. (B)
Single cells were identified by FSC-A vs FSC-H to remove doublets (n>8000 after gating), then were
used for DNA content analysis (Left panel). FSC-A vs DNA content scatter plots show the correlation
of size increase with DNA content (Right panel). "T" indicates pre-titanized cells.
799
800 **Fig S2** (B) PAMPs staining flow cytometry analysis of induced cells from Fig 8 with large cells sub-
801 population highlighted in grey and MFI indicating exposure of Chitin (CFW, right); Mannan (ConA,
802 center); and Chitosan (EosinY, left). "T" indicates pre-titanized cells.
803
804 **Fig S3** (A) TEM of Zc8 titanides recovered from mice lungs, scale bar: 500nm. (B) Mating plates
805 from the indicated isolates crossed with KN99 MATa.

806
807 **Fig S4** Live imaging of CNV111, expressing Nde1-GFP and Cse4-mCherry in an H99 background.
808 Cells were incubated overnight in YNB, and then incubated in 10% FCS, 5% CO_2, 37°C at
809 OD_{600}=0.001 for 6 hours. Cells were immobilized in glass-bottom chamber wells using 1:200 dilution
810 of Crp127 and imaged using a Zeiss Axio-imager at 63x, 36°C, 5%CO_2. Images were acquired every
811 10 minutes. The movie is representative of >10 biological repeats.
812
813 **Fig S5**: DNA content of individual colony forming units recovered from host tissue, L-M1: Lung of
814 mice 1, B-M1: Brain of mice 1, dotted lines indicate median 1C MFI. n=24, except H99 samples from
815 brains(n=21) because of limitation of colonies recovered.
816
817

818 **References**
819

1.   Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM.
Estimation of the current global burden of cryptococcal meningitis among persons living with
HIV/AIDS. AIDS (London, England). 2009;23(4):525-30. doi:
10.1097/QAD.0b013e328322ffac.
2.   May RC, Stone NR, Wiesner DL, Bicanic T, Nielsen K. Cryptococcus: from
environmental saprophyte to global pathogen. Nat Rev Microbiol. 2016;14(2):106-17. doi:
10.1038/nrmicro.2015.6. PubMed PMID: 26685750; PubMed Central PMCID:
PMCPMC5019959.
3.   Denham ST, Brown JCS. Mechanisms of Pulmonary Escape and Dissemination by
Cryptococcus neoformans. J Fungi (Basel). 2018;4(1). doi: 10.3390/jof4010025. PubMed
PMID: 29463005; PubMed Central PMCID: PMCPMC5872328.
4.   Zaragoza O, Garcia-Rivera J, Nosanchuk JD, Cuenca-Estrella M, Rodriguez-Tudela J,
Casadevall A. Fungal Cell Gigantism during Mammalian Infection. PLoS Pathogens.
2010;6(6):1-18.
5.   Okagaki LH, Strain AK, Nielsen JN, Charlier C, Baltes NJ, Chrétien F, et al.
Cryptococcal cell morphology affects host cell interactions and pathogenicity. PLoS
Pathogens. 2010;6(6):e1000953. doi: 10.1371/journal.ppat.1000953.
6.   Zaragoza O, Nielsen K. Titan cells in Cryptococcus neoformans: cells with a giant
impact. Curr Opin Microbiol. 2013;16(4):409-13. doi: 10.1016/j.mib.2013.03.006. PubMed
PMID: 23588027; PubMed Central PMCID: PMCPMC3723695.
7.   Mukaremera L, Lee KK, Wagener J, Wiesner DL, Gow NAR, Nielsen K. Titan cell
production in Cryptococcus neoformans reshapes the cell wall and capsule composition
during infection. The Cell Surface. 2018;1(1):15-24. doi: 10.1016/j.tcsw.2017.12.001.
8.   Feldmesser M, Kress Y, Casadevall A. Dynamic changes in the morphology of
Cryptococcus neoformans during murine pulmonary infection. Microbiology.
2001;147(8):2355-65.
9.   Love GL, Boyd GD, Greer DL. Large Cryptococcus neoformans isolated from brain
abscess. J Clin Microbiol. 1985;22(6):1068-70. PubMed PMID: 3905847; PubMed Central
PMCID: PMCPMC271885.
10.   Cruickshank JG, Cavill R, Jelbert M. Cryptococcus neoformans of unusual
morphology. Appl Microbiol. 1973;25(2):309-12. PubMed PMID: 4121033; PubMed Central
PMCID: PMCPMC380794.
11.   Okagaki LH, Nielsen K. Titan cells confer protection from phagocytosis in
Cryptococcus neoformans infections. Eukaryot Cell. 2012;11(6):820-6. doi:
10.1128/EC.00121-12. PubMed PMID: 22544904; PubMed Central PMCID:
PMCPMC3370461.

12.   Gerstein AC, Fu MS, Mukaremera L, Li Z, Ormerod KL, Fraser JA, et al. Polyploid
titan cells produce haploid and aneuploid progeny to promote stress adaptation. MBio.
2015;6(5):e01340-15. doi: 10.1128/mBio.01340-15. PubMed PMID: 26463162; PubMed
Central PMCID: PMCPMC4620463.
13.   Fernandes KE, Brockway A, Haverkamp M, Cuomo CA, van Ogtrop F, Perfect JR, et
al. Phenotypic Variability Correlates with Clinical Outcome in Cryptococcus Isolates
Obtained from Botswanan HIV/AIDS Patients. MBio. 2018;9(5). doi: 10.1128/mBio.02016-
18. PubMed PMID: 30352938; PubMed Central PMCID: PMCPMC6199498.
14.   Dambuza IM, Drake T, Chapuis A, Zhou X, Correia J, Taylor-Smith L, et al. The
Cryptococcus neoformans Titan cell is an inducible and regulated morphotype underlying
pathogenesis. PLoS Pathog. 2018;14(5):e1006978. doi: 10.1371/journal.ppat.1006978.
PubMed PMID: 29775474; PubMed Central PMCID: PMCPMC5959070.
15.   Crabtree JN, Okagaki LH, Wiesner DL, Strain AK, Nielsen JN, Nielsen K. Titan cell
production enhances the virulence of Cryptococcus neoformans. Infect Immun.
2012;80(11):3776-85. doi: 10.1128/IAI.00507-12. PubMed PMID: 22890995; PubMed
Central PMCID: PMCPMC3486048.
16.   García-Rodas R, Casadevall A, Rodríguez-Tudela JL, Cuenca-Estrella M, Zaragoza O.
Cryptococcus neoformans Capsular Enlargement and Cellular Gigantism during Galleria
mellonella Infection. PLOS ONE. 2011;6(9):e24485. doi: 10.1371/journal.pone.0024485.
17.   Chrisman CJ, Albuquerque P, Guimaraes AJ, Nieves E, Casadevall A. Phospholipids
trigger Cryptococcus neoformans capsular enlargement during interactions with amoebae and
macrophages. PLoS Pathog. 2011;7(5):e1002047. doi: 10.1371/journal.ppat.1002047.
PubMed PMID: 21637814; PubMed Central PMCID: PMCPMC3102711.
18.   Wiesner DL, Specht CA, Lee CK, Smith KD, Mukaremera L, Lee ST, et al. Chitin
recognition via chitotriosidase promotes pathologic type-2 helper T cell responses to
cryptococcal infection. PLoS Pathog. 2015;11(3):e1004701. doi:
10.1371/journal.ppat.1004701. PubMed PMID: 25764512; PubMed Central PMCID:
PMCPMC4357429.
19.   Denham ST, Verma S, Reynolds RC, Worne CL, Daugherty JM, Lane TE, et al.
Regulated Release of Cryptococcal Polysaccharide Drives Virulence and Suppresses Immune
Cell Infiltration into the Central Nervous System. Infect Immun. 2018;86(3). doi:
10.1128/IAI.00662-17. PubMed PMID: 29203547; PubMed Central PMCID:
PMCPMC5820953.
20.   Okagaki LH, Wang Y, Ballou ER, O'Meara TR, Bahn Y-S, Alspaugh JA, et al.
Cryptococcal titan cell formation is regulated by G-protein signaling in response to multiple
stimuli. Eukaryotic Cell. 2011;10(10):1306-16. doi: 10.1128/EC.05179-11.
21.   Gish SR, Maier EJ, Haynes BC, Santiago-Tirado FH, Srikanta DL, Ma CZ, et al.
Computational Analysis Reveals a Key Regulator of Cryptococcal Virulence and
Determinant of Host Response. MBio. 2016;7(2):e00313-16. doi: 10.1128/mBio.00313-16.
PubMed PMID: 27094327; PubMed Central PMCID: PMCPMC4850258.
22.   Mukaremera L, McDonald TR, Nielsen JN, Molenaar CJ, Akampurira A, Schutz C, et
al. The Mouse Inhalation Model of Cryptococcus neoformans Infection Recapitulates Strain
Virulence in Humans and Shows that Closely Related Strains Can Possess Differential
Virulence. Infect Immun. 2019;87(5). doi: 10.1128/IAI.00046-19. PubMed PMID: 30833336;
PubMed Central PMCID: PMCPMC6479045.
23.   Gerstein AC, Jackson KM, McDonald TR, Wang Y, Lueck BD, Bohjanen S, et al.
Identification of Pathogen Genomic Differences That Impact Human Immune Response and
Disease during Cryptococcus neoformans Infection. mBio. 2019;10(4). doi:
10.1128/mBio.01440-19. PubMed PMID: 31311883; PubMed Central PMCID:
PMCPMC6635531.

24.   Gago S, Serrano C, Alastruey-Izquierdo A, Cuesta I, Martin-Mazuelos E, Aller AI, et
al. Molecular identification, antifungal resistance and virulence of Cryptococcus neoformans
and Cryptococcus deneoformans isolated in Seville, Spain. Mycoses. 2017;60(1):40-50. doi:
10.1111/myc.12543. PubMed PMID: 27633849.
25.   Desjardins CA, Giamberardino C, Sykes SM, Yu CH, Tenor JL, Chen Y, et al.
Population genomics and the evolution of virulence in the fungal pathogen Cryptococcus
neoformans. Genome Res. 2017;27(7):1207-19. doi: 10.1101/gr.218727.116. PubMed PMID:
28611159; PubMed Central PMCID: PMCPMC5495072.
26.   Vanhove M, Beale MA, Rhodes J, Chanda D, Lakhi S, Kwenda G, et al. Genomic
epidemiology of Cryptococcus yeasts identifies adaptation to environmental niches
underpinning infection across an African HIV/AIDS cohort. Mol Ecol. 2017;26(7):1991-
2005. doi: 10.1111/mec.13891. PubMed PMID: 27862555; PubMed Central PMCID:
PMCPMC5412878.
27.   Zaragoza O. Multiple Disguises for the Same Party: The Concepts of Morphogenesis
and Phenotypic Variations in Cryptococcus neoformans. Front Microbiol. 2011;2:181. doi:
10.3389/fmicb.2011.00181. PubMed PMID: 21922016; PubMed Central PMCID:
PMCPMC3167222.
28.   Alanio A, Vernel-Pauillac F, Sturny-Leclere A, Dromer F. Cryptococcus neoformans
host adaptation: toward biological evidence of dormancy. mBio. 2015;6(2). doi:
10.1128/mBio.02580-14. PubMed PMID: 25827423; PubMed Central PMCID:
PMCPMC4453510.
29.   Trevijano-Contador N, de Oliveira HC, García-Rodas R, Rossi SA, Llorente I,
Zaballos Á, et al. Cryptococcus neoformans can form titan-like cells in vitro in response to
multiple signals that require the activation of several transduction pathways. PLoS Pathog.
2018;14(5):e1007007. doi: 10.1371/journal.ppat.1007007.
30.   Choi J, Vogl AW, Kronstad JW. Regulated expression of cyclic AMP-dependent
protein kinase A reveals an influence on cell size and the secretion of virulence factors in
Cryptococcus neoformans. Mol Microbiol. 2012;85(4):700-15. doi: 10.1111/j.1365-
2958.2012.08134.x. PubMed PMID: 22717009.
31.   Hommel B, Mukaremera L, Cordero RJB, Coelho C, Desjardins CA, Sturny-Leclere
A, et al. Titan cells formation in Cryptococcus neoformans is finely tuned by environmental
conditions and modulated by positive and negative genetic regulators. PLoS Pathog.
2018;14(5):e1006982. doi: 10.1371/journal.ppat.1006982. PubMed PMID: 29775480;
PubMed Central PMCID: PMCPMC5959062.
32.   Dylag M, Colon-Reyes RJ, Kozubowski L. Titan cell formation is unique to
Cryptococcus species complex. Virulence. 2020;11(1):719-29. doi:
10.1080/21505594.2020.1772657. PubMed PMID: 32498590.
33.   Molloy SF, Kanyama C, Heyderman RS, Loyse A, Kouanfack C, Chanda D, et al.
Antifungal Combinations for Treatment of Cryptococcal Meningitis in Africa. N Engl J Med.
2018;378(11):1004-17. doi: 10.1056/NEJMoa1710922. PubMed PMID: 29539274.
34.   Zhou X, Ballou ER. The Cryptococcus neoformans Titan Cell: From In Vivo
Phenomenon to In Vitro Model. Current Clinical Microbiology Reports. 2018;5(4):252-60.
doi: 10.1007/s40588-018-0107-9.
35.   Mylonakis E, Moreno R, El Khoury JB, Idnurm A, Heitman J, Calderwood SB, et al.
Galleria mellonella as a model system to study Cryptococcus neoformans pathogenesis.
Infect Immun. 2005;73(7):3842-50. doi: 10.1128/IAI.73.7.3842-3850.2005. PubMed PMID:
15972469; PubMed Central PMCID: PMCPMC1168598.
36.   Garcia-Rodas R, Casadevall A, Rodriguez-Tudela JL, Cuenca-Estrella M, Zaragoza O.
Cryptococcus neoformans capsular enlargement and cellular gigantism during Galleria

mellonella infection. PLoS One. 2011;6(9):e24485. doi: 10.1371/journal.pone.0024485.
PubMed PMID: 21915338; PubMed Central PMCID: PMCPMC3168503.
37. Mylonakis E, Moreno R, El Khoury JB, Idnurm A, Heitman J, Calderwood SB, et al.
Galleria mellonella as a model system to study Cryptococcus neoformans pathogenesis.
Infection and Immunity. 2005;73(7):3842-50. doi: 10.1128/IAI.73.7.3842-3850.2005.
38. Erwig LP, Gow NA. Interactions of fungal pathogens with phagocytes. Nat Rev
Microbiol. 2016;14(3):163-76. doi: 10.1038/nrmicro.2015.21. PubMed PMID: 26853116.
39. Probert M, Zhou X, Goodall M, Johnston SA, Bielska E, Ballou ER, et al. A
Glucuronoxylomannan Epitope Exhibits Serotype-Specific Accessibility and Redistributes
towards the Capsule Surface during Titanization of the Fungal Pathogen Cryptococcus
neoformans. Infection and Immunity. 2019;87.
40. Owen DM, Rentero C, Magenau A, Abu-Siniyeh A, Gaus K. Quantitative imaging of
membrane lipid order in cells and organisms. Nature Protocols. 2012;7(1):24-35. doi:
10.1038/nprot.2011.419.
41. Delobel P, Tesniere C. A simple FCM method to avoid misinterpretation in
Saccharomyces cerevisiae cell cycle assessment between G0 and sub-G1. PLoS One.
2014;9(1):e84645. doi: 10.1371/journal.pone.0084645. PubMed PMID: 24392149; PubMed
Central PMCID: PMCPMC3879310.
42. Zhang J, Martinez-Gomez K, Heinzle E, Wahl SA. Metabolic switches from
quiescence to growth in synchronized Saccharomyces cerevisiae. Metabolomics.
2019;15(9):121. doi: 10.1007/s11306-019-1584-4. PubMed PMID: 31468142; PubMed
Central PMCID: PMCPMC6715666.
43. Zafar H, Altamirano S, Ballou ER, Nielsen K. A titanic drug resistance threat in
Cryptococcus neoformans. Curr Opin Microbiol. 2019;52:158-64. doi:
10.1016/j.mib.2019.11.001. PubMed PMID: 31765991.
44. Evans RJ, Li Z, Hughes WS, Djordjevic JT, Nielsen K, May RC. Cryptococcal
phospholipase B1 is required for intracellular proliferation and control of titan cell
morphology during macrophage infection. Infect Immun. 2015;83(4):1296-304. doi:
10.1128/IAI.03104-14. PubMed PMID: 25605772; PubMed Central PMCID:
PMCPMC4363446.
45. Robertson EJ, Najjuka G, Rolfes MA, Akampurira A, Jain N, Anantharanjit J, et al.
Cryptococcus neoformans ex vivo capsule size is associated with intracranial pressure and
host immune response in HIV-associated cryptococcal meningitis. J Infect Dis.
2014;209(1):74-82. doi: 10.1093/infdis/jit435. PubMed PMID: 23945372; PubMed Central
PMCID: PMCPMC3864387.
46. Hommel B, Mukaremera L, Cordero RJB, Desjardins CA, Sturny-Leclere A, Perfect
JR ea. Identification of environmental and genetic factors important for Cryptococcus
neoformans titan cell formation using new in vitro inducing conditions. PLoS Pathog. 2018.
47. Fernandes KE, Dwyer C, Campbell LT, Carter DA. Species in the Cryptococcus gattii
Complex Differ in Capsule and Cell Size following Growth under Capsule-Inducing
Conditions. mSphere. 2016;1(6). doi: 10.1128/mSphere.00350-16. PubMed PMID: 28066814;
PubMed Central PMCID: PMCPMC5196034.
48. Aslanyan L, Ekhar VV, DeLeon-Rodriguez CM, Martinez LR. Capsular specific IgM
enhances complement-mediated phagocytosis and killing of Cryptococcus neoformans by
methamphetamine-treated J774.16 macrophage-like cells. Int Immunopharmacol.
2017;49:77-84. Epub 2017/05/25. doi: 10.1016/j.intimp.2017.05.024. PubMed PMID:
28551495.
49. Trevijano-Contador N, Pianalto KM, Nichols CB, Zaragoza O, Alspaugh JA, Pirofski
L-a. Human IgM Inhibits the Formation of Titan-Like Cells in <span class="named-content

genus-species" id="named-content-1">Cryptococcus neoformans. Infection and
Immunity. 2020;88.
50. Litvack ML, Post M, Palaniyar N. IgM Promotes the Clearance of Small Particles and
Apoptotic Microparticles by Macrophages. PLOS ONE. 2011;6(3):e17223. doi:
10.1371/journal.pone.0017223.
51. Semighini CP, Averette AF, Perfect JR, Heitman J. Deletion of Cryptococcus
neoformans AIF ortholog promotes chromosome aneuploidy and fluconazole-resistance in a
metacaspase-independent manner. PLoS Pathogens. 2011;7(11):e1002364. doi:
10.1371/journal.ppat.1002364.
52. Gusa A, Williams JD, Cho JE, Averette AF, Sun S, Shouse EM, et al. Transposon
mobilization in the human fungal pathogen Cryptococcus is mutagenic during infection and
promotes drug resistance in vitro. Proc Natl Acad Sci U S A. 2020. doi:
10.1073/pnas.2001451117. PubMed PMID: 32303657.
53. Morrow CA, Lee IR, Chow EW, Ormerod KL, Goldinger A, Byrnes EJ, 3rd, et al. A
unique chromosomal rearrangement in the Cryptococcus neoformans var. grubii type strain
enhances key phenotypes associated with virulence. mBio. 2012;3(2). doi:
10.1128/mBio.00310-11. PubMed PMID: 22375073; PubMed Central PMCID:
PMCPMC3302566.
54. Hickman MA, Zeng G, Forche A, Hirakawa MP, Abbey D, Harrison BD, et al. The
'obligate diploid' Candida albicans forms mating-competent haploids. Nature.
2013;494(7435):55-9. doi: 10.1038/nature11865. PubMed PMID: 23364695; PubMed
Central PMCID: PMCPMC3583542.
55. Todd RT, Wikoff TD, Forche A, Selmecki A. Genome plasticity in Candida albicans
is driven by long repeat sequences. Elife. 2019;8. doi: 10.7554/eLife.45954. PubMed PMID:
31172944; PubMed Central PMCID: PMCPMC6591007.
56. Forche A, Abbey D, Pisithkul T, Weinzierl MA, Ringstrom T, Bruck D, et al. Stress
alters rates and types of loss of heterozygosity in Candida albicans. mBio. 2011;2(4). doi:
10.1128/mBio.00129-11. PubMed PMID: 21791579; PubMed Central PMCID:
PMCPMC3143845.
57. Forche A, Cromie G, Gerstein AC, Solis NV, Pisithkul T, Srifa W, et al. Rapid
Phenotypic and Genotypic Diversification After Exposure to the Oral Host Niche in Candida
albicans. Genetics. 2018;209(3):725-41. doi: 10.1534/genetics.118.301019. PubMed PMID:
29724862; PubMed Central PMCID: PMCPMC6028260.
58. Ashton PM, Thanh LT, Trieu PH, Van Anh D, Trinh NM, Beardsley J, et al. Three
phylogenetic groups have driven the recent population expansion of Cryptococcus
neoformans. Nature communications. 2019;10(1):2035. doi: 10.1038/s41467-019-10092-5.
PubMed PMID: 31048698; PubMed Central PMCID: PMCPMC6497710.
59. Litvintseva AP, Kestenbaum L, Vilgalys R, Mitchell TG. Comparative analysis of
environmental and clinical populations of Cryptococcus neoformans. J Clin Microbiol.
2005;43(2):556-64. doi: 10.1128/JCM.43.2.556-564.2005. PubMed PMID: 15695645;
PubMed Central PMCID: PMCPMC548066.
60. Kozubowski L, Yadav V, Chatterjee G, Sridhar S, Yamaguchi M, Kawamoto S, et al.
Ordered kinetochore assembly in the human-pathogenic basidiomycetous yeast Cryptococcus
neoformans. MBio. 2013;4(5):e00614-13. Epub 2013/10/03. doi: 10.1128/mBio.00614-13.
PubMed PMID: 24085781; PubMed Central PMCID: PMC3791896.
61. Smith LM, Dixon EF, May RC. The fungal pathogen Cryptococcus neoformans
manipulates macrophage phagosome maturation. Cell Microbiol. 2015;17(5):702-13. Epub
2014/11/15. doi: 10.1111/cmi.12394. PubMed PMID: 25394938.

62.   Narra R, Maeda JM, Temba H, Mghamba J, Nyanga A, Greiner AL, et al. Notes from
the Field: Ongoing Cholera Epidemic - Tanzania, 2015-2016. MMWR Morb Mortal Wkly
Rep. 2017;66(6):177-8. doi: 10.15585/mmwr.mm6606a5. PubMed PMID: 28207686.
63.   Casadevall A, Cleare W, Feldmesser M, Glatman-Freedman A, Goldman DL, Kozel
TR, et al. Characterization of a murine monoclonal antibody to Cryptococcus neoformans
polysaccharide that is a candidate for human therapeutic studies. Antimicrob Agents
Chemother. 1998;42(6):1437-46. PubMed PMID: 9624491; PubMed Central PMCID:
PMCPMC105619.
64.   Gilbert AS, Seoane PI, Sephton-Clark P, Bojarczuk A, Hotham R, Giurisato E, et al.
Vomocytosis of live pathogens from macrophages is regulated by the atypical MAP kinase
ERK5. Sci Adv. 2017;3(8):e1700898. doi: 10.1126/sciadv.1700898. PubMed PMID:
28835924; PubMed Central PMCID: PMCPMC5559206.
1065
1066

A

T-H99 T-Zc1 T-Zc8 T-Zc12

B

C i ii

T-Zc12

T-Zc8

T-Zc1

T-H99

Count

Size (FSC-A)

IgM-FITC

D

Crp127 IgM-FITC binding (MFI)				
	T-H99	T-Zc1	T-Zc8	T-Zc12
(% Neg)	(36.9)	(63.1)	(94.3)	(39.9)
Low (%)	420 (18.9)	528 (9.52)	220 (4.55)	345 (47.5)
Med (%)	1955 (23.8)	1693 (24.6)	1385 (0.07)	1134 (8.78)
High (%)	5601 (18.9)	4533 (1.33)	N/A	5234 (0.72)

A

B

H99

Zc1

$R^2=0.25$

$R^2=0.0006$

Zc8

Zc12

$R^2=0.016$

$R^2=0.008$

p=0.02

Ploidy Relative to Median

Ploidy Relative to Median

Fluconazole MIC relative to Parent

H99 lung, H99 Brain, Zc1 Lung, Zc1 Brain, Zc8 Lung, Zc8 Brain, Zc12 Lung, Zc12 Brain