Crystal and cryo-EM structures of the cytosolic G protein alpha chaperone and guanine

nucleotide exchange factor Ric-8A bound to Gαi1

Levi J. McClelland[1*], Kaiming Zhang[4 *], Tung-Chung Mou[1,3*], Jake Johnston[1,3], Cindee

Yates-Hansen[1], Shanshan Li[4], Celestine J. Thomas[1,†], Tzanko I. Doukov[6], Sarah

Triest[7,8], Alexandre Wohlkonig[7,8], Gregory G. Tall[9], Jan Steyaert[7,8], Wah Chiu[2,5] and

Stephen R. Sprang[1,2,3]

Affilliations

[1]Center for Biomolecular Structure and Dynamics, University of Montana, Missoula, MT,

USA

[2]Graduate Program in Biochemistry and Biophysics, University of Montana, Missoula,

MT, USA

[3]Division of Biological Sciences,University of Montana, Missoula, MT, USA

[4]Departments of Bioengineering and James H. Clark Center, Stanford University,

Stanford, CA, USA

[5]Biosciences Division, SLAC National Accelerator Laboratory, Stanford University,

Menlo Park, CA, USA

[6]Macromolecular Crystallography Group, Stanford Synchrotron Radiation Light Source,

SLAC National Accelerator Laboratory, Stanford University, Stanford, CA, USA

[7]Structural Biology Brussels, Vrije Universiteit Brussel (VUB), Brussels, Belgium.

[8]VIB-VUB Center for Structural Biology, VIB, Brussels, Belgium.

[9]Department of Pharmacology, University of Michigan Medical School, Ann Arbor, MI,

USA

† Present address, Regeneron Pharmaceutical, Inc. USA

Ric-8A is a cytosolic Guanine Nucleotide exchange Factor (GEF) that activates heterotrimeric G protein alpha subunits (Gα)[1]. Ric-8A is essential to life in multicellular eukaryotes by virtue of its chaperone activity that is required for Gα biogenesis and membrane localization[2,3]. Ric-8A adopts an armadillo (ARM)/HEAT repeat domain architecture and is structurally unrelated to G Protein-Coupled Receptors (GPCR)[4]. Both GEF and chaperone activities are stimulated by Casein Kinase II phosphorylation[5]. The mechanisms by which Ric-8A catalyzes GDP release and GTP binding to Gα, or exerts chaperone activity are unknown. Here, we report the structure of the nanobody-stabilized complex of nucleotide-free Gαi1 (isoform 1 of Gα family i) and phosphorylated Ric-8A at near atomic resolution by cryo-electron microscopy and X-ray crystallography. We find that Ric-8A envelops the GTPase domain of Gα, disrupting all three switch regions that convey Gα nucleotide-binding and signaling activity, and displaces the C-terminal helix and helical domain of Gα. These cooperative interactions dismantle the GDP binding site and promote GDP release, while protecting structural elements of Gα that are dynamic in the nucleotide-free state. The structures also show how *in vivo* phosphorylation stabilizes Gα-binding elements of Ric-8A, thereby enhancing its GEF and chaperone activities.

Genetic[6-8] and biochemical data[9] support a role for Ric-8A in G protein-coupled receptor (GPCR)-independent regulation of asymmetric cell division that is essential for embryonic development[10]. Ric-8A exhibits GEF and chaperone activity towards G protein alpha subunits (Gα) of the I, q and 12/13 classes[1], while Ric-8B, performs these functions for Gαs and Gαolf - each in a variety of cellular contexts[11,12]. That the Gα-class specificity of Ric-8A and Ric-8B is the same for GEF and chaperone activities[13] suggests a common mechanistic basis for both.

Biophysical data show that nucleotide-free Gαi1 is structurally dynamic when bound to

Ric-8A[14,15] and shares some properties with GPCR-bound Gα[16]. Namely, Ric-8A

induces rotational dynamics in the Gα helical domain[17] and binds to the C-terminus of

Gα [14,18]. In contrast to GPCRs, there is evidence that Ric-8A forms extensive

interactions with the Gα switch regions that undergo GTP-dependent conformational

changes[1,15,18,19] and with the Gα β-sheet scaffold[15].

To better understand the structural basis of Ric-8A GEF and chaperone activity, we

have determined the structure of the Ric-8A:Gαi1 complex, using both cryo-electron

microscopy (cryo-EM) and X-ray crystallography. To form the complex we used the N-

terminal 491-residue fragment of rat Ric-8A, which is a more active GEF then the full-

length (530 amino acid) protein, and, to reduce conformational heterogeneity, we used

the N-terminal 31-residue truncation mutant of rat Gαi1 (Δ31NGαi1), an efficacious

substrate for Ric-8A[14]. Recombinant Ric-8A(1-491) was phosphorylated by Casein

Kinase II at S435 and T440, which is necessary and sufficient for stimulation of GEF

activity[5] (**Extended data Fig. 1**). Hereafter we refer to phosphorylated Ric-8A(1-491) as

Ric-8A and Δ31NGαi1 as Gα.

To stabilize, and limit the dynamics of the Ric-8A:Gα complex for crystallographic

and cryo-EM experiments, we developed a panel of camelid nanobodies (Nb)[20] that

specifically recognize either Gαi1, Ric-8A or the complex of the two. We formed a series

of Ric-8A:Gαi1:Nb complexes that included from one to four Nbs from this panel. The

stability of the complexes towards cryogenic vitrification and consequently, the quality

and resolution of cryo-EM reconstructions derived from them, was improved in step with

the number of Nbs in the complex. We determined the cryo-EM structure of a complex of

Ric-8A:Gαi1 with three Nbs bound to Ric-8A and one bound to the helical domain of Gα.

Together, these nanobodies do not significantly affect Ric-8A-GEF activity (**Extended**

Data Fig. 2). The structure was determined from 327,493 particles derived from 8,468

movie images, yielding a 3.9Å resolution reconstruction (**Fig. 1, Extended Data Figs. 3**

and 4 and Table 1).

Figure 1. Architecture of the Ric-8A:Gα:4Nb complex. a, Cryo-EM 3D reconstruction of Ric-8A:Gα:4Nb
is shown with Ric-8A colored *wheat*, Gα GTPase and Helical domains colored *cyan* and *green*,
respectively, and nanobodies colored *gray*. **b**, Annotated ribbon and cylinder drawing, colored as in **a**, but
with the β-strands of Gα rendered in *magenta*. (**C**) The Ric-8A:Gαi1:4Nb complex is shown with the
nanobodies removed and the loop segments of the Gα GTPase domain colored green. Segments of Ric-8A
that contact Gα are rendered in *dark brown*. Ric-8A residues 335-340, which include the two
phosphorylation sites are rendered in *red*.
88

We obtained crystals of Ric-8A:Gαi1 bound to the three Ric-8A-specific Nbs used to

generate the cryo-EM structure. Diffraction from these crystals was highly anisotropic

(**Extended Data Fig. 5**), extending to 4.6Å along *a**, and *b*,* and 3.3Å along *c**, affording

measurement of a 90% complete anisotropic dataset to 3.3Å resolution (**Extended Data**

Table 2).   Initial crystallographic phases were determined by molecular replacement

and subsequently used to fit the cryo-EM density map. Iterative cycles of model-building

and refinement utilized both cryo-EM and X-ray diffraction data to generate the final

models (**Extended Data Table S1 and S2**). In the following discussion, we use

prefixes "r" and "g" for residue and secondary structure identifiers of Ric-8A and $G\alpha$,

respectively. ARM/HEAT repeat helices of Ric-8A are designated according to their

position in the repeat: "A", "B" or "C" for ARM repeats or "a" and "b" for HEAT repeats,

followed by the sequence number of the repeat (1 through 9). We use the established

nomenclature for $G\alpha$ secondary structure [21-23]. Descriptions of the crystal structure of

Ric-8A:$G\alpha$ refer to chains A and B, the better ordered of the two complexes in the

asymmetric unit.

The cryo-EM reconstruction reveals Ric-8A residues 2-487 and the whole of $G\alpha$ with

the exception of the disordered linker (residues 50-76) between the Helical and GTPase

domains (**Fig. 1a and 1b and Extended Data Video 1**). The crystal structure of the Ric-

8A:$G\alpha$ complex also reveals continuous electron density for Ric-8A except for the

linkage between 422 and 430. Notably, residues C-terminal to the last HEAT repeat

(r430-r491) are disordered in structures of Ric-8A in which $G\alpha$ is absent[4,18]. In the

crystal structure, which lacks Nb1056, the helical domain of $G\alpha$i1 and its connections to

the GTPase domain, including much of $g\alpha1$ and all of switch I, are disordered.

Otherwise the cryo-EM and X-ray models of Ric-8A:$G\alpha$ are in good agreement, although

certain irregularly-structured and loop regions in both $G\alpha$ and Ric-8A show significant

divergence (**Extended Data Fig. 6**). Small angle X-ray scattering (SAXS)

measurements of the complex are consistent with the X-ray and cryo-EM structures

(**Extended Data Fig. 7**).

Figure 2. Interactions of Ric-8A with Gαi1. Three major Ric-8A:Gα contact surfaces and phosphorylation
sites are highlighted and labeled in the *green*, *yellow*, *pink* and *blue* overlays (upper left) and enlarged in
panels **a**-**d**. Axes adjacent to each panel label indicate the rotation applied to the overall schematic to
generate the panel view. **a,** Interaction between Gα β-sheet and the Ric-8A C-terminal ARM/HEAT repeat
helix rαA9 and reverse turn r451-r457 (rRT) from the crystal structure. Polypeptide backbones are rendered
as tubes, with diameter proportional to B-factor at Cα. Carbon atoms of Ric-8A and Gαi1 are rendered in
deep brown, and green, respectively, and oxygen and nitrogen atoms are respectively colored *red* and *blue*.
b, Acidic Ric-8A peptide with phosphorylated residues rpS335 and rpT440 bound to the positively charged
surface formed by the 8[th] and 9th Ric-8A ARM/HEAT repeats from the crystal structure. **c,** Interaction of
rα11 with gα3 and Switch II from cryo-EM. Putative hydrogen bonds (donor-acceptor contact < 3.5Å) are
depicted with *orange* dashed lines. **d,** Contacts between gα5 and residues in successive ARM/HEAT
repeats of Ric-8A, from the crystal structure.

Figure 3. Conformational changes induced in Gα by its interaction with Ric-8A. In panels **a-c**,
GDP·Pi-bound Gαi1 (PDB ID 1GIT), rendered in *gray* is superimposed on the crystal structure of Ric-8A-
bound Gα using the Cα atoms of Gαi1 residues 219-225 (gβ4), 262-270 (gβ5) and 318-325. **a,**
Conformational changes due to binding of Ric-8A αa9, αb9 and RT to Gαi1. **b,** Ric-8A-induced conformation
changes dismantle the Gα nucleotide binding site (see text). GDP from 1GIT is included as a stick model for
reference. **c,** Displacement of Switch II by r*α11*. The position of GDP bound to Gαi1·GDP is shown as a
stick model. **d,** Ric-8A-induced rotation of the helical domain away from the GTPase domain of Gα: *top*,
Gαi1·GDP (1GIT) rendered with helices as cylinders and β-strands as ribbons and the helical domain
colored green; atoms of GDP are rendered as spheres. *Bottom*, the cryo-EM – derived model of Ric-8A-
bound Gα with GDP from 1GIT shown as a reference point.

Binding of Ric-8A to the GTPase domain of Gα at three non-contiguous contact

surfaces bury more than 3200Å^2 of solvent-accessible surface area (**Fig. 2a,c,d**).

Together, these interactions destabilize the guanine nucleotide binding site (**Fig. 3**).   At

the core of the complex, Ric-8A αb9 (r401-r408) and the reverse turn r451-r457 (rRT)

interact with the gβ4-gβ6 strands of Gα (**Fig. 2a and Extended Data Figs 8a and b**).

Ric-8 αb9 occupies the site of gα5 in GDP-bound Gα, and consequently gα5 is ejected

from the concave surface of the Gα β-sheet (**Fig. 3a**). Ric-8A residues rY412, rA415,

rA416 and rL418 in rαb9 substitute for nonpolar residues of gα5 to stabilize the

hydrophobic surface of the Gα β-sheet core. These residues are conserved in both A

and B isoforms of vertebrate Ric-8 (**Extended Data Fig. 9a**) and form Van der Waals

interactions with residues in gβ4-gβ6 that are conserved in both Gαi and Gαs families

(**Extended Data Fig. 9b**). The rW415A mutation severely impairs GEF activity (**Fig. 4**),

although the Y412A mutation does not. At the same interface, steric interactions with

rαa9 lever the antiparallel gβ2-gβ3 hairpin away from the GTPase core (**Fig. 3a**), thereby

destabilizing and partially disordering gα1. This, as described below, triggers separation

of the Helical and GTPase domains of Gα.

The ~90° rotation of α5 away from the GTPase domain core (**Fig. 3a**) reconfigures

the "TATC" motif in the g(β6-α5) GDP purine binding loop (**Fig. 3b**). This structural

change perturbs the conserved NKKD motif (in gβ4-αG) that confers specificity for

guanosine nucleotides[24]. Displacement of these two loops dismantles the GDP purine

binding site. The hydrophobic contact between the N-terminus of gα4 and the Ric-8A

αB8-αC8 turn (**Extended Data Fig. 10)** is also important, as indicated by the impairment

of GEF activity resulting from the rL383E mutation (**Fig. 4**).

After its ejection from the Gα β–sheet, gα5 is accommodated in a broad trough

formed by helices rαb2 through rαB8 that line the concave surface of ARM/Heat

superhelix of Ric-8A (**Fig 2d and Extended Data Fig. 8g and h**), as observed also in

the structure of Ric-8A bound to the C-terminus of transducin[18]. The predominantly

hydrophobic amino acid residues of gα5 that interact with the Ric-8A trough (e.g. gF336,

gV339, gI343, gI344 and gL348) otherwise pack against the GTPase domain β-sheet of

nucleotide-bound Gα[25]. Most of the Ric-8A residues that contact gα5 are conserved in

Ric-8B (**Extended Data Fig. 9a**), whereas several of the Ric-8A-contacting residues in

gα5 are not conserved among Gα classes (**Extended Data Fig. 9b**).

Figure 4. Mutational analysis of selected residues at the Ric-8A:Gα interface. GDP-GTP
exchange rates were measured by the rate of tryptophan fluorescence increase upon addition of ΔN31Gαi1
to Ric-8A and GTPγS at final concentrations of 2μM Ric-8A, 1μM ΔN31Gαi1 and 10μM GTPγS. Rate
differences between wild-type Ric-8A and all of the Ric-8A mutants with the exception of Y412A (P<0.82),
are significant at $10^{-5} > p > 10^{-12}$. Horizontal bars represent means and 1 standard deviation.
178

Ric-8A phosphorylation acts as an entropic clamp to promote GEF activity. The Ric-

8A αb9 and RT segments that interact with the Gα β-sheet are connected by an

intrinsically disordered sequence, r430-r450[4] followed by rα10. The segment from r430-

r440 is rich in acidic amino acids, and binds within a basic groove formed by rαA8 (r344-

r358) and rαa9 (r401-r410) (**Fig. 2b**). Phosphorylated rpS435 and rpT440, which are

well-ordered (**Extended Data Fig. 8i-l**) and form multiple ion-pair interactions with

conserved lysine and arginine side chains in the Ric-8A electropositive groove, help to

immobilize the r430-450 connector and consequently stabilize rαb9 and rRT that interact

with gβ4-gβ6. Accordingly, we found earlier that charge→neutral mutations of rR345Q

and rK349A, which interact with rpS435 and adjacent acidic residues, reduce both basal

and phosphorylation-stimulated GEF activity of Ric-8A[4].

Ric-8A helix α11 (r471-r491), which is preceded by an elongated peptide (r458-r470)

that forms an extended arch over the α3-β5 loop of Gα, packs between Gα Switch II

and gα3 (**Fig. 2c, and Extended Data Fig 8c-f**). In the GTP-bound state of Gα, these

two elements form the Gα effector protein binding site[26]. rα11 occupies the position of

Switch II in the G protein heterotrimer[27], and also in Gαi1 bound to GTP analogs[25] and in

the Gαi1•GDP•Pi product complex[28] (**Fig. 3c**).   However, Switch II is disordered in

Gαi1•GDP and probably also in nucleotide-free Gα[22]. Ric-8A interactions with switch II

are functionally important, since mutations of two Gα-contacting residues in rα11, rE478

and L482, which are conserved in both Ric-8A isoforms, impair GEF activity (**Fig. 4**).

The conformations of Switch II observed in the X-ray and cryo-EM structures differ from

each other, and from those adopted in nucleotide-bound Gα[26] (**Extended Data Fig.**

8e,f). Remodeling of Switch II may be facilitated by Ric-8A-induced displacement of β2-

β3 (**Fig. 3a**) described above.

Displacement of gα5 and reorientation of gβ2-gβ3 eliminates a nexus of stabilizing

interactions with the C-terminal residues of gα1, which become disordered (**Fig 3a.**). As

a consequence, contacts between the GTPase and helical domains of Gα are disrupted.

As observed in the cryo-EM structure, the helical domain undergoes a clockwise rotation

of ~60° around an axis roughly aligned with αD of the Gα helical domain (**Fig. 3d**). By

this motion, a channel opens between the helical and GTPase domains, providing a path

of egress for the nucleotide. The magnitude of the rotation is less than that observed in

crystal and cryo-EM structures of GPCR:G protein complexes[16,29], or deduced from

double electron-electron resonance studies of Ric-8A:Gα[17], possibly because it is limited

by steric interactions between the nanobodies bound to Gα and Ric-8A. The variability in

the orientation of the helical domain is also suggested by normal-mode analysis of the

SAXS data (**Extended Data Fig. 7**). The coupled interactions between Ric-8A and Gαi1,

described above, induce profound conformational changes that dismantle the Gα

guanine nucleotide-binding site (**Fig. 3b and Extended Data Video 2**).

The basis for the Gα class selectivity of Ric-8 isoforms is not readily apparent.

The great majority of residues at the Gα contact residues are conserved in Ric-8A and

Ric-8B (**Extended Data Fig 9a**). Gα residues that interact with rαB9 and rRT are

conserved in the Gαi and Gαs classes. However, several residues in gα5 differ between

Gαs and Gαi classes (**Extended Data Fig. 9b**), and, as suggested[18], the latter may

engage in more productive interactions with Ric-8A than their Gαs counterparts.

The structure of the complex suggests elements of Ric-8A chaperone activity by

which it promotes folding and stabilization of nucleotide-free Gα[2]. It is remarkable that

the magnitude of conformational changes induced in Gαi1 by Ric-8A far exceed those

sufficient to effect GDP release in GPCR - G protein complexes[16,23] wherein gα5

undergoes a modest rotation and translation and switch II remains protected by Gβγ as

in the intact heterotrimer[27,30]. Rather, Ric-8A stabilizes nucleotide-free Gα subunits in

the absence of Gβγ[14]. Molecular dynamics simulations suggest that contacts between

gα5 and the Gα β-sheet are dynamic in nucleotide-free Gα[31]. Interactions with Ric-8A

would shield these hydrophobic surfaces from exposure in the cytosol. Switch II is also

a dynamic structure, even in the GDP-bound state of Gα[24] and would likewise be

stabilized by Ric-8A α11. Ric-8A(1-491) partially rescues Gαi1 biosynthesis in Ric-8A [-/-]

cells, but not that of Gαq, for which full-length Ric-8A is required[32]. Hence, the C-

terminal ~40 residues of Ric-8A, which harbor three of the five CKII phosphorylation

sites[5], must play a critical role in Ric-8A chaperone activity. Nevertheless, it is clear that

the mechanism by which Ric-8A stabilizes dynamic regions of the nucleotide-free Gα

GTPase domain also underlie its GEF activity.

The extensive interface between Ric-8A and Gα prompts the question how the two

proteins are able to dissociate upon binding of GTP to Gα, a process that is kinetically

facile in comparison to the overall exchange reaction (**Fig. 5a and Extended Data Fig.**

2). Although displaced slightly by steric conflict with rα11 (**Fig. 3b**), the conformation of

the P-loop is largely retained in the nucleotide-free Gα bound to Ric-8A (**Fig. 5b**), thus

providing a preformed platform for subsequent binding of GTP. Electrostatic repulsion

between the C-terminus of r11 and the γ-phosphate of GTP as modeled in **Fig. 5b**, could

promote its release from its binding site between gα3 and Switch II, allowing Switch II to

refold into its native GTP-bound conformation.   Disruption of Switch II interactions with

Ric-8A would restore the native structure of gβ2-gβ3, promote its interaction with gα1

and thereby destabilize the remaining interface of Ric-8A with the GTPase domain. The

dynamics that accompany Ric-8A binding to Gα•GDP and subsequent release of

Gα•GTP remain to be explored.

Figure 5. The P-loop of Gαi1 provides a pre-ordered binding site for GTP in the complex with Ric-
8A. a, Progress curve for GTP (10μM) binding to purified Ric-8A:Gαi1 complex (1μM) (*red*), relative to that
for nucleotide exchange at Gαi1•GDP (1μM). **b**, Superposition of a model of GTP (PDB ID 1CIP) onto the
nucleotide binding site of Ric-8A-bound Gαi1. Serine 47 of α1 is positioned for Mg^{+2} coordination (see text).

References

1   Tall, G. G., Krumins, A. M. & Gilman, A. G. Mammalian Ric-8A (synembryn) is a

heterotrimeric Galpha protein guanine nucleotide exchange factor. *J Biol Chem*

278, 8356-8362 (2003).

2   Chan, P., Thomas, C. J., Sprang, S. R. & Tall, G. G. Molecular chaperoning function

of Ric-8 is to fold nascent heterotrimeric G protein alpha subunits. *Proc Natl*

Acad Sci U S A **110**, 3794-3799, doi:10.1073/pnas.1220943110 (2013).

3   Gabay, M. *et al.* Ric-8 proteins are molecular chaperones that direct nascent G

protein alpha subunit membrane association. *Sci Signal* **4**, ra79, doi:4/200/ra79

[pii]

10.1126/scisignal.2002223 (2011).

4   Zeng, B. *et al.* Structure, Function, and Dynamics of the Galpha Binding Domain

of Ric-8A. *Structure*, doi:10.1016/j.str.2019.04.013 (2019).

5   Papasergi-Scott, M. M. *et al.* Dual phosphorylation of Ric-8A enhances its ability

to mediate G protein alpha subunit folding and to stimulate guanine nucleotide

exchange. *Sci Signal* **11**, doi:10.1126/scisignal.aap8113 (2018).

6   Miller, K. G. & Rand, J. B. A role for RIC-8 (Synembryn) and GOA-1 (G(o)alpha) in

regulating a subset of centrosome movements during early embryogenesis in

Caenorhabditis elegans. *Genetics* **156**, 1649-1660 (2000).

7   Couwenbergs, C., Spilker, A. C. & Gotta, M. Control of embryonic spindle

positioning and Galpha activity by C. elegans RIC-8. *Curr Biol* **14**, 1871-1876,

doi:S0960982204007456 [pii]10.1016/j.cub.2004.09.059 (2004).

8   Afshar, K. *et al.* RIC-8 Is Required for GPR-1/2-Dependent Galpha Function during

Asymmetric Division of C. elegans Embryos. *Cell* **119**, 219-230 (2004).

9   Woodard, G. E. *et al.* Ric-8A and Gi alpha recruit LGN, NuMA, and dynein to the

cell cortex to help orient the mitotic spindle. *Mol Cell Biol* **30**, 3519-3530,

doi:MCB.00394-10 [pii]10.1128/MCB.00394-10 (2010).

10   Tonissoo, T. *et al.* Nucleotide exchange factor RIC-8 is indispensable in

mammalian early development. *Dev Dyn* **239**, 3404-3415,

doi:10.1002/dvdy.22480 (2010).

11   Chan, P., Gabay, M., Wright, F. A. & Tall, G. G. Ric-8B is a GTP-dependent G

protein alphas guanine nucleotide exchange factor. *J Biol Chem* **286**, 19932-

19942, doi:M110.163675 [pii]10.1074/jbc.M110.163675 (2011).

12   Hinrichs, M. V., Torrejon, M., Montecino, M. & Olate, J. Ric-8: different cellular

roles for a heterotrimeric G-protein GEF. *J Cell Biochem* **113**, 2797-2805,

doi:10.1002/jcb.24162 (2012).

13   Papasergi, M. M., Patel, B. R. & Tall, G. G. The G protein alpha chaperone Ric-8 as

a potential therapeutic target. *Mol Pharmacol* **87**, 52-63,

doi:10.1124/mol.114.094664 (2015).

14   Thomas, C. J. *et al.* The nucleotide exchange factor Ric-8A is a chaperone for the

conformationally dynamic nucleotide-free state of Galphai1. *PLoS One* **6**, e23197,

doi:10.1371/journal.pone.0023197PONE-D-11-08669 [pii] (2011).

15   Kant, R., Zeng, B., Thomas, C. J., Bothner, B. & Sprang, S. R. Ric-8A, a G protein

chaperone with nucleotide exchange activity induces long-range secondary

structure changes in Galpha. *Elife* **5**, doi:10.7554/eLife.19238 (2016).

16   Rasmussen, S. G. *et al.* Crystal structure of the beta2 adrenergic receptor-Gs

protein complex. *Nature* **477**, 549-555, doi:nature10361

[pii]10.1038/nature10361 (2011).

17   Van Eps, N., Thomas, C. J., Hubbell, W. L. & Sprang, S. R. The guanine nucleotide

exchange factor Ric-8A induces domain separation and Ras domain plasticity in

Galphai1. *Proceedings of the National Academy of Sciences of the United States*

of America **112**, 1404-1409, doi:10.1073/pnas.1423878112 (2015).

18   Srivastava, D., Gakhar, L. & Artemyev, N. O. Structural underpinnings of Ric8A

function as a G-protein alpha-subunit chaperone and guanine-nucleotide

exchange factor. *Nat Commun* **10**, 3084, doi:10.1038/s41467-019-11088-x

(2019).

19   Nagai, Y., Nishimura, A., Tago, K., Mizuno, N. & Itoh, H. Ric-8B stabilizes the alpha

subunit of stimulatory G protein by inhibiting its ubiquitination. *J Biol Chem* **285**,

11114-11120 (2010).

20   Pardon, E. *et al.* A general protocol for the generation of Nanobodies for

structural biology. *Nat Protoc* **9**, 674-693, doi:10.1038/nprot.2014.039 (2014).

21   Lambright, D. G., Noel, J. P., Hamm, H. E. & Sigler, P. B. Structural determinants

for activation of the alpha-subunit of a heterotrimeric G protein. *Nature* **369**,

621-628 (1994).

22   Mixon, M. B. *et al.* Tertiary and quaternary structural changes in Gi alpha 1

induced by GTP hydrolysis. *Science* **270**, 954-960 (1995).

23   Flock, T. *et al.* Universal allosteric mechanism for Galpha activation by GPCRs.

Nature **524**, 173-179, doi:10.1038/nature14663 (2015).

24   Sprang, S. R. G protein mechanisms: Insights from structural analysis. *Annu Rev*

Biochem **66**, 639-678 (1997).

25   Coleman, D. E. *et al.* Structures of active conformations of Gi alpha 1 and the

mechanism of GTP hydrolysis. *Science* **265**, 1405-1412 (1994).

26   Sprang, S. R., Chen, Z. & Du, X. Structural basis of effector regulation and signal

termination in heterotrimeric Galpha proteins. *Adv Protein Chem* **74**, 1-65

(2007).

27   Wall, M. A. *et al.* The structure of the G protein heterotrimer Gi alpha 1 beta 1

gamma 2. *Cell* **83**, 1047-1058 (1995).

28   Berghuis, A. M., Lee, E., Raw, A. S., Gilman, A. G. & Sprang, S. R. Structure of the

GDP-Pi complex of Gly203-->Ala G_{ia1}: a mimic of the ternary product complex of

G_a-catalyzed GTP hydrolysis. *Structure* **4**, 1277-1290 (1996).

29   Westfield, G. H. *et al.* Structural flexibility of the G alpha s alpha-helical domain

in the beta2-adrenoceptor Gs complex. *Proc Natl Acad Sci U S A* **108**, 16086-

16091, doi:10.1073/pnas.1113645108 (2011).

30   Lambright, D. G. *et al.* The 2.0Å crystal structure of a heterotrimeric G protein.

342   *Nature* **379**, 311-319 (1996).

343   31   Dror, R. O. *et al.* SIGNAL TRANSDUCTION. Structural basis for nucleotide

344   exchange in heterotrimeric G proteins. *Science* **348**, 1361-1365,

345   doi:10.1126/science.aaa5264 (2015).

346   32   Oner, S. S. *et al.* Regulation of the G-protein regulatory-Galphai signaling

347   complex by nonreceptor guanine nucleotide exchange factors. *J Biol Chem* **288**,

348   3003-3015, doi:10.1074/jbc.M112.418467 (2013).

Methods

Preparation of crosslinked Ric-8A and Gαi1 Bis-sulphosuccinimidyl suberate

(BS3, Pierce-Thermo-Fisher Scientific) was dissolved in water to a concentration of 100

mM. K100 reagent (CovalX) is provided as a 2 mg/ml aqueous solution and contained a

proprietary mixture of inert carbon chain spacers (lengths between 8.8-13.2 Å),

separating 1-hydroxyl-7-azabenzotriozole groups. Ric-8A:Gαi1 complex, prepared as

described[1-3] was dialyzed into 20 mM phosphate buffer saline (PBS) pH 7.4, 1 mM DTT,

and final concentration adjusted to 20 mM in the same buffer. For each 100 ml of

complex, 20 ml of K100 (2mg/ml) and 25 ml of BS3 (100 mM) was added, incubated at

room temperature for 30 minutes and the reaction quenched by addition of 10 ml of 1 M

Tris pH 8.5. Samples were centrifuged at 14,000 rpm on a desk-top centrifuge for 10 min

at 4°C, to remove particulate matter, eluted though a tandem 5 ml HiTrap desalting

column (GE Healthcare) to remove excess cross-linking reagents. Proteins were eluted

in 20 mM PBS, concentrated to 0.3 mg/ml and 1 mM fresh DTT added as the final step.

Cross-linked samples were analyzed by Coomassie-stained SDS-PAGE and verified by

Western blotting using an anti-Ric-8A monoclonal antibody[4] and mouse Anti-Gαi1

monocolonal antibody (Enzo Lifesciences). The crosslinked preparation contained

cross-linked Ric-8A:Gαi1, free Gαi1 and Ric-8A in roughly equal proportion.

Development of Camelid nanobodies Llamas (*Lama glama*) were immunized with

the cross-linked Ric-8A:Gαi1 preparation. Peripheral blood lymphocytes were isolated

from the immunized animals for extraction of total RNA, generation of cDNA and PCR

amplification to isolate cDNAs encoding heavy chain variable domains (VHH) for

subcloning into the pCTCON2 vector for display on the cell surface of *Saccharomyces*

cerevisiae EBY100 cells, as described[5-7]. Cells expressing nanobodies that selectively

recognize Gαi1, Ric-8A or the crosslinked Ric-8A:Gαi1 were identified by three-color

FACS sorting on a FACS AriaIII (BD Biosciences), using Dylight 505 (Pierce, Thermo

Scientific)-conjugated Ric-8A, Dylight 488-conjugated Gαi1 and R-phycoerythrin goat

anti-mouse antibody (Lucron) to label nanobody-expressing clones. Three rounds of

selection were conducted using 10mM fluorescent proteins in 20 mM PBS pH 7.4,

100mM NaCl and 2mM DTT. Nanobody sequences from selected clones yielding

nanobodies that exclusively bind to Gαi1, to Ric-8A either free or bound to Gαi1, or

exclusively to the Ric-8A:Gαi1 complex were subcloned in a pMESy4 vector for

periplasmic expression in *E. coli* WK6 cells as described[8]. Nanobody sequences encode

a C-terminal hexahistidine affinity tag.

Of the set of 27 nanobodies discovered in screening that show strong expression,

twenty were capable of binding to either free Ric-8A or to Ric-8A in complex with Gαi1

but not to Gαi1, three bound exclusively to the Ric-8A:Gαi1 complex and three bound to

Gαi1 or its complex with Ric-8A but not to Ric-8A alone. From this set was identified a

set of four nanobodies capable of binding simultaneously to Ric-8A:Gαi1, as determined

by co-elution as a stable hexameric complex from a GE Healthcare HiLoad 16/600

Superdex 200pg column and validated by mass spectrometry on a MicroFlex MALDI-

ToF MS(Brucker Microflex). Of this set, Nb8109, Nb8117 and Nb8119 bind to Ric-8A

and the Ric-8A:Gαi1 complex and Nb9156 binds to Gαi1.

Nanobody expression and purification   Nanobodies (Nb) 8109, 8117, 8119

and 9156 were encoded in pMESy4 vectors for periplasmic expression in WK6

Escherichia coli cells[7]. Briefly, overnight cultures were grown in 50 ml LB media

containing 100 µg/ml Ampicillin, 100 mM Glucose, and 1 mM $MgCl_2$ in shaker flasks at

190 rpm and 37 °C. After ~16 hours cells were pelleted at 4000 rpm (2200xg) for 10

minutes using a benchtop Sorvall Legend RT. Re-suspended pellets were added to 1 L

TB media containing 100 µg/ml Ampicillin, 5 mM glucose, and 1 mM $MgCl_2$ then

incubated at 190 rpm and 37 °C until achieving an OD_{600} of 0.7-1.0, at which point

temperature was lowered to 28 °C and growths were induced with 300 µM isopropyl β-

D-1-thiogalactopyranoside (IPTG). Approximately 16 hours post-induction, cells were

pelleted at 8000 rpm (12000xg) for 15 min in a Sorvall RC 6+ Centrifuge and cell pellets

were stored at – 80 °C. Pellets were re-suspended in 100 ml of TES buffer (0.2 M Tris

pH 8, 0.5 mM EDTA, 0.5 M sucrose) at 4 °C, vortexed and allowed to stir for a minimum

of 1 hour. Cells were then added dropwise to a 200 ml solution of TES/4 buffer (0.05 M

Tris pH 8, 0.125 mM EDTA, 0.125 M sucrose), then stirred for 1 hour at 4 °C. Lysate

was then spun down at 8000 rpm (12000xg) for 30 min and supernatant was loaded on

a gravity column with Ni-NTA agarose resin (Qiagen). Nanobody-bound resin was

washed with 20 column volumes of Wash 1 (0.05 M Tris pH 8, 1 M NaCl), 5 column

volumes of Wash 2 (0.05 M MES pH 6, 1 M NaCl), 10 column volumes of Wash 3 (0.05

M Tris pH 8, 0.5 M NaCl), and then eluted with 3 column volumes of Elution Buffer (0.05

M Tris pH 8, 0.2 M NaCl, 0.5 M imidazole).   Eluted nanobody was dialyzed overnight in

0.05 M Tris pH 8 and 0.2 M NaCl, then concentrated to 5-10 mg/mL before use or

storage at -80 °C. Before use all nanobodies were gel purified using a Superdex 200

10/300 GL size-exclusion column (GE Healthcare) in gel filtration buffer: 50 mM HEPES

pH 8, 150 mM NaCl, and 1 mM TCEP.

Complexes of Ric-8A, ΔN31Gαi1 and nanobodies Nb8109, Nb8117, Nb8119 in the

presence or absence of Nb9156 were formed by incubating Ric-8A, ΔN31Gαi1 and

nanobodies at a 1:2:2(per nanobody) molar ratio on ice for 16 hours and subsequently

purified by gel filtration chromatography using a GE Healthcare HiLoad 16/600 Superdex

200 pg in Gel Filtration Buffer.

Gα and Ric-8A protein expression and purification N-terminally glutathione-S-

transferase tagged Rat Gαi1 with a 31 residue N-terminal truncation (ΔN31Gαi1) was

expressed from a pDest15 vector in Bl21(DE3) RIPL *Escherichia coli* and purified as

previously described[1,9]. N-terminally hexahistidine tagged Rat Ric-8A(residues 1-491,

hereafter Ric-8A) was expressed and purified as previously described in a pET28a

vector with BL21(DE3) RIPL *Escherichia coli* cells[1,3,10]. Following removal of the

hexahistidine tag using Tobacco Etch Virus protease, Ric-8A protein was loaded onto a

Source 15Q column and eluted with a 500 mM NaCl gradient at 180 mM NaCl and

subsequently loaded on a GE Healthcare HiLoad 16/600 Superdex 200 column for size

exclusion chromatography in gel filtration buffer containing 50 mM Tris pH 8, 150 mM

NaCl and 1 mM Tris(2-carboxyethyl)phosphine (TCEP).

Ric-8A was phosphorylated using casein kinase II (New England Biolabs) as

previously described[10,11]. Briefly, purified Ric-8A was mixed 1:1 with a 2X Kinase

Reaction Buffer containing 0.1 M HEPES pH 8, 0.1 M NaCl, 20 mM $MgCl_2$, 2 mM EGTA,

and 1 mM DTT. ATP was added to a final concentration of 5 mM. 160 Units of casein

kinase II were added per mg of Ric-8A, and the reaction was allowed to proceed for ~16

hours at room temperature. Phosphorylated protein was loaded onto a Source 15Q

column pre-equilibrated with 50 mM HEPES pH 8, 25 mM NaCl, and 2 mM β-ME, and

eluted from a 500 mM NaCl gradient at 210 mM NaCl. Phosphorylated Ric-8A exhibits

an approximately 2 mS/cm shift compared to un-phosphorylated Ric-8A (**Extended Data**

Fig. 1b). All procedures described hereinafter were conducted with phosphorylated Ric-

8A comprising residues 1-491 of the full-length protein and referred to as Ric-8A.

Myristoylated Gαi1 (mGαi1), used in nucleotide exchange assays, was prepared as

described previously[10]. Mutants of Ric-8A, used in nucleotide exchange assays, were

constructed via the QuikChange II XL Site-Directed Mutagenesis Kit (Agilent).

Expression, purification and phosphorylation proceeded as described above.

Guanine Nucleotide Exchange assays were conducted by measuring the change

in Gαi1 tryptophan fluorescence in the presence or absence of Ric-8A using either

ΔN31Gαi1 or mGαi1, as described[3]. All proteins were buffer exchanged and assays

were conducted in 50 mM HEPES pH 8, 150 mM NaCl, 10 mM MgCl$_2$, 1 mM

TCEP. Experiments were carried out at 20°C using a LS55 luminescence spectrometer

(Perkin Elmer) with 4 nm slit widths (excitation 295 nm, emission 345 nm). GEF activity

for Ric-8A and Ric-8A mutants were measured by preincubating 450 µl of Ric-8A and

GTPγS in a quartz fluorescent cuvette prior to addition of 50 µl of 10 µM mGαi1 for a

final concentration of 0.5 µM Ric-8A, 1 µM mGαi1, and 10 µM guanosine 5'-O-[gamma-

thio]triphosphate (GTPγS) in 500 µl.   GTPγS binding to size exclusion chromatography

purified complexes of Ric-8A: ΔN31Gαi1 with and without nanobodies were compared to

intrinsic nucleotide exchange of ΔN31Gαi1. GTPγS was added to preincubated

complexes for final concentrations of 1 µM complex (or ΔN31Gαi1) and 10 µM

GTPγS. For each assay, a minimum of 5 technical replicates were taken for each

sample; more replicates were performed if the series of assays was conducted over

several days, to control for changes in sample activity. Progress curves were fit to single

or double (If a slow binding phase was detected) exponential rate models using

SigmaPlot 7. Statistical significance of rate differences between reference and test

samples was determined by a two-tailed Student's t-test. Probability that differences are

derived from a random distribution is reported. All data points are shown in box plots

that show mean and standard deviation for each data set[12].

Crystallization of the ΔN31Gαi1:Ric-8A:3Nb complex Crystallization trials for the

Ric-8A:ΔN31Gαi1:Nb8109:Nb8117:Nb8119 complex were conducted by vapor diffusion

using commercially-available crystallization screening kits. Sitting drops were set on 96-

2 well INTELLI-PLATEs (Art Robbins Instruments) using a Gryphon crystallization robot

(Art Robbins Instruments) at 3-10 mg/mL protein complex at a 1:1 v/v ratio with

precipitation solution. Initial crystallization conditions were identified from hits on the

ShotGun screen (MD1-88 Molecular Dimensions). Further optimization was carried out

by grid screening variations in sodium malonate (Hampton Research) concentration and

pH and by crystal seeding by hanging drop on 24 well VDXm plates (Hampton

Research). Crystal seed stocks were prepared with the Seed Bead kit (Hampton

Research) in 1.4 M sodium malonate pH 6.9. 0.9 μl of protein stock was added to 0.6 μl

of reservoir and 0.3 μl of crystal seed stock and incubated at 12 °C for a minimum of 1-2

weeks. Optimal crystals were obtained from hanging drops containing 3.6 mg/ml Ric-

8A:ΔN31Gαi1:3Nb, 1.4 M sodium malonate pH 6.9 at 12 °C.

Crystallographic data collection and processing Diffraction data were measured

at the NSLS-II FXS beamline from LN$_2$ cryoprotected crystals measuring approximately

50μ X 50μ X 5μ employing a helical data collection mode[13], using X-rays of 0.9793

wavelength. A Si (111) double crystal monochromator was used to focus the X-ray

beam to a spot size of 1μ X 1μ at the sample. Data were measured in the Phi axis

rotation mode in 0.2° fames, at 0.06s/frame at a beam attenuation of 0.7. Diffraction

data were measured on an Eiger 16M detector at a 133 Hz frame rate. Due to

significant (> 0.5Å) differences in unit cell axis dimension between crystals, data from a

single crystal were selected for data processing and structure determination. In view of

the considerable anisotropy of diffraction (approximately 4.6Å along a^* and b^* and 3.3Å

along c^*), three separate data processing strategies were conducted using the

AutoPROC v1.0.2 software toolbox[14] for evaluation in subsequent model-building and

refinement steps. All three utilized XDS[15] for data indexing and initial integration. The

Standard Isotropic protocol uses SCALEA and TRUNCATE from the CCP4[16] to generate

isotropic data with a resolution cutoff (4.6Å) determined by the criteria $R_{pim} \geq 0.6$, $I/\sigma(I) \geq$

2.0 and $CC_{1/2} \geq 0.3$. The Extended Isotropic protocol employed POINTLESS and

AIMLESS scaling and analysis software[17] to generate scaled intensities extending to the

3.2Å resolution with $I/\sigma(I) \geq 1.2$. The Anisotropic Filtering protocol implemented in

STARANISO[18] generated a dataset that incorporates intensities within a locally

averaged value of $I/\sigma(I)$ to define an anisotropic diffraction cut-off surface (**Extended**

data Table 2).

Crystallographic model building and refinement All crystallographic calculations

were conducted using the PHENIX 1.1.6 software package[19] unless otherwise noted.

Initial crystallographic phases for the Ric-8A:ΔN31Gαi1:3NB complex were determined

by Molecular Replacement using atomic coordinates of Ric-8A (residues 1-426; PDB

ID:6NMG), the Ras domain of Gai1:GDP (PDB ID: 1BOF), and an anti-VGLUT

nanobody (PDB ID:5OCL) as search models. An initial atomic model, consisting of

coordinates for Ric-8A residues 1-426, residues 34-65 and 190-320 of ΔN31Gαi1 and

three nanobody backbone models were manually refit into a sigma-weighted 2mFo-DFc

map using Coot v0.8.6[20] and refined using phenix.refine. Completion and refinement of

the crystallographic model followed the following general strategy: 1) model-fitting to the

cryo-EM reconstruction in regions of the structure in which the path of the polypeptide

chain was not defined in (or differed substantially from) the 2mFo-DFc map followed by

real space refinement with secondary structure and geometry restraints using

phenix.real_space_refine[21]; 2) refitting of the latter to the 2mFo-DFc map and

subsequent refinement; 3) refinement of the cryo-EM map (see below) using the

crystallographic model as an alignment reference. In the regions in which the

polypeptide path observed in the cryo-EM-derived and crystallographic models diverged,

no attempt was made to bring them into agreement. Initial cycles of crystallographic

model building and refinement were carried out using Standard Isotropic or Extended

isotropic data sets derived from datasets derived from merging and scaling data from the

two most strongly diffracting crystals. The final rounds of fitting and refinement of the

complete model (excluding the C-terminus of Gα1, the Helical Domain, and Switch I,

residues 51-184) utilized an Anisotropic Extended dataset comprising data from the

single crystal that afforded the strongest diffraction and optimal merging statistics (see

Extended Data Table 2). A TLS model was applied during refinement using

phenix.refine. Model quality and correlation with the refined electron density were

performed using MolProbity[22]. The atomic coordinates for the refined crystallographic

model for the Ric-8A:ΔN31Gαi1:3Nb complex and associated structure factors are

deposited in the RCSB Protein Data Bank[23] (PDB ID 6TYL). Figures depicting atomic

models were rendered using PyMol version 2.3 (Schrodinger, LLC)

Cryo-EM data collection Three microliters of the Ric-8A:ΔN31Gαi1:4Nb complex at

0.4 mg/ml with 0.01% NP40 were applied onto glow-discharged 200-mesh R2/1

Quantifoil grids (Electron Microscopy Sciences). Grids were blotted for 4 s and rapidly

cryocooled in liquid ethane using a Vitrobot Mark IV (Thermo Fisher Scientific) at room

temperature and 100% humidity. The samples were screened using a Talos Arctica

cryo-electron microscope (Thermo Fisher Scientific) operated at 200 kV and then

imaged in a Titan Krios cryo-electron microscope (Thermo Fisher Scientific) with GIF

energy filter (Gatan) at a magnification of 130,000× (corresponding to a calibrated

sampling of 1.06 Å per pixel). Micrographs were recorded using EPU software (Thermo

Fisher Scientific) with a Gatan K2 Summit direct electron detector, where each image is

composed of 30 individual frames with an exposure time of 6 s and a dose rate of 11.5

electrons per second per $Å^2$. A total of 8,670 movie stacks were collected with a defocus

range of -1.5 to -3.5 μm.

Cryo-EM image processing All micrographs were motion-corrected using

MotionCor2[24] and the contrast transfer function (CTF) was determined using

CTFFIND4[25]. All particles were autopicked using the NeuralNet option in EMAN2 v 2.31

and further checked manually, yielding 768,736 particles from selected 8,468

micrographs. Particle coordinates were then imported to Relion v3.0.6[4], wherein multiple

rounds of 2D classification were performed to remove poor 2D class averages.

Meanwhile, ~ 20,000 particle images were selected to build an initial model using the

"ab-initio 3D" program in cryoSPARC v2[26]. A total of 676,130 particles were used for 3D

classification in Relion to remove the no-conforming particles. Next, five rounds of 3D

heterogeneous refinement, CTF refinement were performed in cryoSPARC to further

remove the poor 3D classes. The final 3D refinement with 327,493 particles was

performed in cryoSPARC using the "non-Uniform refinement" option with a soft mask of

the complex density and a 3.85Å resolution map was obtained. The resolution of the

final map was estimated according to the 0.143 FSC criterion. A 3.9Å low-pass filter was

to the final 3D map for better display (See more information in **Extended Data Fig. 2**

and Table I).

Cryo-EM model building The cryo-EM model for the Ric-8A:ΔN31Gαi1:4Nbs

complex was constructed with a refined crystallographic Ric-8A:ΔN31Gαi:3Nb model (in

which only the GTPase domain was modeled) and a model of the helical domain of Gαi1

in complex with Nb9156 from the crystal structure of Gαi:Nb9156 complex (unpublished

data). The final Ric-8A:ΔN31Gαi1:4Nb cryo-EM model was rebuilt after refinement of the

cryo-EM reconstruction using a Ric-8A:ΔN31Gαi1:4NB mask as described above, with

multiple rounds of manual refitting of the cryo-EM density map optimized refinement

using phenix.real_space_refine with secondary structure and geometry restraints[21].

Model quality was evaluated with MolProbity[22].

Small Angle X-ray Scattering data collection, analysis and modeling

SAXS was performed at BioCAT beamline 18ID at the Advanced Photon Source,

Chicago, with in-line size exclusion chromatography (SEC-SAXS) to separate sample

from aggregates and other contaminants. Protein sample (approximately ~5 mg/ml in

50mM HEPES, pH 8.0, 150mM NaCl and 1mM TCEP) was loaded onto a Superdex 200

Increase 10/300 GL column, which was run at 0.7 ml/min in an AKTA Pure FPLC (GE

Healthcare Life Sciences). After detection by an in-line UV monitor, the sample passed

through the SAXS flow cell, a 1.5mm ID quartz capillary with 10µm walls. Scattering

intensity was recorded using a Pilatus3 1M (Dectris) detector which was placed 3.67m

from the sample affording access to a range of momentum transfer (q) from 0.0065Å^{-1}

to 0.35Å^{-1} [$q=4\pi\sin(\theta)/\lambda$]. Exposures of 0.5 s were acquired every 2 seconds during

elution. Data were reduced using BioXTAS RAW v1.6.0[27]. Buffer blanks were created

by averaging regions flanking the elution peak and subtracted from exposures selected

from the elution peak to create the I(q) vs q curves used for subsequent analyses. *Ab*

initio molecular envelopes were computed by the ATSAS v2.6.2 package[28] programs

DAMMIN[29]. Ten bead models were reconstructed in DAMMIF[30], which were aligned and

averaged in DAMAVER[31] (Volkov, 2003). The Molecular envelope was visualized, and

atomic models fit to molecular envelopes using Chimera v1.10.2[32]. The conformational

flexibility of Ric-8A:ΔN31Gαi1 was modeled by coarse-grained fitting with respect to

experimental SAXS data using SREFLEX program in the ATSAS software package[28].

Normal mode analysis was conducted with automatic determination of rigid body units.

The final disposition of rigid body units after application of normal mode projections was

determined by rigid body refinement with respect to the computed SAXS profile[33].

CRYSOL software from the ATSAS software package was used to model scattering

profiles from atomic coordinates.

Amino acid sequence alignments were conducted using Clustal Omega[34] *via* its

web-server (https://www.ebi.ac.uk/Tools/msa/clustalo/).

Data Availability Coordinates of Ric-8A:ΔN31Gαi1:4Nb model from cryo-EM are

deposited in the RCSB Protein Data Bank (PDB) with ID 6UKT. Coordinates of Ric-

8A:ΔN31Gαi1:3Nb model from crystal structure are deposited in the PDB with ID 6YTL.

The cryo-EM reconstruction is deposited in the Electron Microscopy Data Bank

(https://www.ebi.ac.uk/pdbe/emdb/) with id EMD20812. The small angle X-ray scattering

data for the Ric-8A:ΔN31Gαi1 complex is deposited in the Small Angle Scattering

Biological Database (SASBDB)[35] with accession code SASDG95.

Acknowledgements

This research was supported by NIH R01GM105993 (S.R.S); NIH P41GM103832, R01GM079429, and S10OD021600 (W.C.); NSF 1738547 (T-C.M); NIH R01-GM088242 (G.G.T.); resources of Instruct-ERIC, part of the European Strategy Forum on Research infrastructures (ESFRI), the Research Foundation-Flanders (FWO) for their support to Nanobody discovery, and the Strategic Research Program (SRP) of the Vrije Universiteit Brussel (E.P. and J.S.). The Integrated Structural Biology Core at the University of Montana Center for Biomolecular Structure and Dynamics is supported by NIH COBRE award P20GM103546 (S.R.S.). We thank Drs. Wuxian Shi and Martin Fuchs at the National Synchrotron Light Source II (NSLS II) for assistance with data collection at the FMX (17-ID-2) beamline, supported by NIH grant P41GM111244 and the Department of Energy (DOE), KP1605010. Z.D. is supported by the SSRL Structural Molecular Biology Program by the DOE and by NIH grant P41GM103393. Small angle X-ray scattering experiments were conducted at the Advanced Photon Source, operated for the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357 with support of NIH grant P41 GM103622 and 1S10OD018090-01 for purchase of the Pilatus 3 1M detector. We thank Dr. Baisen Zeng for assistance with GEF assays and SPR quantitation of Nb-Ric-8A binding.

Author Contributions

L.J.M. prepared and crystallized the Ric-8A:Gαi1:3Nb complex, assisted with X-ray data collection, conducted mutagenesis studies and analyzed kinetic data.

K.Z. carried out image acquisition and processing, generated and refined cryo-EM reconstructions.

T-C.M. fit and refined atomic models to cryo-EM and crystallographic data and

conducted SAXS analysis.

J.J. prepared and screened samples for cryo-EM data collection,and assisted with data

analysis.

C. Y-H. Conducted crystallization experiments and assisted with protein preparations.

S.L. assisted with cryo-EM data collection and processing.

C.J.T. prepared samples for nanobody development and conducted initial nanobody

characterization.

T.I.Z. advised on with X-ray data collection and processing.

S.T. and A.W. generated nanobodies.

G.G.T. provided general consultation on project.

J.S. oversaw nanobody development.

W.C. oversaw cryo-EM studies.

S.R.S. designed project, provided overall supervision, and wrote the manuscript with

assistance from co-authors.

Competing interests statement

The authors state no conflicting interests.

Supplementary Information

Supplementary Information is available for this paper

Correspondence

Correspondence and requests for materials should be addressed to Stephen Sprang:

Stephen.sprang@umontana.edu

Methods References

Thomas, C. J. *et al.* The nucleotide exchange factor Ric-8A is a chaperone for the conformationally dynamic nucleotide-free state of Galphai1. *PLoS One* **6**, e23197, doi:10.1371/journal.pone.0023197 PONE-D-11-08669 [pii] (2011).

Van Eps, N., Thomas, C. J., Hubbell, W. L. & Sprang, S. R. The guanine nucleotide exchange factor Ric-8A induces domain separation and Ras domain plasticity in Galphai1. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 1404-1409, doi:10.1073/pnas.1423878112 (2015).

Kant, R., Zeng, B., Thomas, C. J., Bothner, B. & Sprang, S. R. Ric-8A, a G protein chaperone with nucleotide exchange activity induces long-range secondary structure changes in Galpha. *Elife* **5**, doi:10.7554/eLife.19238 (2016).

Chan, P., Thomas, C. J., Sprang, S. R. & Tall, G. G. Molecular chaperoning function of Ric-8 is to fold nascent heterotrimeric G protein alpha subunits. *Proc Natl Acad Sci U S A* **110**, 3794-3799, doi:10.1073/pnas.1220943110 (2013).

Chao, G. *et al.* Isolating and engineering human antibodies using yeast surface display. *Nat Protoc* **1**, 755-768, doi:10.1038/nprot.2006.94 (2006).

Young, C. L., Britton, Z. T. & Robinson, A. S. Recombinant protein expression and purification: a comprehensive review of affinity tags and microbial applications. *Biotechnol J* **7**, 620-634, doi:10.1002/biot.201100155 (2012).

Pardon, E. *et al.* A general protocol for the generation of Nanobodies for structural biology. *Nat Protoc* **9**, 674-693, doi:10.1038/nprot.2014.039 (2014).

Abskharon, R. N. *et al.* Probing the N-terminal beta-sheet conversion in the crystal structure of the human prion protein bound to a nanobody. *J Am Chem Soc* **136**, 937-944, doi:10.1021/ja407527p (2014).

684   9   Thomas, C. J., Tall, G. G., Adhikari, A. & Sprang, S. R. Ric-8A catalyzes guanine

685   nucleotide exchange on G alphai1 bound to the GPR/GoLoco exchange inhibitor

686   AGS3. *J Biol Chem* **283**, 23150-23160 (2008).

687   10   Zeng, B. *et al.* Structure, Function, and Dynamics of the Galpha Binding Domain

688   of Ric-8A. *Structure*, doi:10.1016/j.str.2019.04.013 (2019).

689   11   Yu, W., Yu, M., Papasergi-Scott, M. M. & Tall, G. G. Production of

690   Phosphorylated Ric-8A proteins using protein kinase CK2. *Protein Expr Purif*

691   **154**, 98-103, doi:10.1016/j.pep.2018.10.002 (2019).

692   12   Weissgerber, T. L. *et al.* Data visualization, bar naked: A free tool for creating

693   interactive graphics. *J Biol Chem* **292**, 20592-20598,

694   doi:10.1074/jbc.RA117.000147 (2017).

695   13   Polsinelli, I. *et al.* Comparison of helical scan and standard rotation methods in

696   single-crystal X-ray data collection strategies. *J Synchrotron Radiat* **24**, 42-52,

697   doi:10.1107/S1600577516018488 (2017).

698   14   Vonrhein, C. *et al.* Data processing and analysis with the autoPROC toolbox.

699   *Acta Crystallogr D Biol Crystallogr* **67**, 293-302,

700   doi:10.1107/S0907444911007773 (2011).

701   15   Kabsch, W. Xds. *Acta Crystallogr D Biol Crystallogr* **66**, 125-132,

702   doi:10.1107/S0907444909047337 (2010).

703   16   Collaborative Computational Project, N. The CCP4 suite: programs for protein

704   crystallography. *Acta Crystallographica* **D50**, 760-763 (1994).

705   17   Evans, P. R. & Murshudov, G. N. How good are my data and what is the

706   resolution? *Acta Crystallogr D Biol Crystallogr* **69**, 1204-1214,

707   doi:10.1107/S0907444913000061 (2013).

18   Tickle, I. J. *et al.* STARANISO. (2018).

19   Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for

macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-

221, doi:10.1107/S0907444909052925 (2010).

20   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development

of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501,

doi:10.1107/S0907444910007493 (2010).

21   Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and

crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544,

doi:10.1107/S2059798318006551 (2018).

22   Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-

atom structure validation. *Protein Sci* **27**, 293-315, doi:10.1002/pro.3330 (2018).

23   Berman, H. M. *et al.* The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242.

(2000).

24   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion

for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332,

doi:10.1038/nmeth.4193 (2017).

25   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from

electron micrographs. *J Struct Biol* **192**, 216-221, doi:10.1016/j.jsb.2015.08.008

(2015).

26   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC:

algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods*

14, 290-296, doi:10.1038/nmeth.4169 (2017).

27   Hopkins, J. B., Gillilan, R. E. & Skou, S. BioXTAS RAW: improvements to a free

open-source program for small-angle X-ray scattering data reduction and

analysis. *J Appl Crystallogr* **50**, 1545-1553, doi:10.1107/S1600576717011438

(2017).

28   Franke, D. *et al.* ATSAS 2.8: a comprehensive data analysis suite for small-angle

scattering from macromolecular solutions. *J Appl Crystallogr* **50**, 1212-1225,

doi:10.1107/S1600576717007786 (2017).

29   Svergun, D. I. Restoring low resolution structure of biological macromolecules

from solution scattering using simulated annealing. *Biophys J* **76**, 2879-2886

(1999).

30   Franke, D. & Svergun, D. I. DAMMIF, a program for rapid ab-initio shape

determination in small-angle scattering. *J Appl Crystallogr* **42**, 342-346,

doi:10.1107/S0021889809000338 (2009).

31   Volkov, V. V. & Svergun, D. I. Uniqueness of ab initio shape determination in

small-angle scattering. *J Appl Crystallogr* **36**, 860-864 (2003).

32   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory

research and analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084

(2004).

33   Panjkovich, A. & Svergun, D. I. Deciphering conformational transitions of proteins

by small angle X-ray scattering and normal mode analysis. *Phys Chem Chem*

Phys **18**, 5707-5719, doi:10.1039/c5cp04540a (2016).

34   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple

sequence alignments using Clustal Omega. *Mol Syst Biol* **7**, 539,

doi:10.1038/msb.2011.75 (2011).

35   Valentini, E., Kikhney, A. G., Previtali, G., Jeffries, C. M. & Svergun, D. I.

SASBDB, a repository for biological small-angle scattering data. *Nucleic Acids*

Res **43**, D357-363, doi:10.1093/nar/gku1047 (2015).

759
760

Extended Data Table 1
Cryo-EM Data Collection and Refinement Statistics

Data collection and processing	Ric-8A(491):Δ31NGαi1:4Nb
Microscope	Titan Krios
Voltage (kV)	300
Camera	Gatan K2 Summit
Pixel size (Å)	1.06
Total Dose (e-/Å^2)	69
Defocus range (μm)	-1.5 to -3.5
Number of micrographs	8,670
Number of initial particles	768,736
Symmetry	C1
Number of final particles	327,493
Resolution (0.143 FSC, Å)	3.90
Atomic model refinement	
Software	phenix.real_space_refine
Clashscore, all atoms	8.79
Poor rotamers (%)	0.57
Favored rotamers (%)	99.43
Ramachandran favored (%)	84.24
Ramachandran allowed (%)	15.76
MolProbity score	2.15
Bond RMSD (Å)	0.006
Angle RMSD (Å)	1.005

Crystallographic Data Collection and Refinement Statistics

Data collection	Ric-8A(491):Δ31NGαi1:3NB		
Software	AutoPROC/STARANISO	XDS/AIMLESS	SCALA
Reflection scaling method[#]	Anisotropic filtered (extended)	Isotropic (extended)	Isotropic (standard)
Wavelength (Å)	0.979	0.979	0.979
Resolution range (Å)*	39.58 - 3.3 (3.9 - 3.3) [a*, b*=4.6, c*=3.3]	39.65 - 3.3 (3.9 - 3.3)	39.65 - 4.6 (4.7 - 4.6)
Space group	P 2$_1$	P 2$_1$	P 2$_1$
Unit cell dimensions			
a, b, c (Å)	93.0 144.7 114.4	93.0 144.7 114.4	93.0 144.7 114.4
β (°)	94.7	94.7	94.7
Total reflections*	73843 (3449)	160159 (62218)	58730 (2930)
Unique reflections*	20707 (1035)	44840 (17203)	16296 (796)
Redundancy*	3.6 (3.3)	3.6 (3.6)	3.6 (3.7)
Completeness [spherical] (%)*	45.8 (6.0)	99.1 (99.1)	99.1 (99.5)
Completeness [ellipsoidal] (%)*	90.1 (58.2)	98.3 (91.5)	
Mean I/σ (I)*	3.7 (1.7)	1.9 (0.3)	4.1 (2.2)
Wilson B-factor	91.88		
R_{meas}[†,*]	0.21 (0.70)	0.55 (4.65)	0.18 (0.54)
$R_{p.i.m.}$[†,*]	0.11 (0.38)	0.29 (2.42)	0.10 (0.28)
$CC_{1/2}$[†]	0.99 (0.70)	0.97 (0.17)	0.99 (0.86)
Refinement			
R_{work}[†,*]	0.248 (0.319)		
R_{free}[†,*]	0.287 (0.323)		
CC_{work}[†*]	0.907 (0.719)		
CC_{free}[†*]	0.866 (0.893)		
Number of total atoms			
protein	15938		
total protein residues	2021		
RMS deviations			
bond lengths (Å)	0.002		
bond angles (°)	0.55		
Ramachandran favored (%)[††]	89.16		
Ramachandran allowed (%)[††]	10.53		
Ramachandran outliers (%)[††]	0.30		
Rotamer outliers (%)[††]	0.0		
Clash Score[††]	10.32		
Average B-factor			
Macromolecules	109.97		
Number of TLS groups	1		

[#] The standard isotropic scaling method uses the classical isotropic treatment in truncate program where the resolution cut is determined by (1) Rpim ≤ 0.6. average I/σ(I) in a resolution bin ≥ 2.0, and $CC_{1/2}$ ≥ 0.3. The extended isotropic method includes all the measurements within the sphere of the highest observed resolution limit where the local [I/σ(I) ≥ 1.2; the extended anisotropic method uses the anisotropically-filtered reflections by STARANISO with the same resolution limit as the extended isotropic method. * Data for highest resolution shell are given in brackets. [†] $R_{meas}=\sum_{hkl}(n/n-1)^{1/2}\sum_i |I_i(hkl)-\langle I(hkl)\rangle|/ \sum_{hkl}\sum_i I_i(hkl)$, where $I_i(hkl)$ is the ith observation of the intensity of the reflection hkl and $\langle I_{hkl}\rangle$ is the mean over n observations. $R_{p.i.m.}=\sum_{hkl}(1/n-1)^{1/2}\sum_i |I_i(hkl)-\langle I(hkl)\rangle|/ \sum_{hkl}\sum_i I_i(hkl)$. [§] $R_{work}=\sum_{hkl} ||F_{obs}|-|F_{calc}||/ \sum_{hkl} |F_{obs}|$, where F_{obs} and F_{calc} are the observed and calculated structure-factor amplitudes for each reflection hkl. R_{free} was calculated for 5% of the diffraction data that were selected randomly and excluded from refinement. Correlation coefficients: $CC = S_i((x_i - \langle x\rangle)(y_i - \langle y\rangle))/(S_i((x_i - \langle x\rangle)^2(y_i - \langle y\rangle)^2)^{1/2}$, where x_i and y_i are the ith of n observations of quantities whose mean values are $\langle x\rangle$ and $\langle y\rangle$; for $CC_{1/2}$, x_i, and y_i correspond to intensity measurements derived from each of two randomly selected half-data sets from the set of unmerged data; For CC_{work} and CC_{free} x_i and y_i refer to observed structure factor amplitudes and structure factor amplitudes computed from the refined atomic model, respectively, for the working data set and the set used to compute R_{free}. [††]Calculated using MolProbity[39]

a
```
NB_8109  AQVQLQESGGGLEQAGDSLRLSCAASGLIVSNYAMGWFRQAPGKEREFVAYINWNGGVTY 60
NB_9156  -QVQLQESGGGLEQAGDSLRLSCAASGLIVSNYAMGWFRQAPGKEREFVAYINWNGGVTY 59
NB_8117  -QVQLQESGGGLVQPGGSLRLSCAASGIIFRSNGMAWYRQAPGKEREWVASITSFGDA-I 58
NB_8119  -QVQLQESGGGLVQAGGSLRLSCAASGGIVHISSMGWFRQAPGKQRELVATSPSNGDI-R 58
         ********** * *.********** *.  .*.*:*****:** **      *.

NB_8109  YTNSVKGRFTISRDNAKNTVYLQMNSLKPEDTAVYYCARTSRASVTTRVADFGYWGQGTQ 120
NB_9156  YTNSVKGRFTISRDNAKNTVYLQMNSLKPEDTAVYYCARTSRASVTTRVADFGYWGQGTQ 119
NB_8117  YRDSVKGRFTISRDNARNAVSLQTNSLKTEDTAVYYCNTYPV---------NSAWGQGTQ 109
NB_8119  YADSVKGRFTLSRDNAKNTVSLQMNSLEPEDTAVYYCHSFLRHTAS--ASYNNYYGQGTQ 116
         * :*******:*****:*:* ** ***: ********        . :*****

NB_8109  VTVSSHHHHHHEPEA 135
NB_9156  VTVSSHHHHHHEPEA 134
NB_8117  VTVSSHHHHHHEPEA 124
NB_8119  VTVSSHHHHHHEPEA 131
         **************
```

Extended Data Figure 1. Purification of Ric-8A:Gα:Nb complexes. a, Amino acid sequences of nanobodies used to form complexes. **b**, Anion exchange chromatography of phosphorylated (pRic-8A) and unphosphorylated (uRic-8A). **c**, Gel filtration of complexes at the following molar stoichiometric ratios: 1 Ric-8A:2 Gα:2 Nb 8109:2 Nb 8117:2 Nb 8119 (*red*) and 1 Ric-8A:2 Gα:2 Nb 8109:2 Nb 8117:2 Nb 8119:4 Nb 9156 (*blue*) molar stoichiometric ratios, Ric-8A and Ga correspond to the 1-491 residue construct of Ric-8A and the 31N-terminal truncation mutant of Gαi1, respectively, as described in the main text. (see Methods).

Extended Data Figure 2 Effect of nanobodies on GEF activity of Ric-8A. a, GDP-GTP exchange rates were measured by the rate of tryptophan fluorescence increase upon addition of ΔN31Gαi1 to Ric-8A and GTPγS at final concentrations of 2mM Ric-8A, 1mM ΔN31Gαi1 and 10mM GTPγS in the absence (control) or presence of Nb 8109 (4 mM), Nb 8117 (4 mM), Nb 8119 (4 mM) (3Nb) or the latter with the addition of (4 mM) Nb 9156 (4Nb). GTP binding rates were measured by tryptophan fluorescence increase upon addition of GTPγS (10 mM final concentration) to size exclusion chromatography-purified Ric-8A:ΔN31Gαi1:nNb (1 mM). Progress curves for the 3Nb and 4Nb GDP-GTPγS exchange and 4Nb GTPγS binding assays were fit to a single exponential rate, whereas data for controls and the 3Nb GTPγS binding assay were fit to a double exponential model, due to the presence of a slow kinetic phase, and the kinetic constant of the fast rate (k1) reported. In all cases average and standard deviations of measurement are reported for a minimum of 5 experimental replicates. **b**, Sample progress curves for the data reported in panel **a.**

Extended Data Figure 3. Single-particle cryo-EM of Ric-8A:Gα:4Nb complex and Work-flow of cryo-EM image processing. a, Representative motion corrected cryo-EM micrograph. **b**, Reference-free 2D Class averages computed in Relion. **c**, Work-flow for cryo-EM processing (see Methods). **d**, Gold-standard Fourier Shell Correlation plot for 3D reconstruction generated using cryoSPARK.

Extended Figure 4. Real-space refinement of Ric-8A:Gα :cryoEM model. Model-map correlation (filled circles) and B factors (open circles) are plotted with respect to residue number for the macromolecular components of the Ric-8A:Gα:4NB model derived from cryo-EM. Data were generated using the phenix.real_space_refine program from the PHENIX suite.

(0kl)   (h0l)   (hk0)

Extended Figure 5. Projections of diffraction intensities from crystals of the Ric-8A:Gα:3Nb complex along reciprocal cell axes and color-coded by I/σI. Data generated using the STARANISO sever (see Methods).

Extended Data Figure 6. Agreement of cryo-EM and X-ray structures of Ric-8A:Gαi1. a, Deviation between corresponding Cα atoms after least-squares superposition of Ric-8A coordinates. *Red* lines below residue numbers mark regions at which Ric-8A contacts Gai1 (see also ED Figure 10). The RMSD for all 484 common Cα positions is 1.81Å. The PyMOL align routine yields an RMSD of 1.03Å after rejection of 69 outliers. **b**, Deviation between Gαi1 atoms, computed per panel **a**. The RMSD for all 183 common Cα positions is 2.36Å; PyMol align yields an RMSD of 1.49Å after rejection of 21 outliers. **c,** Superposition of cryoEM (*magenta*) and X-ray (*green*) structures of Gαi1. Structural elements that show the largest structural differences are labeled. **d**, Superposition of Ric-8A models, as per panel **c**.

Extended Figure 7. Solution structure and Normal Mode refined model of Ric-8A:Δ31NGαi1 complex from Small-Angle X-ray scattering. a, Solution scattering profile for Ric-8A:ΔN31Gαi1 complex (*closed circles*). Inset shows, associate Guinier plot. CRYSOL and SRELEX from the ATSAS software package were used, respectively, to compute scattering profiles from atomic coordinates, and use normal-mode analysis to generate a refined atomic model. Computed scattering curves for cryo-EM-derived atomic coordinates of Ric-8A:ΔN31Gαi1 complex are shown in *green* and that from normal mode analysis is rendered in *blue*. The χ^2 for the fit of the computed scattering curve for the cryo-EM model is 5.48 and that for the normal-mode refined model is 1.24. The bottom panel shows the error-weighted residual difference plots $\Delta I/\sigma I = (I(q)_{experimental} - CI(q)_{model})/\sigma I(q)$ *vs q*, where C is a normalizing scale factor. **b,** SREFLEX-refined Ric-8A:ΔN31Gαi1 model fit to the *ab initio* molecular envelope calculated from the experimental SAXS scattering curve (*yellow*). **c**, Superposition of the SREFLEX-refined model of Ric-8A:ΔN31Gαi (Ric-8A, *brown*; GTPase domain of ΔN31GαI, *cyan*; Helical domain of ΔN31GαI, *green*) onto the cryo-EM model (*gray*), indicating a mode of helical domain rotation that may be preferred in solution in the absence of nanobodies. Models were superimposed using the Cα atomic positions of Ric-8A.

Extended Data Figure 8. X-ray and cryo-EM density at selected residue ranges. 2DFo-mFc electron density is contoured at 1.0 σ and cryo-electron density is contoured at a threshold of 3.5. Structural elements of Ric-8A and Gαi1 are labeled in bold italics, N- and C-terminal residue numbers of segments are shown underlined, selected residues are labeled. Carbon atoms in Ric-8A are colored *wheat* and in Gai1, *cyan*. Panels **a, c, e, g** and **i** show residues from crystallographic model with corresponding density. Panels **b, d, f, h** and **j** show the same from the cryo-EM model. Panels **k** and **l**, show the crystallographic model and associated density around phospho-serine 435 and phospho-threonine 440, respectively. The 5 s contour level is shown in *red*.

```
Ric8A_RAT     28 EHSQSFTF  71 RILSRDRSCL  121 LCNLVLSSPTAQ  160 VQFFDLRLLFLLTAL
Ric8A_MOUSE   28 EHSQSFTF  71 RILSRDRSCL  121 LCNLVLSSPTAQ  160 VQFFDLRLLFLLTAL
Ric8A_HUMAN   28 EHSQSFTF  71 RILSRDRNCL  121 LCNLVLSSPVAQ  160 VQFFDLRLLFLLTAL
Ric8A_BOVIN   28 ENSQSFTF  71 RILSRDRSCL  121 LCNLVLSSPVAQ  160 VQFFDLRLLFLLTAL
Ric8A_DANRE   28 EKSQCFSF  71 RILSRDKKSL  121 LCNIVFNSEAAQ  160 VRFFDLRLTFLITAL
Ric8A_CHICK   28 EKSQCFTF  71 RILSRDKYCL  121 LCNIVFSSPRAQ  160 VKFFDLRLLFLLTAL
Ric8A_XENTR   28 ENSQCFFF  71 RILSRDKHAL  122 LCNIIYNSVEAQ  161 SKFFDLRLLFLLTAL
Ric8B_DANRE   28 ENSRTFAF  71 RILSRDKHVL  127 LCNVVFNSVPAQ  164 VGLFSLRLLFLLSAL
Ric8B_XENTR   16 QNRTMFKF  59 RILSRDKKVL  109 LCNIVFNSMEAQ  148 IRCFALRLLFLLTLL
Ric8B_CHICK   30 ENRAVRF   73 RILSRDKKVL  123 LCNIIFNSAVAQ  162 IKCFDLRLLFLLSLL
Ric8B_HUMAN   38 KHRATFKF  71 RILSRDKKVL  121 LCNIVFNSQMAQ  160 IKCFDLRLLFLLSLL
Ric8B_BOVIN   28 KHRATFKF  71 RILSRDKKVL  121 LCNIVFNSQMAQ  160 IKCFDLRLLFVLSLL
Ric8B_MOUSE   28 KHRATFKF  71 RILSRDKKIL  121 LCNIVFNSQVAQ  160 IKCFDLRLLFVLSLL
Ric8B_RAT     28 KHRATFKF  71 RILSRDKKIL  121 LCNIVFNSQVAQ  160 IKCFDLRLLFVLSLL
                 ::  * *     ******:  *      ***::  -.*  **      * *** *:::  *
                    αb1          αb2               αC3              αC4

Ric8A_RAT    225 KVLFNITFDSV  267 TEEFHGHTVNLLG  320 LHQTHRLKECVAPVLSVLTE  361 DVRTRPEVGDLLR
Ric8A_MOUSE  225 KVLFNITFDSV  267 TEEFHGHTVNLLG  320 LHQTHRLKECVAPVLNVLTE  361 DVRTRPEVGDLLR
Ric8A_HUMAN  226 KVLFNITLDSI  268 TEEFHGHAVNLLG  321 LHKTHRLKESVAPVLSVLTE  362 DVRTRPEVGEMLR
Ric8A_BOVIN  225 KVLFNITFDSI  267 TEELHGHAVNLLG  320 LHQTHRLKESVAPVLSVLTE  361 DVRTRPEVGELLR
Ric8A_DANRE  237 KILFNITFDSN  280 TEEMHSHTVNLLG  333 LDRGNKLKETLLPSLNLLTE  374 DVKNRPEVGNALR
Ric8A_CHICK  233 KVLFNITFDSS  276 TEEFHGHTVNLLG  329 LDRGHKLKESLTPVLNLLTE  370 DVKNRPEVGNSLR
Ric8A_XENTR  233 KTLFNITFDIS  276 TEEFHGHTVNLLV  329 LDRGHKLRETLTPVLNLLTE  370 DVKNRPEVGNTLR
Ric8B_DANRE  237 KALFNLTMSDS  277 TEETHSHAINLLS  330 IDKGFNFKEGLTPVLSLLTE  371 DVKERPEIGDTIR
Ric8B_XENTR  220 KALFNVTLDSW  272 TEELHSNTVNLLS  315 IDKGCSYREGLTPVLSLLTE  356 DVSNRPEVGMTVR
Ric8B_CHICK  234 KALFNVTLDSW  276 TEELHSNAINLLS  330 IDKGSSYREGLTPVLSLLTE  361 DVSSRPEVGTTVR
Ric8B_HUMAN  232 KALFNVTVDSW  274 TEELHSNAVNLLS  353 IDKGSSYREGLTPVLSLLTE  304 DVTNRPEVGSTVR
Ric8B_BOVIN  232 KALFNVTVDSW  274 TEELHSNAVNLLS  353 IDKGSSYREGLTPVLSLLTE  304 DVTNRPEVGSTVR
Ric8B_MOUSE  232 KALFNVTVDSW  274 TEELHSNAVNLLS  353 IDKGSSYREGLTPVLSLLTE  304 DVTNRPEVGSTVR
Ric8B_RAT    232 KALFNVTVDSW  274 TEELHSNAVNLLS  353 IDKGSSYREGLTPVLSLLTE  304 DVTNRPEVGSTVR
                 * ***:*. .       *** *-:::***       :.:  - :* :-* *.:***      **  ***:*  :*
                    αC5               αb6                    αC7                  αA8

Ric8A_RAT    374 NKLVRLMTHLDTDVKRVAAEFLFVLCSESVPRFIKYTGYGNAAGLLAARGLMAGGRPEGQ 433
Ric8A_MOUSE  374 NKLVRLMTHLDTDVKRVAAEFLFVLCSESVPRFIKYTGYGNAAGLLAARGLMAGGRPEGQ 433
Ric8A_HUMAN  375 NKLVRLMTHLDTDVKRVAAEFLFVLCSESVPRFIKYTGYGNAAGLLAARGLMAGGRPEGQ 434
Ric8A_BOVIN  374 NKLVRLMTHLDTDVKRVAAEFLFVLCSESVPRFIKYTGYGNAAGLLAARGLMAGGRPEGQ 433
Ric8A_DANRE  387 NKLVRLMTHIDTDVKHCAAEFLFVLCKESVSRFIKYTGYGNAAGLLAARGLMRGGRDPGH 446
Ric8A_CHICK  383 NKLVRLMTHIDTDVKHCAAEFLFVLCKESVSRFVKYTGYGNAAGLLAARGLMAGGREEGE 442
Ric8A_XENTR  383 NKLVRLMTHVDTDVKHCAAEFLFVLCKENVSRFVKYTGYGNAAGLLAARGLLAGGRGEGR 442
Ric8B_DANRE  383 NKLVRLMTHVDMAVKQGAAEFLFVLCKESVDNLLKYTGYGNAAGLLMARGLLAGGRGETQ 443
Ric8B_XENTR  369 NKLVRLMTHVDLGVKQIAAEFLFVLCKERVDSLLKYTGYGNAAGLLAARGLLAGGRGDRW 428
Ric8B_CHICK  384 NKLVRLMTHVDLGVKQIAAEFLFVLCKERVDSLLKYTGYGNAAGLLAARGLLAGGRGDHW 443
Ric8B_HUMAN  407 NKLVRLMTHVDLGVKQIAAEFLFVLCKERVDSLLKYTGYGNAAGLLAARGLLAGGRGDNW 466
Ric8B_BOVIN  407 NKLVRLMTHVDLGVKQIAAEFLFVLCKERVDSLLKYTGYGNAAGLLAARGLLAGGRGDNW 466
Ric8B_MOUSE  407 NKLVRLMTHVDLGVKQIAAEFLFVLCKERVDSLLKYTGYGNAAGLLAARGLLAGGRGDNW 466
Ric8B_RAT    407 NKLVRLMTHVDLGVKQIAAEFLFVLCKERVDSLLKYTGYGNAAGLLAARGLLAGGRGDNW 466
                 *********:*--**:-********.* *- ::******************: ***
                    αB8         αC8      αa9          αb9

Ric8A_RAT    444 EYREAKASINPVTGRVEEKPPNPMEGMTEEQKEHEAMKLVNMFDKLSRHR 493
Ric8A_MOUSE  444 EYREAKASINPVTGRVEEKPPNPMEGMTEEQKEHEAMKLVNMFDKLSRHR 493
Ric8A_HUMAN  445 EYKEAKASINPVTGRVEEKPPNPMEGMTEEQKEHEAMKLVTMFDKLSRNR 494
Ric8A_BOVIN  444 EYKEAKASINPVTGRVEEKPPNPMEGMTEEQKEHEAMKLVNMFDKLSRHR 493
Ric8A_DANRE  457 EYREAKPHINPVTGRVEEEQPNPMEGMTDEQKEYEAMKLVNMFDKLSREQ 506
Ric8A_CHICK  453 EYKEAKPNINPVTGRVEEKLPNPMEGMTEEQKEYEAMKLVNMFDKLSREQ 502
Ric8A_XENTR  453 EYREAKPNINPVTGRVEEKQPNPMDGMTEEQKEYEAMKLVNMFDKLSREQ 502
Ric8B_DANRE  454 EYKSVKPFINPITGHVEDPLPNPLDEMTEEQKEYEAQRLVSMIDELSRKE 503
Ric8B_XENTR  439 EYLLAKPNINLITGHVEEPLPNPMDEMTEEQKEYEAMKLVNMFDKLSREE 488
Ric8B_CHICK  454 EYKSAKPNINLITGHLEEPMPNPMDEMTEEQKEYEAMKLVNMFDKLSRDE 503
Ric8B_HUMAN  477 EYKNAKPNINLITGHLEEPMPNPIDEMTEEQKEYEAMKLVNMLDKLSREE 526
Ric8B_BOVIN  477 EYKNAKPNINLITGHLEEPMPNPIDEMTEEQKEYEAMKLVNMLDKLSREE 526
Ric8B_MOUSE  477 EYKNAKPNINLITGHLEEPMPNPIDEMTEEQKEYEAMKLVNMLDKLSREE 526
Ric8B_RAT    477 EYKNAKPNINLITGHLEEPMPNPIDEMTEEQKEYEAMKLVNMLDKLSREE
                 **  .*  **-:**::*:- ***:: **:****:** :**.*:*:***..
                    RT                            α11
```

Extended Data Figure 9a. Amino acid sequences of Ric-8A and Ric-8B across phylogeny near Gα contact residues. Residue numbers are shown at the start and, for long tracts, the end of subsequences that harbor Ric-8A contacting-residues, highlighted in *green* in the sequence of rat Ric-8A, defined as those that include at least one atom within 4.0 of an atom in Gαi1. Amino acids not identical in rat Ric-8A and rat Ric-8B in *red* in the conservation summary at the bottom of each alignment. Central residues of secondary structural elements are shown below the conservation summary.

```
Gs    31 QKDKQVYRATHRLLLLLGAGESGKSTIVKQM 60    212 FQVDKVNFHMFDVGGQRDERRKWIQCFNDVTA 243
Golf 102 REQKRDLQQTHRLLLLLGAGESGKSTIVKQM 131   268 FQVDKVNFHMFDVGGQRDERRKWIQCFNDVTA 299
G12   46 ARERRAVRRLVKILLLGAGESGKSTFLKQM 75    214 FVIKKIPFKMVDVGGQRSQRQKWFQCFDGITS 245
G13   39 SREKTYVKRLVKILLLGAGESGKSTFLKQM 68    211 FEIKNVPFKMVDVGGQRSERKRWFECFDSVTS 242
GQ    30 NRDKDARRELKLLLLGTGESGKSTFIKQM 59    194 FDLQSVIFRMVDVGGQRSERRKWIHCFENVTS 225
G11   30 ERDKRDARRELKLLLLGTGESGKSTFIKQM 59    194 FDLENIIFRMVDVGGQRSERRKWIHCFENVTS 225
GT3   24 QEDAERDARTVKLLLLGAGESGKSTIVKQM 53    189 FSFKDLNFRMFDVGGQRSERKKWIHCFEGVTC 220
GT1   20 KEDAEKDARTVKLLLLGAGESGKSTIVKQM 49    185 FSFKDLNFRMFDVGGQRSERKKWIHCFEGVTC 216
GT2   24 QEDADKEAKTVKLLLLGAGESGKSTIVKQM 53    189 FSVKDLNFRMFDVGGQRSERKKWIHCFEGVTC 220
GO    24 KEDGISAAKDVKLLLLGAGESGKSTIVKQM 53    190 FTFKNLHFRLFDVGGQRSERKKWIHCFEDVTA 221
GI2   24 REDGEKAAREVKLLLLGAGESGKSTIVKQM 53    190 FTFKDLHFKMFDVGGQRSERKKWIHCFEGVTA 221
GI1   24 REDGEKAAREVKLLLLGAGESGKSTIVKQM 53    189 FTFKDLHFKMFDVGGQRSERKKWIHCFEGVTA 220
GI3   24 REDGEKAAKEVKLLLLGAGESGKSTIVKQM 53    189 FTFKELYFKMFDVGGQRSERKKWIHCFEGVTA 220
         .:   ■ ■:****::*******:;***          * ■..: *::.******.:*::*:■.**:.:*.
             β1  P-loop    α1                   β2    β3       Switch II

Gs   268 EALNLFKSIWNNRWLRTISVILFLNK 293    358 YCYPHFTCAVDTENIRRVFNDCRDIIQRMHLRQYELL 394
Golf 324 ESLDLFESIWNNRWLRTISIILFLNK 349    414 YCYPHFTCAVDTENIRRVFNDCRDIIQRMHLKQYELL 450
G12  270 ESMNIFETIVNNKLFFNVSIILFLNK 295    343 PLFHHFTTAIDTENIRFVFHAVKDTILQENLKDIMLQ 379
G13  267 ESLNIFETIVNNRVFSNVSIILFLNK 292    343 PLFHHFTTAIDTENIRFVFHAVKDTILQENLKDIMLQ 379
GQ   250 ESKALFRTIITYPWFQNSSVILFLNK 275    341 PLYHHFTTAINTENIRLVFRDVKDTILHDNLKQLMLQ 377
G11  250 ESKALFRTIITYPWFQHSSVILFLNK 275    323 IIYSHFTCATDTENIRFVFAAVKDTILQLNLKEYNLV 359
GT3  245 ESLHLFNSICNHKYFATTSIVLFLNK 270    318 EIYSHMTCATDTQNVKFVFDAVTDIIIKENLKDCGLF 354
GT1  241 ESLHLFNSICNHRYFATTSIVLFLNK 266    314 EIYSHMTCATDTQNVKFVFDAVTDIIIKENLKDCGLF 350
GT2  245 ESLHLFNSICNHKFFAATSIVLFLNK 270    318 EIYSHMTCATDTQNVKFVFDAVTDIIIKENLKDCGLF 354
GO   246 ESLMLFDSICNNKFFIDTSIILFLNK 271    318 EIYCHMTCATDTNNIQVVFDAVTDIIIANNLRGCGLF 354
GI2  246 ESMKLFDSICNNKWFTDTSIILFLNK 271    319 EIYTHFTCATDTKNVQFVFDAVTDVIIKNNLKDCGLF 355
GI1  245 ESMKLFDSICNNKWFTDTSIILFLNK 270    318 EIYTHFTCATDTKNVQFVFDAVTDVIIKNNLKDCGLF 354
GI3  245 ESMKLFDSICNNKWFTDTSIILFLNK 270    318 EIYTHFTCATDTKNVQFVFDAVTDVIIKNNLKDCGLF 354
         *:   :* :*  .  ■ :  ■■■*::*****      ■  ■■*:* *■:*:*:■  **  ■■■■* *■   :*: ■■*■
              α3     β5         p1              β6    p2        α5
```

Extended Data Figure 9b. Sequence of rat Gα isoforms within Ric-8A contact regions.
Residue numbers are shown at the start and end of subsequences that harbor Ric-8A contacting-
residues, highlighted in *green* in the sequence of Gαi1, defined as those that include at least one
atom within 4.0 of an atom in Ric-8A. Amino acids not identical in Gαi and Gαs are highlighted in
red, in the conservation summary at the bottom of each alignment. Central residues of secondary
structural elements are shown below the conservation summary. Symbols p1 and p2 denote
guanine nucleotide purine binding sites formed at the β5-αG and β6-α5 loops, respectively.

Extended Data Figure 10. Detail of Ric-8A:Gαi1 contact involving Gαi1 α4 and β6 with Ric-8A αB8-αC8. The crystal structure is shown with Gαi1 backbone in *cyan* and Ric-8A backbone in *wheat* with side-chain atom colors, carbon, *green*, nitrogen, *blue*, oxygen, *red* and sulfur, *yellow*. Mutation of rL383 to glutamate reduces GEF activity to 50% that of wild-type Ric-8A.