1   **A deep dive into the ancestral chromosome number of flowering plants**

3   Angelino Carta*, Gianni Bedini, Lorenzo Peruzzi

4   Dipartimento di Biologia, Botany Unit, University of Pisa, via Derna 1, I-56126 Pisa, Italy

6   * Corresponding author: angelino.carta@unipi.it

Abstract

Chromosome rearrangements are a well-known evolutionary feature in eukaryotic organisms[1],

especially plants. The remarkable diversity of flowering plants (angiosperms) has been

attributed, in part, to the tremendous variation in their chromosome number[2]. This variation has

stimulated a blossoming number of speculations about the ancestral chromosome number of

angiosperms[2-7], but estimates so far remain equivocal and relied on algebraic approaches lacking

an explicit phylogenetic framework. Here we used a probabilistic approach to model haploid

chromosome number (n) changes[8] along a phylogeny embracing more than 10 thousands taxa,

to reconstruct the ancestral chromosome number of the common ancestor of extant angiosperms

and the most recent common ancestor for single angiosperm families.

Bayesian inference revealed an ancestral haploid chromosome number for angiosperms $n = 7$,

reinforcing previous hypotheses[2-7] that suggested a low ancestral basic number. Inferred n for

single families, more than half of which are provided here for the first time, are mostly

congruent with previous evaluations. Chromosome fusion (loss) and duplication (polyploidy)

are the predominant transition types inferred along the phylogenetic tree, emphasising the

importance of both dysploidy[6,9,10] and genome duplication[2,7,11-13] in chromosome number

evolution. Significantly, while dysploidy is equally distributed early and late across the whole

phylogeny, polyploidy is detected mainly towards the tips of the tree. Therefore, little evidence

exists for a link between ancestral chromosome numbers and putative ancient polyploidization

events[14], suggesting that further insights are needed to elucidate the organization of genome

packaging into chromosomes.

Introduction

Each eukaryotic organism has a characteristic chromosome complement, its karyotype, which represents the highest level of structural and functional organization of the nuclear genome[15]. Karyotype constancy ensures the transfer of the same genetic material to the next generation, while karyotype variation provides genetic support to ecological differentiation and adaptation[2,15]. Cytogenetic studies have shown that the tremendous inter- and intra-taxonomic variation of chromosome number documented in flowering plants[2,5,16] is mostly driven by two major mechanisms: a) increases through polyploidy (which may entail a Whole Genome Duplication [WGD] or an increase by half of the genome, demi-duplication[8]); b) decreases or increases through structural chromosomal rearrangements like chromosome fusion, i.e. descending dysploidy, and chromosome fission, i.e. ascending dysploidy.

Polyploidy is a common and ongoing phenomenon, especially in plants[13], that has played an important role in many lineages, with evidence of several rounds of both ancient and recent polyploidization[11,17,18], albeit its distribution in time remains contested[14]. Indeed, although the crucial role of polyploidy in plant diversification on small timescales is widely accepted[2,6], the evolutionary significance of polyplodization for the long-term diversity of angiosperms is still controversial[12]. On the other hand, while dysploidy is more frequent than polyploidy in angiosperms[6], its adaptive consequences have been mostly unexamined[19], until recent studies demonstrated its high evolutionary impact[9].

Chromosome number variation across angiosperm lineages spans two orders of magnitude[15], from $2n = 4$ to $2n = 640$. Previous hypotheses[2-7] of the ancestral basic (monoploid)[20] chromosome number p in angiosperms suggest low numbers, between $p = 6$ and $p = 9$. These hypotheses placed particular attention to 'primitive' extant angiosperms[5] to estimate putative ancestral basic chromosome numbers. More recently, an ancestral chromosome number has been reconstructed using a maximum parsimony approach[7]. However, although parsimony has been widely used to infer ancestral chromosome numbers, it carries significant shortcomings[8],

and more rigorous and complex models to infer chromosome number evolution are currently

available [8,21,22,23].

Here we use probabilistic models, accounting for various types of chromosome number

transitions, to reconstruct the ancestral haploid chromosome number and the occurrence of

chromosome change events across the most massive data set ever assembled linking

chromosome numbers to a phylogeny, sampling 10,766 taxa from 59 orders (92%) and 318

families (73%) of angiosperms.

Chromosome numbers were extracted from the Chromosome Counts DataBase[24], and the

analyses were conducted using pruned versions of two recently published, dated mega-trees for

seed plants[25], the first one (GBMB) constructed with a backbone based on Magallón et al.[26], and

the second one (GBOTB) grounded on Open Tree of Life version 9.1. In addition, we conducted

all analyses again using a different ultrametric tree of 1,559 taxa extracted from a recently

published plastid phylogenomic angiosperm (PPA) tree[27].

Results and Discussion

Regardless of the three alternative phylogenies, $n = 7$ was inferred as the ancestral haploid

chromosome number with the highest posterior probability (Table 1) and likelihood (Table 2).

The ancestral haploid chromosome number $n = 7$ was remarkably stable in the deepest part of

the phylogeny (Fig. 1), while slight variations ($\pm$ 1) in n were inferred at the base of some

lineages. Greater variations were shown in the ancestral haploid chromosome number of many

angiosperm families (see Supplementary Table 1). Monocots exhibited the largest variation of

inferred n among Most Recent Common Ancestors (MRCAs) of plant families (Fig. 2b),

paralleled by a considerable variation in current haploid chromosome numbers (Fig. 2a). Over

70% of inferred n in the 158 families for which previous inferences were available are in line

with previous proposals. For the remaining 160 families (50.3%) the first inferences are

presented here. Discrepancies at the family level among inferences obtained in this study and

those from previous literature are possibly due to the use of routine algebraic approaches, instead of phylogenetic models, to infer chromosome number changes[28]. For example, the inferred *n* of MRCAs of Brassicaceae, Lamiaceae, and Rosaceae are respectively 7, 8, and 7 in our study, but were previously inferred as 12, 14, and 9, respectively[5]. Indeed, even in the presence of a strong phylogenetic signal (e.g., closely related species sharing similar chromosome numbers)[29,30], algebraic inferences of chromosome numbers become increasingly difficult with increasing phylogenetic depth, as identical chromosome numbers will occur in unrelated lineages[19]. The dataset analysed here is the most extensive ever used for inferring ancestral haploid number in angiosperms, but it still poses challenges concerning incomplete taxon sampling and phylogenetic resolution at the family level.

For both GBMB and GBOTB phylogenies, the best model considers up to six parameters (Table 3), i.e. chromosome gain, loss, duplication, demi-duplication rates and rates of gain and loss linearly dependent on the current chromosome number. Our results support the conclusion that genome duplication and dysploidy were critical events in the evolution of angiosperms. Specifically, descending dysploidy, most likely through chromosome fusion, was the most common cytogenetic mechanism of chromosome number change during the evolution of flowering plants, and this is inferred both on branches leading to major clades and on terminal branches (Supplementary Figs. 1-3). Our results emphasize the importance of dysploidy in the evolution of chromosome numbers in angiosperms[9]. Interestingly, demi-duplication events, associated with hybridisation between different ploidy levels and with allopolyploidization processes, were also inferred in a significant number of events, albeit mainly in terminal branches.

Whilst polyploidization is the second most frequent transition type, ancient polyploidy events are underrepresented. The absence of polyploidization events at the base of the tree is in agreement with the maintenance of the ancestral haploid chromosome number *n* = 7 inferred in the deepest part of the phylogeny. Polyploidization events were instead inferred mainly toward

the tips of the tree (Supplementary Fig. 1,2,3), partially supporting previous evidence revealing

independent genome duplications near the base of several families[9,18,28], leading to high haploid

chromosome numbers. Indeed, this may be the case of many families in the Magnoliid clade.

Our results provide no direct insight for a link among some of the most extensive plant

radiations and ancient polyploidization rounds[11]. However, our analyses do not contradict

evidence of WGD events at the origin of angiosperms or before it[14], but rather highlight that

genome size may vary independently of chromosome number[7]. Inferring ancient polyploidy

events from cytological data is indeed a challenging task, because genome rearrangements[31]

following polyploidisation gradually can hide signals of genome duplication over time[9].

Phylogenomic analyses interpret gene duplications as the result of a shared duplication event

occurring in a common ancestor, while models of chromosome number evolution consider

WGDs as separately occurring in different lineages[31].

The main results presented here were drawn using the largest dated mega-tree currently

available for seed plants. We explored the sensitivity of our results by conducting all analyses

again using a different ultrametric tree[26], including a lower number of sampled taxa but allowing

to consider intraspecific chromosome number variation for each taxon. We found only minor

differences at the root (Tables 1,2) and at some internal nodes (Fig. 1).

Reconstructing the ancestral chromosome number is difficult, because there are no suitable

outgroups for direct comparison[32], and because extant early branching angiosperms (e.g.,

Amborella and Nymphaeales) are not necessarily holding plesiomorphic character-states. With

these limitations in mind, we made inferences based on the distribution of n in extant

angiosperms, and using probabilistic models accounting for various types of chromosome

number transitions. Our study is not able to address the origin of chromosome number of the

first angiosperms. Instead, it provides novel, detailed, and well-supported inference of ancestral

haploid chromosome number of the common ancestor of all extant angiosperms, as well as the

earliest steps of the subsequent chromosome number transitions, including n inferences for 318

angiosperm families. Interestingly, our inferred ancestral state for the haploid number *n*

coincides with the ancestral basic chromosome number *p* previously proposed for angiosperms

based on empirical counts[2-7] or paleogenomic approaches[33].

Our study allowed to clarify a long-standing question[2], but such reconstruction necessarily

comes with limitations. Nevertheless, this is a major step forward in understanding the ancestral

chromosome number for angiosperms, and we believe that this issue should be added to the

angiosperm macroevolutionary agenda[34]. Progress in reconstructing the ancestral chromosome

number may require the development of models that include heterogeneity in the patterns of

chromosome evolution across a phylogenetic tree[23], along with a deeper insight into genome and

karyotype evolution.

Methods

Phylogenetic reconstruction

We used two recently published[25] dated megaphylogenies for seed plants, GBMB and GBOTB,

as backbones to generate two alternative phylogenies for angiosperms included in the dataset.

GBMB and GBOTB were constructed using 79,874 and 79,881 taxa, respectively, available in

GenBank and in a backbone provided either by Magallón et al.[26] (GBMB) or by Open Tree of

Life, version 9.1 (GBOTB). In addition, we also used a different ultrametric tree provided by a

recently published plastid phylogenomic angiosperm (PPA) tree[27].

Chromosome numbers collection

The haploid chromosome numbers (*n*) of the species were obtained from the Chromosome

Counts Database (CCDB[24]; http://ccdb.tau.ac.il/; last accessed May 2019) using the R package

chromer[35]. CCDB contains records from original sources that have irregularities of chromosome

counts, so that the ca. 150,000 records were curated semiautomatically using the CCDBcurator

package[36] and custom R scripts. After a first round of automatic cleaning, we examined results

by hand and corrected records where needed.

Species with unknown chromosome counts were pruned from the trees, thus we collected

chromosome numbers for 10,766 taxa included in the GBMB and GBOTB phylogenetic trees,

and for 1,559 taxa included in the PPA tree. In cases where multiple chromosome numbers were

reported for a given taxon, the modal number was used[8,37]. For taxa with numbers suggesting

different ploidy levels, we used the lowest haploid chromosome number available[38]. This coding

scheme allowed us to deal with the problem of the existence of different ploidy levels in a taxon

and also with the low-density sampling conducted in most taxa[38]. Analyses conducted using the

PPA tree encountered less computation limitations, so that we were able to perform them by

explicitly considering intraspecific polymorphism, allowing several chromosome numbers,

together with their respective frequencies, to be set for each taxon[21].

Analyses

The evolution of haploid chromosome numbers of angiosperms was inferred using chromEvol[21]

software v.2.0 (http://www.tau.ac.il/~itaymay/cp/chromEvol/index.html). This software

determines the likelihood of a model to explain the given data along the phylogeny, based on the

combination of two or more of the following parameters: dysploidisation (ascending,

chromosome gain rate λ; descending, chromosome loss rate δ), polyploidisation (chromosome

number duplication with rate ρ, demi-polyploidisation or triploidisation with rate μ) and

incremental changes to the basic number with regard to a rate of multiplication that is different

from a regular duplication[8]. Two additional parameters (λ_1, δ_1) detect linear dependency

between the current haploid number and the rate of gain and loss of chromosomes. We tested 10

models based on a different combination of the parameters above. Four of these models consider

only constant rates (Mc1, Mc2, Mc3, and Mc0), whereas the other four include two linear rate

parameters (Ml1, Ml2, Ml3, and Ml0; Table 3). Both sets have a null model (Mc0 and Ml0) that

assumes no polyploidisation events. Finally, two models (Mb1 and Mb2) consider that the

evolution of chromosome number can also be influenced by the basic number (β) and by its

transition rates (ν).

The minimum chromosome number allowed in the analyses was set to 2, whereas the maximum

number was set to 5 units higher than the highest chromosome number found in the empirical

data. We removed all counts $n > 43$ from the analysis, because for many lineages the sampling

was inadequate to reconstruct such a drastic change in chromosome number[38-39] and because of

computation limitations[23]. The branch lengths were scaled according to the software author's

instructions. The null hypothesis (no polyploidy) was tested with likelihood ratio tests using the

Akaike information criterion (AIC)[40]. To compute the expected number of changes along each

branch, as well as the ancestral haploid chromosome numbers at internal nodes, the best fitted

model for both data sets was rerun using 1,000 simulations. The best model was plotted on the

trees using the ChromEvol functions v0.9-1 elaborated by N. Cusimano

(http://www.sysbot.biologie.uni-muenchen.de/en/people/cusimano/use_r.html) in R.

To test which ancestral haploid chromosome number is most likely fort the root of angiosperms,

the following haploid chromosome numbers were fixed at the root and the likelihood of the

resulting models was compared: n = 4,5,6,7,8,9. These numbers were tested either because

considered putative ancestral character-states[2-7], or because they were identified as the

chromosome numbers showing the highest PP under our Bayesian analysis. All analyses were

performed in the high-performance computing cluster at the University of Pisa.

References

1.   Coghlan, A., Eichler, E.E., Oliver, S.G., Paterson, A.H. & Stein, L. Chromosome evolution

in eukaryotes: a multi-kingdom perspective. *Trends Genet*. **21**, 673–682 (2005).

2.   Stebbins, G.L. *Chromosomal Evolution in Higher Plants*. (Edward Arnold, London, 1971).

3.   Ehrendorfer, F., Krendl, F., Habeler, E., & Sauer, W. Chromosome numbers and evolution

in primitive angiosperms. *Taxon* **17**, 337–353 (1968).

4. Walker, J.W. Chromosome numbers, phylogeny, phytogeography of the Annonaceae and
their bearing on the (original) basic chromosome number of angiosperms. Taxon **21**, 57–65
(1972).

5. Raven, P.H. The bases of angiosperm phylogeny: cytology. *Ann. Missouri Bot. Gard*. **62**,
724–764 (1975).

6. Grant, V. *Plant Speciation* (ed. 2) (Columbia University Press, New York, 1981).

7. Soltis, D.E., Soltis, P.S., Endress, P.K. & Chase, M.W. in *Phylogeny and Evolution of*
Angiosperms (ed. Soltis, D.E., Soltis, P.S., Endress, P.K. & Chase, M.W.) 287–302
(Sinauer Associates, Sunderland, 2005).

8. Mayrose, I., Barker, M.S. & Otto, S.P. Probabilistic models of chromosome number
evolution and the inference of polyploidy. *Syst. Biol*. **59**, 132–144 (2009).

9. Escudero, M. et al. Karyotypic changes through dysploidy persist longer over evolutionary
time than polyploid changes. *PLOS One* **9**, e85266 (2014).

10. Guerra, M. Chromosome numbers in plant cytotaxonomy: concepts and implications.
Cytogenet. Genome Res. **120**, 339–350 (2008).

11. Jiao, Y. et al. Ancestral polyploidy in seed plants and angiosperms. Nature **473**, 97 (2011).

12. Mayrose, I. et al. Recently formed polyploid plants diversify at lower rates. *Science* **333**,
1257–1257 (2011).

13. Wood, T.E. et al. The frequency of polyploid speciation in vascular plants. *Proc. Nat. Acad.*
Sci. USA 106, 13875–13879 (2009).

14. Ruprecht, C. et al. (2017) Revisiting ancestral polyploidy in plants. *Science Adv*. **3**,
e1603195 (2017).

15. Stace, C.A. Cytology and cytogenetics as a fundamental taxonomic resource for the 20th
and 21st centuries. *Taxon* **49**: 451–477 (2000).

16. Levitzky, G.A. The karyotype in systematics. *Bull. Appl. Bot. Genet. Plant Breed*. **27**, 220–
240 (1931).

17. Li, Z. et al. Early genome duplications in conifers and other seed plants. *Sci. Adv*. **1**,
e1501084 (2015).

18. Leebens-Mack, J.H., Barker, M.S., Carpenter, E.J. et al. One thousand plant transcriptomes
and the phylogenomics of green plants. *Nature* **574**, 679–685 (2019).

19. Weiss-Schneeweiss, H., & Schneeweiss, G.M. in *Plant Genome Diversity* Vol. 2 (eds.
Leitch, I.J., Greilhuber, J., Dolezel, J.W.J) 209–230 (Springer, Vienna, 2013).

20. Peruzzi, L. "x" is not a bias, but a number with real biological significance. Plant Biosyst.
147, 1238–1241 (2013).

21. Glick, L. & Mayrose, I. ChromEvol: assessing the pattern of chromosome number evolution
and the inference of polyploidy along a phylogeny. *Mol. Biol. Evol*. **31**, 1914–1922 (2014).

22. Freyman, W.A. & Höhna, S. Cladogenetic and anagenetic models of chromosome number
evolution: a Bayesian model averaging approach. *Syst. Biol*. **67**, 195–215 (2018).

23. Zenil-Ferguson, R., Burleigh, J.G. & Ponciano, J.L. Chromploid: an R package for
chromosome number evolution across the plant tree of life. *Appl. Plant Sci*. **6**, e1037
10.1002/aps3.1037 (2018).

24. Rice, A. et al. The Chromosome Counts Database (CCDB)–a community resource of plant
chromosome numbers. *New Phytol*. **206**, 19–26 (2015).

25. Smith, S.A. & Brown, J.W. Constructing a broadly inclusive seed plant phylogeny. *Am. J.*
Bot. **105**, 1–13 (2018).

26. Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L.L. & Hernández-Hernández, T. A
metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity.
New Phytol. **207**, 437–453 (2015).

27. Li, H.T. et al. Origin of angiosperms and the puzzle of the Jurassic gap. *Nature Plants*, **5**,
461 (2019).

28. Cusimano, N., Sousa, A., & Renner, S.S. Maximum likelihood inference implies a high, not

a low, ancestral haploid chromosome number in Araceae, with a critique of the bias

introduced by 'x'. *Ann. Bot.* **109**, 681–692 (2012).

29. Escudero, M. et al. Selection and inertia in the evolution of holocentric chromosomes in

sedges (*Carex*, Cyperaceae). *New Phytol.* **195**, 237–247 (2012).

30. Schubert, I., & Lysak, M.A. Interpretation of karyotype evolution should consider

chromosome structural constraints. *Trends Gen.* **27**, 207–216 (2011).

31. Mandakova, T. & Lysak, M.A. Post-polyploid diploidization and diversification through

dysploid changes. *Curr. Opinion Plant Biol.* **42**, 55–65 (2018).

32. Doyle, J.A. Molecular and fossil evidence on the origin of angiosperms. *Ann. Rev. Earth*

Planet. Sci. **40**, 301–326 (2012).

33. Salse, J. In silico archeogenomics unveils modern plant genome organization, regulation

and evolution. *Curr. Opinion Plant Biol.* **15**, 122–130 (2012).

34. Sauquet, H. & Magallón, S. Key questions and challenges in angiosperm macroevolution.

New Phytol. **219**, 1170–1187 (2018).

35. Pennell, M.W. Chromer: Interface to Chromosome Counts Database API. R package

version 0.1.2.9000 (2016).

36. Rivero, R., Sessa, E. B., & Zenil-Ferguson, R. EyeChrom and CCDB curator: Visualizing

chromosome count data from plants. *Applications in Plant Sciences*, **7**, e01207 (2019).

37. Salman-Minkov, A., Sabath, N. & Mayrose, I. Whole-genome duplication as a key factor in

crop domestication. *Nature Plants* **2**, 16115 (2016).

38. Márquez-Corro, J.I., Martín-Bravo, S., Spalink, D., Luceño, M. & Escudero, M. Inferring

hypothesis-based transitions in clade-specific models of chromosome number evolution in

sedges (Cyperaceae). *Mol. Phylogenet. Evol.* **135**, 203–209 (2019).

39. Barrett, C.F. et al. Ancient polyploidy and genome evolution in palms. *Genome Biol. Evol.*

11, 1501–1511 (2019).

40. Burnham, K.P. & Anderson, D.R. Model inference. Understanding AIC and BIC in model

selection. *Socio. Meth. Res. Int. J. Bot. Res*. **33**, 261–304 (2004).

Table 1. Summary of chromosome number evolutionary models and inferred ancestral

haploid chromosome number (*n*) in Angiosperms under the best-fitting model.

tree	Best model	LogLik	AIC	Rates						Events inferred with PP > 0.5				Chromosome no. at root node		
				λ	δ	ρ	μ	λ_1	δ_1	Gain	Losses	Duplications	Demi	Bayes (PP): Best p	Bayes(PP): 2nd best p	ML
GBMB (10,766 taxa)	Ml3	-1812 0.0	3625 0.0	0.00 81	0.01 13	0.01 31	0.00 51	0.00 51	0.00 07	50 9.7	162 7.5	1438.1	37 6.1	7 (0.9 7)	8 (0.02)	5
GBOTB (10,766 taxa)	Ml3	-1811 0.0	3624 0.0	0.01 06	0.00 96	0.01 30	0.00 49	0.00 01	0.00 08	58 9.5	162 5.3	1435.3	36 3.3	7 (0.9 8)	8 (0.01)	5
PPA (1559 taxa)	Ml2	-3788. 0	7586 .0	0.01 01	0.00 44	0.00 78	-	-0.00 02	0.00 12	14 5.0	528. 1	244.6	19 1.9	7 (0.2 4)	5 (0.21)	2

Only the best-fitting models are shown. Tree refers to the three alternative phylogenies used. Best model, Ml3 (linear rate model with duplication rate ρ and demi-duplication rate μ) Ml2 (linear rate model with equal duplication and demi-duplication rates); Logarithmic likelihood (LogLik) and AIC scores; rate parameters (*λ* = chromosome gain rate, *δ* = chromosome loss rate, *ρ* = duplication rate, *μ* = demi-duplication rate, *λ₁*= linear chromosome gain, *δ₁* = linear chromosome loss); frequency of the four possible event types with a posterior probability (PP) > 0.5; haploid chromosome number inferred at the root node under Bayesian optimization with the respective PP, and under maximum likelihood (ML).

289   **Table 2. Testing hypotheses about root ancestral haploid chromosome number (n).**

Tree	GBMB		GBOTB		PPA	
	LogLik	AIC	LogLik	AIC	LogLik	AIC
Root fixed at $n = 4$	-18127.1	36266.2	-18129.9	36271.7	-3788.8	7587.6
Root fixed at $n = 5$	-18128.1	36268.2	-18142.6	36297.2	-3787.2	7584.5
Root fixed at $n = 6$	-18123.6	36259.2	-18124.2	36260.4	-3786.7	7583.5
Root fixed at $n = 7$	**-18117.1**	**36246.3**	**-18115.5**	**36243.0**	**-3786.2**	**7582.4**
Root fixed at $n = 8$	-18119.4	36250.9	-18127.8	36267.6	-3787.6	7585.1
Root fixed at $n = 9$	-18122.2	36256.4	-18121.3	36254.7	-3788.4	7586.9

AIC and LogLik values obtained by fixing the root with given ancestral haploid chromosome number. The lowest AIC and LogLik values are shown in bold.

291 **Table 3. Goodness of fit of the 10 different models of chromosome number evolution**

292 **applied to the three alternative phylogenies used.**

Model	GBMB			GBOTB			PPA			Parameters
	LogLik	AIC	AIC$_w$	LogLik	AIC	AIC$_w$	LogLik	AIC	AIC$_w$	
Mc1	-19870	39750	0.0000000000	-19900	39800	0.0000000000	-4032	8070	0.0000000000	$\lambda; \delta; \rho$
Mc2	-18280	36560	0.0000000000	-18270	36550	0.0000000000	-3804	7613	0.0000013710	$\lambda; \delta; \rho=\mu$
Mc3	-18140	36290	0.0000000021	-18140	36280	0.0000000021	-3804	7616	0.0000003059	$\lambda; \delta; \rho; \mu$
Mc0	-47870	95740	0.0000000000	-49060	98120	0.0000000000	-5135	10270	0.0000000000	$\lambda; \delta$
Ml1	-19670	39350	0.0000000000	-19660	39320	0.0000000000	-3999	8008	0.0000000000	$\lambda; \delta; \rho; \lambda_1; \delta_1$
Ml2	-18260	36520	0.0000000000	-18250	36510	0.0000000000	**-3788**	**7586**	**0.9999999979**	$\lambda; \delta; \rho=\mu; \lambda_1; \delta_1$
Ml3	**-18120**	**36250**	**0.9999999979**	**-18110**	**36240**	**0.9999999979**	-3787	7587	0.6065306585	$\lambda; \delta; \rho; \mu; \lambda_1; \delta_1$
Ml0	-45260	90530	0.0000000000	-44910	89820	0.0000000000	-4918	9843	0.0000000000	$\lambda; \delta; \lambda_1; \delta_1$
Mb1	-18650	37310	0.0000000000	-18470	36950	0.0000000000	-4001	8011	0.0000000000	$\lambda; \delta; \beta; \nu$
Mb2	-21790	43580	0.0000000000	-21830	43670	0.0000000000	-4287	8581	0.0000000000	$\lambda; \delta; \rho; \beta; \nu$

Mc indicate models with constant rates, Ml models that include linear rate parameters and Mb models that include base number (not the chromosome number at the root of the phylogeny) parameters[8-21]. Logarithmic likelihood (LogLik), AIC and relative weights scores (AICw). In bold, the lowest AIC value for each phylogeny indicates the best model. The last column indicates the parameter estimates included in each model (see Methods for details).

Figure 1 Reconstruction of ancestral haploid chromosome number (*n*) of angiosperms with the best-fitting model on the different types of trees used (GBMB and PPA). Please note that the plotted trees depict ordinal phylogenetic relationships (sub-ordinal topologies were collapsed to build the figure), and are shown without branch length information. Pie charts at nodes represent the probability of the ancestral haploid chromosome numbers inferred under Bayesian estimation; the numbers at nodes are those with the highest probability. Pie charts and numbers at the tips are the three best inferred ancestral haploid chromosome numbers per each angiosperm order. Black lines, difference in phylogenetic position between GBMB and PPA trees.

Figure 2 Density plots of haploid chromosome numbers (*n*) and of inferred ancestral haploid chromosome number (*n*) for each angiosperm family in four major angiosperm clades (APG IV). a, we identified the number of unique chromosome counts per taxon, i.e. cytotypes, from the original dataset, after excluding counts with $n > 60$, to focus on the most frequent *n* and their putative relation with inferred *p*. **b**, density plots were scaled by the Bayesian posterior probability (PP) estimated for each inferred *p*.

Acknowledgements

The authors thank Marcial Escudero and Itay Mayrose for their help with ChromEvol analyses.

Author contributions

A.C. planned and designed the research, analysed the data and wrote the manuscript. G.B.

assisted in chromosome numbers acquisition. L.P. and G.B. contributed to successive versions

of the manuscript and in solving theoretical and nomenclatural issues. All authors read and

approved the final manuscript.