A deep dive into the ancestral chromosome number and genome size of flowering plants

Angelino Carta*, Gianni Bedini, Lorenzo Peruzzi

Department of Biology, Botany Unit, University of Pisa, via Derna 1, I-56126 Pisa, Italy

Author for correspondence:

Angelino Carta

Tel: +39 050 2211314

Email: acarta@biologia.unipi.it

Total word count: 3903

Introduction word count: 874

M&M word count: 1297

Results word count: 634

Discussion word count: 1069

Number of figures: 3 (should be published in colour)

Number of tables: 4

Supporting information: Tables S1 and Figure S1-S8

Summary

•   Chromosome number and genome variation in flowering plants has stimulated a
blossoming number of speculations about the ancestral chromosome number of
angiosperms, but estimates so far remain equivocal.
•   We used a probabilistic approach to model haploid chromosome number (n) changes
along a phylogeny embracing more than 10 thousands taxa, to reconstruct the ancestral
chromosome number of the common ancestor of extant angiosperms and the most recent
common ancestor for single angiosperm families. Independently, we carried out an
analysis of 1C genome size evolution, including over 5 thousands taxa.
•   Our inferences revealed an ancestral haploid chromosome number for angiosperms $n = 7$,
a diploid status, and an ancestral 1C = 1.73 pg. For 160 families, inferred ancestral n are
provided for the first time.
•   Both descending dysploidy and polyploidy played crucial roles in chromosome number
evolution. While descending dysploidy is equally distributed early and late across the
phylogeny, polyploidy is detected mainly towards the tips. Similarly, also 1C genome
size significantly increases (or decreases) in late-branching lineages. Therefore, no
evidence exists for a clear link between ancestral chromosome numbers and ancient
polyploidization events, suggesting that further insights are needed to elucidate the
organization of genome packaging into chromosomes.

Key words: Angiosperms, chromosome evolution, haploid chromosome number, genome
size, dysploidy, polyploidy

Introduction

Chromosome rearrangements are a well-known evolutionary feature in eukaryotic organisms (Coghlan *et al.,* 2005), especially in plants. The remarkable diversity of flowering plants (angiosperms) has been attributed, in part, to the tremendous variation in their chromosome number (Stebbins, 1971). This variation has stimulated a blossoming number of speculations about the ancestral chromosome number of angiosperms (Ehrendorfer *et al.,* 1968; Stebbins, 1971; Walker 1972; Raven, 1975; Grant, 1981; Soltis *et al.,* 2005), but estimates so far remain equivocal.

Each eukaryotic organism has a characteristic chromosome complement, its karyotype, which represents the highest level of structural and functional organization of the nuclear genome (Stace, 2000). Karyotype constancy ensures the transfer of the same genetic material to the next generation, while karyotype variation provides genetic support to ecological differentiation and adaptation (Stebbins, 1971, Stace, 2000). Cytogenetic studies have shown that the tremendous inter- and intra-taxonomic variation of chromosome number documented in flowering plants (Levitzky, 1931; Stebbins, 1971; Raven, 1975) is mostly driven by two major mechanisms: a) increases through polyploidy (which may entail a Whole Genome Duplication [WGD] or an increase by half of the genome, demi-duplication [Mayrose *et al.,* 2009]); b) decreases or increases through duplication or deletion of single chromosome pairs or structural chromosomal rearrangements like chromosome fusion, i.e. descending dysploidy, and chromosome fission, i.e. ascending dysploidy.

Polyploidy is a common and ongoing phenomenon, especially in plants (Wood *et al.,* 2009), that has played an important role in many lineages, with evidence of several rounds of both ancient and recent polyploidization (Jiao *et al.,* 2011; Li *et al.,* 2015; Leebens-Mack *et al.,* 2019), albeit its distribution in time remains contested (Ruprecht *et al.,* 2017). Indeed, although the crucial role of polyploidy in plant diversification on small timescales is widely accepted (Stebbins, 1971; Grant, 1981), the evolutionary significance of polyplodization for the long-term diversity of angiosperms is still controversial (Mayrose *et al.,* 2011; Soltis *et al.,* 2014; Han *et al.,* 2020). On the other hand, while dysploidy is more frequent than polyploidy in angiosperms (Grant, 1981), its adaptive consequences have been mostly unexamined (Weiss-Schneeweiss & Schneeweiss, 2013), until recent studies demonstrated its high evolutionary impact (Escudero *et al.,* 2014).

In other seed plants, Li et al. (2015) detected two WGDs in the crown node of conifer clades, namely Pinaceae and Cupressaceae, contrary to a previous study suggesting a slow and

steady rate of genome size increase mainly due to the accumulation of repetitive DNA

(Nystedt et al., 2013). Several rounds of polyploidy, followed by diploidization, also occurred

in monilophyte ancestors (Clark et al., 2016), albeit these authors managed a limited

taxonomic sampling. Diploid chromosome numbers range from $2n = 14$ to 66 in extant

gymnosperms, and from $2n = 18$ to 1440 in monilophytes.

Chromosome number variation across angiosperm lineages spans two orders of magnitude

(Stace, 2000), from $2n = 4$ to $2n = 640$. Previous hypotheses (Ehrendorfer *et al.,* 1968;

Stebbins, 1971; Walker 1972; Raven, 1975; Grant, 1981; Soltis *et al.,* 2005) of the ancestral

basic (monoploid; see Peruzzi, 2013) chromosome number p in angiosperms suggest low

numbers, between $p = 6$ and $p = 9$. Genome size also vary tremendously across the flowering

plants (Leitch *et al*., 2005), and it is has been estimated that the ancestral genome size of

angiosperms was very small ($1C < 1.4$ pg, Leitch *et al*., 2005). Early hypotheses to estimate

putative ancestral basic chromosome numbers placed particular attention to 'primitive' extant

angiosperms (Raven, 1975). More recently, an ancestral chromosome number has been

reconstructed using a maximum parsimony approach (Soltis *et al.,* 2005). However, although

parsimony has been widely used to infer ancestral chromosome numbers, it carries significant

shortcomings (Mayrose *et al.,* 2009), and more rigorous and complex models to infer

chromosome number evolution are currently available (Mayrose *et al.,* 2009; Glick &

Mayrose, 2014; Freyman & Höhna, 2018; Zenil-Ferguson *et al.,* 2018). Here we use

probabilistic models, accounting for various types of chromosome number transitions, to

reconstruct the ancestral haploid chromosome number and the occurrence of chromosome

change events across the most massive data set ever assembled linking chromosome numbers

to a phylogeny, sampling 10,766 taxa from 318 families (73%) and 59 orders (92%) of

angiosperms. Chromosome numbers were extracted from the Chromosome Counts DataBase

(Rice *et al.,* 2015), and the analyses were conducted using pruned versions of two recently

published, dated mega-trees for seed plants (Smith & Brown, 2018). In addition, we explored

the sensitivity of our results by conducting all analyses again using a different ultrametric tree

of 1,559 taxa extracted from a recently published alternative angiosperm tree (Li *et al.,* 2019).

As WGD and post-polyploid genome diploidization (PPD) are widespread across

angiosperms (the latter gradually reverting the polyploid genome to one functionally diploid-

like through chromosomal rearrangements, see Mandakova & Lysak, 2018), it is critical to

consider changes in ploidy level besides chromosome number change dynamics. To this aim,

we compared our results about chromosome number evolution with estimated rates of

different shifts in ploidy level (including diploidization events, Zenil-Ferguson *et al.,* 2018)

with inferred ancestral states in genome size sampling over 5,000 taxa from the Plant DNA

C-values database (https://cvalues.science.kew.org/search/angiosperm). Ancestral genome

size inference taken in combination with chromosome number evolution, could allow us to

address genome rearrangements associated with genome size duplication across angiosperm

history.

Materials and Methods

Phylogenetic reconstruction

We used two recently published (Smith & Brown, 2018) dated megaphylogenies for seed

plants, GBMB and GBOTB, as backbones to generate two alternative phylogenies for

angiosperms included in the dataset. GBMB and GBOTB were constructed using 79,874 and

79,881 taxa, respectively, available in GenBank and in a backbone provided either by

Magallón et al. 2015 (GBMB) or by Open Tree of Life, version 9.1 (GBOTB). In addition,

we also used a different ultrametric tree provided by a recently published plastid

phylogenomic angiosperm (PPA) tree (Li *et al.,* 2019).

Chromosome numbers collection

The haploid chromosome numbers (*n*) of the species were obtained from the Chromosome

Counts Database (CCDB [Rice *et al.,* 2015]; http://ccdb.tau.ac.il/; last accessed May 2019)

using the R package chrome (Pennell *et al.,* 2016). CCDB contains records from original

sources that have irregularities of chromosome counts, so that the ca. 150,000 records were

curated semiautomatically using the CCDBcurator package (Rivero *et al.,* 2019). After a first

round of automatic cleaning, we examined results by hand using custom R scripts and

corrected records where needed.

Species with unknown chromosome counts were pruned from the trees, thus we collected

chromosome numbers for 10,766 taxa included in the GBMB and GBOTB phylogenetic

trees, and for 1,559 taxa included in the PPA tree. In cases where multiple chromosome

numbers were reported for a given taxon, the modal number was used (Mayrose *et al.,* 2009;

Salman-Minkov *et al.,* 2016). For taxa with numbers suggesting different ploidy levels, we

used the lowest haploid chromosome number available (Márquez-Corro *et al.,* 2019). This

coding scheme allowed us to deal with the problem of the existence of different ploidy levels

in a taxon and also with the low-density sampling conducted in most taxa (Márquez-Corro *et*

al., 2019). Analyses conducted using the PPA tree encountered less computation limitations,

so that we were able to perform them by explicitly considering intraspecific polymorphism,

allowing several chromosome numbers, together with their respective frequencies, to be set

for each taxon (Glick & Mayrose, 2014).

Genome size collection

Genome size data (1C, prime estimates) for angiosperms were taken from the Plant DNA C-

values database (https://cvalues.science.kew.org/search/angiosperm, release 7.1, March

2020), managed by the Royal Botanic Gardens, Kew (Pellicer & Leitch, 2019). We gathered

5,581 and 661 taxa, respectively matching the GBMB and the PPA trees. Genome size data

were log_{10} transformed to ensure the data conformed to Brownian motion evolution (Beaulieu

et al., 2012; Carta & Peruzzi, 2016).

Analyses

The evolution of haploid chromosome numbers of angiosperms was inferred using

chromEvol software v.2.0 (http://www.tau.ac.il/~itaymay/cp/chromEvol/index.html; Glick &

Mayrose, 2014). This software determines the likelihood of a model to explain the given data

along the phylogeny, based on the combination of two or more of the following parameters:

dysploidization (ascending, chromosome gain rate λ; descending, chromosome loss rate δ),

polyploidization (chromosome number duplication with rate ρ, demi-polyploidization or

triploidization with rate μ) and incremental changes to the basic number (β) with regard to a

rate of multiplication (v) that is different from a regular duplication (Mayrose *et al.,* 2009).

Two additional parameters (λ_1, δ_1) detect linear dependency between the current haploid

number and the rate of gain and loss of chromosomes. We tested 10 models based on a

different combination of the parameters above. Four of these models consider only constant

rates (Mc1, Mc2, Mc3, and Mc0), whereas the other four include two linear rate parameters

(Ml1, Ml2, Ml3, and Ml0; Table 3). Both sets have a null model (Mc0 and Ml0) that assumes

no polyploidisation events. Finally, two models (Mb1 and Mb2) consider that the evolution

of chromosome number can also be influenced by the basic number and by its transition rates.

The minimum chromosome number allowed in the analyses was set to 2, whereas the

maximum number was set to 5 units higher than the highest chromosome number found in

the empirical data. We removed all counts $n > 43$ from the analysis (5,467 counts [3,001

taxa]), because for many lineages the sampling was inadequate to reconstruct such a drastic

change in chromosome number (Márquez-Corro *et al.,* 2019; Barrett *et al.,* 2019) and

because of computation limitations (Zenil-Ferguson *et al.,* 2018). The branch lengths were

scaled according to the software author's instructions. The null hypothesis (no polyploidy)

was tested with likelihood ratio tests using the Akaike information criterion (AIC; Burnham

& Anderson, 2004). To compute the expected number of changes along each branch, as well

as the ancestral haploid chromosome numbers at internal nodes, the best fitted model for both

data sets was rerun using 1,000 simulations. The best model was plotted on the trees using the

ChromEvol functions v0.9-1 elaborated by N. Cusimano in R.

To test which ancestral haploid chromosome number is most likely at the root of

angiosperms, the following haploid chromosome numbers were fixed at the root and the

likelihood of the resulting models was compared: $n = 4,5,6,7,8,9$. These numbers were tested

either because considered putative ancestral character-states (Ehrendorfer *et al.,* 1968;

Stebbins, 1971; Walker 1972; Raven, 1975; Grant, 1981; Soltis *et al.,* 2005), or because they

were identified as the chromosome numbers showing the highest PP under our Bayesian

analysis.

To further support our results, we also modelled haploid chromosome number evolution

using the function Q_chromevolM3 in the R package chromploid (Zenil-Ferguson *et al.*,

2018). We then modeled ploidy change using the function PloidEvol and a subset of 3,759

taxa used in the ChromEvol analysis that have ploidy level information in

https://cvalues.science.kew.org/search/angiosperm. The PlodiEvol model, implemented in the

package chromploid, is a continuous time Markov chain model of ploidy level evolution, that

estimates the rates of different shifts in ploidy level, including diploidization, across a

phylogenetic tree, by analyzing the transitions of ploidy level among species. Optimizations

were performed using a subplex algorithm (nloptr package, Ypma 2014), where the

convergence criterion was based on the changes in the precision of the negative log

likelihood of no more of 10^{-8}. The procedure was run for at least 1,000 iterations.

Ancestral genome sizes were reconstructed using maximum likelihood and visualized on the

tree with the contmap function in the R package phytools (Revell, 2012). Models of adaptive

evolution at macroevolutionary scales typically rely on the Ornstein-Uhlenbeck (OU) model

of trait evolution (Hansen, 1997). This model assumes that an optimum trait value identifies a

selective regime acting on a lineage over the course of its history. The rate of adaptive

evolution towards an optimum trait value θ is governed by α, while the constant σ^2 describes

the rate of stochastic evolution away from the optimum. Here, discrete regime shifts in

genome size on the phylogeny were estimated from the data using a Bayesian reversible-

jump multi-optima OU model in the R package bayou (Uyeda & Harmon, 2014). Two

independent chains were run for at least 200,000 generations, with a thinning interval of 20

and the first 60,000 generations discarded as burn-in. To check the convergence of the chains
we used the Gelman's R. A half-Cauchy distribution was assigned to α and σ^2, a normal
distribution with mean and standard deviation equivalent to the distribution of values for
genome size was assigned to θ, and a uniform distribution was assigned to the probability
that a change in optima occurs on a given branch, with the restriction that only one change
can occur on any branch. Mean genome size was estimated for each species, while within-
species variation to be incorporated into the phylogenetic analysis was not estimated
separately for each species, because within-species samples were limited to a few samples.
OU models are particularly affected by measurement error. Hence, we estimated the pooled
variance across the species and used it, weighted by sample size, to estimate the observation
variance of the individual species. All analyses were performed in the high-performance
computing cluster at the University of Pisa.

Results

Regardless of the three alternative phylogenies, $n = 7$ was inferred as the ancestral haploid
chromosome number with the highest posterior probability (Table 1) and likelihood (Table
2). The ancestral haploid chromosome number $n = 7$ was remarkably stable in the deepest
part of the phylogeny (Fig. 1), while slight variations ($\pm$ 1) in n were inferred at the base of
some lineages. Greater variations were shown in the ancestral haploid chromosome number
of many angiosperm families (see Supplementary Table S1). Monocots exhibited the largest
variation of inferred n among Most Recent Common Ancestors (MRCAs) of plant families
(Fig. 2b), paralleled by a considerable variation in current haploid chromosome numbers
(Fig. 2a). Over 70% of inferred n in the 158 families for which previous inferences were
available are in line with previous proposals. The most interesting disagreements among our
inferences and those previously available in literature are reported in Table 3. For the
remaining 160 families (50.3%) the first inferences are presented here.
For both GBMB and GBOTB phylogenies, the best model considers up to six parameters
(Table 4), i.e. chromosome gain, loss, duplication, demi-duplication rates and rates of gain
and loss linearly dependent on the current chromosome number. Specifically, descending
dysploidy, most likely through chromosome fusion, was the most common cytogenetic
mechanism of chromosome number change during the evolution of flowering plants, and this
is inferred both on branches leading to major clades and on terminal branches

(Supplementary Figs. 1-3). Polyploidization events were also inferred in a significant

number, albeit mainly on terminal branches (Supplementary Figs. 1-3). PloidEvol, having the

ability to define a wider range of increasing ploidy level change events than chromosome

number-based models, detected a small rate of pure duplication for even ploidy levels (ρ =

0.024). Instead, PloidEvol indicate that other two parameters associated with the formation of

even ploidy levels are the largest: the rates of even-to-odd increase (α = 0.697) and of

diploidization (δ = 0.471) and even-to-even transition (ε = 0.116). The stationary distribution

under Ploidevol model indicates a diploid status of the root, with highest probability (0.74)

followed by a tetraploid status (0.18). Analyses conducted using the PPA tree, include a

lower number of sampled taxa but allowed to consider intraspecific chromosome number

variation for each taxon. Nevertheless, results were consistent with those obtained using the

GBMB and GBOTB phylogenies. We found indeed only minor differences at the root

(Tables 1,2) and at some internal nodes (Fig. 1).

The results above are also confirmed by those obtained through the chromploid software,

concerning both the ancestral haploid chromosome number marginal probability (n = 7, 0.54;

n = 8, 0.43; n = 9, 0.01) and the rate parameters, with descending dysploidy (δ = 0.017) and

chromosome doubling (ρ = 0.012) largely exceeding chromosome gain (λ = 0.009) and

demiploidization (μ = 0.005).

The reconstruction of genome size variation across the evolutionary history of angiosperms

indicates that the ancestral holoploid genome size for angiosperms was small (1C = 1.73 pg)

and mostly stable in the basal part of the tree, showing instead decreasing or increasing

values along terminal branches (Fig. 3 and Figs. S4–S8). In addition, the multi-optima OU

model identified several significant shifts in genome size, particularly an increase in genome

size at the base of Monocots and a downsizing at base of Rosids (Figs. S6-7).

Inferred ancestral 1C values for angiosperm families are reported in Table S1. Overall, the

majority of nodes (57%) underwent a decrease in genome size with respect to the ancestral

value of 1C = 1.73 pg (0.24 in Fig. 3). In addition, a value 2-fold the ancestral one (0.48 in

Fig. 3) was reached for the first time only after more than 30 million years of angiosperm

evolutionary history (line in Fig. 3), in the lineage giving rise to Monocots.

Discussion

The main results presented here were drawn using the largest dated mega-tree currently

available for seed plants (but see Janssens *et al.*, 2020). Bayesian inference revealed an

ancestral haploid chromosome number for angiosperms $n = 7$, reinforcing previous

hypotheses (Ehrendorfer *et al.,* 1968; Stebbins, 1971; Walker 1972; Raven, 1975; Grant,

1981; Soltis *et al.,* 2005) that suggested a low ancestral basic number.

As a by-product of our study, our analyses allowed to infer n for single families, more than

half of which are provided here for the first time, while the others are mostly congruent with

previous evaluations. However, were able to highlight a number of discrepancies in ancestral

chromosome numbers previously inferred for a number of families (Table 3), possibly due to

traditional routine algebraic approaches, instead of phylogenetic models, to infer

chromosome number changes (see Cusimano *et al.,* 2012 for further discussion), or to the

analysis of more limited datasets.. For example, the inferred n of MRCAs of Brassicaceae,

Lamiaceae, and Rosaceae are respectively 7, 8, and 7 in our study, but were previously

inferred as 12, 14, and 9, respectively (Raven, 1975). Indeed, even in the presence of a strong

phylogenetic signal (e.g., closely related species sharing similar chromosome numbers;

Escudero *et al.,* 2012; Carta *et al.,* 2018), algebraic inferences of chromosome numbers

become increasingly difficult with increasing phylogenetic depth, as identical chromosome

numbers will occur in unrelated lineages (Weiss-Schneeweiss & Schneeweiss, 2013). The

dataset analysed here is the most extensive ever used for inferring ancestral haploid number

in angiosperms, but it still poses challenges concerning incomplete taxon sampling and

phylogenetic resolution at the family level.

Our chromosome number evolution results support the conclusion that genome duplication

and descending dysploidy were critical events in the evolution of angiosperms (Escudero *et*

al., 2014). Whilst polyploidization is the second most frequent transition type, ancient

polyploidy events are underrepresented. The absence of polyploidization events at the base of

the tree is in agreement with the maintenance of the ancestral haploid chromosome number n

$= 7$ inferred in the deepest part of the phylogeny, with the inferred ancestral diploid status and

with the very low ancestral genome size. Polyploidization events were instead inferred

mainly toward the tips of the tree, partially supporting previous evidence revealing

independent genome duplications near the base of several families (Escudero *et al.,* 2014;

Leebens-Mack *et al.,* 2019; Cusimano *et al.,* 2012), leading to high haploid chromosome

numbers. Indeed, this may be the case of many families in the Magnoliid clade. Our results

do not support a link among some of the most extensive plant radiations and ancient

polyploidization rounds (Jiao *et al.,* 2011). Nevertheless, the macroevolutionary dynamics of

ploidy level changes here presented, emphasize the importance of diploidization events,

suggesting that both WGD and PPD are widespread phenomena in angiosperms. Patterns of

early WGD were suggested also in gymnosperms (Li *et al*., 2015), where however polyploidy

is rare (Rastogi & Ohri, 2020). On the contrary, repeated WGD events in the ancestors of

monilophytes have contributed to their diversity and high chromosome numbers (Clark *et al*.,

2016). Inferring ancient polyploidy events from cytological data is a challenging task,

because diploidization events (Schubert & Lysak, 2011; Mandakova & Lysak, 2018)

following polyploidisation gradually can hide signals of genome duplication over time

(Mayrose *et al*., 2009). Yet, in our analyses, descending dysploidy was indeed the most

common cytogenetic mechanism of chromosome number change, and this phenomenon is

often regarded as a part of the diploidization process (Mandakova & Lysak, 2018). Our

results on chromosome number evolution do not support WGD events at the origin of

angiosperms or before it (Ruprecht *et al.,* 2017), but rather highlight the importance of WGD

events later in the evolution of angiosperms. This is further corroborated by our results on

genome size evolution, which clearly indicate that a) the ancestral genome size was small (1C

= 1.73 pg, a value slightly higher than the inference by Leitch *et al*., 2005) and mostly stable

in the basal part of the tree; b) WGD likely did not occur in angiosperms for the first 30

million years of their evolutionary history (Fig. 3); and c) the majority of angiosperm nodes

(57%) underwent a general genome size reduction.

Reconstructing the ancestral chromosome number is difficult, because there are no suitable

outgroups for direct comparison (Doyle, 2012), and because extant early branching

angiosperms (e.g., *Amborella* and Nymphaeales) are not necessarily holding plesiomorphic

character-states. With these limitations in mind, we made inferences based on the distribution

of *n* and genome size in extant angiosperms, and using probabilistic models accounting for

various types of chromosome number transitions. Our study is not able to address the origin

of either the chromosome number or genome size of the first angiosperms. Instead, it

provides novel, detailed, and well-supported inference of ancestral haploid chromosome

number and genome size of the common ancestor of all extant angiosperms. Interestingly, our

inferred ancestral state for the haploid number *n* coincides with the ancestral basic

chromosome number *p* previously proposed for angiosperms based on empirical counts

(Ehrendorfer *et al.,* 1968; Stebbins, 1971; Walker 1972; Raven, 1975; Grant, 1981; Soltis *et*

al., 2005) or paleogenomic approaches (Salse *et al.,* 2012). In gymnosperms, the

hypothesised ancestral number *n* = 12 (Nystedt *et al*., 2013), based on the most frequent

chromosome number in extant species, is partly supported by the reconstructed ancestral

number in *Juniperus*, *n* = 11 (Farhat *et al*., 2019). Not surprisingly, a higher ancestral

chromosome number has been reconstructed for monilophytes, namely *n* = 22 (Clark *et al*.,

2016). According to Leitch *et al.* (2005) both gymnosperms and monilophytes showed an
"intermediate" ancestral genome size (1C values ranging between 3.5 and 14.0 pg).
Altogether, it appears that spermatophytes may have a lower ancestral chromosome number,
compared to monilophytes, suggesting differences between the genomic evolutionary
trajectories of these groups (Clark *et al.*, 2016).

Our study allowed to clarify a long-standing question (Stebbins, 1971), but such
reconstruction necessarily comes with limitations. We do believe our results allowed a major
step forward in understanding the ancestral chromosome number for angiosperms, and we
believe that this issue should be added to the angiosperm macroevolutionary agenda (Sauquet
& Magallón, 2018). Further progress in reconstructing the ancestral genome organisation in
angiosperms may require the development of models that include heterogeneity in the
patterns of chromosome evolution across a phylogenetic tree (Zenil-Ferguson *et al.,* 2018),
along with a deeper insight into genome and karyotype evolution.

Acknowledgements
The authors thank Marcial Escudero and Itay Mayrose for their help with ChromEvol
analyses. We are also grateful to Rosana Zenil-Ferguson for her help with the chromploid
package.

Author contributions
A.C. planned and designed the research, analysed the data and wrote the manuscript. G.B.
assisted in chromosome numbers acquisition. L.P. and G.B. contributed to successive
versions of the manuscript and in solving theoretical and nomenclatural issues. All authors
read and approved the final manuscript.

Data Availability
All data analysed during this study are included in this published article and its
supplementary information.

References
Barrett CF, McKain MR, Sinn BT, Ge XJ, Zhang Y, Antonelli A, Bacon CD. 2019.
Ancient polyploidy and genome evolution in palms. *Genome Biology and Evolution* **11**:
1501–1511.

Beaulieu JM, Jhwueng DC, Boettiger C, O'Meara BC. 2012. Modeling stabilizing selection: expanding the Ornstein–Uhlenbeck model of adaptive evolution. *Evolution* **66**: 236–2383.

Burnham KP, Anderson DR. 2004. Model inference. Understanding AIC and BIC in model selection. *Sociological Methods and Research* **33:** 261–304.

Carta A, Peruzzi L. 2016. Testing the large genome constraint hypothesis: plant traits, habitat and climate seasonality in Liliaceae. *New Phytologist* **210**: 709–716.

Carta A, Bedini G, Peruzzi L. 2018. Unscrambling phylogenetic effects and ecological determinants of chromosome number in major angiosperm clades. *Scientific Reports* **8**: 1–14.

Clark J, Hidalgo O, Pellicer J, Liu H, Marquardt J, Robert Y, Christenhusz M, Zhang S, Gibby M, Leitch IJ, Schneider H. 2016. Genome evolution of ferns: evidence for relative stasis of genome size across the fern phylogeny. *New Phytologist* **210**: 1072–1082.

Coghlan A, Eichler EE, Oliver SG, Paterson AH, Stein L. 2005. Chromosome evolution in eukaryotes: a multi-kingdom perspective. *Trends in Genetics* **21**, 673–682.

Crowl AA, Miles NW, Visger CJ, Hansen K, Ayers T , Haberle R , Cellinese N. 2016. A global perspective on Campanulaceae: Biogeographic, genomic, and floral evolution. *American Journal of Botany* **103**: 233–245.

Cusimano N, Sousa A, Renner SS. 2012. Maximum likelihood inference implies a high, not a low, ancestral haploid chromosome number in Araceae, with a critique of the bias introduced by 'x'. *Annals of Botany* **109**: 681–692.

Doyle JA. 2012. Molecular and fossil evidence on the origin of angiosperms. *Annual Review of Earth and Planetary Sciences* **40**: 301–326.

Ehrendorfer F, Krendl F, Habeler E, Sauer W. 1968. Chromosome numbers and evolution in primitive angiosperms. *Taxon* **17**: 337–353 (1968).

Escudero M, Martín-Bravo S, Mayrose I, Fernández-Mazuecos M, Fiz-Palacios O, Hipp AL et al. 2014. Karyotypic changes through dysploidy persist longer over evolutionary time than polyploid changes. *PLOS One* **9**, e85266.

Escudero, M, Hipp AL, Hansen TF, Voje KL, Luceño M. 2012. Selection and inertia in the evolution of holocentric chromosomes in sedges (*Carex*, Cyperaceae). *New Phytologist* **195**: 237–247.

Farhat P, Hidalgo O, Robert T, Siljak-Yakovlev S, Leitch IJ, Adams RP, BouDagher-Kharrat M. 2019. Polyploidy in the Conifer Genus *Juniperus*: An Unexpectedly High Rate. *Frontiers in Plant Sciences* 10, doi: 10.3389/fpls.2019.00676.

Freyman WA, Höhna S. 2018. Cladogenetic and anagenetic models of chromosome number evolution: a Bayesian model averaging approach. *Systematic Biology* **67**: 195–215.

Glick L, Mayrose I. 2014. ChromEvol: assessing the pattern of chromosome number evolution and the inference of polyploidy along a phylogeny. *Molecular Biology and Evolution* **31**: 1914–1922.

Grant V. 1981. *Plant Speciation* (ed. 2). New York: Columbia University Press.

Guerra M. 2008. Chromosome numbers in plant cytotaxonomy: concepts and implications. *Cytogenetic and Genome Research* **120**: 339–350.

Han TS, Zheng QJ, Onstein RE, Rojas⬜Andrés BM, Hauenschild F, Muellner⬜Riehl AN, Xing YW. 2020. Polyploidy promotes species diversification of *Allium* through ecological shifts. *New Phytologist* **225**: 571–583.

Hansen TF. 1997. Stabilizing selection and the comparative analysis of adaptation. *Evolution* 51, 1341–1351.

Janssens S, Couvreur TL, Mertens A, Dauby G, Dagallier LP, Abeele SV et al. 2020. A large-scale species level dated angiosperm phylogeny for evolutionary and ecological analyses. *Biodiversity Data Journal*, **8**: e39677.

Jiao Y, Wickett NJ, Ayyampalayam S, Chanderbali AS, Landherr L, Ralph PE et al. 2011. Ancestral polyploidy in seed plants and angiosperms. *Nature* **473**: 97.

Leebens-Mack JH, Barker MS, Carpenter EJ et al. 2019. One thousand plant transcriptomes and the phylogenomics of green plants. *Nature* **574**: 679–685.

Leitch IJ, Soltis DE, Soltis PS, Bennett MD. 2005. Evolution of DNA amounts across land plants (Embryophyta). *Annals of Botany* **95**: 207–217.

Levitzky GA. 1931. The karyotype in systematics. *Bulletin of Applied Botany of Genetics and Plant Breeding* **27**: 220–240.

Li HT, Yi TS, Gao LM, Ma PF, Zhang T, Yang JB, et al. 2019. Origin of angiosperms and the puzzle of the Jurassic gap. *Nature Plants* **5**: 461–470.

Li Z, Baniaga AE, Sessa EB, Scascitelli M, Graham SW, Rieseberg LH, Barker MS. 2015. Early genome duplications in conifers and other seed plants. *Science Advances* **1**: e1501084.

Lysak MA. 2018. Brassicales: an update on chromosomal evolution and ancient polyploidy. *Plant Systematics and Evolution* **304**: 757–762.

Magallón S, Gómez⬜Acevedo S, Sánchez⬜Reyes LL, Hernández⬜Hernández TA. 2015. A metacalibrated time⬜tree documents the early rise of flowering plant phylogenetic diversity. *New Phytologist* **207**: 437–453.

Mandakova T, Lysak MA. 2018. Post-polyploid diploidization and diversification through dysploid changes. *Current Opinion in Plant Biology* **42**: 55–65.

Márquez-Corro JI, Martín-Bravo S, Spalink D, Luceño M, Escudero M. 2019. Inferring hypothesis-based transitions in clade-specific models of chromosome number evolution in sedges (Cyperaceae). *Molecular Phylogenetic and Evolution* **135**: 203–209.

Mayrose I, Zhan SH, Rothfels CJ, Magnuson-Ford K, Barker MS, Rieseberg LH, Otto SP. 2011. Recently formed polyploid plants diversify at lower rates. *Science* **333**: 1257–1257.

Mayrose I, Barker MS, Otto SP. 2009. Probabilistic models of chromosome number evolution and the inference of polyploidy. *Systemtic Biology* **59**: 132–144.

Nystedt B, Street NR, Wetterbom A, Zuccolo A, Lin Y-C, Scofield DG et al. 2013. The Norway spruce genome sequence and conifer genome evolution. *Nature* **497**: 579–584.

Oliveira Marinho MA, Souza G, Felix LP, De Carvalho F. 2019. Comparative cytogenetics of the ACPT clade (Anacampserotaceae, Cactaceae, Portulacaceae, and Talinaceae): a very diverse group of the suborder Cactineae, Caryophyllales. *Protoplasma* **256**: 805–814.

Ørgaard M. 1991. The genus *Cabomba* (Cabombaceae)–a taxonomic study. *Nordic Journal of Botany* **11**: 179–203.

Pellicer J, Kelly LJ, Leitch IJ, Zomlefer WB, Fay MF. 2014. A universe of dwarfs and giants: genome size and chromosome evolution in the monocot family Melanthiaceae. *New Phytologist* **201**: 1484–1497.

Pellicer J, Kelly LJ, Magdalena C, Leitch IJ. 2013. Insight into the dynamics of genome size and chromosome evolution in the early diverging angiosperm lineage Nymphaeales (water lilies). *Genome* **56**: 437–439.

Pellicer J, Leitch IJ. 2019. The Plant DNA C☐values database (release 7.1): an updated online repository of plant genome size data for comparative studies. *New Phytologist* doi: 10.1111/nph.16261

Pennell MW. 2016. Chromer: Interface to Chromosome Counts Database API. R package version 0.1.2.9000.

Peruzzi L. 2013. "x" is not a bias, but a number with real biological significance. *Plant Biosystems* 147: 1238–1241.

Rastogi S, Ohri D. 2020. Chromosome Numbers in Gymnosperms - An Update. *Silvae Genetica* 69: 13–19.

Raven PH. 1975. The bases of angiosperm phylogeny: cytology. *Annals of the Missouri Botanical Garden* **62**: 724–764.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in ecology and evolution* **3**: 217–223.

Rice A, Glick L, Abadi S, Einhorn M, Kopelman NM, Salman☐Minkov A et al. 2015. The Chromosome Counts Database (CCDB)–a community resource of plant chromosome numbers. *New Phytol*. **206**: 19–26.

Rivero R, Sessa EB, Zenil☐Ferguson R. 2019. EyeChrom and CCDB curator: Visualizing chromosome count data from plants. *Applications in Plant Sciences*, **7**, e01207.

Ruprecht C, Lohaus R, Vanneste K, Mutwil M, Nikoloski Z, Van de Peer Y, Persson S. 2017. Revisiting ancestral polyploidy in plants. *Science Advances* **3**: e1603195.

Salman-Minkov A, Sabath N, Mayrose I. 2016. Whole-genome duplication as a key factor in crop domestication. *Nature Plants* **2**: 16115.

Salse J. 2012. In silico archeogenomics unveils modern plant genome organization, regulation and evolution. *Current Opinion in Plant Biology* **15**: 122–130.

Sauquet H., Magallón S. 2018. Key questions and challenges in angiosperm macroevolution. *New Phytologist* **219**: 1170–1187.

Schubert I, Lysak MA. 2011. Interpretation of karyotype evolution should consider chromosome structural constraints. *Trends in Genetics* **27**: 207–216.

Smith SA, Brown JW. 2018. Constructing a broadly inclusive seed plant phylogeny. *American Journal of Botany* **105**: 1–13.

Soltis DE, Segovia☐Salcedo MC, Jordon☐Thaden I, Majure L, Miles NM, Mavrodiev EV et al. 2014. Are polyploids really evolutionary dead☐ends (again)? A critical reappraisal of Mayrose et al.(2011). *New Phytologist* **202**: 1105–1117.

Soltis DE, Soltis PS, Endress PK, Chase MW. 2005. *Phylogeny and Evolution of Angiosperms*. Sunderland: Sinauer Associates.

Stace CA. 2000. Cytology and cytogenetics as a fundamental taxonomic resource for the 20th and 21st centuries. *Taxon* **49**: 451–477.

Stebbins GL. 1971. *Chromosomal Evolution in Higher Plants*. London: Edward Arnold.**Uyeda JC, Harmon LJ. 2014.** A novel Bayesian method for inferring and interpreting the dynamics of adaptive landscapes from phylogenetic comparative data. *Systematic Biology* **63**: 902–918.

Walker JW. 1972. Chromosome numbers, phylogeny, phytogeography of the Annonaceae and their bearing on the (original) basic chromosome number of angiosperms. Taxon **21**: 57–65.

Weiss-Schneeweiss H, Schneeweiss GM. 2013. in *Plant Genome Diversity* Vol. 2 (eds.

Leitch, I.J., Greilhuber, J., Dolezel, J.W.J) 209–230. Vienna: Springer.

Wood, T. E., Takebayashi, N., Barker, M. S., Mayrose, I., Greenspoon, P. B., &

Rieseberg, L. H. 2009. The frequency of polyploid speciation in vascular plants.

Proceedings of the National Academy of Sciences **106**: 13875–13879.

Ypma J. 2014. Nloptr: R Interface To Nlopt. R Package Version, 1. CRAN Repositories.

Zenil-Ferguson R, Burleigh JG, Ponciano JL. 2018. Chromploid: an R package for

chromosome number evolution across the plant tree of life. *Applications in Plant Science* **6**:

e1037 10.1002/aps3.1037.

Figure 1 Reconstruction of ancestral haploid chromosome number (*n*) of angiosperms with

the best-fitting model on the different types of trees used (GBMB and PPA). Please note that

the plotted trees depict ordinal phylogenetic relationships (sub-ordinal topologies were

collapsed to build the figure), and are shown without branch length information. Pie charts at

nodes represent the probability of the ancestral haploid chromosome numbers inferred under

Bayesian estimation; the numbers at nodes are those with the highest probability. Pie charts

and numbers at the tips are the three best inferred ancestral haploid chromosome numbers per

each angiosperm order. Black lines, difference in phylogenetic position between GBMB and

PPA trees.

Figure 2 Density plots of haploid chromosome numbers (n) and of inferred ancestral haploid chromosome number (n) for each angiosperm family in four major angiosperm clades (APG IV). **a**, we identified the number of unique chromosome counts per taxon, i.e. cytotypes, from the original dataset, after excluding counts with $n > 60$, to focus on the most frequent n and their putative relation with inferred p. **b**, density plots were scaled by the Bayesian posterior probability (PP) estimated for each inferred p.

Figure 3 Projection of the tree into phenotype space (traitgram) for genome size evolution in angiosperms (GBMB tree). Ancestral states for the projection and discrete shifts in genome size on the phylogeny were estimated using a Bayesian reversible-jump multi-optima OU model. Branches are coloured according to their median value of theta, with the larger phenotypic values for optima being indicated by blue, green and orange while smaller optima indicated by pink. The straight dashed line coincides with a 2-fold ancestral genome size.

552 **Table 1.** Summary of chromosome number evolutionary models and inferred ancestral haploid chromosome number (*n*) in Angiosperms under

553 the best-fitting model.

tree	Best model	LogLik	AIC	Rates						Events inferred with PP > 0.5				Chromosome no. at root node		
				λ	δ	ρ	μ	λ_l	δ_l	Gain	Losses	Duplications	Demi	Bayes (PP): Best *p*	Bayes(PP): 2nd best *p*	ML
GBMB (10,766 taxa)	Ml3	-18120.0	36250.0	0.0081	0.0113	0.0131	0.0051	0.0051	0.0007	509.7	1627.5	1438.1	376.1	7 (0.97)	8 (0.02)	5
GBOTB (10,766 taxa)	Ml3	-18110.0	36240.0	0.0106	0.0096	0.0130	0.0049	0.0001	0.0008	589.5	1625.3	1435.3	363.3	7 (0.98)	8 (0.01)	5
PPA (1559 taxa)	Ml2	-3788.0	7586.0	0.0101	0.0044	0.0078	-	-0.0002	0.0012	145.0	528.1	244.6	191.9	7 (0.24)	5 (0.21)	2

Only the best-fitting models are shown. Tree refers to the three alternative phylogenies used. Best model, Ml3 (linear rate model with duplication rate ρ and demi-duplication rate μ) Ml2 (linear rate model with equal duplication and demi-duplication rates); Logarithmic likelihood (LogLik) and AIC scores; rate parameters (λ = chromosome gain rate, δ = chromosome loss rate, ρ = duplication rate, μ = demi-duplication rate, λ_l= linear chromosome gain, δ_l = linear chromosome loss); frequency of the four possible event types with a posterior probability (PP) > 0.5; haploid chromosome number inferred at the root node under Bayesian optimization with the respective PP, and under maximum likelihood (ML).

555 **Table 2.** Testing hypotheses about root ancestral haploid chromosome number (n).

Tree	GBMB		GBOTB		PPA	
	LogLik	AIC	LogLik	AIC	LogLik	AIC
Root fixed at $n = 4$	-18127.1	36266.2	-18129.9	36271.7	-3788.8	7587.6
Root fixed at $n = 5$	-18128.1	36268.2	-18142.6	36297.2	-3787.2	7584.5
Root fixed at $n = 6$	-18123.6	36259.2	-18124.2	36260.4	-3786.7	7583.5
Root fixed at $n = 7$	**-18117.1**	**36246.3**	**-18115.5**	**36243.0**	**-3786.2**	**7582.4**
Root fixed at $n = 8$	-18119.4	36250.9	-18127.8	36267.6	-3787.6	7585.1
Root fixed at $n = 9$	-18122.2	36256.4	-18121.3	36254.7	-3788.4	7586.9

AIC and LogLik values obtained by fixing the root with given ancestral haploid chromosome number.
The lowest AIC and LogLik values are shown in bold.

568 **Table 3**. Inferred ancestral n chromosome numbers of most recent common ancestors (MRCAs) for selected angiosperm families. For each

569 family, the three most supported numbers are given (the first most supported in boldface). Only those 52 out of 318 families for which previous -

570 and different - hypotheses were postulated in literature are reported. For further details see Table S1.

family	*n* inferred in this study	*n* (literature)	notes
Acanthaceae	**9**=0.991, 10=0.007, 8=0.002	7	Raven (1975) hypothesized $n = 7$.
Amborellaceae	**7**=0.974, 8=0.024, 6=0.002	13	Raven (1975) hypothesized $n = 13$, which coincides with the current *x* typical of *Amborella trichopoda*, the sole species belonging to this family.
Anacardiaceae	**15**=0.602, 16=0.138, 14=0.106	7	Raven (1975) hypothesized $n = 7$.
Araliaceae	**8**=0.809, 6=0.119, 9=0.031	12	Raven (1975) hypothesized $n = 12$.
Arecaceae	**9**=0.779, 8=0.109, 10=0.101	18, 16	Raven (1975) hypothesized $n = 18$. This number, together with $n = 16$, is also inferred by Cusimano *et al.* (2012). The relevant difference with the latter authors may be due to differences in the number of outgroup taxa, much more limited in Cusimano *et al.* (2012).
Austrobaileyaceae	**7**=0.726, 6=0.145, 8=0.099	22	Raven (1975) hypothesized $n = 22$.
Bignoniaceae	**9**=0.974, 10=0.023, 18=0.002	20	Raven (1975) hypothesized $n = 20$.
Brassicaceae	**7**=0.590, 8=0.340, 6=0.035	12	Raven (1975) hypothesized $n = 12$.
Cabombaceae	**7**=0.378, 8=0.189, 6=0.179	13	Ørgaard (1991) hypothesized $n = 13$.
Campanulaceae	**8**=0.773, 9=0.146, 7=0.073	7, 9	Raven (1975) hypothesized $n = 7$ (our third most likely inferred *n*), while more recently Crowl *et al.* (2016) hypothesized $n = 9$ (our second most likely inferred *n*).
Canellaceae	**7**=0.806, 6=0.108, 8=0.049	14, 13	Raven (1975) hypothesized $n = 14, 13$.
Caricaceae	**7**=0.479, 8=0.187, 6=0.114	9	Both Raven (1975) and Lysak (2018) hypothesized $n = 9$.
Cercidiphyllaceae	**9**=0.159, 10=0.138, 8=0.120	19	Raven (1975) hypothesized $n = 19$.
Circaeasteraceae	**7**=0.477, 8=0.217, 6=0.146	15	Raven (1975) hypothesized $n = 15$.
Cyrillaceae	**6**=0.368, 7=0.247, 8=0.157	10	Raven (1975) hypothesized $n = 10$.
Degeneriaceae	**7**=0.331, 6=0.231, 5=0.111	12	Raven (1975) hypothesized $n = 12$.
Dioncophyllaceae	**6**=0.544, 7=0.292, 5=0.099	18	Raven (1975) hypothesized $n = 18$.
Escalloniaceae	**12**=0.991, 13=0.005, 6=0.002	8, 9	Raven (1975) hypothesized $n = 8, 9$.
Eucommiaceae	**9**=0.270, 8=0.263, 11=0.164	17	Raven (1975) hypothesized $n = 17$.
Fouquieriaceae	**6**=0.538, 7=0.159, 8=0.125	12	Raven (1975) hypothesized $n = 12$.
Gomortegaceae	**11**=0.250, 12=0.177, 22=0.136	21	Raven (1975) hypothesized $n = 21$.

Hamamelidaceae	**6=0.483**, 7=0.323, 8=0.140	12, 16	Raven (1975) hypothesized *n* = 12, 16.
Hernandiaceae	**7=0.299**, 6=0.240, 12=0.154	20, 15	Raven (1975) hypothesized *n* = 20, 15.
Himantandraceae	**7=0.331**, 6=0.231, 5=0.111	12	Raven (1975) hypothesized *n* = 12.
Hydrangeaceae	**12=0.418**, 9=0.256, 13=0.200	7	Raven (1975) hypothesized *n* = 7.
Hypericaceae	**9=0.411**, 10=0.146, 7=0.138	12	Raven (1975) hypothesized *n* = 12.
Juncaginaceae	**12=0.273**, 8=0.235, 11=0.165	6, 10	Raven (1975) hypothesized *n* = 6, while Cusimano *et al.* (2012) inferred *n* = 10.
Lamiaceae	**8=0.765**, 9=0.234, 7=0.000	14	Raven (1975) hypothesized *n* = 14.
Lardizabalaceae	**8=0.345**, 11=0.157, 7=0.153	16, 15, 14	Raven (1975) hypothesized *n* = 16, 15, 14.
Loasaceae	**12=0.599**, 13=0.180, 11=0.121	7	Raven (1975) hypothesized *n* = 7.
Loranthaceae	**6=0.822**, 7=0.083, 8=0.064	12	Raven (1975) hypothesized *n* = 12.
Melanthiaceae	**7=0.335**, 8=0.297, 6=0.229	9	Pellicer *et al.* (2014) hypothesized *n* = 9.
Moringaceae	**7=0.479**, 8=0.187, 6=0.114	14	Lysak (2018) hypothesized *n* = 14.
Myricaceae	**8=0.981**, 9=0.014, 7=0.005	16	Raven (1975) hypothesized *n* = 16.
Nymphaeaceae	**7=0.536**, 8=0.184, 6=0.128	12	While Raven (1975) hypothesized *n* = 12, more recently Pellicer *et al.* (2013) inferred *n* = 17.
Ochnaceae	**14=0.676**, 13=0.222, 7=0.045	7	Raven (1975) hypothesized *n* = 7.
Onagraceae	**10=0.216**, 8=0.210, 7=0.202	11	Raven (1975) hypothesized *n* = 11.
Orobanchaceae	**9=0.918**, 8=0.074, 10=0.006	7, 12	Raven (1975) hypothesized *n* = 7, 12.
Pandanaceae	**30=0.441**, 16=0.126, 18=0.123	30	Raven (1975) hypothesized *n* = 30.
Papaveraceae	**7=0.552**, 8=0.313, 6=0.086	9, 10	Raven (1975) hypothesized *n* = 9, 10.
Polygalaceae	**12=0.477**, 11=0.284, 13=0.077	6	Raven (1975) hypothesized *n* = 6.
Portulacaceae	**12=0.920**, 13=0.038, 11=0.020	9	Oliveira Marinho *et al.* (2019) inferred *n* = 9.
Potamogetonaceae	**13=0.506**, 14=0.379, 12=0.070	10	Cusimano *et al.* (2012) inferred *n* = 10.
Rosaceae	**7=0.935**, 8=0.034, 6=0.028	9	Raven (1975) hypothesized *n* = 9.
Salicaceae	**10=0.640**, 11=0.133, 5=0.072	19	Raven (1975) hypothesized *n* = 19.
Sapindaceae	**10=0.584**, 8=0.183, 9=0.148	7	Raven (1975) hypothesized *n* = 7.
Scheuchzeriaceae	**7=0.394**, 6=0.219, 8=0.135	12	Cusimano *et al.* (2012) inferred *n* = 12.
Schisandraceae	**7=0.816**, 8=0.103, 6=0.049	14, 13	Raven (1975) hypothesized *n* = 14, 13.
Simaroubaceae	**8=0.488**, 9=0.369, 7=0.031	16	Raven (1975) hypothesized *n* = 16.
Tofieldiaceae	**15=0.360**, 16=0.265, 14=0.109	8	Cusimano *et al.* (2012) inferred *n* = 8.
Urticaceae	**7=0.923**, 8=0.023, 6=0.019	14	Raven (1975) hypothesized *n* = 14.
Vitaceae	**13=0.551**, 10=0.192, 14=0.077	12	Raven (1975) hypothesized *n* = 12.

Model	GBMB			GBOTB			PPA			Parameters
	LogLik	AIC	AIC$_w$	LogLik	AIC	AIC$_w$	LogLik	AIC	AIC$_w$	
Mc1	-19870	39750	0.0000000000	-19900	39800	0.0000000000	-4032	8070	0.0000000000	$\lambda; \delta; \rho$
Mc2	-18280	36560	0.0000000000	-18270	36550	0.0000000000	-3804	7613	0.0000013710	$\lambda; \delta; \rho=\mu$
Mc3	-18140	36290	0.0000000021	-18140	36280	0.0000000021	-3804	7616	0.0000003059	$\lambda; \delta; \rho; \mu$
Mc0	-47870	95740	0.0000000000	-49060	98120	0.0000000000	-5135	10270	0.0000000000	$\lambda; \delta$
Ml1	-19670	39350	0.0000000000	-19660	39320	0.0000000000	-3999	8008	0.0000000000	$\lambda; \delta; \rho; \lambda_1; \delta_1$
Ml2	-18260	36520	0.0000000000	-18250	36510	0.0000000000	**-3788**	**7586**	**0.9999999979**	$\lambda; \delta; \rho=\mu; \lambda_1; \delta_1$
Ml3	**-18120**	**36250**	**0.9999999979**	**-18110**	**36240**	**0.9999999979**	-3787	7587	0.6065306585	$\lambda; \delta; \rho; \mu; \lambda_1; \delta_1$
Ml0	-45260	90530	0.0000000000	-44910	89820	0.0000000000	-4918	9843	0.0000000000	$\lambda; \delta; \lambda_1; \delta_1$
Mb1	-18650	37310	0.0000000000	-18470	36950	0.0000000000	-4001	8011	0.0000000000	$\lambda; \delta; \beta; \nu$
Mb2	-21790	43580	0.0000000000	-21830	43670	0.0000000000	-4287	8581	0.0000000000	$\lambda; \delta; \rho; \beta; \nu$

572 **Table 4.** Goodness of fit of the 10 different models of chromosome number evolution applied to the three alternative phylogenies used.

Mc indicate models with constant rates, Ml models that include linear rate parameters and Mb models that include base number (not the chromosome number at the root of the phylogeny) parameters. Logarithmic likelihood (LogLik), AIC and relative weights scores (AICw). In bold, the lowest AIC value for each phylogeny indicates the best model. The last column indicates the parameter estimates included in each model (see Methods for details).

574