Title:

Lumbar corticospinal tract in rodents modulates sensory inputs but does not convey motor command

Authors:

Yunuen Moreno-Lopez[1], Charlotte Bichara[1], Gilles Delbecq, Philippe Isope, Matilde Cordero-Erausquin

[1]These authors contributed equally to this work

Affiliation (+ present address):

Institut des Neurosciences Cellulaires et Intégratives, CNRS UPR 3212, Université de Strasbourg

8 Allée du Général Rouvillois
67000 STRASBOURG
France

M.C.E. : https://orcid.org/0000-0002-1623-1773

P.I.: https://orcid.org/0000-0002-0630-5935

Y.M.L.'s present address: Burke Neurological Institute, White Plains, New York, NY 10605, United States

C.B 's present address: Neuro-Electronics Research Flanders (NERF), 3001 Leuven, Belgium

Corresponding Author :

Matilde Cordero-Erausquin
Tel: (+33) 3 88 45 66 60
Fax: (+33) 3 88 60 16 64
cordero@unistra.fr

Institut des Neurosciences Cellulaires et Intégratives - CNRS UPR 3212
8 allée du Général Rouvillois
67000 Strasbourg cedex, France

Keywords : corticospinal, sensory gating, primary afferent depolarization, motor control

Abstract

It is generally assumed that the main function of the corticospinal tract (CST) is to convey motor commands to bulbar or spinal motoneurons. Yet the CST has also been shown to modulate sensory signals at their entry point in the spinal cord, through presynaptic inhibition. By sequentially investigating different routes of corticofugal pathways through electrophysiological recordings and an intersectional viral strategy, we here demonstrate that motor and sensory modulation commands in mice belong to segregated paths within the CST. Sensory modulation is excuted exclusively by the CST via a population of lumbar interneurons located in the deep dorsal horn. In contrast, the cortex conveys the motor command via a relay in the upper spinal cord or supraspinal motor centers. At lumbar level, the main role of the CST is thus the modulation of sensory inputs, which is an essential component of the selective tuning of sensory feedback, to ensure well-coordinated and skilled movement.

Impact statement

While the corticospinal tract is often considered exclusively as a motor path, this study demonstrates that, in the mouse lumbar cord, its main role is the modulation of sensory inputs.

Text

 Introduction

Several brain and spinal structures have been involved in the generation and control of movement, in

a task-dependent manner. The primary motor cortex is classically associated to forelimb skilled

movements (Starkey et al., 2005; Guo et al., 2015; Miri et al., 2017; Galinanes et al., 2018) and to

complex locomotor tasks (obstacles crossing, leaning scale or beam, etc.) (Beloozerova and Sirota,

1993; DiGiovanna et al., 2016; Wang et al., 2017; Bieler et al., 2018) while its role in rythmic

locomotion appears, if anything, minor (Beloozerova and Sirota, 1993; DiGiovanna et al., 2016). A

coordinated movement relies not only on a proper motor command, but also on the online analysis

of an ongoing multi-sensory feedback indicating whether the movement has reached the planned

objective or requires correction (Akay et al., 2014; Fink et al., 2014; Azim and Seki, 2019). To optimize

the analysis of the overwhelming quantity of reafferent information, specific tuning of peripheral

feedback is possible through inhibition of sensory information at different stages of the sensory

pathway, a phenomenon known as "sensory gating" (Seki and Fetz, 2012; McComas, 2016).

At the spinal cord level, sensory gating can be mediated by presynaptic inhibition of sensory inputs

through primary afferent depolarization (PAD), a powerful mechanism controlled both by supraspinal

centers and by adjacent primary afferents (Eccles et al., 1962). While multiple mechanisms have

been demonstrated (Shreckengost et al., 2021), the most studied cause of PAD is presynaptic GABA-

A mediated depolarization. Primary afferents have indeed a high chloride content and respond to

GABA by a depolarization that has been shown to reduce the amplitude of incoming action potentials

and thus inhibit synaptic transmission (Prescott and De Koninck, 2003; Doyon et al., 2011). PAD has

been involved in the adjustment of reflex excitability (Rudomin and Schmidt, 1999), and its absence

results in a pathological context associated with spasticity (Caron et al., 2020). Proprioceptive

feedback is required for skilled locomotion (Akay et al., 2014), and presynaptic inhibition of sensory

feedback is essential for performing smooth forelimb skilled movements in mice (Akay et al., 2014;

Fink et al., 2014). Interestingly, PAD can also be evoked by stimulation of the sensorimotor cortex in

primates, cat and rats (Andersen et al., 1962; Carpenter et al., 1963; Andersen et al., 1964;

Abdelmoumene et al., 1970; Eguibar et al., 1994; Wall and Lidierth, 1997). This cortically-evoked PAD

may underlie cortical sensory gating at the spinal level and contribute to the cortical involvement in

coordinated movements, in addition to the classical drive of motoneurons. In this study, we aimed at

elucidating the neuronal pathways conveying the cortical motor command and sensory control to the

hindlimb in mice.This cortical control of the spinal cord, either to trigger motoneurons or to

modulate sensory information, can rely on multiple pathways, the corticospinal tract (CST) being the

most direct one. Its involvement in motor control is long acknowledged, although its precise

contribution might vary depending on the motor task (locomotion vs. skilled movement) or targeted

limb (forelimb vs. hindlimb) (Miri et al., 2017; Wang et al., 2017; Karadimas et al., 2019). The CST has

also been involved in cortically-evoked PAD in cats as a pyramidotomy abolishes it (Rudomin et al.,

1986). However, discriminating sensory vs motor contribution has been hampered by the lack of

proper tools to segregate functionally or anatomically overlapping paths.

Importantly, none of the CST functions can rely exclusively on direct CST contacts onto motoneurons

(MN) as these represent the exception rather than the rule. Indeed, monosynaptic CST-MN contacts

are present only in some higher primates, while CST-interneuronal connections are preponderant (in

adults) across species (rodents and primates) (Alstermark et al., 2011; Ebbesen and Brecht, 2017;

Ueno et al., 2018). In the recent years, specific genetic factors have enabled the identification of

several interneuronal populations, and of their inputs, including those from the cortex (Hantman and

Jessell, 2010; Bourane et al., 2015; Abraira et al., 2017; Ueno et al., 2018). But neurons with identical

genetic markers do not necessary share identical function: in these genetically distinct populations,

up to 65% of the neurons receive CST inputs (Ueno et al., 2018), while there is a significant fraction of

neurons that is not targeted by the CST. Although manipulating these populations has been

extremely informative to better understand their role, it remains unclear whether the changes

observed upon these manipulations can be specifically attributed to the neuronal subset that

receives inputs from the CST or to the subset that does not. Therefore, it is still difficult to assess the

specific role of the targets of the CST in movements.

In this study, we have investigated the corticofugal paths conveying motor command and modulation

of sensory inputs through PAD. We have in particular interrogated the contribution of the CST and its

direct lumbar targets through the development of an innovative intersectional viral strategy. We

show that cortically-evoked PAD in lumbar primary afferents is conveyed exclusively by the CST

through a population of lumbar interneurons. In contrast, our results suggest that motor command

to the hindlimbs is relayed either through supraspinal motor centers, or through the CST with a

propriospinal relay in the upper cord. The role of the lumbar CST thus appears to be mainly the

modulation of sensory inputs, which may in turn selectively increases sensory gain and refines motor

control during movement.

Results

PAD and muscular contractions originate in the same cortical area

The rodent sensorimotor cortex has a highly somatotopic organization (Ayling et al., 2009). Mainly

studied in a motor perspective, it has been repeatedly mapped for its ability to provoke motor

contraction (Li and Waters, 1991; Ayling et al., 2009), but rarely for its ability to induce PAD (Wall and

Lidierth, 1997). Cortically-evoked PAD may inhibit the transmission of sensory information from

primary afferents to the central nervous system and can be experimentally assessed by recording

dorsal root potentials (DRPs, Fig. 1A). We thus recorded DRPs *in vivo* in lumbar roots L4-L6 (conveying

hindlimb sensory inputs) of Thy1::ChR2 mice, expressing channelrhodopsin2 in most neuronal cells

including layer V cortical neurons (Arenkiel et al., 2007). DRP recordings were performed in isoflurane-

anesthetized mice, and trains of photostimulations (5x 8 msec-pulses, 1 msec apart) were applied to

activate ChR2 at different positions of the surface of the sensorimotor cortex according to a 500 μm-

spaced grid. The area inducing the DRPs was centered at AP=-0.75 mm, L=1.5 mm (Fig. 1C, n=7 mice).

Similar photostimulation of this area in mice expressing only the fluorescent reporter EGFP in

corticospinal (CS) neurons lead to no DRP signal (see below and Figure 3 – Supplementary figure 1). On

a different group of animals, electromyographic (EMG) recordings were performed in the tibialis

anterior (*TA*) under ketamine/xylazine anesthesia because EMGs cannot be evoked from cortical

stimulation under isoflurane anesthesia (see Methods). The sensorimotor cortex was similarly mapped

with trains of photostimulations (6x 1 msec-pulses, 2 msec apart) and the area inducing EMG was

centered at : AP=-0.75 mm, L=1.75mm (n=5 mice; Fig. 1C). Again, similar photostimulation of this area

in mice expressing only EGFP in CS neurons lead to no EMG signal (see below and Figure 3 –

Supplementary figure 1). We observed that the two zones are concentric, demonstrating that

cortically-evoked PAD and motor command of the hindlimb can originate from the same cortical area.

Spinal targets of the CST

We next investigated the spinal circuit underlying cortically-evoked DRPs. DRPs can be segmentally-

evoked by the activation of a neighboring root, dampening sensory inputs predominantly through

GABA-dependent primary-afferent depolarization (Rudomin and Schmidt, 1999). GABAergic terminals

presynaptic to proprioceptive fibers arise from GAD65 interneurons (Hughes et al., 2005; Betley et al.,

2009) and their activation produces DRPs (Fink et al., 2014). We investigated whether the CST directly

targets GAD65-expressing spinal lumbar neurons that could in turn lead to inhibition of primary

afferents. In TdTomato-flex mice crossed with GAD65::GFP mice, we injected a monosynaptic

anterograde transsynaptic virus (AAV1-CBA::WGA-Cre) encoding the Cre-recombinase fused to the

wheat germ agglutinin (WGA) (Libbrecht et al., 2017) in the area inducing hindlimb DRPs and EMG

(Figure 1 – Figure Supplement 1). The fusion with WGA provides transsynaptic properties to the Cre

recombinase, which is then transferred to the cortex's monosynaptic targets where it triggers the

expression of TdTomato (see Methods, Figure 1 - Figure Supplement 2). We evaluated the proportion

of GAD65+ neurons among the spinal targets of the CST revealed by Td-Tomato expression (Fig. 1D-E):

16.4±3.2% of spinal Td-Tomato positive neurons were GAD65::GFP+ (69 neurons out of 427, n=3 mice),

suggesting that the cortically-evoked DRP may be mediated directly by the activation of a GAD65+

spinal target of the CST. In the spinal dorsal horn, cholinergic neurons are a subpopulation of

GABAergic neurons (Todd, 1991; Mesnage et al., 2011), and cholinergic terminals are known to be

presynaptic to primary afferents (Ribeiro-da-Silva and Cuello, 1990; Pawlowski et al., 2013) and have

recently been shown to modulate primary afferent inputs (Hochman et al., 2010; Shreckengost et al.,

2021). By performing a similar experiment in ChAT::EGFP mice, we showed that only 2.2±1.8% of CST

targets were ChAT::EGFP positive (1 out of 45 neurons, n=4 mice), limiting the likelihood of a direct

major involvement of a cholinergic mechanism (Figure 1 - Figure Supplement 3).

Segregation of pathways for cortically-evoked DRPs and EMGs

Although lumbar DRPs and tibialis-EMGs (TA-EMGs) are elicited by the same area in the mouse

sensorimotor cortex, whether they share the same corticofugal pathway and spinal circuits remains

unknown. We first interrogated the contribution of indirect cortical-to-spinal pathways to these two

functions (Fig. 2). Since broad photostimulation of the sensorimotor cortex in Thy1::ChR2 activates an

heterogeneous population of layer V cortical neurons (Arenkiel et al., 2007) we selectively lesioned the

direct CST (Fig. 2B) using an acute electrolytic lesion at the level of the pyramidal decussation

(pyramidotomy) (Figure 2 - Figure Supplement 1A, C). In this study, we have consistently performed a

caudal pyramidotomy, sparing CST collaterals targeting brainstem motor nuclei that branch more

rostrally (Akintunde and Buxton, 1992; Alstermark and Pettersson, 2014) as well as rubrospinal or

reticulospinal tracts that can be activated by direct corticorubral or corticoreticular neurons (Fig. 2B).

Cortically evoked DRPs were completely abolished by this pyramidotomy (DRP amplitude before: 3.8

µV± 0.32 µV; after: 0.30 µV± 0.12 µV , n=3 mice, p=0.006), whereas cortically-evoked EMGs were hardly

affected (S/N Ratio 3.3± 0.61 before, 2.2 ± 0.1 after, n=4 mice, p=0.517, all Z-Scores above significance

after lesion, Figure 2 - Figure Supplement 2B) (Fig. 2C-D). This demonstrates that indirect cortical-to-

spinal pathways (involving supraspinal motor centers) do not encode cortically-evoked DRPs but have

a major role for cortically-evoked motor contraction of the hindlimb. These results show that, in the

mouse lumbar cord, the CST mediates cortically-evoked DRPs and hence modulates sensory inputs in

a presynaptic manner.

Although the contribution of the rubrospinal or reticulospinal tracts to motor command is well

acknowledged in both rodents and primates, the CST is also believed to encode motor command

(Wang et al., 2017). This could have been underrated by simultaneous stimulation of the indirect

cortical-to-spinal pathways in Thy1::ChR2 mice. We thus interrogated the specific contribution of the

CST to motor contraction by targeting exclusively these neurons through injection of a ChR2-encoding

retrograde virus (AAVrg-CAG-hChR2-H134R-tdTomato) in the lumbar spinal cord (Fig. 3A). As expected,

the infected CS neurons were located in the area delimited by the previous functional mapping of

hindlimb muscle contraction and DRP (Fig. 3B-C). Photostimulation of CS neurons (at the cortical level)

induced a robust EMG signal (S/N Ratio 2.4±0.52, n=5 mice, Fig. 3E, upper), demonstrating that CS

neurons evoke motor contraction. In agreement with the pyramidotomy results (Fig. 2C), the

photostimulation of CS neurons also induced DRPs (mean amplitude = 14.5±4.0μV, n=5), while

photostimulation in control mice expressing GFP in CS neurons induced neither an EMG (all Z-Scores

below significance) nor a DRP signal (mean amplitude = 0.3±0.0 μV, n=4, Figure 3 – Figure supplement

1). Because some CS neurons send collaterals to supraspinal regions involved in motor command

(Akintunde and Buxton, 1992), we next addressed whether the motor command traveled through the

direct cortical-to-spinal branch of CS neurons or via supraspinal relays. A pyramidotomy, similar to the

ones performed above (Figure 2 - Figure Supplement 1C), completely abolished the EMG signal evoked

by cortically activating CS neurons (S/N Ratio 0.9±0.12; n=3 mice, p=0.013; all Z scores below

significance after pyramidotomy, Figure 2 - Figure Supplement 2C) (Fig. 3D-E). This demonstrates that,

while supraspinal collaterals of the CST do not trigger motor output, direct spinal branches can drive

motor command. However, this contribution does not seem significant when other cortical neurons

are co-activated.

Spinal targets of the CST evoke DRPs but not muscular contraction

Because both DRPs and EMGs of a given limb are encoded by the same sensorimotor cortical area and

can be conveyed by the spinal branch of the CST, we next interrogated whether these functions are

segregated at the spinal level. CS neurons projecting to the lumbar spinal cord can give off collaterals

at the cervico-thoracic level (Kamiyama et al., 2015; Karadimas et al., 2019) and contact cervical

propriospinal neurons (Ueno et al., 2012; Ni et al., 2014) that in turn project to lumbar motoneurons

to relay motor command (Ni et al., 2014). Indeed, many experiments demonstrated that these neurons

are involved in segmental coordination (Miller et al., 1975; Nathan et al., 1996; Reed et al., 2006; Reed

et al., 2009). In order to test the contribution of the lumbar branch of CS neurons and rule out

antidromic stimulation of rostral collaterals, we combined the transynaptic tool used in Figure 1 with

an intersectional approach at the lumbar level. We first identified the lumbar segments containing CST

targets by using cortical injection of AAV1-CBA::WGA-Cre in TdTomato-flex mice (Fig. 4A): CST targets

were concentrated in spinal lumbar segments L2-L3, rostral to the large pools of motoneurons. They

were largely located in the dorsal horn (85,4% were dorsal to the central canal), in particular in the

contralateral deep dorsal horn (laminae IV to VII) (Fig. 4B, Fig. 4 – Figure supplement 1). Consistently

with the monosynaptic nature of the transynaptic labeling, their number was not correlated with the

delay between the infection and the histological analysis (Fig. 4 – Figure supplement 1). CST targets

were mostly located in the medial zone, lateral to the dorsal funiculus, where CST fibers penetrate the

grey matter. We thus combined the cortical injection of AAV1-CBA::WGA-Cre with the spinal (L4)

injection of an AAV (AAV9-Ef1a-DIO-ChETA-EYFP) encoding for a Cre-dependent form of ChETA (a ChR2

variant (Gunaydin et al., 2010)) (Fig. 4C). ChETA-EYFP expression was therefore restricted to the lumbar

targets of the CST (Fig. 4C, D). Our transectional approach induced ChETA-EYFP expression even in the

deepest targets of the CST, in the ventral horn (Fig. 4 – Figure supplement 2). There was no ChETA-

EYFP labeling in the dorsal funiculus, demonstrating that the intraspinal AAV injection did not lead to

retrograde infection of CS neurons (Fig. 4 – Figure supplement 2). Intraspinal injections did not affect

the general behavior of mice (e.g. no visible motor deficits nor pain-like behaviors) nor induced long-

lasting weight loss. As surface photostimulation of the spinal cord in Thy1::Cre animals produced an

EMG signal and directly activated neurons even in the deep dorsal horn (Fig. 4 – Figure supplement 3),

we performed an identical photostimulation in the animals expressing ChETA in the targets of the CST.

This stimulation induced a spinal LFP observable as deep as 1150 µm from the surface (Fig. 4 – Figure

supplement 3). It also induced DRPs in the ipsilateral lumbar root of 8 out of 9 mice (2.93 ± 0.48 µV,

n=8 mice, Fig. 4E-G), demonstrating that DRPs can be directly controlled by the CST projecting to the

lumbar cord through local interneurons. However, photostimulation of these same lumbar targets of

the CST (same animals) failed to elicit hindlimb movements or muscle contraction as attested by EMG

recordings (S/N Ratio 1.0±0.02; n=7 mice, Z-Scores below significance, Figure 2 - Figure Supplement

2D) (Fig. 4E-G). Together with the results presented in Fig. 3, we can conclude that CS neurons are able

to evoke motor contraction by activation of their spinal targets, but that these targets are not located

in the lumbar cord but rather at another spinal level. Cortically-evoked DRPs and EMGs thus follow a

similar corticofugal pathway that segregates at the spinal level.

Discussion

In this study, we have evaluated the contribution of the CST to the cortical control of movements,

either through its ability to convey the muscle contraction command or to modulate sensory inputs.

We found that the modulation of sensory inputs from the hindlimb (DRP in lumbar sensory roots) is

exclusively carried out by the CST via a population of lumbar interneurons located in the deep dorsal

horn. In contrast, the cortex induces muscle contraction of the hindlimb (*TA*) through a relay in the

upper spinal cord or supraspinal motor centers. The two mechanisms are therefore segregated, and

the main role of the lumbar CST appears to be the modulation of sensory inputs.

Reliability of circuit investigations

We have identified the cortical area involved in these two mechanisms through functional means:

cortically-evoked DRPs and EMGs. For ethical and technical reasons, most of these recordings were

performed on different animals and conditions. First, EMG recordings after cortical stimulation

required a light anesthesia level (Tennant et al., 2011), only possible using Ketamine/Xylazine. Such a

light anesthesia is not ethically compatible with the large and invasive surgery required for recording

cortically-evoked DRPs. DRPs were thus recorded under isoflurane anesthesia under which no

cortically-evoked EMG could be recorded. We therefore acknowledged that the cortical zone

responsible for DRPs, smaller than the one responsible for EMGs, might have been underestimated

by the deeper anesthesia. We thus did not attempt to quantitatively compare EMG vs DRP map

extents, but concluded that they partially overlap allowing us to stimulate at the center of two maps.

Although the different experimental approaches might also have differentially affected the

recruitment threshold for EMG vs DRP, we do not compare these thresholds in between

stimulation/recording configurations. Rather, we analyze the amplitude of a given signal (EMG *or*

DRP) with a given experimental approach in different animal models (THY1, ChR2-retro,

pyramidotomy, transynaptically labeled CS targets).

Circuit investigation also rely on the efficiency of pyramidotomies which affect the spinal branch of

CS tract while sparing CST collaterals targeting brainstem motor nuclei, that are branching more

rostrally (Akintunde and Buxton, 1992). We performed similar pyramidotomies in two different

group of mice, either Thy1::ChR2 or ChR2-retro group (Fig. 1, Fig. 2, Fig. 1 - Figure Supplement 3). In

the latter group, in which the CST expressed ChR2, we observed a complete loss of cortically-evoked

EMGs after pyramidotomy. These results confirm that the pyramidotomy was efficient to block CST-

induced EMG. Therefore, the remaining EMG signal after pyramidotomy in Thy1::ChR2 mice is likely

the result of a non-CST pathway.

Finally, an intriguing result we report is the absence of EMG signal after photostimulating the targets

of the CST at the lumbar level. The reliability of this result relies on multiple technical controls. First,

the lack of muscle contraction cannot be attributed to damages of spinal microcircuits due to

intraspinal AAV injection. None of the mice presented obvious alteration in the behavior nor a long-

lasting weight loss. In addition, similar intraspinal injections were performed for the ChR2-retrograde

group, and in these animals, EMG was systematically observed after cortical stimulation,

demonstrating the integrity of required spinal circuits. Second, the intraspinal AAV injection

efficiently infected neurons in the dorsal horn, and even in the dorsal part of the ventral horn, i.e. in

an area comprising 97% of the lumbar targets of the CST. Third, the optic fiber positioned at the

surface of the cord evoked an intraspinal LFP as deep as 1150 μm in the double-infected mice, but

also evoked EGMs in THY1::ChR2 mice, as previously reported (Caggiano et al., 2016). We also

demonstrate direct photoactivation of neurons located up to 700 μm from the surface with the short

1 ms stimulus used for EMG recordings. As 73% of the targets of the CST are located within 700 μm

from the surface, this demonstrate that the large majority of this population can be directly activated

by this procedure. Finally, using a similar surface photostimulation with powers equal or lower than

in the present study, Caggiano and colleagues were able to induce motor contraction in ChAT::ChR2

animals while considering that they were directly activating motoneurons located in the deep ventral

horn (Caggiano et al., 2016). We thus consider very likely that the surface illumination we used could

reach most of the lumbar CS-postsynaptic neurons, and that the absence of EMG in this situation is

biologically relevent.

Anatomical definition of the hindlimb sensorimotor cortex

In our study, the cortical area inducing hindlimb contraction is similar to that previously reported in

mice (Li and Waters, 1991; Ayling et al., 2009). This cortical area (centered around 1mm caudal and

1,5mm lateral to bregma) spans over the motor cortex and also partly over the primary

somatosensory cortex according to some atlases and studies (Kirkcaldie et al., 2012; Liu et al., 2018;

Karadimas et al., 2019). We thus use the conservative term "hindlimb sensorimotor cortex".

Interestingly, we found that stimulation of this area also induced PAD at the lumbar level. This

overlap might be a specificity of the "hindlimb area" of the cortex (HLA), and may not generalize to

the "forelimb areas" (RFA and CFA). Indeed, the CS neurons from the forelimb areas are spatially

segregated (forming clusters projecting either to the dorsal or to the intermediate-ventral cervical

cord), while those from the HLA are largely intermingled (Olivares-Moreno et al., 2017). In addition,

our results demonstrate that the same cortical area is labelled after retrograde tracing from the

lumbar cord (Fig. 3C) excluding the existence of a different pool of cortico-lumbar neurons.

Pathways conveying the cortical motor command

Our study concludes that a large fraction of the cortical motor command to the hindlimb is conveyed

through non-CST pathways. These could involve the reticular formation or the red nucleus which

receive cortical innervation, project directly or indirectly onto spinal motoneurons, and control

different parameters of movements (Lavoie and Drew, 2002; Zelenin et al., 2010; Esposito et al.,

2014). Our recordings also suggest that the cortex conveys a motor command to the hindlimb

through the CST projecting to a non-lumbar spinal segment as direct stimulation of the CS lumbar

targets failed to produce muscle contraction. We have discussed above possible technical caveats

and concluded that our approach most likely induced activation of the vast majority of the lumbar CS

targets labelled by the transynaptic tracing. We cannot rule out that some of these lumbar targets

contribute to convey the cortical motor command to the hindlimb but were not labeled in large

enough quantities through our transynaptic approach. The efficiency of WGA-Cre transynaptic

tracing might be target specific (Libbrecht et al., 2017), and the number of labelled neurons in the

spinal cord suggest that not all targets were labeled. However, this would still implicate that these

neurons are distinct from the ones involved in the generation of a DRP, and that these two functions

use segregated lumbar pathways. While this hypothesis cannot be excluded, the recent report by

Karadimas and collaborators (Karadimas et al., 2019) strongly supports the hypothesis that the motor

cortical command conveyed by the CST relays in rostral spinal segments. Indeed, they demonstrate

that the cortical motor command to the hindlimb can be mediated by CS neurons projecting to the

cervical cord, and then relayed by cervical-to-lumbar propriospinal neurons (Karadimas et al., 2019).

Although we selectively study CS projecting at least to the lumbar cord (through lumbar retrograde

infection), many CS neurons have collaterals at different spinal segments (Kamiyama et al., 2015).

Therefore, their photostimulation at the sensorimotor cortex level would activate cervical pathways

in addition to lumbar ones.

Pathways conveying the cortical sensory control

Beyond inducing hindlimb muscle contraction, we have characterized the ability of the cortex to

induce DRPs, which is a measure of PAD(Wall and Lidierth, 1997). We demonstrate that it is

exclusively relayed through the CST and provide some information on spinal interneurons potentially

mediating this effect. The most studied mechanism of segmental PAD involves (Prescott and De

Koninck, 2003; Doyon et al., 2011)a presynaptic GABAergic last order neuron (mostly expressing

GAD65 (Betley et al., 2009)), located in laminae V-VI, under the control of a first order excitatory

interneuron which receives peripheral and descending (including cortical) inputs (Eccles et al., 1962;

Rudomin and Schmidt, 1999). (Betley et al., 2009)The targets of the CST identified by our

transynaptic labeling are indeed predominantly located in laminae IV-VI, and include a small

population of GAD65-expressing neurons. They represent a potential substrate for Cx-DRP that

would thus not require a spinal first order excitatory interneuron.

In addition to the sensorimotor cortex, stimulation of other supraspinal nuclei are known to induce

PAD. This is the case of the red nucleus and the reticular formation (Jimenez et al., 1987; Sirois et al.,

2013), both of which also receive inputs from the sensorimotor cortex. The corticorubral and

corticoreticular tracts were preserved by the pyramydotomy, yet the cortically-induced PAD was

completely lost. This strongly suggests that the ability of the red nucleus and reticular formation to

induce PAD does not rely on a sensorimotor cortical command. The control of incoming sensory

information through PAD plays an important role in the generation of coordinated movements and

processing of sensory information, including nociceptive information (McComas, 2016). GAD65-

expressing neurons are essential for smoothed forelimb movements (Fink et al., 2014), but whether

they require peripheral or cortical information to do so is yet unknown. Similarly, recordings in

primates demonstrated that sensory information is inhibited in a modality- and phase-dependent

manner during forelimb voluntary movements (Seki et al., 2003, 2009; Seki and Fetz, 2012)

suggesting that this inhibition is somehow related to the motor command, but the exact underlying

mechanism remains to be elucidated. Importantly, the change in afferent fiber excitability (Wall,

1958) accompanying the sensory presynaptic inhibition in primates was even smaller (<2μV) (Seki et

al., 2003) than the DRPs we report as a consequence of cortical stimulation, suggesting that Cx-DRPs

might be sufficient to significantly impact the transmission of sensory information. Here, we

demonstrate that CS neurons from the same sensorimotor cortex area (but not necessarily a single

population) are able to relay both motor command and sensory modulation to hindlimb, but through

segregated pathways. Future studies on awake animals and/or focusing on specific sensory

modalities are needed to further elucidate the timing and specificity of the sensory control we have

described.

Significance of the results

In contrast with the present results on hindlimbs, specific stimulation of the CST at the cervical level

induces forelimb contraction (Gu et al., 2017), suggesting a direct relay of the cortical motor

command at this level (although antidromic activation of a collateral pathway cannot be excluded).

Relative to the forelimb, mice do not display an extensive repertoire of skilled hindlimb movements

and a difference in the level of cortical control is expected. The functional segregation we report here

suggests that, while the sensory control is selectively targeted at the lumbar level, the hindlimb

motor command is relayed in the upper cord. The fact that the upper cord (possibly the cervical cord

according to (Karadimas et al., 2019)), receiving the motor command from the forelimb sensorimotor

cortex, also receives the information of hindlimb motor command offers a substrate for the

coordinated control of fore- and hindlimb, which is essential for most, if not all, skilled movements.

The organization of the CST greatly differs between species. From an anatomical point of view, this

tract travels mainly in the dorsal funiculus in rodents, while it is located in the dorsolateral funiculus

in primates, and its projection area within the spinal cord accordingly differs (Lemon, 2008). The

evolution of the CST has been linked to the development of the most dexterous movements that can

be performed by primates, with the appearance of direct cortico-motoneuronal (CMN) contacts that

are absent in rodents. From an evolutionary perspective, it is interesting to note that the lumbar

branch of the CST, which will later acquire a direct motor command role (including with direct CMN

contacts in primates (Porter, 1985)) is specialized in mice in the modulation of sensory inputs.

Importantly, primate CS neurons don't only form CMN contacts but also abundantly project onto

spinal interneurons, and the sensory control function of the CST we report has also been proposed in

primates (Lemon, 2019). While extrapolation of our results to primates difficult in view of the large

species differences, mice are the main and foremost animal model, including for pathologies

affecting the CST such as spinal cord injury or amyotrophic lateral sclerosis. Better understanding the

differential roles of the mouse CST in the cervical and lumbar cord is thus an essential endeavor to

properly interpret data from these models.

Conclusion

Altogether, our results support a segregation of pathways involved in cortically-evoked sensory

modulation vs. motor control. The direct CST is able to induce motor contraction, independently of

its supraspinal collaterals, through spinal targets possibly located at the cervico-thoracic level.

However, we show that motor command is largely mediated by non-CST pathways, most likely

cortico-rubral, or -reticular ones. On the other hand, the major and essential role of the lumbar

rodent CST appears to be hindlimb sensory modulation at the primary afferent level, through a

population of lumbar target neurons whose activation is sufficient to produce DRPs. (Lemon,

2008)This forces re-interpretation of previous studies aiming at promoting CST function in a

therapeutic perspective, in mouse injury or neurodegenerative models. While most of these studies

have exclusively considered motor command aspects, our results provide a new perspective to

analyze these efforts by considering the primary role of rodent lumbar CST, i.e. sensory modulation.

MATERIALS AND METHODS

Animal models. This study was carried out in strict accordance with the national and international laws for laboratory animal welfare and experimentation and was approved in advance by the Ethics Committee of Strasbourg (CREMEAS; CEEA35; agreement number/reference protocol: APAFIS# 12982 - 2017122217349941 v3). The following mice strains were used (adult males and females): Thy1::ChR2 (Cg-Tg(Thy1-COP4/EYFP)18Gfng/J, Jackson Laboratory), TdTomato-Flex (*Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze}*/J, Jackson Laboratory), ChAT::EGFP(von Engelhardt et al., 2007), GAD65::GFP(Lopez-Bendito et al., 2004). All lines except GAD65::GFP were back-crossed (at least 10 generations) on a CD1 background. GAD65::GFP were on a C57Bl6/J background. The mice were housed in the Chronobiotron facility (UMS3415, CNRS, University of Strasbourg) in accordance with the European convention 2010/63/EU on the protection of animals used for scientific purposes.

Electrophysiological recordings.

EMG recording: Ketamine/Xylazine anesthesia was chosen to minimally affect muscular tone. After the initial anesthesia (150 mg/kg of ketamine, Imalgene 1000 and 10 mg/kg of Xylazine, Rompun 20%), complementary doses of Ketamine were given when the animal was showing signs of awakening (additional doses bridged by short exposure to Isoflurane 1–1.5%)(Tennant et al., 2011). Concentric EMG needle electrodes (Myoline Xp) were inserted in the *tibialis anterior (TA)* muscle of the animals and a ground electrode was inserted in the tail. Responses were collected with a recording amplifier (IR183A, Cygnus technology) in series with a second amplifier (Brownlee Precision, model440). The traces were filtered with a bandwidth of 0.1 Hz–10 kHz, recorded with Spike 2 software (version 8.00, CED, Cambridge, UK), and analyzed off line with Clampfit (pCLAMP, version 10.7) and home-made python routines (WinPython 2.7.10, Python Software Fundation). EMG responses were recorded after cortical or spinal photostimulation (see below), and the response to 3 consecutive stimulations was averaged for analysis. The area under the curve of the rectified average trace was calculated in a

window of 30ms, either 40 ms before (baseline noise, N) or 10 ms after (signal, S) the onset of the stimulation. The EMG response was presented as the S/N ratio. In CST lesion experiments, all S/N ratios were expressed as a % of the control S/N (average of 3 responses before the lesion). The noise level (S/N of 1) was similarly expressed as a % of the above defined control S/N.

DRP recording: Under isofluorane (1.5-2%) anesthesia, the mice spinal cord was fixed with the spinal cord unit of a stereotaxic frame (Narishige instruments). A laminectomy was performed to expose the surface of lumbar L3-L6 spinal cord segments. A dorsal root (L4-L6) was dissected, cut before the DRG, and a rootlet was suctioned with a glass pipet (sometime two rootlets were suctioned together). An agar pool was created on the exposed spinal cord and filled with NaCl 0.9%. The amplifiers/filters/software used were the same as for the EMG recordings. DRPs in response to 60 successive cortical or spinal photostimulations (see below) were averaged for analysis. The amplitude of the response from the onset to the peak was measured using Clampfit 10.0. The amplitude of the noise was similarly measured on a window preceding photostimulation. For spinal photostimulation, DRPs and EMGs recordings of 3 series of photostimulation were averaged.

For electrophysiological recordings, we excluded animals that had a constant decreasing signals (run-down) in otherwise stable conditions. For DRP recordings that required a delicate surgery, animals were excluded from the study if the surgical preparation / recovery was compromised or if scaring was abnormal after intraspinal AAV injection.

Single Unit extracellular recording: The animal was prepared and anesthetized as for DRP recordings (except the dorsal root dissection). Single-unit extracellular recordings were made with a glass electrode (Harvard Apparatus, Holliston, MA, USA) filled with 0.5M NaCl and 0.06M HEPES, pH adjusted at 7,3 (resistance ~19 MΩ). A motorized micromanipulator (Narishige, Tokyo, Japan) was used to gradually insert the electrode with 4 µm steps. The amplifiers and software used were the same as for the DRP and EMG recordings, but the signal was at 0.1-3 kHz. The signal was analyzed with Python routines for spike detection (amplitude threshold of 0.30 mV) and delay measure (from onset of photostimulation to spike peak).

Cortical photostimulation. Photostimulation of the cortex was performed on anesthetized animals while recording EMG or DRP. For the cortex, a craniotomy of approximately 9mm^2 was performed under stereo microscope on the contralateral side from the recording site. Muscle contraction, measured by EMG, was evoked by photostimulation with a 250 µm-probe (473 nm laser source PSU-III-FDA, 56mW/cm²) using the following protocol: trains of 6 pulses, 1 ms duration, 2 ms interval; 15 s minimum between each train (Li and Waters, 1991; Carmel and Martin, 2014). Cortically-evoked DRPS typically require trains of electrical stimulations (Andersen et al., 1962, 1964; Eguibar et al., 1994; Wall and Lidierth, 1997), a protocol that we adapted for photostimulation using a 105 µm-probe (460 nm LED source, Prizmatix, 3.36mW/cm²): trains of 5 pulses of 8 ms, 1 ms interval and 2 s between each train. In both cases, the probe was moved at the surface of the cortex along a 500 µm-spaced grid. Isopotential contour plots were created, using for EMGs the S/N and for DRPs the peak amplitude (both normalized to the largest values in each animal) at each photostimulation point in the cortex, using a linear interpolation on a set of X, Y, Z triples of a matrix. For EMG, only S/N ratio above significance (Z-Score>3, Figure 2 - Figure Supplement 2A) were considered for the analysis. Then the isopotentials contours were superimposed on the metric planes of the top view of the cortex (Kirkcaldie et al., 2012).

Spinal cord photostimulation. Photostimulation of the spinal cord was performed on anesthetized mice after laminectomy (described above for DRPs recordings), using a 1,1 mm probe (LED source, 42 mW/cm²) first placed on the surface of the lumbar spinal cord (on top of the injection site). The stimulation trains were those used for cortical stimulation. If no signal was observed when stimulating on top of the injection site, the optical fiber was then swept rostrally and caudally on top the whole lumbar segment, in order to try and obtain a signal. EMG and DRP recordings were sequentially performed and required to alternate between different types of anesthesia: isoflurane (craniotomy and laminectomy surgery), ketamine/xylazine (EMG recording), isoflurane 1.5–2 % (DRPs recordings).

Only the mice filling the two following criteria were kept for analysis: (i) TdTomato$^+$/EYFP$^+$ neurons

were found in the stimulated area; (ii) the response was not precisely associated to the onset or offset

of optical pulses (which would be expected for a photoelectric artefact).

For spinal single-unit extracellular recordings, the photostimulation protocol consisted in either a

single 1ms pulse, or in the same train used to test EMG responses (6 pulses, 1 ms duration, 2 ms

interval).

Pyramidotomy. Electrolytic lesion of the CST was performed in a subset of THY1::ChR2 and ChR2

retrogradely labeled mice, with 200 ms pulses, 30 mA using a constant current stimulator (Digitimer

Ltd thorug) through a silver bipolar electrode inserted in the pyramidal decussation (3.5-4 mm caudal

to Bregma, 6 mm deep). The brain was removed for histological analysis at the end of the experiment.

Cortical and spinal injections.

Brain injections (1.5–2 mm lateral, 0.5–1 mm caudal to Bregma, 0.5 mm deep) were performed as

previously described(Cetin et al., 2006) under isoflurane anesthesia (2–3%). Briefly, 90 to 270 nL of

virus was injected by manual pressure using a 5 mL syringe coupled to a calibrated glass capillary, under

visual control. Spinal injections were performed using a similar manual pressure protocol. The pipette

was inserted in the exposed space between two vertebrae (T13-L1, corresponding to spinal L4), as

previously described (Tappe et al., 2006). 0.45 µL of virus was injected 300 µm lateral to the midline

and 300-400 µm deep. In both cases, Manitol 12.5% was injected i.p. (0.2 – 0.5 ml) after the surgery

to enhance vector spread and improve transduction (Tjolsen et al., 1992). After spinal injection,

transfected neurons were found on multiple segments centered in L4 (depending on the animals, up

to the whole lumbar enlargement).

The animals were kept 2 weeks for the retroAAV injections and a minimum 5 weeks for dual injections

before in vivo DRPs and/or EMGs recordings or histological analysis. Animals for which post hoc

histological analysis demonstrated inappropriate injections coordinates were excluded from the

analysis.

Viruses. AAV2/1-CBA-WGA-CRE-WPRE was purchased at the molecular tools platform at the Centre de

recherche CERVO (Québec, Canada), and was used at a titer of 8×10^{12} vg/ml. AAV-Ef1a-DIO ChETA-

EYFP was a gift from Karl Deisseroth (Gunaydin et al., 2010) (Addgene viral prep # 26968-AAV9;

http://n2t.net/addgene:26968 ; RRID:Addgene_26968) and was used at a titer of 1×10^{13} vg/mL. AAV-

CAG-hChR2-H134R-tdTomato was a gift from Karel Svoboda (Mao et al., 2011) (Addgene viral prep #

28017-AAVrg; http://n2t.net/addgene:28017 ; RRID:Addgene_28017), and was used at a titer of

7×10^{12} vg/mL. pAAV-hSyn-EGFP was a gift from Bryan Roth (Addgene viral prep # 50465-AAVrg;

http://n2t.net/addgene:50465 ; RRID:Addgene_50465) and was used at a titer of 1×10^{13} vg/mL.

Histology

Mice were transcardially perfused with PB followed by 4% paraformaldehyde (PFA) in PB 0.1M, or, if

histological analysis followed electrophysiological recordings, the brain and spinal cord were post-fixed

overnight in PFA 4% in PB 0.1M. Serial 50 µm brain (coronal or sagittal) and spinal (transverse or

sagittal) sections were performed on a vibratome (Leica VT1000 S) and mounted using a DAPI staining

mounting medium (Vectashield, Vector laboratories).

Image analysis. The extent of the cortical injection site was estimated by measuring the spread of

reporter protein fluorescence (TdTomato) in the layer V of the cortex. The intensity of fluorescence

was analyzed as previously described(Lorenzo et al., 2008; Mesnage et al., 2011) on evenly spaced (250

µm apart) transverse brain sections imaged with a microscope (Axio Imager 2, Zeiss). Briefly, the "Plot

Profile" function of ImageJ software (W. Rasband, National Institutes of Health) was used to measure

the intensity of fluorescence along the horizontal axis of a 6000x250 µm rectangle containing the layer

V of the cortex (Figure 1 - Figure Supplement 1). These values were normalized to the highest intensity

for each animal, and used to plot a density map of each injection site. The contour of the injection area

was defined as 30% of the maximal intensity, corresponding to the approximate intensity of individual

neurons.

Spatial distribution of corticospinal post-synaptic neurons: Spinal_lumbar sections (50µm) were

mounted serially and imaged using a Zeiss epifluorescence microscope or a Leica SP5 II confocal

microscope. Images were aligned manually with Photoshop using the central canal and the dorsal

funiculus as landmarks. Each labeled neuron was assigned coordinates corresponding to its distance

to the center of the central canal (X and Y) and the index of the slice containing it (Z). This was used to

calculate the number of labeled neurons every 100 µm x 100 µm bins (XxY) (or 200 µm x200 µm X x Z)

in order to build density plots. The neuronal distribution was plotted using the "spline 16" interpolation

method of the matplotlib library in a homemade python script.

Quantification and statistical analysis. No prior sample size calculation was performed. The mean and

standard deviation of the Noise were measured from the 3 Noise values obtained from each EMG

recording, and used to calculate the Z-scores. The Z-score of each EMG response to the stimulation

was calculated using the following equation:

$$\text{Z-score} = \frac{\text{Signal- mean Noise}}{\text{standard deviation of Noise}}$$

A Z-score of 1.96 or 3 corresponding to a significance level of 0.05 or 0.001 was chosen to discriminate

significant responses from non-significant ones. The conservative choice of threshold was chosen in

agreement with visual inspection of the traces, to ensure that any EMG responses was not simply due

to noise. For each tests, the normality and variance equality were verified by SigmaPlot (Version 13,

Copyright © 2014 Systat Softwara, Inc) using Shapiro-Wilk and Brown-Forsythe's methods, before we

applied the parametric tests. All the data were analyzed using a one-way ANOVA and a post-hoc test

was performed only if an effect showed statistical significance. The P-values for multiple comparison

were measured using Holm-Sidak's method. Error bars in all Figures represent mean ± s.e.m., *P <

0.05, NS, no statistical significance (P ≥ 0. 05). The power of the tests was systematically attested to be

above the desired power of 0.80.

Acknowledgments

We gratefully acknowledge the support from the CNRS and Strasbourg University; University of

Strasbourg Institute for Advanced Study (USIAS) and ANR- 13-JSV4-0003-01 grants to MCE; ANR - 15-

CE37-0001-01 CeMod, ANR - 19-CE37-0007-03 MultiMod, and ANR - 19-CE16-0019-01 NetOnTime to

PI. YML was supported by a postdoctoral fellowships from the Fondation pour la Recherche Médicale

(SPF20160936264) and CB was funded by a fellowship from the Ministère de la Recherche and by the

ARSLA (Association pour la Recherche sur la SLA). CB was a student of the EURIDOL graduate School

of Pain (University of Strasbourg and ANR-17-EURE-0022). We also thank the following for

providing/generating specific mouse lines: Gábor Szabó for GAD65::GFP, Jakob von Engelhardt for

ChAT::EGFP. We thank Dr. Sophie Reibel-Foisset and the staff of the animal facility (Chronobiotron,

UMS 3415 CNRS and Strasbourg University) for technical assistance. We also thank Yves De Koninck,

Frédéric Doussau and Didier Desaintjan for critical reading of the manuscript.

References

Abdelmoumene M, Besson JM, Aleonard P (1970) Cortical areas exerting presynaptic inhibitory

action on the spinal cord in cat and monkey. Brain Res 20:327-329.

Abraira VE et al. (2017) The Cellular and Synaptic Architecture of the Mechanosensory Dorsal Horn.

Cell 168:295-310 e219.

Akay T, Tourtellotte WG, Arber S, Jessell TM (2014) Degradation of mouse locomotor pattern in the

absence of proprioceptive sensory feedback. Proc Natl Acad Sci U S A 111:16877-16882.

Akintunde A, Buxton DF (1992) Origins and collateralization of corticospinal, corticopontine,

corticorubral and corticostriatal tracts: a multiple retrograde fluorescent tracing study. Brain Res

586:208-218.

Alstermark B, Pettersson LG (2014) Skilled reaching and grasping in the rat: lacking effect of

corticospinal lesion. Front Neurol 5:103.

Alstermark B, Pettersson LG, Nishimura Y, Yoshino-Saito K, Tsuboi F, Takahashi M, Isa T (2011) Motor

command for precision grip in the macaque monkey can be mediated by spinal interneurons. J

Neurophysiol 106:122-126.

Andersen P, Eccles JC, Sears TA (1962) Presynaptic inhibitory action of cerebral cortex on the spinal

cord. Nature 194:740-741.

Andersen P, Eccles JC, Sears TA (1964) Cortically Evoked Depolarization of Primary Afferent Fibers in

the Spinal Cord. J Neurophysiol 27:63-77.

Arenkiel BR, Peca J, Davison IG, Feliciano C, Deisseroth K, Augustine GJ, Ehlers MD, Feng G (2007) In

vivo light-induced activation of neural circuitry in transgenic mice expressing channelrhodopsin-2.

Neuron 54:205-218.

Ayling OG, Harrison TC, Boyd JD, Goroshkov A, Murphy TH (2009) Automated light-based mapping of

motor cortex by photoactivation of channelrhodopsin-2 transgenic mice. Nat Methods 6:219-224.

Azim E, Seki K (2019) Gain control in the sensorimotor system. Curr Opin Physiol 8:177-187.

Beloozerova IN, Sirota MG (1993) The role of the motor cortex in the control of accuracy of

locomotor movements in the cat. J Physiol 461:1-25.

Betley JN, Wright CV, Kawaguchi Y, Erdelyi F, Szabo G, Jessell TM, Kaltschmidt JA (2009) Stringent

specificity in the construction of a GABAergic presynaptic inhibitory circuit. Cell 139:161-174.

Bieler L, Grassner L, Zaunmair P, Kreutzer C, Lampe L, Trinka E, Marschallinger J, Aigner L, Couillard-

Despres S (2018) Motor deficits following dorsal corticospinal tract transection in rats: voluntary

versus skilled locomotion readouts. Heliyon 4:e00540.

Bourane S, Grossmann KS, Britz O, Dalet A, Del Barrio MG, Stam FJ, Garcia-Campmany L, Koch S,

Goulding M (2015) Identification of a spinal circuit for light touch and fine motor control. Cell

160:503-515.

Caggiano V, Cheung VC, Bizzi E (2016) An Optogenetic Demonstration of Motor Modularity in the

Mammalian Spinal Cord. Sci Rep 6:35185.

Carmel JB, Martin JH (2014) Motor cortex electrical stimulation augments sprouting of the

corticospinal tract and promotes recovery of motor function. Front Integr Neurosci 8:51.

Caron G, Bilchak JN, Cote MP (2020) Direct evidence for decreased presynaptic inhibition evoked by

PBSt group I muscle afferents after chronic SCI and recovery with step-training in rats. J Physiol

598:4621-4642.

Carpenter D, Lundberg A, Norrsell U (1963) Primary Afferent Depolarization Evoked from the

Sensorimotor Cortex. Acta Physiol Scand 59:126-142.

Cetin A, Komai S, Eliava M, Seeburg PH, Osten P (2006) Stereotaxic gene delivery in the rodent brain.

Nat Protoc 1:3166-3173.

DiGiovanna J, Dominici N, Friedli L, Rigosa J, Duis S, Kreider J, Beauparlant J, van den Brand R,

Schieppati M, Micera S, Courtine G (2016) Engagement of the Rat Hindlimb Motor Cortex across

Natural Locomotor Behaviors. J Neurosci 36:10440-10455.

Doyon N, Prescott SA, Castonguay A, Godin AG, Kroger H, De Koninck Y (2011) Efficacy of synaptic

inhibition depends on multiple, dynamically interacting mechanisms implicated in chloride

homeostasis. PLoS Comput Biol 7:e1002149.

Ebbesen CL, Brecht M (2017) Motor cortex - to act or not to act? Nat Rev Neurosci 18:694-705.

Eccles JC, Kostyuk PG, Schmidt RF (1962) Central pathways responsible for depolarization of primary

afferent fibres. J Physiol 161:237-257.

Eguibar JR, Quevedo J, Jimenez I, Rudomin P (1994) Selective cortical control of information flow

through different intraspinal collaterals of the same muscle afferent fiber. Brain Res 643:328-333.

Esposito MS, Capelli P, Arber S (2014) Brainstem nucleus MdV mediates skilled forelimb motor tasks.

Nature 508:351-356.

Fink AJ, Croce KR, Huang ZJ, Abbott LF, Jessell TM, Azim E (2014) Presynaptic inhibition of spinal

sensory feedback ensures smooth movement. Nature 509:43-48.

Galinanes GL, Bonardi C, Huber D (2018) Directional Reaching for Water as a Cortex-Dependent

Behavioral Framework for Mice. Cell Rep 22:2767-2783.

Gu Z, Kalambogias J, Yoshioka S, Han W, Li Z, Kawasawa YI, Pochareddy S, Liu F, Xu X, Wijeratne HRS,

Ueno M, Blatz E, Salomone J, Kumanogoh A, Rasin MR, Gebelein B, Weirauch MT, Sestan N, Martin

JH, Yoshida Y (2017) Control of species-dependent cortico-motoneuronal connections underlying

manual dexterity. Science 357:400-404.

Gunaydin LA, Yizhar O, Berndt A, Sohal VS, Deisseroth K, Hegemann P (2010) Ultrafast optogenetic

control. Nat Neurosci 13:387-392.

Guo JZ, Graves AR, Guo WW, Zheng J, Lee A, Rodriguez-Gonzalez J, Li N, Macklin JJ, Phillips JW,

Mensh BD, Branson K, Hantman AW (2015) Cortex commands the performance of skilled movement.

Elife 4:e10774.

Hantman AW, Jessell TM (2010) Clarke's column neurons as the focus of a corticospinal corollary

circuit. Nat Neurosci 13:1233-1239.

Hochman S, Shreckengost J, Kimura H, Quevedo J (2010) Presynaptic inhibition of primary afferents

by depolarization: observations supporting nontraditional mechanisms. Ann N Y Acad Sci 1198:140-

152.

Hughes DI, Mackie M, Nagy GG, Riddell JS, Maxwell DJ, Szabo G, Erdelyi F, Veress G, Szucs P, Antal M,

Todd AJ (2005) P boutons in lamina IX of the rodent spinal cord express high levels of glutamic acid

decarboxylase-65 and originate from cells in deep medial dorsal horn. Proc Natl Acad Sci U S A

102:9038-9043.

Jimenez I, Rudomin P, Solodkin M (1987) Mechanisms involved in the depolarization of cutaneous

afferents produced by segmental and descending inputs in the cat spinal cord. Exp Brain Res 69:195-

207.

Kamiyama T, Kameda H, Murabe N, Fukuda S, Yoshioka N, Mizukami H, Ozawa K, Sakurai M (2015)

Corticospinal tract development and spinal cord innervation differ between cervical and lumbar

targets. J Neurosci 35:1181-1191.

Karadimas SK, Satkunendrarajah K, Laliberte AM, Ringuette D, Weisspapir I, Li L, Gosgnach S, Fehlings

MG (2019) Sensory cortical control of movement. Nat Neurosci.

Kirkcaldie M, Watson c, Paxinos G, Franklin K (2012) Straightening out the mouse neocortex. In:

Australian Neuroscience Society Annual Conference. Goald Coast, QLD.

Lavoie S, Drew T (2002) Discharge characteristics of neurons in the red nucleus during voluntary gait

modifications: a comparison with the motor cortex. J Neurophysiol 88:1791-1814.

Lemon R (2019) Recent advances in our understanding of the primate corticospinal system. F1000Res

8.

Lemon RN (2008) Descending pathways in motor control. Annu Rev Neurosci 31:195-218.

Li CX, Waters RS (1991) Organization of the mouse motor cortex studied by retrograde tracing and

intracortical microstimulation (ICMS) mapping. Can J Neurol Sci 18:28-38.

Libbrecht S, Van den Haute C, Malinouskaya L, Gijsbers R, Baekelandt V (2017) Evaluation of WGA-

Cre-dependent topological transgene expression in the rodent brain. Brain Struct Funct 222:717-733.

Liu Y, Latremoliere A, Li X, Zhang Z, Chen M, Wang X, Fang C, Zhu J, Alexandre C, Gao Z, Chen B, Ding

X, Zhou JY, Zhang Y, Chen C, Wang KH, Woolf CJ, He Z (2018) Touch and tactile neuropathic pain

sensitivity are set by corticospinal projections. Nature 561:547-550.

Lopez-Bendito G, Sturgess K, Erdelyi F, Szabo G, Molnar Z, Paulsen O (2004) Preferential origin and

layer destination of GAD65-GFP cortical interneurons. Cereb Cortex 14:1122-1133.

Lorenzo LE, Ramien M, St Louis M, De Koninck Y, Ribeiro-da-Silva A (2008) Postnatal changes in the

Rexed lamination and markers of nociceptive afferents in the superficial dorsal horn of the rat.

JComp Neurol 508:592-604.

Mao T, Kusefoglu D, Hooks BM, Huber D, Petreanu L, Svoboda K (2011) Long-range neuronal circuits

underlying the interaction between sensory and motor cortex. Neuron 72:111-123.

McComas AJ (2016) Hypothesis: Hughlings Jackson and presynaptic inhibition: is there a big picture? J

Neurophysiol 116:41-50.

Mesnage B, Gaillard S, Godin A, Rodeau JL, Hammer M, von Engelhardt J, Wiseman P, De Koninck Y,

Schlichter R, Cordero-Erausquin M (2011) Morphological and functional characterization of

cholinergic interneurons of the dorsal horn of the mouse spinal cord. J Comp Neurol 519:3139-3158.

Miller S, Van Der Burg J, Van Der Meche F (1975) Coordination of movements of the hindlimbs and

forelimbs in different forms of locomotion in normal and decerebrate cats. Brain Res 91:217-237.

Miri A, Warriner CL, Seely JS, Elsayed GF, Cunningham JP, Churchland MM, Jessell TM (2017)

Behaviorally Selective Engagement of Short-Latency Effector Pathways by Motor Cortex. Neuron

95:683-696 e611.

Nathan PW, Smith M, Deacon P (1996) Vestibulospinal, reticulospinal and descending propriospinal

nerve fibres in man. Brain 119 (Pt 6):1809-1833.

Ni Y, Nawabi H, Liu X, Yang L, Miyamichi K, Tedeschi A, Xu B, Wall NR, Callaway EM, He Z (2014)

Characterization of long descending premotor propriospinal neurons in the spinal cord. J Neurosci

34:9404-9417.

Olivares-Moreno R, Moreno-Lopez Y, Concha L, Martinez-Lorenzana G, Condes-Lara M, Cordero-

Erausquin M, Rojas-Piloni G (2017) The rat corticospinal system is functionally and anatomically

segregated. Brain Struct Funct 222:3945-3958.

Pawlowski SA, Gaillard S, Ghorayeb I, Ribeiro-da-Silva A, Schlichter R, Cordero-Erausquin M (2013) A

novel population of cholinergic neurons in the macaque spinal dorsal horn of potential clinical

relevance for pain therapy. J Neurosci 33:3727-3737.

Porter R (1985) The corticomotoneuronal component of the pyramidal tract: corticomotoneuronal

connections and functions in primates. Brain Res 357:1-26.

Prescott SA, De Koninck Y (2003) Gain control of firing rate by shunting inhibition: roles of synaptic

noise and dendritic saturation. Proc Natl Acad Sci U S A 100:2076-2081.

Reed WR, Shum-Siu A, Onifer SM, Magnuson DS (2006) Inter-enlargement pathways in the

ventrolateral funiculus of the adult rat spinal cord. Neuroscience 142:1195-1207.

Reed WR, Shum-Siu A, Whelan A, Onifer SM, Magnuson DS (2009) Anterograde labeling of

ventrolateral funiculus pathways with spinal enlargement connections in the adult rat spinal cord.

Brain Res 1302:76-84.

Ribeiro-da-Silva A, Cuello AC (1990) Choline acetyltransferase-immunoreactive profiles are

presynaptic to primary sensory fibers in the rat superficial dorsal horn. J Comp Neurol 295:370-384.

Rudomin P, Schmidt RF (1999) Presynaptic inhibition in the vertebrate spinal cord revisited. Exp Brain

Res 129:1-37.

Rudomin P, Solodkin M, Jimenez I (1986) PAD and PAH response patterns of group Ia- and Ib-fibers to

cutaneous and descending inputs in the cat spinal cord. J Neurophysiol 56:987-1006.

Seki K, Fetz EE (2012) Gating of sensory input at spinal and cortical levels during preparation and

execution of voluntary movement. J Neurosci 32:890-902.

Seki K, Perlmutter SI, Fetz EE (2003) Sensory input to primate spinal cord is presynaptically inhibited

during voluntary movement. Nat Neurosci 6:1309-1316.

Seki K, Perlmutter SI, Fetz EE (2009) Task-dependent modulation of primary afferent depolarization

in cervical spinal cord of monkeys performing an instructed delay task. J Neurophysiol 102:85-99.

Shreckengost J, Halder M, Mena-Avila E, Garcia-Ramirez DL, Quevedo J, Hochman S (2021) Nicotinic

receptor modulation of primary afferent excitability with selective regulation of Adelta-mediated

spinal actions. J Neurophysiol 125:568-585.

Sirois J, Frigon A, Gossard JP (2013) Independent control of presynaptic inhibition by reticulospinal

and sensory inputs at rest and during rhythmic activities in the cat. J Neurosci 33:8055-8067.

Starkey ML, Barritt AW, Yip PK, Davies M, Hamers FP, McMahon SB, Bradbury EJ (2005) Assessing

behavioural function following a pyramidotomy lesion of the corticospinal tract in adult mice. Exp

Neurol 195:524-539.

Tappe A, Klugmann M, Luo C, Hirlinger D, Agarwal N, Benrath J, Ehrengruber MU, During MJ, Kuner R

(2006) Synaptic scaffolding protein Homer1a protects against chronic inflammatory pain. Nat Med

12:677-681.

Tennant KA, Adkins DL, Donlan NA, Asay AL, Thomas N, Kleim JA, Jones TA (2011) The organization of

the forelimb representation of the C57BL/6 mouse motor cortex as defined by intracortical

microstimulation and cytoarchitecture. Cereb Cortex 21:865-876.

Tjolsen A, Berge OG, Hunskaar S, Rosland JH, Hole K (1992) The formalin test: an evaluation of the

method. Pain 51:5-17.

Todd AJ (1991) Immunohistochemical evidence that acetylcholine and glycine exist in different

populations of GABAergic neurons in lamina III of rat spinal dorsal horn. Neuroscience 44:741-746.

Ueno M, Hayano Y, Nakagawa H, Yamashita T (2012) Intraspinal rewiring of the corticospinal tract

requires target-derived brain-derived neurotrophic factor and compensates lost function after brain

injury. Brain 135:1253-1267.

Ueno M, Nakamura Y, Li J, Gu Z, Niehaus J, Maezawa M, Crone SA, Goulding M, Baccei ML, Yoshida Y

(2018) Corticospinal Circuits from the Sensory and Motor Cortices Differentially Regulate Skilled

Movements through Distinct Spinal Interneurons. Cell Rep 23:1286-1300 e1287.

von Engelhardt J, Eliava M, Meyer AH, Rozov A, Monyer H (2007) Functional characterization of

intrinsic cholinergic interneurons in the cortex. JNeurosci 27:5633-5642.

Wall PD (1958) Excitability changes in afferent fibre terminations and their relation to slow

potentials. J Physiol 142:1-21.

Wall PD, Lidierth M (1997) Five sources of a dorsal root potential: their interactions and origins in the

superficial dorsal horn. J Neurophysiol 78:860-871.

Wang X, Liu Y, Li X, Zhang Z, Yang H, Zhang Y, Williams PR, Alwahab NSA, Kapur K, Yu B, Chen M, Ding

H, Gerfen CR, Wang KH, He Z (2017) Deconstruction of Corticospinal Circuits for Goal-Directed Motor

Skills. Cell 171:440-455 e414.

Zelenin PV, Beloozerova IN, Sirota MG, Orlovsky GN, Deliagina TG (2010) Activity of red nucleus

neurons in the cat during postural corrections. J Neurosci 30:14533-14542.

Figure Legends

Figure 1. The same cortical area evokes dorsal root potentials and muscular contractions of the hindlimbs.

A. Schematic experimental design for cortically-evoked (Cx-) DRP and EMG recordings elicited by photostimulation of the contralateral sensorimotor cortex of Thy1::ChR2 mice. DRPs correspond to the presynaptic depolarization of primary afferents propagating antidromically to the suction recording electrode containing a lumbar dorsal root, while EMGs are recorded in the *TA* muscle.

B. Representative traces of a cortically-evoked DRP (top red trace, average of 30 sweeps) and EMG (bottom blue trace, average of 3 sweeps) after photostimulation (blue window) of the contralateral cortex.

C. *Left,* Heatmaps of the amplitude of the responses of cortically-evoked DRPs (top) and EMGs (bottom) in % or the maximum response. *Right*, Overlap of the two maps (red: DRP, blue: EMG) presented as isopotential contour plots (three color grades corresponding to 37%, 50% and 63% of maximum value). Coordinates of the cortex are expressed in mm and centered on *Bregma*. M1: primary motor cortex, M2: secondary motor cortex, and S1: primary sensory cortex, according to the Paxinos atlas (Kirkcaldie et al., 2012).

D. Schematic experimental design to identify GAD65-expressing neurons amongst corticospinal neurons targets: AAV1-CBA-WGA-CRE was injected in the sensorimotor cortex of TdTomato-flex X GAD65::GFP mice. Analysis of extend of the injection site is presented in Figure 1 – Figure Supplement 1, and of the monosynaptic nature of the transynaptic tracing in Figure 1 - Figure Supplement 2.

E. Photomicrographs (z projection of confocal images) of a GAD65::GFP mouse lumbar dorsal horn (localization of the view indicated in the inset) after transynaptic labeling from the

hindlimb sensorimotor contralateral cortex. A target of the CST (expressing TdTomato after

transynaptic transfer of WGA-Cre) also expresses GAD65::GFP. CST: corticospinal tract. Similar

experiment in ChAT::EGFP mice is presented in Figure 1 – Figure Supplement 3.

Figure 2. The corticospinal tract is essential for cortically-evoked inhibition of primary afferents.

A. *Left*, diagram showing direct and indirect cortical-to-spinal paths. In green: paths potentially

activated by photostimulation in the cortex of Thy1::ChR2 mice. *Right*, experimental design.

B. *Left*, drawing showing the extent of the pyramidal lesion (in grey, pyx) in the frontal plane

(distance to *Bregma* indicated) in two animals whose recordings are respectively illustrated in

C and D. *Right*, diagram showing the indirect cortical-to-spinal paths spared by the

pyramidotomy. Systematic histological analysis of the pyramidotomies is presented in Figure

2 – Figure Supplement 1.

C. *Left*, Representative traces of cortically-evoked DRPs recording before (red) and after (grey)

CST lesion (average of 30 traces). *Right*, the pyramidotomy abolishes cortically-evoked DRPs

(p= 0.006, one-way ANOVA, n= 3 mice). Post-hoc test Holm-Sidak, *P < 0.05

D. *Left*, Representative traces of cortically-evoked EMG recording before (blue) and after (grey)

pyramidal lesion (average of 3 traces). *Right*, There is no significant change in the S/N ratio of

the EMG response before and after the pyramidal lesion (93,8%, 96,9%, 109,3% prior to

pyramidotomy to 77.2% after, p=0.517, one-way ANOVA, n= 4 mice). Grey zone: noise level

(see Methods). EMG Z-scores for individual mice before and after pyramidotomy are

presented in Figure 2 – Figure Supplement 2.

Figure 3. The corticospinal tract can encode muscular contractions.

A. *Left*, diagram illustrating corticospinal neurons and their collaterals expressing ChR2 after

retrograde infection in the lumbar cord with a ChR2-retro AAV. *Right*, experimental design:

the photostimulation and EMG recording session took place at least 3 weeks after the

infection.

B. Sagittal image from the sensorimotor cortex (Lateral= 0.68 mm), showing retrogradely labeled

corticospinal neurons (expressing ChR2-TdTomato).

C. Drawing of a cortex top view, showing the localization of retrogradely labeled corticospinal

neurons in the 5 animals (solid lines). They are within the area functionally determined as

inducing DRPs and EMGs in the hindlimb (dashed lines).

D. *Left*, drawing showing the extent of the pyramidal lesion in the animal whose recordings are

illustrated in E. *Right*, diagram showing the supraspinal collaterals of ChR2-expressing CS

neurons spared by the pyramidotomy. Systematic histological analysis of the pyramidotomies

is presented in Figure 2 – Figure Supplement 1.

E. *Left*, cortically evoked EMG recordings before (blue) and after (grey) the pyramidal lesion*.

Right,* the pyramidotomy abolishes cortically-evoked EMG ($p=0.012$, one-way ANOVA, n= 5

mice). Post-hoc test Holm-Sidak, *$P < 0.05$. Grey zone: noise level (see Methods). EMG Z-

scores for individual mice before and after pyramidotomy are presented in Figure 2 – Figure

Supplement 2.

The blue window on recording traces indicates photostimulation. SMC: sensorimotor cortex,

BSMN: brain stem motor nuclei, C-Th: cervico-thoracic spinal cord, L-SC: Lumbo-sacral spinal cord.

Figure 4. Lumbar corticospinal postsynaptic neurons encode DRPs but no movements.

A. Experimental design: the targets of the CST are labeled through a transynaptic approach

consisting of AAV1-CBA-WGA-CRE injection in the hindlimb sensorimotor cortex of TdTomato-

flex mice.

B. Localization of the spinal targets of the CST: heatmap showing the distribution of the neurons

in the lumbar cord (6 mm long) projected into the transverse plane (average of 9 mice) (*left*)

or horizontal plane (average of 6 mice) (*right*) plane.

C. *Left*, Diagram illustrating that only lumbar direct targets of the CST express CHETA in the

following experiment. *Right*, Experimental design: TdTomato-flex mice received an injection

of AAV1-CBA-WGA-CRE in the hindlimb sensorimotor cortex and an injection of AAV1-Efl-flex-

CHETA-eYFP in the L4 spinal segment.

D. Photomicrographs (z projection of confocal images) from the dorsal horn of the spinal cord

(laminae V/VI); the arrows point at two targets of the CST expressing TdTomato+ and CHETA-

eYFP.

E. Experimental design illustrating spinal photostimulation of the lumbar targets of the CST.

F. Representative traces of DRP (red trace, average of 60 traces) and EMG (blue trace, average

of 3 traces) recordings from the same animal after photostimulation of the lumbar targets of

the CST (blue window).

G. Photostimulation of the lumbar targets of the CST induces DRPs (left) but no EMG signal (right).

Grey zone: noise level (see Methods). EMG Z-scores for individual mice are presented in Figure

2 – Figure Supplement 2.

SMC: sensorimotor cortex, BSMN: brain stem motor nuclei, C-Th: cervico-thoracic spinal cord, L-

SC:Lumbo-sacral spinal cord.

Figure Supplement material:

Figure 1 - Figure Supplement 1. Sensorimotor injection site analyses : cortical layer V fluorescence

quantification.

Figure 1 - Figure Supplement 2. Monosynaptic transynaptic tracing from the cortex

Figure 1 - Figure Supplement 3. Postsynaptic corticospinal neurons do not colocalize with ChAT.

Figure 2 - Figure Supplement 1. Histology of the pyramidotomies.

Figure 2 Figure Supplement 2. Z-score of individual EMG responses assessing the significance of the

corresponding S/N ratio

Figure 3 - Figure Supplement 1. DRPs are induced by photostimulation of ChR2-expressing CS

neurons (but not by GFP-expressing CS neurons)

Figure 4 - Figure Supplement 1. Strategy to label the spinal targets of corticospinal neurons

Figure 4 - Figure Supplement 2. Strategy to stimulate exclusively the spinal targets of corticospinal

neurons

Figure 4 - Figure Supplement 3. Efficacy of the surface spinal photostimulation

Figure 1. The same cortical area evokes dorsal root potentials and muscular contractions of the hindlimbs.

A. Schematic experimental design for cortically-evoked (Cx-) DRP and EMG recordings elicited by photostimulation of the contralateral sensorimotor cortex of Thy1::ChR2 mice. DRPs correspond to the presynaptic depolarization of primary afferents propagating antidromically to the suction recording electrode containing a lumbar dorsal root, while EMGs are recorded in the *TA* muscle.

B. Representative traces of a cortically-evoked DRP (top red trace, average of 30 sweeps) and EMG (bottom blue trace, average of 3 sweeps) after photostimulation (blue window) of the contralateral cortex.

C. *Left*, Heatmaps of the amplitude of the responses of cortically-evoked DRPs (top) and EMGs (bottom) in % or the maximum response. *Right*, Overlap of the two maps (red: DRP, blue: EMG) presented as isopotential contour plots (three color grades corresponding to 37%, 50% and 63% of maximum value). Coordinates of the cortex are expressed in mm and centered on *Bregma*. M1: primary motor cortex, M2: secondary motor cortex, and S1: primary sensory cortex, according to the Paxinos atlas (Kirkcaldie et al., 2012).

D. Schematic experimental design to identify GAD65-expressing neurons amongst corticospinal neurons targets: AAV1-CBA-WGA-CRE was injected in the sensorimotor cortex of TdTomato-flex X GAD65::GFP mice. Analysis of extend of the injection site is presented in Figure 1 – Figure Supplement. 1, and of the monosynaptic nature of the transynaptic tracing in Figure 1 - Figure Supplement 2.

E. Photomicrographs (z projection of confocal images) of a GAD65::GFP mouse lumbar dorsal horn (localization of the view indicated in the inset) after transynaptic labeling from the hindlimb sensorimotor contralateral cortex. A target of the CST (expressing TdTomato after transynaptic transfer of WGA-Cre) also expresses GAD65::GFP. CST: corticospinal tract. Similar experiment in ChAT::EGFP mice is presented in Figure 1 – Figure Supplement 3.

Figure 2. The corticospinal tract is essential for cortically-evoked inhibition of primary afferents.

A. *Left*, diagram showing direct and indirect cortical-to-spinal paths. In green: paths potentially activated by photostimulation in the cortex of Thy1::ChR2 mice. *Right*, experimental design.

B. *Left*, drawing showing the extent of the pyramidal lesion (in grey, pyx) in the frontal plane (distance to *Bregma* indicated) in two animals whose recordings are respectively illustrated in C and D. *Right*, diagram showing the indirect cortical-to-spinal paths spared by the pyramidotomy. Systematic histological analysis of the pyramidotomies is presented in Figure 2 – Figure Supplement 1.

C. *Left*, Representative traces of cortically-evoked DRPs recording before (red) and after (grey) CST lesion (average of 30 traces). *Right*, the pyramidotomy abolishes cortically-evoked DRPs (p= 0.006, one-way ANOVA, n= 3 mice). Post-hoc test Holm-Sidak, *P < 0.05

D. *Left*, Representative traces of cortically-evoked EMG recording before (blue) and after (grey) pyramidal lesion (average of 3 traces). *Right*, There is no significant change in the S/N ratio of the EMG response before and after the pyramidal lesion (p=0.517, one-way ANOVA, n= 4 mice). Grey zone: noise level (see Methods). EMG Z-scores for individual mice before and after pyramidotomy are presented in Figure 2 – Figure Supplement 2.

Figure 3. The corticospinal tract can encode muscular contractions.

A. *Left*, diagram illustrating corticospinal neurons and their collaterals expressing ChR2 after retrograde infection in the lumbar cord with a retroAAV. *Right*, experimental design: the photostimulation and EMG recording session took place at least 3 weeks after the infection.

B. Sagittal image from the sensorimotor cortex (Lateral= 0.68 mm), showing retrogradely labeled corticospinal neurons (expressing ChR2-TdTomato fluorescence).

C. Drawing of a cortex top view, showing the localization of retrogradely labeled corticospinal neurons in the 5 animals (solid lines). They are within the area functionally determined as inducing DRPs and EMGs in the hindlimb (dashed lines).

D. *Left*, drawing showing the extent of the pyramidal lesion in the animal whose recordings are illustrated in E. *Right*, diagram showing the supraspinal collaterals of ChR2-expressing CS neurons spared by the pyramidotomy. Systematic histological analysis of the pyramidotomies is presented in Figure 2 – Figure Supplement 1.

E. *Left*, cortically evoked EMG recordings before (blue) and after (grey) the pyramidal lesion. *Right,* the pyramidotomy abolishes cortically-evoked EMG (p=0.012, one-way ANOVA, n= 5 mice). Post-hoc test Holm-Sidak, *P < 0.05. Grey zone: noise level (see Methods). EMG Z-scores for individual mice before and after pyramidotomy are presented in Figure 2 – Figure Supplement 2.

The blue window on recording traces indicates light stimulation. SMC: sensorimotor cortex, BSMN: brain stem motor nuclei, C-Th: cervico-thoracic spinal cord, L-SC: Lumbo-sacral spinal cord.

Figure 4. Lumbar corticospinal postsynaptic neurons encode DRPs but no movements.

A. Experimental design: the targets of the CST are labeled through a transynaptic approach consisting of AAV1-CBA-WGA-CRE injection in the hindlimb sensorimotor cortex of TdTomato-flex mice.

B. Localization of the spinal targets of the CST: heatmap showing the distribution of the neurons in the lumbar cord (6 mm long) projected into the transverse plane (average of 9 mice) (*left*) or horizontal plane (average of 6 mice) (*right*) plane.

C. *Left*, Diagram illustrating that only lumbar direct targets of the CST express CHETA in the following experiment. *Right*, Experimental design: TdTomato-flex mice received an injection of AAV1-CBA-WGA-CRE in the hindlimb sensorimotor cortex and an injection of AAV1-Efl-flex-CHETA-eYFP in the L4 spinal segment.

D. Photomicrographs (z projection of confocal images) from the dorsal horn of the spinal cord (laminae V/VI); the arrows point at two targets of the CST expressing TdTomato+ and CHETA-eYFP.

E. Experimental design illustrating spinal photostimulation of the lumbar targets of the CST.

F. Representative traces of DRP (red trace, average of 60 traces) and EMG (blue trace, average of 3 traces) recordings from the same animal after photostimulation of the lumbar targets of the CST (blue window).

G. Photostimulation of the lumbar targets of the CST induces DRPs (left) but no EMG signal (right). Grey zone: noise level (see Methods). EMG Z-scores for individual mice are presented in Figure 2 – Figure Supplement 2.

SMC: sensorimotor cortex, BSMN: brain stem motor nuclei, C-Th: cervico-thoracic spinal cord, L-SC:Lumbo-sacral spinal cord.