1

Mechanical control of morphogenetic robustness in an inherently challenging environment

Emmanuel Martin[1], Sophie Theis[1,2], Guillaume Gay[2], Bruno Monier[1], Christian Rouvière[3] and Magali

Suzanne[1]

[1] LBCMCP, Centre de Biologie Intégrative (CBI), Université de Toulouse, CNRS, UPS, France

[2] Morphogénie Logiciels, 32110 St Martin d'Armagnac, France

[3] Image Processing Facility, Centre de Biologie Intégrative (CBI), Université de Toulouse, CNRS, UPS, France

Correspondence: magali.suzanne@univ-tlse3.fr; guillaume@damcb.com;

Lead Contact: magali.suzanne@univ-tlse3.fr

Abstract

16   **Epithelial sheets undergo highly reproducible remodeling to shape organs. This**

17   **stereotyped morphogenesis depends on a well-defined sequence of events leading to the**

18   **regionalized expression of developmental patterning genes that finally triggers downstream**

19   **mechanical forces to drive tissue remodeling at a pre-defined position. However, how tissue**

20   **mechanics controls morphogenetic robustness when challenged by intrinsic perturbations in**

21   **close proximity has never been addressed.**

22   **Here, we show that a bias in force propagation ensures stereotyped morphogenesis**

despite the presence of mechanical noise in the environment. We found that knockdown of the

Arp2/3 complex member Arpc5 specifically affects fold directionality without altering neither

the developmental nor the force generation patterns. By combining *in silico* modeling,

biophysical and ad hoc genetic tools, our data reveal that junctional Myosin II planar polarity

favors long-range force channeling and ensures folding robustness, avoiding force scattering and

thus isolating the fold domain from surrounding mechanical perturbations.

Introduction

An intriguing characteristic of biology is the remarkable reproducibility in size and shape of a
given structure from one individual to another. This trait, called robustness, is characterized by a low
level of variation of a given phenotype when subjected to environmental or genetic variations
(Waddington, 1942).

An important factor ensuring robustness is the existence of redundant mechanisms that
appeared to be frequently used as a way to ensure the generation of a specific trait. It has been
observed at different levels, between different genes, but also between different regulatory sequences
governing gene expression with the discovery of shadow enhancers or more recently between
mechanical networks governing tissue shape generation (Frankel et al., 2010; Perry et al., 2010; Smith
et al., 2018; Yevick et al., 2019). These discoveries highlight the importance of backup mechanisms to
ensure a correct shape. However, they did not inform on the process of canalization of a particular trait
and the protection mechanisms taking place to avoid phenotypic variation in front of environmental
challenges.

The establishment of precise patterning constitutes the initial step for the development of
anatomical structures. Not surprisingly, it has been the main focus in the field to unravel the
mechanisms responsible for the high level of precision observed in terms of growth, scale and
patterning of tissue and organs (Félix and Barkoulas, 2015; Lander, 2011).
Pioneer studies came from the study of signaling gradients in Drosophila, highlighting the high level
of precision of morphogen gradients giving rise to precise boundary of target genes expression, even
when challenged by fluctuation in gene dosage or morphogen production rate (Eldar et al., 2004;
Gregor et al., 2007; Hardway et al., 2008; He et al., 2008). It then became apparent that even if
morphogen gradients appear amazingly robust, a certain degree of variability exists from one cell to
another. This stochasticity in molecular and cellular processes emerged recently as an inherent part of
biological systems, opening a whole field of research focused on how a developing organism can deal
with such intrinsic noise and form stereotyped shapes (Ebert and Sharp, 2012; Gursky et al., 2012;
Hong et al., 2018; Lander, 2011).

Numerous studies highlighted the importance to buffer this noise. Indeed, even in situations
where, counter intuitively, heterogeneity has been shown to play an important role in shape robustness
such as in sepal formation, heterogeneity has to be buffered over time to ensure the formation of
regular and stereotyped organs (Hong et al., 2018). Different mechanisms of noise buffering have been
identified, mainly involved in fine-tuning the expression of genes involved in positional information
(Ebert and Sharp, 2012; Gursky et al., 2012; Herranz and Cohen, 2010; Lott et al., 2007; Manu et al.,
2009; Sato, 2018).

Downstream of this positional information and the establishment of tissue coordinates,
important drivers of tissue shape are mechanical forces, which govern cell shape remodeling and cell

rearrangement processes (Smith et al., 2018). The study of mechanical forces constitutes a growing

field in morphogenesis, underlying their crucial role in tissue dynamics. Indeed, cells and tissues are

physical entities, whose shape is determined by structural components, such as cytoskeletal proteins.

Among them, the molecular motor non-muscle myosin II and the filamentous actin associate to create

a dynamic network, which drives force generation or governs cell architecture. The re-distribution of

acto-myosin within specific subcellular domains drives specific changes in cell shape. These forces,

generated at a single cell level can propagate from one cell to another through adherens junction or

diffusible biomechanical signals (Lecuit and Yap, 2015), ultimately leading to large-scale changes and

tissue shape modification, although long-range propagation and channeling of mechanical forces

remains largely unexplored.

If the cellular machinery responsible for force generation is a key player in the construction of

a particular shape, only a few works addressed the question of mechanical contribution to robustness.

On the one hand, Hong and colleagues (Hong et al., 2018) proposed, based on theoretical modeling,

that tissue mechanics could buffer the heterogeneity observed at a single cell level in term of growth

and stiffness. More recently, this idea has been tested experimentally and mechanical forces have been

shown to buffer local heterogeneity both in zebrafish and in Drosophila (Akieda et al., 2019; Eritano

et al., 2020). On the other hand, Yevick and colleagues (Yevick et al., 2019) identified redundant

mechanical networks involved in the construction of a particular shape. However, despite an important

amount of works on mechanical forces as essential bricks in the construction of tissue shape, how

mechanics affects the degree of variability of a given phenotype in challenging environmental

conditions remain mostly unexplored. Indeed, while the generation and the propagation of forces are

well recognized as an important property of cells and tissues, they are mainly viewed as key

executioners of a pre-established developmental program or as a refining mechanism of intrinsic noise

rather than a guardian of the stereotypic nature and robustness of tissue shape in front of external

challenges.

Here, focusing on tissue invagination as a model of morphogenetic robustness, we identified

a genetic condition in which fold orientation, an extremely robust trait, becomes highly variable. This

variability appears to be independent of tissue patterning; with both tissue regionalization and intrinsic

mechanical signals occurring normally, while fold directionality is perturbed. These deviations point

at regions of high tension, which appear randomly around the fold domain, thus revealing the presence

of mechanical noise in the local environment. Finally, we found that Myosin II planar polarity is both

necessary and sufficient to ensure the robustness of fold directionality, favoring force channeling, and

thus protecting the invagination from neighboring mechanical noise.

Results

Morphogenesis variability in Arpc5 knockdown.

To address the role of tissue mechanics in morphogenetic robustness, we used Drosophila leg

development, a model particularly appropriate since it undergoes a highly stereotyped morphogenesis

with four parallel folds formed during development, in the most distal part of the leg or tarsal region

(Fig1a). To quantify the robustness of this morphogenetic process in a control situation, we measured

the variability of fold positioning (distance from the predicted fold), fold orientation (angle formed

with the proximal-distal axis of the leg) and further quantified the relative orientation of tarsal folds

between them or fold parallelism (Fig1c-f, control). Both fold positioning and orientation appear to be

extremely robust in the control situation, as shown by the low standard deviation observed for each of

these parameters.

To decipher how this particular shape is established and maintained from one individual to

another, we selected, from an unbiased screen expressing a library of RNAi in the tarsal domain of the

leg (manuscript in preparation), candidates affecting specifically fold positioning. Interestingly, we

found a puzzling phenotype, never described before, of misoriented folds. This phenotype was found

for several components of the Arp2/3 complex known to regulate the branched actin network,

including Arpc5, Arpc3a and Arp2 (Fig1b, S1b) (Chesarone and Goode, 2009; Pollard and Beltzner,

2002; Robinson et al., 2001). Interestingly, defects show a wide range of variability and ranged from a

complete absence of folds to fold deviation to normal folding (FigS1b-c). Since these proteins are all

part of the core Arp2/3 complex and their RNAi give similar phenotypes (FigS1b-c), we focused our

analysis on one of them, Arpc5, whose inactivation by RNAi in our experimental conditions resulted

in a 50% reduction of its mRNA level (FigS1a). The implication of Arp2/3 in fold directionality was

unexpected. To further characterize this new phenotype, we first compared the defects observed

during fold formation and at the end of the process. We observed an increase in the proportion of fold

deviation over time, going from 1/3 to 3/4. These results indicate that the deviations are not transient

but stable defects. We further observed that the absence of folding becomes less frequent with time,

suggesting together with the increase of fold deviations that a proportion of folds are delayed and

finally misoriented (FigS1c). To further characterize this mutant phenotype, we choose to focus on

mid-fold formation, a stage at which fold positioning can be defined more accurately. We then

measured the variability of fold positioning, fold orientation and fold parallelism of the subpopulation

of discs (about 2/3) displaying normal or misoriented folds (Fig1c-f), and excluding unfolded legs

since fold direction is impossible to address in these ones (mentioned as flat in FigS1c). For each of

these parameters, standard deviation was significantly higher than in the control, highlighting the high

degree of variability observed in the Arpc5 knockdown condition (Fig1c-f), a phenotype characteristic

of a failure of morphogenetic robustness.

Developmental patterning in Arpc5 knockdown

Although the regulation of fold orientation has never been addressed, fold positioning is
known to be determined by the sequential establishment of positional information along the
developing leg tissue, starting with the restricted expression of morphogens such as *wingless* and
decapentaplegic (Lecuit and Cohen, 1997), the subsequent proximal-distal regionalization through the
expression of patterning genes such as *Hth*, *Dac* and *Dll* (Wu and Cohen, 1999), followed by the
segmental activation of the Notch pathway in the distal part of each segment (de Celis et al., 1998;
Rauskolb and Irvine, 1999). Finally, in the tarsal region, pro-apoptotic genes are expressed in a few
rows of cells in the distal part of each segment (Manjón et al., 2007). This positional information is
then translated into a "patchy" pattern of apoptotic cells in the predicted fold domain. Previous work
that reconstituted the localization of apoptotic cells in fixed samples over time showed that apoptotic
cells appear first on the ventral part, then in the lateral domains, and finally in the most dorsal domain
of this ring-shaped predicted fold (Monier et al., 2015). Before their elimination, these apoptotic cells
exert apico-basal forces, which constitute the initiator mechanical signals for tissue remodeling. These
forces are transmitted to their neighbors leading to an increase in apical myosin accumulation, apical
constriction and eventually tissue folding, a process that lasts 3-4h *in vivo* (Monier et al., 2015).
Since robustness has been shown to rely on the establishment of robust positional information,
we first analyzed the expression pattern of genes known to be involved in fold positioning.
Importantly, in the Arpc5 knockdown condition, the segmental activation of Notch, characterized by
the expression of the Notch target gene *Deadpan* (*Dpn*), is maintained (Fig2a). Indeed, on top of *Dpn*
expression in neurons (see asterisks in Fig2a), consistent with its identification as a pan-neural gene
(Younger-Shepherd et al., 1992), *Dpn* is expressed in stripes in the distal leg, both in the control and
Arpc5 RNAi condition. We further characterized *Dpn* expression domain and found that the width of
its domain is slightly smaller in Arpc5 RNAi (FigS2d), consistent with the role played by Arp2/3 in
Notch activation in other contexts (Rajan et al., 2009). However, the orientation of *Dpn* stripes in
relation to the proximal-distal axis, their parallelism and their curvature were unaffected in Arpc5
knockdown condition (Fig2c; FigS2a,c). Fold deviation in the Arpc5 knockdown appears to result
from a partial dissociation of the folding process from the positional information. Thus, while folds
follow the segmental stripes of Notch activation domain in the control, folds deviate from these
positional cues in the Arpc5 knockdown (Fig2e-f, MovieS1 and S2). We next analyzed the apoptotic
pattern in the Arpc5 knockdown, which appears intact as shown by the unperturbed expression of the
pro-apoptotic gene *reaper* (Fig2b,d; FigS2b,e,f) and the frequency of apoptotic cells in the fold
domain (Fig2g). Finally, the ability of dying cells to generate mechanical signals is identical to the
control, as shown by the formation of apico-basal transient structures of Myosin II in dying cells
(Fig2h), their ability to deform the apical surface (Fig2h) and to generate apico-basal tension (Fig2i,
MovieS3). Thus, while folds deviate in Arpc5 knockdown condition, positional information resulting
from the developmental patterning and the subsequent mechanisms known to be involved in the fold
formation are unaffected. This surprising result prompted us to revisit the prevailing model of

morphogenetic robustness relying on strict regulation of morphogen gradients or key identity genes
(Gilmour et al., 2017). We hypothesized that fold deviation could either be due to the appearance of a
new source of perturbation or by an increased sensitivity to existing perturbations in the knockdown
condition.

180   -

Leg morphogenesis occurs in a mechanically noisy environment.

We next asked if folds deviate towards particular regions of the tissue in Arpc5 knockdown.
We noticed that misoriented folds deviate from the predicted fold domain to more proximal or distal
regions (Fig3a). Proximally, deviated folds often joined a straight cellular alignment present at the
distal border of Notch activation (Fig3b, FigS3a) while distally deviated folds frequently headed
towards apico-basal structures of Myosin II (Fig3d) that are not associated with apoptosis (FigS3b)
and are present at various distances from the predicted fold domain. Importantly, both regions are
strongly enriched in Myosin II, either apically or apico-basally, which usually coincides with high
tension. Laser ablation experiments showed that tension was indeed significantly higher at the "Notch
border" and at non-apoptotic apico-basal myosin structures (Fig3c, e), compared to the neighboring
tissue in Arpc5 knockdown condition. Altogether, these data indicate that, when Arpc5 function is
reduced, folds head towards regions of high tension. This suggests that these regions of high tension
could create mechanical perturbations in close vicinity to the fold domain.

To figure out whether the potential mechanical interference associated with structures under
high tension are either ectopically created or increased following the Arpc5 knockdown, or,
alternatively, present yet masked during normal development, we analyzed the Myosin II pattern in
control flies. Interestingly, both the "Notch border" and non-apoptotic apico-basal myosin structures
were present in the control (Fig3b,d, FigS3) and the tension borne by these structures was comparable
to that measured in Arpc5 knockdown condition (Fig3c,e, MovieS4). To further characterize tissue
mechanics around the fold region, we mapped these regions of high tension and found that these
mechanical perturbations are frequent and located at variable positions around the fold domain both in
the control and in Arpc5 knockdown condition, suggesting the existence of a mechanical noise during
fold morphogenesis (Fig3f). Altogether, these data suggest that leg fold morphogenesis is permanently
challenged by surrounding remodeling events and becomes more sensitive to mechanical perturbations
in the Arpc5 knockdown.

To test the impact of mechanical perturbations on fold formation, we turned to *in silico*
modeling. We previously developed a vertex model able to reproduce leg fold formation both in terms
of tissue shape and cellular organization (Monier et al., 2015). However, this morphogenetic event
was considered as an isolated process in these simulations, and the only mechanical forces applied
were those originating from the apoptotic cells located in the predicted fold domain. Therefore, we

implemented the model (see M&M and FigS4) and integrated random mechanical perturbations in
close vicinity to the predicted fold domain, mimicking the mechanical noise observed *in vivo* (Fig3h).
Interestingly, mechanical noise appears sufficient to induce fold deviations in the simulations,
mimicking the Arpc5 knockdown condition (Fig3g). We further quantified the variability of the
phenotype observed when a random pattern of mechanical perturbations was integrated in the model
and found that, while fold formation appeared robust in the absence of potential interferences, fold
positioning and orientation become consistently more variable in the presence of mechanical
perturbations in the vicinity of the predicted fold domain (Fig3i-j).

219   Altogether, these results strongly suggest that some kind of isolation is required between the
220 fold domain and the neighboring tissue to avoid surrounding mechanical interference and ensure
221 morphogenetic robustness.

Arp2/3 controls Myosin II planar polarization

224   We next asked how the predicted fold domain could become insensitive to nearby mechanical
noise. Strikingly, in the control we noticed that Myosin II bore a polarized junctional distribution
specifically in the fold domain, with a stronger accumulation in the cellular junctions parallel to the
future folds (or circumferential junctions) than in perpendicular ones (or proximal-distal junctions).
This polarity, although already present at the onset of fold formation, is accentuated during fold
progression and coincides with an increase of cell anisotropy (Fig4a-d). Importantly, while the total
amount of Myosin II was unchanged in Arpc5 knockdown condition (FigS5a), Myosin II planar
polarity was lost and cell anisotropy was no more increased in the fold domain (Fig4a-d). These
observations show that folding robustness coincides with planar polarization of Myosin II in the fly
leg.

234   To decipher how Arp2/3 could drive Myosin II planar polarity, we first characterized the
spatial distribution of the F-actin network (using phalloidin) and of the Arp2/3 complex (using UAS-
Arp3-GFP, a construct whose expression does not induce any visible defect, see FigS5b).
Interestingly, we observed a polarized and anti-correlative distribution between F-actin and the Arp3-
GFP fusion protein. While F-actin was preferentially accumulated in junctions parallel to the fold like
Myosin II, Arp3-GFP was mainly present in junctions perpendicular to the fold (Fig5a-d).
Interestingly, the polarized distribution of F-actin is lost in Arpc5 RNAi condition, indicating a role of
Arp2/3 in F-actin polarity.

242   We further characterized the flows of the medio-apical Myosin II and observed a slight bias in
its directionality in the control, with more movement in the proximo-distal axis than in the
circumferential one (FigS5e). Interestingly this bias is lost in Arpc5 knockdown (FigS5f), suggesting
that Arp2/3 could influence Myosin II polarity through the regulation of medio-apical myosin flow.

Finally, we observed a slight polarity in the distribution of adherens junctions (using β-
catenin-GFP), with enrichment similar to Arp2/3. However, this polarity is unperturbed in Arpc5
knockdown (FigS5c-d).

Planar polarity ensures morphogenetic robustness

To test if perturbing Myosin II planar polarity was sufficient to alter fold robustness *in vivo*, we
decided to disrupt Myosin II polarity by an independent method. We generated a new variant of the
nanobody-based GFP-trap technique previously described (Harmansa et al., 2017; Harmansa et al.,
2015), which traps the endogenous Myosin II-GFP fusion protein (Myosin Regulatory Light Chain,
MRLC-GFP) at adherens junctions (Fig6a-b, FigS6a, see M&M). We found that the distribution of
junctional Myosin II is mainly isotropic in this condition, although not totally homogeneous.
Interestingly, using conditional expression of this AJs-GFP-trap construct in the presence of Myosin
II-GFP, we frequently observed deviated folds, mimicking the defects observed in the Arpc5
knockdown (Fig6c), while expression of AJs-GFP-trap in the absence of any GFP does not alter leg
morphogenesis (FigS6b). We also observed that the variability of fold positioning, fold orientation and
fold parallelism was significantly increased in this condition of Myosin II polarization defect
compared to the control (Fig6d-f). These experiments show that decreasing or abolishing Myosin II
planar polarity through two independent strategies (Arpc5 knockdown and AJs-GFP-trap/Myosin II-
GFP) leads to an increase of variability of fold orientation in the fly leg.

We then wondered whether restoring Myosin II polarity in a non-polarized tissue could rescue
the defects of fold deviation. Although not feasible *in vivo*, we could address this question using *in*
silico modeling, asking whether fold deviation caused by local mechanical perturbations (see Fig3g)
could be rescued by the introduction of planar polarized junctional Myosin II. In order to integrate
Myosin II planar polarity in the model, we first checked if tension pattern could be inferred by myosin
distribution in this tissue. Using laser ablation, we found that circumferential junctions bear more
tension than proximal-distal junctions in the control, while tension appears independent of junction
orientation in Arpc5 knockdown (FigS7a-b), consistent with the respective distributions of Myosin II
observed in these conditions. We then mimicked Myosin II planar polarity and the associated tension
anisotropy by the attribution of different values of junctional tension depending on junction orientation
in our model (see M&M and FigS7c-e). Fold robustness was unaffected by the integration of tension
polarity in the model in the absence of external perturbations (FigS7f), while interestingly, gradual
increase of tissue polarity favors fold straightness and insensitivity to surrounding perturbations
(Fig6g). We further quantified fold morphogenetic robustness in our theoretical model for different
degrees of polarity. Interestingly, increasing polarity decreases the degree of deviation of the fold and
thus decreases the variability in fold directionality (Fig6h-i and FigS7g-h). Altogether, *in vivo*

manipulations and *in silico* modeling indicate that the planar polarization of tissue tension favors
mechanical isolation of fold formation, which ultimately ensures morphogenetic robustness.

Planar polarity favors directional force transmission.

We then asked how planar polarity of Myosin II and the associated polarized tension could
protect morphogenesis from surrounding mechanical disturbances, ensuring robust fold formation.

We first performed laser ablation at the level of a tricellular junction in the fold domain and
followed the recoil of vertices in the neighboring tissue, along several cell diameters (Fig7a). We
observed stronger recoil in the direction of the fold in the control (Fig7b-c). On the contrary, in the
Arpc5 knockdown condition, recoil of vertices up to 3-4 cell diameters becomes similar irrespective of
the direction (Fig7d-e). This experiment reveals the existence of an anisotropic multicellular
mechanical coupling, which results in tissue-scale tension anisotropy.

We hypothesized that Myosin II polarized distribution could generate a bias in long-range
force transmission, avoiding force scattering across the tissue. To specifically test how discrete forces,
generated locally, are transmitted along the tissue, we used wound healing as a way to generate local
forces in a spatio-temporally controlled manner (Fernandez-Gonzalez and Zallen, 2013). We next
analyzed the propagation of forces across the tissue. We wounded the tissue locally by laser ablation at
the level of a tricellular junction, waited for the tissue to repair, and as soon as the healing started
generating local forces, performed a circular cut around the healing region and observed the recoil
(Fig7f-g, + ectopic force, bottom). A similar experiment in the absence of a preliminary wound gave
access to the steady state pattern of tensions within the leg tissue (Fig7f-g, steady state, top row).
Thus, in the absence of wound, the internal release (d*) depends on the tension, or residual stress,
present in the circular isolated tissue fragment, while the external release (d) depends on the tension
present in the rest of the tissue. In the presence of wound-healing traction force, the internal release
dw* mostly depends on both the residual stress present in the circular tissue fragment (Fint) plus the
wound-healing traction force (Fw), while the external release dw mostly depends on both residual
stress in the whole tissue (Fext) plus force transmission (FWtr). This allows us to estimate the wound-
healing traction force by comparing dw* and d*, while we can estimate the transmission of forces
created in response to healing by comparing dw and d.

We first analyzed internal recoil and observed that while the recoil is isotropic in the absence
of Myosin II planar polarity (compare dw^*_{PrDi} and dw^*_{Circ} in Arpc5 RNAi, Fig7h), it is anisotropic in
the presence of Myosin II planar polarity (compare dw^*_{PrDi} and dw^*_{Circ} in control, Fig7h). This
anisotropy suggests that the traction force due to the healing (Fw) could be anisotropic in the control.
Regarding the external recoil, without wound healing, circular cutting induced an ovoid-shaped recoil,

indicating that tissue tension was stronger in the circumferential axis in the wildtype condition, while it results in circular recoil in Arpc5 RNAi condition, indicating that tension anisotropy was lost (compare d_{PrDi} to d_{Circ} in control and d_{PrDi} and d_{Circ} in Arpc5 RNAi in Fig7i). In the presence of wound healing, the tendency is the same (compare d_{WPrDi} and d_{WCirc} in control and d_{WPrDi} and d_{WCirc} in Arpc5 RNAi in Fig7i). Interestingly, while an increase in internal recoil is visible both in proximo-distal and circumferential axis in the presence of wound (compare $d*_{PrDi}$ with $dw*_{PrDi}$ and $d*_{Circ}$ with $dw*_{Circ}$ in the control and Arpc5 RNAi in Fig7h), the external recoil is specifically accentuated along the circumferencial axis in the presence of Myosin II planar polarity (compare d_{Circ} with dw_{Circ} and d_{PrDi} with dw_{PrDi} in control, Fig7i). Thus, even if initial wound healing traction force might be anisotropic, these experiments allow us to propose that in the presence of Myosin II planar polarity, forces (both traction forces and forces transmitted in response) are transmitted preferentially in the circumferential direction (in dark grey in Fig7h-i). On the contrary, in the absence of Myosin II planar polarity, no significant increase in external recoil is observed (compare d_{Circ} and dw_{Circ} as well as d_{PrDi} and dw_{PrDi} in Arpc5 RNAi, Fig7i), showing that forces might spread homogeneously around the wound healing region in this condition (in dark blue in Fig7h-i).

Altogether, these experiments indicate that forces are not homogeneously transmitted in the junctional plane of the tissue but are rather channeled in the presence of Myosin II planar polarity.

Discussion

Our study provides direct evidence of a mechanism protecting morphogenesis from environmental perturbations, ensuring tissue shape robustness on top of developmental patterning through the channeling of mechanical forces. This mechanism involves Arp2/3 complex, which controls junctional tension anisotropy through Myosin II planar polarity, and avoid apical force scattering, thus favoring force transmission along the circumferential axis. This ultimately provides resistance to mechanical perturbations that appear randomly in the surrounding tissue and create a mechanically noisy environment.

Our work reveals that Arp2/3 complex plays a crucial role in establishing a planar polarized distribution of Myosin II in the developing leg tissue. We explored several hypotheses to explain the involvement of Arp2/3 in Myosin II planar polarity. Arp2/3 has been related to E-cadherin endocytosis (Georgiou et al., 2008). By favoring E-cadherin endocytosis at junctions, it could indirectly impact Myosin II polarity. However, adherens junction polarity is not perturbed in Arp2/3 knockdown, ruling out this hypothesis. Myosin II planar polarity in the embryo depends on medial-apical flows of acto-myosin (Rauzi et al., 2010). Consistently, we observed a slight bias in the directionality of medial-apical myosin flow in the control that is lost in Arpc5 knockdown. This indicates that Arp2/3 influences Myosin flow, however, these results are complex and require further investigation. Finally,

Arp2/3, which is involved in the formation of branched actin network, is enriched in the junctions that
are perpendicular to the fold, while F-actin is more abundant in circumferential junctions. This
suggests that the distribution and the density of F-actin networks is itself polarized and that proximo-
distal junctions are enriched in branched actin, while circumferential junctions would be enriched in
linear actin. Since Myosin II has been observed to preferentially accumulate with linear actin network
(Michelot and Drubin, 2011), this polarity of the actin networks would favor the accumulation of
Myosin II along circumferential junctions and thus drive Myosin II planar polarity.

This work also highlights that Myosin II planar polarity could protect a morphogenetic process
from the surrounding noise, by playing a role in long-range force chanelling and forming a sort of
mechanical fence. Myosin II planar polarity has been described in different contexts: germ band
extension in Drosophila embryo where it drives tissue elongation (Bertet et al., 2004), but also neural
tube closure in chicken (Nishimura et al., 2012), where it favors polarized junction shortening and
tissue folding, or at compartment boundaries in Drosophila and zebrafish where it maintains a straight
borders (Calzolari et al., 2014; Major and Irvine, 2006; Monier et al., 2010). However, its role in the
protection of morphogenesis from external perturbations was unexpected. Indeed, so far, most of the
studies related to morphogenetic robustness converge on the regulation of gene expression patterns in
response to morphogen gradients and consider acto-myosin either as a component of the "core
toolbox" responsible for building a new shape or as a mechanism to buffer local heterogeneity in
positional information by exerting a feedback on cell-fate decisions (Gilmour et al., 2017). However,
the influence of tissue mechanics in maintaining low variability of a specific shape when challenged
by intrinsic perturbations in close proximity was unexplored.

Interestingly, two recent papers addressed the robustness of tissue invagination. While the
work of Yevick et al highlights the importance of mechanical redundancy to resist to accidental
damage (Yevick et al., 2019), Eritano et al reveal the role of mechanical coupling as an intrinsic
property of morphogenesis to buffer small variability in gene expression patterns (Eritano et al., 2020).
The present study appears complementary to these previous works, showing how morphogenesis is
naturally protected from mechanical perturbations occurring randomly in the surroundings, by creating
a fence through Arp2/3-dependent junctional myosin II planar polarity. It further reveals that tissue
mechanics not only buffer genetic information but can take over the positional information given by
patterning genes since folds can considerably deviate from the pre-established program if forces are
not properly channeled.

Finally, this work further reveals that fold morphogenesis in the Drosophila developing leg occurs
in the presence of mechanical noise, as shown by mechanical perturbations randomly distributed in the
developing tissue at the time of fold formation. Since the occurrence of several developmental
processes in the same time window is frequently observed during development, it is tempting to

speculate that mechanical noise could be a general feature of morphogenesis and that mechanical isolation could be required in a wide variety of morphogenetic processes to avoid force scattering and maintain morphogenetic robustness. Thus, this process of force channeling through Myosin II polarized distribution could be a general way to isolate a particular morphogenetic process from surrounding events, preventing any interference between closely located events and favoring robustness.

Acknowledgements

We thank Michel Labouesse and Yohanns Bellaiche for their constructive comments on the manuscript. MS's lab is supported by grants from the European Research Council (ERC) under the European Union Horizon 2020 research and innovation program (grant number EPAF: 648001), from the Institut National de la Santé et de la Recherche Médicale (Inserm, Plan cancer 2014-2019) and from the association Toulouse Cancer Santé (TCS, ApoMacImaging: 171441). EM has a post doc fellowship from the Association pour la Recherche contre le Cancer (ARC). ST has a CIFRE fellowship from the ANRT.

Author contributions

EM conceived and performed the experiments in fly leg discs. ST participated in modeling conception and realized all the simulations. EM and ST analyzed and quantified the data. BM initiated the project, participated in experimental conception and article writing. GG supervised the modeling conception. MS supervised the project, wrote the paper and provided the funding. CR helped with the analysis of Rpr and Dpn pattern, and PIV analysis.

Competing interest declaration

The authors declare no competing financial interests.

Corresponding authors

Correspondence and requests should be addressed to M.S. (magali.suzanne@univ-tlse3.fr) for biology and materials and to G.G. (guillaume@morphogenie.fr) for modeling.

Main figure titles and legends

Figure 1: Fold morphogenetic robustness is lost in Arpc5 knockdown (related to FigS1).

a-b, 3D reconstructions of a *control (Dll-Gal4; arm-GFP)* (**a**) or *Arpc5* RNAi *(Dll-Gal4; arm-GFP, UAS Arpc5RNAi)*
(**b**) pupal leg discs at 1h30 after puparium formation (APF), showing tarsal folds morphology in blue. **c,** Dot
plots showing the maximal distance between the real folds and the corresponding predicted fold (highlighted
in yellow) in *control* and *Arpc5* RNAi leg discs (n=35, 63, respectively). **d,** Dot plots showing the angle of the
folds relative to the proximal-distal axis in *control* and *Arpc5* RNAi discs (n=35 in both cases). **e,** Dot plots
quantifying the fold parallelism (ratio between the angles of two different folds) in *control* and *Arpc5* RNAi leg
discs (n=35 in both cases). In **c-e**, a *F*-test of equality of variances was used. ns, not significant; ***, p-value <
0.001. Black lines represent the mean ± SD. Genotypes correspond to *sqh-RFPt[3B]; Dll-Gal4, UAS-GC3Ai*
uncrossed or crossed to *UAS Arpc5RNAi*.

Figure 2: Developmental patterning is unaffected in Arpc5 knockdown (Related to FigS2 and Movie S1-S3)

a, Z-sections of *control (sqh-GFP[29B]; Dll-Gal4)* and *Arpc5* RNAi *(sqh-GFP[29B]; Dll-Gal4; UAS-Arpc5RNAi)* leg
discs stained with phalloidin (white) and showing Z-projection of anti-Deadpan (magenta). **b,** Z sections of
control (rpr-lacZ; Dll-Gal4) or *Arpc5* RNAi *(rpr-lacZ; Dll-Gal4; UAS-Arpc5RNAi)* leg discs stained with phalloidin
(white) and showing a Z-projection of rpr-lacZ (yellow). **a,b,** Yellow, blue and green arrowheads respectively
indicate predicted fold, real fold and perfect match between them. Please note that both Dpn and rpr-lacZ are
also expressed in some neurons, indicated by asterisks. Cartoons on the right recapitulate the location of Notch
signaling (purple) and pro-apoptotic genes expression (yellow) in *control* and *Arpc5* RNAi conditions. **c-d,** Dot
plots showing the angle of the stripes of expression of *Deadpan* (c) or *reaper* (d) in the folds relative to the
proximal-distal axis in *control (rpr-lacZ; Dll-Gal4)* and *Arpc5* RNAi *(rpr-lacZ; Dll-Gal4; UAS-Arpc5RNAi)* leg discs
(n=13 and 11 respectively). A *F*-test of equality of variances was used. ns, not significant. Black lines represent
the mean ± SD. **e-f,** 3D-reconstructions of *control* (e) and *Arpc5* RNAi (f) leg discs showing the expression of
Deadpan (magenta, arrows). Yellow and blue domain respectively highlight the predicted and the real fold.
Asterisks point out *Dpn*-expressing neurons. **g,** Average number of dying cells inside or outside the predicted
distal-most tarsal fold in *control* and *Arpc5* RNAi leg discs (n=20 and 28, respectively). Bar graphs indicate the
mean ± SEM. Cartoons on the left recapitulate the regions inside (yellow) and outside (brown) of the predicted
fold. **h,i,** Sagittal views showing apoptotic myosin cables (h) and dot plots showing the initial velocity of
apoptotic cable recoil after laser microdissection (i) in *control (sqh-GFP[29B]; Dll-Gal4)* and *Arpc5* RNAi *(sqh-*
GFP[29B]; Dll-Gal4; UAS-Arpc5RNAi) leg discs (n=6 and 10, respectively). Black line indicates the median. In **g**
and **i**, the statistical significance was calculated using Mann-Whitney U test. ns, not significant. Scale bars are
20 µm in **a**, **b**, **e** and **f**; 2 µm in **h**.

Figure 3: Leg morphogenesis occurs in a mechanically noisy environment (Related to FigS3, FigS4 and Movie
S4).

a, Fold morphology in *control* or *Arpc5* RNAi arm-GFP pupal leg discs and corresponding schemes. **b,d,** Confocal
images showing the 'Notch border' (red arrowheads) (**b**) or the non-apoptotic apico-basal myosin structure
(black arrowheads) (**d**) in *control (sqh-GFP[29B]; Dll-Gal4)* and *Arpc5* RNAi *(sqh-GFP[29B]; Dll-Gal4; UAS-*
Arpc5RNAi) conditions. **c,** Dot plots of the initial recoil velocity observed after microdissection of adherens
junctions at the 'Notch border' or adjacent junctions in *control (sqh-RFPt[3B]; Dll-Gal4; arm-GFP)* and *Arpc5*
RNAi *(sqh-RFPt[3B]; Dll-Gal4; arm-GFP, UAS-Arpc5RNAi)* leg discs (n=11, 12, 10 and 12 respectively). **e,** Dot
plots of the initial recoil velocity of non-apoptotic apico-basal myosin structure or lateral membrane after laser
dissection in *control (sqh-GFP[29B]; Dll-Gal4)* and *Arpc5* RNAi *(sqh-GFP[29B]; Dll-Gal4; UAS-Arpc5RNAi)* leg discs
(n=14, 12, 10 and 10 respectively). In **c**, **e** black lines indicate the median. Statistical significance has been
calculated using Mann-Whitney U test. ns, not significant; *, p-value < 0.05; **, p-value < 0.01; ***, p-value <
0.001. **f,** Rolled-out maps of the fold domain indicating the locations of mechanical perturbations observed in

close vicinity of the predicted fold (yellow) in *control* and *Arpc5* RNAi leg discs (n=14 and 14, same genotypes as
d). **g,** *In silico* simulations at the maximal fold depth without (left) or with (right) mechanical perturbation
(challenged cells are indicated in black). **h,** Rolled-out map of mechanical perturbations random locations (grey)
from 25/55 *in silico* simulations. 3 perturbations were integrated for each simulation. **i,** Dot plots showing the
maximal distance between the real fold and the predicted fold (highlighted in yellow) in *in silico* simulations
without (left) or with (right) mechanical perturbations. **j,** Dot plots showing the angle of the fold relative to the
proximal-distal axis in *in silico* simulations without or with mechanical perturbations. In **a**, **b**, **d**, **g**, predicted fold
is highlighted in yellow, real fold in blue and perfect match between them in green. In i and j, a Levene's test
was used. ***, p-value < 0.001. Scale bar represents 20 µm in a, b, d.

Figure 4: The leg epithelial tissue exhibits myosin planar polarity. (Related to FigS5)

a-d, Close up views of confocal images showing the distribution of β-catenin-GFP and Myosin II (*sqh-RFPt[3B]*)
in *control* and *Arpc5* RNAi leg discs at the onset of fold formation (a,b) or at mid-fold stage (c,d) in the fold
domain (a,c) or in the adjacent domain (b,d); **a'-d'**, Quantification of the Myosin II distribution at junctions. **a"-**
d", Quantification of the cell shape anisotropy. In **a-d**, red and blue arrowheads indicate proximal-distal and
circumferential junctions, respectively. In **a'-d'**, graph bars correspond to the mean ± SEM. In **a"-d"** black lines
indicate the median. Statistical significance has been calculated using Mann-Whitney U test. ns, not significant;
***, p-value < 0.001. n= 752 and n= 590 junctions in a'; n= 153 and n= 197 cells in a"; n= 603 and n= 645
junctions in b'; n= 209 and n= 254 cells in b"; n= 1107 and n= 1223 junctions in c'; n= 301 and n= 364 cells in c";
n= 1087 and n= 980 junctions in d'; n= 381 and n= 384 cells in d".

Figure 5: Polarized Arp2/3 drives polarized F-actin distribution in the leg epithelial tissue (Related to FigS5)

a-b, Confocal images (**a**) showing the distribution of actin in *control* (*Dll-Gal4; arm-GFP*) or *Arpc5* RNAi *(Dll-*
Gal4; arm-GFP, UAS-Arpc5RNAi) leg discs in the fold domain and quantification (**b**) in both the fold domain
(left; n= 1004 and n= 981 junctions respectively) and the adjacent domain (right; n= 823 and n= 672 junctions
respectively). **c-d,** Confocal images (**c**) showing the distribution of Arp3 (*ap-Gal4, UAS α-catenin-TagRFP, UAS*
Arp3-GFP) and quantification (**d**) in the fold domain and in the adjacent domain (n= 582 and n= 432 junctions
respectively).

Figure 6: Planar polarity is required for morphogenetic robustness (Related to FigS6 and FigS7).

a, Confocal images showing the distribution of E-cadherin, α-catenin and Myosin II in *control* (*sqh-GFP[29B];*
tubG80ts; ap-Gal4, UAS α-catenin-TagRFP) or *AJs GFP-trap* (*sqh-GFP[29B]; tubG80ts; ap-Gal4, UAS vhhGFP4-α-*
catenin-TagRFP) leg discs. **b,** Quantification of Myosin II distribution at junctions in leg discs in both conditions
(n= 1009 and n= 564 junctions respectively). **c,** Confocal images showing fold morphology in *control* or *AJs GFP-*
trap leg discs (deviated fold represents 0/15 and 8/12, respectively). Predicted fold domain is highlighted in
yellow, real fold in blue and perfect match between them in green. Scale bar represents 20 µm. **d,** Dot plots
showing the relative maximal distance between the real fold and the predicted fold (highlighted in yellow) in
control or *AJs GFP-trap* leg discs (n=15 and 12 legs, respectively). **e,** Dot plots showing the angle of T3-T4 and
T4-T5 folds relative to the proximal-distal axis in *control* or *AJs GFP-trap* leg discs (n=15 and 12 legs,
respectively). **f,** Dot plots showing the fold parallelism in *control* or *AJs GFP-trap* leg discs (n=15 and 12 legs,
respectively). in **d-f**, A *F*-test of equality of variances has been used to compare the phenotypic variances. ***,
p-value < 0.001. **g,** *In silico* simulations including mechanical perturbations (challenging cells are shown in
black) for different values of junction weight ratio (i.e. tension anisotropy). Predicted fold domain is highlighted
in yellow, real fold in blue and perfect match between them in green. **h,** Dot plots showing the maximal
distance between the real fold and the predicted fold (highlighted in yellow) for different values of junction
weight ratio (polarity). (n= 55 simulations in each cases). **i,** Dot plots showing the angle of the fold relative to
the proximal-distal axis for different values of junction weight ratio (polarity). (n= 55 simulations in each cases).

In **h** and **i**, a Levene's test has been used to compare the variance of phenotypes. *, p-value < 0.05; **, p-value
< 0.01.

Figure 7. Planar polarity favors force channeling.

a, Schematics showing the impact of the wound (left), the forces in presence (F), and the maximum
displacement of vertices at various cell diameters (d). **b,d**, Z-projections of *control* (*Dll-Gal4; arm-GFP*) and
Arpc5 RNAi (*Dll-Gal4; arm-GFP, UAS-Arpc5RNAi*) leg discs before (left) and after (middle) the cut. Overlay are
shown on the right. Red circles indicate site of ablation. Scale bar represents 2 μm. **c,e,** Average displacement
of vertices (represented by vectors) at different distances and angles from the cut site (center) in *control* (**b**)
and *Arpc5* RNAi (**c**) (n= 8, 10 legs respectively). Distance from the cut is color coded as in a. **f**, Schematics
indicating the absence (top) or the presence (bottom) of wound and the forces in presence at the level of the
circular cut (left) and the external and internal maximal recoils after circular ablation (right). In the absence of
wound (top) the internal release d* depends on the residual stress present in the circular isolated tissue
fragment, while the external release d depends on the tension present in the rest of the tissue. In the presence
of wound-healing traction force (bottom), the internal release dw* depends on both the residual stress present
in the circular tissue fragment (Fint) plus the wound-healing traction force (Fw), while the external release dw
depends on both residual stress in the whole tissue plus the force transmitted in response to the wound-
healing traction force (Fwtr). **g**, Confocal images of the *control* and *Arpc5* RNAi leg discs (same genotypes as in
b). External and internal maximal recoils of the tissue are indicated by the green and purple lines respectively,
while the initial positioning of the circular ablation is indicated by a dashed red line. Scale bar represents 5 μm.
h-i, Quantifications of internal (h) and external (i) maximal recoils of the tissue in the absence (light grey and
light blue) or the presence (dark grey and dark blue) of wound in *control* and *Arpc5* RNAi leg discs (n= 10, 10,
12, 10 respectively). Statistical significance has been calculated using Mann-Whitney U test. ns, not significant;
*, p-value < 0.05; **, p-value < 0.01; *** p-value < 0.001. Graph bars correspond to the mean ± SEM.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rat anti-deadpan antibody (1:100)	Abcam	Cat#ab195172
Rabbit anti-cleaved dcp1 (1:200)	Cell Signaling Technology	Cat#9578; RRID: AB_2721060
Rat anti-E-cadherin(1:50)	DSHB	DCAD2; RRID: AB_528120
chicken β-galactosidase (1:1000)	GeneTex	Cat# GTX77365, RRID:AB_379834
Goat anti-rat IgG 647	Thermo Fisher Scientific	Cat# A-21247,RRID:AB_141778
Goat anti-rat IgG 555	Thermo Fisher Scientific	Cat# A-21434 RRID:AB_2535855
Goat anti-rabbit IgG 555	Thermo Fisher Scientific	Cat# A-21428 RRID:AB_2535849
Goat anti-chicken IgG 488	Thermo Fisher Scientific	Cat# A-11039, RRID:AB_2534096
Bacterial and Virus Strains		
Chemicals, Peptides, and Recombinant Proteins		
Phalloidin-Rhodamine	Thermo Fisher Scientific	Cat#R415 RRID:AB_2572408
Phalloidin-Alexa488	Thermo Fisher Scientific	Cat# A12379
Vectashield	Vector Laboratories	Cat#H-1000
TRIzol Reagent	ThermoFisher Scientific	Cat#15596026
20-hydroxyecdysone	Sigma-Aldrich	Cat# H5142
Critical Commercial Assays		
Luna Script reverse transcriptase SuperMix kit	NEB	Cat#E3010S
Luna Universal qPCR Master Mix	NEB	Cat#M3003S
Experimental Models: Organisms/Strains		
D. melanogaster: w; Dll-Gal4[em212]	Gift from G. Morata M. Calleja et al, 1996	FBti0007187

D. melanogaster: ap-Gal4: y[1] w[1118]; P{w[+mW.hs]=GawB}ap[md544]/CyO	BDSC	RRID: BDSC_3041
D. melanogaster: hh-Gal4; UAS-GFP	Gift from C. Benassayag	Centre de Biologie Intégrative (CBI), LBCMCP
D. melanogaster: rpr-4kb-lacZ	Gift from Gines Morata (stock referenced in Flybase)	Centro de Biologia Molecular Severo Ochoa (CBM SO), Madrid
D. melanogaster: w; ; ap[42B11]-Gal4	Gift from C. Estella	Centro de Biologia Molecular Severo Ochoa (CBM SO), Madrid
D. melanogaster: tubG80ts: w[*]; P{w[+mC]=tubP-GAL80[ts]}20; TM2/TM6B, Tb[1]	BDSC	RRID: BDSC_7019
D. melanogaster: arm-GFP: w[*]; P{w[+mW.hs]=arm-GFP.P}57	BDSC	RRID: BDSC_8556
D. melanogaster: Arp3-GFP: w[*]; P{w[+mC]=UASp-Arp3.GFP}2	BDSC	RRID: BDSC_39723
D. melanogaster: Myosin-GFP: w, sqh{TI}-eGFP [29B]	Ambrosini et al, 2019	Centre de Biologie Intégrative (CBI), LBCMCP
D. melanogaster: Myosin-RFP: w, sqh{TI}-TagRFPt [3B]	Ambrosini et al, 2019	Centre de Biologie Intégrative (CBI), LBCMCP
D. melanogaster: ECad-GFP: w ; shg{TI}-eGFP	BDSC	RRID:BDSC_60584
D. melanogaster: w; UAS-alpha-Catenin-TagRFP	Gift from K. Sugimura Ishihara and Sugimura, 2012	FBal0279996 Kyoto University, iCeMS
D. melanogaster: cytoplasmic GFP apoptosensor: *w; UAS-GC3Ai [G16]*	Schott et al, 2017	Centre de Biologie Intégrative (CBI), LBCMCP
D. melanogaster: AJ GFP-trap: w; UASt-vhhGFP4-alpha-Catenin-TagRFP	This study	N/A
D. melanogaster: RNAi of Arpc3A: y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02370}attP2	BDSC	RRID:BDSC_27044
D. melanogaster: RNAi of Arpc5: y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03147}attP2	BDSC	RRID:BDSC_28720
D. melanogaster: RNAi of Arp2: y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02785}attP2/TM3, Sb[1]	BDSC	RRID:BDSC_27705
D. melanogaster: LacZ: w[*]; P{w[+mC]=UAS-lacZ.Exel}2	BDSC	RRID:BDSC_8529
Oligonucleotides		

Primer vhhGFP4-F_InF: GGCCGCGGCTCGAGGATGGTCCAACTGGTGGAGTCTG	This study	N/A
Primer vhhGFP4-R+linker: AAAGATCCTCTAGAGGTACCATGAACTCGCCGCTGCC GGCGGCGCTGCCGGCGCTGCCGCTGGAGACGGTGAC CT	This study	N/A
Primer a-catTagRFP-F_InF#2: AGCGGCGAGTTCATgTTAAAACCTGATAAAATGGGCA CG	This study	N/A
Primer a-catTagRFP-R_InFusion: AAAGATCCTCTAGAGTCAATTAAGTTTGTGCCCCAGT	This study	N/A
Arpc5_FW: GAGGATGATGGGGTGGAGAG	This study	N/A
Arpc5_REV: GTCCATTTGCGTCGACTTGA	This study	N/A
RP49_FW: GACGCTTCAAGGGACAGTATCTG	This study	N/A
RP49_REV: AAACGCGGTTCTGCATGAG	This study	N/A
GAPDH_FW: CGTTCATGCCACCACCGCTA	This study	N/A
GAPDH _REV: CCACGTCCATCACGCCACAA	This study	N/A
Recombinant DNA		
pUAS-Dαcatenin-TagRFP plasmid	Gift from K. Sugimura Ishihara and Sugimura, 2012	Kyoto University, iCeMS
nanoGFP-cry2 plasmid	Gift from R. Le Borgne	IGDR, Rennes
pUASt-attB	DGRC	Cat#1419
Software and Algorithms		
GraphPad Prism 8	GraphPad Software	RRID: SCR_002798
Adobe Illustrator CS5	Adobe	RRID:SCR_010279
Imaris 8.4.1	Bitplane	RRID: SCR_007370
Fiji	https://fiji.sc/	RRID: SCR_002285
ZEN Blue	ZEISS	RRID:SCR_013672
Black Zen software	ZEISS	RRID:SCR_018163
ROE Syscon	Rapp OptoElectronic	N/A
CFX Manager 3.1	Bio-Rad	RRID:SCR_017251
Python 3.7	http://www.python.org/	RRID:SCR_008394
MatPlotLib	http://matplotlib.sourceforge.net	RRID:SCR_008624

Vertex model tissue v0.7.1	https://zenodo.org/badge/latestdoi/32533164	N/A
OpenPIV v0.23.1	https://doi.org/10.5281/zenodo.3930343	N/A
Other		
Schneider's insect medium	Sigma-Aldrich	Cat#S0146
Halocarbon oil	Sigma-Aldrich	Cat#H8773
120 μm deep Secure-Seal™	Sigma-Aldrich	Cat#GBL654008

Contact for Reagent and Resource Sharing

536 Further information and requests for resources and reagents should be directed to and will be

537 fulfilled by the Lead Contact, Magali Suzanne (magali.suzanne@univ-tlse3.fr).

Experimental model and subject details

Experimental Animals

The animal model used here is Drosophila melanogaster, in a context of in vivo/ex vivo

experiments. In order to respect ethic principles, animals were anesthetized with CO2 (adults)

before any manipulation. To avoid any release of flies outside the laboratory, dead flies were

frozen before throwing them. Stocks of living flies were conserved in incubators, either at 18

or 25 degrees to maintain the flies in optimal condition. Genotypes and developmental stages

are indicated below. Experiments were performed in both males and females indifferently.

Loss of function experiments using RNAi were carried out at 30 degrees.

Drosophila melanogaster

ap^{md544}-Gal4, arm-arm-GFP, tubG80ts, UAS-Arp3-GFP and E-Cad-GFP knock-in were

obtained from Bloomington Drosophila Stock Center (BDSC). sqh-eGFP^{KI}[29B], sqh-

RFPt^{KI}[3B] and the GFP apoptosensor UAS-GC3Ai were described previously (Ambrosini et

al., 2019; Schott et al., 2017). hh-Gal4; UAS-GFP is a gift from C. Benassayag. Dll^{EM212}-Gal4

and rpr-4kb-lacZ are gifts from G. Morata. ap^{42B11}-Gal4 and UAS-α-catenin-TagRFP are

gifts from C. Estella and K. Sugimura.

RNAi experiments were realized using UAS-Arpc3A-RNAi (JF02370), UAS-Arpc5-RNAi

(JF03147), UAS-Arp2 RNAi (JF02785), UAS-lacZ, obtained from BDSC.

Method details

AJ GFP-trap construct

To trap Myosin-GFP at junctions, we constructed a p*UASt–vhhGFP4-α-catenin-TagRFP* (AJs

GFP-trap). To do so, *α-catenin-TagRFP* was amplified by PCR with specific primers from a

pUAS-Dαcatenin-TagRFP (kindly provided by Dr K. Sugimura). vhhGFP4 was amplified by

PCR from nanoGFP-cry2 plasmid (gift from Dr Le Borgne). A 36nt sequence was added in

the reverse primer to create a GSAGSAAGSGEF linker between the GFP-trap sequence and

α-catenin-TagRFP sequence. These two fragments were successively cloned using InFusion

technology in pUASt-attB plasmid cut with KpnI.

The resulting p*UASt–vhhGFP4-α-catenin-TagRFP* plasmid injection was performed by the

CBMSO Drosophila Transgenesis Service (Madrid, Spain) using flies carrying attP2 landing

sites on the third chromosome to produce transgenic flies.

RNA and RT-qPCR analyses

RNAi efficiency was assessed by mRNA phenol/chloroform extraction and reverse

transcription was done using the Luna Script reverse transcriptase SuperMix kit (NEB –

E3010S). cDNAs were quantified by qPCR using the Luna Universal qPCR Master Mix

(NEB – M3003S) (primer sequences: Arpc5_FW GAGGATGATGGGGTGGAGAG and

Arpc5_REV GTCCATTTGCGTCGACTTGA) and normalized against RP49 and GAPDH

cDNA levels (primer sequences: RP49_FW GACGCTTCAAGGGACAGTATCTG ;

RP49_REV AAACGCGGTTCTGCATGAG and GAPDH_FW

CGTTCATGCCACCACCGCTA ; GAPDH _REV CCACGTCCATCACGCCACAA). The

relative normalized cDNA expression levels were calculated using the DeltaDeltaCt method

from Bio-Rad CFX Manager 3.1 software.

Immunofluorescence

Imaginal leg discs were dissected after 48h at 29°C at prepupal stage (0, 1.5 or 2.5 hours after

pupae formation – APF) in PBS 1X. Imaginal discs were fixed 20' in paraformaldehyde 4%

diluted in PBS 1X, then washed in PBS 1X and either mounted in Vectashield (Vectors

laboratories) or extensively washed in PBS-Triton 0.3%-BSA 1% (BBT) and incubated

overnight at 4°C with appropriate dilutions of primary antibodies in BBT. Rat anti-deadpan

antibody (Abcam – ab195172) was used at 1:100 dilution, rabbit anti-dcp1 (Cell Signaling

Technology – 9578S) at 1:200, rat anti-E-cadherin antibody (Developmental Studies

Hybridoma Bank – DCAD2) at 1:50 and chicken β-galactosidase (GeneTex – GTX77365) at

1:1000. After washes in BBT, imaginal leg discs were incubated at room temperature for 2 h

with 1:200 anti-rat IgG 647, anti-rat IgG 555, anti-rabbit IgG 555 or anti-chicken IgG 488

(obtained from Interchim) with or without phalloidin (Alexa488 at 1:500 or rhodamin at

1:500 – Fisher Scientific). Then, samples were washed in PBS-Triton 0.3%, suspended in

Vectashield (Vectors laboratories) and mounted on slides.

Ex vivo culture of leg imaginal disc

Imaginal leg discs were dissected from prepupal stage (1.5 hours APF at 29°C) in Schneider's

insect medium (Sigma-Aldrich) supplemented with 15 % fetal calf serum and 0.5 %

penicillin-streptomycin as well as 20-hydroxyecdysone at 2 µg/mL (Sigma-Aldrich, H5142).

Leg discs were transferred on a slide in 12 µL of this medium in a well formed by a 120 µm-

deep double-sided adhesive spacer (Secure-SealTM from Sigma-Aldrich). A coverslip was

then placed on top of the spacer. Halocarbon oil was added on the sides of the spacer to

prevent dehydration.

Confocal imaging

Samples were analyzed using a LSM880 confocal microscope fitted with a Fast Airyscan

module (Carl Zeiss) and equipped with a Plan-Apochromat 40x/NA 1.3 Oil DIC UV-IR M27

objective. Z-stacks were acquired using either the laser scanning confocal mode or the High

Resolution mode (Airyscan) with a pixel size of 0.046 µm/pixel and a *z*-step of 0.220 um.

Airyscan Z-stacks were processed in ZEN software using the automatic strength (6 by

default) and the 3D method.

Laser ablation

Laser ablation experiments were performed using a pulsed DPSS laser (532 nm, pulse length

1.5 ns, repetition rate up to 1 kHz, 3.5 µJ/pulse) steered by a galvanometer-based laser

scanning device (DPSS-532 and UGA-42, from Rapp OptoElectronic, Hamburg, Germany)

and mounted on a LSM880 confocal microscope (Carl Zeiss) equipped with a 63x C-

Apochromat NA 1.2 Water Corr objective (Carl Zeiss). Photo-ablation of apical junction was

done in the focal plane by illuminating at 70 % laser power during 1 s. This focal plane was

acquired every 0.551 s, during 10 s before and at least 45 s after ablation, with a pixel size of

0.13 µm/pixel. Photo-ablation of apico-basal Myosin II enrichment was done in the focal

plane by illuminating at 100 % laser power during 2-2.5 s along a 45-50 pixels line. This focal

plane was acquired every 0.551 s, during 5 s before and at least 45 s after ablation, with a

pixel size of 0.13 µm/pixel. Data analysis was performed with the ImageJ software using a
homemade macro.

For experiments on tissue scale tension anisotropy (Fig7a-e), a Z-stack of 7 slices was
acquired every 1.774s during 10s before and 3 minutes after ablation, with a pixel size of 0.13
µm/pixel. Photo-ablation of a vertex was done during the stack acquisition by illuminating at
75 % laser power during 2 s in a 6 pixels radius circle. Data analysis was performed with the
ImageJ software by measuring the displacement (distance and orientation) reached at the
maximal recoil of all vertices in a 10 µm radius circle around the ablated vertex, from their
initial location (before cut), using Manual tracking plugin.

For circular photo-ablation, the focal plane was illuminated during 4-5 s along a 45 pixels
radius circle (Fig7f-i). This focal plane was acquired every 0.551 s, during 10 s before and at
least 45 s after ablation, with a pixel size of 0.13 µm/pixel. For the second set of experiments
(wound healing), tissue was first wounded at the level of a tricellular junction by illuminating
during 4-5 s along a 10 pixels radius circle and then cut circularly during the healing phase
using the circular shape described above (Fig7f-i). Data analysis was performed with the
ImageJ software. Briefly, the distance between the location of the circular cut and the
maximal recoil induced by this ablation was measured at 0° (proximal-distal axis) and 90°
(circular axis) using the line tool.

The Vertex Model

Initial tissue generation

We modeled the most distal part of the leg imaginal disc (Fig1a), before the T4-T5 fold
formation begins, as a 2D meshwork around a cylinder capped by two hemispheres, oriented
along the proximal-distal axis. We started with a mostly hexagonal lattice with a perimeter of
23 cells and a length of 15 cells. We perform two rounds of cell divisions with a random
division axis to randomize cell side number and create a meshwork of approximately 50 by 30
cells. Diversity of cell shapes is increased by adding variability to the cells' preferred areas,
normally distributed with an 8% variance. The initial tissue has 1652 cells, is 200 µm long in
its proximal-distal axis and 100 µm in diameter.

Mechanical model

The epithelium shape is given by the quasi-static equilibrium of a potential energy dependent
on the junctional mesh geometry, following our previous work (Gracia et al., 2019; Monier et
al., 2015).

This energy is given in equation (1) and is comprised of three cell-level terms and two

constrain terms. At the cell level, apical shape is governed by area and perimeter elasticity

terms, following (Bi et al., 2015). To these terms, we add a linear apico-basal tension for the

apoptotic cells, dependent on cell height. Two more terms ensure maintenance of the overall

tissue shape. First, the total volume of the tissue is maintained by an elastic constrain. Second,

an external barrier is modeled as a sphere surrounding the tissue; when the distance of a

vertex from the sphere center is higher than the sphere radius, it is pulled back to this radius

by an elastic force.

$$E = \sum_{\alpha} \left(\frac{K_\alpha}{2} \left(A_\alpha - A_{\alpha,0} \right)^2 + \frac{K_p}{2} (P - P_{\alpha,0})^2 + T_\alpha^{\mathrm{ab}} h_\alpha \right) + \frac{K_V}{2} (V - V_0)^2 + \sum_i \frac{K_b}{2} \delta^+ R_i^2 \quad (1)$$

The value of (KV) was chosen at the lowest value such that compression of the tissue by

cellular contractility and the effect of the lumen on the fold formation was minimal and kept

the tissue integrity (see FigS6). Here $\sum_\alpha$ marks the sum over every polygonal cell and $\sum_i$

a sum over every vertex. Apico-basal tension is exerted from the vertices towards the

proximal-distal axis (center of the cylinder). We consider an anchor point i' as the protection

of the vertex i onto the proximal-distal axis. i' is rigidly fixed to this axis. The penetration

depth $\delta^+ R_i$ is defined by:

$$\delta^+ R_i = \begin{cases} R_i - R_p & \text{if } R_i > R_b \\ 0 & \text{if } R_i \leq R_b \end{cases} \quad (2)$$

Energy minimum is reached through a gradient descent strategy using the Broyden-Fletcher-

Goldfarb-Shanno bound constrained minimization algorithm from the scipy library (van der

Walt et al., 2011).

Tissue anisotropy

In vivo, cell polarity translates in different mechanical properties for different cell junctions.

Here, in order to create the cell anisotropy (the ratio between the long axis and the

perpendicular axis of the apical surface of the cell), we added a weight on each cell junction

in the calculation of cell perimeter. The modified perimeter is calculated as the weighted sum

$$P^* = \frac{\sum_{ij \in \alpha} \omega_{ij} l_{ij}}{\sum_{ij \in \alpha} \omega_{ij}} n_\alpha \quad (3)$$

Where $\sum_{ij\in\alpha}$ denotes a sum over all edges of cell α and ω_{ij} is the weight of the junction ij,

l_{ij} is the length of the junction ij and n_α is the number of sides of cell α. Note that this

weighted sum is equal to the actual perimeter when all weights are equals, and allows to

model polarity without any modification of the other dynamical parameters.

We set two different values of the weight depending on the orientation of the edge. Weight is

higher for circumferential edges ($\omega_{//}$) than for proximal-distal ones ($\omega_\perp$). (FigS7c-h). As cell

shapes are on average hexagonal, we set the boundary between parallel and perpendicular

junctions at $\pi/3$.

Apoptotic process

Around 30 apoptotic cells are chosen randomly in the fold region according to a probability

density described in our previous work (Monier et al., 2015). The tissue deforms

progressively as apoptotic cells undergo apical constriction and apico-basal traction through

gradual changes in their mechanical parameters, while other cells passively follow the

deformation.

A cell starts apoptosis at time step t_i such that $t_i = t_f \frac{\phi_\alpha}{\pi}$ where ϕ_α is the angle of the cell's

apical surface face center with respect to the dorso-ventral axis and t_f the final simulation

time.

Apoptosis is modeled as a sequence of apical constriction and apico-basal traction.

Apical constriction consists in a reduction of the cell's preferred perimeter $P_{\alpha,0}$ at a constant

rate $\sqrt{\tau_\alpha}$ and a reduction of its preferred area $A_{\alpha,0}$ at constant rate τ_α until the cell area

reaches a threshold A_c as given in equation (4). Once the critical area is reached, preferred

area and preferred perimeters are maintained constant.

$$\begin{cases} P_{\alpha,0}(t) = \dfrac{P_{(\alpha,0)}(t-1)}{\sqrt{\tau_\alpha}} \\[2em] A_{\alpha,0}(t) = \dfrac{A_{(\alpha,0)}(t-1)}{\tau_\alpha} \end{cases} \qquad if\ A_{\alpha,t} > A_c \qquad\qquad (4)$$

700   During the apical constriction phase, contraction is propagated to neighboring cells.

701   Contraction rate τ_β of a neighboring cell β decreases linearly as the cell is farther away from

702   the apoptotic cell:

$$\tau_\beta = \tau_\alpha - \frac{r}{r_{\max}}(\tau_\alpha - \tau_{\min}) \qquad (5)$$

Here $r = 1$ if the cell β is a direct neighbour of the apoptotic cell, 2 if it's a second order

neighbor, and so on. $r_{\max}$ is the span of the propagation and τ_{min} the contraction rate for cells

at r_{max} from the apoptic cell.

Each apoptotic cell can develop an apico-basal tension, during and after the constriction

phase, with probability $p = \exp(-A_\alpha/A_c)$. Traction takes place for N_t time steps, during

which the apico-basal tension exerted on the face is T^{ab}

Mechanical perturbation

We modeled mechanical perturbations during fold formation as three "disrupting cells"

placed randomly at a maximal distance of 40 µm (7 cells distance) from the middle of the

predicted fold position. The "disrupting cell" dynamics is similar to apoptosis, with apical

constriction and apical-basal tension, however, apical constriction starts at the beginning of

the simulation, and when the critical area is reached, apico-basal traction is applied until the

end of the simulation. For every simulation, apart from the time span for which the

mechanical perturbation is exerted, the parameters for the perturbing cell are identical to the

parameters for an apoptotic cell.

Choice of parameter values.

The unit energy (denoted by u) is defined so that the area elasticity modulus K_α equals

1 $u/\mu m^4$. To model lumen incompressibility, lumen volume elasticity K_V is such that apical

contraction compresses the super-ellipsoid by 5% in volume ($K_V = 1.10^{-5}u/\mu m^6$ (FigS4).

Preferred area A_0 and preferred perimeter P_0 are chosen to have a constant $P_0/\sqrt{A_0}$ ratio of 2

throughout the simulation, this corresponds to a stiff tissue in (Bi et al., 2015) framework.

With the above value of $P_0/\sqrt{A_0}$, we choose the apical perimeter elasticity K_p to allow cell

shape changes upon apical constriction.

The code used for modelling is publicly available: https://github.com/DamCB/tyssue and

https://github.com/suzannelab/polarity.

Quantification and statistical analysis

Quantitative analysis of the fold variability

We characterized the variability of fold formation *in vivo* by measuring the maximal distance

between the real fold from the predicted fold, the angle of the fold relative to the proximal-

distal axis and the parallelism between folds, for *control* and *Arpc5 RNAi* expressing leg discs

(genotypes are indicated in each figure legends), as follow:

- The maximal distance between real fold and predicted fold was measured in two steps. First,

we measured the distance between the Notch signaling (labeled by Deadpan immunostaining)

and the position of maximal deviation of the fold using the straight-line tool in ImageJ.

Second, we calculated the mean of maximal distance in the control and subtracted the value

obtained to each measure to normalize them independently from the genotype. Standard

deviation of control legs was measured and used to define predicted fold (± 1.30 µm from the

mean), which is highlighted in yellow on the graph in Figure 1c. Because the width of the

apterous domain is more variable in AJs GFP-trap experiments, in Figure 6d we measured the

relative maximal distance rather than the absolute maximal distance, by dividing the distance

between the T3-T4 and the T4-T5 fold by the width of the *apterous* domain.

- The angle formed between the fold and the proximal-distal axis, was measured in ImageJ

using the angle tool.

- The fold parallelism was calculated as the ratio between angles of two folds of the same leg.

Quantitative analysis of Dpn and Rpr expression domain

Dpn and Rpr stripes of expression have been characterized using Image J software. The

stripes have been outlined to define ROIs. From each ROI, the main axis (or ferret) has been

defined and the orientation of this axis relative to the proximo-distal axis defined. Then, the

voronoi (corresponding to the points equidistant from the proximal and distal border of the

domain) was obtained to determine the mean width of the domain and the curvature.

Cartography of mechanical perturbations *in vivo*

Zones of high apico-basal tension, or "mechanical perturbations", were spotted on 3D

reconstructions of (*sqh-GFP[29B]; Dll-Gal4*) and (*sqh-GFP[29B]; Dll-Gal4; UAS-*

Arpc5RNAi) leg discs using Imaris and located on corresponding positions on rolled-out maps

of the fold regions (Fig3f). To each perturbation corresponds a set of coordinates defined (in

x) by the angle formed between the line going from the center of the leg to the perturbation

and the DV axis and (in y) by the distance between the perturbation and the predicted fold.

Analysis of Junctional Intensities

Using the surface tool on Imaris (Bitplane), a mask was created from the junctional labelling

(arm-GFP, α-catenin-RFP or E-cadherin antibody) to quantify Myosin II, F-actin, adherens

junctions or Arp3 present at junctions. From this new Z-stack, the maximum of intensity

(MaxProj) and the sum of intensity (SumProj) were projected using Fiji (ImageJ 1.51s –

NIH). Then, from the MaxProj, a skeleton was created and each junction was individualized

by suppressing nodes. This template was used to identify each junction as a region of interest

(ROI). From these ROIs, the angle – formed with the proximal-distal axis – and the raw

fluorescence intensity of each junction were measured on the SumProj (allowing to sum the

full quantity of Myosin II present at a given xy position). Manual correction was done if

required.

The junctional mean of fluorescence intensity $\overline{I_j}$ along junction j was normalized to the

average junction intensity over all the junctions analyzed (fold+adjacent domain) $\frac{1}{n}\sum_{i=0}^{n}\overline{I_i}$:

$$I_{Rj} = \frac{\overline{I_j}}{\frac{1}{n}\sum_{i=0}^{n}\overline{I_i}}$$

Analysis of the cell aspect ratio

Using Fiji software, the maximal intensity projection was done from a Z-stack and a skeleton

was created. This template was used to identify each cell as a region of interest (ROI). From

these ROIs, the aspect ratio (major axis / minor axis) was measured in order to define the

anisotropy of the cell.

Quantitative analysis of the apical myosin

Using ImageJ software, a segmented line was drawn along the apical surface of the posterior

(hh-Gal4; UAS-GFP) and the anterior (control) domains on five different z sections for each

leg. For each z section, the mean of fluorescence intensity was measured and the ratio

between posterior and anterior domains was calculated. The mean of these five ratios per leg

is represented as one spot on the graph in Figure S5a.

Quantitative analysis of medio-apical myosin flow

Particle image velocimetry from time-lapse movies of E-cadh-GFP; sqh-RFPt leg discs was

quantified using the OpenPIV Python package. Images were acquired on a Zeiss LSM 880

using High Resolution Airy Scan with a time frame of 15". Flows were analyzed in 2D on

maximum projection of standard deviation. Each cell was isolated using Image J software. The displacement of particles was tracked between successive frames and the mean displacement is presented in a polar charts in FigS5e-f. n=14 for control and n=20 for Arpc5 RNAi.

Cartography of mechanical perturbations *in silico*

In the simulations, each cell has assigned coordinates, allowing us to spot the positions of each mechanical perturbation (randomly generated in the model) in a rolled-out map of the fold domain.

Measure of *in silico* fold deviation

We characterized the variability of fold orientation in the different sets of simulations: with or without mechanical perturbations and with or without tissue polarity. For each simulation, we spotted the position of maximal depth automatically as the point located at the minimal distance from the central axis of the tissue. We then defined its coordinates in relation to the predicted fold position and the DV axis. To calculate the real maximal deviation from the predicted fold domain, we subtracted the maximal deviation obtained without perturbation from the one obtained with perturbations for a given pattern of apoptosis. To calculate the angle formed between the fold and the proximal-distal axis, we measured the angle between the predicted fold and the line formed from fold position on the ventral side and the position of maximal depth. To calculate the real maximal angle from the predicted fold domain, we subtracted the angle without perturbation from the one obtained with perturbations for a given pattern of apoptosis.

Statistical Analysis

The normality of the data sets was determined using Prism 8 (Graph Pad).

A Mann-Whitney U-test was used to assess the significance of differences in tissue recoil after laser ablation (Fig2i; Fig3c,e; Fig7h-i and FigS7a-b) or in apical intensity of MyoII (FigS5a), considering legs as independent from each other. The null hypothesis was that measurements were samples from the same distribution. Tests were performed using Prism 8 (Graph Pad).

Variances of the phenotypes observed *in vivo* were compared using the F-test of equality of variances in Prism 8 (Graph Pad), considering that different data sets follow a normal distribution. Variances from simulated data were compared using the Levene's test in Python

3.7, considering that different data sets do not fit with a normal distribution. The null

hypothesis was that variances of population were equal.

Spearman correlation coefficients (FigS7g-h) and associated p-values were computed online

(www.wessa.net/rwasp_spearman.wasp).

Supplemental video titles

Movie 1. 3D reconstruction of a control leg disc showing the expression of *Deadpan* and the fold domain
(related to Fig2).

Movie 2. 3D reconstruction of an Arpc5 RNAi leg disc showing the expression of *Deadpan* and the fold domain
(related to Fig2).

Movie 3. Laser ablation experiments of apoptotic myosin II cables in *control* and *Arpc5* RNAi leg discs (Related
to Fig2).

Movie 4. Laser ablation experiments of non-apoptotic apico-basal structures of Myosin II in *control* and *Arpc5*
RNAi leg discs (Related to Fig3).

References

Akieda, Y., Ogamino, S., Furuie, H., Ishitani, S., Akiyoshi, R., Nogami, J., Masuda, T.,
Shimizu, N., Ohkawa, Y., and Ishitani, T. (2019). Cell competition corrects noisy Wnt morphogen
gradients to achieve robust patterning in the zebrafish embryo. Nat Commun *10*, 4710.
Ambrosini, A., Rayer, M., Monier, B., and Suzanne, M. (2019). Mechanical Function of
the Nucleus in Force Generation during Epithelial Morphogenesis. Dev Cell *50*, 197-211.e195.
Bertet, C., Sulak, L., and Lecuit, T. (2004). Myosin-dependent junction remodelling
controls planar cell intercalation and axis elongation. Nature *429*, 667-671.
Bi, D., Lopez, J.H., Schwarz, J.M., and Manning, M.L. (2015). A density-independent
rigidity transition in biological tissues. Nature Physics *11*, 1074-1079.
Calzolari, S., Terriente, J., and Pujades, C. (2014). Cell segregation in the vertebrate
hindbrain relies on actomyosin cables located at the interhombomeric boundaries. EMBO J *33*, 686-
701.
Chesarone, M.A., and Goode, B.L. (2009). Actin nucleation and elongation factors:
mechanisms and interplay. Curr Opin Cell Biol *21*, 28-37.
de Celis, J.F., Tyler, D.M., de Celis, J., and Bray, S.J. (1998). Notch signalling mediates
segmentation of the Drosophila leg. Development *125*, 4617-4626.
Ebert, M.S., and Sharp, P.A. (2012). Roles for microRNAs in conferring robustness to
biological processes. Cell *149*, 515-524.
Eldar, A., Shilo, B.Z., and Barkai, N. (2004). Elucidating mechanisms underlying
robustness of morphogen gradients. Curr Opin Genet Dev *14*, 435-439.
Eritano, A.S., Bromley, C.L., Bolea Albero, A., Schütz, L., Wen, F.L., Takeda, M.,
Fukaya, T., Sami, M.M., Shibata, T., Lemke, S., *et al.* (2020). Tissue-Scale Mechanical Coupling
Reduces Morphogenetic Noise to Ensure Precision during Epithelial Folding. Dev Cell.
Fernandez-Gonzalez, R., and Zallen, J.A. (2013). Wounded cells drive rapid epidermal
repair in the early Drosophila embryo. Mol Biol Cell *24*, 3227-3237.
Frankel, N., Davis, G.K., Vargas, D., Wang, S., Payre, F., and Stern, D.L. (2010).
Phenotypic robustness conferred by apparently redundant transcriptional enhancers. Nature *466*,
490-493.
Félix, M.A., and Barkoulas, M. (2015). Pervasive robustness in biological systems. Nat
Rev Genet *16*, 483-496.
Georgiou, M., Marinari, E., Burden, J., and Baum, B. (2008). Cdc42, Par6, and aPKC
regulate Arp2/3-mediated endocytosis to control local adherens junction stability. Curr Biol *18*, 1631-
1638.
Gilmour, D., Rembold, M., and Leptin, M. (2017). From morphogen to
morphogenesis and back. Nature *541*, 311-320.
Gracia, M., Theis, S., Proag, A., Gay, G., Benassayag, C., and Suzanne, M. (2019).
Mechanical impact of epithelial-mesenchymal transition on epithelial morphogenesis in Drosophila.
Nat Commun *10*, 2951.
Gregor, T., Tank, D.W., Wieschaus, E.F., and Bialek, W. (2007). Probing the limits to
positional information. Cell *130*, 153-164.
Gursky, V.V., Surkova, S.Y., and Samsonova, M.G. (2012). Mechanisms of
developmental robustness. Biosystems *109*, 329-335.
Hardway, H., Mukhopadhyay, B., Burke, T., James Hitchman, T., and Forman, R.
(2008). Modeling the precision and robustness of Hunchback border during Drosophila embryonic
development. J Theor Biol *254*, 390-399.
Harmansa, S., Alborelli, I., Bieli, D., Caussinus, E., and Affolter, M. (2017). A
nanobody-based toolset to investigate the role of protein localization and dispersal in. Elife *6*.

885   Harmansa, S., Hamaratoglu, F., Affolter, M., and Caussinus, E. (2015). Dpp spreading
886   is required for medial but not for lateral wing disc growth. Nature *527*, 317-322.

887   He, F., Wen, Y., Deng, J., Lin, X., Lu, L.J., Jiao, R., and Ma, J. (2008). Probing intrinsic
888   properties of a robust morphogen gradient in Drosophila. Dev Cell *15*, 558-567.

889   Herranz, H., and Cohen, S.M. (2010). MicroRNAs and gene regulatory networks:
890   managing the impact of noise in biological systems. Genes Dev *24*, 1339-1344.

891   Hong, L., Dumond, M., Zhu, M., Tsugawa, S., Li, C.B., Boudaoud, A., Hamant, O., and
892   Roeder, A.H.K. (2018). Heterogeneity and Robustness in Plant Morphogenesis: From Cells to Organs.
893   Annu Rev Plant Biol *69*, 469-495.

894   Lander, A.D. (2011). Pattern, growth, and control. Cell *144*, 955-969.

895   Lecuit, T., and Cohen, S.M. (1997). Proximal-distal axis formation in the Drosophila
896   leg. Nature *388*, 139-145.

897   Lecuit, T., and Yap, A.S. (2015). E-cadherin junctions as active mechanical integrators
898   in tissue dynamics. Nat Cell Biol *17*, 533-539.

899   Lott, S.E., Kreitman, M., Palsson, A., Alekseeva, E., and Ludwig, M.Z. (2007).
900   Canalization of segmentation and its evolution in Drosophila. Proc Natl Acad Sci U S A *104*, 10926-
901   10931.

902   Major, R.J., and Irvine, K.D. (2006). Localization and requirement for Myosin II at the
903   dorsal-ventral compartment boundary of the Drosophila wing. Dev Dyn *235*, 3051-3058.

904   Manjón, C., Sánchez-Herrero, E., and Suzanne, M. (2007). Sharp boundaries of Dpp
905   signalling trigger local cell death required for Drosophila leg morphogenesis. Nat Cell Biol *9*, 57-63.

906   Manu, Surkova, S., Spirov, A.V., Gursky, V.V., Janssens, H., Kim, A.R., Radulescu, O.,
907   Vanario-Alonso, C.E., Sharp, D.H., Samsonova, M., *et al.* (2009). Canalization of gene expression in
908   the Drosophila blastoderm by gap gene cross regulation. PLoS Biol *7*, e1000049.

909   Michelot, A., and Drubin, D.G. (2011). Building distinct actin filament networks in a
910   common cytoplasm. Curr Biol *21*, R560-569.

911   Monier, B., Gettings, M., Gay, G., Mangeat, T., Schott, S., Guarner, A., and Suzanne,
912   M. (2015). Apico-basal forces exerted by apoptotic cells drive epithelium folding. Nature *518*, 245-
913   248.

914   Monier, B., Pélissier-Monier, A., Brand, A.H., and Sanson, B. (2010). An actomyosin-
915   based barrier inhibits cell mixing at compartmental boundaries in Drosophila embryos. Nat Cell Biol
916   *12*, 60-65; sup pp 61-69.

917   Nishimura, T., Honda, H., and Takeichi, M. (2012). Planar cell polarity links axes of
918   spatial dynamics in neural-tube closure. Cell *149*, 1084-1097.

919   Perry, M.W., Boettiger, A.N., Bothma, J.P., and Levine, M. (2010). Shadow enhancers
920   foster robustness of Drosophila gastrulation. Curr Biol *20*, 1562-1567.

921   Pollard, T.D., and Beltzner, C.C. (2002). Structure and function of the Arp2/3
922   complex. Curr Opin Struct Biol *12*, 768-774.

923   Rajan, A., Tien, A.C., Haueter, C.M., Schulze, K.L., and Bellen, H.J. (2009). The Arp2/3
924   complex and WASp are required for apical trafficking of Delta into microvilli during cell fate
925   specification of sensory organ precursors. Nat Cell Biol *11*, 815-824.

926   Rauskolb, C., and Irvine, K.D. (1999). Notch-mediated segmentation and growth
927   control of the Drosophila leg. Dev Biol *210*, 339-350.

928   Rauzi, M., Lenne, P.F., and Lecuit, T. (2010). Planar polarized actomyosin contractile
929   flows control epithelial junction remodelling. Nature *468*, 1110-1114.

930   Robinson, R.C., Turbedsky, K., Kaiser, D.A., Marchand, J.B., Higgs, H.N., Choe, S., and
931   Pollard, T.D. (2001). Crystal structure of Arp2/3 complex. Science *294*, 1679-1684.

932   Sato, A. (2018). Chaperones, Canalization, and Evolution of Animal Forms. Int J Mol
933   Sci *19*.

934   Schott, S., Ambrosini, A., Barbaste, A., Benassayag, C., Gracia, M., Proag, A., Rayer,
935   M., Monier, B., and Suzanne, M. (2017). A fluorescent toolkit for spatiotemporal tracking of
936   apoptotic cells in living. Development *144*, 3840-3846.

937   Smith, S.J., Rebeiz, M., and Davidson, L. (2018). From pattern to process: studies at
938 the interface of gene regulatory networks, morphogenesis, and evolution. Curr Opin Genet Dev *51*,
939 103-110.
940   van der Walt, S., Colbert, S.C., and Varoquaux, G. (2011). The NumPy Array: A
941 Structure for Efficient Numerical Computation. Computing in Science and Engineering *13*, 22-30.
942   Waddington (1942). Canalisation of development and the inheritance of acquired
943 characters, pp. 563 – 565.
944   Wu, J., and Cohen, S.M. (1999). Proximodistal axis formation in the Drosophila leg:
945 subdivision into proximal and distal domains by Homothorax and Distal-less. Development *126*, 109-
946 117.
947   Yevick, H.G., Miller, P.W., Dunkel, J., and Martin, A.C. (2019). Structural Redundancy
948 in Supracellular Actomyosin Networks Enables Robust Tissue Folding. Dev Cell *50*, 586-598.e583.
949   Younger-Shepherd, S., Vaessin, H., Bier, E., Jan, L.Y., and Jan, Y.N. (1992). deadpan,
950 an essential pan-neural gene encoding an HLH protein, acts as a denominator in Drosophila sex
951 determination. Cell *70*, 911-922.

Developmental patterning

Notch activation   Pro-apoptotic genes

control

Arp2/3 knockdown

planar polarized Myosin II

↓

Force channeling

↓

STEREOTYPED MORPHOGENESIS

unpolarized Myosin II

↓

Force scattering

↓

VARIABLE MORPHOGENESIS

pulling apoptotic cell

mechanical perturbation

anisotropic force transmission

anisotropic force transmission

isotropic force transmission

isotropic force transmission

a *control*

β-cat

T1 T2 T3 T4 T5

T1-T2 T2-T3 T3-T4 T4-T5

Folds

Tarsal domain

Circ.

Pr-Di

b *Arpc5 RNAi*

T1 T2 T3 T4 T5

incompletely formed

c Maximal distance from predicted fold (µm)

F-test

*** *** ***

control

Arpc5 RNAi

−10   0   +10   −10   0   +10   −10   0   +10

T2-T3   T3-T4   T4-T5

d Angle from proximal-distal axis (°)

F-test

*** ns ***

control

Arpc5 RNAi

110   90   70   110   90   70   110   90   70

T2-T3   T3-T4   T4-T5

e Fold parallelism

F-test

*** *** ***

control

Arpc5 RNAi

0.5   1.0   1.5   0.5   1.0   1.5   0.5   1.0   1.5

$\dfrac{\theta T2\text{-}T3}{\theta T3\text{-}T4}$ $\dfrac{\theta T2\text{-}T3}{\theta T4\text{-}T5}$ $\dfrac{\theta T3\text{-}T4}{\theta T4\text{-}T5}$

f *control*   *Arpc5 RNAi*

d   d

control   *Arpc5 RNAi*

Pr-Di (0°)   θ   Pr-Di (0°)   θ

control   *Arpc5 RNAi*

θ   θ

T2-T3 T3-T4 T4-T5   T2-T3 T3-T4 T4-T5

a Notch signalling

control   *Arpc5 RNAi*

dpn   Actin

b Pro-apoptotic gene (*rpr*)

control   *Arpc5 RNAi*

Actin

control

Arpc5 RNAi

c Angle of dpn expression stripes from proximal-distal axis (°)

F-test   ns   ns   ns

control

Arpc5

110   90   70   110   90   70   110   90   70

T2-T3   **T3-T4**   **T4-T5**

d Angle of rpr expression stripes from proximal-distal axis (°)

F-test   ns   ns   ns

control

Arpc5

110   90   70   110   90   70   110   90   70

T2-T3   **T3-T4**   **T4-T5**

e *control*

Predicted fold   Fold

Notch signaling (dpn)

f *Arpc5 RNAi*

Predicted fold   Fold

Notch signaling (dpn)

g *control*

Arpc5 RNAi

Apoptotic cells

ns   ns

control   Arpc5   control   Arpc5

Inside   Outside

predicted fold domain

h Apoptosis

control   *Arpc5 RNAi*

Ap

B

Apoptosensor / MyoII

i Apoptotic forces

Velocity (µm/s)

ns

control   Arpc5

a

Normal fold = Predicted fold
Deviated fold ≠ Predicted fold

Robust folding (control)

Variable folding (Arpc5 RNAi)

Distal (66%)   Proximal (34%)

b Deviation towards apical structure
control   Arpc5

Circ.
└ Pr-Di   MyoII

c apical cut
1 ←→ 2

Velocity (µm/s)
ns * ** ns
control Arpc5 control Arpc5
(1) - control   (2) - 'N border'

d Deviation towards apico-basal structure
control   Arpc5

Circ.
└ Pr-Di   MyoII

e Ap-B cut
3 ←→ 4

Velocity (µm/s)
ns *** *** ns
control Arpc5 control Arpc5
(3) - control   (4) - Ap-B structure

f
V   D   V   V   D   V
control   Arpc5 RNAi

Dist
Ant ┼ Post.
Prox

µm

predicted fold   folding domain   Ap-B structure   N border

g
No mechanical perturbation   Mechanical perturbation

Circ.
└ Pr-Di

h
V. Ant ┼ Dist / V. Post
Prox

predicted fold
Mechanical perturbation

i
Maximal distance from predicted fold (AU)
***
Mechanical perturbation − +

j
Angle from proximo-distal axis (°)
***
Mechanical perturbation − +

a

control · Arpc5 RNAi

β-cat::GFP Phalloidin β-cat::GFP Phalloidin

Actin

Circ.
(90°)

Pr-Di
(0°)

c

α-cat · Arp3

b

□ *control* ■ *Arpc5 RNAi*

Actin
Rel. Fluo. Int

fold domain

adjacent domain

Pr-Di 0°-15° 15°-30° 30°-45° 45°-60° 60°-75° 75°-90° Circ.
Pr-Di 0°-15° 15°-30° 30°-45° 45°-60° 60°-75° 75°-90° Circ.

d

■ *Fold* ■ *Adjacent*

Arp3
Rel. Fluo. Int

Pr-Di 0°-15° 15°-30° 30°-45° 45°-60° 60°-75° 75°-90° Circ.

a Distance from the cut site (Cell diameter)

0 — 4

Before cut

After cut

F

vertices displacement

d

b control

Before cut | After cut | Overlay

Before | After

c

90° | 180° | 0° | 270°

d Arpc5 RNAi

Before cut | After cut | Overlay

Before | After

e

90° | 180° | 0° | 270°

f

Before circular excision | After circular excision

Steady state

-

F_{ext} | F_{int}

Circular cut

Maximal recoil

d | d*

+ Ectopic force

Wound

F_{Wtr} | F_{ext} | F_{int} | F_w

Circular cut during healing

Maximal recoil

d_w | d_w*

Circ. (90°)

Pr-Di (0°)

g

control | Arpc5 RNAi

Steady state

+ Ectopic force

h Internal recoil

control | Arpc5 RNAi

Maximal recoil (μm)

2,5 | 2 | 1,5 | 1 | 0,5 | 0

ns | *** | * | ns | ** | ns | **

PrDi | Circ. | PrDi | Circ.

wound − | + | − | +

d* | dw* | d* | dw*

i External recoil

control | Arpc5 RNAi

Maximal recoil (μm)

2,5 | 2 | 1,5 | 1 | 0,5 | 0

*** | ns | * | *** | ns | ns | ns

PrDi | Circ. | PrDi | Circ.

wound − | + | − | +

d | dw | d | dw