Title: *GmSALT3* confers shoot Na⁺ and Cl⁻ exclusion in soybean via two distinct processes

Yue Qu[1†], Rongxia Guan[2†], Jayakumar Bose[1], Sam W. Henderson[1], Stefanie Wege[1*], Li-Juan Qiu[2*], Matthew Gilliham[1*]

[1]Australian Research Council Centre of Excellence in Plant Energy Biology, School of Agriculture, Food and Wine & Waite Research Institute, University of Adelaide, Glen Osmond, SA, Australia

[2]The National Key Facility for Crop Gene Resources and Genetic Improvement, Institute of Crop Science, Chinese Academy of Agricultural Sciences, Beijing, China

[*]For correspondence (e-mails matthew.gilliham@adelaide.edu.au; qiulijuan@caas.cn; stephanie.wege@adelaide.edu.au).

[†]These authors have contributed equally to this work.

Funding information

Natural Science Foundation of China, grant number 31830066; Scientific Innovation Project of Chinese Academy of Agricultural Sciences; ARC Centre of Excellence funding CE140100008; ARC Fellowships FT130100709, DE160100804 and DE170100346.

Abstract

Soybean (*Glycine max*) yields are threatened by multiple stresses including soil salinity. *GmSALT3* confers net shoot exclusion for both Na⁺ and Cl⁻ to soybean and improved salt tolerance; however, how the ER-localised GmSALT3 achieves this is unknown. Here, GmSALT3 function was investigated in heterologous systems, and in near isogenic lines that contained the full-length *GmSALT3* gene (NIL-T; salt-tolerant) or a truncated transcript (NIL-S; salt-sensitive). A time course analysis of the NILs confirmed shoot Cl⁻ exclusion breaks down days prior to Na⁺ exclusion. Shoot Na⁺ exclusion occurs via a root xylem-based mechanism in NIL-T lines, as is common amongst plants, with Na⁺ content significantly greater in NIL-S stem xylem sap. In contrast, NIL-T plants exhibited significantly greater Cl⁻ content in both the stem xylem and phloem sap compared to NIL-S, indicating that shoot Cl⁻ exclusion likely depends upon a novel phloem-based Cl⁻ recirculation back to NIL-T roots. In

heterologous expression systems, GmSALT3 restored growth of K^+-uptake-defective *E. coli*, whilst in *Xenopus laevis* oocytes, GmSALT3 contributed to net influx and accumulation of Na^+, K^+, and Cl^-. Overall, these findings provide new insights on the impact of ER-localised GmSALT3 on plasma membrane ion fluxes and reveals a novel mechanism for shoot Cl^- exclusion.

Highlight

GmSALT3 is a candidate gene that impacts soybean salt tolerance. Here, we found GmSALT3 can affect Na^+, K^+, and Cl^- transport in heterologous systems, and is likely to contribute to soybean ionic toxicity tolerance through phloem-based Cl^- recirculation from shoot and root xylem-based Na^+ exclusion.

Keywords: Salinity tolerance, abiotic stress, soybean, GmSALT3, CHX, ion transporter, shoot ion exclusion

Acknowledgments:

This work was funded by the Natural Science Foundation of China, grant number 31830066 (RG); Scientific Innovation Project of Chinese Academy of Agricultural Sciences (L-J Q). YQ and SH were supported by ARC Centre of Excellence funding awarded to MG (CE140100008). ARC Fellowships supported MG (FT130100709), SW (DE160100804) and JB (DE170100346). We thank Yue (Crystal) Wu for generating the *AtCBL1n* entry vector; Dr. Sunita Ramesh and Assoc. Prof Zhonghua Chen for providing *AtKAT1* construct used in this study; Dr. Gwenda Mayo from Adelaide Microscopy for providing assistance in TEM.

Word count: 6133 (Abstract, Introduction, Methods, Results, Discussion, Acknowledgements)

Introduction

There is a large market demand for soybean (*Glycine max*) as a staple edible oil and food crop; soybean is additionally an important component of intercropping and sequential cropping systems (Singh, 2010). As such, soybean production demand has been projected to increase by nearly 80 percent to 390 Mt by 2050 (Alexandratos & Bruinsma, 2012). However, salinity stress threatens soybean production; soil salinity of 18–20 dS/m (approximately 180 – 200 mM NaCl) has been shown to reduce yield by 61% – from 2261 ± 438 kg/ha under control conditions to 881 ± 260 kg/ha (Chang, Chen, Shao, & Wan, 1994). Nevertheless, not all soybean cultivars are equally as sensitive to salinity stress, with some rare soybean germplasm found to be relatively salt-tolerant. In a large-scale evaluation of soybean salinity tolerance, less than 3% of the 10,128 evaluated soybean germplasm exhibited salt tolerance at both the germination stage and seedling stage, only 83 were found to be highly tolerant to salinity at the vegetative stage only (Shao, Wan, Chang, & Chen, 1993). It has been observed that salt-tolerant soybean cultivars often have superior agronomic performance to salt-sensitive cultivars (Phang, Shao, & Lam, 2008), indicating breeding for stress tolerance soybean may improve overall yields even under standard conditions.

In soybean, a major quantitative trait locus (QTL) for salinity tolerance has been consistently mapped to chromosome 3 (Abel, 1969; Ha et al., 2013; Hamwieh et al., 2011; Hamwieh & Xu, 2008; Lee et al., 2004). Since 2014, several studies have been published focusing on the same dominant gene, identified in this chromosome 3 QTL in wild (*GmCHX1*) and cultivated soybean (*GmNcl/GmSALT3*) (Do et al., 2016; Guan et al., 2014; Qi et al., 2014). This gene, *Glyma03g32900* in reference soybean genome of the William 82 variety (William 82. a1.v1.1), is the best candidate for the *Ncl* locus reported by Abel (1969). It should be noted here that in the William 82 genome a2.v1, *Glyma03g32900*, appears to be incorrectly annotated as two transcripts (Glyma.03g171600 and Glyma.03g171700) rather than the full-length coding sequence obtained consistently from cDNA and encoding a full-length CHX protein. Here, we refer to the full-length gene obtained as *GmSALT3*.

According to phylogenetic analyses, the full-length GmSALT3 protein is closely related to *Arabidopsis thaliana* AtCHX20 (a Cation/Proton Exchanger), which is a functionally characterised member of the CPA2 (Cation/Proton Antiporter 2) family of transporters (Guan et al., 2014; Padmanaban et al., 2007). Almost all CHXs that have been functionally characterised have been done so only in heterologous systems. A connection to their roles in plants or their association with phenotypes in loss of function mutants is sparse; GmSALT3 is

the only CHX so far connected to salinity tolerance (Sze & Chanroj, 2018). Functional studies

of AtCHXs have indicated that they might play role in modulating cation and pH homeostasis

within the endomembrane system (Chanroj et al., 2011; Czerny et al., 2016; Padmanaban et al.,

2007). The ER-localised AtCHX20 was, for example, suggested to be an endomembrane K^+

transporter involved in the osmoregulation of guard cells (Padmanaban et al., 2007).

The salt-tolerant soybean cultivar Tiefeng 8 contains a full-length *GmSALT3*, while the

salt sensitive cultivar 85-140 contains a 3.78-kb copia retrotransposon insertion in exon 3 of

GmSALT3 that truncates the transcript with a premature stop codon (making the 85-140

essentially a *Gmsalt3* knockout). *GmSALT3* is predominantly expressed in root cells associated

with the phloem and xylem (Guan et al., 2014); expression of full-length *GmSALT3* in roots

leads to reduced shoot accumulation of Na^+ (Guan et al., 2014; Qi et al., 2014) and Cl^- (Liu et

al., 2016) and increased crop yield in saline conditions (Do et al., 2016; Liu et al., 2016).

GmSALT3 has been localized to the endoplasmic reticulum (ER), similar to AtCHX20. This

differs from most other proteins known to be involved in shoot Na^+ and Cl^- exclusion, which

have mainly been localised to the plasma membrane and tonoplast.

Shoot salt exclusion requires the co-ordinated activity of many ion transporters that

may: 1) limit the net entry of salt into the roots, 2) compartmentalise salt in the roots; and or 3)

retrieve salt from the root xylem or shoots through the phloem. Examples of transporters that

have been implicated in salt tolerance in soybean include: the tonoplast localised GmNHX1

(Na^+/H^+ antiporter 1) (Li et al., 2006a); and the plasma membrane localised GmSOS1 (salt

overly sensitive 1, Na^+/H^+ antiporter) and GmCAX1 (Ca^{2+}/H^+ antiporter 1) (Luo et al., 2005;

Phang et al., 2008). These transporters are proposed to be involved in net exclusion of salt entry

into roots (GmSOS1), its compartmentation (*GmNHX1*) or signalling (*GmCAX1*). Significantly,

the transporters contributing to reduced net transfer of salt to the shoot, the mechanism by

which shoot salt accumulation is commonly regulated in other species, are yet to be resolved

in soybean (Cao, Li, Liu, Kong, & Tran, 2018; Munns & Tester, 2008). For instance, in wheat,

rice and Arabidopsis the HKT1;5-like (high affinity K^+ transport) proteins that localise to cells

that surround the root xylem facilitate Na^+ retrieval back into the root (Rana Munns et al., 2012;

Møller et al., 2009; Ren et al., 2005; Uozumi et al., 2000). GmSALT3 differs from these

previous proteins in being ER-localised and so is unlikely to directly confer shoot salt exclusion

via retrieval from the xylem sap. Therefore, the mechanism by which GmSALT3 imparts

improved salt tolerance is yet to be determined.

Here, we investigated the function of GmSALT3 in heterologous systems and *in planta*

to gain insights into the mechanisms by which GmSALT3 contributes to salt tolerance. In

heterologous systems, *GmSALT3* complemented K$^+$ uptake deficient *Escherichia coli*; while

in *Xenopus laevis* oocytes, *GmSALT3* expression resulted in increased net influx of Na$^+$, K$^+$

and Cl$^-$. Phenotypes of soybean NILs differing in *GmSALT3* alleles revealed that plants

containing full-length *GmSALT3,* compared to truncated *Gmsalt3,* are likely to employ two

distinct mechanisms to achieve shoot Na$^+$ and Cl$^-$ exclusion by facilitating net xylem retrieval

of Na$^+$ and greater phloem Cl$^-$ retranslocation from shoots to roots. As such, we propose that

the endomembrane GmSALT3 has a complex physiological role which ultimately impacts

multiple ion fluxes across the plasma membrane.

Materials and Methods

Plant materials and growth conditions

Soybean plants were grown in a greenhouse (28 °C day, 16 h light and 25 °C night) at

the Plant Research Centre, Waite Campus, University of Adelaide, Australia. Soybean seeds

(Tiefeng 8, salt-tolerant parent; 85-140, salt-sensitive parent; NIL-T, near isogenic line with

GmSALT3; NIL-S, near isogenic line with *Gmsalt3*) were imported from China through

quarantine. Soybean seeds were germinated in pots containing a 1:1 mixture of

perlite:vermiculite as described by Obermeyer & Tyerman (2005).

cDNA cloning and plasmid preparation

To synthesize *GmSALT3* and *Gmsalt3* cDNA (2436 and 1131 nucleotides, respectively),

total RNA was isolated from roots of 4-week old soybean plants using the TRIzol method (Shi

& Bressan, 2006). First-strand cDNA was synthesized using a Thermoscript RT III kit

(Invitrogen, USA). Gene specific primers (GmSALT3_gF and GmSALT3_gR, Gmsalt3_gF

and Gmsalt3_gR; Table S1) were used to amplify *GmSALT3* cDNA with Phusion® High-

Fidelity DNA Polymerase (New England Biolabs) using 35 cycles (98 °C 30 s, 65 °C 30 s, and

72 °C 150 s). Gel-purified PCR products were A-tailed using Taq polymerase (New England

Biolabs) for 30 min at 72 °C, and then recombined into Gateway® entry vector

PCR8/GW/TOPO (Invitrogen) for subsequent applications. Resulting clones were sequenced

using internal primers (GmSALT3_F1, R1, F2, F3, F4, and F5, Gmsalt_F1, F2, and R1; Table

S1). The sequence of *GmSALT3-YFP* was PCR amplified from a previously constructed

plasmid *pBS-35S::GmSALT3-YFP*; the sequence of *GmSALT3-TM10* (first 10 transmembrane

domain of *GmSALT3*) was PCR amplified from *PCR8-GmSALT3*. Gel-purified PCR products

of *GmSALT3-YFP, GmSALT3-TM10* were recombined into entry vector PCR8 as described

above.

GmSALT3 and *Gmsalt3* CDS, *GmSALT3-YFP, GmSALT3-TM10, AtCHX20 and*

AtKAT1 within entry vector PCR8 were cloned into the pGEM-HE vector and a modified

version of pET-DEST42 vector using Gateway ® LR Clonase® (Invitrogen, USA), for *X. laevis*

oocyte expression and *E. coli* expression, respectively. The *E. coli* strain TK2463 lacks the T7

RNA polymerase, therefore, the T7 promoter in pET-DEST42 was replaced with the TAC

promoter. Inverse PCR was performed with 5 ng of plasmid, 50 nM of forward and reverse

primer (SH_215 and SH_216, Table S1), and 0.25 units Phusion High Fidelity Polymerase

(New England Biolabs) in a final volume of 25 µL. Reactions were treated with Cloning

Enhancer (Clontech Laboratories Inc) to remove circular template, and the plasmid was re-

circularised using In-Fusion HD enzyme (Clontech Laboratories Inc).

Growth assay in Escherichia coli

Competent cell preparation and *E. coli* transformation were conducted as described by

Chanroj et al. (2011). The resulting transformants were grown on KML media (10 g/L tryptone,

5 g/L yeast extract, 10 g/L KCl) supplemented with 100 µg/mL carbenicillin and incubated at

37 ºC overnight; positively transformed colonies were identified by colony PCR with gene

specific primers (GmSALT3_gF and _gR; Gmsalt3_gF and gR). For *E. coli* growth

experiments, freshly transformed cells were first grown overnight in 5 ml KML at pH 7.2. Cell

cultures were replenished (OD_{600} = 0.5) and grown for 3 h in KML, and then washed with low

potassium media (10 g/L tryptone, 2 g/L yeast extract, 100 mM D-mannitol) three times. Cells

were normalized to OD_{600} 0.5 for a 96-well plate assay. In each well, 20 µL of normalized cell

suspension was added to 180 µL growth solutions. All test media were supplemented with 100

µg/mL carbenicillin, with or without 0.5 mM IPTG (Isopropyl β-D-1-thiogalactopyranoside).

The 96-well plates were inserted into FLUOstar Omega Fluorescence microplate reader (BMG

LABTECH) set to 37 ºC and A_{600} was measured every 15 min for 18 h with continuous shaking

at 200 rpm.

Characterization of GmSALT3 in X. laevis oocytes

pGEMHE-DEST containing *GmSALT3* was linearized using SphI-HF (New England

BioLabs); cRNA was synthesized using mMESSAGE mMACHINE T7 Kit (Ambion). 46

nL/23 ng of cRNA or equal volumes of RNase-free water were injected into oocytes with a

Nanoinject II microinjector (Drummond Scientific). Oocytes were incubated for 48 h in

Calcium Ringer's solution (96 mM NaCl, 2 mM KCl, 5 mM MgCl$_2$, 5 mM HEPES, 0.6 mM

CaCl$_2$). All solution osmolalities were measured with a Vapor pressure osmometer (Wescor)

and adjusted to 240–260 mOsmol kg^{-1} using mannitol. Measurement of ion profiles in oocytes

followed Munns et al. (2012) with the following modifications. Six replications of 3 grouped

oocytes were used for flame photometry (Sherwood 420), and Cl$^-$ was measured by manual

titrimetric Cl$^-$ measurement using a chloride analyser (Sherwood® Chloridometer 926S). For

YFP imaging, oocytes were analysed with a Zeiss LSM 5 Pascal Confocal Laser-Scanning

Microscope equipped with an argon laser. Excitation/emission conditions for YFP were 514

nm/523 to 600 nm.

MIFE (Microelectrode Ion Flux Estimation) from oocytes

The MIFE technique allows noninvasive concurrent quantification of net fluxes of

several ions using protocols outlined by Shabala et al. (2013) with oocytes being used rather

than plant tissues in our experiments; sodium, potassium, and chloride commercial ionophore

cocktails (93314, 99311, and 99408, respectively; all from Sigma-Aldrich) were used as LIX

(liquid Ion Exchanger) when measuring Na$^+$, K$^+$, and Cl$^-$ fluxes. Oocytes were washed in ND96

(96 mM NaCl, 1 mM KCl, 1 mM MgCl$_2$, 5 mM HEPES, pH 7.5). After incubation in ND96

for 72 hours, fluxes were measured after being transferred to BSM (5 mM NaCl, 0.2 mM KCl,

0.2 mM CaCl$_2$, 5 mM HEPES, pH 7.5).

TEM (Transmission Electron Microscopy)

Fresh soybean NIL-T and NIL-S roots were sectioned to 1 mm (length), stems were

cross sectioned (1 mm in length and 1 mm in radius). Root and stem samples were incubated

overnight in 1.5 ml Eppendorf tubes with fixative (2.5% Glutaraldehyde, 4% Formaldehyde,

4% Sucrose, 0.1 M Phosphate buffer). Samples were washed three times in 1X PBS and then

washed with osmium (263257, Sigma-Aldrich) for 4 hours. Sections were then washed three

times in 1X PBS and soaked in 1X PBS for 20 mins and then embedded in 1% agarose gel.

Agarose embedded soybean root were cut into blocks (1 cm long). A series of dehydration

steps were performed after embedding, using increasing concentrations of ethanol: 50%, 70%,

90% and 95% (each for 30 mins), and 100% ethanol for overnight dehydration. Samples were

then incubated in increasing concentrations of resin (Spurrs): 5%, 10%, 15%, 20%, 25%, 30%,

40%, and 50% (each for 1 h), 50% overnight, 75% (4 h), 100% (4 h), and 100% (overnight).

Resin wells containing the agarose blocks were then incubated at 60 °C for 3 days for

polymerization. Polymerised resin capsules were cut to 70 nm thickness using a diamond knife and visualized under transmission electron microscopy (TEM; PHILIPS CM100).

NaCl treatment and ion accumulation

Soybean plants were treated with 100 mM NaCl every 2 days (1.5 L solution per tray) at 14 days after sowing; the second dose of 1.5 L 100 mM NaCl was applied 16 days after sowing; the same volume of RO water was applied to the tray of the control plants. Saline solutions or RO water were applied into trays that contain pots with soybean plants. Plant tissues were harvested and dried in oven overnight at 60°C or freeze-dried overnight. Dry weight was recorded. Then dried samples were digested in 10 mL 1% (v/v) nitric acid overnight at 65°C or freeze-thawed in Milli-Q water 3 times. Oven-dried samples were utilized to test Na^+ and K^+ accumulation with flame photometry (Sherwood 420), Cl^- was measured by titrimetric Cl^- measurement by using chloride analyser (Sherwood® Chloridometer 926S). Freeze-dried samples were used to measure NO_3 accumulation. The measurement method was modified according to Qiu, Henderson, Tester, Roy, & Gilliham (2016). Supernatant (50 μL) of freeze-dried samples in water was added into 200 μL of 5% (W/V) salicylic acid/H_2SO_4 and incubated at room temperature for 20 min, then 50 μL of this mixture was transferred into 950 μL of 2 M NaOH and incubated at room temperature for at least 20 min (to cool down to room temperature). Of this, 250 μL was loaded into separate wells of a flat-bottom transparent 96-well plate (Greiner Bio-One, Austria). Absorbance was measured at 410 nm. Standards measured included 0, 0.5, 1, 2, 2.5, 5, 7.5, and 10 mM KNO_3.

Phloem and xylem sap extraction

Phloem sap was extracted according to the method of Rupassara (2008) and Ren et al. (2005). Soybean plant petioles were cut at the base (approx. 0.5 cm above the main stem) and stems were cut approx. 2 cm above ground (upper part), and immediately dipped in 1.5 mL of 0.1 mM EDTA solution (pH adjusted to 8 with NaOH) in a 2 mL microcentrifuge tube for 20 min, then the tube was immediately dipped in liquid nitrogen and stored in -80 °C until analysis. The glutamine concentration in the extract was measured using a Glutamine Assay Kit (EGLN-100, EnzyChrom™, BioAssay Systems). Xylem sap was extracted using a pressure chamber. Lower parts of the cut stems with roots were transferred into the pressure chamber. Pressure was increased gradually (0.05 MPa increments) until the xylem sap presented. The first two drops emerging were discarded using a micropipette to reduce contamination from damaged

cells or phloem sap (Berthomieu et al., 2003). Xylem sap was then collected during the

following 5 min, and stored at -20 $^{\circ}$C until analysis. Ion contents was measured as stated above.

Results

GmSALT3 modulates Na$^+$, K$^+$, and Cl$^-$ accumulation in soybean

Liu et al. (2016) previously used an extreme salt treatment (200 mM NaCl) that

typically results in the death of even the salt tolerant germplasm after a prolonged exposure.

Using this method, it was observed that full-length *GmSALT3* improved shoot Na$^+$ and Cl$^-$

exclusion over 10 days, with higher shoot Cl$^-$ accumulation occurring prior to shoot Na$^+$

accumulation in NIL-S (Liu et al., 2016). Here, we used a lower salt concentration (100 mM)

that soybean plants are more likely to encounter in the field and which typically does not result

in death of the plants prior to seed production. Furthermore, we studied the time course of Na$^+$,

Cl$^-$ and K$^+$ accumulation in NIL-T and NIL-S for different leaf types, stems and hypocotyls

(Fig. 1). We confirmed that Cl$^-$ accumulates to higher concentrations in NIL-S shoots

compared to those of NIL-T (Fig. 1b). Additionally, also without salt treatment, we found a

higher Cl$^-$ concentration in NIL-S aerial tissues (leaves and stems) without salt treatment;

whereas significantly higher Na$^+$ accumulation was detected in aerial tissues only after 3 days

of NaCl treatment in NIL-S, and higher Cl$^-$ accumulation continued (Fig. 1a,b; Fig. S1, S2).

We further observed that K$^+$ content was increased in NIL-S compared to NIL-T in these tissues

following NaCl treatment (Fig. 1c; Fig. S3). The observed accumulation of K$^+$ showed a similar

trend to Na$^+$, but to a lesser extent, only increasing from 35.7±1.2 mg/g (day 0) to 51.2±1.2

mg/g (day 10), compared to 0.3±0.1 mg/g and 45.3±5.6 mg/g for Na$^+$ in the same period (Fig.

1a; 1c). Therefore, although shoot K$^+$ increased in NIL-S, the ratio of K$^+$/Na$^+$ was significantly

higher in NIL-T leaves compared to NIL-S when measured after day 3 (Fig. 1d). Salt treatment

affected biomass production; after 10 days NaCl treatment the total dry weight of NIL-T was

significantly higher, as well as all individual dry weights of leaves, roots, and stem (Fig. 1e, f).

After establishing the timeline for shoot ion accumulation in 100 mM NaCl, to ensure

we saw differences between the NILs we chose a 4-day time point to further examine where

salt was differentially accumulated between NIL-T and NIL-S (Fig. 2, Fig. S4). We examined

the ion concentrations in a range of tissue types, and in the vascular sap. After 4 days, Na$^+$

accumulated to a greater extent in all aerial tissues in NIL-S, including leaves (both first

trifoliate leaf, FL, and youngest trifoliate leaf, YL), petioles of the first trifoliate leaf (FLP) and

youngest trifoliate leaf (YLP), higher stem (HS), lower stem (LS), and Hypocotyl (Hy), which
is consistent with what we observed for whole shoot ion content analysis. No significant
differences were observed between the NILs for Na^+ in roots under salt treatment (primary root,
PR and secondary root, SR; Fig. 2a). Under control conditions (4 days treated with RO water),
Na^+ content was significantly more, but still low, in secondary root (SR) of NIL-T compared
to NIL-S, with no significant differences in other tissues (Fig. S4). Detailed analysis of the
accumulation of K^+ under salt treatment revealed that K^+ was only significantly increased in
leaves of NIL-S and not in any other of the investigated aerial tissues or roots (Fig. 2b). No
significant differences between NIL-T and NIL-S were observed for K^+ under control
conditions (Fig. S4).

Under saline conditions, the Cl^- accumulation pattern was similar to that of Na^+
accumulation in aerial parts and hypocotyls; more Cl^- accumulated in leaves, stems and
hypocotyls of NIL-S compared to NIL-T (Fig. 2c). However, in roots (PR and SR), NIL-T
accumulated significantly more Cl^- than NIL-S (Fig. 2c). Under control conditions, Cl^- content
was significantly elevated in leaves and stem of NIL-S, with no differences for hypocotyl and
roots (Fig. S4). In addition, NO_3^- content was also measured under salt treated conditions and
accumulated more in leaves (both FL and YL) and roots in NIL-T compared to NIL-S; the ratio
of Cl^- to NO_3^- was significantly lower in NIL-T leaves compared to NIL-S, and no significant
difference was observed in roots (Fig. 2d). In summary, significantly less Na^+, K^+, and Cl^-
accumulated in NIL-T shoots than NIL-S, which in many plant species is achieved by retention
of the ions in the root to avoid accumulation in the xylem sap flow to the shoot. Xylem and
phloem sap ion concentrations were therefore determined.

Ion concentrations within soybean stem phloem and xylem exudates were examined at
the 4-day time point following NaCl treatment. Glutamine contents within the phloem sap was
used to normalize these measures to avoid differences in sap volumes leading to artefactual
results, as glutamine concentration within the phloem remains constant throughout the day
(Corbesier, Havelange, Lejeune, Bernier, & Périlleux, 2001). It was found that the contents of
Na^+ and K^+ were not significantly different between the phloem sap of NIL-T and -S plants,
but Cl^- concentration was significantly higher in NIL-T stem phloem sap (Fig. 3a). This
indicates an increased flow of Cl^- from NIL-T shoots to their roots compared to that found for
NIL-S plants; Cl^- is therefore likely to be recirculated from the stem to the roots via the phloem
in NIL-T plants. In stem xylem sap, the Na^+ concentration within NIL-S was significantly
greater compared to NIL-T, indicating less Na^+ is loaded into the NIL-T xylem – or more is
retrieved. Interestingly, in contrast to the accumulation data, the Cl^- concentrations were lower

in NIL-S xylem than NIL-T (Fig. 3b). Under control conditions (4 days with RO water), no

significant differences were observed between NIL-T and NIL-S for the ion concentrations

within stem phloem and xylem sap (Fig. S4d), which may be related to the detection levels of

the assays.

We hypothesized a link between the localization of GmSALT3 in the ER (Guan et al., 2014)

and its role in the phloem, as the ER reaches into phloem sieve elements and is directly exposed

to the phloem sap (Turgeon & Wolf, 2009). This localisation may enable a direct ion transport

from and into the sap of phloem cells and may alter the phloem morphology. In an attempt to

investigate this we examined the phloem ultrastructure of three-week old NIL-T and NIL-S

roots and stems following 100 mM NaCl treatment for four days. TEM (Transmission Electron

Microscopy) imaging did not reveal obvious morphological differences between NIL-T and

NIL-S in root phloem cells (Fig. S5), indicating lack of full-length *GmSALT3* does not result

in a disruption to sub-cellular morphology.

GmSALT3 facilitates Na$^+$, K$^+$, and Cl$^-$ transport in heterologous systems

We functionally characterised GmSALT3 in heterologous systems, including the *E.*

coli strain TK2463 (*trkAΔ, kup1Δ, kdpABCDEΔ*), which is defective in native K$^+$ uptake

mechanisms (Epstein, Buurman, McLaggan, & Naprstek, 1993) (Fig. 4) and shows a reduced

growth trajectory in media with low K$^+$ (Epstein et al., 1993). In low K$^+$-containing medium

(~2 mM K$^+$), IPTG (Isopropyl β-D-1-thiogalactopyranoside)-induced expression of full length

GmSALT3 in TK2463 improved its growth compared to bacteria expressing the truncated

Gmsalt3 and the negative control (AtCBL1n, a 12 aa peptide with the sequence

MGCFHSKAAKEF; Batistič, Waadt, Steinhorst, Held, & Kudla, 2010) (Fig. 4a; 4b).

Arabidopsis AtCHX20 (Chanroj et al., 2011) and AtKAT1 (Anderson, Huprikar, Kochian,

Lucas, & Gaber, 1992), a known plasma membrane potassium channel from Arabidopsis, were

included as positive controls, and showed similar or superior growth to the *GmSALT3*-

expressing bacteria (Fig. 4a; 4b). As we had previously used GmSALT3-YFP for determining

the subcellular localisation of GmSALT3 (Guan et al., 2014), we investigated if this

translational fusion of GmSALT3 with YFP on the C-terminus is functional and could

complement TK2463 *E. coli*. Additionally, we were interested if the last transmembrane

domain of GmSALT3 is important for functionality - truncated *Gmsalt3* from NIL-S contains

only 9 out of the 10 predicted alpha-helices. We therefore constructed GmSALT3-TM10, that

has all 10 TM alpha -helices with only the hydrophilic C-terminus removed. Both, YFP-tagged

GmSALT3 and GmSALT3-TM10 increased *E. coli* growth to that of GmSALT3 and
significantly greater than that of Gmsalt3 (Fig. 4a; 4b). None of the test constructs or controls
used had a significant impact on bacterial growth rates on media high K^+ (Fig. S6); on media
with low K^+ and no IPTG GmSALT3 did not show any significant growth differences from
Gmsalt3 which was no different from the negative control (Fig. S6). In summary, bacterial
growth assays suggest that in addition to GmSALT3, YFP-tagged GmSALT3 and GmSALT3-
TM10 are functional. This indicates all transmembrane domains are important for GmSALT3
transport function, but the large (383 aa) hydrophilic C-terminus can be truncated or added to
without abolishing K^+ transport.

As the presence of functional GmSALT3 in soybean causes shoot ion exclusion for Na^+,
Cl^- and K^+, we were interested to see if the presence of the protein in cells influences the flux
of all three ions across the PM. To investigate this, we expressed YFP-tagged and untagged
GmSALT3 in *X. laevis* oocytes. Although GmSALT3 is localised to the ER in plant cells and
no PM-localisation was observed (Guan et al., 2014), we detected fluorescence at the PM in
GmSALT3-YFP expressing oocytes, similar to the fluorescence of oocytes expressing the PM-
localised positive control *TmHKT1;5-A*-YFP (Fig. 5a) (*Triticum monococcum* High-Affinity
K^+ Transporter 1; Munns et al., 2012). It is not uncommon for plant endomembrane proteins
to at least partially localise to the PM in frog oocytes (Maurel, Reizer, Schroeder, & Chrispeels,
1993). Two electrode voltage-clamp electrophysiology revealed that the resting membrane
potential of *GmSALT3*-injected oocytes was more positive compared to H_2O-injected oocytes
when incubated in ND96 media (Fig. 5b), but no consistent current differences were detected.
Therefore, we chose to conduct MIFE (Microelectrode Ion Flux Estimation) experiments to
measure ion fluxes. MIFE results indicated that in *GmSALT3*-injected oocytes net efflux of
Na^+, K^+, and Cl^- was reduced compared to H_2O-injected oocytes (Fig. 5c). Ion content analysis
of oocytes revealed that, in agreement with the reduced net ion efflux in oocytes, *GmSALT3*-
injected oocytes contained significantly more K^+, Na^+, and Cl^- compared to H_2O-injected
oocytes (Fig. 5d). To test whether GmSALT3 induced differences in ion concentration could
be blocked in oocytes, we used the Na^+ channel and Na^+/H^+ exchanger inhibitor amiloride
hydrochloride (Darley et al., 2000). Oocytes were transferred from ND96 to BSM (low Na^+,
K^+ and Cl^-) and incubated overnight. Oocytes incubated without amiloride, showed a smaller
relative K^+, Na^+, and Cl^- content decrease in *GmSALT3*-injected oocytes compared to H_2O-
injected oocytes as expected (Fig. 5d). However, no difference was observed between control
and *GmSALT3*-injected oocytes when the Na^+ transporter inhibitor amiloride was present (Fig.
5d), confirming GmSALT3 could affect transport of K^+, Na^+, and Cl^- across the PM in oocytes.

Discussion

The effect of GmSALT3 on ion accumulation *in planta* following salt treatment

Salt resistance includes tolerance to elevated levels of both Cl^- and Na^+. In many plant species one of the two ions is more deleterious to the plant, however, the situation for soybean is currently unclear with evidence for both ions being more toxic than the other (Läuchli, 1984). Whilst our work does not answer which ion causes the most damage to soybean, it does clearly demonstrate that Na^+ and Cl^- long distance transport, and its accumulation in shoots, is uncoupled in soybean. It is clear that Na^+ and Cl^- long distance transport is under the control of distinct mechanisms even if the same gene appears to impact the transport of both ions. We gathered the evidence to demonstrate this when we treated plants with 100 mM NaCl to mimic a more physiologically relevant salt treatment than has previously been used on NIL plants (i.e. one that would be found in the field). We confirmed in these conditions that shoot exclusion of Cl^- broke down prior to exclusion of Na^+ in NIL-S when compared with NIL-S plants (Fig. 1).

Na^+ and Cl^- are transported to shoots via the xylem transpiration stream. Under saline conditions, Na^+ was accumulated to greater concentrations in the aerial parts of NIL-S but no difference was observed in the roots compared to NIL-T (Fig. 2b); in stem xylem sap, there was a lower Na^+ concentration in NIL-T compared to NIL-S but no significant difference in stem phloem sap (Fig. 3). This indicates more Na^+ is transported in the NIL-S xylem stream. As such net loading of Na^+ to NIL-T xylem is likely to be restricted compared to NIL-S plants. As for Cl^-, more Cl^- is transported to the shoots in NIL-T xylem, but it clearly does not stay in there, instead, the higher phloem sap concentrations indicate that Cl^- most likely is recirculated to the root (Fig 3). *GmSALT3* might therefore impact Cl^- distribution in the plant via phloem recirculation of Cl^-, resulting eventually in higher Cl^- accumulation in NIL-T roots. In addition, when irrigated with RO (Reverse osmosis) water, NIL-S plants had significantly higher accumulation of Cl^- in aerial tissues (leaves and stems) than NIL-T plants, indicating an intrinsic difference in net loading of the shoot even under non-saline conditions (Fig. S4). No differences in phloem Cl^- concentration or shoot Na^+ content could be detected under these control conditions which is possibly the result of the low signal for each parameter and differences might fall under the detection limit of the method (Fig. S4). These data clearly show there are distinct mechanisms operating for the control of long-distance transport of Na^+ and Cl^-.

NO$_3^-$ has previously been shown to be an effective competitor of Cl$^-$ uptake to alleviate
ion toxicities in soybean under salt stress, indicating that leaf Cl$^-$/NO$_3^-$ ratio may be a
significant index for evaluating salt tolerance, as with K$^+$/Na$^+$ (Guo, Zhou, Li, Yu, & Luo,
2017). After salt treatment (100 mM for 4 days), NIL-T had a much lower Cl$^-$/NO$_3^-$ ratio in
leaves than NIL-S (Fig. 2d), indicating that NIL-T may reduce potential chloride toxicity in
shoots by maintaining a low Cl$^-$/NO$_3^-$ ratio (Li et al., 2017; Wege, Gilliham & Henderson,
2017).

In NIL-S, without GmSALT3, K$^+$ concentration in shoots increased during the salt
treatment (simultaneously with Na$^+$), this did not occur in NIL-T (Fig. 1c), and might be a
compensatory effect in NIL-S to maintain a higher K$^+$/Na$^+$ ratio. Interestingly, NIL-T leaves
maintained a constant K$^+$ concentration throughout ten days of salt treatment even though Na$^+$
and Cl$^-$ increased after salt treatment (Fig. 1). No significant difference could be detected for
K$^+$ concentration between roots of NIL-T and NIL-S plants (Fig. 2c). This indicates that
GmSALT3 also affects shoot K$^+$ accumulation in soybean.

Transport function of GmSALT3

Based on the ER-localisation of GmSALT3 (Guan et al., 2014) and the complex and
distinct processes underlying the different salt ion accumulation phenotypes of the two NILs,
we propose that GmSALT3 affects processes at the ER and is not directly mediating ion
transport across the PM or tonoplast itself – this makes it unlike most other transport proteins
identified as important for salt tolerance.

A close homolog of *GmSALT3*, *AtCHX20* in *Arabidopsis thaliana*, is preferentially
expressed in stomatal guard cells, and also localises to the ER like GmSALT3, with an ER-
localised GFP-tagged version complementing the *Atchx20* knockout phenotype (Padmanaban
et al., 2007). *AtCHX20* was shown to enhance growth of *E. coli* deficient in K$^+$ uptake (LB2003)
(Chanroj et al., 2011). We confirmed this result with a different K$^+$ transport deficient *E. coli*
strain (TK2463), and showed that GmSALT3 had a similar effect on *E. coli* growth as
AtCHX20.

Plant CPA2 (Cation-Proton Antiporter 2)/CHX (Cation/H$^+$ Exchangers) have not been
characterised in *X. laevis* oocytes before, different from some CPA2 members from the animal
field. Two *Drosophila* CPA2 transporters, NHA1 and NHA2, have been suggested to act as a
Na$^+$/H$^+$ exchanger and a H$^+$/Cl$^-$ cotransporter, respectively (Chintapalli et al., 2015), suggesting
that proteins from this family can putatively act as either, cation or anion transporters. Unlike
in planta, GmSALT3 targeted, at least partially, to the PM of *X. laevis* oocytes (Fig. 5a).

GmSALT3 protein expression led to a depolarization of the membrane potential, similar to

what was reported for NHA injected oocytes (Chintapalli et al., 2015). We also observed

GmSALT3 expression in oocytes led to a lower net ion efflux of all three ions, K^+, Na^+ and Cl^-

(Fig. 5c). Interestingly, the reduced efflux effect was inhibited with the Na^+ channel and Na^+/H^+

exchanger inhibitor amiloride hydrochloride. Whilst this inhibitor is thought to be selective for

Na^+ transporters, it had an impact on the efflux rate of all three ions measured, Na^+, K^+ and Cl^-.

It is possible that observed differences in ion flux and accumulation in heterologous systems

are the result of changing cytosolic ion concentration which then impacts ion gradients across

the PM and influences the activity of native ion transporters. In such a scenario the fluxes

detected across the PM of oocytes is the combination of GmSALT3 and native transporters;

and the inhibition of one component leads to the inhibition of the entire mechanism.

Putative *in planta* mechanisms for GmSALT3 regulating long distance ion transport

We propose that *in planta* ER-localised GmSALT3 influences ion gradients across the

PM through currently unknown downstream effects. This might be through impacting the

activity of PM-localised transporters, by changing their PM-abundance or changing cytosolic

ion supply, or even more complex effects. For example, if the ionic milieu within the ER is

affected this could impact the function or sorting of other membrane transporters or regulatory

proteins, which then impacts PM fluxes. Such a mechanism has been suggested for prevacuolar

compartment (PVC) localised AtCHX17 and ER localised AtCHX20 (Chanroj et al., 2011).

The complexity of this novel salt tolerance mechanisms is further highlighted by the previous

observation that root GmSALT3 is sufficient to mediate shoot salt exclusion, using NIL-T and

NIL-S grafts (Guan et al., 2014). Further experiments will need to be carried out to decipher

the novel mechanism imparted by GmSALT3 that is important for salt tolerance. This might

enable us to identify similar mechanisms in other (crop) plants.

Phloem translocation of Cl^- appears to be conferred by GmSALT3. Phloem

retranslocation of NaCl in saline conditions has been observed before, for example, it was

found that in maize approximately 13-36% of the Na^+ and Cl^- imported to leaves through the

xylem was retranslocated to the root again in the phloem (Lohaus et al., 2000). However, no

transport protein could be linked to this phenomenon. Until now, no reports have shown

phloem recirculation of Cl^- to be an important salt tolerance mechanism; *GmSALT3* presents

as a very promising candidate for impacting Cl^- long-distance re-translocation.

Conclusion

To summarize, this work has provided further insights into the salinity tolerance mechanisms imparted by GmSALT3 *in planta* and in heterologous systems. We propose that, in NIL-T, the presence of full-length *GmSALT3* mediates Na^+ and Cl^- exclusion from shoots through restricting net xylem loading of Na^+, while Cl^- is retranslocated from the shoots back into the roots via the phloem. In heterologous systems, GmSALT3 is involved in K^+ uptake in *E. coli* cells, and its expression in *X. laevis* oocytes mediates net import of Na^+, K^+, and Cl^-. However, as it is an endomembrane-localised protein, how GmSALT3 contributes to soybean salinity tolerance mechanisms is still not clear, and how its expression directly impacts salt movement to and from shoots is unresolved. As the heterologous expression of *GmSALT3* led to transport differences in all three ions that are affected in distribution in the plant it is tempting to speculate that GmSALT3 may be actively involved in their distribution in tissues. However, as the effect on Cl^- and Na^+ in the xylem and phloem differs, the impact of other processes including other transport proteins are likely to be involved in the NIL phenotypes. RNA-sequencing using NIL-T and NIL-S roots may be beneficial in investigating if GmSALT3 confers salinity tolerance in soybean roots via influencing transcription, and what distinctive pathways and genes have been significantly changed in NIL-T and NIL-S roots under saline conditions.

Acknowledgements

We thank Yue (Crystal) Wu for generating the *AtCBL1n* entry vector; Sunita Ramesh and Zhonghua Chen for providing *AtKAT1* construct used in this study; CSIRO, Australia for donating *E. coli* strain TK2463. We thank Adelaide Microscopy, especially Gwenda Mayo for the use of TEM and other imaging facilities. This work was funded by the Natural Science Foundation of China, grant number 31830066 (RG). YQ and SH were supported by ARC Centre of Excellence funding awarded to MG (CE140100008). ARC Fellowships supported MG (FT130100709), SW (DE160100804) and JB (DE170100346).

Author contributions

MG, YQ and SW designed experiments, with input from LQ and RG. LQ and MG directed the project. YQ performed ion accumulation measurement, TEM, IRGA, *E. coli* and oocyte expression, analysed the data and drafted the manuscript. JB performed MIFE in oocytes. SH assisted with experimental design and provided the *E. coli* expression vectors. YQ, MG and SW edited the manuscript. All authors commented on the manuscript.

References

Abel G. (1969). Inheritance of the capacity for chloride inclusion and chloride exclusion by soybeans. *Crop Science*, 9, 697-698.

Alexandratos N. & Bruinsma J. (2012). World agriculture towards 2030/2050: the 2012 revision: ESA Working paper No.12-03. Rome, FAO.

Anderson J.A., Huprikar S.S., Kochian L.V., Lucas W.J. & Gaber R.F. (1992). Functional expression of a probable *Arabidopsis thaliana* potassium channel in *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences*, 89, 3736-3740.

Batistič O., Waadt R., Steinhorst L., Held K. & Kudla J. (2010). CBL-mediated targeting of CIPKs facilitates the decoding of calcium signals emanating from distinct cellular stores. *The Plant Journal*, 61, 211-222.

Berthomieu P., Conéjéro G., Nublat A., Brackenbury W.J., Lambert C., Savio C., Uozumi N., Oiki S., Yamada K. & Cellier F. (2003). Functional analysis of AtHKT1 in Arabidopsis shows that Na^+ recirculation by the phloem is crucial for salt tolerance. *The EMBO Journal*, 22, 2004-2014.

Cao D., Li Y., Liu B., Kong F., & Tran L.S.P. (2018). Adaptive mechanisms of soybean grown on salt-affected soils. *Land degradation & development*, 29(4), 1054-1064.

Chang R.Z., Chen Y.W., Shao G.H., Wan C.W. (1994). Effect of salt stress on agronomic characters and chemical quality of seeds in soybean. *Soybean Science*, 13, 101-105.

Chanroj S., Lu Y., Padmanaban S., Nanatani K., Uozumi N., Rao R. & Sze H. (2011). Plant-specific cation/H^+ exchanger 17 and its homologs are endomembrane K^+ transporters with roles in protein sorting. *Journal of Biological Chemistry*, 286, 33931-33941.

Chintapalli V.R., Kato A., Henderson L., Hirata T., Woods D.J., Overend G., Davies S.A., Romero M.F. & Dow J.A. (2015). Transport proteins NHA1 and NHA2 are essential for survival, but have distinct transport modalities. *Proceedings of the National Academy of Sciences*, 112, 11720-11725.

Corbesier L., Havelange A., Lejeune P., Bernier G. & Périlleux C. (2001). N content of phloem and xylem exudates during the transition to flowering in *Sinapis alba* and *Arabidopsis thaliana*. *Plant, Cell & Environment*, 24, 367-375.

Czerny D.D., Padmanaban S., Anishkin A., Venema K., Riaz Z. & Sze H. (2016). Protein architecture and core residues in unwound α-helices provide insights to the function of a plant transporter AtCHX17. *Biochimica et Biophysica Acta (BBA)-Biomembranes,* 1858, 1983-1998.

Do T.D., Chen H., Hien V.T.T., Hamwieh A., Yamada T., Sato T., Yan Y., Cong H., Shono
M. & Suenaga K. (2016). Ncl synchronously regulates Na^+, K^+, and Cl^- in soybean and
greatly increases the grain yield in saline field conditions. *Scientific Reports*, 6, 19147.

Epstein W., Buurman E., McLaggan D. & Naprstek J. (1993). Multiple mechanisms, roles and
controls of K^+ transport in *Escherichia coli*: Portland Press Limited.

FAO. (2017). *The future of food and agriculture – Trends and challenges*, Rome: FAO.

Gilliham M., Able J.A. & Roy S.J. (2017). Translating knowledge about abiotic stress tolerance
to breeding programmes. *The Plant Journal*. 90, 898-917.

Guan R., Qu Y., Guo Y., Yu L., Liu Y., Jiang J., Chen J., Ren Y., Liu G. & Tian L. (2014).
Salinity tolerance in soybean is modulated by natural variation in *GmSALT3*. *The Plant
Journal*, 80, 937-950.

Guo J., Zhou Q., Li X., Yu B. & Luo Q. (2017). Enhancing NO_3^- supply confers NaCl tolerance
by adjusting Cl^- uptake and transport in *G. max & G. soja*. *Journal of Soil Science and
Plant Nutrition*, 17, 194-202.

Ha B.K., Vuong T.D., Velusamy V., Nguyen H.T., Shannon J.G. & Lee J.D. (2013). Genetic
mapping of quantitative trait loci conditioning salt tolerance in wild soybean (*Glycine
soja*) PI 483463. *Euphytica*. 193, 79-88.

Hamwieh A., Tuyen D.D., Cong H., Benitez E.R., Takahashi R. & Xu, D.H. (2011).
Identification and validation of a major QTL for salt tolerance in soybean. *Euphytica,
179:,451-459.

Hamwieh A. & Xu D. (2008). Conserved salt tolerance quantitative trait locus QTL) in wild
and cultivated soybeans. *Breeding Science*, 58, 355-359.

Ikeda M. (2005). Distribution of K^+, Na^+ and Cl^- in root and leaf cells of soybean and cucumber
plants grown under salinity conditions. *Soil Science and Plant Nutrition*, 51, 1053-1057.

Lee G.J., Carter J.T.E., Villagarcia M.R., Li Z., Zhou X., Gibbs M.O. & Boerma H.R. (2004).
A major QTL conditioning salt tolerance in S-100 soybean and descendent cultivars.
Theoretical and Applied Genetics, 109, 1610-1619.

Lenis J., Ellersieck M., Blevins D., Sleper D., Nguyen H., Dunn D., Lee J. & Shannon J. (2011).
Differences in ion accumulation and salt tolerance among *Glycine* accessions. *Journal
of Agronomy and Crop Science*. 197, 302-310.

Li B., Tester M. & Gilliham M. (2017). Chloride on the Move. *Trends in Plant Science*, 22,
236-248.

Li W.Y., Wong F.L., Tsai S.N., Phang T.H., Shao G. & Lam H.M. (2006a). Tonoplast-located
GmCLC1 and *GmNHX1* from soybean enhance NaCl tolerance in transgenic bright
yellow (BY)-2 cells. *Plant Cell & Environment*, 29, 1122-1137.

Li X.J., An P., Inanaga S., Eneji A.E. & Tanabe K. (2006b). Salinity and defoliation effects on
soybean growth. *Journal of Plant Nutrition*, 29, 1499-1508.

Liu Y., Yu L., Qu Y., Chen J., Liu X., Hong H., Liu Z., Chang R., Gilliham M., Qiu L. & Guan
R. (2016). GmSALT3, which confers improved soybean salt tolerance in the field,
increases leaf Cl^- exclusion prior to Na^+ exclusion but does not improve early vigor
under salinity. *Frontiers in Plant Science*, 7, 1485

Lohaus G., Hussmann M., Pennewiss K., Schneider H., Zhu J.J. & Sattelmacher B. (2000).
Solute balance of a maize (*Zea mays* L.) source leaf as affected by salt treatment with
special emphasis on phloem retranslocation and ion leaching. *Journal of Experimental
Botany*, 51, 1721-1732.

Luo G.Z., Wang H.W., Huang J., Tian A.G., Wang Y.J., Zhang J.S. & Chen S.Y. (2005a). A
putative plasma membrane cation/proton antiporter from soybean confers salt tolerance
in Arabidopsis. *Plant Molecular Biology*, 59, 809-820.

Luo Q., Yu B. & Liu Y. (2005b). Differential sensitivity to chloride and sodium ions in
seedlings of *Glycine max* and *G. soja* under NaCl stress. *Journal of Plant Physiology*,
162, 1003-1012.

Läuchli A. (1984). Salt exclusion: An adaptation of legumes for crops and pastures under saline
conditions. In *Salinity Tolerance in Plants: Strageiges for Crop Improvement* (Staples,
R.C. and Toenniessen, G.H. eds). New York: Wiley, pp. 171-187.

Maurel C., Reizer J., Schroeder J.I. & Chrispeels M.J.. (1993). The vacuolar membrane protein
gamma-TIP creates water specific channels in *Xenopus oocytes*. *The EMBO Journal*,
12, 2241-2247.

Munns R. & Gilliham M. (2015). Salinity tolerance of crops–what is the cost? *New Phytologist*,
208, 668-673.

Munns R., James R.A., Xu B., Athman A., Conn S.J., Jordans C., Byrt C.S., Hare R.A.,
Tyerman S.D. & Tester M. (2012). Wheat grain yield on saline soils is improved by an
ancestral Na^+ transporter gene. *Nature Biotechnology*, 30, 360-364.

Munns R. & Tester M. (2008). Mechanisms of salinity tolerance. *Annural Review Plant
Biology*, 59, 651-681.

Møller I.S., Gilliham M., Jha D., Mayo G.M., Roy S.J., Coates J.C., Haseloff J. & Tester M.
(2009). Shoot Na^+ exclusion and increased salinity tolerance engineered by cell type-
specific alteration of Na^+ transport in *Arabidopsis*. *Plant Cell*, 21, 2163-2178.

Obermeyer G. & Tyerman S.D. (2005). NH_4^+ currents across the peribacteroid membrane of
soybean. Macroscopic and microscopic properties, inhibition by Mg^{2+}, and temperature
dependence indicate a SubpicoSiemens channel finely regulated by divalent cations.
Plant Physiology, 139, 1015-1029.

Padmanaban S., Chanroj S., Kwak J.M., Li X., Ward J.M. & Sze H. (2007). Participation of
endomembrane cation/H^+ exchanger *AtCHX20* in osmoregulation of guard cells. *Plant*
Physiology, 144, 82-93.

Phang T.H., Shao G. & Lam H.M. (2008). Salt tolerance in soybean. *Journal of Integrative*
Plant Biology, 50, 1196-1212.

Phang T.H., Shao G., Liao H., Yan X. & Lam H.M. (2009). High external phosphate (Pi)
increases sodium ion uptake and reduces salt tolerance of 'Pi-tolerant'soybean.
Physiologia Plantarum, 135, 412-425.

Qi X., Li M.W., Xie M., Liu X., Ni M., Shao G., Song C., Yim A.K.Y., Tao Y., Wong F.L. &
Isobe S. (2014). Identification of a novel salt tolerance gene in wild soybean by whole-
genome sequencing. *Nature Communications*, 5, 4340.

Qiu J., Henderson S.W., Tester M., Roy S.J. & Gilliham M. (2016). SLAH1, a homologue of
the slow type anion channel SLAC1, modulates shoot Cl^- accumulation and salt
tolerance in *Arabidopsis thaliana*. *Journal of Experimental Botany*, 67, 4495-4505.

Ren Z.H., Gao J.P., Li L.G., Cai X.L., Huang W., Chao D.Y., Zhu M.Z., Wang Z.Y., Luan S.
& Lin H.X. (2005). A rice quantitative trait locus for salt tolerance encodes a sodium
transporter. *Nature Genetics*, 37, 1141-1146.

Roy S.J., Negrão S. & Tester M. (2014). Salt resistant crop plants. *Current Opinion in*
Biotechnology, 26, 115-124.

Rupassara S.I. (2008). *Metabolite profiling of leaves and vascular exudates in soybean grown*
under free-air concentration enrichment. University of Illinois at Urbana-Champaign.

Rus A., Baxter I., Muthukumar B., Gustin J., Lahner B., Yakubova E. & Salt D.E. (2006).
Natural variants of AtHKT1 enhance Na^+ accumulation in two wild populations of
Arabidopsis. *PLoS Genetics*, 2, 210.

Shabala S., Shabala L., Bose J., Cuin T. & Newman I. (2013). Ion flux measurements using
the MIFE technique. *Plant Mineral Nutrients: Methods and Protocols*, 171-183.

Shao G.H., Wan C.W., Chang R.Z. & Chen Y.W. (1993). Preliminary study on the damage of
plasma membrane caused by salt stress. *Crops,* 1, 39-40.

Shi H., Bressan R.. (2006). RNA extraction. *Methods in Molecular Biology* 323: 345.

Singh G. (2010). *The soybean: botany, production and uses* Oxfordshire, UK: CABI
Publishing.

Sze H. & Chanroj S. (2018). Plant endomembrane dynamics: Studies of K^+/H^+ antiporters
provide insights on the effects of pH and ion homeostasis. *Plant Physiology*, 177:875-
95.

Tester M. & Davenport R. (2003). Na^+ tolerance and Na^+ transport in higher plants. *Annals of*
Botany, 91, 503-527.

Tester M. & Langridge P. (2010). Breeding technologies to increase crop production in a
changing world. *Science*, 327, 818-822.

Tian H., Baxter I.R., Lahner B., Reinders A., Salt D.E. & Ward J.M. (2010). Arabidopsis
NPCC6/NaKR1 is a phloem mobile metal binding protein necessary for phloem
function and root meristem maintenance. *The Plant Cell*, 22, 3963-3979.

Turgeon R. & Wolf S. (2009). Phloem transport: cellular pathways and molecular trafficking.
In *Annual Review of Plant Biol*ogy, 60, 207-221.

Uozumi N., Kim E.J., Rubio F., Yamaguchi T., Muto S., Tsuboi A., Bakker E.P., Nakamura T.
& Schroeder J.I. (2000). The Arabidopsis *HKT1* gene homolog mediates inward Na^+
currents in *Xenopus laevis* oocytes and Na^+ uptake in *Saccharomyces cerevisiae. Plant*
Physiology, 122, 1249-1260.

Wege S., Gilliham M. & Henderson, S.W. (2017). Chloride: not simply a 'cheap osmoticum',
but a beneficial plant macronutrient. *Journal of Experimental Botany*, 68, 3057-3069.

Xu G.H., Magen H., Tarchitzky J. & Kafkafi U. (2000). Advances in chloride nutrition of
plants. San Diego: Academic Press Inc., pp. 97-150.

Figure Legends

Figure 1 Time-course ion concentration (mg DW^{-1}) in NIL-T and NIL-S leaves during 10 days 100 mmol L^{-1} NaCl stress. Na$^+$ content (**a**), Cl$^-$ content (**b**), K$^+$ (**c**) content, and K$^+$ to Na$^+$ ratio (**d**) in leaves of NIL-T (blue data) and NIL-S (red data). Dry weight of different tissues (**e**) and all tissues (**f**). Observations are means of four replicates ± SEM. Asterisks indicates a significant difference between NIL-T and NIL-S at *p <0.05, **p < 0.01 according to the LSD test. DW, dry weight.

Figure 2 Ion concentration in different tissues of NIL-T and -S after 4 days of 100 mmol L^{-1} NaCl treatment. (**a**) Tissue content of Na$^+$ in leaf, hypocotyl and root of NILs. (**b**) K$^+$ content in leaf, hypocotyl and root of NILs. (**c**) Cl$^-$ content in leaf, hypocotyl and root of NILs. (**d**) NO^{3-} content, Cl$^-$ content and Cl$^-$/NO^{3-} ratio in leaf and root of NILs. Ion contents are expressed as tissue dry weight. Data are means of 3-5 replicates ± SE. Asterisks indicate a significant difference between NIL-T (blue) and NIL-S (red) at *p <0.05, **p < 0.01 according to the LSD test (ion content data in control plants with water supply can be found in Figure S4). FL, first trifoliate leaves; YL, youngest trifoliate leaves; FLP, first trifoliate leaves petiole; YLP, youngest trifoliate leaves petiole; HS, higher stem; LS, lower stem; PR, primary root; SR, secondary root; Hy, hypocotyl. Data shown in (**a**), (**b**), and (**c**) were from one of the two independent experiments with similar results. Data shown in (**d**) were from a separate experiment and repeated twice.

Figure 3 Ion concentration in NIL-T and -S stem phloem and xylem sap after 4 days of 100 mmol L^{-1} NaCl stress. (**a**) Na$^+$, K$^+$, or Cl$^-$ content to glutamine ratio in NIL-T and NIL-S stem phloem sap. (**b**) Na$^+$, K$^+$, and Cl$^-$ concentrations in NIL-T (blue) and NIL-S (red) stem xylem sap. Data are means of three replicates ± SEM. Asterisks indicate a significant difference between NIL 820-T and 820-S at *p <0.05, **p < 0.01 according to the LSD test. SPS, stem phloem sap; SXS, stem xylem sap. Data shown here were from one of the three independent experiments with similar results.

Figure 4 Functional test of *GmSALT3* in *E. coli*. *E. coli* strain TK2463 (defective in potassium uptake, Trk, Kup, and Kdp) harbouring a modified pET-DEST42 vector with *GmSALT3, Gmsalt3, CBL1n, GmSALT3-YFP, GmSALT3-TM10, AtCHX20 and AtKAT1* were grown in different media. (**a**) Low potassium medium (10 g/L tryptone, 2 g/L yeast extract, 100 mM D-mannitol; ~2 mM K$^+$) with additional 0.5 mM Isopropyl β- D-1-thiogalactopyranoside (IPTG). All the test media were supplemented with 100 μg/ml carbenicillin. OD$_{600}$ was monitored every 15 min in 96-well microplate reader. Bars are mean ± SEM of 3 replicates. Growth rates (OD$_{600}$ per hour) of *E.coli* cells within the log phase are

indicated between dotted lines, and their slopes are shown in (**b**) (low potassium medium with

IPTG). Different letters indicate statistically significant differences between *E. coli* cells

harbouring different constructs (one-way ANOVA followed by Fisher's LSD test, $p < 0.05$).

Figure 5 Functional characterization of *GmSALT3* in *X. laevis* oocytes. (a) Plasma

membrane localisation of expressed *GmSALT3-YFP* in oocytes. H_2O-injected and *TmHKT1;5-*

A-YFP (positive control) expressed oocytes are also shown. Scale bar = 0.5 mm. Oocytes were

analysed by confocal laser-scanning microscopy. (**b**) Resting membrane potentials of

GmSALT3- and H_2O-injected oocytes after incubation in ND96 for 72 hours. Values are means

± SEM (n = 8), Asterisks indicate a significant difference between GmSALT3- and H_2O-

injected oocytes at *$p < 0.01$ according to the LSD test. (**c**) Net ion fluxes were measured on

oocytes plasma membrane using MIFE (Microelectrode Ion Flux Estimation). Oocytes were

incubated in ND96 for 72 hours, and washed in ND96 in (96 mM NaCl, 1 mM KCl, 1 mM

$MgCl_2$, 5 mM HEPES, pH 7.5). Measurements were done in BSM (Basal Salty Medium; 5 mM

NaCl, 0.2 mM KCl, 0.2 mM $CaCl_2$, 5 mM HEPES, pH 7.5) and exchanged after time zero to

BSM with sufficiently low concentrations of Na^+, K^+ and Cl^- to ensure a gradient for efflux for

target ions and an improved signal:noise ratio. Values are means ± SEM (n = 4 - 6). (**d**) Ion

(Na^+, K^+, and Cl^-) concentrations in *GmSALT3* (blue) and H_2O (Red) injected oocytes after

incubation in ND96 (72 hours), BSM (overnight), BSM with 100 μM amiloride (overnight).

Values are means ± SEM (n = 4-7). Asterisks indicate a significant difference between

GmSALT3-injected and H_2O-injected oocytes at *$p < 0.05$, **$p < 0.01$ according to the LSD

test.

Supporting Information

Fig. S1. Time-course ion concentration (Na$^+$) in NIL 820-T and 820-S leaves, stems,
hypocotyls, and roots during 10 days NaCl stress.

Fig. S2. Time-course ion concentration (K$^+$) in NIL 820-T and 820-S leaves, stems,
hypocotyls, and roots during 10 days NaCl stress.

Fig. S3. Time-course ion concentration (Cl$^-$) in NIL 820-T and 820-S leaves, stems,
hypocotyls, and roots during 10 days NaCl stress.

Fig. S4. Ion content and ion flux in NIL 820-T and 820-S different tissues after 4 days in
control conditions.

Fig. S5. Ultrastructure of NIL-*GmSALT3* and NIL-*Gmsalt3* root cross sections with 4 days
NaCl stress.

Fig. S6. Functional test of *GmSALT3* in *E. coli* with low potassium and KML medium.

Table S1. Primers used in this study. Primers were designed and adjusted using Primer3.

Figures

Figure 1 Time-course ion concentration (mg DW^{-1}) in NIL-T and NIL-S leaves during 10 days 100 mmol L^{-1} NaCl stress. Na$^+$ content (**a**), Cl$^-$ content (**b**), K$^+$ (**c**) content, and K$^+$ to Na$^+$ ratio (**d**) in leaves of NIL-T (blue data) and NIL-S (red data). Dry weight of different tissues (**e**) and all tissues (**f**). Observations are means of four replicates ± SEM. Asterisks indicates a significant difference between NIL-T and NIL-S at *$p < 0.05$, **$p < 0.01$ according to the LSD test. DW, dry weight.

Figure 2 Ion concentration in different tissues of NIL-T and -S after 4 days of 100 mmol L^{-1} NaCl treatment. (a) Tissue content of Na$^+$ in leaf, hypocotyl and root of NILs. **(b)** K$^+$ content in leaf, hypocotyl and root of NILs. **(c)** Cl$^-$ content in leaf, hypocotyl and root of NILs. **(d)** NO$_3^-$ content, Cl$^-$ content and Cl$^-$/NO$_3^-$ ratio in leaf and root of NILs. Ion contents are expressed as tissue dry weight. Data are means of 3-5 replicates ± SE. Asterisks indicate a significant difference between NIL-T (blue) and NIL-S (red) at *p <0.05, **p < 0.01 according to the LSD test (ion content data in control plants with water supply can be found in Figure S4). FL, first trifoliate leaves; YL, youngest trifoliate leaves; FLP, first trifoliate leaves petiole; YLP, youngest trifoliate leaves petiole; HS, higher stem; LS, lower stem; PR, primary root; SR, secondary root; Hy, hypocotyl. Data shown in **(a)**, **(b)**, and **(c)** were from one of the two independent experiments with similar results. Data shown in **(d)** were from a separate experiment and repeated twice.

Figure 3 Ion concentration in NIL-T and -S stem phloem and xylem sap after 4 days of
100 mmol L^{-1} NaCl stress. (a) Na$^+$, K$^+$, or Cl$^-$ content to glutamine ratio in NIL-T and NIL-S
stem phloem sap. **(b)** Na$^+$, K$^+$, and Cl$^-$ concentrations in NIL-T (blue) and NIL-S (red) stem
xylem sap. Data are means of three replicates ± SEM. Asterisks indicate a significant difference
between NIL 820-T and 820-S at *p <0.05, **p < 0.01 according to the LSD test. SPS, stem
phloem sap; SXS, stem xylem sap. Data shown here were from one of the three independent
experiments with similar results.

Figure 4 Functional test of *GmSALT3* in *E. coli*. *E. coli* strain TK2463 (defective in

potassium uptake, Trk, Kup, and Kdp) harbouring a modified pET-DEST42 vector with

GmSALT3, Gmsalt3, CBL1n, GmSALT3-YFP, GmSALT3-TM10, AtCHX20 and AtKAT1 were

grown in different media. (**a**) Low potassium medium (10 g/L tryptone, 2 g/L yeast extract,

100 mM D-mannitol; ~2 mM K^+) with additional 0.5 mM Isopropyl β- D-1-

thiogalactopyranoside (IPTG). All the test media were supplemented with 100 μg/ml

carbenicillin. OD_{600} was monitored every 15 min in 96-well microplate reader. Bars are mean

± SEM of 3 replicates. Growth rates (OD_{600} per hour) of *E.coli* cells within the log phase are

indicated between dotted lines, and their slopes are shown in (**b**) (low potassium medium with

IPTG). Different letters indicate statistically significant differences between *E. coli* cells

harbouring different constructs (one-way ANOVA followed by Fisher's LSD test, $p < 0.05$).

Figure 5 Functional characterization of *GmSALT3* in *X. laevis* oocytes. (a) Plasma
membrane localisation of expressed *GmSALT3-YFP* in oocytes. H_2O-injected and *TmHKT1;5-
A*-YFP (positive control) expressed oocytes are also shown. Scale bar = 0.5 mm. Oocytes were
analysed by confocal laser-scanning microscopy. (**b**) Resting membrane potentials of
GmSALT3- and H_2O-injected oocytes after incubation in ND96 for 72 hours. Values are means
± SEM (n = 8), Asterisks indicate a significant difference between GmSALT3- and H_2O-
injected oocytes at *$p < 0.01$ according to the LSD test. (**c**) Net ion fluxes were measured on
oocytes plasma membrane using MIFE (Microelectrode Ion Flux Estimation). Oocytes were
incubated in ND96 for 72 hours, and washed in ND96 in (96 mM NaCl, 1 mM KCl, 1 mM
$MgCl_2$, 5 mM HEPES, pH 7.5). Measurements were done in BSM (Basal Salty Medium; 5 mM
NaCl, 0.2 mM KCl, 0.2 mM $CaCl_2$, 5 mM HEPES, pH 7.5) and exchanged after time zero to
BSM with sufficiently low concentrations of Na^+, K^+ and Cl^- to ensure a gradient for efflux for

target ions and an improved signal:noise ratio. Values are means ± SEM (n = 4 - 6). **(d)** Ion

(Na^+, K^+, and Cl^-) concentrations in *GmSALT3* (blue) and H_2O (Red) injected oocytes after

incubation in ND96 (72 hours), BSM (overnight), BSM with 100 μM amiloride (overnight).

Values are means ± SEM (n = 4-7). Asterisks indicate a significant difference between

GmSALT3-injected and H_2O-injected oocytes at *p <0.05, **p < 0.01 according to the LSD

test.