Whole genome comparisons of *Staphylococcus agnetis* isolates from cattle and chickens.

Abdulkarim Shwani,[a] Pamela R. F. Adkins,[b] Nnamdi S. Ekesi,[a] Adnan Alrubaye,[a] Michael J.

Calcutt,[c] John R. Middleton,[b] Douglas D. Rhoads[a,#]

[a]Cell and Molecular Biology program, University of Arkansas, Fayetteville, Arkansas, USA

[b]Dept. of Veterinary Medicine and Surgery, University of Missouri, Columbia, Missouri, USA

[c]Dept. of Veterinary Pathobiology, University of Missouri, Columbia, Missouri, USA

Running Head: *S. agnetis* phylogenomics

[#]Address correspondence to Douglas Rhoads, drhoads@uark.edu

Keywords: Staphlococcus, Broiler, Mastitis, Lameness, Phylogeny

Abstract

We report the comparisons of genomes of eleven isolates of *Staphylococcus agnetis* obtained

from dairy cattle and chickens. *S. agnetis* has been previously associated with subclinical or

clinically mild cases of mastitis in dairy cattle and is one of several Staphylococcal species

isolated from the bone and blood of lame broilers. We have previously shown that a chicken

isolate 908 can induce bacterial chondronecrosis with osteomyelitis (BCO) in chickens with

incidences of lameness over 50%. We have also assembled and analyzed the genome of isolate

908. To better understand the relationship between dairy cattle and broiler isolates, we

assembled genomes for *S. agnetis* isolated from an additional chicken BCO case, and from milk,

mammary gland secretions or udder skin from the collection at the University of Missouri. To

trace phylogenetic relationships, we constructed phylogenetic trees based on multi-locus

sequence typing, and Genome-to-Genome Distance Comparisons. Chicken isolate 908 clustered

with two of the cattle isolates along with isolates from chickens in Denmark and an isolate of *S.*

agnetis we isolated from a BCO lesion on a commercial farm in Arkansas. We used a number of

BLAST tools to compare the chicken isolates to those from cattle and identified 98 coding

sequences distinguishing isolate 908 from the cattle isolates. None of the identified genes

explain the differences in host or tissue tropism. These analyses are critical to understanding

how Staphylococci colonize and infect different hosts and potentially how they can transition to

alternative niches (bone vs dermis).

Importance

Staphylococcus agnetis has been recently recognized as associated with disease in dairy cattle

and meat type chickens. The infections appear to be limited in cattle and systemic in broilers.

This report details the molecular relationships between cattle and chicken isolates in order to

understand how this recently recognized species infects different hosts with different disease

manifestations. The data show the chicken and cattle isolates are very closely related but the

chicken isolates all cluster together suggesting a single jump from cattle to chickens.

Introduction

In the US, skeletal problems are estimated to cost the broiler industry more than 100 million

dollars annually (1-5). Lameness is an important chicken industry issue and a wide array of

bacterial genera have been associated with bacterial chondronecrosis with osteomyelitis (BCO)

in poultry (5-25). *Staphylococcus agnetis*, a coagulase-variable, Gram-positive bacterium has

been found to cause infections of the bones and blood of broilers leading to BCO (26, 27). BCO

primarily affects the growth plate in the proximal femur and tibia, the fast-growing leg bones.

We have shown that an isolate of *S. agnetis* (strain 908) obtained from BCO chickens can induce

BCO lameness at levels greater than 50% of the population when administered in a single dose in

drinking water (26, 27). Previously, *S. agnetis* has also been associated with subclinical or mild

cases of clinical mastitis in dairy cattle (28-31). There are very few reports of *S. agnetis* in

poultry and we have speculated that the virulent strain we isolated may be the result of prolonged

selection resulting from years of inducing BCO lameness at our research farm. Genome

sequence analysis of multiple isolates of *S. agnetis* from the University of Arkansas research

farm have revealed little sequence variation and thus they appear to be clonal (unpublished data).

The annotated complete genome of strain 908 has been published (26). Draft genomes of a cattle

isolate, *S. agnetis* CBMRN20813338 (32), and chicken isolates (33) have been deposited in the

NCBI genome databases. To better understand the phylogenomic relationships between dairy

cattle and broiler isolates, we have generated genome assemblies for multiple cattle isolates and

an additional chicken isolate of *S. agnetis*. We used multi-locus sequence typing (MLST) and

genome distance comparisons to develop phylogenetic trees. We also performed reciprocal

BLAST and BLASTX comparisons to identify genes and gene islands that distinguish the

chicken and cattle isolates. The goal was to determine the phylogenetic relationships between

cattle and chicken isolates, and whether there were easily discernable genes responsible for the

virulence of isolate 908, or species-specific pathogenesis.

Results.

Staphylococcus agnetis **genomes assemblies.**

Sources (host, tissue, disease) for the *S. agnetis* isolates used in our analyses are presented in

Table 1. For this work, we generated draft genomes for eight cattle isolates (1383, 1384, 1385,

1387, 1389, 1390, 1391, 1392) from 2x251 paired end MiSeq reads (Table S1), and we generated

one finished genome for one cattle isolate (1379). These cattle isolates were cultured from skin

swabs (1385, 1389), milk (1379, 1383, 1384, 1387, 1391, 1392), or pre-partum mammary gland

secretions (1390). The new, draft, *de novo* assemblies for the eight cattle isolates ranged from 43

to 328 contigs comprising 2.381 to 2.581 Mbp (Table S1). The hybrid assembly from long and

short reads (see Materials and Methods for details) for cattle isolate 1379 produced a single

chromosome of 2.45 Mbp. We identified *S. agnetis* isolate 1416 from a BCO lesion in a

necropsy sampling of BCO birds on a commercial broiler farm in Arkansas. The hybrid

assembly of the 1416 genome produced a 2.45 Mbp chromosome and plasmids of 59 and 28 kbp.

We included the draft genome for cattle isolate CBMRN20813338 as it was the first *S. agnetis*

genome characterized (32). We had earlier published the finished genome for chicken isolate *S.*

agnetis 908, a *de novo* assembly of Pacific Biosciences long reads, with subsequent correction

with MiSeq reads (26). This assembly includes a single 2.47 Mbp chromosome and a 29 kbp

plasmid. Recently, we identified two additional plasmids of 3.0 and 2.2 kbp (unpublished) from

the assembly data that we have included in our genome comparisons.

Phylogenetic analyses. To begin to trace the phylogenomic relationship between the cattle *S.*

agnetis isolates and those from chicken, we first generated MLST phylogenetic trees. We

included a total of 13 isolates, including the published genome for cattle isolate *S. agnetis*

CBMRN20813338, the 9 new cattle isolate assemblies, chicken isolates 908 and 1416, and *S.*

agnetis isolate 12B from NCBI. We included the genome from strain 12B, isolated from the

milk of a buffalo with bubaline mastitis. The dendrogram from genome BLAST on the NCBI

genome page for *S. agnetis*, presents 12B as the closest genome for a bovine isolate to the

genome for chicken isolate 908. *S. hyicus* ATCC11249[T] (34) from swine exudative dermatitis

was used as the outgroup. Figure 1 presents a tree based on seven housekeeping genes (*ackA,*

fdhD, fdhF, grol, purA, tpiA, tuf), where orthologs could be identified in each of the assemblies,

and the genes are dispersed throughout the 908 chromosome. From the MLST phylogenetic tree,

we see two chicken isolates (908 and 1416) cluster within the cattle *S. agnetis* isolates within a

clade with bovine isolates 1379, 1387 and 12B. MLST analysis with seven virulence genes

(encoding five distinct fibronectin binding proteins and two exotoxins) identified in all of the

assemblies produced a tree with a very similar topology (data not shown).

Since we began our analyses, additional genomes for isolates of *S. agnetis* have been deposited

in NCBI. The NCBI dendrogram based on genomic BLASTN

(https://www.ncbi.nlm.nih.gov/genome/?term=agnetis) for 26 *S. agnetis* assemblies, indicates

that our 908 chicken isolate clusters with three Danish chicken isolates and 5 bovine isolates

(12B, SUNC_2265, SNUC_4805, SNUC_5151, SUC_3261) in a single clade relative to

CBMRN20813338, and 16 additional bovine isolates. In order to expand on the MLST analyses

and all 36 genomes (26 in NCBI and our 10 new assemblies, including 9 isolates of bovine origin

and 1 isolate of chicken origin) we used the Genome-to-Genome Distance Calculator (GGDC) to

generate a phylogenetic tree based on genetic distances computed from whole genome BLASTN

comparisons (35). This included our two chicken isolates (908, 1416) and three chicken isolates

(NEDS, NEFX, NDYM) from organs from two deceased broilers on a farm in Denmark (33).

The phylogenetic tree based on genome distances (Fig. 2) shows that four of the genomes from

chicken isolates (908, NEDS, NEFX, NDYM) cluster within the cattle isolates with the Denmark

chicken isolates being most similar to our chicken isolate 908. Chicken isolate 1416 is in a sister

branch clustered with 7 bovine isolates including isolate 12B from the milk of a buffalo in

Argentina with mastitis. The data are consistent with five, or potentially six, different clades

within the *S. agnetis* species group with the five chicken isolate genomes all within one clade.

The nine new genomes for mastitis-related isolates of *S. agnetis* from the USA are distributed

across all branches of the tree. There is no indication of geographic restriction of particular

genotypes for *S. agnetis* isolated from the bovine mammary gland. Nor is there a particularly

noticeable separation of the chicken isolate genomes from the cattle isolate genomes.

Genome Comparison. We used CGView Server (36) to perform and visualize comparisons of

the 2.47 Mbp chromosome from chicken isolate 908 to three cattle isolate genomes; the finished

1379 isolate genome and draft genomes for isolates 1387 and 1385 (Fig. 3). We selected the

1379 isolate genome for production of a finished genome based on being one of the closest

genomes to the chicken isolates (Fig. 2). We included the draft 1387 isolate genome from the

same branch as the chicken isolates, and 1385 as the largest assembly of the other cattle isolates.

The CGView in Figure 3 identified five gene islands which appear to distinguish chicken isolate

908, from the three cattle isolates. The five islands were also visible when we compared chicken

isolate 908 to our other new draft cattle assemblies or the buffalo isolate 12B (data not shown).

We hypothesized that these islands could potentially contain sequences related to host

adaptation. We annotated the genes in these five islands using BLASTP and further evaluated for

presence in the other four chicken isolate genomes, or any of the currently available 36 bovine

isolate genomes (Table S2). Regions in the 908 genome not represented in the cattle isolates

according to the CGView are located approximately as follows: island 1 for 167-235 kbp; island

2 for 978-1021 kbp; island 3 for 1162-1177 kbp ; island 4 for 1831-1848 kbp; and island 5 for

2007-2018 kbp. Thus, approximately 154 kbp out of 2474 kbp are distinct from the cattle

isolates.

Analysis of the 908 2.47 Mbp chromosome for prophage using the PHASTER website (data not

shown) identifies island 1 as containing two intact Staphylococcal prophage

(Staphy_EW_NC_007056 and Staphy_IME_SA4_NC_029025) from 170.1 to 232.7 kbp.

Islands 2 and 3 are identified as questionable for being complete prophage. Island 2 is

Staphy_2638A_NC_007051 from 980.8 to 1020.9 kbp and island 3 is most similar to a

Clostridium phage, phiMMP04_NC_019422, from 1156.8 to 1178.5 kbp. Island 4 contains

genes indicative of a conjugative transposon, with sequence similarity to Tn*6012* of *S. aureus,*

inserted in an intergenic region approximately 170 bp upstream of the *bioD* gene. The shortest of

the five blocks is island 5 (~11 kbp), which contains an apparent operon encoding a strain

variable Type 1 DNA restriction-modification system (*hsdMSR*). Therefore, 124.4 of the

estimated 154 kbp in the five islands represents prophage sequences, while the other two contain

a probable transposon and a restriction-modification operon. The most similar match to island 4

and 5 in BLAST searches at NCBI were to *S. aureus* genomes. Figure 4 further relates these 5

islands as candidates for host adaptation by comparing the 908 isolate 2.47 Mbp chromosome to

the finished 1379 bovine isolate genome and draft assemblies of chicken isolates NEDS from the

a deceased broiler in Denmark and the finished genome assembly of isolate 1416 from a BCO

broiler on a commercial farm in Arkansas. From these comparisons we conclude that islands 1,

2, 4, and 5 are, for the most part, present in at least one of the other two chicken isolates, but

none of the islands is in both of the other chicken isolates (i.e., specific to all chicken isolates).

There is the caveat that the 908 isolate and 1416 isolate genomes are finished genomes and the

NEDS genome is only a draft genome so any island not found in the NEDS assembly could

potentially be an assembly issue. Close inspection of the TBLASTN analysis of the proteins

from the islands for presence in any of the bovine isolate genomes (Table S2) 164 of the 908

predicted polypeptides have significant matches in at least one of the cattle isolates, while only

32 are not found in any of the cattle isolates. For those 32 polypeptides, 31 are also not found in

either of the four other chicken isolates (1416 or the three Danish isolates). Polypeptide 217 is

the only polypeptide not identified in any bovine isolate genome but is identified in the 1416

genome. Polypeptide 217 is a 47 amino acid hypothetical protein, so we see no real islands of

polypeptides (i.e., chicken specific pathogenicity island) that distinguish chicken isolate genomes

from the bovine isolate genomes.

We have assembled sequences of three plasmids in isolate 908 (29, 3 and 2.2 kbp). None of

these plasmids is found in any of our nine newly assembled cattle isolate listed in Table 1. There

are presently 26 genome assemblies for *S. agnetis* in the NCBI database. Only two assemblies

are listed as "completed" (i.e., finished), our assembly for chicken isolate 908 (26), and isolate

12B from buffalo milk in Argentina (unpublished). The other 24 are draft assemblies. The NCBI

genomes include 21 isolates from cattle, one from buffalo, and four chicken isolates; 908 and the

three chicken isolates from Denmark (NEDS, NEFX, and NDYM). We performed a "highly

similar" BLASTN search for the three plasmids from isolate 908 plasmids in the 26 assemblies

already in NCBI. The 29 kbp plasmid identified one 4729 base contig in the NEFX assembly

with 22% query coverage in 10 different regions, with the largest region comprising 3317

identities over 3331 bases. The 3 kbp plasmid matched 450 out of 478 bases in 565 bp contigs in

all three of the Danish chicken isolate assemblies (NDYM, NEDS, NEFX). The 2.2 kbp plasmid

matched 2071 out of 2080 bases in a 2304 base contig in the chicken NEFX assembly. We

therefore conclude that none of the plasmids in isolate 908 appear to correspond to a chicken

host specialization determinant, but some genes in the 29 kbp plasmid could potentially be

relevant to chicken pathogenesis.

To screen at higher resolution, we used the SEED Viewer sequence comparison tool to compare

entire assemblies for individual polypeptide coding sequences for four isolates: chicken isolate

908 (including the 3 plasmids), chicken isolate 1416, bovine isolate 1379 and the bovine

CBMRN isolate. We used isolates 908 and 1416 as finished assemblies of two chicken isolates,

1379 as a finished assembly of a cattle isolate, and CBMRN as the original draft cattle isolate.

The BLASTP comparison results were then filtered for isolate 908 polypeptides with >90%

identity for polypeptides in isolate 1416, but <50% identity for isolate 1379 and the CBMRN

isolate. This filter identified 99 polypeptides (Table S3). To predict functions of these 99

polypeptides we used both the RAST annotation and the NCBI Prokaryote Genome Annotation

Pipeline (PGAP) to categorize the potential function of these 99 polypeptides. We identified 75

polypeptides as hypothetical or of unknown function, 11 phage related, and 8 involved in mobile

elements or plasmid maintenance. The remaining five polypeptides, listed under "other", are:

deoxyuridine 5'-triphosphate nucleotidohydrolase (EC 3.6.1.23), hypothetical SAR0365 homolog

in superantigen-encoding pathogenicity islands SaPI, ribosyl nicotinamide transporter PnuC-like,

aspartate aminotransferase (EC 2.6.1.1), and N-acetyl-L,L-diaminopimelate aminotransferase

(EC 2.6.1.-). If we relaxed the cutoff for the cattle isolates to <70% identity in cattle isolates we

identified 9 additional polypeptides which added one additional hypothetical polypeptide, four

additional phage related polypeptides, and four additional polypeptides involved in mobile

element or plasmid maintenance. There were no additional polypeptides in the "other" category,

only the five described above (Table S3).

The dUTP nucleotidohydrolase (Gene ID 209; 191,107-191,625 bp) is annotated as a phage

related protein with roles in viral replication for reducing incorporation of uracil in viral DNA

and is located within the Staphy_EW_NC_007056 prophage in island 1 described above, so this

gene is likely to function primarily in the biology of that prophage.

The SAR0365 homolog (Gene ID 1037; 1,018,987-1,020,928 bp) is encoded in island 2 within

the Staphy_2638A_NC_007051 prophage. SAR0365 polypeptide is a hypothetical protein that

the NCBI Prokaryotic Genome Annotation Pipeline annotates as a toxin in the PemK/MazF type

II toxin-antitoxin system. Four of the seven protein entries for SAR0365 homologs in NCBI are

associated with superantigen-encoding pathogenicity islands (SaPI) in clinical isolates of

Staphylococcus aureus and two are associated with *S. aureus* phages. Mobilization of SaPI has

been associated with temperate phage replication (37). The 908 isolate genome contains

additional hypothetical genes annotated as SaPI-associated homologs (Gene ID 1184 1,173,209-

1,173,412 bp; Gene ID 1185 1,173,409-1,173,714 bp; Gene ID 1938 1,957,610-1,959,703 bp;

Gene ID 1939 1,959,935-1,961,452 bp; Gene ID 2116 2,136,487-2,136,603 bp). Mobilization

depends on a terminase (37), but the only SaPI associated terminase is Gene ID 2092 (2,118,053-

2,118,355 bp). We had previously described a cluster of five exotoxin/superantigen-like proteins

from 1,956,884 to 1,968,958 bp (26). Therefore, the only potential superantigen-containing

pathogenicity island would approximate from 1.95 to 2.12 Mbp which would be larger than the

prototypical 15-18 kbp SaPI (37). A BLASTP of the *S. agnetis* protein database at NCBI with

the predicted protein for Gene ID 1037 (SAR0365 homolog) identified the isolate 908 entry, as

well as identical entries in all three Danish chicken isolates (NEDS, NDYM, NEFX), but no

entries in any of the 20 cattle *S. agnetis* isolates in NCBI. Expanding the BLASTP to all

Staphylococcaceae identified highly similar matches (92% identity, 100% query coverage) in

Staphylococcus hominis and less similar (50% identity, 98% query coverage) in *S. aureus*.

Further analyses and additional samples would be required to speculate further regarding the role

of this SAR0365 homolog as a virulence factor in chicken tropism.

The genes for ribosyl nicotinamide transporter (Gene ID 2466), aspartate aminotransferase (Gene

ID 2469) and N-acetyl-L,L-diaminopimelate aminotransferase (Gene ID 2470) are located in a

five gene region on the 29 kbp plasmid, with the other 2 genes encoding hypothetical

polypeptides. We performed BLASTP of the Staphylococcaceae proteins at NCBI. The 89

amino acid ribosyl nicotinamide transporter matched multiple entries from *S. aureus* and all three

Danish *S. agnetis* isolates from chicken. Many of the BLASTP hits for this polypeptide are

annotated as an AbrB family transcriptional regulator by the NCBI Prokaryotic Genome

Annotation Pipeline (PGAP). The 64 amino acid hypothetical polypeptide for Gene ID 2467 is

only conserved in two entries from the Danish *S. agnetis* broiler isolates. The 197 residue

polypeptide from Gene ID 2468 is well conserved in a broad swath of staphylococci and PGAP

annotates this polypeptide as an IS*6* family transposase. We note that Gene ID 2469 and 2470

are close to each other and in different reading frames suggestive of a possible frameshift

introduced as an assembly error. Indeed, if we join the predicted polypeptides of these two

ORFs, BLASTP analysis identifies *Staphylooccus* protein entries that match over the span of the

merged polypeptides. However, we have evaluated this hypothesis further by templated

assembly of the 908 MiSeq data onto the 29 kbp plasmid sequence. The MiSeq reads all agree

with the assembly as presented in our NCBI submission. Therefore, these two ORFs in the 29

kbp plasmid may be a frameshifted pseudogene or, if translated may have alternate functions for

this organism.

Finally, to determine whether there were any regions in the cattle isolates that are not found in

the chicken isolate genome assemblies, we performed a CGView analysis with the finished

genome from isolate 1379 as the reference (Fig. 5). We included cattle isolate CBMRN and

compared to the finished genomes of chicken isolates 908 (chromosome plus three plasmids),

and 1416. There were cattle isolate regions that appeared to be absent from one of the chicken

isolates but there were no regions found in both cattle isolates that were missing from both

chicken isolates.

Discussion.

The staphylococcal genus not only includes a number of pathogenic species infecting vertebrate

animals worldwide, but also includes many saprophytic or commensal species (38-40). *S. agnetis*

is closely related to *S. hyicus* and *S. chromogenes* and was only described as a distinct

staphylococcal species in 2012, based on DNA sequence differences of rDNA and two protein

coding genes in isolates from mastitis in dairy cattle (41). *S. agnetis* cannot be easily

differentiated from *S. hyicus* using routine speciation techniques, e.g. partial 16S rDNA

sequencing, MALDI-TOF, or fermentation methods (28-30, 42). Hence, *S. agnetis* has either

escaped recognition due to misclassification or is an emerging pathogen in some agricultural

animal species. While *S. agnetis* was originally reported in cattle mastitis (41), it has more

recently been reported in chicken bone infections and in multiple internal organs from deceased

broilers (33). Metagenomics also detected *S. agnetis* 16S rDNA sequences in the gut of a sheep

scab mite (43). Phylogenetic analyses based on 16S rRNA sequences cluster *S. agnetis* very

close to *S. hyicus* (26) with a group of staphylococci associated with domestic vertebrate species

(e.g., cattle, swine, dog) (28-31, 34, 44-52). Most of these species are associated with dermal or

epithelial infections, such as exudative dermatitis (28, 29, 31, 34, 46, 48-50, 52-54) and not with

osteomyelitis as we have seen with chicken isolate 908 (26, 27). The more phylogenetically

distant taxon, *S. aureus*, is prominently known for osteomyelitis in humans (55-57). That the

three Danish *S. agnetis* isolates, and our isolates 908 and 1416, are all closely related and within

a clade of the cattle *S. agnetis* isolates suggests a recent expansion of the host range (i.e. from

cattle to chickens) as seen for a human-specific clade of *S. aureus* that "jumped" to chickens in

the United Kingdom (17). The Danish broiler chicken isolates were from multiple tissues from

deceased birds and we have no information about possible osteomyelitis. A single radiation out

of the cattle group also argues against *S. agnetis* jumping back and forth between cattle and

chickens. We have previously reported that isolate 908 can produce a bacteremia in the latter

stages of BCO development before the birds are overtly lame (5, 26, 27, 58). We do not know if

the Danish isolates can induce the BCO lameness that we have demonstrated for isolate 908 (26,

27). Our isolate 908 appears to represent a hypervirulent clone expanded through years of

inducing high levels of BCO lameness on our research farm (5, 26, 27, 59-62) and could have

evolved through selection from less virulent *S. agnetis* in broiler populations. Therefore, our

genomic comparisons have been directed towards identification of any gene(s) that *S. agnetis*

908 could have acquired that facilitate the switch from involvement in cattle epithelial and

mammary gland colonization and infection, to bone infections in chickens.

None of the gene islands (Table S2) or individual genes (Table S3) we identified as

distinguishing isolate 908 from closely related cattle isolates is currently recognizable as a

virulence marker, or that mediates tissue tropism. Previously we had identified 44 virulence

genes in our annotation of the isolate 908 genome (26) and none of these genes is in the regions

distinguishing the chicken and cattle isolates. The genomic analyses of the human-to-chicken

jump for *S. aureus* was associated with acquisition of new mobile elements, inactivation of

several virulence determinants important to human pathogenesis, and enhanced resistance to

chicken neutrophils (17). Most of the distinguishing gene islands in isolate 908 contain genes

associated with mobile elements (prophage), but none are virulence determinants. We have

unpublished evidence that isolate 908 is highly resistant to an immortalized chicken macrophage

and are pursuing the genes for macrophage resistance. Since we have failed to identify unique

virulence genes that distinguish the chicken isolate 908 from the cattle isolates we conclude that

the basis for the jump from cattle to chickens is most likely the result of small alterations (i.e.,

missense or regulatory mutations) in a few virulence-associated factors. Hypervirulence of

isolate 908 in chickens could be from a single amino acid change. Hypervirulence of isolates of

Campylobacter jejuni, were demonstrated to result from a single substitution in an outer

membrane protein, resulting in induction of spontaneous abortions in sheep (63). Therefore,

further fine-level comparisons or directed genome evolution (63) will be required to dissect how

this emerging pathogen has evolved and diversified from cattle mastitis to chicken bone

pathogen.

Materials and Methods

Reference genomes. Isolate designations and host sources are provided in Table 1 and the

details of the genome assemblies and accessions in NCBI are in Table S1. Chicken isolate 908

was from necrotic femoral lesions, while NDYM, NEDS and NEFX were from tissue samples

from deceased broilers in Denmark. Isolates NDYM and NEDS were from the same broiler.

The cattle isolate CBMRN was isolated from milk of a cow with subclinical mastitis that was

enrolled in the Canadian Bovine Mastitis Research Network (CBMRN) cohort study (32).

ATCC11249 represents the *S. hyicus* type strain isolated from a pig exudative epidermititis

lesion (34).

Bacterial strains. Genomes for ten *S. agnetis* isolates were newly assembled (Table S1),

including nine cattle isolates, and one chicken isolate. Chicken isolate 1416 was isolated from a

necrotic femoral lesion of a lame bird in a commercial broiler operation in Arkansas. The nine

cattle isolates were from a collection at the University of Missouri. Two isolates were skin

isolates, one isolate was obtained from a pre-partum mammary secretion, and six isolates were

obtained from the milk of cows with subclinical mastitis. All cattle isolates had been previously

identified as *S. agnetis* based on partial DNA sequence of either elongation factor Tu (*tuf*) or 3-

dehydroquinate dehydratase (*aroD*) (30). All isolates were archived at -80 ^{o}C in 20-40%

glycerol, maintained on Tryptic Soy Agar slants, and grown in Tryptic Soy Broth (TSB; Difco,

Becton, Dickinson and Company, Franklin Lakes, NJ).

Genome sequencing and assembly. DNA isolation was based on the method described by Dyer

and Iandolo (64). Isolates were grown to mid log phase in TSB (40 ml) at 37 ^{o}C with shaking,

pelleted, and resuspended in 2.5 ml 30 mM TrisCl, 3 mM EDTA, 50 mM NaCl, 50 mM glucose,

pH 7.5. Lysostaphin (Sigma-Aldrich, St. Louis, MO) was added to 20 µg/ml, and incubated at 37

^{o}C for 40-60 min. SDS was added to 0.5%, then the lysate was treated with RNAseA (Sigma-

Aldrich) at 20 ug/ul for 30 min at 37 ^{o}C, then Pronase E (Sigma-Aldrich) at 20 ug/ul for 30 min

at 37 ^{o}C. The lysate was then extracted successively with 50:48:2 phenol:CHCl$_3$:isoamyl

alcohol, and 24:1 CHCl$_3$:isoamyl alcohol. DNA was then collected by ethanol precipitation.

DNA was quantified by Hoechst 33258 fluorometry in a GloMax®-Multi Jr. (Promega Corp.,

Madison, WI), and DNA integrity verified by agarose gel electrophoresis. Purified DNA from

each isolate was submitted to the Research Technology Support Facility Genomics Core at

Michigan State University for barcoded-library construction, pooled and subjected to 2x251

sequencing on an Illumina MiSeq. For draft genome assemblies the MiSeq reads were assembled

using the *de novo* pipeline in Lasergene NGen ver. 13.0 (DNAStar, Madison, WI). For isolates

1379 and 1416 we produced finished genomes by hybrid assemblies of MiSeq and Oxford

Nanopore MinION long reads. Long reads were either from barcoded or rapid kit libraries

prepared and sequenced on Minion v9.3 flow cells (Oxford Nanopore Technologies, Oxford

Science Park, UK) according to the manufacturer's recommendations. Minion reads were

filtered with a custom script to filter reads for length and average Q-score, prior to assembly.

For isolate 1379, we filtered for length >=2000 bases and Qscore >=13. For isolate 1416, we

filtered for length >=5000 bases and Qscore >16. Nanopore reads and MiSeq reads were

assembled using the Unicycler ver. 0.4.6.0 pipeline on Galaxy (https://usegalaxy.org) using the

Bold bridging mode. All assemblies and sequence reads have been deposited in NCBI and the

accession and biosample identifiers are in Table S1.

Genome annotation and phylogenetic comparison. The assembled genome sequences were

compared with chicken isolate 908 using BLASTN implemented in CGViewer (36) to identify

regions missing in one or more genome. Specific gene regions were annotated using either the

BASys server at http://www.basys.ca (65) or the Rapid Annotation using System Technology

(RAST) server at http://rast.nmpdr.org (66). Unique genes were verified by TBLASTN

comparisons and reciprocal gene-by-gene BLASTP comparisons using the SEED server (67).

Unique genes were further annotated using the KEGG (Kyoto Encyclopedia of Genes and

Genomes) website at https://www.genome.jp/kegg (68). Prophage identification was performed

using PHASTER (PHAge Search Tool Enhanced Release) at http://phaster.ca (69).

Phylogenetic Analyses. For MLST analysis gene coding sequences were aligned and trimmed in

MegAlign (DNAStar) then concatenated. Clustal Omega implemented in MegAlignPro

(DNAStar) was used to generate phylogenetic trees. Consensus neighbor-joining trees with 2500

bootstrap replications were constructed based on the alignments. Genome-to-Genome Distance

Calculator (GGDC) http://ggdc.dsmz.de/ggdc.php (35) was used to generate whole genome

BLAST distance values. These distance calculations were used to generate a phylogenetic tree

using the neighbor-joining method as implemented at http://trex.uqam.ca. Trees were rendered

and rerooted in Archeopteryx 0.9901 (70).

Acknowledgements

369   This research was partially supported by grants from the Arkansas Biosciences Institute, Chr

370   Hansen, Zinpro LLC, and Cobb Vantress, Inc. The funders had no role in the design of this

371   study, the interpretation of the results, or the contents of this manuscript.

Literature Cited

1.   Whitehead C. 1992. Bone biology and skeletal disorders in poultry. Carfax Publishing

Co.

2.   Sullivan TW. 1994. Skeletal problems in poultry: Estimated annual cost and descriptions.

Poult Sci 73:879-882.

3.   Kestin SC, Su G, Sorensen P. 1999. Different commercial broiler crosses have different

susceptibilities to leg weakness. Poult Sci 78:1085-1090.

4.   Kestin SC, Gordon S, Su G, Sorensen P. 2001. Relationship in broiler chickens between

lameness, liveweight, growth rate and age. Vet Rec 148:195-197.

5.   Wideman RF. 2016. Bacterial chondronecrosis with osteomyelitis and lameness in

broilers: a review. Poult Sci 95 325-344.

6.   Butterworth A. 1999. Infectious components of broiler lameness: a review. . World

Poultry Sci J 55:327-352.

7.   Duff S. 1990. Do different forms of spondylolisthesis occur in broiler fowls? Avian

Pathol 19:279-294.

8.   Duff S, Randall C. 1987. Observations on femoral head abnormalities in broilers. Res Vet

Sci 42:17-23.

389 9. Duff SRI. 1990. Diseases of the musculoskeletal system. *In* Jordan F (ed), Poultry

390   Diseases 3rd ed. Bailliere, Tindall. UK.

391 10. Emslie KR, Nade S. 1983. Acute hematogenous Staphylococcal osteomyelitis: a

392   description of the natural history in an avian model. Am J Pathol 110:333-345.

393 11. Griffiths G, Hopkinson W, Lloyd J. 1984. Staphylococcal necrosis of the head of the

394   femur in broiler chickens. Aust Vet J 61:293-293.

395 12. Julian RJ. 1985. Osteochondrosis, dyschondroplasia, and osteomyelitis causing femoral

396   head necrosis in turkeys. Avian Dis 29:854-866.

397 13. Julian RJ. 2005. Production and growth related disorders and other metabolic diseases of

398   poultry - A review. Vet J 169:350-369.

399 14. Kense MJ, Landman WJM. 2011. *Enterococcus cecorum* infections in broiler breeders

400   and their offspring: Molecular epidemiology. Avian Pathol 40:603-612.

401 15. Kibenge FS, Wilcox G, Pass D. 1983. Pathogenicity of four strains of Staphylococci

402   isolated from chickens with clinical tenosynovitis. Avian Pathol 12:213-220.

403 16. Li Y, Chen L, Wu X, Huo S. 2015. Molecular characterization of multidrug-resistant

404   avian pathogenic *Escherichia coli* isolated from septicemic broilers. Poult Sci 94:601-

405   611.

406 17. Lowder BV, Guinane CM, Ben Zakour NL, Weinert LA, Conway-Morris A, Cartwright

407   RA, Simpson AJ, Rambaut A, Nübel U, Fitzgerald JR. 2009. Recent human-to-poultry

408   host jump, adaptation, and pandemic spread of *Staphylococcus aureus*. PNAS USA

409   106:19545-19550.

18.   Marek A, Stepień-Pyśniak D, Pyzik E, Adaszek Ł, Wilczyński J, Winiarczyk S. 2016.

Occurrence and characterization of Staphylococcus bacteria isolated from poultry in

Western Poland. Berl tierarztl Wschr 129:147-152.

19.   McNamee PT, Smyth JA. 2000. Bacterial chondronecrosis with osteomyelitis ('femoral

head necrosis') of broiler chickens: a review. Avian Pathol 29:477-495.

20.   Nicoll TR, Jensen MM. 1987. Preliminary studies on bacterial interference of

Staphylococcosis of chickens. Avian Dis 31:140-144.

21.   Packialakshmi B, Rath N, Huff W, Huff G. 2015. Poultry femoral head separation and

necrosis: A review. Avian Dis 59:349-354.

22.   Skeeles KJ. 1997. Staphylococcosis, p 247-253. *In* B.W. Calnek, Barnes HJ, Beard CW,

McDougald LR, Saif YM (ed), Diseases of Poultry 10 ed. Iowa State University Press,

Ames, IA, USA.

23.   Smeltzer M, Gillaspy A. 2000. Molecular pathogenesis of Staphylcoccal osteomyelitis.

Poult Sci 79:1042-1049.

24.   Smith HW. 1954. Experimental Staphylococcal infection in chickens. J Pathol Bacteriol

67:81-87.

25.   Ytrehus B, Carlson C, Ekman S. 2007. Etiology and pathogenesis of osteochondrosis.

Vet Pathol 44:429-448.

26.   Al-Rubaye AAK, Couger MB, Ojha S, Pummill JF, Koon JA, II, Wideman RF, Jr.,

Rhoads DD. 2015. Genome analysis of *Staphylococcus agnetis*, an agent of lameness in

broiler chickens. PLoS ONE 10:e0143336.

27.   Al-Rubaye AAK, Ekesi NS, Zaki S, Emami NK, Wideman RF, Rhoads DD. 2017.

Chondronecrosis with osteomyelitis in broilers: Further defining a bacterial challenge

model using the wire flooring model. Poult Sci 96:332-340.

28.   Adkins PRF, Dufour S, Spain JN, Calcutt MJ, Reilly TJ, Stewart GC, Middleton JR.

2018. Molecular characterization of non-aureus Staphylococcus spp. from heifer

intramammary infections and body sites. J Dairy Sci 101:5388-5403.

29.   Adkins PRF, Dufour S, Spain JN, Calcutt MJ, Reilly TJ, Stewart GC, Middleton JR.

2018. Cross-sectional study to identify staphylococcal species isolated from teat and

inguinal skin of different-aged dairy heifers. J Dairy Sci 101:3213-3225.

30.   Adkins PRF, Middleton JR, Calcutt MJ, Stewart GC, Fox LK. 2017. Species

identification and strain typing of *Staphylococcus agnetis* and *Staphylococcus hyicus*

isolates from bovine milk by use of a novel multiplex PCR assay and pulsed-field gel

electrophoresis. J Clin Microbiol 55:1778-1788.

31.   Åvall-Jääskeläinen S, Taponen S, Kant R, Paulin L, Blom J, Palva A, Koort J. 2018.

Comparative genome analysis of 24 bovine-associated Staphylococcus isolates with

special focus on the putative virulence genes. PeerJ 6:e4560.

32.   Calcutt MJ, Foecking MF, Fry PR, Hsieh H-Y, Perry J, Stewart GC, Scholl DT, Messier

S, Middleton JR. 2014. Draft genome sequence of bovine mastitis isolate *Staphylococcus*

agnetis CBMRN 20813338. Genome Announc 2.

33.   Poulsen LL, Thøfner I, Bisgaard M, Olsen RH, Christensen JP, Christensen H. 2017.

Staphylococcus agnetis, a potential pathogen in broiler breeders. Vet Microbiol 212:1-6.

34.   Calcutt MJ, Foecking MF, Hsieh H-Y, Adkins PRF, Stewart GC, Middleton JR. 2015.

Sequence analysis of *Staphylococcus hyicus* ATCC 11249[T], an etiological agent of

exudative epidermitis in swine, reveals a type VII secretion system locus and a novel

116-kilobase genomic island harboring toxin-encoding genes. Genome Announc

3:e01525-14.

35.   Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based

species delimitation with confidence intervals and improved distance functions. BMC

Bioinformatics 14:60.

36.   Grant JR, Stothard P. 2008. The CGView Server: a comparative genomics tool for

circular genomes. Nucl Acid Res 36:W181-W184.

37.   Ruzin A, Lindsay J, Novick RP. 2001. Molecular genetics of SaPI1 – a mobile

pathogenicity island in *Staphylococcus aureus*. Molecular Microbiology 41:365-377.

38.   Becker K, Heilmann C, Peters G. 2014. Coagulase-negative Staphylococci. Clin

Microbiol Rev 27:870.

39.   Cox HU, Newman SS, Roy AF, Hoskins JD. 1984. Species of Staphylococcus isolated

from animal infections. The Cornell veterinarian 74:124-135.

40.   Mathema B, Mediavilla J, Chen L, Kreiswirth B. 2009. Evolution and taxonomy of

staphylococci, p 31-64. *In* Crossley K, Jefferson K, Archer G, Folwer V (ed),

Staphylococci in Human Disease, 2nd Edition. John Wiley & Sons, Oxford, UK.

41.   Taponen S, Supré K, Piessens V, Van Coillie E, De Vliegher S, Koort JMK. 2012.

Staphylococcus agnetis sp. nov., a coagulase-variable species from bovine subclinical

and mild clinical mastitis. Int J Syst Evol Microbiol 62:61-65.

42.   Zadoks RN, Middleton JR, McDougall S, Katholm J, Schukken YH. 2011. Molecular

epidemiology of mastitis pathogens of dairy cattle and comparative relevance to humans.

Journal of Mammary Gland Biology and Neoplasia 16:357-372.

43.   Hogg JC, Lehane MJ. 1999. Identification of bacterial species associated with the sheep

478   scab mite (*Psoroptes ovis*) by using amplified genes coding for 16S rRNA. Appl Environ

479   Microbiol 65:4227-4229.

480   44.   Bicart-See A, Rottman M, Cartwright M, Seiler B, Gamini N, Rodas M, Penary M,

481   Giordano G, Oswald E, Super M, Ingber DE. 2016. Rapid Isolation of *Staphylococcus*

482   *aureus* Pathogens from Infected Clinical Samples Using Magnetic Beads Coated with Fc-

483   Mannose Binding Lectin. PLoS ONE 11:e0156287.

484   45.   Christensen GD, Baddour LM, Simpson WA. 1987. Phenotypic variation of

485   *Staphylococcus epidermidis* slime production in vitro and in vivo. Infect Immun 55:2870-

486   2877.

487   46.   Stepień-Pyśniak D, Marek A, Rzedzicki J. 2009. Occurrence of bacteria of the genus

488   Staphylococcus in table eggs descended from different sources. Pol J Vet Sci 12:481-484.

489   47.   Devriese LA. 1977. Isolation and identification of *Staphylococcus hyicus*. Am J Vet Res

490   38:787-792.

491   48.   Devriese LA, Hajek V, Oeding P, Meyer SA, Schleifer KH. 1978. *Staphylococcus hyicus*

492   (Sompolinsky 1953) comb. nov. and *Staphylococcus hyicus* subsp. chromogenes subsp.

493   nov. Int J Syst Evol Microbiol 28:482-490.

494   49.   Devriese LA, Laevens H, Haesebrouck F, Hommez J. 1994. A simple identification

495   scheme for coagulase negative staphylococci from bovine mastitis. Res Vet Sci 57:240-

496   244.

50.   Foster AP. 2012. Staphylococcal skin disease in livestock. Vet Dermatol 23:342-e63.

51. Tse H, Tsoi HW, Leung SP, Urquhart IJ, Lau SKP, Woo PCY, Yuen KY. 2011.

Complete genome sequence of the veterinary pathogen *Staphylococcus pseudintermedius*

strain HKU10-03, isolated in a case of canine pyoderma. J Bact 193:1783-1784.

52. Zakour NLB, Bannoehr J, van den Broek AHM, Thoday KL, Fitzgerald JR. 2011.

Complete genome sequence of the canine pathogen *Staphylococcus pseudintermedius*. J

Bact 193:2363-2364.

53. Le Maréchal C, Seyffert N, Jardin J, Hernandez D, Jan G, Rault L, Azevedo V, François

P, Schrenzel J, van de Guchte M, Even S, Berkova N, Thiéry R, Fitzgerald JR, Vautor E,

Le Loir Y. 2011. Molecular basis of virulence in *Staphylococcus aureus* mastitis. PLoS

ONE 6:e27354.

54. Witte W, Hummel R, Meyer W, Exner H, Wundrak R. 1977. Ecology of *Staphylococcus*

aureus: Characterization of strains from chicken. Z Allg Mikrobiol 17:639-646.

55. Aanensen DM, Feil EJ, Holden MTG, Dordel J, Yeats CA, Fedosejev A, Goater R,

Castillo-Ramírez S, Corander J, Colijn C, Chlebowicz MA, Schouls L, Heck M, Pluister

G, Ruimy R, Kahlmeter G, Åhman J, Matuschek E, Friedrich AW, Parkhill J, Bentley

SD, Spratt BG, Grundmann H. 2016. Whole-genome sequencing for routine pathogen

surveillance in public health: a population snapshot of invasive *Staphylococcus aureus* in

Europe. mBio 7:e00444-16.

56. Loughran AJ, Gaddy D, Beenken KE, Meeker DG, Morello R, Zhao H, Byrum SD,

Tackett AJ, Cassat JE, Smeltzer MS. 2016. Impact of **sarA** and phenol-soluble modulins

on the pathogenesis of osteomyelitis in diverse clinical Isolates of *Staphylococcus aureus*.

Infect Immun 84:2586-2594.

57.   Powers ME, Wardenburg JB. 2014. Igniting the fire: *Staphylococcus aureus* virulence

factors in the pathogenesis of sepsis. PLoS Pathog 10:e1003871.

58.   Mandal RK, Jiang T, Al-Rubaye AA, Rhoads DD, Wideman RF, Zhao J, Pevzner I,

Kwon YM. 2016. An investigation into blood microbiota and its potential association

with bacterial chondronecrosis with osteomyelitis (BCO) in broilers. Sci Rep 6:25882.

59.   Wideman RF, Al-Rubaye A, Gilley A, Reynolds D, Lester H, Yoho D, Hughes JM,

Pevzner I. 2013. Susceptibility of 4 commercial broiler crosses to lameness attributable to

bacterial chondronecrosis with osteomyelitis. Poult Sci 92:2311-2325.

60.   Wideman RF, Al-Rubaye A, Kwon YM, Blankenship J, Lester H, Mitchell KN, Pevzner

IY, Lohrmann T, Schleifer J. 2015. Prophylactic administration of a combined prebiotic

and probiotic, or therapeutic administration of enrofloxacin, to reduce the incidence of

bacterial chondronecrosis with osteomyelitis in broilers. Poult Sci 94:25-36.

61.   Wideman RF, Al-Rubaye A, Reynolds D, Yoho D, Lester H, Spencer C, Hughes JD,

Pevzner IY. 2014. Bacterial chondronecrosis with osteomyelitis in broilers: Influence of

sires and straight-run versus sex-separate rearing. Poult Sci 93:1675-1687.

62.   Wideman RF, Hamal KR, Stark JM, Blankenship J, Lester H, Mitchell KN, Lorenzoni G,

Pevzner I. 2012. A wire-flooring model for inducing lameness in broilers: Evaluation of

probiotics as a prophylactic treatment. Poult Sci 91:870-883.

63.   Wu Z, Periaswamy B, Sahin O, Yaeger M, Plummer P, Zhai W, Shen Z, Dai L, Chen SL,

Zhang Q. 2016. Point mutations in the major outer membrane protein drive

hypervirulence of a rapidly expanding clone of *Campylobacter jejuni*. PNAS USA

113:10690-10695.

64.   Dyer DW, Iandolo JJ. 1983. Rapid isolation of DNA from *Staphylococcus aureus*. Appl

543   Environ Microbiol 46:283-285.

544   65.   Van Domselaar GH, Stothard P, Shrivastava S, Cruz JA, Guo A, Dong X, Lu P, Szafron

545   D, Greiner R, Wishart DS. 2005. BASys: a web server for automated bacterial genome

546   annotation. Nucl Acid Res 33:W455-W459.

547   66.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S,

548   Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil

549   LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O,

550   Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid annotations using

551   subsystems technology. BMC Genomics 9:75.

552   67.   Goesmann A, McHardy AC, Osterman A, Hanson A, Linke B, Rückert C, Iwata-Reuyl

553   D, Rodionov DA, Frank ED, Glass EM, Meyer F, Olsen G, Pusch GD, Chuang H-Y,

554   Neuweger H, Thiele I, Steiner J, Choudhuri JV, Krause L, Cohoon M, Fonstein M, Kubal

555   M, Diaz N, Jamshidi N, Larsen N, Vassieva O, Zagnitko O, Butler RM, Stevens R,

556   Edwards R, Olson R, Overbeek R, Jensen R, Gerdes S, Begley T, Disz T, de Crécy-

557   Lagard V, Portnoy V, Vonstein V, Ye Y. 2005. The subsystems approach to genome

558   annotation and its use in the project to annotate 1000 genomes. Nucl Acid Res 33:5691-

559   5702.

560   68.   Morishima K, Tanabe M, Furumichi M, Kanehisa M, Sato Y. 2018. New approach for

561   understanding genome variations in KEGG. Nucl Acid Res 47:D590-D595.

562   69.   Pon A, Marcu A, Arndt D, Grant JR, Sajed T, Liang Y, Wishart DS. 2016. PHASTER: a

563   better, faster version of the PHAST phage search tool. Nucl Acid Res 44:W16-W21.

70. Han MV, Zmasek CM. 2009. phyloXML: XML for evolutionary biology and

comparative genomics. BMC Bioinformatics 10:356.

71. Naushad S, Barkema H, Luby C, Condas L, Nobrega D, Carson D, De Buck J. 2016.

Comprehensive phylogenetic analysis of bovine mon-aureus Staphylococci species based

on whole-genome sequencing. Front Microbiol 7.

Table 1. Bacterial genomes utilized in these analyses. Genomes are separated by species

designation and host source. For each genome the isolate designation is given as well as our

abbreviation for this manuscript where indicated. The Isolate Source indicates tissue or sample

source for the bacterial isolate. Genome status indicates whether the genome is considered

finished or draft. Citation is the publication source. Further information on the genome

assemblies is provided in Table S1.

Strain Designation (Abbreviation)	Isolate Source	Genome Status	Citation
S. hyicus isolate from swine			
ATCC 11249 (ATCC)	swine exudative epidermitis	Finished	(34)
S. agnetis isolates from chickens			
1416	broiler femoral BCO lesion; Arkansas broiler commercial farm	Finished	This work
722_230714_2_5_spleen (NEDS)	broiler spleen, deceased; Denmark	Draft	(33)
722_260714_1_8_heart (NDYM)	broiler heart, deceased; Denmark	Draft	
723_310714_2_2_spleen (NEFX)	broiler spleen, deceased; Denmark	Draft	
908	broiler femoral BCO lesion; UA Research farm	Finished	(26)
S. agnetis isolates from bovine			
12B	buffalo milk		none
1379	bovine mammary gland - milk	Finished	This work
1383	bovine mammary gland - milk	Draft	
1384	bovine mammary gland - milk	Draft	
1385	bovine teat skin	Draft	
1387	bovine mammary gland – milk	Draft	
1389	bovine inguinal skin	Draft	
1390	bovine mammary gland – pre-partum mammary gland secretion	Draft	
1391	bovine mammary gland – milk	Draft	
1392	bovine mammary gland - milk	Draft	
33	bovine	Draft	none
3682	bovine milk	Draft	
43	bovine	Draft	
59 (59a)	bovine	Draft	
59 (59b)	bovine	Draft	

Strain Designation (Abbreviation)	Isolate Source	Genome Status	Citation
6	bovine	Draft	none
CBMRN 20813338 (CBMRN)	bovine mammary gland - milk	Draft	(32)
DSM_23656	bovine mastitic milk	Draft	none
SNUC_1371	Holstein clinical mastitis	Draft	(71)
SNUC_1383	Holstein clinical mastitis	Draft	
SNUC_2265	Holstein subclinical mastitis	Draft	
SNUC_2493	Holstein subclinical mastitis	Draft	
SNUC_3261	Holstein subclinical mastitis	Draft	
SNUC_3610	Holstein subclinical mastitis	Draft	
SNUC_3836	Holstein subclinical mastitis	Draft	
SNUC_4051	Holstein subclinical mastitis	Draft	
SNUC_4805	Holstein subclinical mastitis	Draft	
SNUC_5151	Holstein subclinical mastitis	Draft	
SNUC_5631	Holstein subclinical mastitis	Draft	
SNUC_719	Holstein subclinical mastitis	Draft	
SNUC_725	Holstein subclinical mastitis	Draft	

Figure 1. MLST phylogeny two chicken and eleven bovine, isolates of *S. agnetis*, with a swine *S. hyicus* (ATCC) isolate as the out-group. The tree is based on seven housekeeping genes (see text).

Figure 2. Genome to Genome Distance phylogenetic

tree for 35 *S.agnetis* genomes. GGDC data (formula 2)

was used to construct a neighbor-joining tree for

comparison of five chicken (G_) and 31 bovine (B_)

isolates, of *S. agnetis.* Isolates are described in the text

and Table S1. *S. hyicus* (ATCC) from swine (S_) was

the out group.

Figure 3. CGView blastn comparison of *S. agnetis* 908, 1379, 1387 and 1385 genomes.

The *S. agnetis* 908 2.4 Mbp chromosome was the reference for blastn comparisons with three *S. agnetis* cattle isolates, 1379 (pink), 1387 (green), and 1385 (blue). Parameters were Query size = 10000, overlap 5000, expect=0.0001. Intensity of the color is indicative of the blastn score. The outer two rings show the annotated genes for isolate *S. agnetis* 908. The innermost ring indicates Mbp, the second most innner ring is the GC skew (magenta GC skew-; green GC skew+), and the third most inner ring plots GC content. The Numbered boxes indicate the locations of the 5 gene islands discussed in the text and are not scaled to the size of the island.

Figure 4. CGView blast comparison of *S. agnetis* 908, 1379, NEDS, and 1416 genomes.
The *S. agnetis* 908 2.4 Mbp chromosome was the reference for blastn comparisons with *S. agnetis* cattle isolate 1379 (pink), and chicken isolates; NEDS (green), and 1416 (blue). All other details are as in Figure 3 legend.

Figure 5. CGView blastn comparison of *S. agnetis* **1379, CBMRN, 908 and 1416**

genomes. The *S. agnetis* 1379 2.4 Mbp chromosome was the reference for blastn comparisons

with *S. agnetis* cattle isolate CBMRN (pink), and chicken isolates; 908 (green), and 1416

(blue). All other details are as in Figure 3 legend.