Targeted and passive environmental DNA approaches outperform

established methods for detection of quagga mussels, *Dreissena*

rostriformis bugensis in flowing water

Author details:

Rosetta C Blackman[1,2,3*], Kar Keun Sean Ling [3], Lynsey R Harper[3,4], Peter Shum[3,4],

Bernd Hänfling[3] and Lori Lawson Handley[3]

1. Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic Ecology,

Überlandstrasse 133, CH-8600 Dübendorf, Switzerland.

2. Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstr.

190, CH-8057 Zürich, Switzerland.

3. Evolutionary and Environmental Genomics Group (EvoHull), Department of Biological and Marine

Sciences, University of Hull, Cottingham Road, Hull, HU6 7RX, UK

4. School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, L3 3AF,

UK

*Corresponding author: Rosetta Blackman rosiecblackman@gmail.com

Keywords: eDNA, early detection, invasive non-native species, targeted assay, metabarcoding, PCR,

qPCR

Abstract

1. The early detection of invasive non-native species (INNS) is important for informing management actions. Established monitoring methods require the collection or observation of specimens, which is unlikely at the beginning of an invasion when densities are likely to be low. Environmental DNA (eDNA) analysis is a highly promising technique for the detection of INNS – particularly during the early stages of an invasion.

2. Here, we compared the use of traditional kick-net sampling with two eDNA approaches (targeted detection using both conventional and quantitative PCR, and passive detection via metabarcoding with conserved primers) for detection of quagga mussel, *Dreissena rostriformis bugensis;* a high priority INNS, along a density gradient on the River Wraysbury, UK.

3. All three molecular tools outperformed traditional sampling in terms of detection. Conventional PCR and qPCR both had 100% detection rate in all samples, and outperformed metabarcoding when the target species was at low densities. Additionally, quagga mussel DNA copy number (qPCR) and relative read count (metabarcoding) were significantly influenced by both mussel density and distance from source population, with distance being the most significant predictor.

4. *Synthesis and application.* All three molecular approaches were more sensitive than traditional kick-net sampling for the detection of the quagga mussel in flowing water, and both qPCR and metabarcoding enabled estimates of relative abundance. Targeted approaches were more sensitive than metabarcoding, but metabarcoding has the advantage of providing information on the wider community, and consequently impacts of INNS.

44 **Introduction**

Invasive non-native species (INNS) have wide ranging effects on ecosystems, including biodiversity loss

(Doherty et al., 2016; Gallardo et al., 2016) and economic impacts including damage to infrastructure

(Pimentel et al., 2005; Connelly et al., 2007). Eradication or containment methods are most effective

during the early stages of an invasion, both in terms of success and financial cost (Hulme, 2006; Mehta

et al., 2007). Early detection is particularly challenging for species that are small, elusive or cryptic and

these species often go unnoticed until they become established (Simmons et al., 2015; Blackman et al.,

2017). Monitoring organisms such as freshwater invertebrates generally relies on specimen collection

and morphological identification, and newly invading species can be easily overlooked. Alternative

methods that offer rapid and cost-effective detection of new INNS are crucial at a time when rapid

increase in trade and transport has caused a surge in the number of successful invasions to new

environments (Hulme, 2006).

The use of molecular methods to identify DNA taken from environmental samples (i.e. environmental

DNA or "eDNA") has been a major research focus in biodiversity monitoring over the last 10 years,

including the detection of INNS (Ficetola et al., 2008, Thomsen et al., 2012, Blackman et al., 2018). Two

broad approaches can be applied to detect a species of interest. The "targeted" approach utilises

species-specific primers with conventional PCR (cPCR), quantitative PCR (qPCR) or droplet digital PCR

(ddPCR) to detect a single species. By contrast, metabarcoding can be considered a "passive approach"

as it does not focus on single species (Lawson Handley, 2015; Creer et al., 2016; Deiner et al., 2017).

Metabarcoding utilises conserved primers to amplify DNA from a group of taxa e.g. fish and other

vertebrates (e.g. Kelly et al., 2014; Miya et al., 2015; Simmons et al., 2015), zooplankton (e.g. Brown et

al., 2016), or insects (e.g. Yu et al., 2012., Elbrecht et al., 2016). Amplicons are then sequenced using

High Throughput Sequencing (HTS) platforms and bioinformatically assigned to taxa. Metabarcoding

provides an opportunity to detect multiple species simultaneously, including those, which have not been

listed as priority INNS (Simmons et al., 2015; Blackman et al., 2017). Comparisons between traditional

methods for biodiversity monitoring and targeted (Dejean et al., 2012; Tréguier et al., 2014) or passive

(Smart et al., 2015; Hänfling et al., 2016; Wilcox et al., 2016) molecular approaches have demonstrated

an increased probability of detection using eDNA methods.

The targeted approach was first used by Ficetola et al. (2008) to detect invasive American bullfrog

Lithobates catesbeianus, in lentic waterbodies and has since been applied to over 100 target INNS

(Blackman et al., 2018). Conventional PCR is a well-established and widely practiced method for

detecting presence/absence of target DNA by visualisation of PCR products on agarose gels. However,

quantitative PCR (qPCR) has on many occasions been shown to be more sensitive than PCR (e.g. Nathan

et al., 2014; Simmons et al., 2015; De Ventura et al., 2016) and therefore more suitable for detecting

INNS at low density, e.g. the early stages of invasion. Quantitative PCR has the added advantage of

providing the number of DNA copies in a sample, which has been successfully linked to biomass or

abundance of some taxa (e.g. Amphibians, Thomsen et al., 2012; Fish, Takahara et al., 2012; Gastropods,

Goldberg et al., 2013). However, qPCR requires the construction of calibration curves from known

standard concentrations and is substantially more expensive and time-consuming than cPCR (Nathan et

al., 2014) and maybe less accessible to end-users than cPCR (Blackman et al., 2020).

The potential of eDNA metabarcoding for INNS monitoring has already been shown as an effective tool

(Simmons et al., 2015; Brown et al., 2016; Blackman et al., 2017; Borrell et al., 2017; Holman et al.,

2019). These studies include the detection of new INNS, which had been previously overlooked by

traditional methods. For example, Blackman et al. (2017) used a metabarcoding approach to investigate

macroinvertebrate diversity and discovered a non-native Gammaridae species, *Gammarus fossarum*,

which traditional methods had failed to identify for over 50 years in the UK. Similarly, Simmons et al.

(2015) detected the Northern Snakeshead, *Channa argus*, in the Muskingum River Watershed, Ohio,

which had not been included on INNS priority lists. Metabarcoding also has the potential to act as a

surveillance tool and has been applied to monitor high-risk pathways, such as ballast water (Ardura et

al., 2015; Zaiko et al., 2015), the live bait trade (Mahon et al., 2014) and ports (Brown et al., 2016;

Borrell et al., 2017). However, comparisons of targeted and passive approaches for INNS or other focal

species have been exemplified in the bighead carp, *Hypophthalmichthys nobilis* (Simmons et al., 2015)

and great crested newts, *Triturus cristatus* (Harper et al., 2018). These studies both showed that a

targeted approach had a higher sensitivity than the passive approach. Possible causes of this include

primer choice and PCR bias, i.e. species successfully amplifying and masking others in the PCR reaction

during metabarcoding (Harper et al., 2018). Further comparisons of these methods covering different

target taxa are required to better inform management strategies.

In this study we focus on the quagga mussel, *Dreissena rostriformis bugensis*, a highly invasive bivalve

from the Ponto-Caspian region, which has a significant impact on all trophic levels within the

environments it invades (Karatayev et al., 2007; Roy et al., 2014). In a horizon scanning exercise from

2014, the quagga mussel scored the highest out of 591 species evaluated for its potential to invade,

establish and have negative impact in the UK (Roy et al., 2014). Consistent with predictions, quagga

mussels were subsequently detected later the same year (Aldridge et al., 2014). Due to its wider

tolerance to environmental conditions, it has the potential to spread rapidly, and occupy a greater range

of habitat than the closely related zebra mussel, *Dreissena polymorpha,* which has already spread

throughout much of Europe and North America (Nalepa et al., 2010; Gallardo and Aldridge 2013; Quinn

et al., 2014). Motivated by a need for cost effective tools for quagga mussel detection, we previously

designed a cPCR assay, which provided 100% detection rate in field trials with quagga mussel

populations (Blackman et al., 2020). Here we further developed and validated this assay for abundance

estimation using a dye-based qPCR approach in mesocosm experiments and field trials. Secondly, we

directly compared the three molecular methods (cPCR, qPCR and metabarcoding) to traditional kick-net

sampling for detection of quagga mussels in the field. Finally, we investigated whether qPCR and

metabarcoding are suitable for estimating relative abundance of quagga mussels by sampling a density

gradient away from the main population. We hypothesized that: (1) the probability of detecting quagga

mussels will be higher for eDNA methods than traditional methods, (2) targeted methods will have a

higher probability of detecting quagga mussel than the passive method, (3) qPCR will be more sensitive

than cPCR for quagga mussel detection and (4) both read count and DNA copy number will decline with

distance downstream of the main population.

Methods

Field sample collection, DNA capture and extraction

Quagga mussels were first detected in the UK in Wraysbury Reservoir, Berkshire, in October 2014. This

site is considered the founding population of quagga mussels in the UK. The mussels spread through the

outfall and along the adjacent River Wraysbury, which is a tributary of the River Colne. The species

currently has a restricted distribution in the South East of England but has also been recorded in

reservoirs adjacent to the River Lee, north of London, and in the River Thames. Samples were collected

from Wraysbury Reservoir and River in May 2016, working in an upstream direction to ensure DNA

availability and inhibitor levels were not altered by sampling activity. In order to compare established

and molecular methods directly, we collected eDNA and kick-net samples at six sites across the river at

each of three main sampling locations downstream of Wraysbury Reservoir: Wraysbury Weir (WW, 0.61

km downstream of Wraysbury Reservoir), Wraysbury Bridge (WB, 1.10 km), and Wraysbury Gardens

(WG, 2.7 km) (orange diamonds, Fig. 1 and Supplementary Information I Table S1). Water samples from

these three locations consisted of three 500 ml replicates, collected at each of six sites across the width

of the river (n = 54). eDNA samples were collected from the river surface, using sterile gloves and

Gosselin™ HDPE plastic bottles (Fisher Scientific UK Ltd, UK) prior to kick-net sampling. Three-minute

kick-net samples were collected from the same six sites at each of these three locations, in accordance

with established methods (Murray-Bligh, 1999). eDNA sampling of three 500 ml replicates was also

carried out at four additional locations along Wraysbury River (Reservoir Outfall, RO, 0.2 km; Moor Lane,

ML, 1.7 km; Hale Street, HS, 2.72 km; and Upper Thames Confluence, UT, 2.75 km, blue diamonds, Fig. 1,

Supplementary Information I Table S1). These additional eDNA samples were collected to provide

additional fine-scale resolution in the eDNA analysis, and were analysed with PCR and qPCR only,

whereas samples from the three main locations were analysed with PCR, qPCR and metabarcoding.

Sampling locations within the river were chosen based primarily on accessibility, in accordance with

standard macroinvertebrate monitoring procedures carried out by UK agencies. We did not target

specific locations with previous records of quagga mussels, or their preferred habitat. At the time of

sampling, quagga mussels had not been reported upstream of the reservoir. We therefore collected an

eDNA sample upstream of the reservoir with the expectation that this would be negative for quagga

mussel (Waterside Car Park, n = 3). Sample bottles filled with ddH$_2$O were also taken into the field as

filtering blanks (n = 3). Hence the total number of eDNA samples collected in the field was n = 72. All

eDNA samples and blanks were stored in a cool box with ice packs and transported to the laboratory for

filtration within 24 hours. Water samples were filtered through 0.45 μm cellulose nitrate membrane

filters attached to a vacuum pump and DNA extractions performed following Brolaski et al. (2008) with

minor modifications (see Supplementary Information I and II for full details). Kick-net samples were

placed in individual 42 fluid oz. sterile Whirl-pak® bags (Cole-Palmer, Hanwell, London) and stored in a

cool box before being frozen at -20°C in the laboratory. Each sample was thawed at room temperature

and rinsed with ddH$_2$O separately prior to analysis. Samples were then sorted for the presence of *D. r.*

bugensis only and all specimens were counted and subsequently stored at -20°C.

Performance of targeted molecular detection

The performance of targeted DNA assays for the detection of invasive quagga mussel DNA was first

examined through laboratory-based mesocosm experiments. The objectives of these experiments were

a) to validate species-specific DNA assays, using both conventional and quantitative PCR, to target eDNA

detection of quagga mussels; b) to assess the duration and persistence of eDNA signal over a 42-day

period; and c) to examine a correlation between eDNA concentration and known densities of quagga

mussel. The mesocosm experiment was set up to monitor DNA production of mussels at three densities

(1, 5 and 20 individuals, with total and individual biomass weights recorded) over 21 days, with three

replicates of each density and a control tank with no mussels present. Fifteen litre plastic tanks with

individual aeration stones were set up in a climate controlled facility where temperature averaged 16 °C

(range 14–18 °C), with light:dark cycles of 16 h:8 h. Sampling events took place over 42 days at: 0hrs

(prior to specimens being added), 4 hrs, 8 hrs, 24 hrs, 48 hrs, 72 hrs, 7 days, 15 days and 21 days. On day

21, the specimens were removed, and sampling continued at: 22 days, 28 days, 35 days, 42 days (n =

130). Two hundred millilitre water samples were collected at each time point and each sample was

filtered through a single filter membrane following the same protocol as the field samples (Blackman et

al., 2020).

Conventional PCR (cPCR) assay: We used a cPCR assay to detect the quagga mussel, *D. r.*

bugensis, using a species-specific primer pair targeting a 188 bp fragment of the cytochrome oxidase I

(COI) gene (DRB1_F – 5'-GGAAACTGGTTGGTCCCGAT3'- and DRB1_R 5'-GGCCCTGAATGCCCCATAAT-3',

see Supplementary Material I for full PCR protocol). The DRB1 PCR assay has shown to be highly

sensitive (positive detection with 0.001 ng of target tissue DNA per reaction) and does not cross amplify

closely related Dreissenid or native mussel species (Blackman et al., 2020). All mesocosm (n=130), field

samples (n = 66) and control samples (field (3), filter (3) and PCR (6) blanks) were tested with the DRB1

assay. A tissue DNA sample from closely related zebra mussel was included as an additional negative

control.

Quantitative PCR (qPCR) assay: The DRB1 assay was further developed using fluorescent dye-

based qPCR (PowerUp SYBR green, Thermo Fisher, UK, see Supplementary Material I for full details).

Briefly, we performed serial dilutions of DNA from tissue samples (from 1 to 1×10^{-7} ng/µl) to estimate

the LoD, and performed the DRB1 qPCR assay on samples collected from the mesocosm experiment,

field an control samples. Zebra mussel tissue DNA was used as a negative control in the qPCR assays.

Amplification curves, Cq values and melt curves were analysed using StepOne-PlusTM software 2.0.

Quantification calibration curves using a synthesized gBlock of the target fragment was used to estimate

the DNA copy number for each sample (See supplementary Information I and IV for full details).

Passive detection: eDNA metabarcoding

Degenerate metazoan primers that target the conserved COI region were used for metabarcoding

(Geller et al., 2013 and Leray et al., 2013): jgHCO2198: TAIACYTCIGGRTGICC RAARAAYCA and mICOIintF:

GGWACWGGWTGAACWGTWTAYCCYCC. These primers were chosen as they i) target the same

mitochondrial gene as our targeted assays, ii) are the most commonly used COI metabarcoding primers

and are good candidates for studying the entire river community, and iii) we confirmed in previous work

that they amplify Dreissenid mussels. We used universal primers rather than mollusc specific

metabarcoding primers (which are also available, Klymus et al., 2017), since universal primers are more

likely to be useful in routine community monitoring of INNS. We confirmed amplification of the target

taxa from tissue extractions prior to library preparation. Full details of the library preparation are

provided in Supplementary Information I. In brief, the 54 field samples from WW, WB and WG (Fig. 1)

were sequenced, with the addition of 3 field blanks, 3 filter blanks, 6 PCR blanks and 3 positive non-

target tissue samples (2 each of *Triops cancriformis and Osmia bicornis total n* = 69). A two-step PCR

protocol was used for library preparation: first the target region was amplified for each sample and a

second amplification was performed to add the Illumina adaptor sequences onto the initial amplicons.

Samples were normalized following the second PCR using SequalPrep Normalization plates (Invitrogen,

UK) and samples were then pooled in equimolar concentrations. The final pooled library was

concentrated, cleaned with a QIAquick Gel Extraction Kit (Qiagen, UK), quantified and sequenced at 13

pM concentration with 20% PhiX, on an Illumina MiSeq platform using 2 x 250bp v2 chemistry at the

University of Hull.

Bioinformatics and data analysis

Full details of the bioinformatics pipeline are provided in Supplementary Material I. In brief, processing

of Illumina data and taxonomic assignment were performed using a custom pipeline, MetaBEAT

(metaBarcoding and eDNA Analysis Tool) v 0.8 (https://github.com/HullUnibioinformatics/metaBEAT)

but with minor modifications including filtering for the presence of the target taxa, *D. r. bugensis,* by

comparing against curated reference sequences from GenBank, listed in Supplementary Material I.

We normalised the metabarcoding data to account for uneven sequencing depth between samples, by

dividing the number of reads for each OTU within a sample by the total number of reads in that sample

and multiplying by 100 (hereafter referred to as relative read count).We used Generalized Linear Models

(GLMs) to investigate 1) the influence of mussel density, biomass and time on the DNA copy number

(qPCR) since the start of the mesocosm experiment (before the mussels were removed from the tanks

only), and 2) the influence of mussel density and distance from source population on the DNA copy

number (qPCR) from the field samples and relative read count (metabarcoding). Initial model tests

showed all data sets were overdispersed, therefore a negative binomial GLM with a log link function was

used to resolve this issue. To check for model suitability, we checked the residual deviance of each

model fitted a chi square distribution (see Supplementary Information III Tables S2, S3 and S4 for GLM

results). Finally, we assessed the relationship between DNA copy number and relative read count with

the distance along the river using all seven sites included in the qPCR experiment, using a Spearman's

Rank Correlation since the data were not normally distributed. All calculations and visualizations of the

data were carried out in R (R-Core-Team 2017), with GLMs performed using the MASS package

(Venables et al., 2002).

Results

We successfully adapted the DRB1 primer pair for detection of quagga mussel with the addition of SYBR

green. Across both traditional and eDNA field experiments we were able to detect quagga mussel with

kick-net sampling, cPCR, qPCR and metabarcoding. All negative controls (field blanks, filtration blanks,

PCR blanks and non-target tissue blanks were negative for cPCR and qPCR. Minor contamination was

detected in 3 field blanks with metabarcoding, with low reads of *Homo sapiens (5 and 14 reads), Triops*

cancriformis (the positive tissue extract) (10 reads) and unassigned reads (42-189 reads). We therefore

applied a 0.2% read threshold to remove possible contamination from the metabarcoding dataset (i.e.

sequences that had a frequency of 0.2% or less in each sample were removed).

qPCR assay development and validation in mesocosms

The limit of detection (LoD) determined from the serial dilutions of tissue DNA was 1×10^{-4} ng/µl per

reaction with both cPCR and qPCR, which equates to approximately 5 copies per reaction (See

Supplementary Information III Fig. S1 and Fig. S2 for amplification plots). Rapid DNA accumulation in the

first two days of the mesocosm experiments was followed by a depletion in DNA concentration until

removal of mussels on day 21 of the experiment (Fig. 2). This pattern of DNA accumulation and

depletion is highly consistent across all three density treatments. Time since the beginning of the

experiment (until removal of the mussels) had a significant effect on the DNA copy number in each

mesocosm experiment when all density treatments were analysed together (GLM, Z = -7.224, P <

0.0001, AIC = 912.95). There is still some overdispersion with this model, however the overdispersion

parameter indicates this is not substantial (overdispersion statistic = 1.3, Residual Deviance 91.142 on 70

df, χ^2 P = 0.0457). There is substantial overlap between the DNA copy number estimates for the three

density treatments (Fig. 2) and neither density nor total biomass significantly explained the number of

DNA copies amplified by qPCR (Density Z = -1.019, P = 0.308; Total biomass Z = -0.776, P = 0.438). DNA

was still detectable 24 hours after removal of the mussels, but not at 7 days after removal (Day 28) (Fig.

2). All blanks and mesocosm control tank samples were negative of target DNA throughout the

experiment.

Detection of D. r. bugensis by traditional and molecular approaches in the field

Kick-net sampling

The number of *D. r. bugensis* specimens varied from 14 at Wraysbury Weir, the location closest to the

reservoir outfall (source population), to a single specimen at Wraysbury Gardens, the location furthest

from the outfall (Table 1). Detection rate was also highest at Wraysbury Weir (mussels detected in 4/6

sampling events). Mussels were detected in 1/6 samples at each of the other two sampling locations

(Table 1).

Targeted approaches

Detection rate was 100% in all three field locations and all replicates tested for both targeted

approaches (cPCR and qPCR) (Table 1 and Fig. S3). Fig. 3A shows the DNA copy number (log_{10}) of the

qPCR assay for *D. r. bugensis* found at each of the sampling locations. Both distance from source and

number of mussels found at the three locations, significantly influenced the DNA copy number, but

distance was the strongest predictor (Density Z = -3.367, p = 0.000761; Distance Z = -17.223, p < 0.0001;

AIC: 1392.1, Residual Deviance 57.621 on 51 df, χ^2 P = 0.244). A Spearman's Rank correlation between

the DNA copy number and distance from the source population at each of the seven locations showed a

weak correlation (ρ = -0.25, p = 0.5948). However, the low copy numbers found immediately

downstream of the reservoir outfall (Fig. 3A) affected this. If this site is treated as an outlier and

removed from the analysis, the correlation is significant (ρ = -1, p = 0.002778).

Passive approach

The total number of reads produced following quality control and chimera removal was 4,928,265, of

which 576,365 reads were assigned to the samples in this study. A total of 5.19% reads (29,936) were

assigned to *D. r. bugensis.* After applying a 0.2% threshold to remove minor contamination the average

read depth per sample was 2921 over all taxa, and 548 for *D. r. bugensis* (excluding controls). The

clustering procedure identified 81 OTUs revealing a range of macroinvertebrate and algae/diatom

species (Supplementary information III, Fig. S4). The proportion of reads assigned to *D. r. bugensis* from

each sample within each location varied greatly as seen in Fig. 3B (n = 54). The target species was

detected in all samples from both Wraysbury Weir and Wraysbury Bridge locations, while detection at

Wraysbury Gardens was much lower with only 3 of the 18 replicates present for *D. r. bugensis* (Table 1).

Detection rate was therefore 86%, and is lower than the targeted methods. The average raw read count

of *D. r. bugensis* was 843 and 801 at Wraysbury Bridge and Wraysbury Weir respectively, and only 1

read on average at Wraysbury Gardens. Distance was the strongest predictor of the relative read count

(Z = 13.47, p < 0.0001; AIC = 301.86, Residual Deviance 54.989 on 52 df, χ^2 P = 0.36) in agreement with

the qPCR results and when compared to the distance and density model (Distance- Z = -4.707, p <

0.0001; Density Z = -3.773, p = 0.0001; AIC = 258.64, Residual Deviance 49.696 on 51 df, χ^2 P = 0.820),

and density on its own (which was overdispersed).

312 **Discussion**

This study investigates the detection, persistence and quantification of the invasive non-native quagga

mussel (*D. r. bugensis*) using kick-net sampling and environmental DNA (eDNA). Through experimental

mesocosms, we validated a species-specific DNA assay (cPCR) to detect quagga mussel and further

developed a dye-based quantitative PCR (qPCR) assay to evaluate the performance of both targeted

approaches. We carried out an eDNA survey of water samples and compared targeted (cPCR and qPCR)

and passive (DNA metabarcoding) approaches to traditional kick-net sampling for qualitative and semi-

quantitative detection in the River Wraysbury, UK. In line with our first hypothesis, detection rate was

higher for all three eDNA methods than with the traditional method. Secondly, targeted approaches

were more sensitive than the passive approach (100% detection rates versus 86%). Contrary to our third

hypothesis, we observed no increased sensitivity using a dye-based qPCR over cPCR. Lastly, both density

and distance from the source population were significant predictors of relative read count

(metabarcoding) and copy number (qPCR) indicating that both approaches could provide estimates of

quagga mussel abundance in the field.

qPCR validation

Mesocosm validation provides an opportunity to test the LoD of eDNA assays and to investigate the

relationship between density, time and DNA copy number (Dejean et al., 2011; Thomsen et al., 2012;

Blackman et al., 2020). DNA was detected with qPCR at all three densities (1, 5 and 20 individuals) within

4 hours of being present, in agreement with previous cPCR results (Blackman et al., 2020). The LoD from

serial dilutions of tissue DNA were also identical for cPCR and qPCR (1×10^{-4} ng/µl). Surprisingly, the

mesocosm experiments highlighted unexpected dynamics of DNA in closed systems. Firstly, time since

the start of the experiment was a significant predictor of DNA copy number, but rather than increasing

over time or reaching a plateau, DNA accumulated quickly in the first 48 hours then rapidly depleted,

and this trend was seen in all three density treatments. A possible explanation for this is stress induced

on the mussel caused by a change in environment, as reported in other eDNA studies (Li et al., 2019).

Dreissenid mussels, however, are known to "shut-down" in times of stress, such as chemical dosing, and

can remain with their shells closed for up for three weeks (Aldridge et al., 2006). A further explanation

for the rapid accumulation in DNA may be due to the production of protein based byssal threads which

allow the mussels to attach to the substrate (Ricciardi et al., 1998; Aldridge et al., 2004; Timar and

Phaneuf, 2009; Peyer et al., 2009). During the first 48 hours of the experiment, the mussels are likely to

secure themselves to the new substrate. All mussels were found to have done this when they were

removed at the end of the experiment. The mussels are also filter-feeders and were actively feeding (on

microorganisms found in the river water) throughout the experiment producing sloughed cells, faeces

and pseudofaeces (Sansom and Sassoubre, 2017). This feeding behaviour is likely to be a major

contributor to successful detection in the field, and in a closed system, effect the DNA copy number

detected with the qPCR assay.

The second surprising result from the mesocosm experiments was that we found no effect of density or

biomass on DNA copy number. This contradicts several previous studies that have demonstrated a

relationship between biomass or density and DNA copy number (Thomsen et al., 2012; Takahara., et al

2012; Doi et al., 2015). This could in part be due to the physiology and feeding behaviour of mussels,

since eDNA is being consumed as well as produced (Friebertshauser et al., 2019). The complex pattern

of eDNA accumulation and depletion together with the lack of discrimination between different density

treatments presents challenges for estimation of quagga mussel abundance. However, eDNA is unlikely

to accumulate in the same way in natural (particularly lotic) environments, and this case highlights that

mesocosm experiments may poorly reflect the true relationship between DNA copy number and mussel

density in the field. It would be worthwhile to examine whether this pattern is also observed with other

filter feeding species. In keeping with previous studies (e.g. Thomsen et al., 2012), rapid degradation of

eDNA was observed in mesocosm experiments, with a complete loss of signal by one week after mussel

removal. Unfortunately, our experiments did not allow us to fully investigate the degradation rate of

eDNA in mesocosms since our sampling was too limited. We recommend future studies carry out more

intensive sampling once specimens have been removed.

Comparison of eDNA and kick-net sampling for quagga mussel detection

The lower sensitivity of kick-net sampling is highlighted by the failure to collect a *D. r. bugensis* specimen

in any of the first kick-net sampling attempts at each site. As demonstrated in a previous study, species

in low abundance are likely to be missed, with only 62% of macroinvertebrate families detected in a

three-minute kick-net sample (Furse et al., 1981). This highlights the need for multiple replicates for

detection of low abundance species if using kick-net sampling. It should be noted that surveillance

methods specifically for *D. r. bugensis* would be better focussed on the preferred habitat of mussels i.e.

hard substrates such as man-made structures (Aldridge et al., 2014), which is more likely to detect

quagga mussels than the randomised method used here. However, randomised kick-net sampling along

a range of substrate types is the most widely used sampling method for freshwater macroinvertebrate

monitoring, and therefore our experiment reflects the most likely methods to detect new INNS through

routine monitoring schemes.

Of the three molecular approaches compared in this study, we show that both targeted approaches

were more sensitive for detecting *D. r. bugensis* than metabarcoding, but that metabarcoding still

performed encouragingly well, given that the assay targeted metazoan in general rather than the focal

species. Few studies have compared targeted approaches with metabarcoding; however, our findings

are similar to those of Simmons et al., (2015) and Harper et al., (2018), who showed that metabarcoding

using conserved primers was less sensitive than targeted approaches. In this study, there was no

increase in the detection of *D. r. bugensis* when the 0.2% contamination threshold was removed.

However, it would be pertinent to examine the raw output of metabarcoding runs before thresholds are

applied in order to identify any potential INNS with low read counts. As this form of INNS monitoring is

still in its infancy, quality control measures to ensure these detections are correct are yet to be put in

place (Darling et al., 2020). We would therefore recommend any new INNS detections with

metabarcoding are followed up with species-specific primers or collection of specimens prior to

management actions to verify the detection.

A significant advantage of metabarcoding with conserved primers is the added information from non-

target species detection: in our study a further 81 OTUs were identified, and although no additional

INNS were detected, previous studies have demonstrated the power of this approach for detecting new

and unexpected INNS (Blackman et al., 2017). Metabarcoding also provides the opportunity to monitor

changes in community composition as the result of INNS or other ecological stressors (Simmons et al.,

2015; Hänfling et al., 2016), which is extremely promising for better understanding INNS impacts for

more effective prioritisation and management. This information on the wider community is a

justification for the use of universal primers rather than more taxon-specific metabarcoding assays, but

if more targeted information is needed, more specific metabarcoding primers can be developed. Indeed,

specific mollusc metabarcoding primers have recently been developed, and are very promising for more

targeted investigation of mollusc communities (Klymus et al., 2017). We hypothesize that more focussed

metabarcoding assays will yield even higher detection rates than the one reported here, reducing the

difference in sensitivity between targeted and passive approaches. It could also be possible to improve

the power of metabarcoding by increasing spatial sampling. Future studies should investigate how many

samples are needed in order to maximise the probability of detecting rare species with metabarcoding

and also the required read depth needed per sample in order to maximise species detection at a given

sampling site.

Conventional PCR and dye-based qPCR had identical detection rates for *D. r. bugensis* in our study. This

is somewhat surprising as qPCR is generally considered to be more sensitive than cPCR (e.g. Thomsen et

al., 2012, Nathan et al., 2014). Conventional PCR has the advantage of being quick and cost-effective,

requiring limited lab equipment and up to half the time for sample processing than qPCR (Davison et al.,

2016). Drawbacks to cPCR are typically lower sensitivity, and gel-quality issues that make interpretation

of results more subjective than with qPCR. This is dependent on the primer specificity and optimisation

of the PCR. Our conventional *D. r. bugensis* assay is highly specific and fully optimised, yielding clear

bands on agarose gels even in samples with low concentration of target species DNA (as inferred from

qPCR). However, we acknowledge that our experience with this assay may be unusual. An obvious

advantage to qPCR is the quantitative information it provides, in addition to increased sensitivity. The

added information on DNA concentration from qPCR in this study yielded important insights into the

mussel density and distance from the source population (see below), which was not possible with cPCR.

An important consideration is whether the assay could be improved by using probe-based rather than

dye-based qPCR. Addition of a probe is generally considered beneficial to prevent the possibility of non-

target amplification. Given the high specificity of the DRB1 primer pair described here, we believe the

addition of a probe is not necessary in this case, but we stress that adding a probe should be considered

if an extra layer of specificity is needed and/or if non-target amplification is possible.

Quagga mussel eDNA declines with distance from the source and decreasing mussel density

Results from field trials for both dye-based qPCR and metabarcoding demonstrate a significant decrease

in DNA copy number and relative read count of quagga mussel, respectively, with increasing distance

from the main source population (Wraysbury reservoir) and with decreasing mussel density. Kick-net

sample data also indicates that mussel density decreases along the course of the river, with only one

mussel found at Wraysbury Gardens, 2.7 km downstream of the reservoir. The only site that did not fit

this trend was the first site, 20 m downstream of the outfall from Wraysbury reservoir, which had a very

low DNA copy number. The decrease in concentration of quagga mussel eDNA detected with distance

from Wraysbury Reservoir, confirmed with both qPCR and metabarcoding, reflects both the decreasing

density of mussels and a gradient in the transport of DNA downstream. The fact that eDNA

concentration is not homogeneously distributed in the river suggests that it may be possible, at least in

some cases, to infer the source population in cases where it is unknown. This is somewhat surprising

given that our sampling took place after a period of heavy rain and eDNA distribution in rivers is often

considered to be uniform and heavily confounded by flow rate and mixing (e.g Jane et al., 2015).

Characteristics including discharge (Jane et al. 2015), physical structure and substrate type (Wilcox et al.,

2016; Shogren et al., 2017) and water chemistry (Seymour et al., 2018) all influence the transport and

retention of eDNA in lotic systems. Further empirical studies in rivers of all sizes and characteristics,

ideally in combination with hydrological modelling (Carraro et al., 2018), are needed to fully understand

the influence of flow dynamics and substrate type on retention and transport of eDNA in rivers.

Estimating abundance with eDNA is challenging because of the combined physical, chemical and

biological factors that influence production, transport and retention of DNA in a water body (Barnes et

al., 2014, Barnes and Turner 2015). Abundance estimation is expected to be particularly challenging in

rivers due to their complex hydrodynamics, and the transport of eDNA from upstream (e.g. Jerde et al.,

2016). Abundance estimation with metabarcoding is also controversial as read counts are influenced by

bias during sampling, laboratory and bioinformatics steps (Valentini et al., 2016). Despite these

challenges, our data add to the growing body of evidence that both qPCR and metabarcoding can yield

insight into relative biomass, which has rarely been demonstrated in lotic environments (but see e.g. Doi

et al., 2016, Akamatsu et al., 2020). Temporal sampling and controlling for environmental variables,

combined with improved statistical approaches such as site occupancy modelling, which is becoming

increasingly popular in eDNA studies (e.g. Schmidt et al., 2013, Pilliod et al., 2013, Chambert et al., 2018,

Sales et al., 2020) or spatially explicit approaches that take hydrological conditions into account (e.g.

Shogren et al., 2017, Carraro et al., 2018) will provide better quantitative information, and allow

changes in biomass to be monitored.

Conclusions and practical considerations for future INNS biomonitoring with eDNA

We have demonstrated that eDNA from quagga mussels can be successfully detected using cPCR, qPCR

and metabarcoding in a UK river. Both qPCR and metabarcoding provide quantitative information, with

DNA copy number and relative read count significantly decreasing with a combination of distance from

the source population and mussel density. In our previous work we demonstrated that metabarcoding is

an effective tool for passive detection of a previously unrecorded cryptic non-native species, *Gammarus*

fossarum (Blackman et al. 2017). The present study provides further demonstration that metabarcoding

with conserved metazoan primers is suitable as an early warning tool for non-targeted monitoring or

routine surveillance even at a very low population density. The optimal approach for detection of INNS

using eDNA will depend on the objectives of the research or monitoring programme. For early detection

of previously unrecorded INNS, we recommend a comprehensive, randomised sampling design,

combined with metabarcoding using universal primers, which can be supplemented with targeted

approaches if additional verification is needed. Targeted qPCR assays provide optimal sensitivity and

specificity and are particularly suited to surveys of INNS that have been previously recorded. In terms of

future monitoring of quagga mussels, we believe that the sampling and laboratory methods adopted in

the present study are widely applicable, but a temporal study would be beneficial to determine the

optimum timing for sampling, and greater spatial coverage is needed to investigate the influence of

environmental variables on detection rates.

Acknowledgments

This project was funded by the Freshwater Biological Association Gilson Le Cren Award 2017 and the UK

Environment Agency. We would like to thank staff from the Environment Agency - Drs Alice Hiley and

Kerry Walsh for their advice and support and Dr Patrycja Meadows and Colette Sales for their help to

identify field sites, arrange access and collect specimens. We also thank two anonymous reviewers for

helpful and constructive suggestions on a previous version of our manuscript.

Author contributions

The design and concept of this study was developed by RCB and LLH. RCB and LH carried out the field

work. RCB, KKSL and PS performed the lab work. RCB analysed the data. RCB and LLH wrote the paper.

All authors have approved the final manuscript.

Data availability statement

Data generated in this study is available from the GitHub repository:

https://github.com/RosettaBlackman/Blackman_et_al_quagga

References

Akamatsu, Y., Kume, G., Gotou, M., Kono, T., Fujii, T., Inui, R. & Kurita, Y. (2020). Using environmental

DNA analyses to assess the occurrence and abundance of the endangered amphidromous fish

Plecoglossus altivelis ryukyuensis. *Biodiversity data journal*, 8, p.e39679. doi: 10.3897/BDJ.8.e39679.

Aldridge, D.C., P. Elliott, P., & Moggridge. G.D. (2004). The Recent and Rapid Spread of the Zebra Mussel

(*Dreissena Polymorpha*) in Great Britain. Biological Conservation, 119 (2):253–61. doi:

10.1016/j.biocon.2003.11.008

Aldridge, D.C., Elliott P., & Moggridge G.D. (2006). Microencapsulated BioBullets for the Control of

Biofouling Zebra Mussels. Environmental Science & Technology, 40 (3): 975–79. doi:

10.1021/es050614%2B

Aldridge, D. C., Ho, S., & Froufe. E. (2014). The Ponto-Caspian Quagga Mussel, *Dreissena Rostriformis*

Bugensis (Andrusov, 1897), Invades Great Britain. Aquatic Invasions / European Research Network on

Aquatic Invasive Species. doi: 10.3391/ai.2014.9.4.11

Ardura, A., Zaiko, A., Martinez, J.L., Samuiloviene, A., Borrell, Y., & Garcia-Vazquez. E. 2015

Environmental DNA Evidence of Transfer of North Sea Molluscs across Tropical Waters through Ballast

Water. The Journal of Molluscan Studies 81 (4). Oxford University Press: 495–501. doi:

10.1093/mollus/eyv022

Blackman, R.C., Constable, D., Hahn, C., Sheard, A., Durkota, J., Hänfling, B., & Lawson Handley. L.

(2017). Detection of a New Non-Native Freshwater Species by DNA Metabarcoding of Environmental

Samples — First Record of *Gammarus Fossarum* in the UK. Aquatic Invasions 12 (2):177–89. doi:

10.3391/ai.2017.12.2.06

Blackman, R.C., Hänfling, B., & Lawson Handley. L. (2018). The use of environmental DNA as an early

warning tool in the detection of new freshwater invasive non-native species. CAB Reviews Vol 12, 010,

pp1-15. doi: 10.1079/PAVSNNR201813010

Blackman, R.C., Benucci, M., Donnelly, R.C., Hänfling, B., Harper, L.R., Sellers, G.S. & Lawson Handley, L.

(2020) Simple, sensitive and species-specific assays for detecting quagga and zebra mussels (*Dreissena*

rostriformis bugensis and *D. polymorpha*) using environmental DNA. Management of Biological

Invasions. 11 (2), 218-236.

Borrell, Y.J., Miralles, L., Do Huu, H., Mohammed-Geba, K., & Garcia-Vazquez. E. (2017). DNA in a

bottle—Rapid Metabarcoding Survey for Early Alerts of Invasive Species in Ports. PloS One 12 (9). Public

Library of Science: e0183347. doi: 10.1371/journal.pone.0183347

Brown, E.A., Chain, F.C.C., Zhan, A., MacIsaac, H.J., & Cristescu. M.C. (2016). Early Detection of Aquatic

Invaders Using Metabarcoding Reveals a High Number of Non-Indigenous Species in Canadian Ports.

Diversity & Distributions. doi: 10.1111/ddi.12465

Carraro, L. Hartikaninen, H., Jokela, J., Bertuzzo, E. & Rinaldo, A. (2018) Estimating species distribution

and abundance in river networks using environmental DNA. *Proceedings of the National Academy of*

Sciences of the United States of America, 115(46), 11724–11729. doi: 10.1073/pnas.1813843115.

Chambert, T., Pilliod, D.S., Goldbrg, C.S., Doi, H. & Takahara, T. (2018) An analytical framework for

estimating aquatic species density from environmental DNA. *Ecology and evolution*, 8(6), 3468–3477.

doi: 10.1002/ece3.3764

Connelly, N.A., O'Neill, C.R., Knuth, B.A., & Brown, T.L. (2007). Economic impacts of zebra mussels on

drinking water treatment and electric power generation facilities. Environmental management, 40(1)

pp.105–112. doi: 10.1007/s00267-006-0296-5

Creer, S, Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, W.K., … & Bik. H.M. (2016). The

Ecologist's Field Guide to Sequence-Based Identification of Biodiversity. Methods in Ecology and

Evolution / British Ecological Society 7 (9):1008–18. doi: 10.1111/2041-210X.12574

Davison, P. I., Créach, V., Liang, W-J., Andreou, D., Britton, J.R., & Copp, G.H.. (2016). Laboratory and

Field Validation of a Simple Method for Detecting Four Species of Non-Native Freshwater Fish Using

eDNA. Journal of Fish Biology 89 (3):1782–93. doi: doi.org/10.1111/jfb.13086

Darling, J. A., Pochon, X., Abbott, C.L., Inglis, G.J. & Zaiko, A. (2020) The risks of using molecular

biodiversity data for incidental detection of species of concern. Diversity and Distributions. doi:

doi.org/10.1111/ddi.13108

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., & Creer, S. (2017).

Environmental DNA Metabarcoding: Transforming How We Survey Animal and Plant Communities.

Molecular Ecology 26 (21):5872–95. doi: 10.1111/mec.14350

Dejean, T., Valentini, A., Duparc, A., Pellier-Cuit, S., Pompanon, F., Taberlet, P., & Miaud. C. (2011).

Persistence of Environmental DNA in Freshwater Ecosystems. PloS One 6 (8) doi:

10.1371/journal.pone.0023398

Dejean, T., Valentini, A., Miquel, C., Taberlet, P., Bellemain, E., & Miaud. C. (2012). Improved Detection

of an Alien Invasive Species through Environmental DNA Barcoding: The Example of the American

Bullfrog Lithobates Catesbeianus. The Journal of Applied Ecology 49 (4) 953–59. doi: 10.1111/j.1365-

2664.2012.02171.x

De Ventura, L., Kopp, K., Seppälä, K., & Jokela. J. (2017). Tracing the Quagga Mussel Invasion along the

Rhine River System Using eDNA Markers: Early Detection and Surveillance of Invasive Zebra and Quagga

Mussels. Vol 8 (1) 101–112. doi:10.3391/mbi.2017.8.1.10

Doherty, T.S., Glen, A.S., Nimmo, D. G., Ritchie, E. G., & Dickman, C. R. (2016). Invasive predators and

global biodiversity loss. Proceedings of the National Academy of Sciences of the United States of

America, 113(40), pp.11261–11265. doi: 10.1073/pnas.1602480113

Doi, H., Uchii, K., & Takahara T, (2015). Use of droplet digital PCR for estimation of fish abundance and

biomass in environmental DNA surveys. PloS one 10(3). doi: 10.1371/journal.pone.0122763

Doi, H., Inui, R., Akamatsu, Y., Kanno K., Yamanaka, H., Takahara, T. & Minamoto, T. (2017)

Environmental DNA analysis for estimating the abundance and biomass of stream fish. *Freshwater*

biology, 62(1), 30–39. doi: 10.1111/fwb.12846.

Elbrecht, V., & Leese. F. (2017). Validation and Development of Freshwater Invertebrate Metabarcoding

COI Primers for Environmental Impact Assessment. Frontiers in Freshwater Science. 5(11), 1– 11. doi:

10.3389/fenvs.2017.00011

Ficetola, G.F., Miaud, C., Pompanon, F., & Taberlet. P. (2008). Species Detection Using Environmental

DNA from Water Samples. Biology Letters 4 (4):423–25. doi: 10.1098/rsbl.2008.0118

Friebertshauser R, Shollenberger K, Janosik A, Garner JT, Johnston C (2019) The effect of bivalve

filtration on eDNA-based detection of aquatic organisms. PLoS ONE 14(11): e0222830. doi:

10.1371/journal.pone.0222830

Furse, M. T., Wright, J.F., Armitage, P.D., & Moss, D. (1981). An Appraisal of Pond-Net Samples for

Biological Monitoring of Lotic Macro-Invertebrates. Water Research 15 (6):679–89. doi: 10.1016/0043-

1354(81)90160-3

Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial

cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys.

Molecular ecology resources, 13, 851–861. doi: 10.1111/1755-0998.12138

Gallardo, B., & Aldridge, D.C. (2013) The 'dirty Dozen': Socio-Economic Factors Amplify the Invasion

Potential of 12 High-Risk Aquatic Invasive Species in Great Britain and Ireland. The Journal of Applied

Ecology 50 (3):757–66. doi: 10.1111/1365-2664.12079

Gallardo, B., Clavero, M., Sánchez, M. I., & Vilà, M. (2016). Global ecological impacts of invasive species

in aquatic ecosystems. Global change biology, 22(1), pp.151–163. doi: 10.1111/gcb.13004

Goldberg, C S., Sepulveda, A., Ray, A., Baumgardt, J., & Waits. L.P. (2013). Environmental DNA as a New

Method for Early Detection of New Zealand Mudsnails (Potamopyrgus Antipodarum). Freshwater

Science 32 (3) 792–800. doi: 10.1899/13-046.1

Hänfling, B., Lawson Handley, L., Read, D.S., Hahn, C., Li, J., Nichols, P., … Winfield. I. (2016).

Environmental DNA Metabarcoding of Lake Fish Communities Reflects Long-Term Data from Established

Survey Methods. Molecular Ecology 25 (13):3101–19. doi: 10.1111/mec.13660

Harper, LR., Lawson Handley, L., Hahn, C., Boonham. N., Rees, HC., Gough, KC., … Hänfling, B. (2017).

Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of great

crested newt (Triturus cristatus). Ecology and Evolution 8 (12) 6330-6341. doi: 10.1002/ece3.4013

Holman, L.E., de Bruyn, M., Creer, S., Carvalho., Robidart., J., & Rius, M. (2019). Detection of introduced

and resident marine species using environmental DNA metabarcoding of sediment and water. *Scientific*

reports, 9(1), p.11559. doi: 10.1038/s41598-019-47899-7

Hulme, P.E., (2006). Beyond Control: Wider Implications for the Management of Biological Invasions.

The Journal of Applied Ecology 43 (5): 835–47. Doi: 10.1111/j.1365-2664.2006.01227.x

Jane, S. F., Wilcox, T.M., McKelvey, K.S., Young, M.K., Schwartz, M.K., Lowe, W.H., Letcher, B.G. &

Whiteley. A.R. (2015). Distance, Flow and PCR Inhibition: eDNA Dynamics in Two Headwater Streams.

Molecular Ecology Resources 15 (1):216–27. doi: 10.1111/1755-0998.12285

Jerde, C.L., Olds, B.P., Shogren, A.J., Andruszkiewicz, E.A., Mahon, A. R., Bolster, D. & Tank, J.L. (2016)

Influence of Stream Bottom Substrate on Retention and Transport of Vertebrate Environmental DNA.

Environmental science & technology, 50(16), 8770–8779. doi: 10.1021/acs.est.6b01761.

Karatayev, A.Y., Padilla, D.K., Minchin, D., Boltovskoy, D., & Burlakova. L.E. (2007). Changes in Global

Economies and Trade: The Potential Spread of Exotic Freshwater Bivalves. Biological Invasions 9 (2).

161–80. doi: 10.1007/s10530-006-9013-9

Kelly, R. P., Port, J.A., Yamahara, K.M., & Crowder, L.B. (2014). Using Environmental DNA to Census

Marine Fishes in a Large Mesocosm. PloS One 9 (1). doi: 10.1371/journal.pone.0086175

Klymus, K.E., Marshall, N.T. & Stepien, C.A. (2017) Environmental DNA (eDNA) metabarcoding assays to

detect invasive invertebrate species in the Great Lakes. *PloS one*, 12(5), p.e0177643. doi:

10.1371/journal.pone.0177643.

Lawson Handley, L. (2015). How Will the 'molecular Revolution' Contribute to Biological Recording?

Biological Journal of the Linnean Society. Linnean Society of London 115 (3):750–66. doi:

10.1111/bij.12516

Leray, M., Yang, J.Y., Meyer, C.P., Mills, S.C., Agudelo, N., Ranwez, V., … Machida, R.J. (2013). A New

Versatile Primer Set Targeting a Short Fragment of the Mitochondrial COI Region for Metabarcoding

Metazoan Diversity: Application for Characterizing Coral Reef Fish Gut Contents. Frontiers in Zoology

10:34. doi: 10.1186/1742-9994-10-34

Li, J., Lawson Handley, L., Harper, L.R., Brys, R., Watson, H.V., Di Muri, C., … Hänfling, B. (2019). "Limited

Dispersion and Quick Degradation of Environmental DNA in Fish Ponds Inferred by Metabarcoding."

Environmental DNA 1 (3): 238–50.

Mahon, A.R., Nathan, L.H., & Jerde, C.L. (2014). Meta-Genomic Surveillance of Invasive Species in the

Bait Trade. Conservation Genetics Resources 6. doi: 10.1007/s12686-014-0213-9

Mehta, S.V., Haight, R. G., Homans, F. R, Polasky, S & Venette, R.C. (2007). Optimal detection and

control strategies for invasive species management. Ecological economics: the journal of the

International Society for Ecological Economics, 61(2), pp.237–245. doi: 10.1016/j.ecolecon.2006.10.024

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., … Iwasaki, W. (2015). MiFish, a Set of

Universal PCR Primers for Metabarcoding Environmental DNA from Fishes: Detection of More than 230

Subtropical Marine Species. Royal Society Open Science 2 (7). doi: 10.1098/rsos.150088

Murray-Bligh J.A.D. (1999). Procedure for Collecting and Analysing Macroinvertebrate Samples,

Environment Agency, Bristol, UK.

Nathan, L.M., Simmons, M., Wegleitner, B.J., Jerde, C.L., & Mahon, A.R., (2014). Quantifying

Environmental DNA Signals for Aquatic Invasive Species across Multiple Detection Platforms.

Environmental Science & Technology 48 (21):12800–806. doi: 10.1021/es5034052

Nalepa, T. F., Fanslow, D.L., & Pothoven, S.A. (2010). Recent Changes in Density, Biomass, Recruitment,

Size Structure, and Nutritional State of Dreissena Populations in Southern Lake Michigan. Journal of

Great Lakes Research 36 (January):5–19. doi: 10.1016/j.jglr.2010.03.013

Peyer, S. M., McCarthy, A.J., & Lee, C.E. (2009). Zebra Mussels Anchor Byssal Threads Faster and Tighter

than Quagga Mussels in Flow. The Journal of Experimental Biology 212 (Pt 13): 2027–36. doi:

10.1242/jeb.028688

Pilliod, D.S., Goldberg, C.S., Arkle, R.S. & Waits, L.P. (2013) Estimating occupancy and abundance of

stream amphibians using environmental DNA from filtered water samples. *Canadian journal of fisheries*

and aquatic sciences. Journal canadien des sciences halieutiques et aquatiques, 70(8), 1123–1130. doi:

10.1139/cjfas-2013-0047.

Pimentel, D., Zuniga, R., & Morrison, D. (2005). Update on the environmental and economic costs

associated with alien-invasive species in the United States. Ecological economics: the journal of the

International Society for Ecological Economics, 52(3), pp.273–288. doi: 10.1016/j.ecolecon.2004.10.002

Quinn, A., Gallardo, B., & Aldridge, D.C. (2014). Quantifying the Ecological Niche Overlap between Two

Interacting Invasive Species: The Zebra Mussel (*Dreissena Polymorpha*) and the Quagga Mussel

(*Dreissena Rostriformis Bugensis*). Aquatic Conservation: Marine and Freshwater Ecosystems 24 (3):324–

37. doi: 10.1002/aqc.2414

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Ricciardi, A., Neves, R.J., & Rasmussen. J.B. (1998). Impending Extinctions of North American Freshwater

Mussels (Unionoida) Following the Zebra Mussel (*Dreissena Polymorpha*) Invasion. The Journal of Animal

Ecology 67 (4). 613–19. doi: 10.1046/j.1365-2656.1998.00220.x

Roy, H.E., Peyton, J., Aldridge, D.C., Bantock, T., Blackburn, T.M., Britton, R., & Clark P. (2014). Horizon

Scanning for Invasive Alien Species with the Potential to Threaten Biodiversity in Great Britain. Global

Change Biology 20 (12):3859–71. doi: 10.1111/gcb.12603

Schmidt, B.R., Kéry, M., Ursenbacher, S., Hyman, O.J & Collins, J.P. (2013) Site occupancy models in the

analysis of environmental DNA presence/absence surveys: a case study of an emerging amphibian

pathogen. *Methods in ecology and evolution / British Ecological Society*, 4(7), 646–653. doi:

10.1111/2041-210X.12052.

Sales, N.G. McKenzie MB, Drake J, et al (2020) Fishing for mammals: Landscape-level monitoring of

terrestrial and semi-aquatic communities using eDNA from riverine systems B. Mosher, ed. *The Journal*

of applied ecology, 57(4), 707–716. doi: 10.1111/1365-2664.13592.

Sansom, B.J., & L.M. Sassoubre. (2017). Environmental DNA (eDNA) Shedding and Decay Rates to Model

Freshwater Mussel eDNA Transport in a River. Environmental Science & Technology. 51 (24):14244-

14253. doi: 10.1021/acs.est.7b05199.

Seymour, M., Durance, I., Cosby, B.J. *et al.* (2018) Acidity promotes degradation of multi-species

environmental DNA in lotic mesocosms. *Communications Biology*, 1(1), 4. doi: 10.1038/s42003-017-

0005-3.

Simmons, M., Tucker, A., Chadderton, L.W., Jerde, C.L., & Mahon. A.R. (2015). Active and Passive

Environmental DNA Surveillance of Aquatic Invasive Species. Canadian Journal of Fisheries and Aquatic

Sciences. Journal Canadien Des Sciences Halieutiques et Aquatiques 73 (1). NRC Research Press: 76–83.

doi: 10.1139/cjfas-2015-0262

Shogren, A. J., Tank, J.L., Andruszkiewicz, E., Olds, B., Mahon, A.R., Jerde, C.L., & Bolster. D. (2017).

Controls on eDNA Movement in Streams: Transport, Retention, and Resuspension. Scientific Reports 7

(1):5065. doi: 10.1038/s41598-017-05223-1

Smart, A.S., Tingley, R., Weeks, A.R., van Rooyen, A.R., & McCarthy, M.A. (2015). Environmental DNA

Sampling Is More Sensitive than a Traditional Survey Technique for Detecting an Aquatic Invader.

Ecological Applications: A Publication of the Ecological Society of America 25 (7). Ecological Society of

America: 1944–52. doi: 10.1890/14-1751.1

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata,Z. (2012). Estimation of Fish Biomass

Using Environmental DNA. PloS One 7 (4). doi: 10.1371/journal.pone.0035868

Timar, L., & Phaneuf, D.J. (2009). Modeling the Human-Induced Spread of an Aquatic Invasive: The Case

of the Zebra Mussel. Ecological Economics: 68 (12). Elsevier: 3060–71. doi:

10.1016/j.ecolecon.2009.07.011

Thomsen, P.F., Kielgast, J., Lønsmann Iversen, L., Wiuf, C., Rasmussen, M., Gilbert, TP., …Willerslev. E.

(2012). Monitoring Endangered Freshwater Biodiversity Using Environmental DNA. Molecular Ecology

21 (11):2565–73. doi: 10.1111/j.1365-294X.2011.05418.x

Tréguier, A., Paillisson, J., Dejean, T., Valentini, A., Schlaepfer, M.A., & Roussel. J. (2014). Environmental

DNA Surveillance for Invertebrate Species: Advantages and Technical Limitations to Detect Invasive

Crayfish Procambarus Clarkii in Freshwater Ponds. The Journal of Applied Ecology 51 (4):871–79. doi:

10.1111/1365-2664.12262

Venables, W. N., & Ripley, B. D. (2002) Modern Applied Statistics with S. Fourth edition. Springer.

Wilcox, T.M., McKelvey, K.S., Young, M.K., Sepulveda, A.J., Shepard, B.B., Jane, S.F., … Schwartz. M.K.

(2016). Understanding Environmental DNA Detection Probabilities: A Case Study Using a Stream-

Dwelling Char Salvelinus Fontinalis. Biological Conservation 194:209–16. doi:

10.1016/j.biocon.2015.12.023

Yu, D.W., Ji, Y., Emerson, B.C., Wang, X., Ye, C., Yang, C., & Ding, Z. (2012). Biodiversity Soup:

Metabarcoding of Arthropods for Rapid Biodiversity Assessment and Biomonitoring. Methods in Ecology

and Evolution / British Ecological Society 3 (4): 613–23. doi: 10.1111/j.2041-210X.2012.00198.x

Zaiko, A., Martinez, J.L., Schmidt-Petersen, J., Ribicic, D., Samuiloviene, A., & Garcia-Vazquez. E. (2015).

Metabarcoding Approach for the Ballast Water Surveillance--an Advantageous Solution or an Awkward

Challenge? Marine Pollution Bulletin 92 (1-2):25–34. doi: 10.1016/j.marpolbul.2015.01.008

Figures and Table:

Figure 1: Sampling locations on the Wraysbury River. Diamond symbols represent the sampling

locations: Reservoir Outfall (RO, 0.2 km downstream of the reservoir), Wraysbury Weir (WW, 0.61 km),

Wraysbury Bridge (WB, 1.10 km), Moor Lane (ML, 1.7 km), Wraysbury Gardens (WG, 2.7 km), Hale

Street (HS, 2.72 km), and Upstream Thames confluence (UT, 2.75 km). eDNA and kick-net samples were

collected from sites with an orange outline and sites sampled for eDNA only have a blue outlined symbol

(Full grid references are provided in Supplementary Information I Table S1).

Figure 2: *Dreissena rostriformis bugensis* **qPCR mesocosm assay validation.** The graph shows the mean

and standard deviation of DNA copy number recorded for each density treatment during the mesocosm

experiment: red line – 1 specimen, green line – 5 specimens, blue line – 20 specimens and the black line

is the mean copy number for all three densities. The vertical dashed line indicates when the mussels

were removed from the mesocosm experiment.

Figure 3: *Dreissena rostriformis bugensis* DNA signal recorded at field sites. A - qPCR DNA copy number

per µl (log^{10}) at all 7 sample sites. B – Normalised relative read count of taxa assigned to quagga mussel

at the three locations sequenced.

Table 1 A - C: Comparison of methods for the detection of *D. r. bugensis* at three locations on the

Wraysbury River. "+" indicates a positive detection and "−" indicates no detection for each technical

replicate (i.e. three per sample for the eDNA methods and one for the kick sample). The numbers in

parentheses under Kick Sampling correspond to the number of individuals found in each sample.

A. Wraysbury Weir (WW)

	Kick sampling	Environmental DNA		
Sample No.	No. of individuals	PCR	qPCR	Metabarcoding
1	-	+ + +	+ + +	+ + +
2	+ (4)	+ + +	+ + +	+ + +
3	+ (4)	+ + +	+ + +	+ + +
4	-	+ + +	+ + +	+ + +
5	+ (3)	+ + +	+ + +	+ + +
6	+ (3)	+ + +	+ + +	+ + +

812 B. Wraysbury Bridge (WB)

	Kick sampling	Environmental DNA		
Sample No.	No. of individuals	PCR	qPCR	Metabarcoding
1	-	+ + +	+ + +	+ + +
2	-	+ + +	+ + +	+ + +
3	-	+ + +	+ + +	+ + +
4	-	+ + +	+ + +	+ + +
5	+ (4)	+ + +	+ + +	+ + +
6	-	+ + +	+ + +	+ + +

814   C. Wraysbury Gardens (WG)

	Kick sampling		**Environmental DNA**	
Sample No.	No. of individuals	PCR	qPCR	Metabarcoding
1	-	+ + +	+ + +	- + -
2	-	+ + +	+ + +	- - +
3	+ (1)	+ + +	+ + +	- - +
4	-	+ + +	+ + +	- - -
5	-	+ + +	+ + +	- - -
6	-	+ + +	+ + +	- - -