Title:

Environmental DNA consistently outperforms established methods for detection of quagga mussels,

Dreissena rostriformis bugensis in flowing water.

Author details:

Rosetta C Blackman[1,2,3*], Kar Keun Sean Ling [3], Lynsey R Harper[3], Peter Shum[3,4], Bernd Hänfling[3] and

Lori Lawson-Handley[3]

[1] Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic Ecology,

Überlandstrasse 133, CH-8600 Dübendorf, Switzerland.

[2] Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstr.

190, CH-8057 Zürich, Switzerland.

[3] Evolutionary and Environmental Genomics Group (EvoHull), School of Environmental Sciences,

University of Hull, Hull, UK

[4] Hopkins Marine Station, Stanford University, Pacific Grove, CA, USA

Corresponding author: Rosetta Blackman rosiecblackman@gmail.com

Abstract

1. The early detection of invasive non-native species (INNS) is important to allow for informed management actions. Established monitoring methods require the collection or observation of specimens, which is unlikely at the beginning of an invasion when densities are likely to be low. Environmental DNA (eDNA) analysis is a highly promising technique for the detection of INNS – particularly during the early stages of an invasion.

2. Here, we compared the use of traditional kick-net sampling with two eDNA approaches (targeted detection using both conventional and quantitative PCR, and passive detection via metabarcoding) for detection of quagga mussel, *Dreissena rostriformis bugensis;* a high priority INNS, along a density gradient on the River Wraysbury, UK.

3. All three molecular tools outperformed traditional sampling in terms of detection. Conventional PCR and qPCR both had 100% detection rate in all samples, and outperformed metabarcoding when the target species was at low densities. Additionally, quagga mussel DNA copy number (qPCR) and read count (metabarcoding) had a significant negative relationship with distance from source population.

4. *Synthesis and application.* All three molecular approaches were more sensitive than traditional kick-net sampling for the detection of the quagga mussel in flowing water, and both qPCR and metabarcoding enabled estimates of relative abundance. Targeted approaches were more sensitive than metabarcoding, but metabarcoding has the advantage of providing information on the wider community, and consequently impacts of INNS.

Keywords: early detection, invasive non-native species, targeted assay, metabarcoding, PCR, qPCR

Introduction

Invasive non-native species (INNS) have wide ranging effects on ecosystems, from biodiversity loss

(Doherty et al., 2016; Gallardo et al., 2016) to economic impacts including damage to infrastructure

(Pimentel et al., 2005; Connelly et al., 2007). Eradication or containment methods are most effective

during the early stage of an invasion, both in terms of success and financial cost (Hulme, 2006; Mehta et

al., 2007). Early detection is particularly challenging for species that are small, elusive or cryptic and

these species often go unnoticed until they become established (Simmons et al., 2015; Blackman et al.,

2017). Monitoring organisms such as freshwater invertebrates generally relies on specimen collection and

morphological identification, and newly invading species can be easily overlooked. Alternative methods

that offer rapid and cost-effective detection of new INNS are crucial at a time when rapid increase in

trade and transport has caused a surge in the number of successful invasions to new environments

(Hulme, 2006).

The use of molecular methods to identify DNA taken from environmental samples (i.e. environmental

DNA or "eDNA") has been a major research focus in biodiversity monitoring over the last 10 years,

including the detection of INNS (Ficetola et al., 2008, Thomsen et al., 2012). Two broad approaches can

be applied to detect a species of interest. The "targeted" approach utilises species-specific primers with

conventional PCR, quantitative PCR (qPCR) or droplet digital PCR (ddPCR) to detect a single species.

By contrast, metabarcoding can be considered a "passive approach" as it does not focus on single species

(Lawson Handley, 2015; Creer et al., 2016; Deiner et al., 2017). Metabarcoding utilises conserved

primers to amplify DNA from a group of taxa e.g. fish and other vertebrates (e.g. Kelly et al., 2014; Miya

et al., 2015; Simmons et al., 2015), zooplankton (e.g. Brown et al., 2016), insects (e.g. Yu et al., 2012.,

Elbrecht et al., 2016). Amplicons are then sequenced using High Throughput Sequencing (HTS)

platforms and bioinformatically assigned to taxa. Metabarcoding provides an opportunity to detect

multiple species simultaneously, including those, which have not been listed as priority INNS (Simmons

et al., 2015; Blackman et al., 2017). Comparisons between traditional methods for biodiversity

monitoring and targeted (Dejean et al., 2012; Tréguier et al., 2014) or passive (Smart et al., 2015;

Hänfling et al., 2016; Wilcox et al., 2016) molecular approaches have demonstrated an increased

probability of detection using eDNA methods.

The targeted approach was first used by Ficetola et al. (2008) to detect invasive *Lithobates catesbeianus*,

in lentic waterbodies and has since been applied to over 100 target INNS (Blackman et al., 2018).

Conventional PCR is a well-established and widely practiced method for detecting presence/absence of

target DNA by visualisation of PCR products on agarose gels. However, qPCR has on many occasions

been shown to be more sensitive than PCR, (e.g. Nathan et al., 2014; Simmons et al., 2015; De Ventura et

al., 2016) and therefore more suitable for detecting INNS at low density, e.g. the early stages of invasion.

Quantitative PCR has the added advantage of providing the number of DNA copies in a sample, which in

previous studies has been linked to biomass or abundance of some taxa (e.g. Amphibians, Thomsen et al.,

2012; Fish, Takahara et al., 2012; Gastropods, Goldberg et al., 2013). With the addition of a probe-based

qPCR assay there is added specificity not found with conventional or dye-based qPCR. However, qPCR

requires the construction of calibration curves from known standard concentrations and is substantially

more expensive and time-consuming than conventional PCR (Nathan et al., 2014) and maybe less

accessible to end-users than conventional PCR.

The potential of eDNA metabarcoding for INNS monitoring has already been shown as an effective tool

(Simmons et al., 2015; Brown et al., 2016; Blackman et al., 2017; Borrell et al., 2017; Holman et al., 2019).

These studies include the detection of new INNS, which had been previously overlooked by traditional

methods or targeted molecular tools. For example, Blackman et al. (2017) used a metabarcoding approach

to investigate macroinvertebrate diversity and discovered a non-native Gammaridae species, *Gammarus*

fossarum, which traditional methods had failed to identify for over 50 years in the UK. Similarly, Simmons

et al. (2015) detected the Northern Snakeshead, *Channa argus*, in the Muskingum River Watershed, Ohio,

which had not been included on INNS priority lists. Metabarcoding also has the potential to act as a

surveillance tool and has been applied to monitor high-risk pathways, such as ballast water (Ardura et al.,

2015; Zaiko et al., 2015), the live bait trade (Mahon et al., 2014) and ports (Brown et al., 2016; Borrell et

al., 2017). However, comparisons of targeted and passive approaches for INNS or other focal species are

limited to a few exceptions including bighead carp, *Hypophthalmichthys nobilis* (Simmons et al., 2015) and

great crested newts, *Triturus cristatus* (Harper et al., 2018). These studies both showed that a targeted

approach had a higher sensitivity than the passive approach. Possible causes of this include primer choice

and PCR bias, i.e. species successfully amplifying and masking others in the PCR reaction during

metabarcoding (Harper et al., 2018), but further comparisons of these methods that include a more diverse

set of target taxa are required to better inform management strategies.

In this study we focus on the quagga mussel, *Dreissena rostriformis bugensis*, a highly invasive bivalve

from the Ponto-Caspian region, which has a significant impact on all trophic levels within the

environments it invades (Karatayev et al., 2007; Roy et al., 2014). The quagga mussel scored the highest

for its potential invasions to the UK in terms of impact, risk of invading and establishing in a study from

2014 (Roy et al., 2014), and was later detected that same year (Aldridge et al., 2014). Due to its wider

tolerance to environmental conditions, it has the potential to spread rapidly, and occupy a greater range of

habitat than the closely related zebra mussel, *Dreissena polymorpha,* which has already spread throughout

much of Europe and North America (Nalepa et al., 2010; Gallardo and Aldridge 2013; Quinn et al.,

2014). Motivated by a need for cost effective tools for quagga mussel detection, we previously designed a

conventional PCR assay, which provided 100% detection rate in field trials with quagga mussel

populations (Blackman et al., *in press*). Firstly, we further developed and validated this assay for

abundance estimation using a dye-based qPCR approach for quagga mussels in mesocosm experiments

and field trials. Secondly, we directly compared the three molecular methods (conventional PCR, qPCR

and metabarcoding) to traditional kick-net sampling for detection of quagga mussels in the field. Finally,

we investigated whether qPCR and metabarcoding are suitable for estimating relative abundance of

quagga mussels; by sampling a density gradient away from the main population. We hypothesized that:

(1) the probability of detecting quagga mussels will be higher for eDNA methods than traditional

methods, (2) targeted methods will have a higher probability of detecting quagga mussel than the passive

method, (3) qPCR will be more sensitive than conventional PCR for quagga mussel detection and (4) both

read count and DNA copy number will decline with distance downstream of the source population in the

Wraysbury Reservoir.

Methods

Field sample collection, DNA capture and extraction

Wraysbury Reservoir, Berkshire, is considered the founding population of quagga mussels in the UK. The

mussels spread through the outfall and along the adjacent River Wraysbury. Samples were collected by

working in an upstream direction to ensure DNA availability and inhibitor levels were not altered by

sampling activity. Environmental DNA samples comprised of 3 x 500 ml water samples collected from

the surface at points across the river width and combined into sterile sampling bottles (i.e. 1.5 L in total

per sample). In order to compare established and molecular methods directly, we collected six of each

sample type (6 x 1.5 L water samples and 6 x kick-net samples) at three main sampling sites along the

river. These sites were Wraysbury Weir (WW, 0.61 km downstream of Wraysbury Reservoir), Wraysbury

Bridge (WB, 1.10 km), and Wraysbury Gardens (WG, 2.75 km) (See Fig. 1 and Supplementary

Information I Table S1) (n = 54). In addition, we collected four eDNA samples for conventional and

qPCR analysis only. These sites were: Reservoir Outfall (RO, 0.2 km downstream from Wraysbury

Reservoir), Moor Lane (ML, 1.7 km), Hale Street (HS, 2.72 km), and Upstream Thames confluence (UT,

2.75 km) (See Fig. 1 and Supplementary Information I Table S1 for further sampling site details) (n = 12,

n = 66 total). An upstream sample from above the reservoir outfall acted as a negative control. Sample

bottles filled with ddH_2O were taken into the field to be used as sample blanks (n = 6). (See

Supplementary Information II and III for water filtration and DNA extraction methods details)

Kick-net sampling was performed using a standard three minute kick-net sampling method followed by a

one minute hand search (Murray-Bligh, 1999). Kick-net samples were placed in labelled 42 fluid oz.

sterile Whirl-pak® bag (Cole-Palmer, Hanwell, London) and stored in a cool box before being frozen at

the laboratory. Each sample was defrosted and rinsed with ddH$_2$O separately prior to analysis. Samples

were then sorted for the presence of *D. r. bugensis* only and all specimens were counted and subsequently

stored at -20°C.

Figure 1: Sampling sites on the Wraysbury River, downstream of the main source population of *D. r. bugensis* in the Wraysbury Reservoir. Sampling sites were the Reservoir Outfall (RO, 0.2 km downstream of the reservoir), Wraysbury Weir (WW, 0.61 km), Wraysbury Bridge (WB, 1.10 km), Moor Lane (ML, 1.7 km), Wraysbury Gardens (WG, 2.7 km), Hale Street (HS, 2.72 km), and Upstream Thames confluence (UT, 2.75 km). Full grid references are provided in Supplementary Information I Table S1.

<u>Conventional PCR</u>

We used the species-specific primer set published by Blackman et al. (*in press*) for PCR amplification,

which targets a 188 bp fragment of the cytochrome oxidase I (COI) gene (DRB1_F -

GGAAACTGGTTGGTCCCGAT and DRB1_R GGCCCTGAATGCCCCATAAT). We previously

demonstrated that *D. r. bugensis* can be detected in mesocosms using this assay after four hours and at

three densities (1, 5, and 20 individuals in 15 L water, Blackman et al., *in press*). This assay is highly

sensitive for PCR (Limit of Detection [LoD] 0.0005 ng/μl) and does not cross amplify closely related

Dreissenid or native mussel species (Blackman et al., *in press*). All samples (n = 66), sampling blanks (n

= 6), filter blanks (n = 6), and PCR blanks (n = 6) were tested with the DRB1 assay, including tissue

samples from the closely related zebra mussel, acting as an additional control and to check for cross

contamination (See Supplementary Information II for full details).

<u>qPCR assay development</u>

The DRB1 assay was further developed using fluorescent dye-based qPCR. We carried out a serial

dilution of DNA from tissue samples (from 1 to 1×10^{-7} ng/μl) to estimate the Limit of Detection (LoD),

and tested the assay in the same mesocosm experiment previously used for validating the conventional

PCR assay (Blackman et al., *in press*) to ensure the assay detected *D. r. bugensis* DNA only with no cross

amplification. In brief, the mesocosm experiment monitored DNA production from mussels in three

densities (1, 5 and 20 individuals) over 21 days, with 3 replicates of each density and a control tank with

no mussels present (See Blackman et al., *in press* for further details). Fifteen litre tanks were sampled

over 21 days at: 0hrs (prior to specimens being added), 4hrs, 8hrs, 24hrs, 48 hrs, 72 hrs, 7 days, 15 days

and 21 days. On day 21, the specimens were removed, and sampling continued at: 22 days, 28 days, 35

days, 42 days (n = 130). Two hundred millilitres of water was collected at each time point and each

sample was filtered through a single filter membrane following the same protocol as the field samples.

All samples (n = 66), sampling blanks (n = 6), filter blanks (n = 6), and PCR blanks (n = 6) were tested

with DRB1 qPCR assay. Each qPCR plate had the addition of tissue DNA from a zebra mussel to act as

an additional negative control (See supplementary Information II for full qPCR protocol). Amplification

curves, Cq values and melt curves were analysed using StepOne-PlusTM software 2.0. DNA copy

number for each sample was calculated using Cq value and standard curve results (See supplementary

Information II and IV for full details).

Passive detection: eDNA metabarcoding

To match the region used for conventional and qPCR we chose the COI region for metabarcoding, which

also has the broadest taxonomic coverage for macroinvertebrates in public sequence databases. The

metabarcoding library was constructed using universal primers described by Geller et al., 2013 and Leray

et al., 2013. We confirmed amplification of the target taxa with tissue extractions prior to library

preparation. A two-step protocol was used for library preparation: briefly, this method amplifies the target

region during the first PCR and sample identification tags are added during a reduced cycle second PCR,

full details of the workflow are in Supplementary Information II. In total 54 samples were sequenced,

with the addition of 6 field blanks, 6 PCR blanks and 4 positive non-target tissue samples (n= 70).

Bioinformatics pipeline

The bioinformatics approach as described in Blackman et al., (2017) was followed with minor

modifications (See Supplementary Information II). In brief, sequences were filtered for the presence of

the target taxa only, *D. r. bugensis,* using GenBank sequences: EF080862, JX945980, EF080861,

U47651, AF495877, EU484436, KJ881409 (See Supplementary Information II for full details).

Data analysis

In order to normalise the sequencing depth of each sample from the metabarcoding data the number of

sequences assigned to *D. r. bugensis* was divided by the total number of sequences from that sample. We

used Generalized Linear Models (GLMs) to investigate 1) the influence of mussel density, biomass and

time on the DNA copy number (qPCR) since the start of the mesocosm experiment (before the mussels

were removed from the tanks only), and 2) the influence of mussel density and distance from source

population on the DNA copy number (qPCR) from the field samples and read count (metabarcoding).

Initial model tests showed all data sets were overdispersed, therefore a negative binomial GLM with a log

link function was used to resolve this issue. To check for model suitability, we checked the residual

deviance of each model fitted a chi square distribution (see Supplementary Information I for GLM

results). Finally, we assessed the relationship between DNA copy number and read count with the

distance along the river using all seven sites included in the qPCR experiment, using a Spearman's Rank

Correlation since the data were not normally distributed. All calculations and visualizations of the data

were carried out in R (R-Core-Team 2017), with GLMs performed using the MASS package (Venables et

al., 2002).

Results

qPCR assay development and validation in mesocosms

The LoD determined from the serial dilutions of tissue DNA was 1×10^{-4} ng/µl per reaction with both

conventional PCR and qPCR, which equates to approximately 5 copies per reaction (See Supplementary

Information I Fig. S1 and Fig. S3 for amplification plots). We previously demonstrated that *D. r. bugensis*

can be detected in mesocosms by conventional PCR after four hours and at all three densities using the

same mesocosm experiment as described in this study (Blackman et al., *in press*). This result was

confirmed here using qPCR (Fig. 2). Rapid DNA accumulation in the first two days of the mesocosm

experiments was followed by a depletion in DNA concentration until removal of mussels on day 21 of the

experiment. This pattern of DNA accumulation and depletion is highly consistent across all three density

treatments (Fig. 2). Time since the beginning of the experiment (until removal of the mussels) had a

significant effect on the DNA copy number in each mesocosm experiment when all density treatments

were analysed together (GLM, Z = -7.224, P < 0.0001, AIC = 912.95). There is still some overdispersion

with this model, however the overdispersion parameter indicates this is not substantial (overdispersion

statistic = 1.3, Residual Deviance 91.142 on 70 df, χ^2 P = 0.0457). There is substantial overlap between

the DNA copy number estimates for the three density treatments (Fig. 2) and neither density nor total

biomass significantly explained the number of DNA copies amplified by qPCR (Density Z = -1.019, P

= 0.308; Total biomass Z = -0.776, P = 0.438). DNA was still detectable 24 hours after removal of the

mussels, but not at the at 7 days after removal (Day 28) (Fig. 2). All blanks and mesocosm control tank

samples were clear of target DNA throughout the experiment.

Figure 2: *Dreissena rostriformis bugensis* **qPCR mesocosm assay validation.** The graph shows the

mean and standard deviation of DNA copy number recorded for each density treatment during the

mesocosm experiment: red line – 1 specimen, green line – 5 specimens, blue line – 20 specimens and the

black line is the mean copy number for all three densities. The vertical dashed line indicates when the

mussels were removed from the mesocosm experiment.

Detection of D. r. bugensis by traditional and molecular approaches in the field

Kick-net sampling

The number of *D. r. bugensis* specimens varied from 14 at Wraysbury Weir, the site closest to the

reservoir outfall (source population), to a single specimen at Wraysbury Gardens, the site furthest from

the outfall (Table 1). Detection rate was also highest at Wraysbury Weir (mussels detected in 4/6

sampling events). Mussels were detected in 1/6 samples at each of the other two sampling sites (Table 1).

Targeted approaches

Detection rate was 100% in all three field sites and all replicates tested for both targeted approaches

(conventional PCR and qPCR) (Table 1 and Fig. S2). For both conventional PCR and qPCR, all negative

and non-target tissue samples were clear of amplification for *D. r. bugensis* DNA. Fig. 3A shows the

DNA copy number (log_{10}) for *D. r. bugensis* found at each of the sampling sites. Both distance from

source and number of mussels found at the three sites, significantly influenced the DNA copy number,

but distance was the strongest predictor (Density Z = -3.367, P = 0.000761; Distance Z = -17.223, P < 2e-

16; AIC: 1392.1, Residual Deviance 57.621 on 51 df, χ^2 P = 0.244). A Spearman's Rank correlation

between the DNA copy number at each of the seven sites screened by qPCR and distance from the source

population showed a weak correlation (ρ = -0.25, P = 0.5948). However, the low copy numbers found

immediately downstream of the reservoir outfall (Fig. 3A) likely affected this. If this site is treated as an

outlier and removed from the analysis, the correlation is significant (ρ = -1, P = 0.002778).

Passive approach

Metabarcoding was less sensitive than the targeted methods at the lowest quagga mussel density

(detection rate = 86%). The total assigned read count passing quality control after chimera removal was

279 4,928,265, of which 0.6% of reads (i.e. 29,570 reads) were assigned to *D. r. bugensis* after application of

280 a 0.2% threshold to remove minor contamination. The clustering procedure identified 281 OTUs in total

281 including a range of macroinvertebrate and algae/diatom species (Supplementary information I, Fig. S4).

282 The proportion of reads assigned to *D. r. bugensis* from each sample within each site varied greatly as

283 seen in Fig. 3B (n=54). The target species was detected in all samples from both Wraysbury Weir and

284 Wraysbury Bridge sites, while detection at Wraysbury Gardens was much lower with only 3 of the 18

285 replicates having *D. r. bugensis* reads (Table 1). Furthermore, the raw number of *D. r. bugensis* reads was

286 substantially lower with average read counts of 843 and 801 at Wraysbury Bridge and Wraysbury Weir,

287 respectively and only 1 read on average at Wraysbury Gardens. The read count shows a strong

288 relationship with both the number of mussels collected at the three sites and the distance from the source

289 populations, but distance was the strongest best predictor (Distance $Z = -4.707$, $P < 0.0001$; Density $Z = -$

290 3.773, $P = 0.0001$; AIC $= 258.64$, Residual Deviance 49.696 on 51 df, χ^2 $P = 0.820$), in agreement with

291 the qPCR results.

Table 1 A - C: Comparison of methods for the detection of *D. r. bugensis* at three sites on the

Wraysbury River. "+" indicates a positive detection and "–" indicates no detection for each technical

replicate (i.e. three per sample for the eDNA methods and one for the kick sample). The numbers in

parentheses under Kick Sampling correspond to the number of individuals found in each sample.

A. Wraysbury Weir (WW)

Sample No.	Kick sampling	PCR	qPCR	Metabarcoding
1	-	+ + +	+ + +	+ + +
2	+ (4)	+ + +	+ + +	+ + +
3	+ (4)	+ + +	+ + +	+ + +
4	-	+ + +	+ + +	+ + +
5	+ (3)	+ + +	+ + +	+ + +
6	+ (3)	+ + +	+ + +	+ + +

B. Wraysbury Bridge (WB)

Sample No.	Kick sampling	PCR	qPCR	Metabarcoding
1	-	+ + +	+ + +	+ + +
2	-	+ + +	+ + +	+ + +
3	-	+ + +	+ + +	+ + +
4	-	+ + +	+ + +	+ + +
5	+ (4)	+ + +	+ + +	+ + +
6	-	+ + +	+ + +	+ + +

C. Wraysbury Gardens (WG)

Sample No.	Kick sampling	PCR	qPCR	Metabarcoding
1	-	+ + +	+ + +	- + -
2	-	+ + +	+ + +	- - +
3	+ (1)	+ + +	+ + +	- - +
4	-	+ + +	+ + +	- - -
5	-	+ + +	+ + +	- - -
6	-	+ + +	+ + +	- - -

Figure 3: *Dreissena rostriformis bugensis* DNA signal recorded at field sites. A - qPCR DNA copy

number per μl (log[10]) at all 7 sample sites. B - Metabarcoding read count per sample at each of the three

sites sequenced. The graph shows the read number for each site normalised over the total number of reads

obtained from each sample.

Discussion

In this study, we developed and validated a dye-based qPCR assay for *D. r. bugensis* then compared

targeted (conventional PCR and qPCR) and passive (metabarcoding) approaches to traditional kick-net

sampling for qualitative and semi-quantitative detection in the field. In line with our first hypothesis,

detection rate was higher for all three eDNA methods than the traditional method. Secondly, targeted

approaches were more sensitive than the passive approach. Contrary to our third hypothesis, we saw no

increased sensitivity using a dye-based qPCR over conventional PCR. Lastly, both density and distance

from the source population were significant predictors of read count (metabarcoding) and copy number

(qPCR) indicating that both approaches could provide some indication of quagga mussel abundance in the

field.

qPCR validation

Mesocosm validation provides an opportunity to test the LoD of eDNA assays and to investigate the

relationship between density, time and DNA copy number (Dejean et al., 2011; Thomsen et al., 2012;

Blackman et al., *in press*). DNA was detected with qPCR at all three densities (1, 5 and 20 individuals)

within 4 hours of being present, in agreement with previous conventional PCR results (Blackman et al., *in*

press). The LoD from serial dilutions of tissue DNA were also identical for conventional and qPCR (1 x

10^{-4} ng/µl). Surprisingly, the mesocosm experiments highlighted unexpected dynamics of DNA in closed

systems. Firstly, time since the start of the experiment was a significant predictor of DNA copy number,

but rather than increasing over time or reaching a plateau, DNA accumulated quickly in the 48 hours then

rapidly depleted, and this trend was replicated in all three density treatments. A possible explanation for

this is stress induced on the mussel caused by a change in environment, as reported in other eDNA studies

(Li et al., 2019). Dreissenid mussels, however, are known to "shut-down" in times of stress, such as

chemical dosing, and can remain with their shells closed for up for three weeks (Aldridge et al., 2006). A

further explanation for the rapid accumulation in DNA may be due to the production of protein based

byssal threads which allow the mussels to attach to the substrate (Ricciardi et al., 1998; Aldridge et al.,

2004; Timar and Phaneuf, 2009; Peyer et al., 2009). During the first 48 hours of the experiment, the

mussels are likely to secure themselves to the new substrate, all mussels were found to have done this

when they were removed at the end of the experiment. The mussels are also filter-feeders and were

actively feeding (on microorganisms found in the river water) throughout the experiment producing

sloughed cells, faeces and pseudofaeces (Sansom and Sassoubre., 2017). This feeding behaviour is likely

to be a major contributor to successful detection in the field and in a closed system effect the amount

DNA copy number detected with the qPCR assay.

The second surprising result from the mesocosm experiments was that we found no effect of density or

biomass on DNA copy number. This contradicts several previous studies that have demonstrated a

relationship between biomass or density and DNA copy number (Thomsen et al., 2012; Takahara., et al

2012; Doi et al., 2015). This could in part be due to the physiology and feeding behaviour of mussels,

since eDNA is being consumed as well as produced. The complex pattern of eDNA accumulation and

depletion together with the lack of discrimination between different density treatments presents

challenges for estimation of quagga mussel abundance. However, eDNA is unlikely to accumulate in the

same way in natural (particularly lotic) environments, and this case highlights that mesocosm experiments

may poorly reflect the true relationship between DNA copy number and mussel density in the field. It

would be worthwhile to test whether this result is confirmed with other filter feeding species. More in

keeping with previous studies (e.g. Thomsen et al., 2012), rapid degradation of eDNA was observed in

mesocosm experiments, with a complete loss of signal by one week after mussel removal.

Comparison of eDNA and kick-net sampling for quagga mussel detection

The lower sensitivity for kick-net sampling is highlighted by the failure to collect a *D. r. bugensis*

specimen in any of the first kick-net sampling attempts at each site. As demonstrated in a previous study,

species in low abundance are likely to be missed with only 62% of macroinvertebrate families detected in

a three minute kick-net sample (Furse et al., 1981), and therefore highlights the need for multiple

replicates for detection of low abundance species if using kick-net sampling. It should be noted that

surveillance methods for *D. r. bugensis* should be focussed on the preferred habitat of mussels i.e. hard

substrates such as man-made structures (Aldridge et al., 2014), which is more likely to detect quagga

mussels than the randomised method used here. However, randomised kick-net sampling along a range of

substrate types is the most widely used sampling method for freshwater macroinvertebrate monitoring,

and therefore the most likely to detect new INNS.

Of the three molecular approaches compared in this study, we show that both targeted approaches were

more sensitive for detecting *D. r. bugensis* than metabarcoding. Few studies have compared targeted

approaches with metabarcoding; however, our findings are similar to those of Simmons et al., (2015) and

Harper et al., (2018), who showed that metabarcoding was less sensitive than targeted approaches. In this

study, there was no increase in the detection of *D. r. bugensis* without a contamination threshold applied

(0.2%). However, it would be pertinent to examine the raw output of metabarcoding runs before

thresholds are applied in order to identify any potential INNS with low read counts, and to verify

detection of priority INNS with a targeted assay. A significant advantage of metabarcoding is the added

information it provides on non-target species. A further 281 OTUs were identified in this dataset, and

although no additional INNS were detected, previous studies have demonstrated the power of this

approach for detecting new and unexpected INNS (Blackman et al., 2017). Metabarcoding also provides

the opportunity to monitor changes in community composition as the result of INNS or other ecological

stressors (Simmons et al., 2015; Hänfling et al., 2016), which is extremely promising for better

understanding INNS impacts for more effective prioritisation and management. It could also be possible

to improve the power of metabarcoding by increasing spatial sampling. Future studies should investigate

how many samples are needed in order to maximise the probability of detecting rare species with

metabarcoding.

Conventional PCR and dye-based qPCR had identical performance for *D. r. bugensis* detection in our

study. To increase specificity eDNA qPCR studies often use the addition of a probe to prevent non-target

amplification. However, given the already high specificity of the DRB1 primer set described here, we

believe the addition of a probe is not necessary for INNS monitoring needs. The chosen approach will

depend on the specific requirements of the monitoring programme. If quantitative information is not

required, conventional PCR is easily accessible to end-users. It offers a quick and cost-effective approach

for the detection of new INNS, requiring limited lab equipment and up to half the time for sample

processing than qPCR (Davison et al., 2016). If quantitative information is required, for example when

estimating population sizes and monitoring areas where eradication attempts have been carried out, qPCR

will be more appropriate.

Estimating quagga mussel abundance with eDNA

Results from field trials for both dye-based qPCR and metabarcoding demonstrate a significant decrease

in DNA copy number and read count, respectively, with increasing distance from the main source

population (Wraysbury reservoir). Kick-net sample data also indicates that mussel density decreases

along the course of the river, with only one mussel found at Wraysbury Gardens, 2.7 km downstream of

the reservoir. Quantitative data from qPCR and metabarcoding is particularly informative in this case as

the concentration gradient suggests that the source population may be inferred in cases where it is

unknown. The only site that did not fit this trend was the first site, 20 m downstream of the outfall from

Wraysbury reservoir, which had a very low DNA read and number.

There are two main processes which are known to influence the transport of DNA: firstly, the effect of

stream discharge on transport (Jane et al., 2015), and secondly, the fundamental structure of the river

(Wilcox et al., 2016; Shogren et al., 2017). During this study, sampling took place at a period of increased

flow (following heavy rain). In contrast to the findings of Jane et al., (2015), the DNA signal was not

uniform along the stretch as expected by higher flows and increased mixing. It is therefore likely other

factors were influencing the DNA signal across the river length. Shogren et al., (2017) in particular,

summarised the elements most influential on the movement of DNA as "Transport, Retention and

Resuspension". Our data indicates distance has a significant correlation with DNA copy number along the

2.75 km stretch of river. It is highly likely that our data are reflect both a decrease in DNA being shed by

populations of decreasing size as well as the effect of DNA being retained by the substrate in slower flow

areas (slacks, glides and pools) and re-suspended in higher velocity flows (riffles and runs). To what

extent these factors are influencing the DNA signal is currently unclear, and further studies during

different flow rates are needed in order to explore these variables further.

Practical consideration for future INNS biomonitoring and conclusions

1.   Successful early detection of INNS monitoring requires tools fit for purpose: current methods are

unsuitable for the detection of species in low densities and often rely on expertise, which are unable

to predict or identify new or unexpected INNS. Here we have demonstrated that eDNA is a viable

alternative for both targeted and community surveys.

2.   Here we have also shown that dependant on the information required (quantitative,

presence/absence or community level surveillance) we are able to tailor molecular tools to answer

different questions about INNS dynamics.

3.   The optimal approach we believe is to combine both molecular approaches: carry out routine

surveillance monitoring for those new or unexpected INNS using a metabarcoding approach

followed by verification with targeted assays. Targeted assays should be applied for those INNS

expected to arrive or those that have established already and are at risk of spreading.

Acknowledgment

This project was funded by the Freshwater Biological Association Gilson Le Cren Award 2017 and the

UK Environment Agency. We would like to thank staff from the Environment Agency - Drs Alice Hiley

References

Aldridge, D.C., P. Elliott, P., & Moggridge. G.D. (2004). The Recent and Rapid Spread of the Zebra

Mussel (*Dreissena Polymorpha*) in Great Britain. Biological Conservation, 119 (2):253–61. doi:

10.1016/j.biocon.2003.11.008

Aldridge, D.C., Elliott P., & Moggridge G.D. (2006). Microencapsulated BioBullets for the Control of

Biofouling Zebra Mussels. Environmental Science & Technology, 40 (3): 975–79. doi:

10.1021/es050614%2B

Aldridge, D. C., Ho, S., & Froufe. E. (2014). The Ponto-Caspian Quagga Mussel, *Dreissena Rostriformis*

Bugensis (Andrusov, 1897), Invades Great Britain. Aquatic Invasions / European Research Network on

Aquatic Invasive Species. doi: 10.3391/ai.2014.9.4.11

Ardura, A., Zaiko, A., Martinez, J.L., Samuiloviene, A., Borrell, Y., & Garcia-Vazquez. E. 2015

Environmental DNA Evidence of Transfer of North Sea Molluscs across Tropical Waters through Ballast

Water. The Journal of Molluscan Studies 81 (4). Oxford University Press: 495–501. doi:

10.1093/mollus/eyv022

Blackman, R.C., Constable, D., Hahn, C., Sheard, A., Durkota, J., Hänfling, B., & Lawson Handley. L.

(2017). Detection of a New Non-Native Freshwater Species by DNA Metabarcoding of Environmental

Samples — First Record of *Gammarus Fossarum* in the UK. Aquatic Invasions 12 (2):177–89. doi:

10.3391/ai.2017.12.2.06

Blackman, R.C., Hänfling, B., & Lawson Handley. L. (2018). The use of environmental DNA as an early

warning tool in the detection of new freshwater invasive non-native species. CAB Reviews Vol 12, 010,

pp1-15. doi: 10.1079/PAVSNNR201813010

Blackman, R.C., Benucci, M., Donnelly, R.C., Hänfling, B., Harper, L.R., Sellers, G.S. & Lawson

Handley, L. Simple, sensitive and species-specific assays for detecting quagga and zebra mussels

(*Dreissena rostriformis bugensis* and *D. polymorpha*) using environmental DNA. Management of

Biological Invasions. *In press.*

Borrell, Y.J., Miralles, L., Do Huu, H., Mohammed-Geba, K., & Garcia-Vazquez. E. (2017). DNA in a

bottle—Rapid Metabarcoding Survey for Early Alerts of Invasive Species in Ports. PloS One 12 (9).

Public Library of Science: e0183347. doi: 10.1371/journal.pone.0183347

Brown, E.A., Chain, F.C.C., Zhan, A., MacIsaac, H.J., & Cristescu. M.C. (2016). Early Detection of

Aquatic Invaders Using Metabarcoding Reveals a High Number of Non-Indigenous Species in Canadian

Ports. Diversity & Distributions. doi: 10.1111/ddi.12465

Connelly, N.A., O'Neill, C.R., Knuth, B.A., & Brown, T.L. (2007). Economic impacts of zebra

mussels on drinking water treatment and electric power generation facilities. Environmental

management, 40(1) pp.105–112. doi: 10.1007/s00267-006-0296-5

Creer, S, Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, W.K., … & Bik. H.M. (2016). The

Ecologist's Field Guide to Sequence-Based Identification of Biodiversity. Methods in Ecology and

Evolution / British Ecological Society 7 (9):1008–18. doi: 10.1111/2041-210X.12574

Davison, P. I., Créach, V., Liang, W-J., Andreou, D., Britton, J.R., & Copp, G.H.. (2016). Laboratory and

Field Validation of a Simple Method for Detecting Four Species of Non-Native Freshwater Fish Using

eDNA. Journal of Fish Biology 89 (3):1782–93. doi: doi.org/10.1111/jfb.13086

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., & Creer, S.

(2017). Environmental DNA Metabarcoding: Transforming How We Survey Animal and Plant

Communities. Molecular Ecology 26 (21):5872–95. doi: 10.1111/mec.14350

Dejean, T., Valentini, A., Duparc, A., Pellier-Cuit, S., Pompanon, F., Taberlet, P., & Miaud. C. (2011).

Persistence of Environmental DNA in Freshwater Ecosystems. PloS One 6 (8) doi:

10.1371/journal.pone.0023398

Dejean, T., Valentini, A., Miquel, C., Taberlet, P., Bellemain, E., & Miaud. C. (2012). Improved

Detection of an Alien Invasive Species through Environmental DNA Barcoding: The Example of the

American Bullfrog Lithobates Catesbeianus. The Journal of Applied Ecology 49 (4) 953–59. doi:

10.1111/j.1365-2664.2012.02171.x

De Ventura, L., Kopp, K., Seppälä, K., & Jokela. J. (2017). Tracing the Quagga Mussel Invasion along

the Rhine River System Using eDNA Markers: Early Detection and Surveillance of Invasive Zebra and

Quagga Mussels. Vol 8 (1) 101–112. doi:10.3391/mbi.2017.8.1.10

Doherty, T.S., Glen, A.S., Nimmo, D. G., Ritchie, E. G., & Dickman, C. R. (2016). Invasive predators

and global biodiversity loss. Proceedings of the National Academy of Sciences of the United States of

America, 113(40), pp.11261–11265. doi: 10.1073/pnas.1602480113

Doi, H., Uchii, K., & Takahara T, (2015a). Use of droplet digital PCR for estimation of fish abundance

and biomass in environmental DNA surveys. PloS one 10(3). doi: 10.1371/journal.pone.0122763

Elbrecht, V., & Leese. F. (2017). Validation and Development of Freshwater Invertebrate Metabarcoding

COI Primers for Environmental Impact Assessment. Frontiers in Freshwater Science. 5(11), 1– 11. doi:

10.3389/fenvs.2017.00011

Ficetola, G.F., Miaud, C., Pompanon, F., & Taberlet. P. (2008). Species Detection Using Environmental

DNA from Water Samples. Biology Letters 4 (4):423–25. doi: 10.1098/rsbl.2008.0118

Furse, M. T., Wright, J.F., Armitage, P.D., & Moss, D. (1981). An Appraisal of Pond-Net Samples for

Biological Monitoring of Lotic Macro-Invertebrates. Water Research 15 (6):679–89. doi: 10.1016/0043-

1354(81)90160-3

Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial

cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys.

Molecular ecology resources, 13, 851–861. doi: 10.1111/1755-0998.12138

Gallardo, B., & Aldridge, D.C. (2013) The 'dirty Dozen': Socio-Economic Factors Amplify the Invasion

Potential of 12 High-Risk Aquatic Invasive Species in Great Britain and Ireland. The Journal of Applied

Ecology 50 (3):757–66. doi: 10.1111/1365-2664.12079

Gallardo, B., Clavero, M., Sánchez, M. I., & Vilà, M. (2016). Global ecological impacts of invasive

species in aquatic ecosystems. Global change biology, 22(1), pp.151–163. doi: 10.1111/gcb.13004

Goldberg, C S., Sepulveda, A., Ray, A., Baumgardt, J., & Waits. L.P. (2013). Environmental DNA as a

New Method for Early Detection of New Zealand Mudsnails (Potamopyrgus Antipodarum). Freshwater

Science 32 (3) 792–800. doi: 10.1899/13-046.1

Hänfling, B., Lawson Handley, L., Read, D.S., Hahn, C., Li, J., Nichols, P., … Winfield. I. (2016).

Environmental DNA Metabarcoding of Lake Fish Communities Reflects Long-Term Data from

Established Survey Methods. Molecular Ecology 25 (13):3101–19. doi: 10.1111/mec.13660

Harper, LR., Lawson Handley, L., Hahn, C., Boonham. N., Rees, HC., Gough, KC., … Hänfling, B.

(2017). Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of

great crested newt (Triturus cristatus). Ecology and Evolution 8 (12) 6330-6341. doi: 10.1002/ece3.4013

Holman, L.E., de Bruyn, M., Creer, S., Carvalho., Robidart., J., & Rius, M. (2019). Detection of

introduced and resident marine species using environmental DNA metabarcoding of sediment and water.

Scientific reports, 9(1), p.11559. doi: 10.1038/s41598-019-47899-7

Hulme, P.E., (2006). Beyond Control: Wider Implications for the Management of Biological Invasions.

The Journal of Applied Ecology 43 (5): 835–47. Doi: 10.1111/j.1365-2664.2006.01227.x

Jane, S. F., Wilcox, T.M., McKelvey, K.S., Young, M.K., Schwartz, M.K., Lowe, … Whiteley. A.R.

(2015). Distance, Flow and PCR Inhibition: eDNA Dynamics in Two Headwater Streams. Molecular

Ecology Resources 15 (1):216–27. doi: 10.1111/1755-0998.12285

Karatayev, A.Y., Padilla, D.K., Minchin, D., Boltovskoy, D., & Burlakova. L.E. (2007). Changes in

Global Economies and Trade: The Potential Spread of Exotic Freshwater Bivalves. Biological Invasions 9

(2). 161–80. doi: 10.1007/s10530-006-9013-9

Kelly, R. P., Port, J.A., Yamahara, K.M., & Crowder, L.B. (2014). Using Environmental DNA to Census

Marine Fishes in a Large Mesocosm. PloS One 9 (1). doi: 10.1371/journal.pone.0086175

Lawson Handley, L. (2015). How Will the 'molecular Revolution' Contribute to Biological Recording?

Biological Journal of the Linnean Society. Linnean Society of London 115 (3):750–66. doi:

10.1111/bij.12516

Leray, M., Yang, J.Y., Meyer, C.P., Mills, S.C., Agudelo, N., Ranwez, V., … Machida, R.J. (2013). A

New Versatile Primer Set Targeting a Short Fragment of the Mitochondrial COI Region for

Metabarcoding Metazoan Diversity: Application for Characterizing Coral Reef Fish Gut Contents.

Frontiers in Zoology 10:34. doi: 10.1186/1742-9994-10-34

Li, J., Lawson Handley, L., Harper, L.R., Brys, R., Watson, H.V., Di Muri, C., … Hänfling, B. (2019).

"Limited Dispersion and Quick Degradation of Environmental DNA in Fish Ponds Inferred by

Metabarcoding." *Environmental DNA* 1 (3): 238–50.

Mahon, A.R., Nathan, L.H., & Jerde, C.L. (2014). Meta-Genomic Surveillance of Invasive Species in the

Bait Trade. Conservation Genetics Resources 6. doi: 10.1007/s12686-014-0213-9

Mehta, S.V., Haight, R. G., Homans, F. R, Polasky, S & Venette, R.C. (2007). Optimal detection and

control strategies for invasive species management. Ecological economics: the journal of the International

Society for Ecological Economics, 61(2), pp.237–245. doi: 10.1016/j.ecolecon.2006.10.024

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., … Iwasaki, W. (2015). MiFish, a Set

of Universal PCR Primers for Metabarcoding Environmental DNA from Fishes: Detection of More than

230 Subtropical Marine Species. Royal Society Open Science 2 (7). doi: 10.1098/rsos.150088

Murray-Bligh J.A.D. (1999). Procedure for Collecting and Analysing Macroinvertebrate Samples,

Environment Agency, Bristol, UK.

Nathan, L.M., Simmons, M., Wegleitner, B.J., Jerde, C.L., & Mahon, A.R., (2014). Quantifying

Environmental DNA Signals for Aquatic Invasive Species across Multiple Detection Platforms.

Environmental Science & Technology 48 (21):12800–806. doi: 10.1021/es5034052

Nalepa, T. F., Fanslow, D.L., & Pothoven, S.A. (2010). Recent Changes in Density, Biomass,

Recruitment, Size Structure, and Nutritional State of Dreissena Populations in Southern Lake Michigan.

Journal of Great Lakes Research 36 (January):5–19. doi: 10.1016/j.jglr.2010.03.013

Peyer, S. M., McCarthy, A.J., & Lee, C.E. (2009). Zebra Mussels Anchor Byssal Threads Faster and

Tighter than Quagga Mussels in Flow. The Journal of Experimental Biology 212 (Pt 13): 2027–36. doi:

10.1242/jeb.028688

Pimentel, D., Zuniga, R., & Morrison, D. (2005). Update on the environmental and economic costs

associated with alien-invasive species in the United States. Ecological economics: the journal of the

International Society for Ecological Economics, 52(3), pp.273–288. doi: 10.1016/j.ecolecon.2004.10.002

Quinn, A., Gallardo, B., & Aldridge, D.C. (2014). Quantifying the Ecological Niche Overlap between

Two Interacting Invasive Species: The Zebra Mussel (*Dreissena Polymorpha*) and the Quagga Mussel

(*Dreissena Rostriformis Bugensis*). Aquatic Conservation: Marine and Freshwater Ecosystems 24

(3):324–37. doi: 10.1002/aqc.2414

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Ricciardi, A., Neves, R.J., & Rasmussen. J.B. (1998). Impending Extinctions of North American

Freshwater Mussels (Unionoida) Following the Zebra Mussel (*Dreissena Polymorpha*) Invasion. The

Journal of Animal Ecology 67 (4). 613–19. doi: 10.1046/j.1365-2656.1998.00220.x

Roy, H.E., Peyton, J., Aldridge, D.C., Bantock, T., Blackburn, T.M., Britton, R., & Clark P. (2014).

Horizon Scanning for Invasive Alien Species with the Potential to Threaten Biodiversity in Great Britain.

Global Change Biology 20 (12):3859–71. doi: 10.1111/gcb.12603

Sansom, B.J., & L.M. Sassoubre. (2017). Environmental DNA (eDNA) Shedding and Decay Rates to

Model Freshwater Mussel eDNA Transport in a River. Environmental Science & Technology. 51

(24):14244-14253. doi: 10.1021/acs.est.7b05199.

Simmons, M., Tucker, A., Chadderton, L.W., Jerde, C.L., & Mahon. A.R. (2015). Active and Passive

Environmental DNA Surveillance of Aquatic Invasive Species. Canadian Journal of Fisheries and

Aquatic Sciences. Journal Canadien Des Sciences Halieutiques et Aquatiques 73 (1). NRC Research

Press: 76–83. doi: 10.1139/cjfas-2015-0262

Shogren, A. J., Tank, J.L., Andruszkiewicz, E., Olds, B., Mahon, A.R., Jerde, C.L., & Bolster. D. (2017).

Controls on eDNA Movement in Streams: Transport, Retention, and Resuspension. Scientific Reports 7

(1):5065. doi: 10.1038/s41598-017-05223-1

Smart, A.S., Tingley, R., Weeks, A.R., van Rooyen, A.R., & McCarthy, M.A. (2015). Environmental

DNA Sampling Is More Sensitive than a Traditional Survey Technique for Detecting an Aquatic Invader.

Ecological Applications: A Publication of the Ecological Society of America 25 (7). Ecological Society

of America: 1944–52. doi: 10.1890/14-1751.1

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata,Z. (2012). Estimation of Fish Biomass

Using Environmental DNA. PloS One 7 (4). doi: 10.1371/journal.pone.0035868

Timar, L., & Phaneuf, D.J. (2009). Modeling the Human-Induced Spread of an Aquatic Invasive: The

Case of the Zebra Mussel. Ecological Economics: 68 (12). Elsevier: 3060–71. doi:

10.1016/j.ecolecon.2009.07.011

Thomsen, P.F., Kielgast, J., Lønsmann Iversen, L., Wiuf, C., Rasmussen, M., Gilbert, TP., …Willerslev.

E. (2012). Monitoring Endangered Freshwater Biodiversity Using Environmental DNA. Molecular

Ecology 21 (11):2565–73. doi: 10.1111/j.1365-294X.2011.05418.x

Tréguier, A., Paillisson, J., Dejean, T., Valentini, A., Schlaepfer, M.A., & Roussel. J. (2014).

Environmental DNA Surveillance for Invertebrate Species: Advantages and Technical Limitations to

Detect Invasive Crayfish Procambarus Clarkii in Freshwater Ponds. The Journal of Applied Ecology 51

(4):871–79. doi: 10.1111/1365-2664.12262

Venables, W. N., & Ripley, B. D. (2002) Modern Applied Statistics with S. Fourth edition. Springer.

Wilcox, T.M., McKelvey, K.S., Young, M.K., Sepulveda, A.J., Shepard, B.B., Jane, S.F., … Schwartz.

M.K. (2016). Understanding Environmental DNA Detection Probabilities: A Case Study Using a Stream-

Dwelling Char Salvelinus Fontinalis. Biological Conservation 194:209–16. doi:

10.1016/j.biocon.2015.12.023

Yu, D.W., Ji, Y., Emerson, B.C., Wang, X., Ye, C., Yang, C., & Ding, Z. (2012). Biodiversity Soup:

Metabarcoding of Arthropods for Rapid Biodiversity Assessment and Biomonitoring. Methods in

Ecology and Evolution / British Ecological Society 3 (4): 613–23. doi: 10.1111/j.2041-

210X.2012.00198.x

Zaiko, A., Martinez, J.L., Schmidt-Petersen, J., Ribicic, D., Samuiloviene, A., & Garcia-Vazquez. E.

(2015). Metabarcoding Approach for the Ballast Water Surveillance--an Advantageous Solution or an

Awkward Challenge? Marine Pollution Bulletin 92 (1-2):25–34. doi: 10.1016/j.marpolbul.2015.01.008