1 **Structure-function relation of the developing calyx of Held synapse *in vivo***

2
**Abbreviated title:** Strength-size relation of the calyx of Held

**Martijn C. Sierksma[1,3], Johan A. Slotman[2], Adriaan B. Houtsmuller[2] and J. Gerard G. Borst[1]**

[1]Department of Neuroscience or [2]Department of Pathology – Optical Imaging Centre, Erasmus MC,
University Medical Center Rotterdam, 3000 CA, Rotterdam, The Netherlands.

[3]Sorbonne Université, Inserm, CNRS, Institut de la Vision, 17 Rue Moreau, F-75012 Paris, France.

Correspondence to g.borst@erasmusmc.nl

17
18

**Conflict of interest**: The authors declare no competing financial interests.

**Contributions**
JGGB and MCS designed experiments. MCS performed *in vivo* electrophysiology and imaging
experiments, analyzed the electrophysiology and the images. JAS performed imaging experiments
and imaging analysis. JAS and ABH designed imaging analysis. MCS drafted the manuscript. JAS, ABH
and JGGB commented on manuscript.

**Acknowledgements**
This work was financially supported by the Earth and Life Sciences-Netherlands Organisation of
Scientific Research (NWO, #823.02.006, 'Development of a Giant Synapse'). We are very grateful for
the help of Elize Haasdijk and Celina Glimmerveen, who performed the immunolabeling. We thank
Marcel van der Heijden, Peter Bremen and Aaron Wong for advice on the analyses.

**Abstract**

In adult rodents, a principal neuron in the medial nucleus of the trapezoid (MNTB) is generally

contacted by a single, giant axosomatic terminal called the calyx of Held, but how this one-on-one

relation is established is still unknown. Anatomical evidence suggests that during development

principal neurons are innervated by multiple calyces, which may indicate calyceal competition, but *in*

*vivo* electrophysiological recordings from principal neurons have indicated that only a single strong

synaptic connection forms per cell. To test whether a mismatch between synaptic strength and

terminal size exists, we compared strength of synaptic inputs during early postnatal development

with the morphology of the synaptic terminals. *In vivo* whole-cell recordings of the MNTB neurons

from newborn rats of either sex were made while stimulating their afferent axons, allowing us to

identify multiple inputs. The strength of the strongest input increased to calyceal levels in a few days

across cells, while the strength of the second strongest input was stable. Cells were subsequently

immuno-labeled for vesicular glutamate transporters (VGluT) to reveal axosomatic terminals.

Synaptic strength of the strongest input was correlated with the contact area of the largest VGluT

cluster at the soma. No clear mismatch was observed between structure and strength. Together, our

data agree with a developmental scheme in which one input strengthens and becomes the calyx of

Held, but not with multi-calyceal competition.

**Significance**

Synapses are usually very small, but they can be as large as a cell body. What is the relation between

the size of a synapse and its strength? We measured in neonatal rats the strength of a specific

synaptic connection that grows in a few days from a small to a giant synapse, called the calyx of Held

synapse. Synaptic strength was correlated with the contact surface of the largest glutamatergic

terminal on the recorded neuron. Our results also suggest that during development only one of the

connections per cell increases in strength and expands over the soma, forming the calyx of Held

synapse.

## Introduction

Synapses come in a large variety of sizes and shapes. Most terminals have a diameter of about one µm, forming a single synaptic contact on a dendrite. In contrast, some synaptic terminals are >10 µm, and contain >100 release sites (Atwood & Karunanithi, 2002; Rollenhagen & Lübke, 2006). These large synapses have facilitated the study of the biophysical properties of synapses. A prime example is the calyx of Held synapse in the auditory brainstem. The calyx spans ~20 µm, large enough to be accessible for patch-clamp electrophysiology (Forsythe, 1994; Borst *et al.*, 1995). Owing to the presence of hundreds of active zones (Sätzler *et al.*, 2002; Taschenberger *et al.*, 2002; Dondzillo *et al.*, 2010), the calyx of Held can rapidly trigger action potentials in its target, a glycinergic neuron in the medial nucleus of the trapezoid body (MNTB), thus functioning as a fast, high-fidelity, inverting relay in the auditory brainstem (Borst & Soria van Hoeve, 2012).

The calyx of Held synapse grows from an axon of a globular bushy cell (GBC) that will initially form small boutons. In rodents these contacts appear before birth (Kandler & Friauf, 1993; Hoffpauir *et al.*, 2006; Rodríguez-Contreras *et al.*, 2008; Hoffpauir *et al.*, 2010; Holcomb *et al.*, 2013). Although the innervation initially is highly divergent, eventually a GBC will give rise to only one or a few calyces, and most adult principal neurons are innervated by a single calyx of Held (Held, 1893; Morest, 1968; Kuwabara *et al.*, 1991; Kandler & Friauf, 1993; Rodríguez-Contreras *et al.*, 2006). The developmental mechanisms that ensure the transition to the one-on-one innervation are largely unclear (Yu & Goodrich, 2014). This transition happens between the second and the fifth postnatal day (P2-5). At P5 the strength of one input overshadows the other ones (Chuhma & Ohmori, 1998; Hoffpauir *et al.*, 2006; Hoffpauir *et al.*, 2010; Sierksma *et al.*, 2016), and one of the terminals has expanded over the soma of the neuron (Hoffpauir *et al.*, 2006; Holcomb *et al.*, 2013). From these studies it was suggested that half of the MNTB neurons are contacted by multiple calyces at this age range (Hoffpauir *et al.*, 2010; Holcomb *et al.*, 2013), in agreement with observations from mouse slice recordings (Bergsman *et al.*, 2004; Hoffpauir *et al.*, 2010; Xiao *et al.*, 2013). However, *in vivo*

physiological evidence for calyceal competition was observed in only very few cases for the rat

(Sierksma *et al.*, 2016).

These findings may indicate a discrepancy between terminal size and synaptic strength. To test this

possibility, one needs to correlate synaptic structures with synaptic strength. Serial sectioning of

tissue combined with EM offers the possibility of fully reconstructing every synapse in a restricted

volume (Denk & Horstmann, 2004; Hoffpauir *et al.*, 2006; Holcomb *et al.*, 2013), and allows the

reconstruction of entire calyces (Sätzler *et al.*, 2002; Holcomb *et al.*, 2013; Qiu *et al.*, 2015).

However, it does not offer a direct estimate of synaptic strength. Whereas slice electrophysiology

has the advantage of allowing simultaneous recordings of pre- and postsynaptic structures (Borst *et*

*al.*, 1995; Rodríguez-Contreras *et al.*, 2008), there is the uncertainty associated with possible cutting

of inputs during slice preparation. Immunolabeling of vesicular glutamate transporters (VGluT)

localizes synapses in the MNTB (Rodríguez-Contreras *et al.*, 2006; Rodríguez-Contreras *et al.*, 2008;

Soria Van Hoeve & Borst, 2010). As Piccolo, an active zone protein (Südhof, 2012; Gundelfinger *et al.*,

2016), is present in young calyces (Dondzillo *et al.*, 2010), the combination of VGluT and Piccolo may

indicate synaptic connectivity in the MNTB. How VGluT relates to synaptic strength is still unclear.

We therefore combined VGluT immunolabeling with *in vivo* whole-cell recordings during which we

recorded the inputs that are regularly active (Lorteije *et al.*, 2009; Sierksma *et al.*, 2016). We applied

electrical stimulation of the afferent axons to further identify the developing inputs. The recorded

cells were subsequently immunolabeled for VGluT and Piccolo, allowing a direct comparison

between synapse strength and structure for the developing calyx of Held synapse.

## Materials & methods

### *Animals*

All procedures conformed to the European legislation and were approved by the local animal ethics committee (EDC, project no. 115-14-11). Wistar dams (WU) were purchased from Charles River and housed within the Erasmus animal facility (EDC). The day of birth or finding the litter was taken as postnatal day (P)0. The dams had *ad libitum* access to food and water, and additional bedding and shelter material was provided for nest building.

### *Surgery and in vivo electrophysiology*

We used a ventral approach to gain access to the auditory brainstem, as described previously (Rodríguez-Contreras *et al.*, 2008; Sierksma *et al.*, 2016). First, we searched for bursting activity with a glass electrode to find the presumed location of the MNTB. When bursting activity was found, the search pipette was retracted and a bipolar stimulation electrode (MicroProbes for Life Science, PI2ST30.1H10) was placed contralateral from the recorded MNTB, straddling the trapezoid body, as described by Crins *et al.* (2011). Successful placement was assessed by measuring the stimulation current threshold and size of the field potential.

Whole-cell recordings with biocytin (2 mg/mL) in the pipette solution were made from principal cells; to be confident that the recording was obtained from the same neuron as the one that was recovered with immunolabeling, we made whole-cell recordings from only a single principal neuron per animal. Whenever a glia cell or non-principal cell was encountered, the pipette was retracted and a new pipette was used. Membrane potentials were compensated prior to the recordings for a liquid junction potential of -11 mV. Drift in membrane potential was <5 mV and remained uncorrected. Recordings were made with a MultiClamp 700B in current-clamp mode, with bridge balance set in the range of 20-70 MΩ and pipette capacitance compensation ~4-6 pF. Signals were low-pass filtered with a 4-pole Bessel filter at 10 kHz and digitized by a DigiData 1440A (Molecular Devices Co.) at 25 kHz. Acquisition was done with Clampex 10.2 running on a Windows XP computer.

Stimulation intensities were manually adjusted during the experiments. First, we quickly assessed at

which stimulation levels new excitatory postsynaptic potentials (EPSPs) were recruited, and at which

polarity most inputs could be discerned. Then, stimulation intensity was lowered to the activation

threshold of the first input, and step-wise increased over time. In some experiments the stimulation

intensity was also stepwise reduced, and then the data were pooled. For every stimulation strength

>30 sweeps were collected at 2 Hz. Stimulation current was typically below 0.4 mA to prevent

damage to the axons; at high stimulation currents (0.3 mA or higher) antidromic APs with latencies

<1 ms were often elicited in the principal neuron. This could still happen even though we used

bipolar stimulation electrodes, placed on the contralateral side from the recorded neuron.

Preferably, the stimulation electrode would be placed at the midline, where all calcigenic axons

converge ventrally. However, to avoid direct postsynaptic activation we placed the stimulation

electrode more laterally. This location may have led to preferential activation of the axon bundles

that run ventrally from the other MNTB (Kandler & Friauf, 1993).

After a recording the animal was deeply anesthetized and perfused with 6-10 mL of cold saline (0.9%

NaCl, w/v), followed by 8-10 mL of 4% of paraformaldehyde (PFA, w/v) dissolved in 0.12 M

phosphate-buffer (PB, pH 7.2-7.4).

***Antibodies***

The following primary antibodies were used: rabbit polyclonal anti-Piccolo (Synaptic Systems

142003, RRID: AB_2160182; 1:1000), guinea-pig polyclonal anti-vesicular glutamate transporter 1

(Millipore AB5905, RRID: AB_2301751; 1:3000) and 2 (Millipore AB2251, RRID: AB_1587626;

1:3000). Secondary antibodies Alexa Fluor 488 against rabbit and Alexa Fluor 645 against guinea pig

were obtained from Jackson (1:200 or 1:400). Streptavidin-Alexa Fluor 594 conjugate (1:200) was

obtained from Thermo Fisher Scientific. Throughout, VGluT labeling refers to both VGluT1 and

VGluT2.

 *Immunolabeling procedure*

Immunolabeling procedure was based on the free-floating method. After perfusion the brain was

carefully removed from the skull, and post-fixed overnight at 4 °C. Then, it was left overnight in 10%

(w/v) sucrose in 0.1 M PB, embedded in 10% (w/v) gelatin and 10% (w/v) sucrose, and again fixed

overnight at 4 °C in 30% (w/v) sucrose and 10% (w/v) formaldehyde. The brain was cryoprotected

with 30% (w/v) sucrose solution in 0.1 M PB for >24 hr at 4 °C. Coronal slices (40 μm) were made on

a freezing microtome, and were collected in 0.1 M PB. The sections were heated for 3 hr to 80 °C in

10 mM sodium citrate (pH 6 at RT), washed four times for 10 min with 0.9% (w/v) NaCl in 0.05 M PB

(PBS). Then, the sections were pre-absorbed for 1 hr at RT with 10% (v/v) normal horse serum and

10% (v/v) Triton X-100 in PBS followed by 36-48 hr incubation under gentle agitation at 4 °C with

primary antibody solution containing 2% (v/v) normal horse serum, 0.4% (v/v) Triton X-100 and the

primary antibodies. The slices were washed four times for 10 min in PBS at RT, and incubated for 2

hr under gentle agitation at RT in the secondary antibody solution containing 2% (v/v) normal horse

serum, 0.4% (v/v) Triton X-100, and the secondary antibodies. For the biocytin-filled cells the

streptavidin conjugate was also added at this step. The slices were washed once for 10 min at RT

with PBS, and incubated for 10 min RT in 0.3 μM DAPI (D3571, Invitrogen) in 0.1 M PB. Sections were

then washed three times at RT with 0.1 M PB, mounted on glass coverslips with gelatin-chrome

alum, air-dried, and closed with Mowiol mounting solution containing 10% (w/v) Mowiol 4-88, 25%

(v/v) glycerol in 0.1 M Tris-buffer (pH 8.5). Sections were kept at 4 °C in the dark until further use.

*Confocal and structured illumination imaging*

Confocal imaging was done on a Zeiss LSM 700 microscope equipped with a plan-apochromat 20x,

0.75 NA and a 63x, 1.4 NA objective, and PMT detectors, or on a Zeiss Elyra PS1 microscope, which is

described below. Confocal images were acquired with optimized settings for laser power, detector

gains and pin hole diameters. Large (2048x2048 pixels), high-resolution images were acquired with a

pixel size of 0.041 μm laterally and 0.110 μm radially to facilitate the comparison with structured-

illumination microscopy (SIM) images. Low-resolution tile images were acquired with a pixel size of

0.274 µm.

Structured-illumination imaging was done on a Zeiss Elyra PS1 system equipped with 488, 561 and

642 nm, 100 mW diode lasers; fluorescence was acquired with a Zeiss plan-apochromat 63x, 1.4 NA

objective and an Andor iXon DU 885 EMCCD camera (1002 x 1004 pixels). Gratings were presented

at 5 phases and 5 rotations for every depth. Sampling interval was 110 nm axially and the

reconstructed super-resolution image has a pixel size of 41 nm. The fluorophore signals were

acquired sequentially. Reconstructions were made with built-in algorithms using ZEN 2012 SP1 black

(v8.1.3.484).

***Electrophysiological analysis***

EPSPs were detected as described previously (Sierksma *et al.*, 2016) based on an EPSP minimal rate

of rise (>0.7 V/s) and an EPSP onset based on the third derivative. EPSPs that were present between

1 to 8 ms post-stimulation were categorized as 'evoked EPSPs'. Next, we grouped the evoked EPSPs

based on stimulus strength, latency and rate of rise. We assumed that with increasing stimulation

intensities previously-recruited EPSPs would still be activated. For consecutive stimulus intensities

Welch's *t* was calculated for the EPSP latency and the rate of rise, and both *t* values were added

together. When the summed *t* value was above 6 and the increase of the response rate of rise was

>20%, we considered the evoked EPSPs different. We visually checked whether this difference was

due to a new input or just a shift in activation probability of a previously-activated input. For a new

input its rate of rise was calculated by subtracting the rates of rise of previously identified inputs

when they were of similar latencies (±0.5 ms). For four recordings we found EPSPs with longer

latencies (Figure 1C), and for these recordings the Welch's *t* test was calculated but restricted to a

latency domain defined by the experimenter. For the example in Figure 1C this was 1-3 ms. Lastly,

the input rate of rise was checked against the spontaneously-occurring EPSP rates of rise, and the

defined input was only accepted if a spontaneous EPSP with a similar rate of rise also occurred.

In addition to the inputs identified by stimulation, we identified inputs within the spontaneous

EPSPs. By comparing the distribution of EPSP rates of rise of all detected EPSPs in a recording with

the evoked EPSPs, we could sometimes identify additional larger inputs (8 large inputs in 8 cells).

Within the spontaneous EPSPs, many were of a small size 0.7-2 V/s and these inputs were therefore

hard to identify. If small EPSPs were not evoked, we visually selected peaks from the histogram of

the EPSP rate of rise (Figure 1C-D) to define these smaller inputs, and added them to the dataset to

acknowledge their existence. However, the inputs with small EPSPs were hard to demarcate and

therefore likely underestimated.

On average, we identified 5.6 inputs per cell. For comparison, prior to synaptic elimination 5-12

axons connect to a principal neuron in EM studies of the mouse MNTB (Hoffpauir *et al.*, 2006). Based

on the comparison of spontaneous and evoked activity, in 8 out of 22 cells we were unable to

activate the cell's largest input. As we were able to distinguish with the stimulation a total of 80

synapses (both large and small) in these 22 cells, at least 1 in 10 axons (8 out of 80), but probably

more on the order of one in three axons (8 out of 22) were not electrically stimulated. Spontaneous

activity partially negated this problem, but may have introduced a bias towards the most active

synapses. To what extent synaptic inputs of the principal neuron are active in every spontaneous

activity burst is unknown, but anecdotal evidence showed extensive co-occurrence of two distinct,

large inputs during spontaneous bursts (Sierksma *et al.*, 2016).

Our method of characterizing EPSP populations in the spontaneous EPSPs has some limitations. We

used the presence of a clear peak in the histogram to indicate the presence of an input. However,

these peaks were generally distinguishable only for the larger EPSPs. Any input giving rise to an EPSP

smaller than 2 V/s would not be distinguishable. Nonetheless, the rates of rise of both the strongest

and second strongest input were very similar when estimated from either spontaneous or evoked

events.

We tested whether the strong inputs could be further subdivided based on the presence of inter-

event intervals that were shorter than the refractory period or the silent period within minibursts

(Sierksma *et al.*, 2016), but this did not lead to the identification of additional connections.

Altogether, we believe that our combined analysis of spontaneous and evoked responses identified

most synaptic inputs, except for some smaller inputs.

***Image analysis***

Analysis of 3D-SIM image stacks was done in ImageJ using the FIJI framework (Schindelin *et al.*, 2012;

Schneider *et al.*, 2012). To determine the area of the postsynaptic cell body covered by the calyx of

Held, we segmented biocytin-stained cell bodies using a custom-designed 3D Sphere Flood fill

algorithm (https://github.com/ErasmusOIC/Sphere_Floodfill). The sphere had a radius of 10 pixels; a

voxel was included in the 3D structure when 40% of its neighboring voxels within the sphere were

above threshold. To account for the width of the synaptic cleft, the segmented cell body was dilated.

Optimal threshold and number of dilation iterations (range 4-19) were determined for each cell by

maximizing the covering ratio. Where needed, axonal protrusions were manually removed from the

segmentation. VGluT-positive structures overlapping with the dilated cell body were segmented in

3D using an automatic threshold (IJ_Isodata) in combination with a regular 3D Flood fill algorithm

(https://github.com/ErasmusOIC/FloodFill3D) using a criterion of 26 neighbors above threshold to

select contiguous structures. The area of the cell body was determined using the outline. The area of

the cell body overlapping with each contiguous VGluT-positive structure was measured. The fraction

of the cell body that was covered by the calyx of Held was calculated as the ratio of the covering

areas and the total area.

All images were pseudo-colored and contrasted in ImageJ. Color gamma enhancement was

performed with Adobe Photoshop 19.1.0. We noticed that the peak intensities of Piccolo and VGluT

did not overlap in SIM images. From line profiles it became clear that Piccolo labeling did overlap

with VGluT at lower intensities, but not at the highest VGluT intensities.

***Statistical analyses***

Linear regression analyses against pup age were performed for the developmental effects. When the

developmental effect was significant, the slope of the fit is reported; when not, the average is

reported, and subsequently the *F*-statistic with the degrees of freedom, and the *p*-value are

reported. Subsequent testing on the same data was Bonferroni-corrected, and are reported as

'corrected *p*'. Effect sizes of a correlation are reported as Pearson's *r*. To compare prespike-

associated strong inputs with the other strong inputs, we performed equal-variances *t*-test for the

EPSP amplitude and latency. EPSP rate of rise and synaptic efficacy to trigger postsynaptic firing ($P_{AP}$)

were correlated with age. Age correction was performed by adding age as an explanatory variable to

the regression analysis. $P_{AP}$ was judged to have non-normally distributed residuals. This was

corrected by performing the regression analysis on the ranks. The *F*-statistic of the regression is

reported and the effect of prespike variable within this regression. Sigmoid and linear fits were

performed with Igor Pro 6.37 (WaveMetrics Inc.). To correlate the competition index with relative

coverage we used Spearman's rank correlation ρ. *p*-values < 0.05 were considered significant.

**Results**

*Postsynaptic responses of* in vivo *stimulated fibers*

We made *in vivo* whole-cell recordings from principal neurons in the rat MNTB around the time the

calyx of Held synapse forms. A bipolar stimulation electrode was placed contralaterally from the

recording site to activate different axonal inputs based on their activation threshold (see Methods;

Figure 1A). Above a threshold of 83 ± 19 μA (mean ± SD, *n* = 30 animals, range: 65-140 μA) a field

potential could be evoked in the MNTB. In a total of 22 cells EPSPs could be evoked, and with

increasing current stimulation we typically observed either the presence of a large, all-or-none,

calyceal-like input, or more graded increases in EPSP size (Figure 1B). In both cases the evoked

responses could reach AP threshold (*n* = 16 of 22 cells). To identify the different inputs, we looked

for jumps in the rates of rise or changes in the latency of the evoked responses following an increase

in stimulus intensity, as detailed in the Methods; these discontinuities were interpreted as the

recruitment of a new input. A comparison of the rates of rise of the evoked EPSP with those of the

spontaneously-occurring EPSPs was used as a further check on the accuracy of the identified inputs

(Figure 1C-D). In 8 out of the 22 cells we were unable to observe the largest spontaneous EPSPs

following the electrical stimulation, suggesting that we did not activate the strongest input for these

cells. EPSP subpopulations were found with typical latencies of 1-3 and 4-8 ms (Figure 1E), as

described in slice studies (Banks & Smith, 1992; Hoffpauir *et al.*, 2006; Marrs & Spirou, 2012). The

large EPSPs always belonged to the subpopulation with 1-3 ms delays (Figure 1E). On average the

EPSP latencies became smaller during the first postnatal week (Figure 1F). In addition to early

myelination (Rozeik & Von Keyserlingk, 1987; Hamano *et al.*, 1996), an increase in axon diameter

and/or conductances may have contributed to this decrease in latency. The different subpopulations

may reflect different developmental stages among the calycigenic axons.

Within this developmental period the principal neuron passes through a stage of exuberant synaptic

connectivity (Holcomb *et al.*, 2013). In addition to the 22 cells in which we recorded both evoked

and spontaneous EPSPs, we recorded in 10 cells only the spontaneous activity. In total, we identified

180 putative inputs in 32 cells, 5.6 ± 0.3 per cell (mean ± SEM). The distribution of input rate of rise

was skewed to the right (Figure 2A). By sorting per cell the inputs on their mean EPSP rate of rise, a

discontinuity was observed at about 10 V/s (Figure 2A-B), which we used as a threshold to

distinguish between weak and strong EPSPs. To have a measure of synaptic competition, we

calculated the competition index, which is defined as the ratio of the rates of rise of the second

strongest and the strongest input (Figure 2C). The competition index decreased during development

($-0.07 \pm 0.02$ day$^{-1}$, $F_{1,30} = 11$, $p = 0.002$). This decrease can signify both an increase of the strongest

input rate of rise and a reduction of the second strongest input. We therefore plotted the two

against each other (Figure 2D). The rate of rise of the strongest input clearly increased during

development ($4.2 \pm 1.1$ V s$^{-1}$ day$^{-1}$, $F_{1,30} = 14$, corrected $p = 0.001$), while the rate of rise of the second

strongest did not change ($3.7 \pm 0.2$ V/s, $F_{1,30} = 2.1$, corrected $p = 0.3$). Inputs identified using

stimulation or spontaneous activity had similar rates of rise for both the strongest (stim.: $19 \pm 3$ V/s,

n = 14; spont.: $16 \pm 2$ V/s, n = 18) and the second strongest input (stim.: $3.9 \pm 0.3$ V/s, n = 13; spont.:

$3.6 \pm 0.3$ V/s, n = 19), suggesting that the selective developmental increase in the rate of rise of the

largest inputs was unrelated to the identification method. In contrast to what may have been

expected for competing inputs, the developmental increase in strength of the strongest input did

not coincide with a decrease in the second strongest input.

***Strong inputs and prespikes***

The effects of electrical stimulation of the calyx of Held synapse have been extensively studied in

slices (Borst & Soria van Hoeve, 2012), but equivalent *in vivo* recordings have not yet been reported.

In nine cells we could activate an input associated with an EPSP >10 V/s that could easily be

distinguished from the other EPSPs. These inputs could be activated with a threshold current of 0.24

± 0.05 mA (mean ± SEM; range 0.12 – 0.40 mA), although the reliability of activation with current

above threshold varied between pups (83 ± 7% of the stimulations, mean ± SEM; range 59-100%).

The average latency of the input's EPSP was $2.1 \pm 0.1$ ms (mean ± SEM, range 1.6 – 2.5 ms) with little

jitter between trials (SD/mean: 2.3 ± 0.4%, range 1.0-3.8%). Given the distance of 500-1000 μm

between the stimulation and recording electrode and a synaptic delay of ~0.5 ms, the propagation

speed can be estimated to be in the order of 1 m/s. The AP travel time became less variable with

development (jitter vs. age, $r$ = -0.75, -0.7 ± 0.2% per day, $F_{1,7}$ = 9.0, $p$ = 0.02). In addition, a decrease

in latency ($r$ = -0.7, -0.18 ± 0.07 ms/day, $F_{1,7}$ = 5.6, $p$ = 0.05) suggested a developmental increase in

conduction speed. This increase is likely underestimated, as we positioned the stimulation electrode

as contralaterally as was permitted by the cranial window to be able to separate the stimulus

artefact from the responses. These developmental changes will contribute towards the precise

timing of the calyx of Held in adult, hearing animals.

The calyx of Held can typically be identified in postsynaptic recordings by the presence of a prespike,

a small deflection preceding the large EPSPs (Forsythe, 1994; Borst *et al.*, 1995; Lorteije *et al.*, 2009).

Prespikes can already be detected in the first postnatal week (Crins *et al.*, 2011; Sierksma *et al.*,

2016). We therefore inspected every strong input whether a prespike was present (Figure 3A). In

four of four cells >P5, six of ten cells of P4-5, and in none of six P2-3 cells large EPSPs were preceded

by a prespike (Figure 3B). Hence, the presence of the prespike before a large EPSP was more likely at

older ages ($r$ = 0.7); a fit with a logistic function yielded a mid-point of 4.1 ± 0.4 days postnatally

(mean ± SD) and a steepness of 0.7 ± 0.4 days, suggesting that within two days most strong inputs

will become associated with a prespike. This switch may be explained by an increase in the currents

over the calyceal membrane facing the synaptic cleft or by an increase in the coupling between the

pre- and the postsynaptic membranes. The currents depend on apposition area (Hoffpauir *et al.*,

2006; Holcomb *et al.*, 2013), time course of the calyceal AP, which accelerates considerably during

early development (Taschenberger & von Gersdorff, 2000; Sierksma & Borst, 2017), isochronicity of

the presynaptic currents, and the subcellular distribution of presynaptic ion channels. In addition,

the coupling depends on synaptic cleft resistivity and postsynaptic admittance (Savtchenko, 2007).

The rapid change in most of these properties during this period should underlie the emergence of

the prespike.

As we observed strong inputs with and without prespike, we asked whether their properties were

different. For the inputs that were electrically stimulated, the strong inputs with and without a

prespike had a latency of 2.1 ± 0.2 ms and 2.2 ± 0.1 ms, respectively (Figure 3C, n = 4 and 5 inputs, $t_7$

= 0.5, $r$ = -0.2, $p$ = 0.6). For the EPSP rate of rise, amplitude and the efficacy to trigger a postsynaptic

AP ($P_{AP}$), we included the strong inputs that were identified in the spontaneous activity. The EPSP

rate of rise was 28 ± 2 V/s versus 19 ± 2 V/s (Figure 3D, age correction, $F_{2,17}$ = 5.3, effect of prespike,

5 ± 4 V/s, $t_{17}$ = 1.2, $p$ = 0.3) and had a coefficient of variation in the range of 7-38%. EPSP amplitude,

which did not significantly correlate with age ($r$ = 0.2), was 19.6 ± 1.1 mV versus 15.6 ± 1.5 mV ($t_{18}$ =

2.1, $r$ = 0.45, $p$ = 0.05). As these EPSPs typically triggered an AP (75 ± 7%, mean ± SEM; range 11-

100%), their amplitudes are underestimated. The fraction of EPSPs that triggered a postsynaptic AP

was 0.88 ± 0.07 versus 0.63 ± 0.11 (Figure 3E, rank-transform and age correction, $F_{2,17}$ = 5.4, effect of

prespike: 7 ± 3, $t_{17}$ = 2.6, $p$ = 0.02). Overall, prespike-associated inputs triggered postsynaptic activity

more effectively. Still, some terminals that did not generate a detectable prespike were comparable

to the calyces in strength and efficacy, as we observed before (Sierksma *et al.*, 2016). Next, we will

discuss the relation between physiological properties and the underlying synaptic structure.

***The active zone protein Piccolo in the neonatal auditory brainstem***

The dramatic developmental change in the glutamatergic innervation of MNTB neurons can be

visualized by VGluT immunolabeling (Rodríguez-Contreras *et al.*, 2008). One limitation is that VGluT

labeling does not mark release sites, which can complicate the identification of the target neuron.

We therefore tested whether Piccolo, an active zone protein known to be already present in P9

calyces (Dondzillo *et al.*, 2010), could be used to visualize the release face of terminals within this

developmental period. Already at P3-4, Piccolo-labeling was found throughout the superior olivary

complex (SOC) in all auditory nuclei ($n$ = 8 rats; Figure 4A). Although the staining was not particularly

strong in the MNTB, it clearly co-labeled large perisomatic structures containing VGluT (Figure 4B).

As early as P2, the Piccolo labeling could delineate an edge of the larger VGluT-clusters (Figure 4C, *n*

= 2 of 5 P2 animals). Piccolo puncta were also found outside the somatic regions of the MNTB, and a

Piccolo punctum was occasionally found within the soma of a principal cell (Figure 4C). Generally,

when imaged with confocal microscopy, the Piccolo edges appeared to be composed of individual

spots, which were close to the diffraction limit (Figure 4C-D). We turned to structured-illumination

microscopy (Gustafsson, 2005), which revealed that the Piccolo clusters were indeed composed of

multiple spots (Figure 4D). These spots were also found in biocytin-filled calyces, indicating that

Piccolo can be present in the active zones of the calyx of Held at P5 ($n$ = 11 of 12 calyces; Figure 4E).

Piccolo thus identifies the release face of synapses and thereby can help to identify the target of the

VGluT cluster in the case of axosomatic terminals (Figure 4C-D). However, dendritic synapses cannot

be reliably identified without labeling the postsynaptic neuron.

***Structure-function correlation within and across cells***

From the 32 recorded cells, we recovered 20 cells with immunolabeling for VGluT (1 P2, 7 P3, 8 P4, 2

P5, 1 P6, 1 P7). For these cells we segmented the soma and the VGluT clusters surrounding the soma

using a new, semi-automatic procedure (see Methods, Figure 5A). This segmentation takes into

account that a VGluT cluster may have multiple contact points with the soma by clustering

connected voxels. A VGluT cluster is thus defined as a contiguous structure of VGluT-positive voxels

and VGluT clusters, by definition, were not in contact with each other. The total somatic surface was

620 ± 60 $\mu m^2$ (mean ± SEM, n = 20), which is comparable to earlier estimates (Sommer *et al.*, 1993;

Hoffpauir *et al.*, 2006). We observed that multiple VGluT clusters contacted the postsynaptic cell, of

which most had a postsynaptic contact area <3 $\mu m^2$ (Figure 5B-C). The total postsynaptic area that

was covered by VGluT ranged from 2.4 $\mu m^2$ to 76.5 $\mu m^2$. As calyces can cover >50% of the soma

(Sätzler *et al.*, 2002; Hoffpauir *et al.*, 2006), our segmentation of VGluT may give us only a lower-

bound approximation of its size. The difference may reflect a delay in VGluT filling after calyx

formation (Rodríguez-Contreras *et al.*, 2008). For each cell we identified the largest and second

largest VGluT cluster. The contact area of the largest cluster on the postsynaptic cell ranged from 0.4

$\mu m^2$ to 59.7 $\mu m^2$ (mean 13.9 ± 3.5 $\mu m^2$). For the second largest cluster it ranged from 0.3 $\mu m^2$ to 10.0

$\mu m^2$ (2.5 ± 0.7 $\mu m^2$). For 3 cells the Piccolo labeling failed. For 9 cells we could see clear Piccolo

labeling coinciding with the largest VGluT cluster directed towards the biocytin-filled cell. For 8 cells

VGluT clusters surrounding the biocytin-labeled soma did not show clear Piccolo labeling, but Piccolo

coincided with VGluT in the surrounding tissue. More in-depth analysis of the Piccolo clusters was

not attempted because of reconstruction-related noise.

If the size of the VGluT cluster is a useful estimate of the size of the terminal, then it should predict

the presence and amplitude of a prespike in our *in vivo* recordings. In five of the recovered cells we

recorded a prespike. The size of the VGluT cluster was moderately correlated with the presence of a

prespike ($\rho$ = 0.63, Figure 5G). The largest VGluT cluster was 30.0 ± 7.5 $\mu m^2$ in the cells in which we

recorded a prespike. This was significantly larger than in the cells without a prespike (8.5 ± 2.9 $\mu m^2$,

$t_{18}$ = 3.3, *p* = 0.004). Next, we compared the prespike peak-to-peak amplitude with the size of the

largest VGluT cluster. We excluded one of the recordings, in which the prespikes had an amplitude

of 15 mV. In this recording a small patch of calyceal membrane may have been present within the

electrode tip (Forsythe, 1994). As the calyx was not stained for biocytin, the membrane of the calyx

was apparently not damaged by the pipette. For the remaining prespikes the average amplitudes

were 0.11, 0.26, 0.30 and 0.37 mV, which corresponded with values of 19.2, 23.1, 25.6 and 59.7 $\mu m^2$

for the largest VGluT cluster (r = 0.8). We cannot exclude that this strong correlation was by chance

($F_{1,2}$ = 3, *p* = 0.2). Also, we observed VGluT clusters with similar contact area among the cells from

which we did not record a prespike. Altogether, the size of the largest VGluT cluster may indicate the

presence of a large terminal, but it is unlikely to reveal the entire contact area made by the terminal.

As synaptic strength depends more on glutamate release than on terminal size, the observed VGluT

cluster may still be a good approximation of synaptic strength. We therefore compared synaptic

strength with the contact area of VGluT clusters (Figure 5E). The size of the strongest input was

correlated to the size of the largest VGluT cluster (log-transform, *r* = 0.8, $F_{1,18}$ = 26, *p* < 0.001). The

structure-strength relation could be approximated by EPSP (V/s) = $\beta$ area$^\alpha$ with $\beta$ = 6.8 ± 0.9 V s$^{-1}$ $\mu m^{-1}$

$^{2\alpha}$ and $\alpha$ = 0.35 ± 0.06. Thus, the input strength was correlated with the cube root of VGluT area. For

the older pups (2 P5, 1 P6 and 1 P7 cell) the structure-function relation was underestimated; three of

four synapses were stronger than expected from their apposition area. This may indicate that there

is a developmental change in the structure-function relation or that the total vesicle-loaded contact

of giant synapses is underestimated as the largest VGluT cluster may be only a small part of the

synapse.

Calyces that innervate a multi-calyceal principal cell are filled with synaptic vesicles (Holcomb *et al.*,

2013). The size of the second largest VGluT cluster was not correlated with the size of the second

strongest input ($r$ = -0.1, $F_{1,16}$ = 0.3, $p$ = 0.6). However, we cannot be sure whether multiple VGluT

clusters are independent inputs or belong to the same synapse.  We therefore compared the

competition index to the ratio of the largest VGluT cluster to the total somatic surface contacted by

VGluT. This ratio ranged from 9-98% (on average 42 ± 7%) and was negatively correlated to the

competition index (Figure 5F, $F_{2,17}$ = 13, $p$ < 0.001, age correction, $\rho$ = -0.7). We thus conclude that

we did not find evidence for large discrepancies between the physiological and morphological size of

inputs at the developing calyx of Held synapse. A large VGluT cluster likely indicates the presence of

a strong input that is much stronger than the other inputs of its target neuron.

In the case of multi-calyceal innervation, the different calyces each occupy a contiguous area,

sometimes at opposite sides of the target neuron (Holcomb *et al.*, 2013). This type of multi-

innervation was not observed here (Figure 5B, C and F).

**Discussion**

In this study we combined *in vivo* electrophysiology with immunolabeling of the developing calyx of

Held synapse. With afferent stimulation we identified multiple inputs for each principal neuron.

Increasing stimulation levels showed either a graded increase or a sudden, large jump in the rate of

rise of the response. The strongest inputs across cells showed a shorter latency. The strongest inputs

were typically at least twice as strong as the other inputs. This divergence increased during

development with little change in the strength of the second strongest input. We found that in the

first postnatal week Piccolo is already present within perisomatic VGluT clusters, which are putative

calyceal terminals. The contact surface of the largest VGluT cluster correlated with the strength of

the strongest input *in vivo*. Furthermore, the relative coverage of the largest VGluT cluster

correlated with the relative strength of the strongest to the second strongest input. Overall, synaptic

structure thus correlated with synaptic strength *in vivo*. The association of a strong input with a

prespike became more likely during development. Prespike-associated inputs were more effective in

triggering postsynaptic activity, but some inputs without a prespike had similar properties.

*Calyceal competition during development?*

The prevalence of multiple calyces on a single principal MNTB neuron is a subject of debate. In slice

recordings multiple strong inputs have been observed (Bergsman *et al.*, 2004; Hoffpauir *et al.*, 2006;

Hoffpauir *et al.*, 2010), and structural evidence for the presence of multiple large terminals on

rodent MNTB neurons has also been obtained (Wimmer *et al.*, 2004; Hoffpauir *et al.*, 2006; Holcomb

*et al.*, 2013; Matho, 2013). Although only a minority (up to 10%) of neurons is persistently

innervated by multiple calyces (Holcomb *et al.*, 2013; Matho, 2013), as many as half of the principal

neurons are putatively contacted during development by multiple large terminals (Holcomb *et al.*,

2013). On the other hand, to our knowledge, not a single conclusive example of multi-calyceal

innervation has been reported for *in vivo* studies of the adult MNTB. In this developmental study,

neither our *in vivo* recordings nor our synapse reconstructions revealed multi-calyceal innervation,

but weak calyces with a low density of VGluT, or inputs that are unable to generate prespikes would

not have been identified as such. In our earlier study, three out of 132 cells were found that each

had two strong inputs that could be separated based on the presence or absence of a prespike

(Sierksma *et al.*, 2016). As we observed large VGluT clusters without the identification of prespike-

associated input for their target cells, these cases may have been examples of multi-calyceal

innervation, but we cannot exclude that one of the inputs was dendritic. From our earlier (3/132

cells) and the current (0/32 cells) experiments, we thus estimate that <10% of rat principal neurons

has multi-calyceal innervation during development.

Initial neural connectivity is extensively remodeled to form the final neural circuitry (Katz & Shatz,

1996). As the remodeling will strengthen proper inputs and eliminate improper ones, it is viewed as

a competition between inputs with their activity and efficacy in triggering postsynaptic APs as

instructive cues. The principal neuron has an extreme developmental outcome by allowing the

formation of only one giant synapse amidst multiple calycigenic inputs (Morest, 1968; Baydyuk *et al.*,

2016). The presence of competition has been evaluated by the (relative) contact surface (Holcomb *et*

*al.*, 2013) or by the (relative) strength of the inputs (Hoffpauir *et al.*, 2010; Sierksma *et al.*, 2016).

Here, we compared a structural and a functional measure for competition, and generally observed a

good match between the two. One limitation is that our immunolabeling did not allow the

unambiguous identification of different inputs based on the VGluT clusters, which limits us to give a

definite measure of structural competition. Still, the observed match between VGluT relative

coverage and the competition index indicates that the physiological and morphological development

are linked.

Two types of competition have been contrasted previously (Holcomb *et al.*, 2013), exemplified by

the neuromuscular junction (NMJ) and the cerebellar climbing fiber-Purkinje cell synapse (CF-PC).

Before mono-innervation is established in the case of the NMJ, one input becomes >4 fold stronger

than the other ones by both an increase of the strongest and a decrease of the other inputs (Colman

*et al.*, 1997). As NMJ inputs compete at and for a small patch of the muscle membrane (Sanes &

Lichtman, 1999; Darabid *et al.*, 2014), the expansion of one NMJ necessarily involves the shrinkage

of another, as reflected in the opposing changes in strength. In contrast, CFs compete at the soma

for the apical dendrite (Watanabe & Kano, 2011). A fourfold strengthening of the strongest input

was observed (Hashimoto & Kano, 2003) while other inputs remained constant (Bosman *et al.*,

2008). Subsequently, one of the immature CFs is allowed to translocate to the apical dendrite

(Hashimoto *et al.*, 2009; Watanabe & Kano, 2011), ending the competition (Carrillo *et al.*, 2013). We

observed that the strongest input increases in strength without a substantial change for the second

strongest. This may indicate that they do not compete for overlapping synaptic domains (Holcomb *et*

*al.*, 2013). In addition, we observed that the strongest input can already be four times stronger than

the second strongest while its coverage is still low. This might indicate that the increase in strength

precedes and instructs the expansion (Hoffpauir *et al.*, 2010). The inverse, a large VGluT cluster with

a high competition index, was not observed, suggesting that VGluT filling of an axosomatic terminal

might be a late stage in maturation (Rodríguez-Contreras *et al.*, 2008).

***Structure-function relation for the developing calyx of Held synapse***

The presence of a prespike indicates the innervation by a giant terminal (Forsythe, 1994). In this

study four cells with a prespike-associated input had a large VGluT cluster and one cell did not. This

may indicate that the terminal was larger than revealed by VGluT labeling (Rodríguez-Contreras *et*

*al.*, 2008). Four cells were observed where a large VGluT cluster did not match with a prespike-

associated input. By recording in each pup only a single neuron, we ensured that the retrieved cell

was the recorded cell, as evidenced by the clear match between synapse strength and VGluT cluster.

The capacitive filtering by the soma may have obscured small prespikes that might have been

detectable in voltage-clamp recordings. There are also a number of biological factors that influence

the prespike amplitude (see Results). For now, the low number of cells with a discrepancy between

structure and morphology does not warrant any firm conclusions. Experimental observations on the

relation between presynaptic AP and the prespike have so far been qualitative (Forsythe, 1994; Borst

*et al.*, 1995; Wang *et al.*, 2017). Experimental and computational studies on the prespike

(Savtchenko, 2007) based on exact morphology (Spirou *et al.*, 2008) may give us an answer under

which conditions a calyx generates a prespike.

From our structure-function analyses we found that VGluT labeling closely follows the strength of

the strongest input. This VGluT cluster likely represents the total pool of glutamatergic vesicles in the

calyx of Held (Rodríguez-Contreras *et al.*, 2008). The relation between the size of this pool and

synaptic strength is complex, depending on among others quantal size, release probability, and

number of release sites (Atwood & Karunanithi, 2002; Schneggenburger & Forsythe, 2006; Borst &

Soria van Hoeve, 2012). We found a cube root relation between VGluT contact surface and EPSP rate

of rise, suggesting that the increase in vesicle pool is relatively ineffective in increasing synaptic

strength. This can be partially explained by homeostatic plasticity of postsynaptic excitability

(Hoffpauir *et al.*, 2010; Rusu & Borst, 2011; Sierksma *et al.*, 2016). Another possibility is that the

input also has formed dendritic synapses (Rodríguez-Contreras *et al.*, 2008; Holcomb *et al.*, 2013).

However, the EPSC-EPSP relation saturates for large EPSCs (Lorteije *et al.*, 2009; Sierksma *et al.*,

2016), indicating that for weak inputs (small clusters) the relation may be more linear.

For the study of the relationship between structure and function, the calyx of Held synapse has the

advantage of an axosomatic, (putative) one-to-one innervation facilitating the identification of the

strongest input *in vivo* with the largest cluster *ex vivo*. An even more direct relationship might be

possible following simultaneous electrophysiology and calcium imaging *in vivo* (Winnubst *et al.*,

2015). Subsequent reconstruction of the synapses after *in vivo* imaging (Liang *et al.*, 2018) would

allow a structure-function description for multiple synapses of a single cell. Such studies may further

define the relationship between morphological and functional properties of the precursors of the

calyx of Held. Finally, longitudinal imaging might reveal some of the dynamics associated with the

formation of the calyx of Held (Rodríguez-Contreras *et al.*, 2008).

**References**

Atwood HL & Karunanithi S. (2002). Diversification of synaptic strength: presynaptic elements. *Nat*

*Rev Neurosci* **3,** 497-516.

Banks MI & Smith PH. (1992). Intracellular recordings from neurobiotin-labeled cells in brain slices of

the rat medial nucleus of the trapezoid body. *Journal of Neuroscience* **12,** 2819-2837.

Baydyuk M, Xu J & Wu LG. (2016). The calyx of Held in the auditory system: Structure, function, and

development. *Hear Res* **338,** 22-31.

Bergsman JB, De Camilli P & McCormick DA. (2004). Multiple large inputs to principal cells in the

mouse medial nucleus of the trapezoid body. *J Neurophysiol* **92,** 545-552.

Borst JGG, Helmchen F & Sakmann B. (1995). Pre- and postsynaptic whole-cell recordings in the

medial nucleus of the trapezoid body of the rat. *J Physiol* **489,** 825-840.

Borst JGG & Soria van Hoeve J. (2012). The calyx of Held synapse: from model synapse to auditory

relay. *Annual Review of Physiology* **74,** 199-224.

Bosman LWJ, Takechi H, Hartmann J, Eilers J & Konnerth A. (2008). Homosynaptic long-term synaptic

potentiation of the "winner" climbing fiber synapse in developing Purkinje cells. *J Neurosci*

**28,** 798-807.

Carrillo J, Nishiyama N & Nishiyama H. (2013). Dendritic translocation establishes the winner in

cerebellar climbing fiber synapse elimination. *J Neurosci* **33,** 7641-7653.

Chuhma N & Ohmori H. (1998). Postnatal development of phase-locked high-fidelity synaptic

transmission in the medial nucleus of the trapezoid body of the rat. *J Neurosci* **18,** 512-520.

Colman H, Nabekura J & Lichtman JW. (1997). Alterations in synaptic strength preceding axon

withdrawal. *Science* **275,** 356-361.

Crins TTH, Rusu SI, Rodríguez-Contreras A & Borst JGG. (2011). Developmental changes in short-term

plasticity at the rat calyx of Held synapse. *J Neurosci* **31,** 11706-11717.

Darabid H, Perez-Gonzalez AP & Robitaille R. (2014). Neuromuscular synaptogenesis: coordinating

partners with multiple functions. *Nature Reviews Neuroscience* **15,** 703.

Denk W & Horstmann H. (2004). Serial block-face scanning electron microscopy to reconstruct three-

dimensional tissue nanostructure. *PLOS Biology* **2,** e329.

Dondzillo A, Sätzler K, Horstmann H, Altrock WD, Gundelfinger ED & Kuner T. (2010). Targeted three-

dimensional immunohistochemistry reveals localization of presynaptic proteins Bassoon and

Piccolo in the rat calyx of Held before and after the onset of hearing. *J Comp Neurol* **518,**

1008-1029.

Forsythe ID. (1994). Direct patch recording from identified presynaptic terminals mediating

glutamatergic EPSCs in the rat CNS, *in vitro*. *J Physiol* **479,** 381-387.

Gundelfinger ED, Reissner C & Garner CC. (2016). Role of Bassoon and Piccolo in assembly and

molecular organization of the active zone. *Frontiers in Synaptic Neuroscience* **7**.

Gustafsson MGL. (2005). Nonlinear structured-illumination microscopy: Wide-field fluorescence

imaging with theoretically unlimited resolution. *Proceedings of the National Academy of*

*Sciences of the United States of America* **102,** 13081-13086.

Hamano K, Iwasaki N, Takeya T & Takita H. (1996). A quantitative analysis of rat central nervous

system myelination using the immunohistochemical method for MBP. *Dev Brain Res* **93,** 18-

22.

Hashimoto K, Ichikawa R, Kitamura K, Watanabe M & Kano M. (2009). Translocation of a "winner"

climbing fiber to the Purkinje cell dendrite and subsequent elimination of "losers" from the

soma in developing cerebellum. *Neuron* **63,** 106-118.

Hashimoto K & Kano M. (2003). Functional differentiation of multiple climbing fiber inputs during

synapse elimination in the developing cerebellum. *Neuron* **38,** 785-796.

Held H. (1893). Die centrale Gehörleitung. *Archiv fur Anatomie und Physiologie, Anatomie Abtheil***,**

201-248.

Hoffpauir BK, Grimes JL, Mathers PH & Spirou GA. (2006). Synaptogenesis of the calyx of Held: rapid

onset of function and one-to-one morphological innervation. *J Neurosci* **26,** 5511-5523.

Hoffpauir BK, Kolson DR, Mathers PH & Spirou GA. (2010). Maturation of synaptic partners:

functional phenotype and synaptic organization tuned in synchrony. *J Physiol* **588,** 4365-

4385.

Holcomb PS, Hoffpauir BK, Hoyson MC, Jackson DR, Deerinck TJ, Marrs GS, Dehoff M, Wu J, Ellisman

MH & Spirou GA. (2013). Synaptic inputs compete during rapid formation of the calyx of

Held: a new model system for neural development. *J Neurosci* **33,** 12954-12969.

Kandler K & Friauf E. (1993). Pre- and postnatal development of efferent connections of the cochlear

nucleus in the rat. *Journal of Comparative Neurology* **328,** 161-184.

Katz LC & Shatz CJ. (1996). Synaptic activity and the construction of cortical circuits. *Science* **274,**

1133-1138.

Kuwabara N, DiCaprio RA & Zook JM. (1991). Afferents to the medial nucleus of the trapezoid body

and their collateral projections. *Journal of Comparative Neurology* **314,** 684-706.

Liang L, Fratzl A, Goldey G, Ramesh RN, Sugden AU, Morgan JL, Chen C & Andermann ML. (2018). A

fine-scale functional logic to convergence from retina to thalamus. *Cell* **173,** 1343-1355

e1324.

Lorteije JAM, Rusu SI, Kushmerick C & Borst JGG. (2009). Reliability and precision of the mouse calyx

of Held synapse. *J Neurosci* **29,** 13770-13784.

Marrs GS & Spirou GA. (2012). Embryonic assembly of auditory circuits: spiral ganglion and

brainstem. *J Physiol* **590,** 2391-2408.

Matho KSH. (2013). Structure and dynamism in a central neural circuit in adulthood and postnatal

development. In *École Doctorale Cerveau, Cognition, Comportement*. Université Pierre et

Marie Curie, Paris, France.

Morest DK. (1968). The growth of synaptic endings in the mammalian brain: a study of the calyces of

the trapezoid body. *Zeitschrift für Anatomie und Entwicklungsgeschichte* **127,** 201-220.

Qiu X, Zhu Q & Sun J. (2015). Quantitative analysis of vesicle recycling at the calyx of Held synapse.

*Proceedings of the National Academy of Sciences* **112,** 4779-4784.

Rodríguez-Contreras A, de Lange RPJ, Lucassen PJ & Borst JGG. (2006). Branching of calyceal

afferents during postnatal development in the rat auditory brainstem. *J Comp Neurol* **496,**

214-228.

Rodríguez-Contreras A, van Hoeve JS, Habets RLP, Locher H & Borst JGG. (2008). Dynamic

development of the calyx of Held synapse. *Proc Natl Acad Sci U S A* **105,** 5603-5608.

Rollenhagen A & Lübke JHR. (2006). The morphology of excitatory central synapses: from structure

to function. *Cell Tissue Res* **326,** 221-237.

Rozeik C & Von Keyserlingk D. (1987). The sequence of myelination in the brainstem of the rat

monitored by myelin basic protein immunohistochemistry. *Developmental Brain Research*

**35,** 183-190.

Rusu SI & Borst JGG. (2011). Developmental changes in intrinsic excitability of principal neurons in

the rat medial nucleus of the trapezoid body. *Dev Neurobiol* **71,** 284-295.

Sanes JR & Lichtman JW. (1999). Development of the vertebrate neuromuscular junction. *Annu Rev*

*Neurosci* **22,** 389-442.

Sätzler K, Söhl LF, Bollmann JH, Borst JGG, Frotscher M, Sakmann B & Lübke JHR. (2002). Three-

dimensional reconstruction of a calyx of Held and its postsynaptic principal neuron in the

medial nucleus of the trapezoid body. *J Neurosci* **22,** 10567-10579.

Savtchenko LP. (2007). Bilateral processing in chemical synapses with electrical 'ephaptic' feedback:

a theoretical model. *Math Biosci* **207,** 113-137.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P & Cardona

A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9,** 676-

682.

Schneggenburger R & Forsythe ID. (2006). The calyx of Held. *Cell Tissue Res* **326,** 311-337.

Schneider CA, Rasband WS & Eliceiri KW. (2012). NIH Image to ImageJ: 25 years of image analysis.

*Nat Methods* **9,** 671-675.

Sierksma MC & Borst JGG. (2017). Resistance to action potential depression of a rat axon terminal in

vivo. *Proceedings of the National Academy of Sciences* **114,** 4249-4254.

Sierksma MC, Tedja MS & Borst JGG. (2016). *In vivo* matching of postsynaptic excitability with

spontaneous synaptic inputs during formation of the rat calyx of Held synapse. *J Physiol* **595,**

207-231.

Sommer I, Lingenhöhl K & Friauf E. (1993). Principal cells of the rat medial nucleus of the trapezoid

body: an intracellular in vivo study of their physiology and morphology. *Experimental Brain*

*Research* **95,** 223-239.

Soria Van Hoeve J & Borst JGG. (2010). Delayed appearance of the scaffolding proteins PSD-95 and

Homer-1 at the developing rat calyx of Held synapse. *J Comp Neurol* **518,** 4581-4590.

Spirou GA, Chirila FV, von Gersdorff H & Manis PB. (2008). Heterogeneous $Ca^{2+}$ influx along the adult

calyx of Held: a structural and computational study. *Neuroscience* **154,** 171-185.

Südhof Thomas C. (2012). The presynaptic active zone. *Neuron* **75,** 11-25.

Taschenberger H, Leão RM, Rowland KC, Spirou GA & von Gersdorff H. (2002). Optimizing synaptic

architecture and efficiency for high-frequency transmission. *Neuron* **36,** 1127-1143.

Taschenberger H & von Gersdorff H. (2000). Fine-tuning an auditory synapse for speed and fidelity:

developmental changes in presynaptic waveform, EPSC kinetics, and synaptic plasticity. *J*

*Neurosci* **20,** 9162-9173.

Wang T, van Woerden GM, Elgersma Y & Borst JGG. (2017). Enhanced transmission at the calyx of

Held synapse in a mouse model for Angelman syndrome. *Front Cell Neurosci* **11,** 418.

Watanabe M & Kano M. (2011). Climbing fiber synapse elimination in cerebellar Purkinje cells. *Eur J*

*Neurosci* **34,** 1697-1710.

Wimmer VC, Nevian T & Kuner T. (2004). Targeted in vivo expression of proteins in the calyx of Held.

*Pflügers Arch* **449,** 319-333.

Winnubst J, Cheyne JE, Niculescu D & Lohmann C. (2015). Spontaneous activity drives local synaptic

plasticity in vivo. *Neuron* **87,** 399-410.

Xiao L, Michalski N, Kronander E, Gjoni E, Genoud C, Knott G & Schneggenburger R. (2013). BMP

signaling specifies the development of a large and fast CNS synapse. *Nat Neurosci* **16,** 856-

864.

Yu WM & Goodrich LV. (2014). Morphological and physiological development of auditory synapses.

*Hear Res* **311,** 3-16.

**Figure 1. Responses to current stimulation of developing calycigenic axons.** (A) Schematic drawing

of the experimental approach. A bipolar stimulation electrode straddling the trapezoid body was

positioned contralaterally from the recorded MNTB, and whole-cell recordings were made from

principal cells of the MNTB. The basilar artery (rostrocaudal, thick red line) and the anterior-inferior

cerebellar artery (mediolateral, red line) served as landmarks. Arrows indicate lateral (L) and rostral

(R) directions. (B) Waterfall plots of two whole-cell recordings, sorted on stimulation intensity (color-

coded as indicated in calibration bars). In the P4 example (left) a gradual increase in the evoked EPSP

was observed, suggesting the recruitment of multiple inputs, while in the P6 example (right) a single,

large EPSP was observed, which elicited a postsynaptic AP. Schematic drawing depicts the assumed

synaptic innervation. Scale bars indicate 25 mV and 2 ms. (C) Identification of synaptic inputs. Rates

of rise of evoked EPSPs vs. stimulation intensity (top) or EPSP latency (middle) for the P4 example

shown in B. The bottom graph shows the histograms of the rates of rise of evoked and spontaneous

EPSPs. Arrow heads on top of the graph indicate the rates of rise of identified inputs. Data points in

the top graph were horizontally scattered by <50 µA for visualization purposes. (D) As C, except for

the P6 example shown in B. (E) Cumulative distribution of latencies of all evoked EPSPs (black) and of

large EPSPs (>10 V/s, blue). Individual cells are shown in light blue and grey. Large EPSPs have a

shorter latency than most other EPSPs. (F) The average cumulative distribution of the evoked EPSP

latencies grouped by age suggests a developmental increase in propagation speed.

Figure 2. Synaptic competition in MNTB follows the expansion of the strongest input without

changing the second strongest input. (A) Histogram of the average rate of rise for all inputs (open)

and the strongest inputs (closed). Line indicates the threshold for 'strong inputs'. (B) For every cell

the inputs are incrementally sorted by their average EPSP rate of rise. Diamonds indicate the inputs

associated with a prespike. Colors indicate the ages as in C. Note the root scale of the ordinate. (C)

Cumulative distributions of the competition index, i.e. the ratio of the rates of rise of the second

strongest to the strongest input. (D) The rate of rise of the second strongest input against the

strongest. Markers correspond to individual principal cells. Closed markers indicate that both inputs

were identified based on spontaneous activity. Line shows identity line. Colors indicate age as in C.

**Figure 3. Comparison of strong inputs with and without prespikes.** (A) Large EPSPs of a strong input

with (left; P6) and of a strong input without (right; P5) a prespike. EPSPs were aligned on their onset.

The insets show the same traces with their $V_m$ offsets subtracted. Scale bars indicate 10 mV and 1

ms, and scale bars in the inset indicate 0.5 mV and 0.2 ms. (B) Developmental changes in the

presence of a prespike preceding a large EPSP. Circles indicate cells where a large EPSP was

recorded. Bars represent age average. Blue line represents the fit of the logistic function with the

values reported in the graph. (C-E) Comparison of strong inputs with or without a prespike. In C only

evoked large EPSPs were analyzed. Circles indicate individual cells. Line represents the average and

SEM.

**Figure 4. Developmental expression of Piccolo in the MNTB.** (A) Fluorescent labeling of Piccolo

(green), VGluT (magenta) and DAPI (blue) in the SOC of a P4 rat. Midline is indicated with a white

line. Auditory nuclei are delineated with dashed lines. (B) Higher magnification of the grey box in A

showing principal cells of the MNTB with large perisomatic clusters of VGluT and Piccolo. (C) A P2

and P5 principal cell with fluorescent labeling of VGluT, Piccolo, and VGluT + Piccolo + DAPI with

confocal microscopy. (D) Comparison of confocal microscopy and structured-illumination microscopy

(SIM) of perisomatic clusters of Piccolo and VGluT of the P2 principal cell shown in C (top), and of the

P5 principal cell shown in C (bottom). (E) SIM image of a P5 biocytin-filled calyx (top, blue) with

VGluT and Piccolo, revealing active zone-like structures in the Piccolo labeling (bottom). Non-

calyceal labeling was masked. Scale bars indicate 200 µm in A, 15 µm in B, 10 µm in C, 2 µm in D, and

10 and 1 µm in E. Images in C and D were background-subtracted. Abbreviations: LSO, lateral

superior olive; MNTB, medial nucleus of the trapezoid body; P, postnatal; SIM, structured-

illumination microscopy; SOC, superior olivary complex; SPON, superior periolivary nucleus; VGluT,

vesicular glutamate transporter 1 and 2; VNTB, ventral nucleus of the trapezoid body.

**Figure 5. Contact area of perisomatic VGluT clusters correlates with synaptic strength *in vivo*.** (A)
Example of segmentation of VGluT clusters around the soma. The image before (top) and after
segmentation (below). Blue: biocytin; Green: VGluT labeling. In the bottom image different VGluT
clusters are depicted by different shades of green. Red line indicates contour of the cell after
dilation. For this cell 1 input was associated with a prespike. (B-C) 3D reconstructions of somata
(blue) with VGluT clusters (green). Dendrites were manually removed. Right image is the same soma
after 180 degrees rotation. Postsynaptic cell in B did not have a prespike-associated input. The VGluT
cluster in C was manually separated from a nearby cluster that terminated on another cell. (D)

Largest VGluT cluster against the presence of a prespike-associated input. (E) Input strength of

strongest input against contact area of the largest VGluT cluster for each cell. Line indicates linear fit

of log-transformed variables. Dotted line indicates 95% confidence intervals. (F) Relation between

area of largest and second largest VGluT cluster. Shaded area: possibly multi-calyceal innervation.

(G) Competition index against relative coverage. Competition index equals the second strongest

input relative to the strongest input. Relative coverage equals the contact area of the largest VGluT

cluster relative to the total area contacted by VGluT clusters. Shaded area indicates region with low

competition index and low relative coverage. Colors in D-G indicate age as in previous figures.

Triangle indicates the data point that corresponds to the cell shown in A. Closed circle and diamond

indicate data points corresponding to the cells shown in B and C, respectively.