Increased antibiotic susceptibility in *Neisseria gonorrhoeae* through

adaptation to the cervical environment

Kevin C Ma[1¶], Tatum D Mortimer[1¶], Allison L Hicks[1], Nicole E Wheeler[2], Leonor Sánchez-Busó[2],

Daniel Golparian[3], George Taiaroa[4], Daniel HF Rubin[1], Yi Wang[1], Deborah A Williamson[4],

Magnus Unemo[3], Simon R Harris[5], Yonatan H Grad[1,6*]

[1] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public

Health, Boston, USA

[2] Centre for Genomic Pathogen Surveillance, Wellcome Sanger Institute, Wellcome Genome

Campus, Hinxton, Cambridgeshire, United Kingdom

[3] WHO Collaborating Centre for Gonorrhoea and other STIs, Swedish Reference Laboratory for

STIs, Faculty of Medicine and Health, Örebro University, Örebro, Sweden

[4] Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology and

Immunology, The University of Melbourne at The Peter Doherty Institute for Infection and

Immunity, Melbourne, Australia

[5] Microbiotica Ltd, Biodata Innovation Centre, Wellcome Genome Campus, Hinxton,

Cambridgeshire, United Kingdom

[6] Division of Infectious Diseases, Brigham and Women's Hospital and Harvard Medical School,

Boston, USA

* Corresponding author

E-mail: ygrad@hsph.harvard.edu (YHG)

¶ These authors contributed equally to this work.

Abstract

Neisseria gonorrhoeae is an urgent public health threat due to rapidly increasing incidence and antibiotic resistance. In contrast with the trend of increasing resistance, clinical isolates that have reverted to susceptibility regularly appear, prompting questions about which pressures compete with antibiotics to shape gonococcal evolution. Here, we used genome-wide association on the largest collection of *N. gonorrhoeae* isolates to date (n=4882) to identify loss-of-function (LOF) mutations in the efflux pump *mtrCDE* operon as a mechanism of increased antibiotic susceptibility and demonstrate that these mutations are overrepresented in cervical isolates relative to urethral isolates (odds ratio (OR) = 3.74, 95% CI [1.98-6.70]). In support of a model in which pump expression incurs a fitness cost in this niche, cervical isolates were also enriched relative to urethral isolates in LOF mutations in the *mtrCDE* activator *mtrA* (OR = 8.60, 95% CI [4.96-14.57]) and in *farA*, a subunit of the FarAB efflux pump (OR = 6.25, 95% CI [3.90-9.83]). In total, approximately 1 in 3 cervical isolates (36.4%) contained a LOF mutation in either the efflux pump components *mtrC* or *farA* or the activator *mtrA*. Our findings extend beyond *N. gonorrhoeae* to other Neisseria: *mtrC* LOF mutations are rare (<1%) in the primarily nasopharyngeal-colonizing *N. meningitidis* in a collection of 14,798 genomes but enriched in a heterosexual urethritis-associated lineage (8.6%, p = 9.90×10^{-5}), indicating that efflux pump downregulation contributes broadly to the adaptation of pathogenic *Neisseria* to the female urogenital tract. Overall, our findings highlight the impact of integrating microbial population genomics with host metadata and demonstrate how host environmental pressures can lead to increased antibiotic susceptibility.

Introduction

Neisseria gonorrhoeae is the causative agent of the sexually transmitted disease gonorrhea. Antibiotics have played a key role in shaping gonococcal evolution[1-3], with *N. gonorrhoeae* gaining resistance to each of the first line antibiotics used to treat it[4-6]. As *N. gonorrhoeae* is an obligate human pathogen, the mucosal niches it infects—most commonly including the urethra, cervix, pharynx, and rectum—must also influence its evolution[7]. The gonococcal phylogeny suggests the interaction of these factors, with an ancestral split between a relatively drug-susceptible lineage circulating primarily in heterosexuals and a relatively drug-resistant lineage circulating primarily in men who have sex with men[3].

Despite the deeply concerning increase in antibiotic resistance reported in gonococcal populations globally[8], some clinical isolates of *N. gonorrhoeae* have become more susceptible to antibiotics[9,10]. This unexpected phenomenon prompts questions about which environmental pressures could be drivers of increased susceptibility and the mechanisms by which suppression or reversion of resistance may occur. To address these questions, we analyzed the genomes of a global collection of clinical isolates together with patient demographic and clinical data to identify mutations associated with increased susceptibility and define the environments in which they appear.

Antibiotic susceptibility is influenced by unknown genetic determinants in *N. gonorrhoeae*

We first assessed how well variation in antibiotic resistance phenotype was captured by the presence and absence of known resistance markers. To do so, we assembled and examined a global dataset comprising the genomes and minimum inhibitory concentrations (MICs) of 4882 isolates collected across 66 countries and 38 years (Figure 1, Supplementary Table 1)[3,9-20]. We modeled log-transformed MICs using multiple regression on a panel of experimentally characterized resistance markers for the three most clinically relevant antibiotics[21,22]. This enabled us to make quantitative predictions of MIC based on known genotypic markers and to assess how well these markers predicted true MIC values. For the macrolide azithromycin, we observed that 367/4461 (8.23%) isolates had predicted MICs that deviated by two dilutions or more from their true reported phenotypic values. The majority (66.8%) of these isolates had MICs that were lower than expected, indicative of susceptibility unexplained by the genetic determinants in our model. Overall MIC variance explained by known resistance mutations was relatively low (adjusted R^2 = 0.681), in agreement with prior studies that employed whole-

genome supervised learning algorithms to predict azithromycin resistance[23]. MIC variance
explained by known resistance mutations was also low for ceftriaxone (adjusted R^2 = 0.719) but
higher for ciprofloxacin (adjusted R^2 = 0.944), with 1.52% and 2.19% of strains, respectively,
exhibiting two dilutions or lower reported MICs compared to predictions, similarly indicating
unexplained susceptibility. The predictive modeling results therefore suggested unknown
modifiers that promote susceptibility for multiple classes of antibiotics in *N. gonorrhoeae*.
91
92

Figure 1 – Population structure and susceptibility profile of *N. gonorrhoeae* global meta-
analysis collection. A midpoint rooted recombination-corrected maximum likelihood phylogeny
of 4882 genomes based on 68697 SNPs (Supplementary Table 1) was annotated with binarized
resistance (ciprofloxacin) or decreased susceptibility (azithromycin, ceftriaxone) values.
Annotation rings are listed in order of ciprofloxacin (purple), ceftriaxone (blue), and azithromycin
(pink) from innermost to outermost. Branch length represents total number of substitutions after
removal of predicted recombination.
102

Microbial genome-wide association identifies a variant that contributes to susceptibility in the operon encoding the MtrCDE efflux pump

To identify novel antibiotic susceptibility loci in an unbiased manner, we conducted a bacterial genome-wide association study (GWAS). We used a linear mixed model framework to control for population structure, and we used unitigs from genome assemblies to capture SNPs, indels, and accessory genome elements[24-26]. We performed a GWAS on the sequences of 4463 isolates with associated azithromycin MICs using a Bonferroni-corrected significance threshold of 2.97×10^{-7}. The linear mixed model adequately controlled for population structure (Supplementary Figure 1), and the proportion of phenotypic MIC variance attributable to genotype (i.e., narrow-sense heritability) estimated by the linear mixed model was high ($h^2 =$ 0.97). In line with this, we observed highly significant unitigs with positive effect sizes corresponding to the known resistance substitutions C2611T and A2059G (*E. coli* numbering) in the 23S ribosomal RNA gene as positive controls (Figure 2)[27]. The next most significant variant was a unitig associated with increased susceptibility that mapped to *mtrC* (β, or effect size, = -2.80, 95% CI [-3.04, -2.56]; p-value = 3.30×10^{-109}). Overexpression of the *mtrCDE* efflux pump operon has been shown to decrease gonococcal susceptibility to a range of hydrophobic acids and antimicrobial agents[4,28], and conversely, knockout of the pump results in multi-drug hypersusceptibility[29]. To assess whether this *mtrC* variant was associated with increased susceptibility to other antibiotics, we performed GWAS for ceftriaxone (for which MICs were available from 4090 isolates) and for ciprofloxacin (4105 isolates). We recovered known ceftriaxone resistance mutations including recombination in the *penA* gene and ciprofloxacin resistance substitutions in DNA gyrase (*gyrA*). In agreement with the known pleiotropic effect of the MtrCDE efflux pump[29], we observed the same *mtrC* unitig at genome-wide significance associated with increased susceptibility to both ceftriaxone (β = -1.15, 95% CI [-1.32, -0.98]; p-value = 2.84×10^{-38}) and ciprofloxacin (β = -1.30, 95% CI [-1.55, -1.05]; p-value = 1.25×10^{-23}) (Figure 2). Across all three drugs, heritability estimates for this *mtrC* variant were comparable to that of prevalent major resistance determinants (azithromycin h^2: 0.323; ceftriaxone h^2: 0.199; ciprofloxacin h^2: 0.156), indicating that unexplained susceptibility in our model could be partially addressed by inclusion of this mutation.

Annotation of the *mtrC* unitig revealed that it represented a two base pair deletion in a 'GC' dinucleotide hexarepeat, leading to early termination of *mtrC* translation and probable loss of MtrCDE activity[30] (Figure 2 inset). A laboratory-generated gonococcal mutant with a four base pair deletion in this same *mtrC* dinucleotide hexarepeat exhibited multi-drug susceptibility[30], and

clinical gonococcal isolates hypersensitive to erythromycin were shown to have mutations

mapping to this locus[31]. To directly test the hypothesis that the two base pair deletion also

contributed to increased susceptibility for the panel of antibiotics we examined, we

complemented the mutation in a clinical isolate belonging to the multidrug-resistant lineage ST-

1901[32] and observed significant increases in MICs for all three antibiotics, as predicted by the

GWAS (Supplementary Table 2).

Figure 2 – GWAS identifies a variant mapping to *mtrC* associated with increased
susceptibility to azithromycin, ceftriaxone, and ciprofloxacin. Negative log$_{10}$-transformed p-
values for unitigs tested in GWAS on MICs to azithromycin (pink, n=4463), ceftriaxone (blue,
n=4090), and ciprofloxacin (purple, n=4105) are shown in the Manhattan plot. The sign of the
regression coefficient β (indicating positive or negative association) is depicted in the legend.
Known influential resistance determinants were annotated and the *mtrC* variant associated with
susceptibility highlighted in gray. Inset: schematic of the *mtr* genetic regulon including structural
genes *mtrCDE*, the activator *mtrA*, and the repressor *mtrR*. The genomic location and
nucleotide change of the *mtrC* GWAS variant relative to the gonococcal NCCP11945 reference
genome is shown.

We searched for additional *mtrC* loss-of-function (LOF) mutations and found six clinical isolates

with genomes encoding indels outside of the dinucleotide hexarepeat that also were associated

with increased susceptibility (Supplementary Figure 2A). Ten isolates that had acquired the two

base pair deletion also have a two base pair insertion elsewhere in *mtrC* that restores the

original coding frame, suggesting that loss of MtrC function may be reverted by further mutation

or recombination (Supplementary Figure 2A). In line with this, *mtrC* LOF mutations have

emerged numerous times throughout the phylogeny (Supplementary Figure 3), indicative of

possible repeated gains and losses of a dinucleotide in the hexarepeat region due to DNA

polymerase slippage[33]. In total, including all strains with *mtrC* frameshift mutations and

excluding revertants, we identified 185 isolates (3.79%) that encoded a LOF allele of *mtrC*.

Presence of the *mtrC* LOF mutation in isolates with known resistance markers was correlated

with significantly reduced MICs (Supplementary Figure 4), and inclusion of *mtrC* LOF mutations

in our linear model increased adjusted R^2 values (azithromycin: 0.681 to 0.739; ceftriaxone:

0.719 to 0.738; ciprofloxacin: 0.944 to 0.947), decreased the proportion of strains with

unexplained susceptibility (azithromycin: 5.49% to 3.09%; ceftriaxone: 1.51% to 1.15%;

ciprofloxacin: 2.19% to 1.80%), and significantly improved model fit (p < $2.2{\times}10^{-16}$ for all three

antibiotics; Likelihood-ratio χ^2 test for nested models). *mtrC* LOF strains were identified in 28 of

the 66 countries surveyed and ranged in isolation date from 1998 to 2017. Our results

demonstrate that a relatively common mechanism of gonococcal acquired antibiotic

susceptibility is a two base pair deletion in *mtrC* and that such mutations are globally and

temporally widespread.

Loss of the MtrCDE efflux pump is associated with cervical infection

The MtrCDE pump has been demonstrated to play a critical role in gonococcal survival in the

presence of human neutrophils and in the female murine genital tract model of gonococcal

infection, and overexpression of *mtrCDE* results in substantial fitness benefits for dealing with

both antimicrobial and environmental pressures[34-37]. The relative frequency of the *mtrC* LOF

mutations we observe (occurring in approximately 1 in every 25 isolates) thus seems unusual

for a mutation predicted to be deleterious for human infection. *mtrC* LOF strains do not grow

more or less quickly *in vitro* than *mtrC* wild-type strains, indicating that this mutation does not

confer a simple fitness benefit due to reduced energetic cost[31,34]. Instead, we hypothesized that

there are unique environments that select for non-functional efflux pump.

We aggregated patient-level metadata across included studies on sex partner preferences and

anatomical site of infection. Sex of sex partner and *mtrC* genotypic information was available for

1975 isolates from individual patients. There was a significant association between *mtrC* LOF

and sex of sex partners (p = 0.04021; Fisher's exact test) (Figure 3a), and *mtrC* LOF occurred

more often in isolates from men who have sex with women (MSW) (28/626, 4.47%) compared

to isolates from men who have sex with men (MSM) (31/1189, 2.61%) (OR = 1.75, 95% CI

[1.00-3.04], p = 0.037; Fisher's exact test). To understand whether anatomical selective

pressures contributed to this enrichment, we analyzed the site of infection and *mtrC* genotypic

information available for 2730 isolates. *mtrC* LOF mutations were significantly associated with

site of infection (p = 6.49×10^{-5}; Fisher's exact test) and were overrepresented particularly in

cervical isolates: 16 out of 129 (12.4%) cervical isolates contained an *mtrC* LOF mutation

compared to 82 out of 2249 urethral isolates (3.65%; OR = 3.74, 95% CI [1.98-6.70], p =

4.71×10^{-5}; Fisher's exact test), 3 out of 106 pharyngeal isolates (2.83%; OR = 4.83, 95% CI

[1.33-26.63], p = 0.00769; Fisher's exact test), and 4 out of 246 rectal isolates (1.63%; OR =

8.52, 95% CI [2.67-35.787], p = 2.39×10^{-5}; Fisher's exact test) (Figure 3b). Because our meta-

analysis collection comprises datasets potentially biased by preferential sampling for drug-

resistant strains, we validated our epidemiological associations on a set of 2186 sequenced

isolates, corresponding to all cultured isolates of *N. gonorrhoeae* in the state of Victoria,

Australia in 2017[38]. We again observed significant associations between *mtrC* LOF and sex of

sex partner (p = 0.0180; Fisher's exact test) as well as anatomical site of infection (p = 0.0256;

Fisher's exact test) (Supplementary Figure 5, Supplementary Text). *mtrC* LOF mutations were

again overrepresented in cervical isolates: 9 out of 227 (3.96%) cervical isolates contained an

mtrC LOF mutation compared to 15 out of 882 urethral isolates (1.70%; OR = 2.38, 95% CI

[0.91-5.91], p = 0.0679; Fisher's exact test), 3 out of 386 pharyngeal isolates (0.78%; OR =

5.26, 95% CI [1.29-30.51], p = 0.0117; Fisher's exact test), and 7 out of 632 rectal isolates

(1.11%; OR = 3.68, 95% CI [1.20-11.78], p = 0.0173; Fisher's exact test). These results indicate

that environmental pressures unique to female urogenital infection may select for loss of the

primary gonococcal efflux pump resulting in broadly increased susceptibility to antibiotics and

host-derived antimicrobial peptides.

Figure 3 – Gonococcal *mtrC*, *mtrA*, and *farA* LOF mutations are associated with patient sex of sex partner and site of infection in the global meta-analysis dataset. a) Sex of sex partner in patients infected with isolates with either intact or LOF alleles of *mtrC* (left), *mtrA* (middle), or *farA* (right). b) Site of infection in patients infected with isolates with either intact or LOF alleles of *mtrC* (left), *mtrA* (middle), or *farA* (right). *mtrA* alleles were predicted as LOF only in the absence of other epistatic Mtr overexpression mutations. Statistical significance between genotype and patient metadata was assessed by Fisher's exact test: * p<0.05, ** p<0.01, and *** p<0.001. WSM = women who have sex with men, MSW = men who have sex with women, MSMW = men who have sex with men and women, MSM = men who have sex with men.

Activator loss-of-function offers an additional level of adaptive regulatory control

The association of *mtrC* LOF mutations with cervical specimens suggests that other mutations that downregulate expression of the *mtrCDE* operon should also promote adaptation to the cervical niche. The MtrCDE efflux pump regulon comprises the MtrR repressor and the MtrA activator (Figure 2 inset), the latter of which exists in two allelic forms: a wild-type functional gene capable of inducing *mtrCDE* expression and a variant rendered non-functional by an 11-bp deletion near the 5' end of the gene[39] (Supplementary Figure 2B). Knocking out *mtrA* has a detrimental effect on fitness in the gonococcal mouse model, and epistatic *mtrR* mutations resulting in overexpression of *mtrCDE* compensate for this fitness defect by masking the effect of the *mtrA* knockout[35]. Prior work assessing the genomic diversity of *mtrA* in a set of 922 primarily male urethral specimens found only four isolates with the 11-bp deletion (0.43%)[40], seemingly in agreement with the *in vivo* importance of *mtrA*. However, in our global meta-analysis dataset, 362/4882 isolates (7.41%) were predicted to be *mtrA* LOF, of which the majority (357/362, 98.6%) were due to the 11-bp deletion. Of all 4882 isolates, 268 (5.49%) had *mtrA* LOF mutations in non-*mtrCDE* overexpression backgrounds (as defined by the absence of known *mtrR* promoter or coding sequence mutations or *mtrCDE* mosaic alleles) and therefore not epistatically masked. We repeated our epidemiological associations on these *mtrA* LOF strains without concurrent overexpression mutations and observed highly significant associations with reported patient sex of sex partner ($p = 1.81\times10^{-11}$; Fisher's exact test) and site of infection ($p = 1.64\times10^{-12}$; Fisher's exact test) (Figure 3). As with *mtrC* LOF mutations, *mtrA* LOF mutations were significantly overrepresented in cervical isolates: 25 out of 129 (19.4%) cervical isolates contained an *mtrA* LOF mutation compared to 61 out of 2248 urethral isolates (2.71%; OR = 8.60, 95% CI [4.96-14.57], $p = 4.60\times10^{-13}$; Fisher's exact test), 4 out of 106 pharyngeal isolates (3.78%; OR = 6.09, 95% CI [2.00-24.93], $p = 0.000240$; Fisher's exact test), and 4 out of 246 rectal isolates (1.63%; OR = 14.43, 95% CI [4.81-58.52], $p = 3.00\times10^{-9}$; Fisher's exact test). In the Australian validation cohort[38], the majority (81/85, 95.3%) of *mtrA* LOF strains had concurrent *mtrCDE* overexpression mutations, so it was not possible to test for these associations. In such genetic backgrounds where overexpression mutations mask the effect of *mtrA* LOF, *mtrC* LOF is the preferred method of efflux pump downregulation: the majority of *mtrC* LOF mutations in both the global dataset (174/180, 96.7%) and the Australian cohort (33/35, 94.3%) occurred in backgrounds with known *mtr* overexpression mutations. Phylogenetic analysis showed that the distribution of *mtrA* LOF differed from that of *mtrC* LOF with fewer introductions but more sustained transmission and that the two mutations were largely non-overlapping (Supplementary Figure 3). Our results indicate that multiple adaptive

paths for MtrCDE efflux pump downregulation exist depending on genetic interactions with other
concurrent mutations in the *mtrCDE* regulon.

A second class of proton-dependent efflux pumps appears to contribute to adaptation
via loss of function mutations
The associations we observed in the *mtrCDE* regulon raised the question of the mechanism by
which the cervical environment could select for pump downregulation. Recent work on
Pseudomonas suggested one possible model: overexpression of homologous *P. aeruginosa*
efflux pumps belonging to the same resistance/nodulation/cell division (RND) proton/substrate
antiporter family as MtrCDE results in a fitness cost due to increased cytoplasmic acidification[41].
This fitness cost was only observed in anaerobic conditions, where aerobic respiration cannot
be used to dissipate excess protons efficiently[41]. Analogous conditions in the female urogenital
tract, potentially augmented by environmental acidity, could create a similar selective pressure
during human infection that leads to pump downregulation or loss.

This model predicts that adaptation to these conditions would similarly result in downregulation
of FarAB, the other proton-substrate antiporter efflux pump in *N. gonorrhoeae*. FarAB is a
member of the major facilitator superfamily (MFS) of efflux pumps and effluxes long-chain fatty
acids[42,43]. In our global dataset, 332/4838 (6.86%) of isolates were predicted to have *farA* LOF
mutations, of which the majority (316/332; 95.2%) were due to indels in a homopolymeric
stretch of eight 'T' nucleotides near the 5' end of the gene (Supplementary Figure 2C). *farA* LOF
mutations were associated with patient sex of sex partner (p = $5.06×10^{-10}$; Fisher's exact test)
and site of infection (p = $1.76×10^{-12}$; Fisher's exact test) and overrepresented in cervical
isolates: 33 out of 129 (25.6%) cervical isolates contained a *farA* LOF mutation compared to
117 out of 2246 urethral isolates (5.21%; OR = 6.25, 95% CI [3.90-9.83], p = $3.24×10^{-13}$;
Fisher's exact test), 3 out of 106 pharyngeal isolates (2.83%; OR = 11.70, 95% CI [3.50-61.61],
p = $3.80×10^{-7}$; Fisher's exact test), and 12 out of 246 rectal isolates (4.88%; OR = 6.66, 95% CI
[3.19-14.80], p = $1.57×10^{-8}$; Fisher's exact test) (Figure 3). *farA* LOF mutations were prevalent
also in our Australian validation dataset[38] (225/2180; 10.32%) and again associated with sex of
sex partner (p < $2.20×10^{-16}$; Fisher's exact test) and site of infection (p < $2.20×10^{-16}$; Fisher's
exact test) (Supplementary Figure 5). The phylogenetic distribution of *farA* LOF mutations
overlapped with that of *mtrA* LOF mutations and indicated sustained transmission
(Supplementary Figure 3). We did not observe frequent LOF mutations in the sodium gradient-
dependent MATE family efflux pump NorM[44] or in the ATP-dependent ABC family pump

MacAB[45] (Supplementary Table 3). The prevalence and cervical enrichment of *farA* LOF

mutations and the relative rarity of LOF mutations in other non-proton motive force-driven

pumps suggests that cytoplasmic acidification may be a mechanism by which the female

urogenital tract selects for efflux pump loss.

Meningococcal evolution of efflux pump loss-of-function is driven by urogenital

adaptation

N. meningitidis, a species closely related to *N. gonorrhoeae,* colonizes the oropharyngeal tract

and can cause invasive disease, including meningitis and septicemia[46]. We characterized *mtrC*

diversity in a collection of 14,798 *N. meningitides* genomes, reasoning that the cervical

environmental pressures that select for efflux pump LOF in the gonococcus will be rarely

encountered by the meningococcus. In agreement with this, the 'GC' hexarepeat associated

with most gonococcal *mtrC* LOF mutations was less conserved in *N. meningitidis*; only

9684/14798 (65.4%) isolates contained an intact hexarepeat compared to 4644/4882 (95.1%) of

N. gonorrhoeae isolates (p < 2.2e-16; Fisher's exact test). In this same collection, we observed

mtrC LOF due to deletions in the hexarepeat region in only 82 meningococcal isolates (0.55%),

with a similar frequency of 25/4059 (0.62%) in a curated dataset comprising all invasive

meningococcal disease isolates collected in the UK from 2009-2013[47]. The observed

interruption of 'GC' dinucleotide repeats, predicted to result in a lower mutation rate[48], and the

relative rarity of *mtrC* LOF mutations suggests that efflux pump loss is not generally adaptive in

N. meningitidis. However, a urogenitally-adapted meningococcal lineage has recently emerged

in the US associated with outbreaks of non-gonococcal urethritis in heterosexual patients[49-51]. In

isolates from this lineage, the prevalence of *mtrC* LOF mutations was 18/207 (8.70%),

substantially higher than typical *N. meningitidis* and comparable to the prevalence of

gonococcal *mtrC* LOF mutations in MSW in our global dataset (4.47%). We compared the

frequency of *mtrC* LOF mutations in the urogenital lineage to geographically and genetically

matched isolates (i.e., all publicly available PubMLST ST-11 North American isolates) and

observed a significant difference in prevalence (18 out of 207 or 8.70% versus 2 out of 248 or

0.81%; OR = 10.74, 95% CI [2.52-96.56], p = 9.90×10^{-5}; Fisher's exact test). Most *mtrC* LOF

mutations occurred due to the same hexarepeat two base pair deletion that we previously

observed for *N. gonorrhoeae*, and in line with this, *mtrC* LOF in this urogenital lineage arose

multiple times independently similarly to gonococcal *mtrC* LOF mutations (Figure 4,

Supplementary Figure 3). *farA* LOF mutations were not observed in this meningococcal lineage.

We conclude that MtrCDE efflux pump LOF is rare in typical meningococcal strains that inhabit

the oropharynx but elevated in frequency in a unique urogenitally-adapted lineage circulating in

heterosexuals, indicative of potential ongoing adaptation to the cervical niche. Our results

suggest that efflux pump loss is broadly adaptive for cervical colonization across pathogenic

Neisseria.

Figure 4 – *mtrC* LOF mutations are enriched in a lineage of ST-11 urogenitally-adapted *N.*
meningitidis. A core-genome maximum likelihood phylogeny based on 25045 SNPs was
estimated of all North American ST-11 *N. meningitidis* strains from PubMLST (n=456; accessed
2019-09-03) rooted with meningococcal reference genome MC58. Membership in ST-11
urogenital clade (blue) defined as in Retchless et al., 2018[50]. Genomes with *mtrC* LOF
mutations are indicated in pink. Branch length represents substitutions per site.

Discussion

In an era in which widespread antimicrobial pressure has led to the emergence of extensively drug-resistant *N. gonorrhoeae*[52], isolates that appear to have reverted to susceptibility still arise[9,10], demonstrating that antibiotic and host environmental pressures interact to shape the evolution of *N. gonorrhoeae*. Here, we showed that frameshift-mediated truncations in the *mtrC* component of the MtrCDE efflux pump are the primary mechanism for epistatic increases in antibiotic susceptibility across a global collection of clinical gonococcal isolates, as suggested by prior work[30,31]. *mtrC* LOF mutations are enriched in cervical isolates and an inactive form of the pump activator MtrA exhibits similar trends, supporting a model in which reduced or eliminated *mtrCDE* efflux pump expression contributes to adaptation to the female genital tract. We hypothesized that the mechanism by which this occurs is through increased cytoplasmic acidification in anaerobic conditions[41] and demonstrated that LOF mutations in *farA*, encoding a subunit of the other proton motive force-driven pump FarAB, were likewise enriched in cervical isolates. *mtrC* and *farA* LOF mutations primarily occurred in short homopolymeric sequences and thus may occur via phase variation at a frequency higher than baseline mutation rate, similar to other resistance suppressor mutations[53]. In total, 36.4% of cervical isolates in the global dataset and 32.6% in the Australian dataset contained a LOF mutation in either *mtrC, farA,* or *mtrA*, indicating that efflux pump downregulation via multiple genetic mechanisms is prevalent in cervical infection. Despite significant associations only a proportion of cervical isolates exhibited these LOF genotypes, suggesting variation in cervix-associated pressures or indicating that cervical culture specimens were obtained before niche pressures could select for pump downregulation. Furthermore, while the cervix is the primary site of infection and source for culture in women, the selective pressures at play may include other sites more broadly in the female genital tract and may be influenced by the presence of other microbial species both pathogenic and commensal. While antibiotic use for gonorrhea in men and women may differ due to variation in rates of symptomatic infections, differences in antibiotic use could not explain the associations we observed for the non-antibiotic substrate efflux pump *farAB* or the increased frequency of *mtrC* LOF mutations in urogenitally-adapted meningococci.

Our model extended to the other pathogenic *Neisseria* species, *N. meningitidis*, in that a urogenital clade transmitting in primarily heterosexual populations appeared to be undergoing further urogenital adaptation via the same *mtrC* frameshift mutation that was most commonly

observed for *N. gonorrhoeae*. In the absence of data on cases of cervicitis, we hypothesized

that for this meningococcal lineage, efflux pump LOF emerged in the female urogenital tract

and was transmitted to heterosexual men resulting in the enrichment we observed. Efflux

pumps are common across Gram-negative bacteria[54], and their loss may be a general

adaptive strategy for species that face similar pressures as *N. gonorrhoeae* and urogenitally-

adapted *N. meningitidis*. In support of this, clinical isolates of *Pseudomonas aeruginosa* with

truncations in genes homologous to *mtrC*[55,56] and antibiotic hypersensitivity have been

obtained from cystic fibrosis patients, in whom the thick mucus in airway environments can

exhibit increased acidity and decreased oxygen availability[57,58].

Our results also suggest potential therapeutic avenues for addressing the emergence of

multidrug-resistant gonococcal strains. Selective knockdown of MtrCDE homologs in other

bacteria via antisense RNA[59] and bacteriophages[60] has successfully re-sensitized resistant

strains and enhanced antibiotic efficacy, and ectopic expression in *N. gonorrhoeae* of the *mtrR*

repressor in a cephalosporin-resistant strain enhances gonococcal killing by β-lactam

antibiotics in the mouse model[61]. Our population-wide estimated effect sizes for *mtrC* LOF

mutations provide a prediction for the re-sensitization effect of MtrCDE knockdown across

multiple genetic backgrounds and suggest particularly strong effects for the macrolide

azithromycin (Supplementary Figure 4).

In summary, by analysis of population genomics and patient clinical data, we have shown that

pathogenic *Neisseria* can use multiple avenues of efflux pump perturbation as an adaptive

strategy to respond to host environmental pressures and illustrate how these host pressures

may result in increased antibiotic susceptibility in *N. gonorrhoeae*.

Methods

Genomics pipeline: Reads for isolates with either associated azithromycin, ciprofloxacin, or ceftriaxone MIC metadata were downloaded from datasets listed in Supplementary Table 1. Reads were inspected using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and removed if GC content diverged notably from expected values (approximately 52-54%) or if base quality was systematically poor. We mapped read data to the NCCP11945 reference genome (RefSeq accession: NC_011035.1) using BWA-MEM (version 0.7.17-r1188)[62,63] and deduplicated reads using Picard (version 2.8.0) (https://github.com/broadinstitute/picard). BamQC in Qualimap (version 2.2.1)[64] was run to identify samples with less than 70% of reads aligned or samples with less than 40X coverage, which were discarded. We used Pilon (version 1.16)[65] to call variants with mindepth set to 10 and minmq set to 20 and generated pseudogenomes from Pilon VCFs by including all PASS sites and alternate alleles with AF > 0.9; all other sites were assigned as 'N'. Samples with greater than 15% of sites across the genome missing were also excluded. We created *de novo a*ssemblies using SPAdes (version 3.12.0)[66] and quality filtered contigs to ensure coverage greater than 10X, length greater than 500 base pairs, and total genome size approximately equal to the FA1090 genome size (2.0 to 2.3 Mbp). We annotated assemblies with Prokka (version 1.13)[67], and clustered core genes using Roary (version 3.12)[68] and core intergenic regions using piggy (version 1.2)[69]. A recombination-corrected phylogeny of all isolates was constructed by running Gubbins (version 2.3.4) on the aligned pseudogenomes and visualized in iTOL (version 4.4.2)[70-72]. All isolates with associated metadata and accession numbers are listed in Supplementary Tables 5 and 6.

Resistance allele calling: Known resistance determinants in single-copy genes were called by identifying expected SNPs in the pseudogenomes. For categorizing mosaic alleles of *mtr*, we ran BLASTn (version 2.6.0)[73] on the *de novo* assemblies using a query sequence from FA1090 (Genbank accession: NC_002946.2) comprising the *mtr* intergenic promoter region and *mtrCDE*. BLAST results were aligned using MAFFT (version 7.450)[74] and clustered into distinct allelic families using FastBAPS (version 1.0.0)[75]. We confirmed that horizontally-transferred *mtr* alleles associated with resistance from prior studies[5] corresponded to distinct clusters in FastBAPS. A similar approach was used to cluster *penA* alleles after running BLASTn with a *penA* reference sequence from FA1090. Variant calling in the multi-copy 23S

rRNA locus was done by mapping to a modified NCCP11945 reference genome containing

only one copy of the 23S rRNA and analyzing variant allele frequencies[76]. We identified

truncated MtrR proteins using Prokka annotations, and mutations in the *mtr* promoter region

associated with upregulation of *mtrCDE* (A deletion and TT insertion in inverted repeat, *mtr*

120) using an alignment of the *mtr* promoter from piggy output.

Phenotype processing and linear models: We doubled GISP azithromycin MICs before

2005 to account for the GISP MIC protocol testing change[77]. Samples with binary resistance

phenotypes (i.e., "SUS" and "RES") were discarded. For samples with MICs listed as above or

below a threshold (indicated by greater than or less than symbols), the MIC was set to equal

the provided threshold. MICs were log2-transformed for use as continuous outcome variables

in linear modeling and GWAS. We modeled transformed MICs using a panel of known

resistance markers[22] and included the recently characterized mosaic *mtrCDE* alleles conferring

azithromycin resistance[5], as well as genetic pairwise interactions and isolate country of origin.

Formulas called by the lm function in R (version 3.5.1) for each drug were (with codon or

nucleotide site indicated after each gene or rRNA, respectively):

Azithromycin: Country + (MtrR 39 + MtrR 45 + MtrR truncated + Mtr promoter + Mtr BAPS +

23S rRNA 2059 + 23S rRNA 2611)^2

Ceftriaxone: Country + (MtrR 39 + MtrR 45 + MtrR truncated + Mtr promoter + PenA BAPS +

PBP1 421 + PBP2 501 + PBP2 542 + PBP2 551 + PorB 120 + PorB 121)^2

Ciprofloxacin: Country + (MtrR 39 + MtrR 45 + MtrR truncated + Mtr promoter + GyrA 91 +

GyrA 95 + ParC 86 + ParC 87 + ParC 91 + PorB 120 + PorB 121)^2

To visualize the continuous MICs using thresholds as on Figure 1, we binarized MICs using

the CLSI resistance breakpoint for ciprofloxacin, the CLSI non-susceptibility breakpoint for

azithromycin, and the CDC GISP surveillance breakpoint for ceftriaxone.

GWAS and unitig annotation: Linear mixed model GWAS was run using Pyseer (version

1.2.0) with unitigs generated from GATB (version 1.3.0). To create the Manhattan plot, we

mapped all unitigs from the GWAS using BWA-MEM (modified parameters: -B 2 and -O 3) to

the pan-susceptible WHO F strain reference genome (Genbank accession:

GCA_900087635.2) edited to contain only one locus of the 23S rRNA. Significant unitigs were

annotated using Pyseer's annotation pipeline. Unitigs mapping to multiple sites in the genome
and in or near the highly variable *pilE* (encoding pilus subunit) or *piiC* (encoding opacity protein
family) genes were excluded, as were unitigs less than twenty base pairs in length. Due to
redundancy and linkage, variants will be spanned by multiple overlapping unitigs with similar
frequencies and p-values. For ease of interpretation, we grouped unitigs within 50 base pairs
of each other and represented each cluster by the most significant unitig. Unitigs with allele
frequency greater than 50% were also excluded as they represented the majority allele. Unitig
clusters were then annotated by gene or adjacent genes for unitigs mapping to intergenic
regions and further analyzed for predicted functional effect relative to the WHO F reference
genome in Geneious Prime (version 2019.2.1).

Identifying LOF and upregulation alleles: To identify predicted LOF alleles of efflux pump
proteins, we ran BLASTn on the *de novo* assemblies using a query sequence from FA1090
(reference genome FA19 was used for *mtrA*). Sequences that were full-length or
approximately full-length (+/- 5 nucleotides) beginning with expected start codons were
translated using Biopython[78]. Peptides shorter than 90% of the expected full-length size of the
protein were further analyzed in Geneious Prime to identify the nucleotide mutations resulting
in predicted LOF. We called *mtrCDE* overexpression status by identifying the presence of any
of the known *mtrR* promoter mutations, MtrR coding sequence mutations, and mosaic *mtrCDE*
alleles.

Experimental validation: *N. gonorrhoeae* culture was conducted on GCB agar (Difco) plates
supplemented with 1% Kellogg's supplements[79] at 37°C in a 5% CO2 atmosphere.
Antimicrobial susceptibility testing was conducted on GCB agar supplemented with 1%
IsoVitaleX (Becton Dickinson) using Etests (bioMérieux) at 37°C in a 5% CO2 atmosphere. We
selected a clinical isolate (NY0195) from the multidrug-resistant lineage ST-1901[32] that
contained an *mtrC* LOF mutation mediated by a two base pair hexarepeat deletion and
confirmed via Etests that its MIC matched, within one dilution, its reported MIC. Isolate
NY0195 contained mosaic *penA* allele XXXIV conferring cephalosporin reduced susceptibility
and the *gyrA* S91F substitution conferring ciprofloxacin resistance[9]. We complemented the
mtrC LOF mutation in this strain by transforming it via electroporation[79] with a 2kb PCR
product containing a *Neisserial* DNA uptake sequence and an in-frame *mtrC* allele, obtained
by colony PCR from a neighboring isolate (GCGS0759). After obtaining transformants by

selecting on an azithromycin 0.05 µg/mL GCB plate supplemented with Kellogg's supplement,

we confirmed successful transformation by Sanger sequencing of the *mtrC* gene. No

spontaneous mutants on azithromycin 0.05 µg/mL plates were observed after conducting

control transformations in the absence of GCGS0759 *mtrC* PCR product. We conducted

antimicrobial susceptibility testing in triplicate using Etests, assessing statistical significance

between parental and transformant MICs by a two-sample t-test.

Metadata analysis: Patient metadata were collected from supplementary tables where

available and values standardized across datasets using a common ontology (i.e., for site of

infection, classification as urethra, rectum, pharynx, or cervix). Fisher's exact test in R (version

3.5.1) was used to infer whether there was nonrandom association between *mtrC* LOF

presence and either anatomical site of infection or sex of sex partner. For sex of sex partner

analysis, isolates cultured from multiple sites on the same patient were counted as only one

data point.

Meningococcal *mtrC* analysis: *mtrC* alleles from *N. meningitidis* assembled genomes were

downloaded from PubMLST (n=14798; accessed 2019-09-03) by setting (species = "Neisseria

meningitidis"), filtering by (Sequence bin size >= 2 Mbp), and exporting sequences for Locus

"NEIS1634"[80]. *mtrC* LOF alleles were identified as described above. We generated a core-

genome maximum likelihood phylogeny of all North American ST-11 *N. meningitidis* strains

from PubMLST (n=456; accessed 2019-09-03) rooted with meningococcal reference genome

MC58 (Genbank accession: AE002098.2) using Roary (version 3.12) and annotated it using

metadata from Retchless et al., 2018[50]. Overrepresentation of *mtrC* LOF alleles in the US

urogenital lineage compared to selected control datasets was assessed using Fisher's exact

test in R (version 3.5.1).

Acknowledgements

This work was supported by the NIH/NIAID grant 1R01AI132606-01 and the Smith Family

Foundation. TDM is additionally supported by the NIH/NIAID 1 F32 AI145157-01, and KCM is

additionally supported by the NSF GRFP. DHFR was supported by award Number

T32GM007753 from the National Institute of General Medical Sciences. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the

National Institute of General Medical Sciences or the National Institutes of Health. DAW is

supported by an Early Career Fellowship from the National Health and Medical Research

Council of Australia (GNT1123854). Portions of this research were conducted on the O2 high-

performance computing cluster, supported by the Research Computing Group at Harvard

Medical School. This publication made use of the Meningitis Research Foundation

Meningococcus Genome Library (http://www.meningitis.org/research/genome) developed by

Public Health England, the Wellcome Trust Sanger Institute, and the University of Oxford as a

collaboration and funded by the Meningitis Research Foundation. The authors additionally thank

Crista Wadsworth, Samantha Palace, and other members of the Grad Lab for helpful comments

during development of the project.

Publication	Study summary	Timespan	Included number
Mortimer et al., 2020	Transmission and AMR surveillance in New York City, USA	2011-2015	888
Sánchez-Busó et al., 2019[3]	Worldwide phylogeography and evolution of gonococcus	1979-2012	408
Yahara et al., 2018[10]	AMR surveillance in Kyoto and Osaka, Japan	1996-2015	260
Ryan et al., 2018[11]	AMR surveillance in Ireland	2012-2016	39
Harris et al., 2018[12]	Genomic survey across European Euro-GASP participant countries (n=20)	2013	1048
Fifer et al., 2018[13]	High-level azithromycin resistance outbreak in UK	2004-2017	50
Lee et al., 2018[14]	Genomic epidemiology in New Zealand	2014-2015	397
Kwong et al., 2017[15]	Transmission among MSM in Melbourne, Australia	2005-2014	94
Eyre et al., 2017[22] and De Silva et al., 2016[20]	Transmission in Brighton, UK	2004-2011	231
Grad et al., 2016[9] and 2014[17]	AMR surveillance across CDC GISP clinics, USA	2000-2013	1100
Demczuk et al., 2016[19]	Azithromycin resistance surveillance in Canada	1991-2014	199
Demczuk et al., 2015[18]	Cephalosporin decreased susceptibility surveillance in Canada	1989-2013	114
Ezewudo et al., 2015[16]	Population structure and AMR surveillance	1982-2011	54

546 **Supplementary Table 1 – Datasets included in global meta-analysis collection.** All isolates
547 that passed genomics quality control filters (see methods) with associated azithromycin, or
548 ceftriaxone, or ciprofloxacin metadata were included (n=4882 in total). Euro-GASP = European
549 Gonococcal Antimicrobial Surveillance Program; CDC GISP = Centers for Disease Control and
550 Prevention Gonococcal Isolate Surveillance Project.

Strain	Azithromycin	Ceftriaxone	Ciprofloxacin
NY0195	0.064, 0.094, 0.094	0.016, 0.023, 0.023	6, 6, 4
NY0195 *mtrC*(in-frame)	0.5, 0.75, 0.75	0.064, 0.064, 0.064	≥ 32, ≥ 32, ≥ 32
	p = 0.0187	p = 0.00289	p = 0.000624

553 **Supplementary Table 2 – Restoration of the *mtrC* coding frame in a clinical isolate by**
554 **transformation increases MICs to azithromycin, ceftriaxone, and ciprofloxacin.** MIC Etests
555 were conducted in triplicate with all results reported. Statistical significance for MIC differences
556 between parental strain and transformant strain was assessed by a two-sample t-test after
557 setting the value of "≥ 32" to 32.

Dataset	Gene	N LOF	N Total	Percentage
Global	*farA*	332	4838	6.86%
Global	*farB*	2	4850	0.04%
Global	*norM*	2	4852	0.04%
Global	*macA*	1	4847	0.02%
Global	*macB*	13	4845	0.27%
Global	*mtrR*	386	4845	7.97%
Global	*mtrA*	268 (362)	4842	5.45% (7.48%)
Global	*mtrC*	185	4847	3.82%
Global	*mtrD*	10	4815	0.21%
Global	*mtrE*	0	4849	0.00%
Australia	*farA*	225	2180	10.32%
Australia	*farB*	1	2186	0.05%
Australia	*norM*	0	2186	0.00%
Australia	*macA*	0	2186	0.00%
Australia	*macB*	0	2186	0.00%
Australia	*mtrA*	4 (85)	2186	0.18% (3.89%)
Australia	*mtrR*	253	2183	11.59%
Australia	*mtrC*	35	2186	1.60%
Australia	*mtrD*	2	2185	0.09%
Australia	*mtrE*	0	2186	0.00%

565 **Supplementary Table 3 – Prevalence of predicted LOF mutations in efflux pump genes**
566 **for both global and Australian datasets.** Counts for *mtrA* LOF mutations in the absence of
567 other epistatic *mtrCDE* overexpression mutations are listed first, with counts for total number of
568 *mtrA* LOF mutations regardless of *mtrCDE* overexpression status listed in parentheses.

569
570
571
572 **Supplementary Figure 1** – **Q-Q plots of expected versus observed p-values for GWAS on**
573 **a) azithromycin, b) ceftriaxone, and c) ciprofloxacin.** Under the null hypothesis of no causal
574 variants and in the absence of confounders such as population structure, p-values are
575 distributed uniformly and would be expected to lie along the red y=x line.
576

a) Mutational diversity in *mtrC*

Wild-type *mtrC*
'GC' deletion in hexarepeat region
'GC' deletion reverted via insertion elsewhere in gene
'A' insertion leading to nonsense mutation
Mosaic *mtrC* with 'GC' deletion

b) Mutational diversity in *mtrA*

Wild-type *mtrA*
11 base pair deletion near 5' end

c) Mutational diversity in *farA*

Wild-type *farA*
Nonsense mutation
'T' deletion in homopolymeric tract
'T' insertion in homopolymeric tract
'G' deletion near homopolymeric tract

577
578
579 **Supplementary Figure 2 – Alignment of various LOF mutations observed in a)** *mtrC*, **b)**
580 *mtrA*, **and c)** *farA* **in the global dataset.** Alignments were visualized in Geneious. Dots
581 indicate nucleotide or amino acid consensus sequence.
582
583

584
585
Supplementary Figure 3 – **Phylogenetic distribution of gonococcal *mtrC*, MtrA, and *farA***
LOF alleles with patient site of infection (n=2742) in global dataset. The recombination-
corrected maximum likelihood phylogeny based on 36347 SNPs is shown annotated with rings
(from innermost to outermost) for site of infection, *mtrCDE* regulon LOF mutations, and *farA*
LOF mutations. Branch length represents total number of substitutions after removal of
predicted recombination.
592

593
594
595 **Supplementary Figure 4 – MIC distributions for isolates with indicated resistance**
596 **determinants for azithromycin, ceftriaxone, and ciprofloxacin respectively, stratified by**
597 *mtrC* **genotypic status.** Statistical significance between *mtrC* intact and LOF MIC distributions
598 was assessed by Mann-Whitney U Test: * p<0.05, ** p<0.01, and *** p<0.001.
599

Supplementary Figure 5 – Gonococcal *mtrC* and *farA* LOF mutations are associated with
sex of sex partner and site of infection in the validation dataset. a) Sex partner information
in patients infected with isolates with either intact or LOF alleles of *mtrC* (left) or *farA* (right).
MSMW patients were labelled as MSM in the validation dataset[38]. b) Site of infection in patients
infected with isolates with either intact or LOF alleles of *mtrC* (left) or *farA* (right) datasets.
Statistical significance between intact versus LOF patient metadata distributions was assessed
by Fisher's exact test: * p<0.05, ** p<0.01, and *** p<0.001. WSM = women who have sex with
men, MSW = men who have sex with women, MSMW = men who have sex with men and
women, MSM = men who have sex with men.

616 **Supplementary Figure 6 – Phylogenetic distribution of gonococcal *mtrC* LOF alleles with**
617 **patient site of infection (n=2186) in the validation dataset.** The recombination-corrected
618 maximum likelihood phylogeny based on 26669 SNPs is shown annotated with rings (from
619 innermost to outermost) for site of infection, *mtrC* LOF mutations, and *farA* LOF mutations.
620 Branch length represents total number of substitutions after removal of predicted recombination.

References

References

1   Olesen, S. W. *et al.* Azithromycin susceptibility in Neisseria gonorrhoeae and seasonal
macrolide use. *J Infect Dis*, doi:10.1093/infdis/jiy551 (2018).
2   Olesen, S. W. & Grad, Y. H. Deciphering the impact of bystander selection for antibiotic
resistance in Neisseria gonorrhoeae. *J Infect Dis*, doi:10.1093/infdis/jiz156 (2019).
3   Sanchez-Buso, L. *et al.* The impact of antimicrobials on gonococcal evolution. *Nat*
Microbiol, doi:10.1038/s41564-019-0501-y (2019).
4   Unemo, M. & Shafer, W. M. Antimicrobial resistance in Neisseria gonorrhoeae in the
21st century: past, evolution, and future. *Clin Microbiol Rev* **27**, 587-613,
doi:10.1128/CMR.00010-14 (2014).
5   Wadsworth, C. B., Arnold, B. J., Sater, M. R. A. & Grad, Y. H. Azithromycin Resistance
through Interspecific Acquisition of an Epistasis-Dependent Efflux Pump Component and
Transcriptional Regulator in Neisseria gonorrhoeae. *MBio* **9**, doi:10.1128/mBio.01419-18
(2018).
6   Rouquette-Loughlin, C. E. *et al.* Mechanistic Basis for Decreased Antimicrobial
Susceptibility in a Clinical Isolate of Neisseria gonorrhoeae Possessing a Mosaic-Like
mtr Efflux Pump Locus. *MBio* **9**, doi:10.1128/mBio.02281-18 (2018).
7   Quillin, S. J. & Seifert, H. S. Neisseria gonorrhoeae host adaptation and pathogenesis.
Nat Rev Microbiol **16**, 226-240, doi:10.1038/nrmicro.2017.169 (2018).
8   Wi, T. *et al.* Antimicrobial resistance in Neisseria gonorrhoeae: Global surveillance and a
call for international collaborative action. *PLoS Med* **14**, e1002344,
doi:10.1371/journal.pmed.1002344 (2017).
9   Grad, Y. H. *et al.* Genomic Epidemiology of Gonococcal Resistance to Extended-
Spectrum Cephalosporins, Macrolides, and Fluoroquinolones in the United States, 2000-
2013. *J Infect Dis* **214**, 1579-1587, doi:10.1093/infdis/jiw420 (2016).
10   Yahara, K. *et al.* Genomic surveillance of Neisseria gonorrhoeae to investigate the
distribution and evolution of antimicrobial-resistance determinants and lineages. *Microb*
Genom **4**, doi:10.1099/mgen.0.000205 (2018).
11   Ryan, L. *et al.* Antimicrobial resistance and molecular epidemiology using whole-genome
sequencing of Neisseria gonorrhoeae in Ireland, 2014-2016: focus on extended-
spectrum cephalosporins and azithromycin. *Eur J Clin Microbiol Infect Dis*,
doi:10.1007/s10096-018-3296-5 (2018).
12   Harris, S. R. *et al.* Public health surveillance of multidrug-resistant clones of Neisseria
gonorrhoeae in Europe: a genomic survey. *Lancet Infect Dis* **18**, 758-768,
doi:10.1016/S1473-3099(18)30225-1 (2018).
13   Fifer, H. *et al.* Sustained transmission of high-level azithromycin-resistant Neisseria
gonorrhoeae in England: an observational study. *Lancet Infect Dis* **18**, 573-581,
doi:10.1016/S1473-3099(18)30122-1 (2018).
14   Lee, R. S. *et al.* Genomic epidemiology and antimicrobial resistance of Neisseria
gonorrhoeae in New Zealand. *J Antimicrob Chemother* **73**, 353-364,
doi:10.1093/jac/dkx405 (2018).
15   Kwong, J. C. *et al.* Whole-genome sequencing reveals transmission of gonococcal
antibiotic resistance among men who have sex with men: an observational study. *Sex*
Transm Infect **94**, 151-157, doi:10.1136/sextrans-2017-053287 (2018).
16   Ezewudo, M. N. *et al.* Population structure of Neisseria gonorrhoeae based on whole
genome data and its relationship with antibiotic resistance. *PeerJ* **3**, e806,
doi:10.7717/peerj.806 (2015).
17   Grad, Y. H. *et al.* Genomic epidemiology of Neisseria gonorrhoeae with reduced
susceptibility to cefixime in the USA: a retrospective observational study. *Lancet Infect*
Dis **14**, 220-226, doi:10.1016/S1473-3099(13)70693-5 (2014).

18   Demczuk, W. *et al.* Whole-genome phylogenomic heterogeneity of Neisseria
676   gonorrhoeae isolates with decreased cephalosporin susceptibility collected in Canada
677   between 1989 and 2013. *J Clin Microbiol* **53**, 191-200, doi:10.1128/JCM.02589-14
678   (2015).
679   19   Demczuk, W. *et al.* Genomic Epidemiology and Molecular Resistance Mechanisms of
680   Azithromycin-Resistant Neisseria gonorrhoeae in Canada from 1997 to 2014. *J Clin*
681   *Microbiol* **54**, 1304-1313, doi:10.1128/JCM.03195-15 (2016).
682   20   De Silva, D. *et al.* Whole-genome sequencing to determine transmission of Neisseria
683   gonorrhoeae: an observational study. *Lancet Infect Dis* **16**, 1295-1303,
684   doi:10.1016/S1473-3099(16)30157-8 (2016).
685   21   Wadsworth, C. B., Arnold, B. J., Abdul Sater, M. R. & Grad, Y. H. Azithromycin
686   resistance through interspecific acquisition of an epistasis dependent efflux pump
687   component and transcriptional regulator in Neisseria gonorrhoeae. . *BioRxiv* (2018).
688   22   Eyre, D. W. *et al.* WGS to predict antibiotic MICs for Neisseria gonorrhoeae. *J*
689   *Antimicrob Chemother* **72**, 1937-1947, doi:10.1093/jac/dkx067 (2017).
690   23   Hicks, A. L. *et al.* Evaluation of parameters affecting performance and reliability of
691   machine learning-based antibiotic susceptibility testing from whole genome sequencing
692   data. *PLoS Comput Biol* **15**, e1007349, doi:10.1371/journal.pcbi.1007349 (2019).
693   24   Earle, S. G. *et al.* Identifying lineage effects when controlling for population structure
694   improves power in bacterial association studies. *Nat Microbiol* **1**, 16041,
695   doi:10.1038/nmicrobiol.2016.41 (2016).
696   25   Lees, J. A., Galardini, M., Bentley, S. D., Weiser, J. N. & Corander, J. pyseer: a
697   comprehensive tool for microbial pangenome-wide association studies. *Bioinformatics*,
698   bty539-bty539, doi:10.1093/bioinformatics/bty539 (2018).
699   26   Jaillard, M. *et al.* A fast and agnostic method for bacterial genome-wide association
700   studies: Bridging the gap between k-mers and genetic events. *PLoS Genet* **14**,
701   e1007758, doi:10.1371/journal.pgen.1007758 (2018).
702   27   Mortimer, T. D. & Grad, Y. H. Applications of genomics to slow the spread of multidrug-
703   resistant Neisseria gonorrhoeae. *Ann N Y Acad Sci*, doi:10.1111/nyas.13871 (2018).
704   28   Veal, W. L., Nicholas, R. A. & Shafer, W. M. Overexpression of the MtrC-MtrD-MtrE
705   efflux pump due to an mtrR mutation is required for chromosomally mediated penicillin
706   resistance in Neisseria gonorrhoeae. *J Bacteriol* **184**, 5619-5624,
707   doi:10.1128/jb.184.20.5619-5624.2002 (2002).
708   29   Golparian, D., Shafer, W. M., Ohnishi, M. & Unemo, M. Importance of multidrug efflux
709   pumps in the antimicrobial resistance property of clinical multidrug-resistant isolates of
710   Neisseria gonorrhoeae. *Antimicrob Agents Chemother* **58**, 3556-3559,
711   doi:10.1128/AAC.00038-14 (2014).
712   30   Veal, W. L. *et al.* Loss-of-function mutations in the mtr efflux system of Neisseria
713   gonorrhoeae. *Microbiology* **144 (Pt 3)**, 621-627, doi:10.1099/00221287-144-3-621
714   (1998).
715   31   Eisenstein, B. I. & Sparling, P. F. Mutations to increased antibiotic sensitivity in naturally-
716   occurring gonococci. *Nature* **271**, 242-244, doi:10.1038/271242a0 (1978).
717   32   Shimuta, K. *et al.* Emergence and evolution of internationally disseminated
718   cephalosporin-resistant Neisseria gonorrhoeae clones from 1995 to 2005 in Japan. *BMC*
719   *Infect Dis* **15**, 378, doi:10.1186/s12879-015-1110-x (2015).
720   33   Bichara, M., Pinet, I., Schumacher, S. & Fuchs, R. P. Mechanisms of dinucleotide repeat
721   instability in Escherichia coli. *Genetics* **154**, 533-542 (2000).
722   34   Handing, J. W., Ragland, S. A., Bharathan, U. V. & Criss, A. K. The MtrCDE Efflux Pump
723   Contributes to Survival of Neisseria gonorrhoeae From Human Neutrophils and Their
724   Antimicrobial Components. *Front Microbiol* **9**, 2688, doi:10.3389/fmicb.2018.02688
725   (2018).

35 Warner, D. M., Folster, J. P., Shafer, W. M. & Jerse, A. E. Regulation of the MtrC-MtrD-
MtrE efflux-pump system modulates the in vivo fitness of Neisseria gonorrhoeae. *J Infect
Dis* **196**, 1804-1812, doi:10.1086/522964 (2007).
36 Warner, D. M., Shafer, W. M. & Jerse, A. E. Clinically relevant mutations that cause
derepression of the Neisseria gonorrhoeae MtrC-MtrD-MtrE Efflux pump system confer
different levels of antimicrobial resistance and in vivo fitness. *Mol Microbiol* **70**, 462-478,
doi:10.1111/j.1365-2958.2008.06424.x (2008).
37 Folster, J. P. *et al.* MtrR modulates rpoH expression and levels of antimicrobial
resistance in Neisseria gonorrhoeae. *J Bacteriol* **191**, 287-297, doi:10.1128/JB.01165-08
(2009).
38 Williamson, D. A. *et al.* Bridging of Neisseria gonorrhoeae lineages across sexual
networks in the HIV pre-exposure prophylaxis era. *Nat Commun* **10**, 3988,
doi:10.1038/s41467-019-12053-4 (2019).
39 Rouquette, C., Harmon, J. B. & Shafer, W. M. Induction of the mtrCDE-encoded efflux
pump system of Neisseria gonorrhoeae requires MtrA, an AraC-like protein. *Mol
Microbiol* **33**, 651-658, doi:10.1046/j.1365-2958.1999.01517.x (1999).
40 Vidyaprakash, E., Abrams, A. J., Shafer, W. M. & Trees, D. L. Whole-Genome
Sequencing of a Large Panel of Contemporary Neisseria gonorrhoeae Clinical Isolates
Indicates that a Wild-Type mtrA Gene Is Common: Implications for Inducible
Antimicrobial Resistance. *Antimicrob Agents Chemother* **61**, doi:10.1128/AAC.00262-17
(2017).
41 Olivares Pacheco, J., Alvarez-Ortega, C., Alcalde Rico, M. & Martinez, J. L. Metabolic
Compensation of Fitness Costs Is a General Outcome for Antibiotic-Resistant
Pseudomonas aeruginosa Mutants Overexpressing Efflux Pumps. *MBio* **8**,
doi:10.1128/mBio.00500-17 (2017).
42 Zgurskaya, H. I. Multicomponent drug efflux complexes: architecture and mechanism of
assembly. *Future Microbiol* **4**, 919-932, doi:10.2217/fmb.09.62 (2009).
43 Lee, E. H. & Shafer, W. M. The farAB-encoded efflux pump mediates resistance of
gonococci to long-chained antibacterial fatty acids. *Mol Microbiol* **33**, 839-845,
doi:10.1046/j.1365-2958.1999.01530.x (1999).
44 Rouquette-Loughlin, C., Dunham, S. A., Kuhn, M., Balthazar, J. T. & Shafer, W. M. The
NorM efflux pump of Neisseria gonorrhoeae and Neisseria meningitidis recognizes
antimicrobial cationic compounds. *J Bacteriol* **185**, 1101-1106,
doi:10.1128/jb.185.3.1101-1106.2003 (2003).
45 Rouquette-Loughlin, C. E., Balthazar, J. T. & Shafer, W. M. Characterization of the
MacA-MacB efflux system in Neisseria gonorrhoeae. *J Antimicrob Chemother* **56**, 856-
860, doi:10.1093/jac/dki333 (2005).
46 Stephens, D. S., Greenwood, B. & Brandtzaeg, P. Epidemic meningitis,
meningococcaemia, and Neisseria meningitidis. *Lancet* **369**, 2196-2210,
doi:10.1016/S0140-6736(07)61016-2 (2007).
47 Hill, D. M. *et al.* Genomic epidemiology of age-associated meningococcal lineages in
national surveillance: an observational cohort study. *Lancet Infect Dis* **15**, 1420-1428,
doi:10.1016/S1473-3099(15)00267-4 (2015).
48 Bichara, M., Schumacher, S. & Fuchs, R. P. Genetic instability within monotonous runs
of CpG sequences in Escherichia coli. *Genetics* **140**, 897-907 (1995).
49 Bazan, J. A. *et al.* Notes from the Field: Increase in Neisseria meningitidis-Associated
Urethritis Among Men at Two Sentinel Clinics - Columbus, Ohio, and Oakland County,
Michigan, 2015. *MMWR Morb Mortal Wkly Rep* **65**, 550-552,
doi:10.15585/mmwr.mm6521a5 (2016).

50 Retchless, A. C. *et al.* Expansion of a urethritis-associated Neisseria meningitidis clade
in the United States with concurrent acquisition of N. gonorrhoeae alleles. *BMC*
Genomics **19**, 176, doi:10.1186/s12864-018-4560-x (2018).
51 Tzeng, Y. L. *et al.* Emergence of a new Neisseria meningitidis clonal complex 11 lineage
11.2 clade as an effective urogenital pathogen. *Proc Natl Acad Sci U S A* **114**, 4237-
4242, doi:10.1073/pnas.1620971114 (2017).
52 Whiley, D. M., Jennison, A., Pearson, J. & Lahra, M. M. Genetic characterisation of
Neisseria gonorrhoeae resistant to both ceftriaxone and azithromycin. *Lancet Infect Dis*
18, 717-718, doi:10.1016/S1473-3099(18)30340-2 (2018).
53 Kime, L. *et al.* Transient Silencing of Antibiotic Resistance by Mutation Represents a
Significant Potential Source of Unanticipated Therapeutic Failure. *MBio* **10**,
doi:10.1128/mBio.01755-19 (2019).
54 Du, D. *et al.* Multidrug efflux pumps: structure, function and regulation. *Nat Rev Microbiol*
16, 523-539, doi:10.1038/s41579-018-0048-6 (2018).
55 Vettoretti, L. *et al.* Efflux unbalance in Pseudomonas aeruginosa isolates from cystic
fibrosis patients. *Antimicrob Agents Chemother* **53**, 1987-1997, doi:10.1128/AAC.01024-
08 (2009).
56 Chalhoub, H. *et al.* Mechanisms of intrinsic resistance and acquired susceptibility of
Pseudomonas aeruginosa isolated from cystic fibrosis patients to temocillin, a revived
antibiotic. *Sci Rep* **7**, 40208, doi:10.1038/srep40208 (2017).
57 Garnett, J. P. *et al.* Hyperglycaemia and Pseudomonas aeruginosa acidify cystic fibrosis
airway surface liquid by elevating epithelial monocarboxylate transporter 2 dependent
lactate-H(+) secretion. *Sci Rep* **6**, 37955, doi:10.1038/srep37955 (2016).
58 Worlitzsch, D. *et al.* Effects of reduced mucus oxygen concentration in airway
Pseudomonas infections of cystic fibrosis patients. *J Clin Invest* **109**, 317-325,
doi:10.1172/JCI13870 (2002).
59 Ayhan, D. H. *et al.* Sequence-Specific Targeting of Bacterial Resistance Genes
Increases Antibiotic Efficacy. *PLoS Biol* **14**, e1002552, doi:10.1371/journal.pbio.1002552
(2016).
60 Chan, B. K. *et al.* Phage selection restores antibiotic sensitivity in MDR Pseudomonas
aeruginosa. *Sci Rep* **6**, 26717, doi:10.1038/srep26717 (2016).
61 Chen, S. *et al.* Could Dampening Expression of the Neisseria gonorrhoeae mtrCDE-
Encoded Efflux Pump Be a Strategy To Preserve Currently or Resurrect Formerly Used
Antibiotics To Treat Gonorrhea? *MBio* **10**, doi:10.1128/mBio.01576-19 (2019).
62 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).
63 Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint **1303.3997** (2013).
64 Garcia-Alcalde, F. *et al.* Qualimap: evaluating next-generation sequencing alignment
data. *Bioinformatics* **28**, 2678-2679, doi:10.1093/bioinformatics/bts503 (2012).
65 Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* **9**, e112963,
doi:10.1371/journal.pone.0112963 (2014).
66 Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *J Comput Biol* **19**, 455-477, doi:10.1089/cmb.2012.0021 (2012).
67 Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-
2069, doi:10.1093/bioinformatics/btu153 (2014).
68 Page, A. J. *et al.* Roary: rapid large-scale prokaryote pan genome analysis.
Bioinformatics **31**, 3691-3693, doi:10.1093/bioinformatics/btv421 (2015).

69   Thorpe, H. A., Bayliss, S. C., Sheppard, S. K. & Feil, E. J. Piggy: a rapid, large-scale
pan-genome analysis tool for intergenic regions in bacteria. *Gigascience* **7**, 1-11,
doi:10.1093/gigascience/giy015 (2018).
70   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res* **47**, W256-W259, doi:10.1093/nar/gkz239 (2019).
71   Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant
bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* **43**, e15,
doi:10.1093/nar/gku1196 (2015).
72   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312-1313, doi:10.1093/bioinformatics/btu033
(2014).
73   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *J Mol Biol* **215**, 403-410, doi:10.1016/S0022-2836(05)80360-2 (1990).
74   Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066,
doi:10.1093/nar/gkf436 (2002).
75   Tonkin-Hill, G., Lees, J. A., Bentley, S. D., Frost, S. D. W. & Corander, J. Fast
hierarchical Bayesian analysis of population structure. *Nucleic Acids Res* **47**, 5539-5549,
doi:10.1093/nar/gkz361 (2019).
76   Johnson, S. R., Grad, Y., Abrams, A. J., Pettus, K. & Trees, D. L. Use of whole-genome
sequencing data to analyze 23S rRNA-mediated azithromycin resistance. *Int J*
Antimicrob Agents **49**, 252-254, doi:10.1016/j.ijantimicag.2016.10.023 (2017).
77   Kersh, E. N. *et al.* Rationale for a Neisseria gonorrhoeae Susceptible Only Interpretive
Breakpoint for Azithromycin. *Clin Infect Dis*, doi:10.1093/cid/ciz292 (2019).
78   Cock, P. J. *et al.* Biopython: freely available Python tools for computational molecular
biology and bioinformatics. *Bioinformatics* **25**, 1422-1423,
doi:10.1093/bioinformatics/btp163 (2009).
79   Dillard, J. P. Genetic Manipulation of Neisseria gonorrhoeae. *Curr Protoc Microbiol*
Chapter 4, Unit4A 2, doi:10.1002/9780471729259.mc04a02s23 (2011).
80   Jolley, K. A., Bray, J. E. & Maiden, M. C. J. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res*
3, 124, doi:10.12688/wellcomeopenres.14826.1 (2018).

856