1
Computed tomography reveals hip dysplasia
2
in the extinct Pleistocene saber-tooth cat *Smilodon*
3

Mairin A. Balisi[1,2*‡], Abhinav K. Sharma[3‡], Carrie M. Howard[1], Christopher A.

Shaw[1,4], Robert Klapper[5], Emily L. Lindsey[1,6,7]

[1] La Brea Tar Pits and Museum, Natural History Museum of Los Angeles County,
Los Angeles, California, United States of America

[2] Department of Life and Environmental Sciences, University of California, Merced,
California, United States of America

[3] David Geffen School of Medicine, University of California, Los Angeles,
California, United States of America

[4] Idaho Museum of Natural History, Idaho State University, Pocatello, Idaho, United
States of America

[5] Orthopaedics, Cedars Sinai Medical Group, Los Angeles, California, United
States of America

[6] Department of Earth Sciences, University of Southern California, Los Angeles,
California, United States of America

[7] Institute of the Environment and Sustainability, University of California – Los
Angeles, Los Angeles, California, United States of America

* Corresponding author

‡ Equal contribution

E-mail: mbalisi@tarpits.org

Abstract

Reconstructing the behavior of extinct species is challenging, particularly for those with no living analogues. However, damage preserved as paleopathologies on bone can record how an animal moved in life, potentially reflecting behavioral patterns. Here, we assess hypothesized etiologies of pathology in a pelvis and associated right femur of a *Smilodon fatalis* saber-toothed cat, one of the best-studied species from the Pleistocene-age Rancho La Brea asphalt seeps, California, USA, using visualization by computed tomography (CT). The pelvis exhibits massive destruction of the right hip socket that was interpreted, for nearly a century, to have developed from trauma and infection. CT imaging reveals instead that the pathological distortions characterize chronic remodeling that began at birth and led to degeneration of the joint over the animal's life. These results suggest that this individual suffered from hip dysplasia, a congenital condition common in domestic dogs and cats. This individual reached adulthood but could not have hunted properly nor defended territory on its own, likely relying on a social group for feeding and protection. While extant social felids are rare, these fossils and others with similar pathologies are consistent with a spectrum of social strategies in *Smilodon* supported by a predominance of previous studies.

Introduction

The saber-toothed cat *Smilodon fatalis* is one of the best-studied apex predators from the late Pleistocene, if not across the entire history of fossil mammals[1]. Much of our knowledge about this species comes from the Rancho La Brea (RLB) asphalt seeps

in Los Angeles, California, United States, which have preserved thousands of

individuals of *Smilodon* from at least 50,000 years ago until their extinction

approximately 11,000 years ago[1]. The seeps primarily functioned as a carnivore trap: a

large herbivore mired in the asphalt inadvertently would attract large carnivores and

scavengers, which themselves would become entrapped in great numbers[2]. Studies of

Smilodon at RLB have enabled reconstruction of its feeding behavior as an ambush

predator specializing on herbivorous megafauna, inferred using independent

approaches ranging from comparative morphology (e.g., [3]) to stable isotopes (e.g., [4]).

As well, the abundant specimens include numerous examples of bone lesions, or

pathologies[5]: a phenomenon that tends to be rare at more typical fossil sites (e.g.,

fluvial deposits, bone beds, ashfalls, caves) but captured by RLB's large sample sizes.

Because bone remodels throughout an animal's life in response to stress, strain,

and injury[6], paleopathologies can preserve a record of realized behavior and

supplement the picture of potential behavior presented by skeletal morphology.

Differences in the distribution of pathologies throughout the skeleton, for example,

distinguish *Smilodon* from a contemporaneous predator, the dire wolf *Aenocyon*

("*Canis*") *dirus*, reflecting interspecific differences in hunting behavior and prey

preference corroborated by independently gathered data[4]. While injuries in the dire wolf

tend to be concentrated around its distal limbs, supporting the hypothesis that it was a

pursuit predator, *Smilodon*'s injuries cluster around its midline, supporting inferences

that it ambushed and grappled with prey[5]. As the aggregate result of how an animal

moved over the course of its life, pathologies present a relatively direct record of the

animal's interactions with its predators, prey, environment, and even conspecifics,

potentially including intraspecific interactions like social behavior.

The current study centers on a *Smilodon* specimen (LACMHC 131) that has

been described as "the most strikingly pathological object in the collection of Rancho La

Brea fossils"[7]. The specimen is a right innominate bone exhibiting massive distortion

and destruction of the hip socket (Fig. 1). An analysis in 1930 by Moodie[7], restricted to

an inspection of the gross morphology, proposed that this pathology had resulted from

infection following violent trauma, possibly during an encounter with a conspecific,

which also led to dislocation of the femur from the hip. Moodie found no pathological

bones potentially associated with the injured innominate; because of disarticulation by

flowing asphalt over thousands of years, associated elements are rarely encountered at

RLB. However, more than half a century later, Shermis[8] described a pathological femur

(LACMHC 6963; Fig. 2) associated with another pathological pelvis. Later, this femur

was determined to be associated instead with the specimen described by Moodie[9],

enabling examination of the effects of a single event on associated skeletal elements.

In the present study, we supplement gross morphology and analyze LACMHC

131 and 6963 using computed tomography to observe the internal bone structure of

these elements. We evaluate the historical inference that this pathology was the result

of trauma and assess different etiologies: traumatic arthritis, infective arthritis, or

degenerative arthritis. We present basic morphometric data, quantitatively compare

these pathological bones to other pathological and non-pathological *Smilodon* elements

at RLB, and examine these specimens in the context of similar pathologies in living

cats. Finally, we explore the implications of the diagnosis on reconstructions of social

behavior in *Smilodon* and the potential contribution of paleopathology to a growing

interdisciplinary body of literature supporting sociality in this extinct predator.

Results

Description

LACMHC 131 is an intact pelvis with all sutures completely fused (Fig. 1;

Supplementary File S1; Supplementary Video S1). The distal end of LACMHC 6963, the

femur, is broken and missing (Fig. 2; Supplementary File S1), precluding verification of

distal epiphyseal fusion. However, the proximal epiphyses are fused to the shaft.

Imaging shows no signs of callus, or bone regeneration and healing, typically

seen following fracture. Rather, osteophytes likely signal bone remodeling secondary to

malformation of the joint. The changes in the right acetabulum and right femur are

consistent with those expected from repetitive subluxation and subsequent necrosis.

The right acetabulum is shallow and elliptical-shaped as opposed to concentric-shaped.

A hole in the bone, likely the result of postmortem asphaltic wear given its sharp edges,

marks the thin medial wall of the acetabulum, which is lined otherwise with exostoses.

The left acetabulum appears non-pathological; however, the ilium anterodorsal to the

socket—origin of the quadriceps femoris muscles—is rugose (Fig. 1; Supplementary

File S1; Supplementary Video S1), contrasting with typical *Smilodon* pelvic specimens

(e.g., Supplementary Fig. S1). The head of the pathological right femur is flattened and

laden with anatomical distortions; as well, the lesser trochanter is enlarged into a round

knob (Fig. 2; Supplementary File S1). In contrast, a non-pathological femur bears a

round head that is appropriately developed (Supplementary Fig. S2). This head would fit

snugly into a concentric-shaped socket such as the left acetabulum of the pathological

pelvis or either acetabulum of a non-pathological pelvis (e.g., Supplementary Fig. S1),

allowing for a normal axis of rotation and movement.

The four cardinal findings of arthritis on imaging are bony sclerosis, osteophytes,

joint space narrowing, and subchondral cysts. The CT images of the pathological

specimens reveal evidence of such degenerative changes and an absence of fractures

from traumatic impact (Figs. 3, 4; Supplementary Video S2; Supplementary Data S1-

S3). The images demonstrate sclerosis and osteophytes in the right acetabulum (Fig.

3), which are changes consistent with degenerative arthritis. Profuse remodeling with

osteophyte formation also marks the right femoral head (Fig. 4), likely in response to the

degenerative process from repeated subluxation and dysplasia. This contrasts with the

relatively smooth, evenly dense cortical bone of the non-pathological femur (Fig. 5).

Altogether, these findings indicate dysplasia of the hip as the pathological cause.

Body-size estimation

Basic morphometric data for the two pathological specimens are presented in

Table 1. The two specimens lie well within the size range of adult *Smilodon* pelves (Fig.

6) and femora (Fig. 7). These size comparisons as well as epiphyseal fusion suggest

that the individual represented by these two specimens was an adult.

Using published regressions of body mass on postcranial measurements in living

big cats[10–12], we estimated the body mass of the individual bearing the pathologies. Our

calculated results range from 114 kg (based on lengths on the right pathological

innominate) to 125 kg (based on lengths on the left non-pathological innominate; likely

an underestimate because big cats like the African lion comprise a minority of the broad

comparative sample[10]), to 229 kg (based on femur circumference[11]) to 309 kg (using a

different equation also based on femur circumference[12]). These estimates, like the

skeletal measurements on which they are based, fall well within the range of body-mass

estimates for non-pathological *Smilodon* from the same fossil deposit: from a median of

106 kg (median absolute deviation 14.15 kg) based on pelvis lengths[10]; to a median of

223 kg (median absolute deviation 24.95 kg) ranging up to 300 kg (median absolute

deviation 36.12 kg) based on femur circumference[11,12].

Prevalence at Rancho La Brea

The focal specimens are from RLB deposit ("pit") 61/67, which has produced 225

cataloged *Smilodon* pelvis specimens (99 left, 115 right, 13 complete; yielding a

minimum number of individuals (MNI) = 128). Of these 225 pelves, 28 (14 left, 11 right,

three complete; MNI = 17) bear lesions notable enough to have warranted inclusion in

the Pathology collection of the museum at La Brea Tar Pits[9]. Three other Pit 61/67

pelves in the Pathology collection show deformations like those on the pelvis central to

this study. These are: LACMHC 8904 (a left innominate; Fig. 13 in [7]), LACMHC 8916 (a

left innominate; Supplementary Fig. S3, this study), and LACMHC 9854 (a pelvis

preserving both sides, described by Shermis[8]). Findings include osteoarthritic alteration

of and chronic traction injuries around a shallow acetabulum, although etiology of these

pathologies would need to be confirmed with CT imaging as in Figs. 4 and 5. Pit 61/67's

total of four potentially dysplastic pelves (MNI = 4) brings the prevalence in this deposit

to a conservative estimate of 4 / 128 = 3.13%. All RLB deposits together catalog 1,624

Smilodon pelvis specimens (744 left, 772 right, 108 complete; MNI = 880); of these, we

found three additional specimens with possible dysplasia (Supplementary Figs. S4-S6).

This brings the dysplasia prevalence across all deposits to a conservative estimate of 7

/ 880 = 0.80%. Other dysplastic specimens may have escaped inclusion in the

Pathology collection, so this inferred percentage represents a minimum potential

incidence of dysplasia among *Smilodon* individuals preserved at RLB.

Similarly, of 714 cataloged *Smilodon* femora (381 left, 333 right; MNI = 381) in Pit

61/67, six specimens (two left, four right) preserve skeletal pathologies notable enough

to have warranted inclusion in the Pathology collection. However, LACMHC 6963—the

pathological femur in the current study—is the only obviously dysplastic femur from Pit

61/67 in the Pathology collection. Indeed, we are not aware of femora in any other RLB

deposit that have a femoral head as misshapen as that of LACMHC 6963. However,

distortion likely would be less apparent in incipient stages of dysplasia, and such

specimens may have escaped notice in the thousands of RLB specimens that have

been curated and databased over the last century.

Discussion

The arthritic degeneration visualized in the pathological *Smilodon* specimens

could have arisen from one of three etiologies: traumatic, infective or degenerative

arthritis. Findings on the specimens make infective or traumatic arthritis less likely. In

the case of infective arthritis, the presupposition is that the animal developed typically

before an insult that led to infection and subsequent obliteration of the hip joint. This

assumption also holds true for the case of traumatic arthritis following an injury or

fracture. However, the anatomical distortions of the right femoral head, in conjunction

with the obliteration of the right acetabulum, suggest chronic changes that led to

degeneration over time (Figs. 3, 4). The degeneration of the femoral head would not be

expected if the degenerative change in the hip joint were due to infection or trauma, as

the development of the pelvis and femur presumably would have been complete before

the insult or injury occurred during the adult cat's life.

Instead, the condition of the right acetabulum and right femoral head

demonstrates anatomy consistent with developmental distortion. Typically, the head of

the femur develops in conjunction with the acetabulum of the pelvis[13]. The spherical

femoral head fits into the concentric-shaped acetabulum to form a ball-and-socket joint

that enables a four-legged mammal to ambulate, lie down, sit down, stand up, and

generally function normally[13]. In developmental hip dysplasia, however, the acetabulum

does not develop appropriately, and the articulation between the femoral head and

acetabulum is lost. An elliptical (as opposed to concentric-shaped) acetabulum causes

progressive subluxation (dislocation) of the femoral head[14], which can result in coxa

plana, or necrosis of the bony nucleus of the femoral head[13]. This subsequent coxa

plana produces flattening and degeneration of the normally spherical femoral head[15].

Proper anatomical development and ossification of the hip joint rely on

continuous and symmetrical pressure of the femoral head on the acetabulum, and

dysplasia results from improper positioning of the femoral head within the acetabulum[13].

Dysplastic hips are characterized by pathological restructuring and accelerated

remodeling of the joint in response to abnormal forces and tensions that create stress.

This produces formation of new bone in some areas and resorption of bone in others,

ultimately causing degenerative joint disease[13]. Dysplastic hips have varying degrees of

deformity and malformation, but typically the acetabula are hypoplastic and deficient in

various planes and dimensions (Supplementary Fig. S7).

This pathology starts to impact movement at the time of first walking, although

minimal pain tends to ensue at this time because of the animal's flexibility at its early

age. As the joint cartilage wears out, however, bone begins to rub on bone. The

resulting forces make the bone stiffer, producing osteophytes or bone spurs as well as

sclerosis that manifests on CT imaging as increased bone density (Figs. 3 and 4;

Supplementary Video S2; Supplementary Data S1-S2). At this point, loading the limb

would cause pain, and range of motion would be limited. Therefore, the animal

examined in this study would have spent as little time as possible on its right hindlimb,

needing to compensate for the handicap by increasing the load on its left hindlimb. This

compensation would explain the exostoses on the left ilium anterodorsal to the non-

pathological acetabulum (Fig. 1; Supplementary File S1; Supplementary Video S1),

indicating abnormal pulling of the quadriceps femoris muscles originating in this area.

Hip dysplasia is a heritable, polygenic condition that affects a range of mammal

species[13], including humans[14]. Canine hip dysplasia (CHD) is one of the most prevalent

orthopedic diseases in domestic dogs (*Canis lupus familiaris*)[16] and is very well studied,

in part because it is similar to developmental dysplasia of the human hip[17]. Feline hip

dysplasia (FHD) has received less clinical attention than CHD, possibly because its

functional impairment is less overt or because domestic cats (*Felis catus*) are able to

compensate for the resulting lameness better than dogs[18,19]. The overall results of

physiological changes from dysplasia are mechanical imbalance and instability in the

hip joint causing displacement due to opposing forces from the acetabulum and femoral

head, and osteophytes in the acetabulum to compensate for cartilage loss[13].

Embryologically, articular joints differentiate from skeletal mesenchyme *in situ*

with the support of surrounding tissues that sustain mechanical and physiological forces

that tend to pull on the joints[13]. In dogs, hip joints are normal at birth, as teratological

factors and mechanical stresses that could displace the femoral head are rare at this

time[13]. Epiphyseal ossification normally begins by 12 days of age; in dogs that

eventually develop CHD, anatomical changes of the femoral head and pelvic socket

begin before week three[20]. In dysplastic hips, the teres ligament, which is crucial for

holding the femoral head in place, is too short; this produces luxation, or dislocation, of

the top of the femoral head, beginning at around seven weeks[13]. This luxation increases

throughout development, degrading the articular cartilage that surrounds the femoral

head, delaying ossification of the femur and acetabulum[13], and shortening the affected

limb, as the femoral head becomes positioned higher in the acetabulum.

In clinical reports of hip dysplasia in domestic cats, osteoarthritis (i.e.,

degenerative joint disease, DJD) of the hip secondary to FHD is well known[21]. For

example, osteoarthritis was recorded in 43 of 45 (95.6%) of cats with FHD[22]. As well, in

5 of 13 (38.5%) cases of hip osteoarthritis with a radiographically or historically

identifiable cause, hip dysplasia was pinpointed as the cause, with the remaining cases

resulting from trauma or equivocal between trauma and dysplasia[18]. A recent study of

FHD in Maine Coons—a large-bodied domestic cat breed in which hip dysplasia is

known to be common—calculated a prevalence of 37.4%, finding severity to increase

with age and body mass[23]. The same study further highlighted a genetic correlation

between FHD and large body size within the Maine Coon[23], inviting inquiry into how

FHD impacts other breeds and non-domestic felid species across a range of body sizes.

Reports of FHD in non-domestic large cats are rarer than in domestic cats.

Captive snow leopards have exhibited hip dysplasia; across 14 zoos, seven cases were

classified as moderate to severe, and at least two individual snow leopards needed total

hip replacement before being able to breed[24,25]. Accounts of hip functional impairment in

other captive large cats have tended to report osteoarthritis, which can be associated

with FHD though may also stem from trauma and increased age[26–28].

For wild-caught large cats, the only comprehensive study of which we are aware

is a survey of 386 individuals (283 wild-caught) across three felid genera mounted as

exhibit skeletons in multiple North American natural history museums[27]. Though not

focusing on hip dysplasia, the study tracked degenerative joint disease, which may be

associated with dysplasia[18,22]. The sample recorded DJD in 9.7% of 31 tigers, 2.3% of

88 African lions, and 5.1% of 59 mountain lions (*Puma concolor*), and none in five other

species of big cat. These frequencies are low compared to domestic cats, perhaps

owing to differences in body size, diet, and lifestyle between large wild cats and

domestic cats, as well as selective breeding constraining genetic variation in domestic

animals. Furthermore, selection against hip dysplasia would be expected in the wild

because hip dysplasia would compromise hunting[21]. Though this study identified

instances of non-inflammatory osteoarthritis in the shoulder, elbow, and stifle joint, it

found none in the hip. However, 4% of all joints afflicted by spondyloarthropathy—a

form of inflammatory arthritis—included the hip[27].

What is the significance of Smilodon, an extinct Pleistocene predator, having the

same congenital defect as living domestic cats and dogs? Previous workers have

inferred social behavior from *Smilodon's* pathologies, interpreting signs of healing as

evidence that the animal continued to live after injury[9]. Given the severity of many

injuries, authors have argued, the animal would have starved to death had it not

operated within a social structure. The present hip dysplasia having manifested from a

young age—hindering the animal's ability to hunt prey and defend a home range over

the course of its life—supports this assertion, although other inferences are possible.

Sociality, the degree to which individuals live with conspecifics in groups[29], is

difficult to infer in *Smilodon* given that it has no living analogues or closely related taxa.

Estimated to have weighed between 160 and 350 kg ([3,11,] this study), *Smilodon* was at

least the size of the Amur tiger (*Panthera tigris altaica*), the largest living cat; some

estimates reach 369 to 469 kg, placing *Smilodon* in the range of the largest extant

ursids[12,30]. No living felid has *Smilodon*'s elongate, knife-like canines or stocky, powerful

build. As well, *Smilodon* (of the extinct felid lineage Machairodontinae) is only distantly

related to extant large felids (Felinae), introducing further uncertainty. Based on its

robust morphology (e.g., [31,32]) and on evidence from stable isotopes (e.g., [4]), it likely

stalked and ambushed prey; therefore, it may have been comparable to the African lion

(*Panthera leo*), which has a similar hunting strategy and is the only truly social extant

felid[33]. Yet sociality varies across felid species, including within a genus; for example,

other extant pantherines like tigers (*P. tigris*) show incipient sociality[34], while jaguars (*P.*

onca) are solitary except for females with cubs. Social strategies also can vary within

species, e.g., between sexes. For instance, African lion females are philopatric and

social throughout their lives, while adult males are often nomadic and solitary until

joining a gregarious pride, which itself usually lasts for only a few years[35]. This social

variation complicates behavioral inferences based on ancestral reconstructions.

Advocates of the solitary-cat hypothesis[36,37] have cited *Smilodon*'s small relative

brain size determined using endocranial casts as support for solitary behavior, because

sociality exerts high cognitive demands. However, in 39 species across nine carnivoran

families, larger relative brain size was found to correlate with problem-solving

capabilities rather than social behavior[38]. Rather than analyses of overall

encephalization across carnivoran families, studies of relative regional brain volume

within families and species have been more informative regarding sociality[39,40]. In both

African lions and cougars (*Puma concolor*, a solitary species), total relative endocranial

volume was not sexually dimorphic; however, relative anterior cerebrum volume was

significantly greater in female African lions than males, a difference absent in cougars[35].

Though regional endocranial studies have yet to be performed on *Smilodon*, the

gregarious-cat hypothesis has drawn support from multiple lines of evidence. One is the

abundance of *Smilodon* relative to prey at RLB[7,8,31], although detractors have pointed

out that some extant large cats aggregate at carcasses despite otherwise being

solitary[37]. A full range of ages is present among RLB *Smilodon*; in contrast, animals

interpreted to be solitary, such as the American lion *Panthera atrox,* are represented

largely by adult individuals[41]. As well, the proportions of social and solitary species at

RLB parallel those drawn to audio recordings of herbivore distress calls in the African

savanna, suggesting that RLB *Smilodon* sample sizes are more consistent with it having

been social rather than solitary[42,43]. The lack of size sexual dimorphism in *Smilodon* is

more typical of modern solitary cats[44] but could also be reflective of monogamy within a

gregarious species, like modern wolves. Most relevant to the current study, the

existence of healed injuries in *Smilodon* also has been interpreted as evidence for

social behavior, with the assumption that surviving long after serious injury would be

difficult if not impossible without cooperative sociality[9]. We now revisit this interpretation

considering the novel diagnosis of hip dysplasia in this study.

Smilodon's large body size necessitated preying on megaherbivores for

adequate sustenance[3]. To do so, like most large cats today, it would have used its

hindlimbs for propulsion and acceleration[45,46], a pounce behavior enabled by its

morphology. *Smilodon's* ratio of total forelimb to hindlimb length is greater while its ratio

of tibia to femur length ranks lower than in living felids[31]. The shorter hindlimbs lacking

the distal limb elongation in cursorial animals suggest that *Smilodon* was an ambush

predator surpassing the ability of felids today[47]. Hunting large prey is dangerous[48]. After

the initial hindlimb-powered leap, *Smilodon* would have grappled with its struggling prey,

as evidenced by traumatic injuries in the rotator cuff and radiating from the ventral

midline dorsolaterally to where the ribs articulate with the spine[5]. As it subdued prey

with robust forelimbs[32,45] under enough torque to injure the lumbar vertebrae[5], *Smilodon*

would have needed to leverage itself against the ground using its hindlimbs. Therefore,

the pelvis and femur would have been critical to multiple phases of its hunting strategy.

A dysplastic individual would have encountered much difficulty hunting in this

manner. Yet, as evidenced by the complete fusion of its pelvic and femoral epiphyses

(Figs. 1, 2) as well as its large body size (Figs. 6, 7), the individual in this study had

reached adult age. (Studies of the detailed timing of epiphyseal fusion in large wild cats

are lacking, but distal femoral epiphyses fuse at around the same time as or soon after

proximal femoral epiphyses in domestic cats and dogs[49,50]. Given this, the broken distal

femur likely had a fused epiphysis, as on its intact proximal end.) Limbs in African lions

completely fuse between 4.5 and 5.5 years[51–53], so it is reasonable to assume that

adulthood in *Smilodon* likely started at around four years old. This estimate is reinforced

by bone histological work quantifying at least four to seven lines of arrested growth

(LAGs; one per growth year) in limb bones with fused epiphyses belonging to *Smilodon*

fatalis from the Talara asphaltic deposits in Peru[54]. Some LAGs in the Talara

histological specimens likely have been masked by secondary bone remodeling, which

may be more extensive in larger-bodied taxa[54], making these specimens possibly older

than the number of visible LAGs suggest. Therefore, four years represents a likely

minimum age for this individual, although it could have been much older.

Ontogenetic growth patterns in teeth and bone further support inferences of

sociality. In *Smilodon*, teeth appear to mature earlier than when sutures and long-bone

epiphyses fuse, suggesting delayed weaning, prolonged juvenile dependence, and

extended familial care until the adult hunting morphology—saber canines and robust

limbs—was complete[44]. At RLB, most sampled *Smilodon* specimens show significant

pulp cavity closure of the lower canine (14 of 19 specimens over approximately 80%

closure), a sign of dental maturation[55]. This contrasts with RLB pantherine pulp cavities,

which are more evenly distributed across the closure percentage range, suggesting that

teeth mature earlier in *Smilodon* than in pantherines. (Other age assessments have

ruled out the possibility that *Smilodon* juveniles were underrepresented relative to

pantherines[41].) At Talara, age determination by dentition yields low estimates of

juveniles (zero based on skulls; 8% based on dentaries), but age determination based

on limb epiphyseal fusion yields higher estimates (41% juveniles)[54]. Histology of Talara

Smilodon long bones reinforces this mismatch, as an apparent adult femur with fused

epiphyses and seven LAGs was found to lack avascular and acellular subperiosteal

lamellar bone[54], suggesting that it had not yet finished growing. Further, prolonged

parental care was interpreted in a recent description, from Pleistocene deposits in

Corralito, Ecuador, of two subadult *Smilodon fatalis* individuals inferred to have been

siblings and associated with an adult that was likely their mother[56]. This scenario of

prolonged parental care, like that in the social African lion, would help explain how the

individual in this current study survived to adulthood given its debilitating handicap.

Novel application of CT visualization to an old question of paleopathology has

enabled diagnosis of hip dysplasia, a lifelong condition, in an individual *Smilodon fatalis*

saber-toothed cat. This individual was likely not the only *Smilodon* afflicted with hip

dysplasia: multiple RLB *Smilodon* pelvic specimens, especially that described by

Shermis[8], exhibit gross morphology similar to the pathological pelvis examined in this

study (Supplementary Figs. S3-S6). The individual examined in this study reached

adulthood (at least four to seven years of age) but could never have hunted nor

defended territory on its own, given its locomotor impairment that would have been

present since infancy. As such, this individual likely survived to adulthood by association

with a social group that assisted it with feeding and protection.

Further conclusions are limited by the lack of a comprehensive and systematic

comparative dataset comprising pathological postcrania from extant species, a

persistent limitation of paleopathological studies[5]. Natural history museums may acquire

cranial remains from zoos or similar institutions but often lack storage to accommodate

postcranial skeletons, especially for large mammals. As well, while radiographic studies

on domestic cats and dogs illustrate the nature of hip dysplasia, these studies tend to

examine pathological bones *in situ*, still embedded in a muscular framework (e.g.,

Supplementary Fig. S7). This is opposed to the bones-only, flesh-free context of

paleopathological specimens. Computed tomography and digital data may be key to

building a comparative paleopathology dataset in the future.

Within the scope of this study, we cannot rule out the hypothesis that the

pathological animal was a scavenger and may have obtained food outside the context

of a social structure. It is also possible that, regardless of its disability, its large size and

fearsome canines made it a strong interference competitor. However, the pathological

specimens examined here are consistent with the predominance of studies supporting a

spectrum of social strategies in this extinct predator. In many extant carnivorans,

sociality offers the benefits of cooperative hunting and rearing of young (e.g., [57]):

benefits that likely also applied to *Smilodon* in the late Pleistocene. As *Smilodon*

coexisted with a rich megafaunal carnivore community including dire wolves (*Aenocyon*

dirus), American lions (*Panthera atrox*), and short-faced bears (*Arctodus simus*),

cooperative sociality may have aided its success as a predator in a crowded field.

Methods

All specimens examined are from the Pleistocene-age Rancho La Brea

collections of the museum at the La Brea Tar Pits (LBTPM), part of the Natural History

Museums of Los Angeles County (NHMLAC; formerly LACM), Los Angeles, California.

At Rancho La Brea, different fossiliferous asphaltic deposits (which became human-

made "pits" during the historical excavation process) resulted from periodic asphalt seep

activity, entrapping organisms across different timespans over the past 55,000 years[58]

with varied depositional environments and taphonomic histories[59]. In this context, we

reduced potential variability in these factors by selecting all specimens from a single

deposit, Pit 61/67. Pit 61/67 is the most recent deposit at Rancho La Brea that precedes

the late Pleistocene megafaunal extinctions at around 11,000 years before present[60], at

which point *Smilodon fatalis* became extinct.

We examined the external surfaces of the pathological pelvis including the right

innominate (LACMHC 131) and associated pathological right femur (LACMHC 6963).

We also inspected in detail an unassociated non-pathological right femur (LACMHC K-

3232) from the same deposit and of similar size and ontogenetic stage. Initial surface

scanning of all specimens was carried out at LBTPM using an Artec Space Spider

(Artec 3D) as a means of digital preservation and to provide a 3D visual with color. The

surface scans were processed in Artec Studio 12 and fused into a model with a

resolution of 0.2 mm. CT imaging of the three specimens was performed at the S. Mark

Taper Foundation Imaging Center, Los Angeles, California, on a GE Revolution (GE

Healthcare, Waukesha, WI) 256-slice scanner with 0.625 mm slice thickness. Imaging

parameters were KVP=120, mA=300, 0.5 second rotation time, and 0.51 pitch using a

medium body FOV. The data were acquired in the axial plane, reformatted into soft

tissue and bone algorithms, and viewed in the axial, coronal, and sagittal planes. CT

images were converted to 3D models using the segmentation software Mimics

(Materialise). Geomagic Freeform (3D Systems) was used to upload and determine

placement of the plane for cross-sections of the 3D reconstructions.

To estimate the individual's body size and the population incidence of this

pathology, we measured pelvis lengths and femur circumference on the pathological

specimens and compared them to non-pathological pelves and femora from Pit 61/67.

Lengths were measured using Mitutoyo calipers. We restricted this comparison to Pit

61/67, source of the pathological specimens, because previous work has revealed

body-size differences among *Smilodon* from different RLB deposits, likely due to climate

adaptation[61]. We estimated body size using published regressions of pelvis lengths[10]

and femur circumference[11,12] against body masses of living felids like the African lion.

While quantitative methods for scoring hip dysplasia exist (e.g., [62]), these

methods were developed for veterinary use on live animals presumably smaller than

Pleistocene megafauna, employed to describe x-rays of articulated limbs (e.g., [23]).

Therefore, this scoring system was inappropriate for the disarticulated and exposed

bony material in the paleontological record, as in this study. However, several RLB

Smilodon pelvic specimens preserve gross morphology consistent with dysplasia, which

we surveyed to calculate prevalence of this pathology in the RLB *Smilodon* population.

No permits were required for the described study.

References

455   1.   *Smilodon: The Iconic Sabertooth*. (Johns Hopkins University Press, 2018).

456   2.   Spencer, L. M., Van Valkenburgh, B. & Harris, J. M. Taphonomic analysis of large

457   mammals recovered from the Pleistocene Rancho La Brea tar seeps.

Paleobiology **29**, 561–575 (2003).

3. Van Valkenburgh, B., Hayward, M. W., Ripple, W. J., Meloro, C. & Roth, V. L. The impact of large terrestrial carnivores on Pleistocene ecosystems. *Proc. Natl. Acad. Sci.* **113**, 862–867 (2016).

4. DeSantis, L. R. G. *et al.* Causes and consequences of Pleistocene megafaunal extinctions as revealed from Rancho La Brea mammals. *Curr. Biol.* **29**, 2488-2495.e2 (2019).

5. Brown, C., Balisi, M. A., Shaw, C. A. & Van Valkenburgh, B. Skeletal trauma reflects hunting behaviour in extinct sabre-tooth cats and dire wolves. *Nat. Ecol. Evol.* **1**, 1–7 (2017).

6. Taylor, D., Hazenberg, J. G. & Lee, T. C. Living with cracks: Damage and repair in human bone. *Nat. Mater.* **6**, 263–268 (2007).

7. Moodie, R. L. Pleistocene luxations. *Am. J. Surg.* **IX**, 348–362 (1930).

8. Shermis, S. Healed massive pelvic fracture in a Smilodon from Rancho La Brea, California. *Paleobios* **1**, 121–126 (1983).

9. Shaw, C. A. & Ware, C. S. Smilodon paleopathology: a summary of research at Rancho La Brea. in *Smilodon: The Iconic Sabertooth* (eds. Werdelin, L., Mcdonald, H. G. & Shaw, C. A.) (Johns Hopkins University Press, 2018).

10. Toledo, N., Cassini, G. H., Vizcaíno, S. F. & Bargo, M. S. Mass estimation of Santacrucian sloths from the early Miocene Santa Cruz Formation of Patagonia, Argentina. *Acta Palaeontol. Pol.* **59**, 267–280 (2014).

11. Christiansen, P. & Harris, J. M. Body size of Smilodon (Mammalia: Felidae). *J. Morphol.* **266**, 369–384 (2005).

481   12.   Anyonge, W. Body mass in large extant and extinct carnivores. *J. Zool.* **231**, 339–

482   350 (1993).

483   13.   Riser, W. H., Rhodes, W. H. & Newton, C. D. Hip dysplasia. in *Textbook of Small*

484   *Animal Orthopaedics* (J.B. Lippincott Company, 1985).

485   14.   Weinstein, S. L. Natural history of congenital dislocation and dysplasia. *Clin.*

486   *Orthop. Relat. Res.* **225**, 62–76 (1987).

487   15.   Creel, R. Congenital hip dysplasia. *Iowa State Univ. Vet.* **23**, 32–33 (1961).

488   16.   Bartolomé, N. *et al.* A genetic predictive model for canine hip dysplasia:

489   integration of Genome Wide Association Study (GWAS) and candidate gene

490   approaches. *PLoS One* **10**, 1–13 (2015).

491   17.   Zhou, Z. *et al.* Differential genetic regulation of canine hip dysplasia and

492   osteoarthritis. *PLoS One* **5**, (2010).

493   18.   Clarke, S. P. *et al.* Prevalence of radiographic signs of degenerative joint disease

494   in a hospital population of cats. *Vet. Rec.* **157**, 793–799 (2005).

495   19.   Perry, K. Feline hip dysplasia: A challenge to recognise and treat. *J. Feline Med.*

496   *Surg.* **18**, 203–218 (2016).

497   20.   Kealy, R. D. *et al.* Evaluation of the effect of limited food consumption on

498   radiographic evidence of osteoarthritis in dogs. *J. Am. Vet. Med. Assoc.* **217**,

499   1678–1680 (2000).

500   21.   Lascelles, B. D. X. Feline degenerative joint disease. *Vet. Surg.* **39**, 2–13 (2010).

501   22.   Keller, G. G., Reed, A. L., Lattimer, J. C. & Corley, E. A. Hip dysplasia: A feline

502   population study. *Vet. Radiol. Ultrasound* **40**, 460–464 (1999).

23.   Low, M. *et al.* Demography, heritability and genetic correlation of feline hip

dysplasia and response to selection in a health screening programme. *Sci. Rep.*

9, 1–9 (2019).

24. Wharton, D. & Mainka, S. A. Management and husbandry of the snow leopard

Uncia uncia. *Int. Zoo Yearb.* **35**, 139–147 (1997).

25. Suedmeyer, W. K., Cook, J. L., Tomlinson, J. L. & Crouch, D. T. Bilateral total hip

replacement in a snow leopard (Uncia uncia) with bilateral hip dysplasia. in

International Association for Aquatic Animal Medicine (2000).

26. Whiteside, D. P., Remedios, A. M., Black, S. R. & Finn-Bodner, S. T. Meloxicam

and surgical denervation of the coxofemoral joint for the treatment of degenerative

osteoarthritis in a Bengal tiger (Panthera tigris tigris). *J. Zoo Wildl. Med.* **37**, 416–

419 (2006).

27. Rothschild, B. M., Rothschild, C. & Woods, R. J. Inflammatory arthritis in large

cats: An expanded spectrum of spondyloarthropathy. *J. Zoo Wildl. Med.* **29**, 279–

284 (1998).

28. Napier, J. E., Lund, M. S., Armstrong, D. L. & McAloose, D. A retrospective study

of morbidity and mortality in the North American Amur leopard (Panthera pardus

orientalis) population in zoologic institutions from 1992 to 2014. *J. Zoo Wildl. Med.*

49, 70–78 (2018).

29. Eisenberg, J. F. The social organisation of mammals. *Handb. der Zool.* **10**, 1–92

(1966).

30. Torregrosa, V., Petrucci, M., Pérez-Claros, J. A. & Palmqvist, P. Nasal aperture

area and body mass in felids: Ecophysiological implications and paleobiological

inferences. *Geobios* **43**, 653–661 (2010).

31. Gonyea, W. J. Behavioral implications of saber-toothed felid morphology.

Paleobiology **2**, 332–342 (1976).

32. Meachen-Samuels, J. A. Morphological convergence of the prey-killing arsenal of

sabertooth predators. *Paleobiology* **38**, 1–14 (2012).

33. Schaller, G. B. *The Serengeti lion: a study of predator-prey relations.* (University

of Chicago Press, 1972).

34. Karanth, K. U. & Sunquist, M. E. Behavioural correlates of predation by tiger

(Panthera tigris), leopard (Panthera pardus), and dhole in Nagarahole, India. *J.*

Zool. London **250**, 255–265 (2000).

35. Arsznov, B. M. & Sakai, S. T. Pride diaries: Sex, brain size and sociality in the

African lion (Panthera leo) and cougar (Puma concolor). *Brain. Behav. Evol.* **79**,

275–289 (2012).

36. McCall, S., Naples, V. & Martin, L. D. Assessing behavior in extinct animals: Was

Smilodon social? *Brain. Behav. Evol.* **61**, 159–164 (2003).

37. Kiffner, C. Coincidence or evidence: was the sabretooth cat Smilodon social?

Biol. Lett. **5**, 561–562 (2009).

38. Benson-Amram, S., Dantzer, B., Stricker, G., Swanson, E. M. & Holekamp, K. E.

Brain size predicts problem-solving abilities in mammalian carnivores. *Proc. Natl.*

Acad. Sci. **113**, 2532–2537 (2016).

39. Sakai, S. T., Arsznov, B. M., Hristova, A. E., Yoon, E. J. & Lundrigan, B. L. Big cat

coalitions: A comparative analysis of regional brain volumes in Felidae. *Front.*

Neuroanat. **10**, 1–12 (2016).

40. Vinuesa, V. *et al.* Inferences of social behavior in bone-cracking hyaenids

(Carnivora, Hyaenidae) based on digital paleoneurological techniques:

Implications for human–carnivoran interactions in the Pleistocene. *Quat. Int.* **413**,

7–14 (2016).

41.   Miller, G. J. On the distribution of Smilodon californicus Bovard from Rancho La

Brea. *Los Angeles Cty. Museum Contrib. Sci.* **131**, 1–19 (1968).

42.   Carbone, C. *et al.* Parallels between playbacks and Pleistocene tar seeps suggest

sociality in an extinct sabretooth cat, Smilodon. *Biol. Lett.* **5**, 81–85 (2009).

43.   Van Valkenburgh, B. *et al.* Sociality in Rancho La Brea Smilodon: arguments

favour 'evidence' over 'coincidence'. *Biol. Lett.* **5**, 563–564 (2009).

44.   Van Valkenburgh, B. & Sacco, T. Sexual dimorphism, social behavior, and

intrasexual competition in large Pleistocene carnivorans. *J. Vertebr. Paleontol.* **22**,

164–169 (2002).

45.   Meachen-Samuels, J. A. & van Valkenburgh, B. Radiographs reveal exceptional

forelimb strength in the sabertooth cat, Smilodon fatalis. *PLoS One* **5**, (2010).

46.   Martín-Serra, A., Figueirido, B. & Palmqvist, P. A three-dimensional analysis of

the morphological evolution and locomotor behaviour of the carnivoran hind limb.

BMC Evol. Biol. **14**, 1–13 (2014).

47.   Anyonge, W. Locomotor behaviour in Plio-Pleistocene sabre-tooth cats: a

biomechanical analysis. *J. Zool.* **238**, 395–413 (1996).

48.   Carbone, C., Teacher, A. & Rowcliffe, J. M. The costs of carnivory. *PLoS Biol.* **5**,

363–368 (2007).

49.   Sumner-Smith, G. Observations on epiphyseal fusion of the canine appendicular

skeleton. *J. Small Anim. Pract.* **7**, 303–311 (1966).

50.   Smith, R. N. Fusion of Ossification Centres in the Cat. *J. Small Anim. Pract.* **10**,

523–530 (1969).

51.   Smuts, G. L., Robinson, G. A. & Whyte, I. J. Comparative growth of wild male and

female lions (Panthera leo). *J. Zool.* **190**, 365–373 (1980).

52.   Kirberger, R. M., du Plessis, W. M. & Turner, P. H. Radiologic anatomy of the

normal appendicular skeleton of the lion (Panthera leo). Part 1: thoracic limb. *J.*

Zoo Wildl. Med. **36**, 29–35 (2005).

53.   Kirberger, R. M., du Plessis, W. M. & Turner, P. H. Radiologic anatomy of the

normal appendicular skeleton of the lion (Panthera leo). Part 2: pelvic limb. *J. Zoo*

Wildl. Med. **36**, 29–35 (2005).

54.   Seymour, K. L., Reynolds, A. R. & Churcher, C. S. Smilodon fatalis from Talara,

Peru: Sex, age, mass, and histology. in *Smilodon: The Iconic Sabertooth* (eds.

Werdelin, L., McDonald, H. G. & Shaw, C.) (Johns Hopkins University Press,

2018).

55.   Meachen-Samuels, J. A. & Binder, W. J. Sexual dimorphism and ontogenetic

growth in the American lion and sabertoothed cat from Rancho La Brea. *J. Zool.*

280, 271–279 (2010).

56.   Reynolds, A. R., Seymour, K. L. & Evans, D. C. Smilodon fatalis siblings reveal

life history in a saber-toothed cat. *iScience* 101916 (2020)

doi:10.1016/j.isci.2020.101916.

57.   Macdonald, D. W. The ecology of carnivore social behaviour. *Nature* **301**, 379–

384 (1983).

58.   O'Keefe, F. R., Fet, E. V & Harris, J. M. Compilation, calibration, and synthesis of

faunal and floral radiocarbon dates, Rancho La Brea, California. *Contrib. Sci.* **518**,

1–16 (2009).

59. Mychajliw, A. M., Rice, K. A., Tewksbury, L. R., Southon, J. R. & Lindsey, E. L.

Exceptionally preserved asphaltic coprolites expand the spatiotemporal range of a

North American paleoecological proxy. *Sci. Rep.* **10**, 1–12 (2020).

60. Fuller, B. T., Harris, J. M., Farrell, A. B., Takeuchi, G. T. & Southon, J. R. Sample

preparation for radiocarbon dating and isotopic analysis of bone from Rancho La

Brea. *Nat. Hist. Museum Los Angeles Cty. Sci. Ser.* **42**, 151–167 (2015).

61. Meachen, J. A., O'Keefe, F. R. & Sadlier, R. W. Evolution in the sabre-tooth cat,

Smilodon fatalis, in response to Pleistocene climate change. *J. Evol. Biol.* **27**,

714–723 (2014).

62. Loder, R. T. & Todhunter, R. J. Demographics of hip dysplasia in the Maine Coon

cat. *J. Feline Med. Surg.* **20**, 302–307 (2018).

Figure Legends

611   **Figure 1.** Photographs of LACMHC 131, a pathological pelvis belonging to

Smilodon fatalis. **(A)** Right lateral view showing destruction of the acetabulum;

anterodorsal end to the right. **(B)** Left lateral view showing the acetabulum intact but

with exostoses around the anterodorsal rim; anterodorsal end to the left. **(C)** Dorsal and

(D) ventral views showing asymmetry in the pelvis; anterior end to the right.

616   **Figure 2.** Photographs of LACMHC 6963, a pathological right femur belonging to

Smilodon fatalis. **(A)** Anterior and **(B)** posterior views of full femur, excluding the distal

epiphysis broken after death; proximal end on the left. **(C)** Anterior and **(D)** posterior

close-up views of the proximal end. **(E)** Dorsal close-up view of the femoral head,

greater trochanter, and lesser trochanter in lower center background. **(F)** Lateral close-

up view of the greater trochanter and lesser trochanter (lower center), which is enlarged

into a round knob. The upper scale bar refers to A and B and the lower scale bar refers

to C, D, E, and F.

Figure 3. Three-dimensional model of the pathological *Smilodon* pelvis and

acetabulum (LACMHC 131) reconstructed from CT images. Horizontal numbered lines

in (A) mark anatomical location of numbered CT slices displayed in (E). **(A)** Ventral view

of pelvis, anterior toward top of image; **(B)** rotated 45 degrees from ventral about the

anteroposterior axis; **(C)** rotated 90 degrees from ventral about the anteroposterior axis;

(D) rotated 135 degrees from ventral about the anteroposterior axis. **(E)** 0.63mm axial

CT images of the pathological pelvis demonstrating a dysplastic right hip socket,

corresponding with the three-dimensional model, posterior view.

Figure 4. Three-dimensional model of the pathological *Smilodon* right femur

reconstructed from CT images. Horizontal numbered lines in (A) mark anatomical

location of numbered CT slices displayed in (E). **(A)** Anterior view of right femur; **(B)**

rotated 90 degrees from anterior about the vertical axis (lateral view); **(C)** rotated 180

degrees from anterior about the vertical axis (posterior view); **(D)** rotated 270 degrees

from anterior about the vertical axis, displaying the femoral head and neck (medial

view). **(E)** 0.63mm axial CT images of the pathological right femur demonstrating

morphological distortion at the femoral head and neck, corresponding with the three-

dimensional model. Each inset image has been rotated so that top corresponds to

anterior, right corresponds to anatomical right, bottom corresponds to posterior, and left

corresponds to anatomical left sides.

Figure 5. Three-dimensional model of a normal *Smilodon* right femur (LACMHC

K-2584) reconstructed from CT images. Horizontal numbered lines in (A) mark

anatomical location of numbered CT slices displayed in (E). **(A)** Anterior view of right

femur; **(B)** rotated 90 degrees from anterior about the vertical axis (lateral view); **(C)**

rotated 180 degrees from anterior about the vertical axis (posterior view); **(D)** rotated

270 degrees from anterior about the vertical axis, displaying the femoral head and neck

(medial view). **(E)** 1.25mm axial CT images of the normal right femur corresponding with

the three-dimensional model. Each inset image has been flipped and rotated so that top

corresponds to anterior, right corresponds to anatomical right, bottom corresponds to

posterior, and left corresponds to anatomical left sides.

Figure 6. Metric data for the pathological *Smilodon* pelvis (LACMHC 131) in

comparison to non-pathological specimens also from Pit 61/67 and potentially dysplastic

pelvis specimens from other deposits. Inset diagram illustrates lengths measured.

Across all measurements, the pathological pelvis lies well within the size range of non-

pathological adult *Smilodon* pelves, suggesting that the animal to which it belonged also

had grown to adult age. Numeric values are in Supplementary Data S4. **(A)** Bivariate

plot showing ilium length (from the anterior-most point of the pelvis to the anterior

border of the acetabulum) against total anteroposterior pelvis length. **(B)** Bivariate plot

showing ischium length (from the posterior-most point of the pelvis to the posterior

border of the acetabulum) against total anteroposterior pelvis length. **(C)** Bivariate plot

showing ischium length against ilium length.

Figure 7. Metric data for the pathological *Smilodon* femur (LACMHC 6963) in

comparison to non-pathological adult specimens also from Pit 61/67. Numeric values

are in Supplementary Data S5. **(A)** Bivariate plot of mediolateral diameter against

anteroposterior diameter of the femoral head. The femoral head of LACMHC 6963 is

mediolaterally elongated and anteroposteriorly compressed compared to non-

pathological specimens. **(B)** Bivariate plot of mediolateral diameter against

anteroposterior diameter of the femoral diaphysis. The diaphysis of LACMHC 6963 is

mediolaterally compressed relative to its anteroposterior diameter. **(C)** Histogram of

femoral diaphysis least circumference, showing LACMHC 6963 (orange line) well within

the range of non-pathological adult specimens and suggesting that the individual with

the pathology also had reached adult age.

Table

Table 1. Basic morphometric data for the two focal pathological *Smilodon fatalis*

specimens: (A) pelvis, LACMHC 131; (B) right femur, LACMHC 6963. Morphometric

data for comparative specimens are in Supplementary Data S4 and S5 and plotted in

Figures 6 and 7.

	Left	Right
	(non-pathological)	(pathological)
A. Pelvis measurements (mm)		
Total anteroposterior length	336.20	340.20
Ilium length (to anterior-most border	172.70	167.74

of acetabulum)		
Ischium length (from posterior-most border of acetabulum)	116.40	120.24
B. Femur measurements (mm)		
Intertrochanteric width		85.08
Head length		49.90
Head width		38.35
Transverse (mediolateral) diameter		32.73
Anteroposterior diameter		32.84
Diaphysis least circumference		109.00

Acknowledgements

We acknowledge that Rancho La Brea is on the traditional territory and

homelands of the Chumash and Gabrielino / Tongva people. We thank Aisling Farrell at

LBTPM for facilitating specimen access and loans; CT segmentation specialists at the

S. Mark Taper Foundation Imaging Center for CT-scanning specimens; Jeff Busey and

Jessica Butler at Zimmer Biomet for aiding design of 3D models; Karin Rice for x-ray

images of Erik the cat; Fred Heald for his work on the LBTPM Pathology collection; and

Andrew Kitchener, Carlo Meloro, Alexis Mychajliw, Blaire Van Valkenburgh, and an

anonymous reviewer for constructive comments. A U.S. National Science Foundation

Postdoctoral Research Fellowship in Biology (DBI-1812301) funded MAB.

Author Contributions

MAB and AKS wrote the manuscript text; CMH surface-scanned specimens and

created videos; MAB, CMH, and AKS prepared figures; MAB collected quantitative data;

CAS described comparative-specimen pathologies; RK and ELL acquired CT scans;

MAB and ELL guided study design. All authors reviewed and approved the manuscript.

Additional Information

Competing interests

The authors declare no competing interests.

A

B

3 cm

C

D

E

F

3 cm

max

*bone
density*

min