Role of the transcriptional regulator SP140 in resistance to bacterial infections via repression of type I interferons

Daisy X. Ji[1*], Kristen C. Witt[1*], Dmitri I. Kotov[1,2], Shally R. Margolis[1], Alexander Louie[1], Victoria Chevée[1], Katherine J. Chen[1,2], Harmandeep S. Dhaliwal[3], Angus Y. Lee[3], Dario S. Zamboni[4], Igor Kramnik[5], Daniel A. Portnoy[1,6,7], K. Heran Darwin[8], Russell E. Vance[1,2,3**]

[1] Division of Immunology and Pathogenesis, Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720 USA

[2] Howard Hughes Medical Institute, University of California, Berkeley, CA 94720 USA

[3] Cancer Research Laboratory, University of California, Berkeley, CA 94720 USA

[4] Department of Cell Biology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil.

[5] The National Emerging Infectious Diseases Laboratory, Department of Medicine (Pulmonary Center), and Department of Microbiology, Boston University School of Medicine, Boston, MA, USA.

[6] Division of Biochemistry, Biophysics and Structural Biology, Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720 USA

[7] Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720 USA

[8] Department of Microbiology, New York University Grossman School of Medicine, New York, NY, USA

* These authors contributed equally to this work.

** e-mail: rvance@berkeley.edu

Impact Statement: Repression of type I interferons by SP140 is essential for resistance to *Legionella pneumophila*, and *Mycobacterium tuberculosis*.

Abstract

Type I interferons (IFNs) are essential for anti-viral immunity, but often impair protective

immune responses during bacterial infections. An important question is how type I IFNs are

strongly induced during viral infections, and yet are appropriately restrained during bacterial

infections. The *Super susceptibility to tuberculosis 1* (*Sst1*) locus in mice confers resistance to

diverse bacterial infections. Here we provide evidence that *Sp140* is a gene encoded within the

Sst1 locus that represses type I IFN transcription during bacterial infections. We generated

Sp140$^{-/-}$ mice and find they are susceptible to infection by *Legionella pneumophila* and

Mycobacterium tuberculosis. Susceptibility of *Sp140$^{-/-}$* mice to bacterial infection was rescued

by crosses to mice lacking the type I IFN receptor (*Ifnar$^{-/-}$*). Our results implicate *Sp140* as an

important repressor of type I IFNs that is essential for resistance to bacterial infections.

Introduction

Type I interferons (IFNs) comprise a group of cytokines, including interferon-β and multiple interferon-α isoforms, that are essential for immune defense against most viruses (Stetson and Medzhitov, 2006). Type I IFNs signal through a cell surface receptor, the interferon alpha and beta receptor (IFNAR), to induce an 'anti-viral state' that is characterized by the transcriptional induction of hundreds of interferon stimulated genes (ISGs) (Schneider et al., 2014). Many ISGs encode proteins with direct anti-viral activities. Type I IFNs also promote anti-viral responses by cytotoxic T cells and Natural Killer cells. Accordingly, $Ifnar^{-/-}$ mice are highly susceptible to most viral infections.

Many ISGs are also induced by IFN-γ (also called type II IFN). However, type I and type II IFNs appear to be specialized for the control of different classes of pathogens (Crisler and Lenz, 2018). Whereas type I IFNs are predominantly anti-viral, the ISGs induced by IFN-γ appears to be especially important for the control of diverse intracellular pathogens, including bacteria and parasites. In contrast, type I IFNs play complex roles during bacterial infections (Boxx and Cheng, 2016; Donovan et al., 2017; McNab et al., 2015; Moreira-Teixeira et al., 2018). Some ISGs induced by type I IFN, most notably certain guanylate binding proteins (GBPs), have anti-bacterial activities (Pilla-Moffett et al., 2016). At the same time, several proteins induced by type I IFNs, including interleukin-10 (IL-10) and IL-1 receptor antagonist (IL-1RA), impair anti-bacterial immunity (Boxx and Cheng, 2016; Ji et al., 2019; Mayer-Barber et al., 2014). As a result, the net effect of type I IFN is often to increase susceptibility to bacterial infections. For example, $Ifnar^{-/-}$ mice exhibit enhanced resistance to *Listeria monocytogenes* (Auerbuch et al., 2004; Carrero et al., 2004; O'Connell et al., 2004) and *Mycobacterium tuberculosis* (Donovan et al., 2017; Dorhoi et al., 2014; Ji et al., 2019; Mayer-Barber et al., 2014; Moreira-Teixeira et al., 2018). Multiple mechanisms appear to explain resistance of $Ifnar^{-/-}$ mice to *L. monocytogenes*, including a negative effect of type I IFNs on protective IFN-γ signaling (Rayamajhi et al., 2010). Likewise, diverse mechanisms underlie the negative effects of type I IFNs during *M. tuberculosis* infection, including alterations of eicosanoid production (Mayer-Barber et al., 2014) and the induction of IL-1Ra (Ji et al., 2019), both of which impair protective IL-1 responses.

As an experimental model for dissecting the mechanisms by which inappropriate type I IFN responses are restrained during bacterial infections, we have compared mice harboring different haplotypes of the *Super susceptibility to tuberculosis 1* (*Sst1*) locus (Pan et al., 2005; Pichugin et al., 2009). The *Sst1* locus encompasses about 10M base pairs of mouse chromosome 1, a region that contains approximately 50 genes. Mice harboring the susceptible (S) haplotype of *Sst1*, derived from the C3H/HeBFeJ mouse strain, succumb relatively rapidly to *M. tuberculosis*

infection as compared to isogenic mice harboring the resistant (R) *Sst1* haplotype (derived from

C57BL/6 mice). Likewise, $Sst1^S$ mice also exhibit enhanced susceptibility to *Listeria*

monocytogenes (Boyartchuk et al., 2004; Pan et al., 2005) and *Chlamydia pneumoniae* (He et al.,

2013). The susceptibility of $Sst1^S$ mice to *M. tuberculosis* was reversed by crossing to $Ifnar^{-/-}$

mice (He et al., 2013; Ji et al., 2019), thereby demonstrating the causative role of type I IFNs in

driving the susceptibility phenotype. Although multiple type I IFN-induced genes are likely

responsible for the detrimental effects of type I IFNs during bacterial infections, heterozygous

deficiency of a single type I IFN-induced gene, *Il1rn* (encoding IL-1 receptor antagonist), was

sufficient to almost entirely reverse the susceptibility of $Sst1^S$ mice to *M. tuberculosis* (Ji et al.,

2019).

The $Sst1^R$ haplotype is dominant over the $Sst1^S$ haplotype, suggesting that $Sst1^R$ likely

encodes a protective factor that is absent from $Sst1^S$ mice (Pan et al., 2005; Pichugin et al., 2009).

By comparing gene expression in $Sst1^R$ versus $Sst1^S$ mice, *Sp110* (also known as *Ipr1*) was

discovered as an *Sst1*-encoded gene that is transcribed selectively in $Sst1^R$ mice (Pan et al.,

2005). Transgenic expression of *Sp110* in $Sst1^S$ mice partially restored resistance to *M.*

tuberculosis and *L. monocytogenes* (Pan et al., 2005). However, the causative role of *Sp110* in

conferring resistance to bacterial infections was not confirmed by the generation of *Sp110*-

deficient B6 mice. Null mutations of human *SP110* are associated with VODI (hepatic veno-

occlusive disease with immunodeficiency syndrome, OMIM 235550), but not mycobacterial

diseases (Roscioli et al., 2006). Some studies have found polymorphisms in *SP110* to be

associated with susceptibility to TB, though not consistently so across different ethnic groups

(Chang et al., 2018; Fox et al., 2014; Lei et al., 2012; Png et al., 2012; Thye et al., 2006; Tosh et

al., 2006; Zhang et al., 2017).

In humans and mice, SP110 is a part of the Speckled Protein (SP) family of nuclear

proteins, consisting of SP100, SP110 and SP140 (and SP140L in humans only) (Perniola and

Musco, 2014). The SP family members also exhibit a high degree of similarity to AIRE, a

transcriptional regulator that promotes tolerance to self-antigens by inducing their expression in

thymic epithelial cells (Anderson and Su, 2016; Perniola and Musco, 2014). All members of the

SP-AIRE family in both mice and humans have an N-terminal SP100 domain that appears to

function as a homotypic protein-protein interaction domain (Fraschilla and Jeffrey, 2020; Huoh

et al, 2020). The SP100 domain is closely related to the Caspase Activation and Recruitment

Domain (CARD), though SP family members are not believed to activate caspases. SP-AIRE

proteins also contain a DNA-binding SAND domain (Bottomley et al., 2001). Certain SP

isoforms, including all human full-length SP family members and mouse SP140, also include a

plant homeobox domain (PHD) and a bromodomain (BRD) (Perniola and Musco, 2014). The

genes encoding SP family proteins are linked in a small cluster in both mouse and human

genomes and are inducible by IFN-γ in a variety of cell lines. The mouse *Sp100/110/140* gene

cluster is adjacent to a highly repetitive 'homogenously staining region' (HSR) of chromosome 1

that remains poorly assembled in the most recent genome assembly due to the presence of as

many as 40 near-identical repeats of *Sp110*-like sequences (Pan et al., 2005; Weichenhan et al.,

2001). Most of these repeated *Sp110*-like sequences in the HSR appear to be either incomplete

copies of *Sp110* or pseudogenes that are not believed to be translated, but their presence has

nevertheless complicated genetic targeting and analysis of the SP gene family.

With the advent of CRISPR–Cas9-based methods (Wang et al., 2013), we were able to

generate *Sp110*$^{-/-}$ mice on the B6 background. Surprisingly, we found that *Sp110*$^{-/-}$ mice do not

phenocopy the susceptibility of *Sst1*^{S} mice to *M. tuberculosis* infection *in vivo*. Upon analysis of

additional candidate genes in the *Sst1* locus, we found that B6.*Sst1*^{S} mice also lack expression of

Sp140. To test whether loss of *Sp140* might account for the susceptibility of *Sst1*^{S} mice to

bacterial infections, we generated *Sp140*$^{-/-}$ mice. We found these mice are as susceptible as

B6.*Sst1*^{S} mice to the intracellular bacterial pathogens *M. tuberculosis* and *Legionella*

pneumophila. Similar to B6.*Sst1*^{S} mice, *Sp140*$^{-/-}$ mice exhibit an exacerbated type I IFN

response after bacterial infection, and the susceptibility of *Sp140*$^{-/-}$ mice is rescued by crosses to

Ifnar$^{-/-}$ mice. Our results suggest that loss of *Sp140* explains the susceptibility to bacterial

infections associated with the *Sst1*^{S} haplotype. These data further suggest that SP140 is a novel

negative regulator of type I IFN responses that is essential for protection against intracellular

bacterial infections.

Results

***Sp110*$^{-/-}$ mice are not susceptible to *M. tuberculosis*.** Loss of *Sp110* expression was proposed to

account for the susceptibility of mice carrying the *Sst1*^{S} haplotype to bacterial infections (Pan et

al., 2005). We first confirmed that bone marrow-derived macrophages (BMMs) from B6.*Sst1*^{S}

mice lack expression of *Sp110* protein (Figure 1A). To determine whether loss of *Sp110* confers

susceptibility to bacterial infections, we used CRISPR–Cas9 to target exon 3 of *Sp110* to

generate *Sp110*$^{-/-}$ mice on the C57BL/6 (B6) background (Figure 1– figure supplement 1). We

generated three independent *Sp110*$^{-/-}$ lines, denoted as lines 61, 65 and 71 (Figure 1A, Figure 1–

figure supplement 1). All three lines lacked expression of *Sp110*, as verified using three different

antibodies (Figure 1A). *Sp110*$^{-/-}$ mice are viable and are born at normal Mendelian ratios and

litter sizes. When aerosol infected with a low-dose of *M. tuberculosis*, *Sp110*$^{-/-}$ mice did not

phenocopy the susceptibility observed in B6.*Sst1*^{S} mice (Figure 1B-D). At day 25 post-infection,

Sp110⁻/⁻ lungs resembled those of wild-type B6 mice (Figure 1B) and harbored fewer bacteria

than the lungs of B6.*Sst1^S* mice, similar to both the B6 and *Sp110$^{+/-}$* littermates (Figure 1C).

Likewise, the survival of infected *Sp110$^{-/-}$* mice was indistinguishable from B6 mice, and mice

of both genotypes survived considerably longer than the B6.*Sst1^S* mice (Figure 1D). Thus,

despite the absence of *Sp110* from *Sst1^S* mice, our results indicate that the loss of *Sp110* is not

sufficient to replicate the susceptibility to *M. tuberculosis* associated with the *Sst1^S* locus.

***Sp140$^{-/-}$* mice are susceptible to bacterial infections.** Given that *Sp110* deficiency did not

phenocopy the susceptibility of *Sst1^S* mice, we asked whether any other genes found within the

Sst1 locus differ in expression between B6 and B6.*Sst1^S* BMMs. We noted that a homolog of

Sp110 called *Sp140* was also reduced in expression in B6.*Sst1^S* cells compared to B6 cells

(Figure 2A). Immunoblot confirmed that neither untreated nor IFN-γ treated BMMs from

B6.*Sst1^S* mice produce SP140 protein (Figure 2B). We therefore used CRISPR–Cas9 to generate

two independent lines of *Sp140$^{-/-}$* mice on a pure B6 background (Figure 2 – figure supplement

1A-C). Our analysis focused primarily on line 1, which we found lacked expression of SP140

protein (Figure 2B) but retains the production of SP110 protein (Figure 2 – figure supplement

1D). Like *Sp110$^{-/-}$* and *Sst1^S* mice, *Sp140$^{-/-}$* mice are viable, fertile and born at the expected

Mendelian ratios. When infected with *M. tuberculosis*, however, *Sp140$^{-/-}$* mice exhibited high

bacterial burdens in their lungs, similar to B6.*Sst1^S* mice and significantly greater than B6,

Sp110$^{-/-}$ or *Sp140$^{+/-}$* littermate mice at day 28 post-infection (Figure 2C, Figure 2 – figure

supplement 1E). *Sp140$^{-/-}$* lungs at 25 days post-infection with *M. tuberculosis* were more similar

to B6.*Sst1^s* lungs than to B6 or *Sp110$^{-/-}$* lungs (Figure 2D, Figure 2 – figure supplement 2). The

increased susceptibility of *Sp140$^{-/-}$* mice was accompanied by significant weight loss and

shortened survival upon infection with *M. tuberculosis*, again phenocopying the B6.*Sst1^S* mice

(Figure 2E-F). Both of the independent lines of *Sp140$^{-/-}$* mice were similarly susceptible to *M.*

tuberculosis (Figure 2 – figure supplement 1E). We also found that both B6.*Sst1^S* and *Sp140$^{-/-}$*

mice were more susceptible to the intracellular Gram-negative bacterium *Legionella*

pneumophila, as compared to the B6 and *Sp110$^{-/-}$* mice (Figure 2G).

173   An important caveat to the use of CRIPSR–Cas9 to generate *Sp140$^{-/-}$* mice is the

presence of an unknown number of nearly identical *Sp140*-like genes in the *Sst1* locus and non-

localized chromosome 1 genome contigs (that presumably map to the adjacent HSR that remains

unassembled by the mouse genome project). It is possible that the guide RNA we used to disrupt

exon 3 of *Sp140* also disrupted these uncharacterized *Sp140*-like genes. However, it is not clear

if these uncharacterized *Sp140*-like genes give rise to functional proteins. Nevertheless, to

identify potential mutated off-target genes in our *Sp140$^{-/-}$* mice, we amplified exons 2/3 of *Sp140*

and any potential paralogs from genomic DNA and from cDNA derived from *M. tuberculosis*-

infected lungs, and subjected the amplicons to deep sequencing (Figure 2 – figure supplement 3).
Although we found evidence for several edited *Sp140*-like exons in our *Sp140*$^{-/-}$ mice, only one
of these edited off-target genes was found to be detectably expressed from analysis of RNA-seq
data from *M. tuberculosis*-infected lungs, and this off-target appeared to be edited only in one of
our founder lines (Figure 2 – figure supplement 3B). Thus, mutation of *Sp140* itself is the most
parsimonious explanation for susceptibility of our *Sp140*$^{-/-}$ mice, a conclusion further supported
by complementation of the mutation in BMMs (see below, Figure 2 – figure supplement 4).
Collectively our results strongly suggest that the lack expression of *Sp140* in B6.*Sst1*^{S} mice
explains the broad susceptibility of these mice to bacterial infections.

Enhanced type I IFN responses in *Sp140*$^{-/-}$ and B6.*Sst1*^{S} mice. We and others previously
reported that TNFα induces higher levels of type I IFN-induced genes in *Sst1*^{S} BMMs as
compared to B6 BMMs (Bhattacharya et al., 2018; Ji et al., 2019). We also observed higher
levels of *Ifnb* transcripts in the lungs of B6.*Sst1*^{S} mice infected with *M. tuberculosis*, as
compared to infected B6 mice (Ji et al., 2019). Similar to B6.*Sst1*^{S} BMMs, *Sp140*$^{-/-}$ BMMs also
exhibited elevated expression of *Ifnb* and interferon-stimulated genes (ISGs) when stimulated
with TNFα (Figure 2 – figure supplement 4A). Importantly, we were also able to complement
the enhanced IFN phenotype of *Sp140*$^{-/-}$ BMMs by transducing *Sp140*$^{-/-}$ BMMs with a
retrovirus expressing a *Sp140* cDNA driven by a minimal CMV promoter (Figure 2 – figure
supplement 4B). Repression of *Ifnb* by overexpression of *Sp140* in *Sp140*$^{-/-}$ BMMs was
selective, as *Sp140* overexpression did not repress the transcription of *Tnfa* induced by TNFα
(Figure 2 – figure supplement 4B).
When infected with *M. tuberculosis*, the lungs of *Sp140*$^{-/-}$ and B6.*Sst1*^{S} mice also
exhibited higher levels of *Ifnb* transcript as compared to B6, *Sp110*$^{-/-}$ and *Sp140*$^{+/-}$ littermate
mice (Figure 3A). Likewise, during *L. pneumophila* infection, *Sp140*$^{-/-}$ mice expressed more
Ifnb in their lungs, as compared to B6 mice. Importantly, elevated *Ifnb* was evident at 48 hours
post-infection when there is no difference in bacterial burdens between the genotypes, and at 96
hours post-infection, when *Sp140*$^{-/-}$ mice have greater bacterial burdens (Figure 3B).

Infected *Sp140*$^{-/-}$ and B6.*Sst1*^{S} lungs show similar gene expression patterns. We used RNA
sequencing to analyze the global gene expression patterns in *M. tuberculosis*-infected lungs of
B6, *Sp110*$^{-/-}$, *Sp140*$^{-/-}$ and B6.*Sst1*^{S} mice at day 28 post-infection (Figure 4). Principal
component analysis revealed that while there is spread among individual samples, the expression
pattern of *Sp140*$^{-/-}$ and B6.*Sst1*^{S} lungs segregates from the expression pattern in B6 and *Sp110*$^{-/-}$
lungs along the PC1 axis (77% of variance) (Figure 4A). Euclidean distance analysis revealed a
similar pattern, with B6.*Sst1*^{S} and *Sp140*$^{-/-}$ mice clustering together, and away from B6 and
Sp110$^{-/-}$ mice (Figure 4B). At the time point analyzed (28 dpi), both *Sp140*$^{-/-}$ and B6.*Sst1*^{S} mice

exhibit higher bacterial burdens than B6 and $Sp110^{-/-}$ mice (Figure 2C). Thus, the similarity of
the gene expression profile of B6.$Sst1^S$ and $Sp140^{-/-}$ lungs may merely reflect increased
inflammation in these lungs. Alternatively, the increased bacterial burdens may be due to a
similarly enhanced type I IFN response in these mice, which leads to secondary bacterial
outgrowth and inflammation. Therefore, we specifically compared the change in expression of
two subsets of genes: (1) hallmark inflammatory response pathway (Figure 4C) and (2) type I
interferon response genes (Figure 4D). This analysis revealed that B6.$Sst1^S$ and $Sp140^{-/-}$ mice
not only show a similarly increased inflammatory gene signature, as expected, but in addition
showed a similarly increased type I IFN gene signature. Only 269 genes were significantly
differentially expressed (adjusted p-value <0.05) between $Sp140^{-/-}$ and B6.$Sst1^S$ samples,
whereas 1520 genes were significantly differentially expressed between $Sp140^{-/-}$ and B6. Within
the 269 genes differentially expressed between $Sp140^{-/-}$ and B6.$Sst1^S$, 62 were immunoglobulin
genes and 62 were annotated as pseudogenes and most differences are only of modest
significance (Figure 4E). At present, we cannot explain these differences, but since these genes
are not linked to the $Sst1$ locus, and since $Sp110^{-/-}$ mice did not exhibit similar changes, we
suspect these expression differences reflect background differences between the $Sst1^S$ and
$Sp140^{-/-}$ strains. Interestingly, the gene most significantly differentially expressed between
B6.$Sst1^S$ and $Sp140^{-/-}$ mice (i.e., with the smallest adjusted p-value) was $Sp110$ (Figure 4E). This
result is expected, given that $Sp110$ is not expressed in B6.$Sst1^S$ but is retained in our $Sp140^{-/-}$
mice (Figure 2 – figure supplement 1D). Together, these results show that while they are not
identical, the transcriptomes of $Sp140^{-/-}$ and B6.$Sst1^S$ mice greatly overlap during *M.*
tuberculosis infection, and importantly, both strains exhibit a similar type I IFN signature. Given
the susceptibility of B6.$Sst1^S$ mice is due to overproduction of type I IFN (Ji et al., 2019), we
hypothesized that type I IFNs might also mediate the susceptibility of $Sp140^{-/-}$ mice.

Susceptibility of $Sp140^{-/-}$ mice to bacterial infections is dependent on type I IFN signaling.
To determine whether type I IFNs exacerbate *M. tuberculosis* infection of $Sp140^{-/-}$ mice, *M.*
tuberculosis-infected $Sp140^{-/-}$ mice were treated with a blocking antibody against IFNAR1.
Compared to mice that only received isotype control antibody, $Sp140^{-/-}$ mice that received the
anti-IFNAR1 antibody had reduced bacterial burdens in their lungs (Figure 5 – figure
supplement 1). We also generated $Sp140^{-/-}Ifnar^{-/-}$ double-deficient mice and infected them with
M. tuberculosis (Figure 5A-B). Loss of *Ifnar* protected $Sp140^{-/-}$ mice from weight loss (Figure
5A) and reduced bacterial burdens at day 25 post-infection, similar to those seen in B6 mice
(Figure 5B). Furthermore, $Sp140^{-/-}Ifnar^{-/-}$ mice were partially protected from *L. pneumophila*
infection, to a similar degree as B6.$Sst1^SIfnar^{-/-}$ mice (Figure 5C-D). These results show that
similar to B6.$Sst1^S$ mice, type I IFN signaling is responsible for the susceptibility of $Sp140^{-/-}$

mice to *M. tuberculosis*, and partially responsible for the susceptibility of $Sp140^{-/-}$ mice to *L.*
pneumophila.

Discussion

Humans and other vertebrates encounter diverse classes of pathogens, including viruses, bacteria,
fungi and parasites. In response, vertebrate immune systems have evolved stereotypical
responses appropriate for distinct pathogen types. For example, type I IFN-driven immunity is
generally critical for defense against viruses (Schneider et al., 2014; Stetson and Medzhitov,
2006), whereas type II IFN (IFN-γ)-driven immunity mediates resistance to intracellular
pathogens (Crisler and Lenz, 2018). Additionally, IL-1 is important for inducing neutrophil and
other responses against extracellular pathogens (Mantovani et al., 2019), and IL-4/-13 (Type 2
immunity) orchestrates responses to helminths and other parasites (Locksley, 1994). Thus, an
important question is how the immune system generates responses that are appropriate for
resistance to a specific pathogen while repressing inappropriate responses. The alternative
strategy of making all types of responses to all pathogens appears not to be employed, possibly
because it would be too energetically costly, or incur too much inflammatory damage to the host.
Although there is still much to be learned, it appears that negative feedback is essential to
enforce choices between possible types of immune responses. For example, IL-4 and IFN-γ have
long been appreciated to act as reciprocal negative antagonists of each other (Locksley, 1994). In
addition, anti-viral type I IFNs have long been appreciated to negatively regulate IFN-γ and IL-1-
driven anti-bacterial responses (Donovan et al., 2017; Moreira-Teixeira et al., 2018). Although
negative regulation of IFN-γ/IL-1 by type I IFN is likely beneficial to limit immunopathology
during viral infections, $Sst1^S$ mice provide an example of how excessive or inappropriate
negative regulation by type I IFN can also be detrimental during bacterial infections (He et al.,
2013; Ji et al., 2019). In this study, we therefore sought to understand the molecular mechanisms
by which wild-type ($Sst1^R$) mice are able to restrain type I IFNs appropriately during bacterial
infections.

Although the *Sst1* locus was first described in 2005 (Pan et al., 2005), further genetic
analysis of the locus has been hindered by its extreme repetitiveness and the concomitant
difficulty in generating specific loss-of-function mutations in *Sst1*-linked genes. In particular, the
loss of *Sp110 (Ipr1)* has long been proposed to explain the susceptibility of *Sst1* mice to bacterial
infections. However, while we could confirm the loss of *Sp110* expression in $Sst1^S$ mice, $Sp110^-$
$^{/-}$ mice were never generated and thus its essential role in host defense has been unclear. The
advent of CRIPSR/Cas9-based methods of genome engineering allowed us to generate $Sp110^{-/-}$

mice. Unexpectedly, we found $Sp110^{-/-}$ mice were fully resistant to *M. tuberculosis* infection,

and we thus conclude that lack of *Sp110* is not sufficient to explain the $Sst1^S$ phenotype. An

important caveat of genetic studies of the *Sst1* locus is that generating specific gene knockouts is

still nearly impossible in this genetic region, even with CRISPR–Cas9. Indeed, the guide

sequence used to target exon 3 of *Sp110* also targets an unknown number of pseudogene copies

of *Sp110*-like genes located within the unassembled adjacent 'homogenously staining region' of

mouse chromosome 1. Thus, we expect that additional off-target mutations are likely present in

our $Sp110^{-/-}$ mutant mice. However, given that the *Sp110* pseudogenes are not known to be

expressed, we consider it unlikely that collateral mutations would affect our conclusions.

Moreover, any off-target mutations should differ among the three founder mice we analyzed and

are thus unlikely to explain the consistent resistant phenotype we observed in all three founders.

Lastly, since we were able to establish that all the founders at a minimum lack SP110 protein,

additional mutations would not affect our conclusion that *Sp110* is not essential for resistance to

M. tuberculosis.

Given that loss of *Sp110* was not sufficient to explain the susceptibility of $Sst1^S$ mice to

bacterial infections, we considered other explanations. We found that $Sst1^S$ mice also lack

expression of *Sp140*, an *Sst1*-linked homolog of *Sp110*. Our data suggest that deletion of *Sp140*

is sufficient to recapitulate the full $Sst1^S$ phenotype including broad susceptibility to multiple

bacterial infections including *M. tuberculosis* and *L. pneumophila*. Importantly, the susceptibility

of $Sp140^{-/-}$ mice to bacterial infection correlates with an exacerbated type I IFN response, as is

also the case for $Sst1^S$ mice. Likewise, as with $Sst1^S$ mice, the susceptibility of $Sp140^{-/-}$ mice was

rescued by antibody blockade or genetic deletion of the type I IFN receptor (*Ifnar*). We therefore

conclude that loss of *Sp140* likely explains the *Sst1* phenotype. It remains possible that the

additional loss of *Sp110* in $Sst1^S$ mice further exacerbates the $Sst1^S$ phenotype as compared to

$Sp140^{-/-}$ mice. However, in our studies, we did not observe a consistent difference between $Sst1^S$

(i.e., $Sp110^{-/-}Sp140^{-/-}$) mice as compared to our $Sp140^{-/-}$ mice.

Another important caveat to our study is that it remains possible that our $Sp140^{-/-}$ mice

carry additional mutations that contribute to, or even fully explain, their observed phenotype.

This concern is somewhat ameliorated by our analysis of two independent $Sp140^{-/-}$ founders,

both of which exhibited susceptibility to *M. tuberculosis* (Figure 2 – figure supplement 1E). We

confirmed there is normal SP110 protein levels in the spleen of uninfected $Sp140^{-/-}$ mice, and

normal levels of *Sp110* and *Sp100* mRNA in the lungs of *M. tuberculosis*-infected mice. Thus,

collateral loss of SP100 or SP110 is unlikely to explain the phenotype of our $Sp140^{-/-}$ mice. To

address the possibility of mutations in unannotated SP-like genes, we used deep amplicon

sequencing of genomic DNA and cDNA from $Sp140^{-/-}$ mice. We confirmed that both founder

lines harbored distinct off-target mutations. Most of the identified off-target mutations are in
previously unidentified sequences that likely originate from *Sp140* paralogs within the unmapped
HSR, and may, like many of the HSR sequences, be pseudogenes. Most off-target mutated genes
also appear to be expressed at a far lower level than *Sp140* in lungs during *M. tuberculosis*
infection. In one of our *Sp140$^{-/-}$* lines, we identified an off-target mutated *Sp140*-like paralog
that was expressed at detectable levels in the lungs of *M. tuberculosis*-infected mice. This
paralog was 100% identical to *Sp140* in the sequenced region and was only distinguished from
Sp140 itself because it lacked the deletion that was introduced into the edited *Sp140* gene.
Importantly this previously undescribed *Sp140*-like expressed sequence was not mutated in our
second *Sp140$^{-/-}$* line and is thus unlikely to explain resistance to *M. tuberculosis* infection. As an
alternative approach to confirm the phenotype of *Sp140$^{-/-}$* mice is due to loss of *Sp140*, we
overexpressed *Sp140* in *Sp140$^{-/-}$* BMMs. Crucially, we found *Sp140* complements aberrant
elevated *Ifnb1* transcription exhibited by *Sp140$^{-/-}$* BMMs upon TNFα stimulation. Lastly, *Sp110*
and *Sp140* are the only two *Sst1*-linked genes that we were able to find to be differentially
expressed between B6 and B6.*Sst1^{S}* mice, and as discussed above, our genetic studies suggest
little role for the loss of *Sp110*. Thus, while it is formally possible that an edited *Sp140* homolog
that was not identified by our amplicon sequencing contributes to the susceptibility to bacterial
infection and elevated type I IFN in *Sp140$^{-/-}$* mice, the most parsimonious explanation of our
data is that deficiency in *Sp140* accounts for the *Sst1^{S}* phenotype. We expect that future
mechanistic studies will be critical to further confirm this conclusion.

Because *Sp140* is inducible by IFN-γ, our results suggest the existence of a novel
feedback loop by which IFN-γ acts to repress the transcription of type I IFNs via SP140. This
feedback loop appears to be essential for host defense against diverse bacterial pathogens. A
major question that remains is how SP140 acts to repress the type I IFN response. SP140
contains DNA/chromatin-binding domains, such as SAND, PHD and Bromodomains, which
suggests the hypothesis that SP140 functions as a direct transcriptional repressor of type I IFN
genes. However, much more indirect mechanisms are also possible. Recent studies suggest that
hyper type I IFN responses in TNF-stimulated B6.*Sst1^{s}* BMMs derive from aberrant oxidative
stress that activates the kinase JNK and ultimately results in a non-resolving stress response that
promotes necrosis (Bhattacharya et al., 2020; Brownhill, 2020). Interestingly, mouse SP140
localizes to nuclear structures called PML bodies. PML bodies are implicated in a variety of cell
processes such as apoptosis, cell cycle, DNA damage response, senescence, and cell-intrinsic
antiviral responses (Scherer and Stamminger, 2016). Whether or not the repressive effects of
SP140 on type I IFN expression occur via the activity of PML bodies is an important outstanding
question. Another major question is whether or how the repression of type I IFNs by SP140 is

specific for bacterial infections and, if not, whether the presence of SP140 impairs anti-viral
immunity. Lastly, polymorphisms in human SP140 are associated with chronic lymphocytic
leukemia (CLL), Crohn's disease and multiple sclerosis (MS) (Franke et al., 2010; Jostins et al.,
2012; Karaky et al., 2018; Matesanz et al., 2015; Slager et al., 2013). Studies using siRNA and
shRNA-mediated knockdown have also implicated SP140 in the repression of lineage-
inappropriate genes in macrophages (Mehta et al., 2017). Our generation of $Sp140^{-/-}$ mice is
therefore important to permit future studies into these alternative roles of SP140.

Materials and Methods

Mice. All mice were specific pathogen-free, maintained under a 12-hr light-dark cycle (7AM to
7PM), and given a standard chow diet (Harlan irradiated laboratory animal diet) *ad libitum*. All
mice were sex and age-matched at 6-10 weeks old at the beginning of infections. C57BL/6J (B6)
and B6(Cg)-*Ifnar1tm1.2Ees*/J (*Ifnar$^{-/-}$*) were originally purchased from Jackson Laboratories
and subsequently bred at UC Berkeley. B6J.C3-*Sst* $^{C3HeB/FeJ}$Krmn mice (referred to as B6.*Sst1^S*
throughout) were from the colony of I. Kramnik at Boston University and then transferred to UC
Berkeley. CRISPR/Cas9 targeting was performed by pronuclear injection of Cas9 mRNA and
sgRNA into fertilized zygotes from colony-born C57BL/6J mice, essentially as described
previously (Wang et al., 2013). Founder mice were genotyped as described below, and founders
carrying *Sp140* mutations were bred one generation to C57BL/6J to separate modified *Sp140*
haplotypes. Homozygous lines were generated by interbreeding heterozygotes carrying matched
Sp140 haplotypes. *Sp140$^{-/-}$Ifnar$^{-/-}$* were generated by crossing the *Sp140$^{-/-}$* and *Ifnar$^{-/-}$* mice in-
house. All animals used in experiments were bred in-house unless otherwise noted in the figure
legends. All animal experiments complied with the regulatory standards of, and were approved
by, the University of California Berkeley Institutional Animal Care and Use Committee.

Genotyping of *Sp110* alleles. Exon 3 and the surrounding intronic regions were amplified by
PCR using the following primers (all 5' to 3'): Sp110 fwd, CTCTCCGCTCGGTGACTAC, and
rev, CTGCACATGTGACAAGGATCTC. The primer combinations were designed to
distinguish *Sp110* from other *Sp110*-like genes. Primers were used at 200nM in each 20μl
reaction with 1x Dreamtaq Green PCR Master Mix (Thermo Fisher Scientific). Cleaned PCR
products were diluted at 1:10 and sequenced using Sanger sequencing (Elim Biopharm).

Genotyping of *Sp140* alleles. Exon 3 and the surrounding intronic regions were amplified by
bracket PCR using the following primers (all 5′ to 3′): Sp140-1 fwd,
ACGAATAGCAAGCAGGAATGCT, and rev, GGTTCCGGCTGAGCACTTAT. The PCR
products are diluted at 1:10 and 2μl were used as template for the second PCR using the

following primers: Sp140-2 fwd, TGAGGACAGAACTCAGGGAG, and rev,

ACACGCCTTTAATCCCAGCATTT. The primer combinations were designed to distinguish

Sp140 from other *Sp140*-like genes. Primers were used at 200nM in each 20μl reaction with 1x

Dreamtaq Green PCR Master Mix (Thermo Fisher Scientific). Cleaned PCR products were

diluted at 1:10 and sequenced using Sanger sequencing (Elim Biopharm). PCRs were performed

as described above for *Sp110* and sequenced using Sanger sequencing (Elim Biopharm).

***Mycobacterium tuberculosis* infections.** *M. tuberculosis* strain Erdman (gift of S.A. Stanley)

was used for all infections. Frozen stocks of this wild-type strain were made from a single

culture and used for all experiments. Cultures for infection were grown in Middlebrook 7H9

liquid medium supplemented with 10% albumin-dextrose-saline, 0.4% glycerol and 0.05%

Tween-80 for five days at 37°C. Mice were aerosol infected using an inhalation exposure system

(Glas-Col, Terre Haute, IN). A total of 9 ml of diluted culture was loaded into the nebulizer

calibrated to deliver ~20 to 50 bacteria per mouse as confirmed by measurement of colony

forming units (CFUs) in the lungs 1 day following infection. Mice were sacrificed at various

days post-infection (as described in figure legends) to measure CFUs and RNA levels. All but

one lung lobe was homogenized in PBS plus 0.05% Tween-80, and serial dilutions were plated

on 7H11 plates supplemented with 10% oleic acid, albumin, dextrose, catalase (OADC) and

0.5% glycerol. CFUs were counted 21 days after plating. The remaining lobe was used for

histology or for RNA extraction. For histology, the sample was fixed in 10% formalin for at least

48 hours then stored in 70% ethanol. Samples were sent to Histowiz Inc for embedding in wax,

sectioning and staining with hematoxylin and eosin. For survival experiments, mice were

monitored for weight loss and were euthanized when they reached a humane endpoint as

determined by the University of California Berkeley Institutional Animal Care and Use

Committee.

***Legionella pneumophila* infections.** Infections were performed using *L. pneumophila* strain

JR32 Δ*flaA* (gift of D.S. Zamboni) as previously described (Goncalves et al., 2019; Mascarenhas

et al., 2015). Briefly, frozen cultures were streaked out on to BCYE plates to obtain single

colonies. A single colony was chosen and streaked on to a new BCYE plate to obtain a 1cm by

1cm square bacterial lawn, and incubated for 2 days at 37°C. The patch was solubilized in

autoclaved MilliQ water and the optical density was measured at 600nm. Culture was diluted to

2.5 x10^6 bacteria/ml in sterile PBS. The mice were first anesthetized with ketamine and xylazine

(90 mg/kg and 5 mg/kg, respectively) by intraperitoneal injection then infected intranasally with

40 μL with PBS containing a final dilution of 1×10^5 bacteria per mouse. For enumerating of

CFU, the lungs were harvested and homogenized in 5 mL of autoclaved MilliQ water for 30

seconds, using a tissue homogenizer. Lung homogenates were diluted in autoclaved MilliQ water

and plated on BCYE agar plates. CFU was enumerated after plates were incubated for 4 days at

37°C.

Bone marrow-derived macrophages (BMMs) and TNF-treatment. Bone marrow was

harvested from mouse femurs and tibias, and cells were differentiated by culture on non-tissue

culture-treated plates in RPMI supplemented with supernatant from 3T3-MCSF cells (gift of B.

Beutler), 10% fetal bovine serum (FBS), 2mM glutamine, 100 U/ml streptomycin and 100 μg/ml

penicillin in a humidified incubator (37°C, 5%CO2). BMMs were harvested six days after

plating and frozen in 95% FBS and 5% DMSO. For *in vitro* experiments, BMMs were thawed

into media as described above for 4 hours in a humidified 37°C incubator. Adherent cells were

washed with PBS, counted and replated at $1.2x10^6 \sim 1.5x10^6$ cells/well in a TC-treated 6-well

plate. Cells were treated with 10 ng/ml recombinant mouse TNFα (410-TRNC-010, R&D

systems) diluted in the media as described above.

Quantitative/conventional RT-PCR. Total RNA from BMMs was extracted using E.Z.N.A.

Total RNA Kit I (Omega Bio-tek) according to manufacturer specifications. Total RNA from

infected tissues was extracted by homogenizing in TRIzol reagent (Life technologies) then

mixing thoroughly with chloroform, both done under BSL3 conditions. Samples were then

removed from the BSL3 facility and transferred to fresh tubes under BSL2 conditions. Aqueous

phase was separated by centrifugation and RNA was further purified using the E.Z.N.A. Total

RNA Kit I (Omega Bio-tek). Equal amounts of RNA from each sample were treated with DNase

(RQ1, Promega) and cDNA was made using Superscript III (Invitrogen). Complementary cDNA

reactions were primed with poly(dT) for the measurement of mature transcripts. For experiments

with multiple time points, macrophage samples were frozen in the RLT buffer (Qiagen) and

infected tissue samples in RNA*later*™ solution (Invitrogen) and processed to RNA at the same

time. Quantitative PCR was performed using QuantiStudio 5 Real-Time PCR System (Applied

Biosystems) with Power Sybr Green PCR Master Mix (Thermo Fisher Scientific) according to

manufacturer specifications. Transcript levels were normalized to housekeeping genes *Rps17,*

Actb and *Oaz1* unless otherwise specified. The following primers were used in this study. *Rps17*

sense: CGCCATTATCCC CAGCAAG; *Rps17* antisense: TGTCGGGATCCACCTCAATG;

Oaz1 sense: GTG GTG GCC TCT ACA TCG AG; *Oaz1* antisense: AGC AGA TGA AAA CGT

GGT CAG; *Actb* sense: CGC AGC CAC TGT CGA GTC; *Actb* antisense: CCT TCT GAC CCA

TTC CCA CC; *Ifnb* sense: GTCCTCAACTGCTCTCCACT; *Ifnb* antisense:

CCTGCAACCACCACTCATTC; *Gbp4* sense: TGAGTACCTGGAGAATGCCCT; *Gbp4*

antisense: TGGCCGAATTGGATGCTTGG; *Ifit3* sense: AGCCCACACCCAGCTTTT; *Ifit3*

antisense: CAGAGATTCCCGGTTGACCT. *Tnfa* sense: TCTTCTCATTCCTGCTTGTGG;

Tnfa antisense: GGTCTGGGCCATAGAACTGA. Conventional RT-PCR shown in Figure 2A

was done using the following primers. Sense: GTCCCTTGGAGTCTGTGTAGG; antisense:
CATCCTGGGGCTCTTGTCTTG.

Immunoblot. Samples were lysed in RIPA buffer with protease inhibitor cocktail (Roche) to
obtain total protein lysate and were clarified by spinning at ~16,000×g for 30 min at 4°C.
Clarified lysates were analyzed with Pierce BCA protein assay kit (Thermo Fisher Scientific)
according to manufacturer specification and diluted to the same concentration and denatured
with SDS-loading buffer. Samples were separated on NuPAGE Bis–Tris 4% to 12% gradient
gels (Thermo Fisher Scientific) following the manufacturer's protocol. Gels were transferred
onto ImmobilonFL PVDF membranes at 35 V for 90 min and blocked with Odyssey blocking
buffer (Li-Cor). Proteins were detected on a Li-Cor Odyssey Blot Imager using the following
primary and secondary antibodies. Rabbit anti-SP110 or SP140 serums were produced by
Covance and used at 1:1000 dilution. Hybridoma cells expressing monoclonal anti-SP110
antibody were a gift of I. Kramnik. Antibodies were produced in-house as previously described
(Ji et al., 2019) and used at 100 ng/ml. Alexa Fluor 680-conjugated secondary antibodies
(Invitrogen) were used at 0.4 mg/ml.

RNA sequencing and analysis. Total RNA was isolated as described above. Illumina-
compatible libraries were generated by the University of California, Berkeley, QB3 Vincent J.
Coates Genomics Sequencing Laboratory. PolyA selection was performed to deplete rRNA.
Libraries were constructed using Kapa Biosystem library preparation kits. The libraries were
multiplexed and sequenced using one flow cell on Novaseq 6000 (Illumina) as 50bp paired-end
reads. Base calling was performed using bcl2fastq2 v2.20. The sequences were aligned to mm10
genome using Kallisto v.0.46.0 using standard parameters (Pimentel et al., 2017) and analyzed
using Deseq2 (Love et al., 2014) and DEVis packages (Price et al., 2019). For Deseq2 and
DEVis analysis, all raw counts were incremented by 1 to avoid excluding genes due to division
by 0 in the normalization process.

Antibody-mediated neutralization. Mice were given anti-IFNAR1 antibody or isotype control
once every 2 days, starting 7 days post-infection. All treatments were delivered by
intraperitoneal injection. Mouse anti-mouse IFNAR1 (MAR1-5A3) and isotype control (GIR208,
mouse anti-human IFNGR-α chain) were purchased from Leinco Technologies Inc. For
injections antibody stocks were diluted in sterile PBS and each mouse received 500 μg per
injection.

Amplicon sequencing and analysis. Amplicons comprising the 5'intron of exon 3 of *Sp140* and
the end of exon 3 were amplified from crude DNA from ear clips of Sp140KO-1 mice (sense:
TCATATAACCCATAAATCCATCATGACA; antisense:

CCATTTAGGAAGAAGTGTTTTAGAGTCT) with PrimeStar PCR components (Takara,

R010b) for 18 cycles according to manufacturer specifications, then diluted 50-fold and

barcoded for an additional 18 cycles with Illumina-compatible sequencing adaptors. Amplicons

of *Sp140* exon 3 (sense: AATATCAAGAAACATGTAAGAACCTGGT; antisense:

CCATTTAGGAAGAAGTGTTTTAGAGTCT) and exon 2-3 (sense:

GCAGAAGTTTCAGGAATATCAAGAAACATGTAAG; antisense:

ACTTCTTCTGTACATTGCTGAGGATGT) were amplified from cDNA generated from lungs

infected with *M. tuberculosis* for 25 cycles with PrimeStar before barcoding. Libraries were

generated by the University of California, Berkeley, QB3 Vincent J. Coates Genomics

Sequencing Laboratory, and were multiplexed and sequenced on an Illumina Miseq platform

with v2 chemistry and 300 bp single-end reads for DNA amplicons, and Illumina Miseq Nano

platform with v3 chemistry for 300bp single end reads for cDNA amplicons. Reads were aligned

with Burrows-Wheeler Aligner (BWA-MEM) with default parameters (Li, *et al*, 2009; Li, 2013)

to chromosome 1 and non-localized genome contigs of the *Mus musculus* genome (assembly

mm10) as well as the *Sp140* gene and transcript X1(XM_030255396.1), converted to BAM files

with samtools (Li, 2009), and visualized in IGV 2.8. Subsets of reads were extracted from

alignment files using the Seqkit toolkit (Shen, 2016).

Retroviral transduction of BMMs. Self-inactivating pTMGP vector (SINV) with either a

minimal CMV promoter driving Sp140 or a minimal CMV promoter and 4 Gal4 binding sites

driving mNeonGreen, and the reporter mAmetrine driven by a PGK promoter were cloned using

Infusion (638910, Takara). pTGMP was a gift from Scott Lowe (Addgene plasmid # 32716).

Virus was harvested from GP-2 cells transfected with SINV vectors and VSV-G grown in

DMEM supplemented with 30% FBS and 2mM glutamine, 100 U/ml streptomycin and 100

µg/ml penicillin (adapted from protocols described in Schmidt *et al*, 2015). Harvested virus was

concentrated 100-fold by ultracentrifugation in RPMI before storage at -80 °C. Virus was thawed

and titrated on bone marrow to optimize transduction efficiency. Bone marrow was harvested as

described above and the entirety of the bone marrow was plated in a non-TC 15 cm plate. The

next day, bone marrow was harvested and transduced with SINV virus on plates coated with 10

µg/cm^2 Retronectin (T100, Takara) for 1.5-2 hrs at 650 xg and 37°C. After 2 days of additional

culture, media was replenished, then transduced bone marrow was cultured for 3 additional days

before sorting. Sorted transduced macrophages were stimulated with 5 ng/mL recombinant

murine IFN-γ (575304, Biolegend) 12-14 hours before stimulation with 10 ng/mL recombinant

TNFα as described above for 4 hours (FBS used in these experiments was from Omega, LOT

721017, CAT# FB-12). RNA isolation and RT-qPCR were performed as described above.

Statistical analysis. All data were analyzed with Mann-Whitney test unless otherwise noted.
Tests were run using GraphPad Prism 5. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$. All error bars
are s.e. Figures show exact p values for $p > 0.0005$.

Data accession. RNA-seq data is available at GEO, accession number GSE166114. Amplicon
sequencing data is available at the SRA, BioProject accession number PRJNA698382.

Acknowledgements: We thank the Stanley and Cox laboratories for support with *M.*
tuberculosis experiments, L. Flores, P. Dietzen and R. Chavez for technical assistance, and
members of the Vance, Barton, Cox, Stanley, and Portnoy labs for advice and discussions.

Author contributions: Conceptualization–D.X.J., K.H.D., A.Y.L., R.E.V.; Methodology–
D.X.J., K.C.W., D.I.K., S.R.M., A.L., K.J.C., A.Y.L., D.S.Z.; Resources–I.K., D.A.P., R.E.V.;
Data curation–D.X.J.; Writing – original draft preparation–D.X.J., R.E.V.; Writing – review &
editing–D.X.J., K.C.W., D.I.K., K.H.D., R.E.V.; Supervision–D.A.P., K.H.D., R.E.V.; Funding
acquisition–D.A.P., K.H.D., R.E.V.

Funding: R.E.V. is supported by an Investigator Award from the Howard Hughes Medical
Institute. This work was also supported by NIH grants R37AI075039 (R.E.V.), P01AI066302
(R.E.V. and D.A.P.). R.E.V. and K.H.D. were Burroughs Wellcome Fund Investigators in the
Pathogenesis of Infectious Disease.

Competing Interests: R.E.V. consults for Ventus Therapeutics.

Ethics Statement: Animal studies were approved by the UC Berkeley Animal Care and Use
Committee.

 Figure Legends

Figure 1. *Sp110⁻/⁻* mice are not susceptible to *M. tuberculosis* infections. (A) BMDMs were
treated with 10 U/ml of IFNγ for 24 hours and cells were lysed with RIPA buffer. Five μg of
total protein was loaded on each lane, and immunoblot was performed with respective antibodies
as shown. Molecular weight standards are shown on the left of each blot in kDa. Individual
membranes were imaged separately. Three independent lines of *Sp100⁻/⁻* mice were analyzed
(denoted lines 61, 65, and 71). **(B-D)**, Lungs of mice infected with *M. tuberculosis* were stained
with hematoxylin and eosin (H&E) for histology **(B)**, measured for CFU at 25 days post-
infection (Mann-Whitney test) **(C)** or, monitored for survival **(D)**. All except B6 mice were bred
in-house, and combined results from the three independent *Sp110⁻/⁻* lines are shown.
Representative of 2 experiments **(B, D)**; combined results of 3 infections **(C)**. *, $P \leq 0.05$; **, P
≤ 0.01; ***, $P \leq 0.005$.

Figure 2. *Sp140⁻/⁻* mice are susceptible to bacterial pathogens. (A) RT-PCR of cDNA from
BMMs of the indicated genotypes. Red arrow indicates band corresponding to a portion of
Sp140, verified by sequencing. **(B)** Immunoblot of lysates from *Sp140⁻/⁻* and WT BMMs treated
with 10 U/ml of recombinant mouse IFNγ for 24 hours. Equal amounts of protein were loaded
for immunoblot with anti-SP140 antibody. **(C-F)** Mice were infected with *M. tuberculosis* and
measured for lung CFU at 28 days post-infection **(C)** body weight over time **(E)**, and survival
(F). Statistics in **(E)** shows comparison to B6 at day 28, and data are from 10 B6, 11 B6.Sst1s,
and *Sp110⁻/⁻*, 14 *Sp140⁻/⁻*, and 6 *Sp140⁺/⁻* mice. **(D)** H&E staining of lungs at 25 days post-
infection with *M. tuberculosis*. **(G)** Mice were infected with *L. pneumophila* and lungs were
measured for CFU at 96 hours post-infection. All mice were bred in-house, *Sp140⁻/⁻* and
Sp140⁺/⁻ were littermates **(C-F)**. **C**, **E**, and **G** are combined results of two independent
infections. **A-D** shows representative analysis of one *Sp140⁻/⁻* line (line 1), whereas **F-G**
includes a mixture of both line 1 and 2. Results of infection of both lines with *M. tuberculosis* is
shown in Figure S2E. **(C, E, F, G)** Mann-Whitney test. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.005$.

Figure 3. *Sp140⁻/⁻* mice have elevated *Ifnb* transcripts during bacterial infection. (A) Mice
were infected with *M. tuberculosis* and at 28 days post-infection lungs were processed for total
RNA, which were used for RT-qPCR. Combined results of 2 independent experiments. **(B)** Mice
were infected with *L. pneumophila* and RT-qPCR was performed on lungs collected at indicated
times. Combined results of 2 independent infections. All mice were bred in-house, *Sp140⁻/⁻* and
Sp140⁺/⁻ were littermates. **(A-B)** Mann-Whitney test. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.005$.

Figure 4. Global gene expression analysis of *Sp110⁻/⁻*, *Sp140⁻/⁻* and B6.*Sst1ˢ* lungs after *M.*
***tuberculosis* infection. (A)** PCA or **(B)** Euclidean distance analysis of all the samples. **(C-D)**

heatmaps of gene expression in $\log_2$-fold change from B6. Genes shown are those significantly

different between $Sp140^{-/-}$ and B6: (**C**) GSEA Hallmark inflammatory response; and (**D**) GO

type I IFN response genes. (**E**) volcano plot comparing $Sp140^{-/-}$ to B6.$Sst1^S$ expression. Dots in

red are 2-fold differentially expressed with adjusted *p*-value ≤ 0.05.

Figure 5. Susceptibility of $Sp140^{-/-}$ to *M. tuberculosis* and *L. pneumophila* is dependent on

type I IFN signaling. (**A-B**) mice were infected with *M. tuberculosis* and measured for body

weight (**A**) and bacterial burdens at day 25 (**B**). Statistics in **A** show comparison to B6; data are

from 9 B6, and 13 $Sp140^{-/-}$ and $Sp140^{-/-}$ *Ifnar*$^{-/-}$ mice. Combined results of 2 experiments. (**C-**

D) bacteria burden in *L. pneumophila*-infected mice at 96 hours. Combined results of 2

experiments. All mice were bred in-house (**A-B**, **D**); all but B6 were bred in-house (**C**). Mann-

Whitney test (**A-D**). *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.005$.

Figure 1 − figure supplement 1. CRISPR–Cas9 targeting strategy for $Sp110^{-/-}$ mice. (**A**)

595   Mouse *Sp110* gene. Guide RNA sequence for CRISPR–Cas9 targeting and protospacer-adjacent

596   motif (PAM) are indicated. (**B-D**) *Sp110* locus in wildtype (WT) and three independent lines.

597   Homozygotes of 2 lines identified by sequencing (**B-C**), and heterozygote of the 3rd line by PCR

598   products separated on an agarose gel (**D**). Arrow indicates the mutant band.

Figure 2 − figure supplement 1. CRISPR–Cas9 targeting strategy for $Sp140^{-/-}$ and

validation of founders. (**A**) Mouse *Sp140* gene. Guide RNA sequence for CRISPR–Cas9

targeting and protospacer-adjacent motif (PAM) are indicated. (**B-C**) *Sp140* locus in wildtype

(WT) and 2 independent founders of $Sp140^{-/-}$ validated by sequencing. (**D**) Immunoblot for

SP110 using BMMs from mice of the indicated genotypes. Intervening lanes have been removed

for clarity (indicated by line in the image). (**E**) *M. tuberculosis*-infected mice were harvested for

CFU at 25 days post-infection. Empty and filled triangles indicate the two independent lines of

$Sp140^{-/-}$ used in this infection. All mice were bred in-house and $Sp140^{+/-}$ were littermates with

$Sp140^{-/-}$ line 2. Mann-Whitney test, *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.005$.

Figure 2 − figure supplement 2. Histology of lungs from B6, B6.$Sst1^s$, $Sp110^{-/-}$, $Sp140^{-/-}$

609   **mice after infection with *M. tuberculosis*.** H&E staining of entire lung sections from mice of

610   indicated genotypes at 25 days post-infection with *M. tuberculosis.* Black squares denote

611   sections shown in Figure 2C. Each image represents a lung section from a different mouse.

612   Borders in background color have been added around each image. Scale bar applies to all

613   images.

Figure 2 − figure supplement 3. Characterization of off-targets in $Sp140^{-/-}$ mice. (**A**)

Schematic of amplicon sequencing strategy for *Sp140* and *Sp140* homologs. (B) Summary of

edited *Sp140* homologs from amplicon sequencing and RNA-seq analysis. SNPs are denoted

based on the *Sp140* X1 transcript. Expression level was roughly estimated from read counts.

Three B6 and 2 *Sp140*$^{-/-}$ mice from each founder line were used as biological replicates for

Sp140 exon 2/3 amplicon sequencing from cDNA, 2 mice per genotype were used for *Sp140*

exon 3 amplicon sequencing from cDNA, and 1 mouse per genotype was used for *Sp140* exon 3

amplicon sequencing from DNA.

Figure 2 – figure supplement 4. Complementation of hyper type I IFN responses in *Sp140*$^{-}$

$^{/-}$ **BMMs.** (**A**) BMMs were left untreated or treated with TNF-α for 24 hours. Total RNA was

used for RT-qPCR. Averages of technical duplicates for one biological replicate are shown. Data

is representative of two independent experiments. (**B**) RT-qPCR of *Sp140*$^{-/-}$ BMMs transduced

with either control SINV-minCMV-GAL4-mNeonGreen (SINV-mNeonGreen) or SINV-

minCMV-Sp140 (SINV-Sp140), primed with 5 ng/mL IFN-γ for 14 hours and treated with 10

ng/mL TNF-α for 4 hours. *, $p \leq 0.05$ calculated with an unpaired t-test with Welch's

correction. Data are representative of two independent experiments.

Figure 5 – figure supplement 1. Antibody blockade of IFNAR1 reduces bacterial burden in

***Sp140*$^{-/-}$ mice during *M. tuberculosis* infection.** Mice were infected with *M. tuberculosis* and

treated with either IFNAR1-blocking antibody or isotype control starting 7 days post-infection.

At 25 days post-infection lungs were harvested to enumerate CFU. Results of one experiment.

All mice were bred in-house. Mann-Whitney test, *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.005$.

References

Anderson, M.S., and Su, M.A. (2016). AIRE expands: new roles in immune tolerance and beyond. Nat Rev Immunol *16*, 247-258.

Auerbuch, V., Brockstedt, D.G., Meyer-Morse, N., O'Riordan, M., and Portnoy, D.A. (2004). Mice lacking the type I interferon receptor are resistant to Listeria monocytogenes. J Exp Med *200*, 527-533.

Bhattacharya, B., Chattrerjee, S., Berland, R., Pichugin, A.V., Gao, Y., Connor, J., Ivanov, A., Yan, B.S., Kobzik, L., and Kramnik, I. (2018). Increased susceptibility to intracellular bacteria and necrotic inflammation driven by a dysregulated macrophage response to TNF. bioRxiv.

Bottomley, M.J., Collard, M.W., Huggenvik, J.I., Liu, Z., Gibson, T.J., and Sattler, M. (2001). The SAND domain structure defines a novel DNA-binding fold in transcriptional regulation. Nat Struct Biol *8*, 626-633.

Boxx, G.M., and Cheng, G. (2016). The Roles of Type I Interferon in Bacterial Infection. Cell Host Microbe *19*, 760-769.

Boyartchuk, V., Rojas, M., Yan, B.S., Jobe, O., Hurt, N., Dorfman, D.M., Higgins, D.E., Dietrich, W.F., and Kramnik, I. (2004). The host resistance locus sst1 controls innate immunity to Listeria monocytogenes infection in immunodeficient mice. J Immunol *173*, 5112-5120.

Carrero, J.A., Calderon, B., and Unanue, E.R. (2004). Type I interferon sensitizes lymphocytes to apoptosis and reduces resistance to Listeria infection. J Exp Med *200*, 535-540.

Chang, S.Y., Chen, M.L., Lee, M.R., Liang, Y.C., Lu, T.P., Wang, J.Y., and Yan, B.S. (2018). SP110 Polymorphisms Are Genetic Markers for Vulnerability to Latent and Active Tuberculosis Infection in Taiwan. Dis Markers *2018*, 4687380.

Crisler, W.J., and Lenz, L.L. (2018). Crosstalk between type I and II interferons in regulation of myeloid cell responses during bacterial infection. Curr Opin Immunol *54*, 35-41.

Donovan, M.L., Schultz, T.E., Duke, T.J., and Blumenthal, A. (2017). Type I Interferons in the Pathogenesis of Tuberculosis: Molecular Drivers and Immunological Consequences. Front Immunol *8*, 1633.

Dorhoi, A., Yeremeev, V., Nouailles, G., Weiner, J., 3rd, Jorg, S., Heinemann, E., Oberbeck-Muller, D., Knaul, J.K., Vogelzang, A., Reece, S.T.*, et al.* (2014). Type I IFN signaling triggers immunopathology in tuberculosis-susceptible mice by modulating lung phagocyte dynamics. Eur J Immunol *44*, 2380-2393.

Fox, G.J., Sy, D.N., Nhung, N.V., Yu, B., Ellis, M.K., Van Hung, N., Cuong, N.K., Thi Lien, L., Marks, G.B., Saunders, B.M.*, et al.* (2014). Polymorphisms of SP110 are associated with both pulmonary and extra-pulmonary tuberculosis among the Vietnamese. PLoS One *9*, e99496.

Franke, A., McGovern, D.P., Barrett, J.C., Wang, K., Radford-Smith, G.L., Ahmad, T., Lees, C.W., Balschun, T., Lee, J., Roberts, R.*, et al.* (2010). Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. Nat Genet *42*, 1118-1125.

Goncalves, A.V., Margolis, S.R., Quirino, G.F.S., Mascarenhas, D.P.A., Rauch, I., Nichols, R.D., Ansaldo, E., Fontana, M.F., Vance, R.E., and Zamboni, D.S. (2019). Gasdermin-D and Caspase-7 are the key Caspase-1/8 substrates downstream of the NAIP5/NLRC4 inflammasome required for restriction of Legionella pneumophila. PLoS pathogens *15*, e1007886.

He, X., Berland, R., Mekasha, S., Christensen, T.G., Alroy, J., Kramnik, I., and Ingalls, R.R.
(2013). The sst1 resistance locus regulates evasion of type I interferon signaling by
Chlamydia pneumoniae as a disease tolerance mechanism. PLoS Pathog 9, e1003569.
Ji, D.X., Yamashiro, L.H., Chen, K.J., Mukaida, N., Kramnik, I., Darwin, K.H., and Vance, R.E.
(2019). Type I interferon-driven susceptibility to Mycobacterium tuberculosis is mediated
by IL-1Ra. Nat Microbiol 4, 2128-2135.
Jostins, L., Ripke, S., Weersma, R.K., Duerr, R.H., McGovern, D.P., Hui, K.Y., Lee, J.C.,
Schumm, L.P., Sharma, Y., Anderson, C.A., et al. (2012). Host-microbe interactions have
shaped the genetic architecture of inflammatory bowel disease. Nature 491, 119-124.
Karaky, M., Fedetz, M., Potenciano, V., Andres-Leon, E., Codina, A.E., Barrionuevo, C., Alcina,
A., and Matesanz, F. (2018). SP140 regulates the expression of immune-related genes
associated with multiple sclerosis and other autoimmune diseases by NF-kappaB
inhibition. Hum Mol Genet 27, 4012-4023.
Lei, X., Zhu, H., Zha, L., and Wang, Y. (2012). SP110 gene polymorphisms and tuberculosis
susceptibility: a systematic review and meta-analysis based on 10 624 subjects. Infect
Genet Evol 12, 1473-1480.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R;
1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map
format and SAMtools. Bioinformatics 25(16):2078-9.
Locksley, R.M. (1994). Th2 cells: help for helminths. J Exp Med 179, 1405-1407.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 15, 550.
Mantovani, A., Dinarello, C.A., Molgora, M., and Garlanda, C. (2019). Interleukin-1 and Related
Cytokines in the Regulation of Inflammation and Immunity. Immunity 50, 778-795.
Mascarenhas, D.P., Pereira, M.S., Manin, G.Z., Hori, J.I., and Zamboni, D.S. (2015). Interleukin
1 receptor-driven neutrophil recruitment accounts to MyD88-dependent pulmonary
clearance of legionella pneumophila infection in vivo. J Infect Dis 211, 322-330.
Matesanz, F., Potenciano, V., Fedetz, M., Ramos-Mozo, P., Abad-Grau Mdel, M., Karaky, M.,
Barrionuevo, C., Izquierdo, G., Ruiz-Pena, J.L., Garcia-Sanchez, M.I., et al. (2015). A
functional variant that affects exon-skipping and protein expression of SP140 as genetic
mechanism predisposing to multiple sclerosis. Hum Mol Genet 24, 5619-5627.
Mayer-Barber, K.D., Andrade, B.B., Oland, S.D., Amaral, E.P., Barber, D.L., Gonzales, J.,
Derrick, S.C., Shi, R., Kumar, N.P., Wei, W., et al. (2014). Host-directed therapy of
tuberculosis based on interleukin-1 and type I interferon crosstalk. Nature 511, 99-103.
McNab, F., Mayer-Barber, K., Sher, A., Wack, A., and O'Garra, A. (2015). Type I interferons in
infectious disease. Nat Rev Immunol 15, 87-103.
Mehta, S., Cronkite, D.A., Basavappa, M., Saunders, T.L., Adiliaghdam, F., Amatullah, H.,
Morrison, S.A., Pagan, J.D., Anthony, R.M., Tonnerre, P., et al. (2017). Maintenance of
macrophage transcriptional programs and intestinal homeostasis by epigenetic reader
SP140. Sci Immunol 2.
Moreira-Teixeira, L., Mayer-Barber, K., Sher, A., and O'Garra, A. (2018). Type I interferons in
tuberculosis: Foe and occasionally friend. J Exp Med 215, 1273-1285.
O'Connell, R.M., Saha, S.K., Vaidya, S.A., Bruhn, K.W., Miranda, G.A., Zarnegar, B., Perry,
A.K., Nguyen, B.O., Lane, T.F., Taniguchi, T., et al. (2004). Type I interferon production
enhances susceptibility to Listeria monocytogenes infection. J Exp Med 200, 437-445.

Pan, H., Yan, B.S., Rojas, M., Shebzukhov, Y.V., Zhou, H., Kobzik, L., Higgins, D.E., Daly,
M.J., Bloom, B.R., and Kramnik, I. (2005). Ipr1 gene mediates innate immunity to
tuberculosis. Nature *434*, 767-772.
Perniola, R., and Musco, G. (2014). The biophysical and biochemical properties of the
autoimmune regulator (AIRE) protein. Biochim Biophys Acta *1842*, 326-337.
Pichugin, A.V., Yan, B.S., Sloutsky, A., Kobzik, L., and Kramnik, I. (2009). Dominant role of
the sst1 locus in pathogenesis of necrotizing lung granulomas during chronic tuberculosis
infection and reactivation in genetically resistant hosts. Am J Pathol *174*, 2190-2201.
Pilla-Moffett, D., Barber, M.F., Taylor, G.A., and Coers, J. (2016). Interferon-Inducible GTPases
in Host Resistance, Inflammation and Disease. J Mol Biol *428*, 3495-3513.
Pimentel, H., Bray, N.L., Puente, S., Melsted, P., and Pachter, L. (2017). Differential analysis of
RNA-seq incorporating quantification uncertainty. Nat Methods *14*, 687-690.
Png, E., Alisjahbana, B., Sahiratmadja, E., Marzuki, S., Nelwan, R., Adnan, I., van de Vosse, E.,
Hibberd, M., van Crevel, R., Ottenhoff, T.H.*, et al.* (2012). Polymorphisms in SP110 are
not associated with pulmonary tuberculosis in Indonesians. Infect Genet Evol *12*, 1319-
1323.
Price, A., Caciula, A., Guo, C., Lee, B., Morrison, J., Rasmussen, A., Lipkin, W.I., and Jain, K.
(2019). DEvis: an R package for aggregation and visualization of differential expression
data. BMC Bioinformatics *20*, 110.
Rayamajhi, M., Humann, J., Penheiter, K., Andreasen, K., and Lenz, L.L. (2010). Induction of
IFN-alphabeta enables Listeria monocytogenes to suppress macrophage activation by
IFN-gamma. J Exp Med *207*, 327-337.
Roscioli, T., Cliffe, S.T., Bloch, D.B., Bell, C.G., Mullan, G., Taylor, P.J., Sarris, M., Wang, J.,
Donald, J.A., Kirk, E.P.*, et al.* (2006). Mutations in the gene encoding the PML nuclear
body protein Sp110 are associated with immunodeficiency and hepatic veno-occlusive
disease. Nat Genet *38*, 620-622.
Scherer, M., and Stamminger, T. (2016). Emerging Role of PML Nuclear Bodies in Innate
Immune Signaling. J Virol *90*, 5850-5854.
Schneider, W.M., Chevillotte, M.D., and Rice, C.M. (2014). Interferon-stimulated genes: a
complex web of host defenses. Annu Rev Immunol *32*, 513-545.
Slager, S.L., Caporaso, N.E., de Sanjose, S., and Goldin, L.R. (2013). Genetic susceptibility to
chronic lymphocytic leukemia. Semin Hematol *50*, 296-302.
Stetson, D.B., and Medzhitov, R. (2006). Type I interferons in host defense. Immunity *25*, 373-
381.
Thye, T., Browne, E.N., Chinbuah, M.A., Gyapong, J., Osei, I., Owusu-Dabo, E., Niemann, S.,
Rusch-Gerdes, S., Horstmann, R.D., and Meyer, C.G. (2006). No associations of human
pulmonary tuberculosis with Sp110 variants. J Med Genet *43*, e32.
Tosh, K., Campbell, S.J., Fielding, K., Sillah, J., Bah, B., Gustafson, P., Manneh, K., Lisse, I.,
Sirugo, G., Bennett, S.*, et al.* (2006). Variants in the SP110 gene are associated with
genetic susceptibility to tuberculosis in West Africa. Proc Natl Acad Sci U S A *103*,
10364-10368.
Wang, H., Yang, H., Shivalila, C.S., Dawlaty, M.M., Cheng, A.W., Zhang, F., and Jaenisch, R.
(2013). One-step generation of mice carrying mutations in multiple genes by
CRISPR/Cas-mediated genome engineering. Cell *153*, 910-918.

Weichenhan, D., Kunze, B., Winking, H., van Geel, M., Osoegawa, K., de Jong, P.J., and Traut,
W. (2001). Source and component genes of a 6-200 Mb gene cluster in the house mouse.
Mamm Genome *12*, 590-594.
Zhang, S., Wang, X.B., Han, Y.D., Wang, C., Zhou, Y., and Zheng, F. (2017). Certain
Polymorphisms in SP110 Gene Confer Susceptibility to Tuberculosis: A Comprehensive
Review and Updated Meta-Analysis. Yonsei Med J *58*, 165-173.
779

Figure 1. *Sp110⁻/⁻* **mice are not susceptible to** *M. tuberculosis* **infections.** (**A**) BMDMs were treated with 10 U/ml of IFNγ for 24 hours and cells were lysed with RIPA buffer. Five µg of total protein was loaded on each lane, and immunoblot was performed with respective antibodies as shown. Molecular weight standards are shown on the left of each blot in kDa. Individual membranes were imaged separately. Three independent lines of *Sp100⁻/⁻* mice were analyzed (denoted lines 61, 65, and 71). (**B-D**), Lungs of mice infected with *M. tuberculosis* were stained with hematoxylin and eosin (H&E) for histology (**B**), measured for CFU at 25 days post-infection (Mann-Whitney test) (**C**) or, monitored for survival (**D**). All except B6 mice were bred in-house, and combined results from the three independent *Sp110⁻/⁻* lines are shown. Representative of 2 experiments (**B**, **D**); combined results of 3 infections (**C**). *, $p ≤ 0.05$; **, $p ≤ 0.01$; ***, $p ≤ 0.005$.

Figure 2. *Sp140⁻/⁻* mice are susceptible to bacterial pathogens. (**A**) RT-PCR of cDNA from BMMs of the indicated genotypes. Red arrow indicates band corresponding to a portion of Sp140, verified by sequencing. (**B**) Immunoblot of lysates from *Sp140⁻/⁻* and wildtype BMMs treated with 10 U/ml of recombinant mouse IFNγ for 24 hours. Equal amounts of protein were loaded for immunoblot with anti-SP140 antibody. (**C-F**) Mice were infected with *M. tuberculosis* and measured for lung CFU at 28 days post-infection (**C**), body weight over time (**E**), and survival (**F**). Statistics in (**E**) shows comparison to B6 at day 28, and data are from 10 B6, 11 B6.*Sst1ˢ* and *Sp110⁻/⁻*, 14 *Sp140⁻/⁻*, and 6 *Sp140⁺/⁻* mice. (**D**) H&E staining of lungs at 25 days post-infection with *M. tuberculosis*. (**G**) Mice were infected with *L. pneumophila* and lungs were measured for CFU at 96 hours post-infection. All mice were bred inhouse, and *Sp140⁻/⁻* and *Sp140⁺/⁻* were littermates (**C-G**). **C**, **E**, and **G** are combined results of two independent infections. **A-E** shows representative analysis of one *Sp140⁻/⁻* line (line 1), whereas **F-G** includes a mixture of both line 1 and 2. (**C**, **E**, **F**, **G**) Mann-Whitney test. *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.005.

Figure 3. *Sp140⁻/⁻* mice have elevated *Ifnb* transcripts during bacterial infection. (**A**) Mice were infected with *M. tuberculosis* and at 28 days post-infection lungs were processed for total RNA, which were used for RT-qPCR. Combined results of 2 independent experiments. (**B**) Mice were infected with *L. pneumophila* and RT-qPCR was performed on lungs collected at indicated times. Combined results of 2 independent infections. All mice were bred in-house, *Sp140⁻/⁻* and *Sp140⁺/⁻* were littermates. (**A**-**B**) Mann-Whitney test. *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.005.

Figure 4. Global gene expression analysis of *Sp110*⁻/⁻, *Sp140*⁻/⁻ and B6.*Sst1*ˢ lungs after *M. tuberculosis* infection. (**A**) PCA or (**B**) Euclidean distance analysis of all the samples. (**C-D**) heatmaps of gene expression in log₂-fold change from B6. Genes shown are those significantly different between *Sp140*⁻/⁻ and B6 (**C**) GSEA Hallmark inflammatory response; and (**D**) GO type I IFN genes response genes. (**E**) volcano plot comparing *Sp140*⁻/⁻ to B6.*Sst1*ˢ expression. Dots in red are 2-fold differentially expressed with adjusted *p*-value ≤ 0.05.

Figure 5. Susceptibility of *Sp140*⁻/⁻ to *M. tuberculosis* and *L. pneumophila* is dependent on type I IFN signaling. (**A**-**B**) Mice were infected with *M. tuberculosis* and measured for body weight (**A**) and bacterial burdens at day 25 (**B**). Statistics in **A** show comparison to B6; data are from 9 B6, and 13 *Sp140*⁻/⁻ and *Sp140*⁻/⁻ *Ifnar*⁻/⁻ mice. Combined results of 2 experiments. (**C**-**D**) bacteria burden in *L. pneumophila*-infected mice at 96 hours. Combined results of 2 experiments. All mice were bred in-house (**A**-**B**, **D**); all but B6 were bred in-house (**C**). Mann-Whitney test (**A**-**D**). *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.005.

Figure 1– figure supplement 1. CRISPR–Cas9 targeting strategy for *Sp110⁻/⁻* mice. (**A**) Mouse *Sp110* gene. Guide RNA sequence for CRISPR–Cas9 targeting and protospacer-adjacent motif (PAM) are indicated. (**B-D**) *Sp110* locus in wildtype (WT) and three independent lines. Homozygotes of 2 lines identified by sequencing (**B**-**C**), and heterozygote of the 3rd line by PCR products separated on an agarose gel (**D**). Arrow indicates the mutant band.

Figure 2 – figure supplement 1. CRISPR–Cas9 targeting strategy for *Sp140⁻ᐟ⁻* and validation of founders. (**A**) Mouse *Sp140* gene. Guide RNA sequence for CRISPR–Cas9 targeting and protospacer-adjacent motif (PAM) are indicated. (**B**-**C**) *Sp140* locus in wildtype (WT) and 2 independent founders of *Sp140⁻ᐟ⁻* validated by sequencing. (**D**) Immunoblot for SP110 using BMMs from mice of the indicated genotypes. Intervening lanes have been removed for clarity (indicated by line in the image). (**E**) *M. tuberculosis*-infected mice were harvested for CFU at 25 days post-infection. Empty and filled triangles indicate the two independent lines of *Sp140⁻ᐟ⁻* used in this infection. All mice were bred in-house and *Sp140⁺ᐟ⁻* were littermates with *Sp140⁻ᐟ⁻* line 2. Mann-Whitney test, *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.005.

Figure 2 – figure supplement 2. Histology of B6, B6.*Sst1ˢ*, *Sp110⁻/⁻*, *Sp140⁻/⁻* lungs infected with *M. tuberculosis*. H&E staining of entire lung sections from mice of indicated genotypes at 25 days post-infection with *M. tuberculosis*. Black squares denote sections shown in Figure 2C. Each image represents a lung section from a different mouse. Borders in background color have been added around each image. Scale bar applies to all images.

A Genomic DNA (B6, *Sp140⁻ᐟ⁻1*)   cDNA from TB-infected mouse lungs (B6, *Sp140⁻ᐟ⁻1, Sp140⁻ᐟ⁻2*)

B Summary of edited *Sp140* homologs in *Sp140⁻ᐟ⁻* mice

Name	Dataset	Distinguishing SNPs from *Sp140* mRNA	Edited in *Sp140⁻ᐟ⁻1*?	Edited in *Sp140⁻ᐟ⁻2*?	Estimated level of expression in TB-infected lungs
LOC100041057	*Sp140* exon 3 amplicons, DNA (*Sp140⁻ᐟ⁻1* and B6 only)	G at 1483, T at 1513	Yes	Unknown (not expressed)	None
Sp140 homolog 1	*Sp140* exon 3 amplicons, DNA (*Sp140⁻ᐟ⁻1* and B6 only)	T at 1482, G at 1483	Yes	Unknown (not expressed)	None
Sp140 homolog 2	*Sp140* exon 2 and 3 amplicons, cDNA; RNA-seq of TB-infected lungs	None	Yes	No	Expressed
Sp140 homolog 3	*Sp140* exon 3 amplicons, cDNA	T at 1462	Yes	Yes	Very low (not detectable by RNA-seq)
Sp140 homolog 4	*Sp140* exon 3 amplicons, cDNA	T at 1500	Yes	Yes	Very low (not detectable by RNA-seq)

Figure 2 – figure supplement 3. Characterization of off-targets in *Sp140⁻ᐟ⁻* mice. (**A**) Schematic of amplicon sequencing strategy for *Sp140* and *Sp140* homologs. (**B**) Summary of edited *Sp140* homologs from amplicon sequencing and RNA-seq analysis. SNPs are denoted based on the *Sp140* X1 transcript. Expression level was roughly estimated from read counts. Two mice per genotype were used as biological replicates for amplicon sequencing for cDNA, and one mouse per genotype was used for amplicon sequencing from DNA.

A

B

Figure 2– figure supplement 4. Complementation of hyper type I IFN responses in *Sp140*–/– BMMs. (A) BMMs were left untreated or treated with TNF-α for 24 hours. Total RNA was used for RT-qPCR. Averages of technical duplicates for one biological replicate are shown. Data is representative of two independent experiments. **(B)** RT-qPCR of *Sp140*–/– BMMs transduced with either control SINV-minCMV-GAL4-mNeonGreen (SINV-mNeonGreen) or SINV-minCMV-Sp140 (SINV-SP140), primed with 5 ng/mL IFN-γ for 14 hours and treated with 10 ng/mL TNF-α for 4 hours. Three biological replicates were used for SINV-mNeonGreen conditions, and 4 biological replicates were used for SINV-Sp140 conditions. *, $p \leq 0.05$ calculated with an unpaired t-test with Welch's correction. Data are representative of two independent experiments.

**Figure 5 – figure supplement 1. Antibody blockade of IFNAR1 reduces bacterial burden in *Sp140*−/−
mice during *M. tuberculosis* infection.** Mice were infected with *M. tuberculosis* and treated with either
IFNAR1-blocking antibody or isotype control starting 7 days post-infection. At 25 days post-infection lungs
were harvested to enumerate CFU. Results of one experiment. All mice were bred in-house. Mann-Whitney
test, *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.005.