1

Local retinoic acid directs emergence of the extraocular muscle functional unit

Glenda Comai[1,#], Marketa Tesarova[2], Valerie Dupé[3], Muriel Rhinn[4], Pedro Vallecillo Garcia[5], Fabio da Silva[6,7], Betty Feret[4], Katherine Exelby[8], Pascal Dollé[4], Leif Carlsson[9], Brian Pryce[10], Francois Spitz[11], Sigmar Stricker[5], Tomas Zikmund[2], Jozef Kaiser[2], James Briscoe[8], Andreas Schedl[6], Norbert B. Ghyselinck[4], Ronen Schweitzer[10] and Shahragim Tajbakhsh[1,#]

Stem Cells & Development Unit, UMR CNRS 3738, Institut Pasteur, Paris, France
Central European Institute of Technology, Brno University of Technology, Brno, Czech Republic
Université de Rennes, CNRS, IGDR, Rennes, France
IGBMC-Institut de Génétique et de Biologie Moleculaire et Cellulaire, Illkirch, France
Institute for Chemistry and Biochemistry, Freie Universität Berlin, D-14195 Berlin, Germany
Université Côte d'Azur, INSERM, CNRS, iBV, Nice, France
Division of Molecular Embryology, German Cancer Research Center (DKFZ), Heidelberg
The Francis Crick Institute, London, NW1 1AT, United Kingdom
Umea Center for Molecular Medicine, Umea University, Umea, Sweden
Research Division, Shriners Hospital for Children, Portland, OR 97209, USA
Genomics of Animal Development Unit, Institut Pasteur, Paris, France
Authors for correspondance

ABSTRACT

Coordinated development of muscles, tendons, and their attachment sites ensures emergence of functional musculoskeletal units that are adapted to diverse anatomical demands among different species. How these different tissues are patterned and functionally assembled during embryogenesis is poorly understood. Here, we investigated the morphogenesis of extraocular muscles (EOMs), an evolutionary conserved cranial muscle group that is crucial for the coordinated movement of the eyeballs and for visual acuity. By means of lineage analysis, we redefined the cellular origins of periocular connective tissues interacting with the EOMs, which do not arise exclusively from neural crest mesenchyme as previously thought. Using 3D imaging approaches, we established an integrative blueprint for the EOM functional unit. By doing so, we identified a developmental time window where individual EOMs emerge from a unique muscle anlage and establish insertions in the sclera, which sets these muscles apart from classical muscle-to-bone type of insertions. Further, we demonstrate that the eyeballs are a source of diffusible retinoic acid that allow their targeting by the EOMs in a temporal and dose dependent manner. Using genetically modified mice and inhibitor treatments, we find that endogenous local variations in the concentration of retinoids contribute to the establishment of tendon condensations and attachment sites that precede the initiation of muscle patterning. Collectively, our results highlight how global and site-specific programs are deployed for the assembly of muscle functional units with precise definition of muscle shapes and topographical wiring of their tendon attachments.

INTRODUCTION

Acquisition of shape and pattern during development depends on the orchestrated crosstalk between a variety of tissues and cell types. While significant knowledge on the mechanisms of differentiation and patterning within individual tissues has been attained, much less is known on how patterning of different adjacent tissues is integrated. The vertebrate musculoskeletal system serves as an ideal model to study these processes as different tissues including muscle, tendon and their attachments need to be articulated in three-dimensions (3D) for proper function (Hasson, 2011; Huang, 2017).

Among the craniofacial muscles, the morphological configuration of the extraocular muscles (EOMs) has been a longstanding challenge in comparative anatomy and evolutionary biology. Besides specialized adaptations, the basic EOM pattern is shared among all vertebrate classes (Noden and Francis-West, 2006; Suzuki et al., 2016; Young, 2008) and includes 4 recti muscles (the superior rectus, the medial rectus, the inferior rectus, and the lateral rectus) and two oblique muscles (superior oblique and inferior oblique) for movement of the eyeball. Most vertebrates have also accessory ocular muscles involved in protecting the cornea (Noden and Francis-West, 2006; Spencer and Porter, 2006): the retractor bulbi, which serves to retract the eye in several groups of vertebrates and the levator palpabrae superioris, which is not directly associated with the eyeball but controls eyelid elevation in mammalian species. As such, the EOMs constitute an archetypal and autonomous functional unit for the study of how muscles, tendons and tendon attachments are integrated with the development of the eyeball, their target organ.

Craniofacial muscles are derived from cranial paraxial and prechordal head mesoderm (Noden and Francis-West, 2006; Ziermann et al., 2018) and the corresponding connective tissues, i.e. tendons, bones and cartilages and muscle connective tissue surrounding myofibres, were reported to be derived from cranial neural crest cells (NCC) (Nassari et al., 2017; Ziermann et al., 2018). Although early myogenesis is NCC-independent, NCC later regulate the differentiation and segregation of muscle precursors, dictate the pattern of muscle fiber alignment, and that of associated skeletal and tendon structures (Bohnsack et al., 2011; Ericsson et al., 2004; Noden, 1983b; Rinon et al., 2007; Tokita and Schneider, 2009; von Scheven et al., 2006). Moreover, studies of mouse mutants in which several genes were deleted in NCC cells have explicitly demonstrated their non-cell autonomous role in muscle morphogenesis at the level of the jaw (McGurk et al., 2017; Rinon et al., 2007; Shimizu et al., 2018), extraocular (Evans and Gage, 2005; Heude et al., 2015) and somitic-derived tongue muscles (Iwata et al., 2013). However, the full series of events driving morphogenesis of craniofacial musculoskeletal functional units is to date unexplored, probably due to the anatomical complexity of their configuration in the head.

Understanding the developmental mechanisms that allow musculoskeletal connectivity is
essential to understand their anatomical diversification during the evolution of the vertebrate
head. Yet, proximate factors that allow the coordinated emergence of the individual muscle
masses with that of their tendons and attachment sites are poorly defined.

To-date, much of the work on the tissue interdependency during musculosketal
development has been done in the developing limb. Lateral plate mesodem derived
connective tissue cells and tendon primordia establish a pre-pattern and restrict the locations
where limb muscle progenitors proliferate and differentiate (Kardon, 1998; Kardon et al.,
2003) prior to segregation (splitting) into individual muscle masses by several extrinsically
controlled mechanisms (Hasson, 2011; Sefton and Kardon, 2019). Limb tendons are initially
established independently of muscle, but their later development is modular, and depend on
muscle or cartilage according to their location in the limb (Huang, 2017). Finally, generation
of bone superstructures enable anchoring of muscles via tendons to the skeleton. Bone
superstructures are initiated independently of muscle, by a unique set of progenitors that co-
express SOX9 and Scleraxis (Scx) (Blitz et al., 2013; Blitz et al., 2009; Sugimoto et al.,
2013), but remarkably, muscle and superstructure patterning seem to be regulated in a
coordinated manner (Colasanto et al., 2016; Swinehart et al., 2013). However, given the
striking differences in gene regulatory networks governing cranial muscle development
(Diogo et al., 2015; Sambasivan et al., 2011), as well as the diverse embryological origins of
connective tissue cells in the head (Nassari et al., 2017; Ziermann et al., 2018), it is unclear if
this established logic of musculoskeletal integration is conserved in the head and for muscles
that do not operate with muscle-to-bone attachments such as the EOMs.

It has been proposed that the developing central nervous system, sensory organs,
pharyngeal endoderm and surface ectoderm provide spatial patterning cues to cranial
connective tissue precursors (Noden and Francis-West, 2006; Ziermann et al., 2018). In this
context, all-trans retinoic acid (ATRA), the biologically active metabolite of retinol (vitamin A),
is a critical morphogen with widespread roles in craniofacial development (Gitton et al., 2010;
Rhinn and Dolle, 2012). ATRA acts as a ligand for nuclear retinoic acid receptors (RARs),
which are ligand-dependent transcriptional regulators that work as heterodimers with retinoid
X receptors (RXRs) (Cunningham and Duester, 2015; Rhinn and Dolle, 2012). ATRA is
synthetized from retinol through two oxidation steps by specific retinol/alcohol
dehydrogenases (RDH/ADH) and retinaldehyde dehydrogenases (ALDH1A1, ALDH1A2 and
ALDH1A3) and catabolized into hydroxylated derivatives by enzymes of the cytochrome
P450 subfamily (CYP26s) (Cunningham and Duester, 2015; Rhinn and Dolle, 2012). ATRA
was shown to be critical for early eye development in several species (Bohnsack et al., 2012;
Harding and Moosajee, 2019; See et al., 2008), where the enzymes responsible of ATRA
metabolism are expressed in the early retina with tight spatiotemporal patterns (Cvekl and

Wang, 2009). Several studies have demostrated that through production of RA, the
developing eye acts as a signaling center nucleating anterior segment morphogenesis, with
the transcription factor *Pitx2* as the potential major downstream effector in periocular NCC
(Bohnsack et al., 2012; Matt et al., 2005; Matt et al., 2008; Molotkov et al., 2006). However,
whether ATRA is required for morphogenesis of the adjacent EOMs and associated
connective tissues remains unexplored.
Here, by a combination of 3D-mesoscopic imaging approaches, we present the first
integrative blueprint for morphogenesis of the EOM functional unit. We provide genetic
evidence for the existence of a retinoic acid signaling module that coordinates the
emergence of individual EOMs, their tendons and insertion sites. We show that the action of
retinoic acid signaling in muscle patterning is mainly non-cell autonomous, through its action
on the neural crest-derived periocular mesenchyme. Furthermore, we characterise the
dynamics of EOM-tendon insertion sites, which are impaired upon ATRA-deficiency. Overall,
our results show that in spite of the distinct embryological origins of these tissues in the
head, interactions between muscles, tendons and their attachments are similar to those
observed in the limb, yet they exhibit specifc hallmarks that are characteristic of this
anatomical location. This includes regionalized apoptotic foci that presage tendon attachment
sites in a context that is distinct from the canonical tendon to bone configuration.
146
Results
Genetic fate mapping of mouse periocular tissues
To investigate the morphogenesis of the EOMs in the mouse, we focused first on
adjacent tissues whose development should be coordinated with that of the EOMs. The
periocular mesenchyme (POM) is a heterogeneous cell population surrounding the optic cup
that gives rise to specialized structures of the anterior segment of the eye and connective
tissues associated with the EOMs (Williams and Bohnsack, 2015). With exception of the
EOMs and endothelial lining of ocular blood vessels (choroid), all connective tissues of the
POM (cartilage, muscle connective tissue, tendons), were reported to be derived from NCC
in zebrafish, chicken and mouse embryos (Couly et al., 1992; Creuzet et al., 2005; Gage et
al., 2005; Grenier et al., 2009; Johnston et al., 1979; Noden, 1983a). However, the extent of
the POM in relation to its different cell populations, has not been fully assessed. Thus, we
investigated the expression pattern of PITX2, a well established marker of both the POM and
EOM progenitors (Gage et al., 2005), *Scx*, a tendon progenitor marker (Schweitzer et al.,
2001) and SOX9, which is expressed in NCC and required for determination of the
chondrogenic lineage (Mori-Akiyama et al., 2003). We simultaneously traced the contribution
of different embryonic populations using *Mesp1^{Cre};R26^{tdTomato}* and *Wnt1^{Cre};R26^{tdTomato}* mice

that label the mesodermal and NCC derivatives (Danielian et al., 1998; Saga et al., 1999) in

combination with the *Tg:Scx-GFP* transgenic line (Pryce et al., 2007).

Coronal sections at the level of the eye at E13.5 showed that *Scx-GFP*, SOX9 and

PITX2 expression delimited a triangular area of cells medial to the eyeball encompasing the

EOMs (dotted lines, Fig 1A-D), however, with variable intensities along the medio-lateral

axis. SOX9 and PITX2 were more strongly expressed laterally between the retina and

ectoderm, where the EOM tendons insert. The *Scx-GFP* reporter labelled the future EOM

tendons and also muscle connective tissue as described for other muscles (Deries et al.,

2010). As expected, EOM tendons and muscle connective tissue at the insertion level were

of NCC origin (*Wnt1^{Cre};R26^{tdTomato}*, Fig 1A-D). Surprisingly, we found that lineage

contributions differed in dorsal sections. The future Annulus of Zinn, the tendinous ring from

which the recti muscles originate at the base of the skull (Spencer and Porter, 2006), as well

as the muscle connective tissue at this level, were of mesodermal origin

(*Mesp1^{Cre};R26^{tdTomato}*, Fig 1E-H), although they also co-expressed *Scx-GFP*, SOX9 and

PITX2 at this stage. The dual mesoderm/NCC contribution to POM connective tissues was

observed from E11.5 (not shown) to E17.5 (Suppl Fig 1A-F). These findings indicate that the

EOMs develop in close association with connective tissues of two distinct embryonic origins:

neural crest laterally (at the EOM insertion level) and mesoderm medially (at the EOM origin

level).

Development of the EOMs and their insertions overlap spatio-temporally

Patterning of the EOMs and their insertions has been largely disregarded due to their

complex anatomy, which is difficult to interpret from tissue sections alone. Therefore, we

established an imaging pipeline that includes whole mount immunofluorescence (WMIF) of

the periocular region, tissue clearing, confocal microscopy and reconstructions of the

obtained images into 3D objects. To visualize the developing EOM we used

MYOD/MYOG/Desmin as myogenic commitment and differentiation markers and MF20 to

label myofibers (Fig 2A-C', Video 1). At E11.75, the EOMs were present as a single anlage

(Fig 2A-A') medial to the eyeball. By E12.5, the EOM anlage split towards the POM-ring into

the submasses corresponding to the future four recti, two oblique muscles and the accesory

retractor bulbi muscle (Fig 2B-B'). Fully separated muscle subsets were evident by E13.5

(Fig 2C-C'). Thus, we conclude that EOM patterning in the mouse occurs by splitting from a

single anlage with major morphogenetic changes taking place between E11.75 and E12.5.

The EOMs insert into the sclera, a dense fibrous layer derived from the POM. Foci of

cell death in the POM were suggested to label the tendon attachment positions of the 4 recti

muscles (Sulik et al., 2001). Thus, to better understand the establishment of the EOM

insertions in the POM within the time window of EOM patterning, we performed whole mount

immunostaining for the POM marker PITX2 and LysoTracker Red staining to detect programmed cell death on live embryos (Fogel et al., 2012) (Fig 2D,E). WMIF for the PITX2 at E11.75 revealed a ring of PITX2^{high} cells immediately adjacent to the retinal pigmented epithelium (RPE) (POM-ring) that formed a continuum with PITX2^{low} cells extending towards the base of the EOM anlage (medial-POM) (Fig 2D). Lysotracker Red staining revealed a highly dynamic pattern in the POM-ring, where four foci were present at E11.75 at the horizons of the eye, which regressed from E12.5 onwards (Fig 2E). To confirm that the apoptotic foci define tendon attachment positions in the POM, we performed whole mount imunostaining for EOM progenitors and GFP (*Tg:SCX-GFP*) on Lysotracker stained embryos. Surprisingly, we observed that even before muscle splitting initiated, *Scx-GFP+* condensations bridged the edges of the EOM anlage and the four Lysotracker+ foci in the POM (E11.75, Video 2), presaging the locales of the future individual muscles (E12.5, Video 3).

To examine the EOM insertions with higher resolution, we immunostained *Wnt1^{Cre};R26^{tdTomato};Scx-GFP* coronal sections with POM markers (Fig 2F-N). At E12, PITX2 and SOX9 showed high expression in the lateral POM at the insertion site, while *Scx-GFP* was mostly expressed in a salt and pepper pattern at this level (Fig 2G-I). Where cells appeared more compact, as assessed by SOX9 and PITX2 staining, the Tomato staining appeared punctate (Fig 2F), reminiscent of apoptotic cells. Cell death was confirmed by LysoTracker Red and TUNEL staining on adjacent sections (Fig 2J, J'). By E13.5, PITX2 continued to mark a large periocular region, but SOX9 and *Scx-GFP* expression patterns were segregated and apoptotic foci undetectable (Fig 2K-N). Sox9 was expressed more laterally in a condensed group of cells probably representing the lateral tendon attachment points (Fig 2M), preceded medially by strong *Scx-GFP* staining at the tendon tips (Fig 2N). By E14.5, lateral EOM tendons expressed *Scx-GFP*, Tenascin and also PITX2 (Fig 2P-R, Suppl Fig 2A-H). At this stage, expression of SOX9 and PITX2 were donwregulated in the POM and reduced to the thin scleral layer (Fig 2R, S, Suppl Fig 2E-L). Altogether, these results show that patterning of the EOMs, their tendons and insertion sites overlap spatio-temporally and thus might be regulated in a coordinated manner. Moreover, similarly to other locations in the body, *Scx* and SOX9 show complementary patterns at the EOM insertion site, but additional specifc hallmarks, notably the presence of apoptotic foci and PITX2 expression, seem to be characteristic of this site.

Abnormal EOM morphogenesis in mutants with ocular malformations

To obtain further insights into the development of the EOM functional unit, we next focused on the role of the target organ, the eyeball, on EOM patterning. To end, we performed micro-computed tomography (micro-CT) scans in mouse mutants with a spectrum

of ocular perturbations. First, we analyzed $Pax6^{Sey/Sey}$ embryos, which are anophthalmic
where eye development is arrested at the optic vesicle stage (Grindley et al., 1995; Kaufman
et al., 1995; Roberts, 1967). In heterozygote embryos, EOM patterning proceeded as in
wildtype controls, where 4 recti and 2 oblique muscles were identifiable by E13.5 (Fig 3A, B).
In contrast, $Pax6^{Sey/Sey}$ mutant EOMs appeared as a unique sheet of tissue on top of a
rudimentary optic vesicle (Fig 3C). Given that Pax6 is expressed in the optic vesicle, giving
rise to the retina and RPE, as well as in the overlying ectoderm that forms the lens and
cornea (Grindley et al., 1995), but not in the EOMs, these observations suggest a non-cell
autonomous role in EOM patterning. Similarly, in $Lhx2^{Cre};Lhx2^{fl/fl}$ embryos, where inactivation
of the $Lhx2$ gene in eye committed progenitor cells leds to a degeneration of the optic vesicle
at E11.5 (Hagglund et al., 2011), EOM patterning was severely affected and few EOM
submasses could be observed (Suppl Fig 3A-B'). Finally, in cyclopic embryos resulting from
inversion of Shh regulatory regions, EOMs underwent splitting and projected towards the
centrally located ectopic eye, although with an abnormal 3D arrangement as reported for
human cyclopia conditions (Suppl Fig 3C-C'') (Ziermann et al., 2018). Together with previous
studies where surgical removal of the developing eye results in smaller EOMs with limited
spatial organization (Bohnsack et al., 2011; Twitty, 1932; von Scheven et al., 2006), our
observations reinforce the notion that the eye is not only a target organ for muscle insertion,
but it plays a critical role as an organizer of EOM patterning. However, as severe eye defects
are associated with abnormal migration of NCC to the periocular region (Langenberg et al.,
2008; Matsuo et al., 1993), it is difficult to pinpoint the underlying cause of the observed EOM
phenotypes. Thus, we decided to assess directly whether the developing eye could act as a
source of molecular signals to induce the geometrically complex EOM pattern and that of
their associated connective tissues.
262
Muscle patterning depends on retinoic acid signaling of neural origin
To study the role of target organ derived cues in EOM patterning, we investigated the
role of retinoic acid signaling, which plays multiple paracrine roles during embryonic eye
development (Cvekl and Wang, 2009). As ALDH1A1-3 are the rate-limiting enzymes in the
production of ATRA (Duester, 2009), we characterised their expression in the periocular
region at the time of EOM patterning. Between E10.5 and E12.5, $Aldh1a1$ was expressed
strongly in the dorsal retina, lens and corneal ectoderm (Suppl Fig 4A), $Aldh1a2$ was
expressed in the temporal mesenchyme (Suppl Fig 4A-A''), and $Aldh1a3$ was mostly
restricted to the ventral retina and corneal ectoderm (Fig 4A, Suppl Fig 4A) (Matt et al., 2005;
Molotkov et al., 2006). Interestingly, ALDH1A3 was the only retinaldehyde dehydrogenase
protein detected in the RPE and optic stalk/optic nerve between E10.5-E12.5 and thus
centrally positioned with respect to EOM development (Fig 4A). To target the retinoic

signaling pathway (Fig 4B), we used *Aldh1a3$^{-/-}$* (Dupe et al., 2003) and *Rdh10$^{-/-}$* (Rhinn et al.,

2011) mutants. We also administered the BMS493 inhibitor (pan-RAR inverse agonist) to

pregnant females every 10-12h between E10.5-E11.75, i.e. preceeding the initiation of

muscle splitting (Fig 2A-C) and once NCC migration to the POM has finalized (Osumi-

Yamashita et al., 1994; Serbedzija et al., 1992). Micro-CT analysis showed that *Aldh1a3$^{-/-}$*

and BMS493-treated embryos displayed eye ventralization and a shortened optic nerve

when compared to controls (Suppl Fig 4B), but they retained the overall organization of the

nasal capsule and orbit (Suppl Fig 4C). Ventral and lateral views of the 3D-reconstructed

EOMs (Fig 4C-E), showed that *Aldh1a3$^{-/-}$* and BMS493-treated embryos lacked the standard

3D arrangement of 4 recti and 2 oblique muscles observed at E13.5 in control embryos.

Nevertheless, in all cases EOMs originated medially from the hypochiasmatic cartilages of

the presphenoid bone, indicating that the overall orientation of the EOMs was preserved (Fig

4D'''). Thus, given that the EOMs are more affected at their insertion than their origin level

upon ATRA deficiency, this finding suggests that EOM patterning is in part modular.

To analyze EOM and tendon patterning with higher resolution, we performed whole

mount immunostainings for differentiated myofibres (MyHC) and tendon (Tenascin) (Fig 4F-

H'). On medial and lateral views of 3D-reconstructed EOMs, only the retractor bulbi and

superior rectus could be clearly identified among the non-segregated muscle fibres in

Aldh1a3$^{-/-}$ embryos (Fig 4G, G'). As expected from global invalidation of retinoic acid

signaling, EOM perturbation was more severe in BMS493-treated embryos (Fig 4H, H'). The

SO was absent, or continuous with the anterior part of the anlage, and the medial portion of

the retractor bulbi was thicker and less clearly isolated from the rest of the anlage. In both

conditions, Tenascin and *Scx-GFP+* cells were present at the tips of the individual, though

mispatterned, muscles at this stage (Fig 4G',H', Video 4). Analysis of *Aldh1a3$^{-/-}$* and

BMS493-treated embryos revealed on average a 26% reduction in the EOM volume

compared to controls (Fig 4I). However, MyHC+ myofibres were present in *Aldh1a3$^{-/-}$* and

BMS493-treated embryos (Fig 4G,H), suggesting that EOM differentiation was not overtly

affected in these conditions. Instead, these observations suggest that EOM fiber alignment

and segregation is dependent on retinoic acid signaling.

As dose and temporal control are critical in the context of retinoic acid signaling

(Chawla et al., 2016; Gitton et al., 2010; See et al., 2008), we performed other BMS493

injection regimes between E10.5 and E12.5 (Table 1). EOM patterning phenotypes

categorised as strong or severe at E13.5 were only obtained when a E10.75 time point of

injection (8PM of day E10.5) was included in the regime (experiment type I-III, Table 1, Suppl

Fig 4D). Surprisingly, even a single injection at this time point resulted in strong phenotypes

(experiment type IV, Table 1, Suppl Fig 4D), while exclusion of this timepoint (experiment

type VI and V, Table 1, Suppl Fig 4D) resulted only in mild phenotypes at best. Therefore, we

identified an early and restricted temporal window where ATRA activity, prior to any sign of
muscle splitting, impacts correct muscle patterning 36-48 hours later. Given the critical action
of RDH10 in generating retinaldehyde, the intermediate metabolite in the biosynthesis of
ATRA, we examined EOM development in *Rdh10* mutant embryos. *Rdh10* is normally
expressed in the optic vesicle and RPE (Sandell et al., 2007), and in *Rdh10*[-/-] embryos, eyes
develop intracranially, close to the diencephalon with a very short ventral retina (Rhinn et al.,
2011). WMIF analysis of *Rdh10* mutant EOMs revealed that the anlage was specified, but
lacked any sign of segmentation, in agreement with an upstream role in ATRA synthesis
(Suppl Fig 4E,F). Taken together, our data show that retinoic acid signaling is essential for
the correct myofiber alignment and segregation of EOM masses in a dose and time
dependent manner.
323

Muscle patterning is dependent on ATRA-responsive cells in the periocular
mesenchyme
To track cells that are responsive to ATRA and responsible of EOM patterning we
used a novel retinoic acid transgenic reporter line (Schedl et al., unpublished) that comprises
three *RARE* elements from the *Rarb* gene fused to the *Hsp68* promoter driving expression of
a tamoxifen-inducible *Cre-ERT2* recombinase. By crossing this line with the $R26^{mTmG}$ reporter
mice (Muzumdar et al., 2007) (Fig 5A, $Rare^{CreERT2};R26^{mTmG}$), we permanently labelled ATRA-
responsive cells and their descendants with membrane GFP expression. We found a greater
amount of responsive cells in the periocular region when Tamoxifen was administered
between E9.75-E10.5 (Suppl Fig 5A-C). To assess if myogenic cells and the adjacent POM
respond to retinoic acid signaling, we scored the co-expression of GFP (reporter) and
myogenic markers (PAX7, MYOD, MYOG) in bulk cell preparations from the periocular
region. Notably, the great majority of reporter positive-responsive cells were not myogenic
and thus were part of periocular conective tissues (88% if induction at E9.75; 98% if
induction at E10.5) (Fig 5B,C). Upon induction of the reporter at E9.75, and analysis on
tissue sections and by WMIF at E11.75 and E12.5, we observed that the mesenchyme
around the optic nerve in close proximity to the developing EOMs was positive for the
reporter (Fig 5D-E',H). Interestingly, the overal number of POM GFP+ cells increased from
E11.75 to E12, which is the temporal window corresponding to EOM splitting (Fig 5D-E').
The SOX9+ POM-ring that includes the EOM attachments also had abundant numbers of
GFP+ cells (Fig 5F-G',H'). Finally, inactivation of downstream retinoic acid signaling with
BMS493, prior and subsequent to the induction of $Rare^{CreERT2};R26^{mTmG}$ mice with Tamoxifen
(Suppl Fig 5D), greatly reduced the responsiveness of the reporter in the POM compared to
non-treated controls and led to muscle mispatterning (Suppl Fig 5E-H, Videos 5 and 6).
Together, these results are in agreement with previous studies (Matt et al., 2005; Matt et al.,

2008; Molotkov et al., 2006) showing that despite a sophisticated pattern of ATRA metabolism in the developing retina, retinoic acid signaling exerts its action mostly non-cell autonomously, in the adjacent POM, and within a temporal window that is crucial for EOM patterning.

Compound *Aldh1a1/Aldh1a3* mutants (Matt et al., 2005; Molotkov et al., 2006), *Rarb/Rarg* mutants (Ghyselinck et al., 1997) and *Rarb/Rarg* NCC-specific mutants (Matt et al., 2008) display similar phenotypes in the anterior segment of the eye and POM. These findings have been attributed to the fact that ATRA synthesized by ALDH1A1 and ALDH1A3 in the neural retina activate RARB and RARG in the POM*,* which work as heterodimers with RXRA (Kastner et al., 1994; Mori et al., 2001). To directly examine whether the role of retinoic acid signaling in muscle patterning is driven non-cell autonomously, through the action of RARB/RARG receptors in the NCC-derived POM, we performed WMIF for muscle in *Wnt1^{Cre};Rarb$^{fl/fl}$;Rarg$^{fl/fl}$* embryos. Accordingly, these NCC-*Rarb/Rarg* mutants showed absence of muscle splitting (Fig 5I-J') as observed in *Rdh10$^{-/-}$* embryos (Suppl Fig 4F). Altogether, these data suggest that synthesis of ATRA from neural derivatives (retina, optic nerve, RPE) at early stages (≈E10.5) is crucial for EOM patterning, through its action on NCC-derived periocular connective tissues.

Defective organization of the EOM insertions in retinoic acid signaling deficient embryos

To examine which connective tissue subpopulations, including the EOM insertions in the POM, were affected by retinoic acid signaling deficiency, we assessed the distribution of POM markers on tissue sections of *Aldh1a3$^{-/-}$* and BMS493-treated embryos. Examination of the distrubution of Collagen XII (Fig 6A-C'), a marker of the sclera and the presumptive corneal epithelium (Oh et al., 1993), showed that ATRA-deficiency led to reduced expression of Collagen XII at E12 at the level of the medial-POM which gives rise to the sclera (Fig 6B-C, brackets, B'-C' insets). In addition, while PITX2 was expressed continuously in the entire POM in controls (Fig 6D, dotted lines, D', inset), expression in the medial-POM was lost upon ATRA-deficiency (Fig 6E-F, brackets, E',F', insets). This observation is in agreement with *Pitx2* being a retinoic acid signaling target in the NCC-derived POM (Matt et al., 2005; Matt et al., 2008; Molotkov et al., 2006) and the sclera being absent at fetal stages in *Rarb/Rarg*-NCC mutants (Matt et al., 2008). Interestingly, POM organization was unaffected in *Myf5$^{nlacZ/nlacZ}$* mutants (Fig 6G, dotted lines, G', inset), where the initial EOM anlage forms but myogenesis is aborted from E11.5 (Sambasivan et al., 2009). Thus, we conclude that intrinsic organization of the POM is dependent on retinoic acid signaling but not on the presence of myogenic cells.

To understand if the abnormal distribution of POM markers translates into EOM
insertion defects, we examined the organization of the POM-ring in 3D in control, *Aldh1a3*[-/-]
and BMS493-treated embryos. We systematically compared E11.75 and E12 embryos to
visualize the organization of these tissues prior to, and during muscle splitting (Fig 2A-C).
Control embryos showed the presence of partially overlapping SOX9 and PITX2 POM-ring
domains, together with 4 apoptotic foci labelling the prospective tendon attachment points of
the 4 recti muscles (Fig 6H-J, Suppl Fig 6A-C',E,F). As expected, ATRA-deficiency perturbed
these patterns in a dose dependent manner. In lateral views of the head of *Aldh1a3*[-/-]
embryos, SOX9 and PITX2 had narrower expression domains than controls, restricted to the
anterior-dorsal and ventro-lateral eye quadrants (Fig 6K-L, Suppl Fig 6G,H). In BMS493-
treated embryos, expression of SOX9 in the POM (Fig 6N, Suppl Fig 6I) and RPE (Suppl Fig
6D,D') was reduced to a minority of cells, and PITX2 expression was restricted ventro-
laterally (Fig 6O, Suppl Fig 6J). Accordingly, the apoptotic foci were lost or disorganized in
both cases of ATRA-deficiency with a single ventro-medial Lysotracker+ spot almost
invariably present in both conditions (Fig 6M,P, Suppl Fig 6G-J). Finally, in medial views of
the reconstructed 3D volume, we saw a continuum of PITX2 expression from the POM-ring
to the medial-POM in control embryos (Suppl Fig 6N), while we noticed loss of medial-POM
PITX2 expression in both cases of ATRA-deficiency (Suppl Fig 6O-P, blue arrowheads) as
observed on tissue sections (Fig 6E',F', brackets). Thus, ATRA is required for the proper
specification/organization of the medial POM and POM-ring.

As tendons were present at muscle tips in E13.5 *Aldh1a3*[-/-] and BMS493-treated
embryos, but located ectopically with respect to a wild-type configuration (Fig 4F'-J'), we
suspected that the initial 3D-organization of *Scx-GFP*+ tendon and connective tissue
progenitors might also be affected. We performed WMIF for differentiated muscle (MyHC)
and tendon (*Tg:SCX-GFP*) and segmented the reconstructed 3D volume. In control embryos
at E12.5, *Scx-GFP*+ condensations projected from the tips of the future 4 recti muscles
towards the POM-ring (Fig 6 Q,Q', Video 7), facing the 4 apoptotic foci (Fig 6J, Video 3).
Moreover, *Scx-GFP*+ cells were also present in POM-muscle connective tissue presaging
the final locales of the recti muscles (Fig 6T). In BMS493-treated embryos at E12, *Scx-GFP*
expression was less organized and diffuse (Fig 6R,R', blue arrowhead) yet *Scx-GFP*+
condensations could be observed at the tips of ectopic muscle masses (Fig 6R',U white
arrowhead, Video 8). In contrast, *Scx-GFP* prepatterning in muscle connective tissues was
preserved in *Myf5*[nlacZ/nlacZ] mutants and *Scx-GFP*+ condensations formed at the positions of
prospective (although absent) recti muscles (Fig 6S,S',V, white arrowheads). This situation is
similar to the presence of individual tendons on the distal autopod of muscle-less limbs
(Huang et al., 2015; Kardon, 1998) and branchial arches (Grenier et al., 2009). However, at
later stages, tendons did not continue their maturation in the absence of EOMs (Suppl Fig

6Q-R', video 9 and 10), reinforcing the notion that, as in most other anatomical locations
(Huang, 2017), tendon differentiation and maintenance appear to ultimately rely on muscle-
tendon interactions. In summary, ATRA produced by the developing eye influences the
organization of the entire neural crest derived POM, including tendon condensations and
EOM insertions in the POM-ring. Moreover, given that retinoic acid signaling exerts this
action with a time interval that precedes and overlaps with that of EOM anlage segregation
into individual muscle masses, we propose that EOM and insertion site patterning are
regulated in a coordinated manner.
430
431
DISCUSSION
The stark differences in embryological origins, function and susceptibility to disease
among cranial and somite-derived muscles (Sambasivan et al., 2011), provides impetus to
study in detail the morphogenesis of different muscle groups and their integration as part of
the musculoskeletal system. Here we focused on EOMs, an evolutionarily conserved group of
muscles that are precisely engineered for fine displacement of the eyeball, and thus crucial
for visual acuity. Using genetic and 3D imaging approaches, we analysed EOM development
from their emergence as a unique anlage to the establishment of a fully formed functional
unit with insertions in the sclera, which sets it apart from classical muscle-to-bone insertions
studied to date. We identified a spatiotemporal window in which retinoic acid signaling from
the target organ is required for patterning of NCC-derived soft tissues of the POM. These
findings provide insights into the deployment of site-specific programs for the establishment
of anatomically distinct muscle functional units, with precise definition of muscle shapes and
topographical wiring of their tendon attachments.
446
Dual origin of EOM insertions
The connective tissues of the rostral cranium (bone, cartilage, tendon and muscle
connective tissue) were reported to be derived from the neural crest (Nassari et al., 2017;
Noden and Francis-West, 2006; Ziermann et al., 2018). However, the hypochiasmatic
cartilages, from which the recti muscles originate in mammals, are an exception to this rule
as they are mesoderm-derived (McBratney-Owen et al., 2008). Here, we found that at this
location, tendon and muscle connective tissue are also of mesodermal origin. As such, EOM
originate in mesoderm-derived bone and insert in NCC-derived fibrous tissue of the sclera
with muscle connective tissue following the embryonic origins of the respective attachments
as observed in other anatomical locations (Heude et al., 2018). Taken together, our
observations redefine a novel boundary for NCC contribution to the connective tissues of the

periocular region and suggest that at their origin, the EOMs and their associated connective
tissues might have evolved neomorphically *in situ* from the same mesodermal source.
460

RA responsiveness on the POM
Previous studies showed that NCC have a critical role in the acquisition of cranial
muscle morphology (Bohnsack et al., 2011; McGurk et al., 2017; Shimizu et al., 2018; Tokita
and Schneider, 2009; von Scheven et al., 2006). Our data using the *Rare^CreERT2* reporter and
following invalidation of *Rarb/Rarg* in NCC derivatives suggest that the action of retinoic acid
signaling on EOM patterning is also indirect, through its action on periocular connective
tissues.
The observation that BMS493 treatments to impair ATRA signaling results in a more
severe phenotype than that observed in *Aldh1a3* mutants could have several explanations.
First, BMS493 is not a RAR antagonist but pan-RAR inverse agonist, which is capable of
repressing RAR basal activity by favoring the recruitment of corepressors and promoting
chromatin compaction (Germain et al., 2009). Second, additional sources of ATRA could
signal to the POM and be inhibited by the BMS493 treatment. Candidates include *Aldh1a1*
(retina), *Aldh1a2* (temporal mesenchyme) and *Cyp1B1*, a member of the cytochrome p450
family of mono-oxygenases which can generate ATRA in the retina in a retinaldehyde
dehydrogenease-independent manner (Chambers et al., 2007). Noneless, the severe
patterning phenotypes observed in the *Aldh1a3* single mutant suggest that the local
synthesis of ATRA in the RPE, optic stalk and ventral retina at early stages (≈E10.5) cannot
be fully compensated by the action of other retinaldehyde dehydrogeneases in adjacent
tissues (this study, (Dupe et al., 2003; Harding and Moosajee, 2019)). Interestingly, the
constitutive *RARE-hsp68-lacZ* transgene displays strong reporter activity in the retina and
RPE (Matt et al., 2005; Rossant et al., 1991). Thus, we cannot exclude the possibility that
some of the phenotypes observed in periocular connective tissues of the *Aldh1a3* mutant are
partly due to changes in signalling pathways downstream of ATRA in the retina and/or RPE,
that subsequently act in the POM.
486

EOM and tendon patterning are concerted processes under RA control
We identified a short temporal window (E10.5-E11.75) where ATRA is essential for the
precise morphogenesis of several elements of the POM, including the formation of tendon
condensations and insertion sites in the POM-ring. These phenotypes are already evident
from E11.75, a stage where the EOM still figures as a muscle anlage. Interestingly, analysis
of chick and mouse limb development also identified a short time window where limb
mesenchyme perturbations result later in muscle and tendon patterning defects (Chevallier
and Kieny, 1982; Hasson et al., 2010). Moreover, concomitant muscle and tendon defects

have been observed in the chick limb and zebrafish jaw upon retinoic acid signaling
misregulation (McGurk et al., 2017; Rodriguez-Guzman et al., 2007). Altogether, these
results suggest that at certain anatomical locations including the EOMs, endogenous local
variations in the concentration of retinoids contribute to the establishment of *Scx-GFP+*
condensations, a critical step in muscle functional unit assembly. As ectopic muscles masses
are associated with ectopic tendons upon ATRA deficiency, one possible explanation is that
retinoic acid signaling directly controls *Scx-GFP+* condensations, which in turn help organize
the adjacent forming muscles. On the other hand, it is also possible that connective tissue
mispatterning upon ATRA deficiency determines the organization of the muscle masses and
the muscle tips direct the organization of the *Scx-GFP+* condensations.

Role of RA in the integration of EOM patterning and insertion site formation

The EOMs insert into the sclera, a non-bone structure in mammals. The generation of
4 attachment points that precisely mirror the position of the 4 recti muscles is a
morphogenesis conundrum, as details on their specification are scarse. Four mesenchymal
condensations have been described at the periphery of the developing optic cup in cat and
human embryos, in positions facing the locations of the recti muscles (Gilbert, 1957; Gilbert,
1947). In the mouse, at equivalent developmental timepoints (E11.5-E12.5) and homologous
positions, apoptotic foci were observed and suggested to provide attachment for the recti
muscles (Sulik et al., 2001). Using 3D-imaging, we showed that at E11.75, the apoptotic foci
are aligned with *Scx-GFP+* condensations that project medially, before any sign of splitting of
the EOM anlage. As the apoptotic foci decrease in size from E12.5 onwards concomittant
with ongoing muscle splitting and tendon formation, it is possible that they initially presage
the POM for future tendon insertions, but are not required for attachment *per se*. These foci
appear in a SOX9+ PITX2+ domain of the future sclera, however, to our knowledge their
presence has not been reported in superstructures at other anatomical locations.

In agreement with the role of RA in the induction of cell death in another locations in
the embryo (Dupe et al., 2003), our study shows that upon perturbation of RA signaling, the
apoptotic foci in the POM are absent or severely reduced. However, it is also possible that
the loss of foci upon RA inhibition is secondary to loss or mispatterning of the SOX9+
PITX2+ POM-ring. Thus, it will be of interest to determine if null or NCC specific mutations in
PITX2 affect the patterns or amount of naturally occurring cell death in the periocular
condensations. Although our work sheds light on the genetic mechanisms that regulate EOM
insertion, the cellular mechanisms underlying this process remains undefined. It is tempting
to speculate that the apoptotic foci are related to the disappearance of signaling centers
(Jernvall et al., 1998; Nonomura et al., 2013) or compaction-mediated cell death and

extrusion (Moreno et al., 2019). Future experiments directed towards modifying the amount
or timing of cell death will be informative.

Few reports have demonstrated that mispatterning of specific muscles is coupled with
aberrant superstructures (Colasanto et al., 2016; Swinehart et al., 2013; Tokita and
Schneider, 2009) or removal of the prospective tendon attachment sites (Rodriguez-Guzman
et al., 2007). Our results, are in agreement with that model given that EOM mispatterning is
concomitant with aberrant EOM insertions. Given that mammals do not develop a cartilage
layer within the sclera (Franz-Odendaal and Vickaryous, 2006; Thompson et al., 2010),
expression of SOX9 in the POM-ring of mouse embryos at the time of EOM patterning is
intriguing. SOX9 expression at this level appears to be dependent on a functional RPE and
ATRA ((Thompson et al., 2010), this study). In this context, our data suggest that transient
SOX9 expression may represent a redeployment of the developmental module for tendon
attachment, despite the fact there is no cartilage in the mammalian sclera. Finally,
transcriptome analysis revealed the existence of global and regional regulatory modules for
superstructure patterning in the limb bones (Eyal et al., 2019), offering a mechanism to
induce variations in attachment sites without having to rewrite the entire skeletogenic
program (Eyal et al., 2019). As several markers of the POM (*Pitx2*, *Foxc1/2*) and retinoic acid
signaling modulators (*Cyp26a1/b1*) were identified as part of specific limb superstructure
signatures (Eyal et al., 2019), it is tempting to speculate that those genes also play a
conserved role in the generation of the attachment module of the EOMs.

Conclusion
The developing eye has been proposed to be key organizer of craniofacial development,
independent of its role in vision (Kish et al., 2011). This notion is based on the role of the eye
for proper NCC migration to the periocular region (Langenberg et al., 2008), the common
association of ocular and craniofacial developmental abnormalities (Zein, 2012) and the
initial development of an eye in blind vertebrate species (Durand, 1976; Kish et al., 2011;
Yamamoto et al., 2004). Moreover, as the appearance of a "camera-type" eye is a vertebrate
innovation (Lamb et al., 2007) and the EOM are already present in lamprey (Suzuki et al.,
2016), it is possible that muscles and their target tissue might have co-evolved. Here, by
characterizing coordinated patterning of the EOMs, their respective tendons and
attachments, our findings illustrate further the role of the developing eye as a signaling center
allowing integration of the EOM functional unit in the POM. Our results show that the tissue
interactions during the development of this craniofacial muscle unit share features with those
described in the limb, but with additional regional properties (e.g. cell death, specification of
attachments in the sclera) that seem to have been specifically incorporated into this group.
Moreover, the capacity to instruct muscle patterning through variations in connective tissue

derivatives, provides a mechanism to explain the plasticity of the musculoskeletal system at
the anatomical and inter-species levels, while ensuring structural and functional integration
during evolution. These findings also imply that the generation of musculoskeletal units do
not require major restructuring of the developmental programs of all the tissues implicated.
Instead, co-option of a general program and simultaneous addition of local features appear
to elaborate musculoskeletal diversity.

Figure Legends
Fig1. Lineage contributions to the EOM functional unit.
(A-H) Neural Crest cell (NCC, *Wnt1^{Cre};R26^{Tom}*) and mesoderm (*Mesp1^{Cre};R26^{Tom}*) lineage
contributions to the periocular region on coronal cryosections of E13.5 embryos, combined
with immunostaining for GFP (*Tg:Scx-GFP* tendon reporter), muscle (PAX7/MYOD/MYOG,
myogenic markers), SOX9 (chondrogenic/scleral marker) and PITX2 (EOM muscle
progenitor and periocular mesenchyme marker). Sections at ventral (A-D) and dorsal (E-H)
levels. Asterisks in 1A-D denote tendon insertion sites. Dashed regions in A-H delimitate the
periocular mesenchyme. Panels A-B and E-F, an anti-DsRed antibody was used to enhance
the R26^{Tom} signal. Panels C-D and G-H, the endogenous reporter signal was used. C,D and
G,H are consecutive sections to the ones shown in panels A,B and E,F respectively. (n=3
per condition).
az, annulus of zinn; ce, corneal ectoderm; eom, extraocular muscles; le, lens; l-pom, lateral
periocular mesenchyme, at the level of the tendon insertions; mct, muscle connective tissue;
m-pom, medial periocular mesenchyme; r, retina; rpe, retinal pigmented epithelium; s, sclera;
t, tendon (at the level of the insertion); ti, tendon insertion site.

Fig2. Developmental timing of EOM functional unit components
(A-C) WMIF for the MYOD/MYOG/Desmin (myogenic differentiation markers) (A, B) and
MF20 (myofibers) (C) at the indicated embryonic stages. Medial views (left eye).
(D) WMIF for MYOD/Desmin (labeling the EOM anlage, shown as isosurface for clarity) and
PITX2 (labelling the POM and EOM anlage) on E11.75 embryos (left eye). Lateral and
medial views as schemes.
(E) Whole Mount LysotrackerRed staining of the periocular region at the indicated stages.
The POM is delimited with a dotted line. Anterior (a), posterior (p), medial (m) and lateral (l)
apototic foci are indicated. Asterisks indicate apoptotic foci with reduced intensity from E12.5
onwards.

(F-I) Immunostaining of lateral POM in coronal sections of E12 *Wnt1*[Cre]*;R26*[Tom]*;Scx-GFP*

embryos. Images correspond to insertion site of superior rectus muscle in POM. G,

immunostaining of section adjacent to the one shown in (F,H,I; channels split for clarity).

(J,J') Tunel, LysotrackerRed and SOX9 staining of area depicted in I. Scale bars 50 microns.

(K,N) Immunostaining of lateral POM in coronal sections of E13.5 *Wnt1*[Cre]*;R26*[Tom]*;Scx-GFP*

embryos. Images correspond to insertion site of superior rectus muscle in POM. L,

immunostaining of section adjacent to the one shown in (K,M,N; channels split for clarity).

(O-S) Immunostaining on coronal sections of E14.5 *Scx-GFP* embryos. Tnc (Tenascin), GFP

and PITX2 co-localize in tendons at level of insertion. MF20 (O-Q) and Smooth muscle actin

(SMA) (R,S) were used to label EOM muscle. Dashes in R,S were drawn according to GFP

labelling on adjacent sections. Asterisks in O denote tendon insertion sites. Low

magnification views of panels P-S are shown in Suppl Fig 2. (n=3 per condition).

anl, anlage; apop, apoptosis spots; eom, extraocular muscles; ir, inferior rectus; le, lens; lg,

lacrimal gland sulcus; t, tendon (at the level of the insertion); lr, lateral rectus; i, inferior EOM

anlage projection; io, inferior oblique; mr, medial rectus; pom-r, periocular mesenchyme ring;

m-pom, medial periocular mesenchyme; rpe, retinal pigmented epithelium; s, sclera; s,

superior EOM anlage projection; sr, superior rectus; so, superior oblique; rb, retractor bulbi.

Fig3. EOM morphogenesis in mutants with eye defects

(A-C) μCT-based 3D reconstruction of EOM, eyeball, optic nerve and lens in E13.5 control

(A), *Pax6*[Sey/+] (B) and *Pax6*[Sey/Sey] embryos (C). Note that in heterozygote embryos EOM

patterning proceeds normally despite having a smaller retina and lens (n=3).

Fig4. Extraocular muscle morphogenesis is dependent on ATRA

(A) Immunostaining for ALDH1A3 on coronal sections of E10.5, E11.5 and E12.5 control

embryos (n=2).

(B) Scheme of retinoic acid signaling pathway with key enzymes for oxidation of retinol and

retinaldehyde (pink) and mutants/inhibitors used in this study (blue).

(C-D''') μCT-based 3D-reconstruction of chondrogenic mesenchymal condensations of nasal

capsule, trabecular cartilage, EOM, eyeball, optic nerve and lens in E13.5 control, *Aldh1a3*[KO]

and BMS treated embryos. EOM visualization in context of whole head (D), nasal capsule

(D'-D''), trabecular capsule (D''').

(E) Raw μCT data of panels in C-D (n=2)

(F-H) EOM 3D-reconstructions of whole mount MyHC immunostaining of E13.5 control (F),

Aldh1a3[KO] (G) and BMS treated embryos (H) (n>9). (1-7) denote non-segregated muscle

masses with differential fiber orientation (see also Suppl Fig 5E). Raw immunostaining data

for MyHC (muscle) and Tenascin (Tnc, tendon) are shown in panels F'-H'.

(I) Relative EOM volume (compared to control) of E13.5 3D-reconstructed whole mount immunostained EOM of E13.5 control, $Aldh1a3^{KO}$ and BMS treated embryos. Each dot represents an individual embryo (n>9).

dr, dorsal retina; hc, hypochiasmatic cartilage; io, inferior oblique; ir, inferior rectus; le, lens; lr, lateral rectus; mr, medial rectus; os, optic stalk; se, surface ectoderm; so, superior oblique; sr, superior rectus; rb, retractor bulbi; rpe, retinal pigmented epithelium; vr, ventral retina

Fig5. Periocular connective tissues are responsive to retinoic acid signaling

(A) Scheme of mouse alleles used. $Tg:RARE^{CreERT2}$ line is composed of three RARE elements from the *Rarb* gene fused to hsp68 promoter driving expression of a tamoxifen-inducible Cre^{ERT2} recombinase. Crossing this line with the $R26^{mTmG}$ reporter line, allows permanent labelling of the ATRA-responsive cells and tracing of their descendents.

(B) Strategy used to determine cell-types reponsive to retinoic acid signaling in $Tg:RARE^{CreERT2}::R26^{mTmG}$ embryos. Tamoxifen was injected to pregnant females at E9.75 or E10.5 and cells of the periocular region isolated at E11.75 or E12.5.

(C) Percentage of recombined (GFP+) cells in connective tissue (CT) and myogenic (Myog) populations for different treatments. Each dot represents an individual embryo (n>4 embryos/condition).

(D-G') Whole mount immunostaining for SMA (smooth muscle actin, differentiated muscle), GFP (ATRA-responsive cells) and SOX9 (periocular mesenchyme) at indicated embryonic stages. (D-E') 3D-reconstructed EOM labelling. Arrowhead indicates an increased number of reporter positive cells surrounding the optic nerve (asterisk) at E12. (F-G') Segmented periocular region. pom-ring (dotted lines) (n=3).

(H-H') Coronal sections at different eye levels of E11.75 $Tg:RARE^{CreERT2}::R26^{mTmG}$ embryos immunostained for GFP (RA-responsive cells) and MYOD/MYOG (muscle). Higher magnification views as insets. Arrowheads mark presence of RA-responsive cells in medial periocular mesenchyme (H) and lateral-POM (H'). Asterisks indicate sporadic labelling in myogenic cells (n=3).

(I-J') EOM 3D-reconstructions of whole mount MF20 immunostaining of E12.5 of control ($Wnt1Cre: Rarb^{fl/+}:Rarg^{fl/+}$) (I,I') and mutant ($Wnt1^{Cre}:Rarb^{fl/fl}:Rarg^{fl/fl}$) (J,J') embryos. Medial and lateral views are shown. Note absence of splitting in mutant EOM. Arrowheads indicate split EOM in controls (n=2 per genotype).

Fig 6. EOM insertions in the POM altered upon ATRA-deficiency

(A-C') Immunostaining for Collagen XII (ColXII) on E12 coronal sections. Reduced ColXII expression domain in $Aldh1a3^{KO}$ (B,B') and BMS493 treated embryos (C,C') (arrowheads).

ColXII expression in medial-pom (future sclera) is lost (brackets). Split ColXII channel as insets in (A'-C').

(D-G) Immunostaining for PITX2 (muscle progenitors, periocular mesenchyme) and MYOD/MYOG (myogenic cells) on coronal E12 sections. Asterisks indicate superior oblique muscle (SO). Dashed line mark periocular mesenchyme (pom). Brackets on *Aldh1a3^{KO}* and BMS493 embryos indicate absence of PITX2 expression in medial-pom. Split PITX2 channel as insets in (D'-G').

(H-P) Lateral views of periocular region of whole mount immunostaining (GFP, SOX9, PITX2) and Lysotracker Red Staining (Lyso, apoptosis) of E12 control (H-J), *Aldh1a3^{KO}* (K-M) and BMS493 treated (N-P) embryos (left eyes). *Aldh1a3^{KO}* and BMS493 embryos do not show a complete pom-ring as controls (dotted lines in I). Arrowheads mark remaining expression of SOX9/PITX2 in the POM of *Aldh1a3^{KO}* (L) or inhibitor treated (O) embryos. Asterisks in (J,M,P) mark apoptosis spots in POM.

(Q-V) WMIF for MF20 (myofibers) and GFP (*Tg:Scx-GFP*) on control (Q,Q',T), BMS493 treated (R,R',U) and *Myf5$^{nlacZ/nlacZ}$* E12 embryos (S,S',V) (left eyes). **(Q-S)** The periocular area was segmented from adjacent structures for ease on visualization (dashed lines). **(T-V)** Single Z-section of the segmented volume. (n=3 per condition)

White arrowheads in Q',S', T,V indicate correct position of tendon condensations for recti muscles. Blue arrowhead in R' mark a diffuse *Scx-GFP+* pattern at the position of the mispatterned MR (medial recti) muscle and white arrowheads in R', U mark an ectopic *Scx-GFP* condensation at the muscle tip.

le, lens; nlg, nasolacrimal gland; eom, extraocular muscles

Supplementary Figure Legends

Supplementary Fig 1. Lineage contributions to the EOM functional unit.

(A-F) Neural Crest cell (NCC, *Wnt1^{Cre};R26^{Tom}*) and mesoderm (*Mesp1^{Cre};R26^{Tom}*) lineage contributions to the periocular region on coronal cryosections of E17.5 embryos, combined with immunostaining for GFP (*Tg:Scx-GFP* tendon reporter) and muscle (Tnnt3). Sections at ventral (A-F) and dorsal (A'-F') levels. Note that lateral tendons at the level of the orbit are NCC-derived (A-C), whereas tendons at the levels of the annulus of zinn (D'-F') are mesoderm-derived.

az, annulus of zinn; eom, extraocular muscles; hc, hypochiasmatic cartilage; lt, lateral tendon (at the level of the insertion); mct, muscle connective tissue

Supplementary Fig 2. Developmental timing of EOM functional unit components

(A-L) Immunostaining for the indicated markers on coronal sections of E14.5 *Tg:Scx-GFP* embryos. Tnc (tenascin) (n=2 per condition).

eom, extraocular muscles; lt, lateral tendon (at the level of the insertion; asteriks); m-pom, medial periocular mesenchyme; mct, muscle connective tissue; on, optic nerve; rpe, retinal pigmented epithelium

Supplementary Fig 3. EOM morphogenesis in mutants with eye defects

(A-B) μCT-based 3D-reconstruction of EOM and eyeball in E18.5 control (A) and *Lhx2Cre:Lhx2fl/fl* (B) embryos. Arrowheads highlight some extent of EOM segregation in the mutant. (A'-B') Coronal sections of control (A') and mutant (B') embryos stained with MyHC. Arrowheads indicate individual EOM masses (n=2).

(C-C'') Analysis of EOM patterning in E15.5 embryos containing inversions of *Shh* genomic regulatory regions (Inv(6-C2)). (C) Skeletal preparation of mutant embryos displaying cyclopia. (C'-C'') μCT-based 3D reconstruction of EOM and eyeball of mutant embryos (n=2).

Supplementary Fig 4. EOM morphogenesis is dependent retinoic acid signaling

(A) Whole mount in situ hybridization for *Aldh1a1*, *Aldh1a2* and *Aldh1a3* in E10.5, E11.5 and E12.5 embryos (n=3).

(A') In situ hybridization for *Aldh1a2* and immunostaining for ALDH1A2 and muscle (MF20) on E12.5 coronal sections. ALDH1A2 is expressed in the temporal mesenchyme and muscle connective tissue (n=3).

(B) μCT-based 3D-reconstruction of eyeball, optic nerve and lens of E13.5 control, *Aldh1a3^{KO}* and BMS treated embryos (n=2 each genotype). The lower row is a scheme of a sphere fitting the eyeball and lens, and a cylinder for the optic nerve. Note ventralization of the eyeball in *Aldh1a3^{KO}* and BMS treated embryos.

(C) μCT-based 3D-reconstruction of the mesenchymal condensations of the nasal capsule and trabecular cartilage of E13.5 control (white), *Aldh1a3^{KO}* (yellow) and BMS-treated (green) embryos.

(D-D') EOM 3D-reconstructions of whole mount MyHC immunostaining of E13.5 of BMS-treated embryos as described in Table S1. The most severe phenotype obtained in each condition is shown. Two different examples of the phenotype observed upon treatment VI (a,b) are shown (most and least severe). An *Aldh1a3^{KO}* embryo is shown as comparison. Asterisk denote ectopic duplicated SO (superior oblique) muscle. (D') Higher magnification views from D with examples of adjacent, non-split muscle masses (1,2) depicted by differential fiber orientation.

(E-F) Whole mount MyHC immunostaining of E13.5 control (G) and *Rdh10^KO* (H) embryos. Note total absence of muscle splitting in mutant (n=3 per condition).

ce, corneal ectoderm; dr, dorsal retina; io, inferior oblique; ir, inferior rectus; le, lens; lgb, lacrimal gland bud; lr, lateral rectus; mr, medial rectus; nlg, nasolacrimal groove; np, nasal pit; os, optic stalk; se, surface ectoderm; so, superior oblique; sr, superior rectus; rb, retractor bulbi; rpe, retinal pigmented epithelium; tm, temporal mesenchyme; vr, ventral retina

Supplementary Fig 5. Retinoic acid signaling responsiveness in the periocular region

(A-C) Macroscopic views of endogenous GFP fluorescence of *Tg:RARE^CreERT2^:R^26mTmG^* embryos. Tamoxifen (Tam) was injected into pregnant females at indicated time points. Arroheads indicate labelling in periocular mesenchyme (POM) (n>3 per condition).

(D) Strategy used to determine responsiveness of *Tg:RARE^CreERT2^* reporter in presence of RA signaling reverse agonist BMS493. BMS493 was injected to pregnant females between E10 and E11.75.

(E-H) Whole mount immunostaining for MyHC (muscle), GFP (ATRA-responsive cells) of control (E, F) and BMS493-treated embryos (G, H). (E, G) Medial views of 3D-reconstructed EOM labelling. BMS493 treatment before and after tamoxifen induction reveals a drastic decrease in GFP+ cells in the periocular region (H) compared to controls (F). Asterisks mark the location of the optic nerve (n=2).

Supplementary Fig 6. EOM insertions in the POM are altered upon ATRA-deficiency

(A-D) Immunostaining for GFP (tendon progenitors), SOX9 (periocular mesenchyme, POM), MF20 (myofibers) together with Lysotracker Red staining (Lyso, apoptosis) on coronal E12 sections of *Tg:Scx-GFP* control (A-B') and BMS493 treated (C-D') embryos. (B',D') higher magnification views of the periocular mesenchyme (POM) region. In BMS493-treated embryos, Lysotracker Red and SOX9 staining are absent in POM (D', asterisk) and SOX9 staining also missing in the RPE (D', double asterisk).

(E-P) Whole mount immunostaining for SOX9 and PITX2, SMA and Lysotracker Red Staining of E11.75 control (E, F, N), *Aldh1a3^KO* (G,H, O) and BMS493-treated (I, J, P) embryos (right eyes). In lateral views, neither *Aldh1a3^KO* or BMS493 embryos show a PITX2+ pom-ring as the controls (F). Arrowheads in (G,H,J) mark remaining expression of SOX9/PITX2 in POM of mutant or inhibitor treated embryos. Asterisks in (F,H,J) mark apoptosis spots in POM.

(N-P) Segmented medial views of periocular region of control embryos. Volumes were truncated in Z for clarity. Full view as schemes below. The PITX2+ pom-ring is continuous with the medial-pom (white arrowhead, N). In mutant and inhibitor treated embryos, residual PITX2 expression in POM is discontinous with the medial-POM.

(Q-R) Whole mount immunostaining for GFP and β-gal (*Myf5^{nlacZ}* reporter, myogenic progenitors) of E14.5 *Tg:Scx-GFP*::*Myf5$^{nlacZ/+}$* (Q, control) and *Tg:Scx-GFP*::*My5$^{nlacZ/nlacZ}$* (R, mutant) embryos (left eyes). Asterisk indicate few remaining β-gal+ cells in mutant. (Q',R') split GFP channel. Arrowheads in R indicate the correct position of tendon condensations for the 4 recti muscles although these are absent in the mutant. Lower panel, higher magnification views.

m, extraocular muscle; anl, anlage; apop, apoptosis; m-pom, medial periocular mesenchyme; l-pom, lateral periocular mesenchyme; rpe, retinal pigmented epithelium.

Tables

Table 1. BMS493 injection regimes. The same BMS493 concentration (Conc BMS) was used across injection regimes, from I to VI. Pink cases mark injection time points in the morning (mo), midday (mid) or evening (eve) between E10.5 and E12.5. The number of embryos with different muscle phenotype severity at E13.5 are shown: (no), no altered phenotype; mild (+), muscle mispatterning but overall organization retained; strong (++), strong mispatterning but with a minumum of 2 muscles split; severe (+++), no or almost absent splitting. n is the total number of embryos analyzed for a certain treatment type (n=38 embryos analyzed in total); N exp, number of times experiment was repeated.

Table 2. Antibodies and resources used in this study.

Videos

Video 1. Temporal sequence of EOM patterning. 3D-recontruction of whole mount immunostaining of the developing EOM in embryos ranging from E11.5 to E14.5. EOM were segmented from adjacent head structures and 3D-reconstructed in Imaris.

Video 2. 3D-recontruction of EOM functional unit at E11.75. Immunostaining for Muscle (MF20, Cyan) and tendon (*Tg:Scx-GFP*, Green). Apoptotic foci are visualised by Lysotracker Red. A clipping plane is added for clarity of visualisation. Note how tendon condensations start to organize radially and towards the apoptotic foci in the POM.

Video 3. 3D-recontruction of EOM functional unit at E12.5. Immunostaining for Muscle (MF20, Cyan) and tendon (*Tg:Scx-GFP*, Green). Apoptotic foci are visualised by Lysotracker Red. A clipping plane is added for clarity of visualisation. Note how tendon condensations of the recti muscles are more refined at this stage and project towards the apoptotic foci in the POM.

Video 4. 3D-recontruction of te EOM functional unit at E13.5 of a BMS treated embryo (E10.5→ E11.75). Immunostaining for Muscle (MyHC, Red) and tendon (*Tg:Scx-GFP*,

Green). Note the presence of tendon condensations at the tips of mispatterned muscle masses.

Video 5. 3D-recontruction of whole mount immunostaining of control EOM (MyHC, Red) and *Tg:Rare^{CreERT2}:R26^{mTmG}* reporter (GFP, Green) at E13.5 (Tamoxifen induction at E10.5). EOMs are shown as isosurface for clarity of visualisation.

Video 6. 3D-recontruction of whole mount immunostaining of EOMs (MyHC, Red) and *Tg:Rare^{CreERT2}:R26^{mTmG}* reporter (GFP, Green) at E13.5 (Tamoxifen induction at E10.5) on BMS-treated embryos (E10.5 → E11.75). EOMs are shown as isosurface for clarity of visualisation.

Video 7. 3D-recontruction of whole mount immunostaining of control EOM (MyHC, Red) and tendon (*Tg:Scx-GFP*, Green) at E12.5. A clipping plane has been added for clarity of visualisation.

Video 8. 3D-recontruction of whole mount immunostaining of EOM (MyHC, Red) and tendons (*Tg:Scx-GFP*, Green) of a BMS-treated embryo (E10.5 → E11.75) at E12.5. A clipping plane has been added for clarity of visualisation. Note that distribution of Scx+ cells is more difuse than for controls (Video 7), but an *Scx-GFP+* condensation is present at the tip of the inferior muscle mass.

Video 9. 3D-recontruction of whole mount immunostaining of control EOM (MYOG/MYOG, Red) and tendon (*Tg:Scx-GFP*, Green) at E14.5.

Video 10. 3D-recontruction of whole mount immunostaining of *Myf5$^{nlacZ/nlacZ}$* EOM (MYOG/MYOG, Red) and tendon (*Tg:Scx-GFP*, Green) at E14.5.

Materials and methods

Mouse strains and animal information

Animals were handled as per European Community guidelines and the ethics committee of the Institut Pasteur (CETEA) approved protocols. The following strains were previously described: *Mesp1^{Cre}* (Saga et al., 1999), *Tg:Wnt1^{Cre}* (Danielian et al., 1998), *Tg:Rare^{CreERT2}* (Schedl et al., unpublished), *R26^{Tom}* (Ai9; (Madisen et al., 2010)), *R26^{mTmG}* (Muzumdar et al., 2007), *Tg:Scx-GFP* (Pryce et al., 2007), *Myf5^{nlacZ}* (Tajbakhsh et al., 1996), *Aldh1a3^{KO}* (Dupe et al., 2003), *Rarb^{flox}* (Chapellier et al., 2002a), *Rarg^{flox}*, (Chapellier et al., 2002b), *Rdh10^{KO}* (Rhinn et al., 2011), *Pax6$^{Sey/Sey}$* (Roberts, 1967). *Rdh10^{KO}* embryos were received from the laboratory of Pascal Dollé. *Wnt1Cre:Rarb$^{fl/fl}$:Rarg$^{fl/fl}$* embryos were received from the laboratory of Valerie Dupé. *Pax6$^{Sey/Sey}$* and control embryos were received from the laboratory of James Briscoe. *Lhx2^{Cre};Lhx2$^{fl/fl}$* (Hagglund et al., 2011) were received from the laboratory of Leif Carlsson. *Shh(invC-6)2* embryos were received from the laboratory of François Spitz.

To generate experimental embryos for *Mesp1^Cre^* or *Wnt1^Cre^* together with *Tg:Scx-GFP* and
*R26^Tom^* lineage tracings, Cre/+ males were crossed with *Tg:Scx-GFP::R26^Tom/Tom^* females.
Mice were kept on a mixed genetic background C57BL/6JRj and DBA/2JRj (B6D2F1, Janvier
Labs). Mouse embryos and foetuses were collected between embryonic day (E) E10 and
E18.5, with noon on the day of the vaginal plug considered as E0.5. Pregnant females were
euthanised by cervical dislocation.

To induce recombination with the *Rare^CreERT2^:R26^mTmG^* line, 5mg of tamoxifen (Sigma
#T5648) were administered by gavage to pregnant females. A 25mg/ml stock solution in 5%
ethanol and 95% sunflower seed oil was prepared by thorough resuspention with rocking at
4°C.

To inhibit retinoic acid signaling, pregnant females of relevant genotypes were injected
intraperitoneally with 10mg/kg of BMS493 (Tocris, 3509), a pan-retinoic acid receptor (pan-
RAR) inverse agonist. A 5mg/ml BMS493 stock solution in DMSO (SIGMA, D2650) was
prepared and stored in single use aliquots at -20°C in tight cap tubes. At the time of injection,
the aliquot was thawed, 200ul of sterile PBS was added per 50ul aliquot and injected
immediately.

Immunofluorescence
Embryos were fixed 2.5h in 4% paraformaldhehyde (PFA; 15710, Electron Microscopy
Sciences) in PBS with 0,2-0,5% Triton X-100 (according to the embryonic stage) at 4°C and
washed overnight at 4°C in PBS. For cryosectioning, embryos were equilibrated in 30%
sucrose in PBS overnight at 4°C and embedded in OCT. Cryosections (16-18μm) were
allowed to dry at RT for 30 min and washed in PBS. Immunostaining was performed as
described in (Comai et al., 2019). TUNEL staining, that marks double strand breaks, was
performed with the In Situ Cell Death Detection Kit/Fluorescein (Roche, 11 684 795 910).
Slides were first pretreated with a 2:1 mix of Ethanol:Acetic Acid 5min -20C, washed twice 20
min with PBS at RT and processed for TUNEL staining as described by the manufacturer.

For whole mount immunostaining, embryos were fixed and washed as above and dehydrated
in 50% Methanol in PBS and twice in 100% Methanol, 30min each at RT and kept at -20°C
till needed. Heads were rehydrated, the periocular region was micro-dissected in PBS and
immunostaining performed as described in (Comai et al., 2019). For embryos older than
E13.5, the alternative pretreatment (containing 0.1%Tween-20, 0.1%TritonX100,
0.1%Deoxycholate, 0.1%NP40, 20%DMSO in PBS) and primary antibody immunolabelling
steps of the idisco protocol (https://idisco.info/idisco-protocol/) were generally used. In all
cases secondary antibodies were applied in blocking buffer as described in (Comai et al.,
2019) for >4 days at 4C with rocking. After immunolabelling, samples washed in
0.1%Tween/PBS, dehydrated in 50% Methanol in PBS, 100% Methanol 10min each at RT,

cleared with a mix benzyl alcohol and benzyl benzoate (BABB) and mounted for imaging as
described in (Yokomizo et al., 2012).
For detection of cell death in whole mount live tissues, LysoTracker Red staining was done,
as it reveals lysosomal activity correlated with increased cell death (Fogel et al., 2012;
Zucker et al., 1999). Embryos were quickly dissected in HBSS (Invitrogen, 14025-092),
incubated in 2ml tubes containing 5μM of LysoTracker™ Red DND-99 (Molecular Probes,
L7528) 45min at 37˚C with rocking in the dark, washed twice in PBS, fixed and processed for
cryosections or whole mount immunostaining as above.
904

Antibodies
Primary and secondary antibodies used in this study are listed in Table 2. To detect
differenting EOM we used: a-smooth muscle actin (SMA), which is transiently expressed in
differentiating myoblasts and myotubes (Babai et al., 1990; Li et al., 1996; Sawtell and
Lessard, 1989); Desmin, an early cytoskeletal muscle protein expressed in myoblasts,
myotubes and myofibers (Babai et al., 1990; Li et al., 1993) and myosin heavy chain (MF20)
to label sarcomeric myosin (Bader et al., 1982).
912

Static Imaging
A Zeiss SteREO Discovery V20 macroscope was used for imaging the endogenous
fluorescence of whole embryos at the time of dissection. For tissue sections and whole
mount immunostaining of cleared embryos a LSM700 and a LSM800 laser-scanning
confocal microscope with ZEN software (Carl Zeiss) were used.
All images were assembled in Adobe Photoshop and InDesign (Adobe Systems). Volume-3D
rendering of the z-stack series was performed in Imaris (version 7.2.1) software (Bitplane).
For ease of EOM visualization, the Z-stack volumes were first manually segmented to define
the EOM or the whole POM area using the Isosurface Imaris function. The signal outside the
isosurface was set to zero, the corresponding channel duplicated and subsequently a new
isosurface was created using automatic thresholding on the new channel.
924

925 **In Situ Hybridization**
926 Whole-mount in situ hybridization with digoxigenin-labeled antisense mRNA probes was
927 performed as described previously (Comai et al., 2014). The Aldh1a1, Aldh1a2, Aldh1a3
928 probes were previously described (Matt et al., 2005; Molotkov et al., 2006).
929

931 **micro-CT (micro computed tomography) analysis**

The tissue contrasting protocol has been adapted from the original protocol developed by (Metscher, 2009) and applied to mouse embryos as described in (Kaucka et al., 2018) and (Heude et al., 2018). For tissue contrasting, E13.5 embryos were stained in 0.5% PTA (phospho-tungstic acid) in 90% Methanol for 4 days, E15.5 embryos were stained in 0.7% PTA in 90% Methanol for 1 week.

The μCT analysis of the embryos was conducted using the GE phoenix v|tome|x L 240 (GE Sensing and Inspection Technologies GmbH, Germany), equipped with a 180 kV/15W maximum power nanofocus X-ray tube and flat panel detector DXR250 with 2048 × 2048 pixel, 200 × 200 μm pixel size. The exposure time of the detector was 900 ms in every position over 360°. Three projections were acquired and averaged in every position for reduction of the noise in μCT data. The acceleration voltage was set to 60 kV and tube current to 200 μA. The radiation was filtered by 0.2 mm of aluminium plate. The voxel size of obtained volumes (depending on a size of an embryo) appeared in the range of 2 μm - 6 μm. The tomographic reconstructions were performed using GE phoenix datos|x 2.0 3D computed tomography software (GE Sensing and Inspection Technologies GmbH, Germany). The EOM, eye and cartilages in the embryo head were segmented by an operator with semi-automatic tools within Avizo - 3D image data processing software (FEI, USA). The 3D segmented region was transformed to a polygonal mesh as a STL file, imported to VG Studio MAX 2.2 software (Volume Graphics GmbH, Germany) for surface smoothing and 3D visualisation.

Cell isolation from the periocular region

The periocular region of *Rare^CreERT2:R26^mTmG* embryos (including the eye itself) was micro-dissected and minced with small scissors inside a 2ml Epperndorf tube. 1ml of TrypLE™ Express (Invitrogen, 12604013) was added to each tube incubated for 15min at 37°C with agitation. Samples were resuspended by gently pipetting up and down 10-15 times using a P1000 pipette. Upon addition of 1ml of culture media containing (10μg/ml) DNaseI (Roche, 11284932001) , samples were spun 15' 500g at RT, pellet resuspended in 400ul of culture media containing of 20% foetal bovine serum (FBS; Gibco), 1% Penicillin-Streptomycin (15140 Gibco), 2% Ultroser G (15950-017, Pall Biosepra) in 50:50 DMEM:F12 (31966 and 31765, Gibco) and plated on individual wells of 8 well glass bottom dishes (Ibidi, 80826) coated with 1 mg/ml of Matrigel (354234, BD Biosciences). Cells were allowed to attach for 8h at 37 °C 5% CO_2, washed with PBS and fixed 15 min RT with 4% PFA/PBS. After fixation, cells were washed in PBS and permeabilized with 0.5% Triton in PBS X-100 5 min at RT. After three washes in PBS (5min each), cells were blocked with 20% goat serum in PBS 1 h at RT. Primary antibodies were added to cells in 2% goat serum in PBS for 2 h at RT or ON

968 at 4C. Cells were washed three times with PBS, incubated with secondary antibodies 1 h at
969 RT, washed in PBS and kept in PBS for imaging.
970

Statistics

The number of embryos of each genotype used for analysis is indicated in the figure legends and Table 1. The graphs were plotted and statistical analyses were performed using Prism8 (GraphPad Software, Inc). All data points are presented as mean ± SEM (error bars). Statistical tests used for analysis are indicated on the respective figure legends. P values less than 0.05 were considered significant ($*p < 0.05$; $**p < 0.01$; $***p < 0.001$).

Acknowledgements

We acknowledge funding support from the Institut Pasteur, Association Française contre le Myopathies, Agence Nationale de la Recherche (Laboratoire d'Excellence Revive, Investissement d'Avenir; ANR-10-LABX-73) and the Centre National de la Recherche Scientifique. We gratefully acknowledge the UtechS Photonic BioImaging (Imagopole), C2RT, Institut Pasteur, supported by the French National Research Agency (France BioImaging; ANR-10–INSB–04; Investments for the Future). MT, TZ and JK acknowledge the project CEITEC 2020 (LQ1601) with financial support from the Ministry of Education, Youth and Sports of the Czech Republic under the National Sustainability Programme II and Ceitec Nano+ project CZ.02.01/0.0./.0.0./16_013/0001728 under the program OP RDE. MT was financially supported by by the Brno City Municipality as a Brno Ph.D. Talent Scholarship Holder. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests

The authors declare no competing interests.

References

Babai, F., Musevi-Aghdam, J., Schurch, W., Royal, A., and Gabbiani, G. (1990). Coexpression of alpha-sarcomeric actin, alpha-smooth muscle actin and desmin during myogenesis in rat and mouse embryos I. Skeletal muscle. Differentiation *44*, 132-142.

Bader, D., Masaki, T., and Fischman, D.A. (1982). Immunochemical analysis of myosin heavy chain during avian myogenesis in vivo and in vitro. The Journal of cell biology *95*, 763-770.

Blitz, E., Sharir, A., Akiyama, H., and Zelzer, E. (2013). Tendon-bone attachment unit is formed modularly by a distinct pool of Scx- and Sox9-positive progenitors. Development *140*, 2680-2690.

Blitz, E., Viukov, S., Sharir, A., Shwartz, Y., Galloway, J.L., Pryce, B.A., Johnson, R.L., Tabin, C.J., Schweitzer, R., and Zelzer, E. (2009). Bone ridge patterning during musculoskeletal assembly is

mediated through SCX regulation of Bmp4 at the tendon-skeleton junction. Developmental cell *17*,
861-873.

Bohnsack, B.L., Gallina, D., Thompson, H., Kasprick, D.S., Lucarelli, M.J., Dootz, G., Nelson, C.,
McGonnell, I.M., and Kahana, A. (2011). Development of extraocular muscles requires early signals
from periocular neural crest and the developing eye. Archives of ophthalmology *129*, 1030-1041.

Bohnsack, B.L., Kasprick, D.S., Kish, P.E., Goldman, D., and Kahana, A. (2012). A zebrafish model of
axenfeld-rieger syndrome reveals that pitx2 regulation by retinoic acid is essential for ocular and
craniofacial development. Investigative ophthalmology & visual science *53*, 7-22.

Chambers, D., Wilson, L., Maden, M., and Lumsden, A. (2007). RALDH-independent generation of
retinoic acid during vertebrate embryogenesis by CYP1B1. Development *134*, 1369-1383.

Chapellier, B., Mark, M., Bastien, J., Dierich, A., LeMeur, M., Chambon, P., and Ghyselinck, N.B.
(2002a). A conditional floxed (loxP-flanked) allele for the retinoic acid receptor beta (RARbeta) gene.
Genesis *32*, 91-94.

Chapellier, B., Mark, M., Garnier, J.M., Dierich, A., Chambon, P., and Ghyselinck, N.B. (2002b). A
conditional floxed (loxP-flanked) allele for the retinoic acid receptor gamma (RARgamma) gene.
Genesis *32*, 95-98.

Chawla, B., Schley, E., Williams, A.L., and Bohnsack, B.L. (2016). Retinoic Acid and Pitx2 Regulate
Early Neural Crest Survival and Migration in Craniofacial and Ocular Development. Birth Defects Res
B Dev Reprod Toxicol *107*, 126-135.

Chevallier, A., and Kieny, M. (1982). On the role of the connective tissue in the patterning of the chick
limb musculature. Wilehm Roux Arch Dev Biol *191*, 277-280.

Colasanto, M.P., Eyal, S., Mohassel, P., Bamshad, M., Bonnemann, C.G., Zelzer, E., Moon, A.M., and
Kardon, G. (2016). Development of a subset of forelimb muscles and their attachment sites requires
the ulnar-mammary syndrome gene Tbx3. Dis Model Mech *9*, 1257-1269.

Comai, G., Heude, E., Mella, S., Paisant, S., Pala, F., Gallardo, M., Langa, F., Kardon, G.,
Gopalakrishnan, S., and Tajbakhsh, S. (2019). A distinct cardiopharyngeal mesoderm genetic
hierarchy establishes antero-posterior patterning of esophagus striated muscle. Elife *8*.

Comai, G., Sambasivan, R., Gopalakrishnan, S., and Tajbakhsh, S. (2014). Variations in the efficiency
of lineage marking and ablation confound distinctions between myogenic cell populations.
Developmental cell *31*, 654-667.

Couly, G.F., Coltey, P.M., and Douarin, N.M.L. (1992). The developmental fate of the cephalic
mesoderm in quail-chick chimeras. Development *114*, 1-15.

Creuzet, S., Couly, G., and Le Douarin, N.M. (2005). Patterning the neural crest derivatives during
development of the vertebrate head: insights from avian studies. J Anat *207*, 447-459.

Cunningham, T.J., and Duester, G. (2015). Mechanisms of retinoic acid signalling and its roles in
organ and limb development. Nature reviews Molecular cell biology *16*, 110-123.

Cvekl, A., and Wang, W.L. (2009). Retinoic acid signaling in mammalian eye development. Exp Eye
Res *89*, 280-291.

Danielian, P.S., Muccino, D., Rowitch, D.H., Michael, S.K., and McMahon, A.P. (1998). Modification of
gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. Curr Biol
8, 1323-1326.

Deries, M., Schweitzer, R., and Duxson, M.J. (2010). Developmental fate of the mammalian myotome.
Developmental dynamics : an official publication of the American Association of Anatomists *239*,
2898-2910.

Diogo, R., Kelly, R.G., Christiaen, L., Levine, M., Ziermann, J.M., Molnar, J.L., Noden, D.M., and
Tzahor, E. (2015). A new heart for a new head in vertebrate cardiopharyngeal evolution. Nature *520*,
466-473.

Duester, G. (2009). Keeping an eye on retinoic acid signaling during eye development. Chem Biol
Interact *178*, 178-181.

Dupe, V., Matt, N., Garnier, J.M., Chambon, P., Mark, M., and Ghyselinck, N.B. (2003). A newborn
lethal defect due to inactivation of retinaldehyde dehydrogenase type 3 is prevented by maternal
retinoic acid treatment. Proceedings of the National Academy of Sciences of the United States of
America *100*, 14036-14041.

Durand, J.P. (1976). Ocular development and involution in the European cave salamander, Proteus
anguinus laurenti. Biol Bull *151*, 450-466.

Ericsson, R., Cerny, R., Falck, P., and Olsson, L. (2004). Role of cranial neural crest cells in visceral
arch muscle positioning and morphogenesis in the Mexican axolotl, Ambystoma mexicanum.
Developmental dynamics : an official publication of the American Association of Anatomists *231*, 237-
247.

Evans, A.L., and Gage, P.J. (2005). Expression of the homeobox gene Pitx2 in neural crest is required
for optic stalk and ocular anterior segment development. Human molecular genetics *14*, 3347-3359.

Eyal, S., Kult, S., Rubin, S., Krief, S., Felsenthal, N., Pineault, K.M., Leshkowitz, D., Salame, T.M.,
Addadi, Y., Wellik, D.M.*, et al.* (2019). Bone morphology is regulated modularly by global and regional
genetic programs. Development *146*.

Fogel, J.L., Thein, T.Z., and Mariani, F.V. (2012). Use of LysoTracker to detect programmed cell death
in embryos and differentiating embryonic stem cells. J Vis Exp.

Franz-Odendaal, T.A., and Vickaryous, M.K. (2006). Skeletal elements in the vertebrate eye and
adnexa: morphological and developmental perspectives. Developmental dynamics : an official
publication of the American Association of Anatomists *235*, 1244-1255.

Gage, P.J., Rhoades, W., Prucka, S.K., and Hjalt, T. (2005). Fate maps of neural crest and mesoderm
in the mammalian eye. Investigative ophthalmology & visual science *46*, 4200-4208.

Germain, P., Gaudon, C., Pogenberg, V., Sanglier, S., Van Dorsselaer, A., Royer, C.A., Lazar, M.A.,
Bourguet, W., and Gronemeyer, H. (2009). Differential action on coregulator interaction defines
inverse retinoid agonists and neutral antagonists. Chem Biol *16*, 479-489.

Ghyselinck, N.B., Dupe, V., Dierich, A., Messaddeq, N., Garnier, J.M., Rochette-Egly, C., Chambon,
P., and Mark, M. (1997). Role of the retinoic acid receptor beta (RARbeta) during mouse development.
The International journal of developmental biology *41*, 425-447.

Gilbert, P. (1957). The origin and development of the human extrinsic ocular muscles. . Contrib
Embryol, 36: 59-78.

Gilbert, P.W. (1947). The origin and development of the extrinsic ocular muscles in the domestic cat.
Journal of morphology *81*, 151-193.

Gitton, Y., Heude, E., Vieux-Rochas, M., Benouaiche, L., Fontaine, A., Sato, T., Kurihara, Y., Kurihara,
H., Couly, G., and Levi, G. (2010). Evolving maps in craniofacial development. Seminars in cell &
developmental biology *21*, 301-308.

Grenier, J., Teillet, M.A., Grifone, R., Kelly, R.G., and Duprez, D. (2009). Relationship between neural
crest cells and cranial mesoderm during head muscle development. PloS one *4*, e4381.

Grindley, J.C., Davidson, D.R., and Hill, R.E. (1995). The role of Pax-6 in eye and nasal development.
Development *121*, 1433-1442.

Hagglund, A.C., Dahl, L., and Carlsson, L. (2011). Lhx2 is required for patterning and expansion of a
distinct progenitor cell population committed to eye development. PloS one *6*, e23387.

Harding, P., and Moosajee, M. (2019). The Molecular Basis of Human Anophthalmia and
Microphthalmia. J Dev Biol *7*.

Hasson, P. (2011). "Soft" tissue patterning: muscles and tendons of the limb take their form.
Developmental dynamics : an official publication of the American Association of Anatomists *240*,
1100-1107.

Hasson, P., DeLaurier, A., Bennett, M., Grigorieva, E., Naiche, L.A., Papaioannou, V.E., Mohun, T.J.,
and Logan, M.P. (2010). Tbx4 and tbx5 acting in connective tissue are required for limb muscle and
tendon patterning. Developmental cell *18*, 148-156.

Heude, E., Bellessort, B., Fontaine, A., Hamazaki, M., Treier, A.C., Treier, M., Levi, G., and Narboux-
Neme, N. (2015). Etiology of craniofacial malformations in mouse models of blepharophimosis, ptosis
and epicanthus inversus syndrome. Human molecular genetics *24*, 1670-1681.

Heude, E., Tesarova, M., Sefton, E.M., Jullian, E., Adachi, N., Grimaldi, A., Zikmund, T., Kaiser, J.,
Kardon, G., Kelly, R.G.*, et al.* (2018). Unique morphogenetic signatures define mammalian neck
muscles and associated connective tissues. Elife *7*.

Huang, A.H. (2017). Coordinated development of the limb musculoskeletal system: Tendon and
muscle patterning and integration with the skeleton. Developmental biology *429*, 420-428.

Huang, A.H., Riordan, T.J., Pryce, B., Weibel, J.L., Watson, S.S., Long, F., Lefebvre, V., Harfe, B.D.,
Stadler, H.S., Akiyama, H.*, et al.* (2015). Musculoskeletal integration at the wrist underlies the modular
development of limb tendons. Development *142*, 2431-2441.

Iwata, J., Suzuki, A., Pelikan, R.C., Ho, T.V., and Chai, Y. (2013). Noncanonical transforming growth
factor beta (TGFbeta) signaling in cranial neural crest cells causes tongue muscle developmental
defects. The Journal of biological chemistry *288*, 29760-29770.

Jernvall, J., Aberg, T., Kettunen, P., Keranen, S., and Thesleff, I. (1998). The life history of an
embryonic signaling center: BMP-4 induces p21 and is associated with apoptosis in the mouse tooth
enamel knot. Development *125*, 161-169.

Johnston, M.C., Noden, D.M., Hazelton, R.D., Coulombre, J.L., and Coulombre, A.J. (1979). Origins of
avian ocular and periocular tissues. Exp Eye Res *29*, 27-43.

Kardon, G. (1998). Muscle and tendon morphogenesis in the avian hind limb. Development *125*, 4019-
4032.

Kardon, G., Harfe, B.D., and Tabin, C.J. (2003). A Tcf4-positive mesodermal population provides a
prepattern for vertebrate limb muscle patterning. Developmental cell *5*, 937-944.

Kastner, P., Grondona, J.M., Mark, M., Gansmuller, A., LeMeur, M., Decimo, D., Vonesch, J.L., Dolle,
P., and Chambon, P. (1994). Genetic analysis of RXR alpha developmental function: convergence of
RXR and RAR signaling pathways in heart and eye morphogenesis. Cell *78*, 987-1003.

Kaucka, M., Petersen, J., Tesarova, M., Szarowska, B., Kastriti, M.E., Xie, M., Kicheva, A., Annusver,
K., Kasper, M., Symmons, O.*, et al.* (2018). Signals from the brain and olfactory epithelium control
shaping of the mammalian nasal capsule cartilage. Elife *7*.

Kaufman, M.H., Chang, H.H., and Shaw, J.P. (1995). Craniofacial abnormalities in homozygous Small
eye (Sey/Sey) embryos and newborn mice. J Anat *186 (Pt 3)*, 607-617.

Kish, P.E., Bohnsack, B.L., Gallina, D., Kasprick, D.S., and Kahana, A. (2011). The eye as an
organizer of craniofacial development. Genesis *49*, 222-230.

Lamb, T.D., Collin, S.P., and Pugh, E.N., Jr. (2007). Evolution of the vertebrate eye: opsins,
photoreceptors, retina and eye cup. Nature reviews Neuroscience *8*, 960-976.

Langenberg, T., Kahana, A., Wszalek, J.A., and Halloran, M.C. (2008). The eye organizes neural crest
cell migration. Developmental dynamics : an official publication of the American Association of
Anatomists *237*, 1645-1652.

Li, L., Miano, J.M., Cserjesi, P., and Olson, E.N. (1996). SM22 alpha, a marker of adult smooth
muscle, is expressed in multiple myogenic lineages during embryogenesis. Circulation research *78*,
188-195.

Li, Z., Marchand, P., Humbert, J., Babinet, C., and Paulin, D. (1993). Desmin sequence elements
regulating skeletal muscle-specific expression in transgenic mice. Development *117*, 947-959.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D.,
Hawrylycz, M.J., Jones, A.R.*, et al.* (2010). A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. Nature neuroscience *13*, 133-140.

Matsuo, T., Osumi-Yamashita, N., Noji, S., Ohuchi, H., Koyama, E., Myokai, F., Matsuo, N., Taniguchi,
S., Doi, H., Iseki, S.*, et al.* (1993). A mutation in the Pax-6 gene in rat small eye is associated with
impaired migration of midbrain crest cells. Nature genetics *3*, 299-304.

Matt, N., Dupe, V., Garnier, J.M., Dennefeld, C., Chambon, P., Mark, M., and Ghyselinck, N.B. (2005).
Retinoic acid-dependent eye morphogenesis is orchestrated by neural crest cells. Development *132*,
4789-4800.

Matt, N., Ghyselinck, N.B., Pellerin, I., and Dupe, V. (2008). Impairing retinoic acid signalling in the
neural crest cells is sufficient to alter entire eye morphogenesis. Developmental biology *320*, 140-148.

McBratney-Owen, B., Iseki, S., Bamforth, S.D., Olsen, B.R., and Morriss-Kay, G.M. (2008).
Development and tissue origins of the mammalian cranial base. Developmental biology *322*, 121-132.

McGurk, P.D., Swartz, M.E., Chen, J.W., Galloway, J.L., and Eberhart, J.K. (2017). In vivo zebrafish
morphogenesis shows Cyp26b1 promotes tendon condensation and musculoskeletal patterning in the
embryonic jaw. PLoS genetics *13*, e1007112.

Metscher, B.D. (2009). MicroCT for developmental biology: a versatile tool for high-contrast 3D
imaging at histological resolutions. Developmental dynamics : an official publication of the American
Association of Anatomists *238*, 632-640.

Molotkov, A., Molotkova, N., and Duester, G. (2006). Retinoic acid guides eye morphogenetic
movements via paracrine signaling but is unnecessary for retinal dorsoventral patterning.
Development *133*, 1901-1910.

Moreno, E., Valon, L., Levillayer, F., and Levayer, R. (2019). Competition for Space Induces Cell
Elimination through Compaction-Driven ERK Downregulation. Curr Biol *29*, 23-34 e28.

Mori, M., Ghyselinck, N.B., Chambon, P., and Mark, M. (2001). Systematic immunolocalization of
retinoid receptors in developing and adult mouse eyes. Investigative ophthalmology & visual science
42, 1312-1318.

Mori-Akiyama, Y., Akiyama, H., Rowitch, D.H., and de Crombrugghe, B. (2003). Sox9 is required for
determination of the chondrogenic cell lineage in the cranial neural crest. Proceedings of the National
Academy of Sciences of the United States of America *100*, 9360-9365.

Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L., and Luo, L. (2007). A global double-fluorescent Cre
reporter mouse. Genesis *45*, 593-605.

Nassari, S., Duprez, D., and Fournier-Thibault, C. (2017). Non-myogenic Contribution to Muscle
Development and Homeostasis: The Role of Connective Tissues. Front Cell Dev Biol *5*, 22.

Noden, D.M. (1983a). The embryonic origins of avian cephalic and cervical muscles and associated
connective tissues. Am J Anat *168*, 257-276.

Noden, D.M. (1983b). The role of the neural crest in patterning of avian cranial skeletal, connective,
and muscle tissues. Developmental biology *96*, 144-165.

Noden, D.M., and Francis-West, P. (2006). The differentiation and morphogenesis of craniofacial
muscles. Developmental dynamics : an official publication of the American Association of Anatomists
235, 1194-1218.

Nonomura, K., Yamaguchi, Y., Hamachi, M., Koike, M., Uchiyama, Y., Nakazato, K., Mochizuki, A.,
Sakaue-Sawano, A., Miyawaki, A., Yoshida, H.*, et al.* (2013). Local apoptosis modulates early
mammalian brain development through the elimination of morphogen-producing cells. Developmental
cell *27*, 621-634.

Oh, S.P., Griffith, C.M., Hay, E.D., and Olsen, B.R. (1993). Tissue-specific expression of type XII
collagen during mouse embryonic development. Developmental dynamics : an official publication of
the American Association of Anatomists *196*, 37-46.

Osumi-Yamashita, N., Ninomiya, Y., Doi, H., and Eto, K. (1994). The contribution of both forebrain and
midbrain crest cells to the mesenchyme in the frontonasal mass of mouse embryos. Developmental
biology *164*, 409-419.

Pryce, B.A., Brent, A.E., Murchison, N.D., Tabin, C.J., and Schweitzer, R. (2007). Generation of
transgenic tendon reporters, ScxGFP and ScxAP, using regulatory elements of the scleraxis gene.
Developmental dynamics : an official publication of the American Association of Anatomists *236*,
1677-1682.

Rhinn, M., and Dolle, P. (2012). Retinoic acid signalling during development. Development *139*, 843-
858.

Rhinn, M., Schuhbaur, B., Niederreither, K., and Dolle, P. (2011). Involvement of retinol
dehydrogenase 10 in embryonic patterning and rescue of its loss of function by maternal retinaldehyde
treatment. Proceedings of the National Academy of Sciences of the United States of America *108*,
16687-16692.

Rinon, A., Lazar, S., Marshall, H., Buchmann-Moller, S., Neufeld, A., Elhanany-Tamir, H., Taketo,
M.M., Sommer, L., Krumlauf, R., and Tzahor, E. (2007). Cranial neural crest cells regulate head
muscle patterning and differentiation during vertebrate embryogenesis. Development *134*, 3065-3075.

Roberts, R.C. (1967). Small eyes—a new dominant eye mutant in the mouse. Genet Res, Cam *9*,
121-122.

Rodriguez-Guzman, M., Montero, J.A., Santesteban, E., Ganan, Y., Macias, D., and Hurle, J.M.
(2007). Tendon-muscle crosstalk controls muscle bellies morphogenesis, which is mediated by cell
death and retinoic acid signaling. Developmental biology *302*, 267-280.

Rossant, J., Zirngibl, R., Cado, D., Shago, M., and Giguere, V. (1991). Expression of a retinoic acid
response element-hsplacZ transgene defines specific domains of transcriptional activity during mouse
embryogenesis. Genes & development *5*, 1333-1344.

Saga, Y., Miyagawa-Tomita, S., Takagi, A., Kitajima, S., Miyazaki, J., and Inoue, T. (1999). MesP1 is
expressed in the heart precursor cells and required for the formation of a single heart tube.
Development *126*, 3437-3447.

Sambasivan, R., Gayraud-Morel, B., Dumas, G., Cimper, C., Paisant, S., Kelly, R., and Tajbakhsh, S.
(2009). Distinct regulatory cascades govern extraocular and pharyngeal arch muscle progenitor cell
fates. Developmental cell *16*, 810-821.

Sambasivan, R., Kuratani, S., and Tajbakhsh, S. (2011). An eye on the head: the development and
evolution of craniofacial muscles. Development *138*, 2401-2415.

Sandell, L.L., Sanderson, B.W., Moiseyev, G., Johnson, T., Mushegian, A., Young, K., Rey, J.P., Ma,
J.X., Staehling-Hampton, K., and Trainor, P.A. (2007). RDH10 is essential for synthesis of embryonic
retinoic acid and is required for limb, craniofacial, and organ development. Genes & development *21*,
1113-1124.

Sawtell, N.M., and Lessard, J.L. (1989). Cellular distribution of smooth muscle actins during
mammalian embryogenesis: expression of the alpha-vascular but not the gamma-enteric isoform in
differentiating striated myocytes. The Journal of cell biology *109*, 2929-2937.

Schweitzer, R., Chyung, J.H., Murtaugh, L.C., Brent, A.E., Rosen, V., Olson, E.N., Lassar, A., and
Tabin, C.J. (2001). Analysis of the tendon cell fate using Scleraxis, a specific marker for tendons and
ligaments. Development *128*, 3855-3866.

See, A.W., Kaiser, M.E., White, J.C., and Clagett-Dame, M. (2008). A nutritional model of late
embryonic vitamin A deficiency produces defects in organogenesis at a high penetrance and reveals
new roles for the vitamin in skeletal development. Developmental biology *316*, 171-190.

Sefton, E.M., and Kardon, G. (2019). Connecting muscle development, birth defects, and evolution:
An essential role for muscle connective tissue. Current topics in developmental biology *132*, 137-176.

Serbedzija, G.N., Bronner-Fraser, M., and Fraser, S.E. (1992). Vital dye analysis of cranial neural
crest cell migration in the mouse embryo. Development *116*, 297-307.

Shimizu, M., Narboux-Neme, N., Gitton, Y., de Lombares, C., Fontaine, A., Alfama, G., Kitazawa, T.,
Kawamura, Y., Heude, E., Marshall, L.*, et al.* (2018). Probing the origin of matching functional jaws:
roles of Dlx5/6 in cranial neural crest cells. Scientific reports *8*, 14975.

Spencer, R.F., and Porter, J.D. (2006). Biological organization of the extraocular muscles. Progress in
brain research *151*, 43-80.

Sugimoto, Y., Takimoto, A., Akiyama, H., Kist, R., Scherer, G., Nakamura, T., Hiraki, Y., and
Shukunami, C. (2013). Scx+/Sox9+ progenitors contribute to the establishment of the junction
between cartilage and tendon/ligament. Development *140*, 2280-2288.

Sulik, K.K., Dehart, D.B., Johnson, C.S., Ellis, S.L., Chen, S.Y., Dunty, W.C., Jr., and Zucker, R.M.
(2001). Programmed cell death in extraocular muscle tendon/sclera precursors. Molecular vision *7*,
184-191.

Suzuki, D.G., Fukumoto, Y., Yoshimura, M., Yamazaki, Y., Kosaka, J., Kuratani, S., and Wada, H.
(2016). Comparative morphology and development of extra-ocular muscles in the lamprey and
gnathostomes reveal the ancestral state and developmental patterns of the vertebrate head.
Zoological Lett *2*, 10.

Swinehart, I.T., Schlientz, A.J., Quintanilla, C.A., Mortlock, D.P., and Wellik, D.M. (2013). Hox11
genes are required for regional patterning and integration of muscle, tendon and bone. Development
140, 4574-4582.

Tajbakhsh, S., Rocancourt, D., and Buckingham, M. (1996). Muscle progenitor cells failing to respond
to positional cues adopt non-myogenic fates in myf-5 null mice. Nature *384*, 266-270.

Thompson, H., Griffiths, J.S., Jeffery, G., and McGonnell, I.M. (2010). The retinal pigment epithelium
of the eye regulates the development of scleral cartilage. Developmental biology *347*, 40-52.

Tokita, M., and Schneider, R.A. (2009). Developmental origins of species-specific muscle pattern.
Developmental biology *331*, 311-325.

Twitty, V.C. (1932). Influence of the eye on the growth of its associated structures, studied by means
of heteroplastic transplantation. Journal of Experimental Zoology *61*, 333-374.

von Scheven, G., Alvares, L.E., Mootoosamy, R.C., and Dietrich, S. (2006). Neural tube derived
signals and Fgf8 act antagonistically to specify eye versus mandibular arch muscles. Development
133, 2731-2745.

Williams, A.L., and Bohnsack, B.L. (2015). Neural crest derivatives in ocular development: discerning
the eye of the storm. Birth defects research Part C, Embryo today : reviews *105*, 87-95.

Yamamoto, Y., Stock, D.W., and Jeffery, W.R. (2004). Hedgehog signalling controls eye degeneration
in blind cavefish. Nature *431*, 844-847.

Yokomizo, T., Yamada-Inagawa, T., Yzaguirre, A.D., Chen, M.J., Speck, N.A., and Dzierzak, E.
(2012). Whole-mount three-dimensional imaging of internally localized immunostained cells within
mouse embryos. Nat Protoc *7*, 421-431.

Young, G.C. (2008). Number and arrangement of extraocular muscles in primitive gnathostomes:
evidence from extinct placoderm fishes. Biology letters *4*, 110-114.

Zein, W.M., Lewanda, A.F., Traboulsi, E.I. and Jabs, E.W. (2012). Ocular Manifestations of
Syndromes with Craniofacial Abnormalities. In Genetic Diseases of the Eye (2 ed) E.I. Traboulsi, ed.
(Oxford University Press).

Ziermann, J.M., Diogo, R., and Noden, D.M. (2018). Neural crest and the patterning of vertebrate
craniofacial muscles. Genesis *56*, e23097.

Zucker, R.M., Hunter, E.S., 3rd, and Rogers, J.M. (1999). Apoptosis and morphology in mouse
embryos by confocal laser scanning microscopy. Methods *18*, 473-480.

1289

Figure 1

Neural Crest: Wnt1^Cre;R26^Tom;Scx-GFP

Muscle	GFP	SOX9	PITX2

Coronal section - ventral

A — E13.5, eom
B — t, mct, rpe, t
C — ti, rpe
D — r, le, eom, ti, ce, 200μm
l-pom, m-pom, mct

Mesoderm: Mesp1^Cre;R26^Tom;Scx-GFP

Muscle	GFP	SOX9	PITX2

Coronal section - dorsal

E — E13.5, eom, az, mct
F — az, mct
G — az
H — eom, az, 200μm

Dorsal (E-H)
Ventral (A-D)

insertion
Figure2 (A-J)
M ← → L

origin

Muscle — Other tissues
Wnt1 Lineage — Mesp1 Lineage
Neural crest derived MCT — Mesoderm derived MCT
Neural crest derived attachment — Mesoderm derived attachment

A

copies of DR5 RARE
of RARβ2 promoter

ROSA26 locus

RA-responsive cells are mGFP+

B

Tamox
or

Bulk
or

E9.75   E10.5   E11.75   E12.5

C

% GFP+ cells

E9.75 → E11.5   E9.75 → E12.5   E10.5 → E12.5

CT   Myog   CT   Myog   CT   Myog

D Rare^CreERT2:R26^mTmG (Tam E9.75)

WM - medial view

D'

E11.75   100μm

E

E12

E'

F Rare^CreERT2:R26^mTmG (Tam E9.75)

WM - lateral view

F'

pom-r   100μm

G

pom-r

G'

3D-rec EOM   GFP   SOX9   GFP

H Rare^CreERT2:R26^mTmG (Tam E9.75)

coronal sections

E11.75   **H'**

100μm

MYOD/MYOG   GFP   Hoechst

I Wnt1^Cre: RARβ^fl/+γ^fl/+

3D-rec EOM

E12.5   100μm
medial   **I'** lateral

J Wnt1^Cre:RARβ^fl/fl γ^fl/fl

medial   **J'** lateral

H-H'

F-G'
I', J'

Tnc GFP Muscle Hoechst

PITX2 GFP Muscle Hoechst

SOX9 GFP Muscle Hoechst

Whole mount

A

Aldh1a1 — E10.5 (dr, le), E11.5 (ce, dr, le), E12.5 (ce, dr, le)

Aldh1a2 — tm

Aldh1a3 — E10.5 (rpe, dr, vr, np), E11.5 (ce, dr, vr, nlg), E12.5 (ce, vr, lgb)

200μm

Section

A' E12.5 *Aldh1a2*

A" MF20 ALDH1A2

B E13.5 — eyeball optic nerve lens

ctrl   Aldh1a3^KO   BMS (E10.5 - E11.75)

100μm

C E13.5 — ctrl Aldh1a3^KO BMS (E10.5 - E11.75)

nasal capsule   trabecular cartilage

ventral   lateral   dorsal   200μm

D I, II, III, IV, V, VIa, VIb, Aldh1a3^KO

3D-rec EOM   3D-rec EOM

100μm

D' 1, 2

E ctrl E13.5 — sr, so, rb, lr, mr, ir, io

F Rdh10^KO

200μm

MyHC

Rare^{CreERT2}:R26^{mTmG}

Tam E9.75	Tam E10.5	Tam E11.5

A E12.5

500μm

B

C

Responsive POM

D Tamox

E10.0 E10.75 E11.0 E11.75 E13.5

BMS 5mg/kg

Dissection
WMIF

Rare^{CreERT2}:R26^{mTmG}

ctrl

E E13.5

F

100μm

BMS
(E10.0 - E11.75)

G

H

3D-rec EOM
medial view

MyHC GFP

ctrl | **BMS493**

coronal sections

A — B — E12 — 100μm

C — D — MF20 — eom

C' — D' — Lyso — rpe — apop — * — **

SCX SOX9 Lyso/MF20

Myf5^nlacZ/+ | **Myf5^nlacZ/nlacZ**

WM - medial view

Q — R — E14.5 — 100μm

Q' — R'

SCX β-gal — 50μm

A-D'
a m / v d / p

E-M

ctrl | **Aldh1a3^KO** | **BMS493**

WM - lateral view

E — G — I — E11.75 — SOX9 Lyso

F — H — J — PITX2 Lyso — 100μm

WM-medial view (half volume)

N — O — P — PITX2 SMA — m-pom — anl — pom-r — 100μm

POM scheme (right eye)

m-pom — pom-r — anl

Table 1

Exp type	Conc BMS	Injections E10.5 mo	E10.5 mid	E10.5 eve	E11.5 mo	E11.5 mid	E11.5 eve	E12.5 mo	E12.5 mid	E12.5 eve	Muscle phenotype (n° embryos) no	+	++	+++	n	N exp
I	10 mg/Kg	■		■	■		■	■		■			3	3	6	4
II		■		■	■		■					1	7	1	9	2
III			■	■	■		■						4	5	9	2
IV				■								2	2		4	1
V			■		■		■	■		■	3	1	1		5	3
VI					■		■	■		■	2	3			5	1

total 38

Table2. Key Resources

REAGENT TYPE OR RESOURCE	DESIGNATION	SOURCE	Identifiers	Additional information
Primary antibody	Rabbit polyclonal anti-Aldh1A3	Sigma	Cat# HPA046271, RRID:AB_10965992	IF (1:100)/Citrate unmasking pH6
Primary antibody	Rabbit polyclonal anti-Aldh1A2	Abcam	Cat# ab96060, RRID:AB_10679336	IF (1:400)/Citrate unmasking pH6
Primary antibody	Chicken polyclonal anti-β-gal	Abcam	Cat. #: ab9361 RRID:AB_307210	IF (1:1000)
Primary antibody	Rabbit polyclonal anti-β-gal	MP Biomedicals	Cat. #: MP 559761 RRID:AB_2687418	IF (1:1500)
Primary antibody	Rabbit polyclonal anti-Collagen XII	from M. Koch	N/A	IF (1:3000)
Primary antibody	Mouse monoclonal IgG1 anti-Desmin	Dako	Cat# M0760, RRID:AB_2335684	IF (1:500)
Primary antibody	Chicken polyclonal anti-GFP	Abcam	Cat# ab13970, RRID:AB_300798	IF (1:1000)
Primary antibody	Rabbit polyclonal anti-GFP	Thermo Fisher Scientific	Cat# A-11122, RRID:AB_221569	IF (1:1000)
Primary antibody	Mouse monoclonal IgG1 anti-Myod	BD Biosciences	Cat# 554130, RRID:AB_395255	IF (1:100)
Primary antibody	Mouse monoclonal IgG1 anti-Myog	DSHB	Cat. #: F5D, RRID:AB_2146602	IF (1:30)
Primary antibody	Mouse monoclonal IgG2b anti-MF20	DSHB	Cat# MF 20, RRID:AB_2147781	IF (1:30)
Primary antibody	Rabbit polyclonal anti-MyHC	from G.Cossu	N/A	IF (1:750)
Primary antibody	Mouse monoclonal IgG1 anti-Pax7	DSHB	Cat. #: AB_528428, RRID:AB_528428	IF (1:20)
Primary antibody	Rabbit polyclonal anti-Pitx2	CapraScience	Cat# PA-1020, RRID:AB_2819162	IF (1:1000)
Primary antibody	Mouse monoclonal IgG2a anti-Smooth muscle actin-Cy3 conjugated	Sigma-Aldrich	Cat# C6198, RRID:AB_476856	IF (1:500)
Primary antibody	Rabbit polyclonal anti-Sox9	Millipore	Cat# AB5535, RRID:AB_2239761	IF (1:1500)
Primary antibody	Rat monoclonal IgG1 anti-Tenascin C	Thermo Fisher Scientific	Cat# MA1-26778, RRID:AB_2256026	IF (1:200)
Primary antibody	Mouse monoclonal IgG1 anti-Tnnt3	Sigma-Aldrich	Cat# T6277, RRID:AB_261723	IF (1:200)
Seconday antibody	Alexa Fluor® 633 Goat Anti-Chicken IgG (H+L)	Thermo Fisher Scientific	Cat# A-21103, RRID:AB_2535756	IF (1:500)
Seconday antibody	Alexa Fluor® 488 Goat Anti-Chicken IgG (H+L)	Thermo Fisher Scientific	Cat# A-11039, RRID:AB_2534096	IF (1:500)
Seconday antibody	Cy3 AffiniPure F(ab)'2 Fragment Donkey Anti-Chicken IgY (IgG)	Jackson ImmunoResearch Labs	Cat# 703-166-155, RRID:AB_2340364	IF (1:500)
Seconday antibody	Alexa Fluor® 488 AffiniPure F(ab)2 Fragment Donkey Anti-Chicken IgY (IgG)	Jackson ImmunoResearch Labs	Cat# 703-546-155, RRID:AB_2340376	IF (1:500)
Seconday antibody	Alexa Fluor® 633 F(ab')2 Fragment of Goat Anti-Rabbit IgG	Thermo Fisher Scientific	Cat# A-21072, RRID:AB_2535733	IF (1:500)
Seconday antibody	Alexa Fluor® 555 F(ab')2 Fragment of Goat Anti-Rabbit IgG	Thermo Fisher Scientific	Cat# A-21430, RRID:AB_2535851	IF (1:500)
Seconday antibody	Alexa Fluor® 488 F(ab')2 Fragment of Goat Anti-Rabbit IgG	Thermo Fisher Scientific	Cat# A-11070, RRID:AB_2534114	IF (1:500)
Seconday antibody	Alexa Fluor® 555 Goat Anti-Rat IgG (H+L)	Thermo Fisher Scientific	Cat# A-21434, RRID:AB_2535855	IF (1:500)
Seconday antibody	Alexa Fluor® 488 Goat Anti-Rat IgG (H+L)	Thermo Fisher Scientific	Cat# A-11006, RRID:AB_2534074	IF (1:500)
Seconday antibody	Alexa Fluor® 633 Goat Anti-Mouse IgG1(γ1)	Thermo Fisher Scientific	Cat# A-21126, RRID:AB_2535768	IF (1:500)
Seconday antibody	Cy3-AffiniPure Goat Anti-Mouse IgG1(γ1)	Jackson ImmunoResearch Labs	Cat# 115-165-205, RRID:AB_2338694	IF (1:500)
Seconday antibody	Alexa Fluor®488 AffiniPure Goat Anti-Mouse IgG1(γ1)	Jackson ImmunoResearch Labs	Cat# 115-545-205, RRID:AB_2338854	IF (1:500)
Seconday antibody	Alexa Fluor® 633 Goat Anti-Mouse IgG2b (γ2b)	Thermo Fisher Scientific	Cat# A-21146, RRID:AB_2535782	IF (1:500)
Seconday antibody	Cy3-AffiniPure Goat Anti-Mouse IgG2b (γ2b)	Jackson ImmunoResearch Labs	Cat# 115-165-207, RRID:AB_2338696	IF (1:500)
Seconday antibody	Alexa Fluor® 488 Goat Anti-Mouse IgG2b (γ2b)	Thermo Fisher Scientific	Cat# A-21141, RRID:AB_2535778	IF (1:500)
Chemical	Tamoxifen	Sigma-Aldrich	Cat#T5648	
Chemical	BMS493	Tocris	Cat#3509	
Commercial assay, kit	In Situ Cell Death Detection Kit - Fluorescein	Roche	Cat#11684795910	
Commercial assay, kit	LysoTracker™ Red DND-99	Thermo Fisher Scientific	Cat#L7528	
strain (*Mus musculus*)	B6D2F1/JRj	Janvier	N/A	
genetic reagent (*M. musculus*)	$R26^{mTmG}$	PMID:17868096	MGI:3716464	(Muzumdar et al., 2007)
genetic reagent (*M. musculus*)	$R26^{Tom}$ (Ai9)	PMID:20023653	MGI:3809523	(Madisen et al., 2010)
genetic reagent (*M. musculus*)	Mesp1Cre	PMID:10393122	MGI:2176467	(Saga et al., 1999)
genetic reagent (*M. musculus*)	Tg:Wnt1Cre	PMID:9843687	MGI:2386570	(Danielian et al., 1998)
genetic reagent (*M. musculus*)	Tg:Scx-GFP	PMID:17497702	MGI:3717419	(Pryce et al., 2007)
genetic reagent (*M. musculus*)	$Myf5^{nlacZ}$		MGI:5581636	(Tajbakhsh et al., 1996)
genetic reagent (*M. musculus*)	$Aldh1a3^{KO}$	PMID:14623956	MGI:3043947	(Dupe et al., 2003)
genetic reagent (*M. musculus*)	$Rarb^{flox}$ (RARbetaL2)	PMID:11857787	MGI:2386108	(Chapellier et al., 2002a)
genetic reagent (*M. musculus*)	$Rarg^{flox}$ (RARgammaL2)	PMID:11857788	MGI:2386111	(Chapellier et al., 2002b)
genetic reagent (*M. musculus*)	$Rdh10^{KO}$ (Rdh10L-)	PMID:21930923	MGI:5294926	(Rhinn et al., 2011)
genetic reagent (*M. musculus*)	$Pax6^{Sey}$	N/A	MGI:1856155	(Roberts, 1967)
genetic reagent (*M. musculus*)	$Lhx2^{Cre}$	PMID:21886788	MGI:5300954	(Hagglund et al., 2011)
	$Lhx2^{flox}$	PMID:20386748	MGI:4453328	(Tornqvist et al., 2010)