General DNA methylation patterns and environmentally-induced differential methylation in the eastern oyster (*Crassostrea virginica*)

Yaamini R. Venkataraman[1], Alan M. Downey-Wall[2], Justin Ries[2], Isaac Westfield[2], Samuel J. White[1], Steven B. Roberts[1], Kathleen E. Lotterhos[2]

[1]School of Aquatic & Fishery Sciences, University of Washington, 1122 NE Boat St, Seattle, WA, USA
[2]Department of Marine and Environmental Sciences, Northeastern University, 430 Nahant Road, Nahant, MA 01908

Abstract

Epigenetic modification, specifically DNA methylation, is one possible mechanism for intergenerational plasticity. Before inheritance of methylation patterns can be characterized, we need a better understanding of how environmental change modifies the parental epigenome. To examine the influence of experimental ocean acidification on eastern oyster (*Crassostrea virginica*) gonad tissue, oysters were cultured in the laboratory under control (491 ± 49 µatm) or high (2550 ± 211 µatm) pCO_2 conditions for four weeks. DNA from reproductive tissue was isolated from five oysters per treatment, then subjected to bisulfite treatment and DNA sequencing. Irrespective of treatment, DNA methylation was primarily found in gene bodies with approximately 22% of the genome predicted to be methylated In response to elevated pCO_2 we found 598 differentially methylated loci primarily overlapping with gene bodies. A majority of differentially methylated loci were in exons (61.5%) with less intron overlap (31.9%). While there was though there was no evidence of a significant tendency for the genes with differentially methylated loci to be associated with distinct biological processes, the concentration of these loci in gene bodies, including genes involved in protein ubiquitination and biomineralization suggests DNA methylation may be important for transcriptional control in response to ocean acidification. Understanding how experimental ocean acidification conditions modify the oyster epigenome, and if these modifications are inherited, allows for a better understanding of how ecosystems will respond to environmental change.

Keywords: eastern oyster, DNA methylation, epigenetics, ocean acidification, protein ubiquitination

Introduction

As increased anthropogenic carbon dioxide is expected to create adverse conditions for calcifying organisms (IPCC 2019), efforts have been made to understand how ocean acidification impacts ecologically and economically important organisms like bivalves (Parker et al., 2013; Ekstrom et al., 2015). Bivalve species are sensitive to reduced aragonite saturation associated with ocean acidification, with larvae being particularly vulnerable (Barton et al., 2012; Waldbusser et al., 2014). Shell structure may be compromised in larvae, juveniles, and adults (Gazeau et al., 2007; Kurihara et al., 2007; Beniash et al., 2010; Ries, 2011). Aside from affecting calcification and shell growth, ocean acidification can impact protein synthesis, energy production, metabolism, antioxidant responses, and reproduction (Tomanek et al., 2011; Timmins-Schiffman et al., 2014; Dineshram et al., 2016; Boulais et al., 2017; Omoregie et al., 2019). Additionally, adult exposure to ocean acidification may impact their larvae (reviewed in (Ross et al., 2016; Byrne et al., 2019). For example, adult Manila clams (*Ruditapes philippinarum*) and mussels (*Musculista senhousia*) reproductively conditioned in high pCO_2 waters yield offspring that exhibit significantly faster development or lower oxidative stress protein activity in those same conditions (Zhao et al., 2018, 2019). In contrast, northern quahog (hard clam; *Mercenaria mercenaria*) and bay scallop (*Argopecten irradians*) larvae may be more vulnerable to ocean acidification and additional stressors when parents are reproductively conditioned in high pCO_2 waters (Griffith and Gobler, 2017). Some species exhibit both positive and negative carryover effects (ex. *Saccostrea glomerata*; (Parker et al., 2012, 2017). Intergenerational effects have also been documented when adult exposure to ocean acidification does not coincide with reproductive maturity (ex. *Crassostrea gigas*; (Venkataraman et al., 2019)). Although intergenerational carryover effects are now at the forefront of ocean acidification research in bivalve species, it is still unclear how these responses manifest.

Epigenetics is the next frontier for understanding how environmental memory may modulate phenotypic plasticity across generations (Eirin-Lopez and Putnam, 2018). Epigenetics refers to changes in gene expression that do not arise from the original DNA sequence, with methylation of cytosine bases being the most studied mechanism (Bird, 2002; Deans and Maggert, 2015). Unlike highly methylated vertebrate genomes, marine invertebrate taxa have spurious methylation throughout their genomes, similar to a mosaic pattern (Suzuki and Bird, 2008). Genes that benefit from stable transcription, such as housekeeping genes, tend to be more methylated, while environmental response genes that are less methylated are prone to more spurious transcription and alternative splicing patterns, thereby possibly increasing phenotypic plasticity (Roberts and Gavery, 2012; Dimond and Roberts, 2016; Gatzmann et al., 2018). Increased levels of DNA methylation can also correlate with increased transcription. Several loci-level resolution studies in *C. gigas* demonstrate a positive association between DNA methylation and gene expression that is consistent across cell types (Roberts and Gavery, 2012; Gavery and Roberts, 2013; Olson and Roberts, 2014). Since DNA methylation could provide a direct link between environmental conditions and phenotypic plasticity via influencing gene activity, elucidating how invertebrate methylomes respond to abiotic factors is crucial for understanding potential acclimatization mechanisms (Bossdorf et al., 2008; Hofmann, 2017).

82
While bivalve species have been used as model organisms to characterize marine invertebrate
methylomes, how ocean acidification affects bivalve DNA methylation is poorly understood.
Methylation responses to ocean acidification have been studied, however, in multiple coral
species. When placed in low pH conditions (7.6-7.35), *Montipora capitata* did not demonstrate
any differences in calcification, metabolic profiles, or DNA methylation in comparison to clonal
fragments in ambient pH (7.9-7.65) (Putnam et al., 2016). DNA methylation increased in another
coral species, *Pocillopora damicornis*, in addition to reduced in calcification and more
differences in metabolic profiles (Putnam et al., 2016). The coral *Stylophora pistillata* also
demonstrates increased global methylation as pH decreases ($pH_{treatment}$ = 7.2, 7.4, 7.8; $pH_{control}$ =
8.0), with methylation reducing spurious transcription (Liew et al., 2018b). Combined whole
genome bisulfite sequencing and RNA sequencing revealed differential methylation and
expression of growth and stress response pathways controlled differences in cell and polyp size
between treatments (Liew et al., 2018b). The association between DNA methylation and
phenotypic differences in these corals demonstrates that epigenetic regulation of genes is
potentially important for acclimatization and adaptation to environmental perturbation. Recent
examination of *C. virginica* methylation patterns in response to a natural salinity gradient
suggests that differential methylation may modulate environmental response in this species
(Johnson and Kelly, 2019). As bivalves may exhibit a similar response to ocean acidification,
continued study could elucidate if DNA methylation could be a mechanism for intergenerational
effects.
103
There is evidence that suggests that methylation patterns can be inherited in marine
invertebrates. For example, purple sea urchin (*Strongylocentrotus purpuratus*) offspring have
methylomes that reflect maternal rearing conditions (Strader et al., 2019). Different parental
temperature and salinity regimes influence larval methylomes in *Platygyra daedalea* (Liew et al.,
2018a). In the Pacific oyster (*Crasostrea gigas*), parental exposure to pesticides influence DNA
methylation in spat, even though the spat were not exposed to these conditions (Rondon et al.,
2017). Methylation changes in reproductive tissue are likely the ones that could be inherited,
and may play a role in carryover effects. Before determining if DNA methylation is a viable
mechanism for altering subsequent generation phenotypes, the epigenome of bivalve
reproductive tissue in response to ocean acidification must be characterized.
114
The present study is the first to determine if ocean acidification induces differential methylation
in reproductive tissue in the eastern oyster (*Crassostrea virginica*). Adult *C. virginica* were
exposed to control or elevated pCO_2 conditions. We hypothesize that ocean acidification will
induce differential methylation in *C. virginica* gonad tissue. Genes that contain differentially
methylated loci will have biological functions that allow for acclimatization to environmental
perturbation. Understanding how experimental ocean acidification conditions modify the oyster
epigenome, and if these modifications are inherited, allows for a better understanding of how
ecosystems will respond to environmental change.

Methods

Experimental Design

Adult *C. virginica* (9.55 cm ± 0.45) were collected from an intertidal oyster reef in Plum Island Sound, MA (42.681764, -70.813498) in mid-July 2016. The oysters were transported to the Marine Science Center at Northeastern University (Nahant, MA), where they were cleaned and randomly assigned to one of six flow-through tanks (50L) maintained at ambient seawater conditions. Oysters were acclimated for 14 days under control conditions (500 µatm; 14-15℃) before initiating a 28-day experimental exposure. Half of the tanks remained at control pCO_2 conditions (500 µatm, $\Omega_{calcite} > 1$), while the other half were ramped up to elevated pCO_2 conditions (2500 µatm, $\Omega_{calcite} < 1$) over 24 hours. This elevated treatment is consistent with observations in other estuarine ecosystems that oysters inhabit (Feely et al., 2010). Five oysters were collected from each treatment at the end of the 28 day exposure. They were immediately dissected with gonadal tissue harvested and immediately flash frozen.

Treatment conditions were replicated across three tanks, with oysters distributed evenly among tanks (1-2 oysters per tank). Each tank had an independent flow-regulator that delivered fresh, natural seawater at approximately 150 ml min_{-1}. Carbonate chemistry was maintained independently for each tank by mixtures of compressed CO_2 and compressed air at flow rates proportional to the target pCO_2 conditions. Gas flow rates were maintained with *Aalborg* digital solenoid-valve-controlled mass flow controllers (Model GFC17, precision = 0.1mL/min). Within a treatment, tanks were replenished with fresh seawater and each tank was independently bubbled with its own mixed gas stream, with partial recirculation and filtration with other tanks in the treatment. As a result, the carbonate chemistry (i.e., the independent variable by which the treatments were differentiated) of the replicate tanks were slightly different from each other, which is evidence of their technical independence. Temperature was maintained at 15℃ using Aqua Euro USA model MC-1/4HP chillers coupled with 50-watt electric heaters. Average salinity was determined by the incoming natural seawater and reflected ambient ocean salinity of Massachusetts Bay near the Marine Science Center (Latitude = 42.416100, Longitude = -70.907737). Oysters were fed 2.81 mL/day of a 10% Shellfish Diet 1800 twice daily following Food and Agriculture Organization's best practices for oysters (Helm and Bourne, 2004).

Measurement and control of seawater carbonate chemistry

The carbonate chemistry of tanks was controlled by bubbling mixtures of compressed CO_2 and compressed air at flow rates proportional to the target pCO_2 conditions. The control pCO_2 treatments were maintained by bubbling compressed ambient air only.

Temperature, pH, and salinity of all replicate tanks was measured three times per week for the duration of the experiment. Temperature was measured using a glass thermometer to 0.1℃ accuracy, pH was measured using an *Accumet* solid state pH electrode (precision = 1mV), salinity was measured using a *YSI 3200* conductivity probe (precision = 0.1 ppt). Every two weeks, seawater samples were collected from each replicate tank for analysis of dissolved inorganic carbon (DIC) and total alkalinity (A_T). Samples were collected in 250 mL borosilicate glass bottles sealed with a greased stopper, immediately poisoned with 100 µL saturated $HgCl_2$ solution, and then refrigerated. Samples were analyzed for DIC via coulomtery and Alk_T via

closed-cell potentiometric Gran Titration with a VINDTA 3C (Marianda Corporation). Other
carbonate system parameters, including $\Omega_{calcite}$, pH, and pCO_2, were calculated from DIC, A_T,
salinity, and temperature using CO2SYS software version 2.1 (Lewis and Wallace, 1998; Van
Heuven et al., 2011), using the seawater pH scale (mol/kg-SW) with K1 and K2 values from
(Roy, Rabindra N Roy, Lakshimi N Vogel, Kathleen M Porter-Moore, C Pearson, Tara Good,
Catherine E Millero, Frank J Campbell, Douglas M, 1993), a KHSO4 value from (Dickson,
1990), and a [B]Tvalue from (Lee, Kitach Tae-Wook, Kim Byrne, Robert H Millero, Frank J
Feely, Richard A Liu, Yong-Ming, 2010).

MBD-BS Library Preparation
DNA was isolated from five gonad tissue samples per treatment using the E.Z.N.A. Mollusc Kit
(Omega) according to the manufacturer's instructions. Isolated DNA was quantified using a
Qubit dsDNA BR Kit (Invitrogen). DNA samples, ranging from 12.8 ng/µL to 157 ng/µL, were
placed in 1.5 mL centrifuge tubes and sonicated using a QSONICA CD0004054245 (Newtown,
CT) in 30 second interval periods over ten minutes at 4 ºC and 25% intensity. Shearing size
(350bp) was verified using a 2200 TapeStation System (Agilent Technologies). Samples were
enriched for methylated DNA using MethylMiner kit (Invitrogen). Library preparation and
sequencing was performed by ZymoReearch using Pico Methyl-Seq Library Prep Kit (Cat.
#D5455) to generate 100bp paired-end reads on the HiSeq1500 sequencer (Illumina, Inc.).

Global Methylation Characterization
Sequences were trimmed with 10 bp removed from both the 5' and 3' ends using TrimGalore!
v.0.4.5 (Martin, 2011). Quality of sequences was assessed with FastQC v.0.11.7 (Andrews,
2010). The *C. virginica* genome (NCBI Accession GCA_002022765.4) was prepared using
Bowtie 2-2.3.4 (Linux x84_64 version; (Langmead and Salzberg, 2012)) within the
`bismark_genome_preparation` function in Bismark v.0.19.0 (Krueger and Andrews, 2011).
Trimmed sample sequences were then aligned to the genome using Bismark v.0.19.0 (Krueger
and Andrews, 2011) with non-directionality specified alignment score set using `-score_min`
`L,0,-1.2`. Deduplicated alignments files (ie .bam) were sorted and indexed using SAMtools
v.1.9 (Li et al., 2009). Methylation calls were extracted from deduplicated files using
`bismark_methylation_extractor`.
195
Various *C. virginica* genome feature tracks were created for downstream analyses using
BEDtools v2.26.0 (Quinlan and Hall, 2010). Genes, mRNA, and exons were derived directly
from the *C. virginica* genome on NCBI (Gómez-Chiarri et al., 2015). The complement of the
exon track was used to identify introns. Putative promoter regions were defined as those 1kb
upstream of transcription start sites. Putative transposable elements were identified using
RepeatMasker (v4.07) with RepBase-20170127 and RMBlast 2.6.0 (Smit et al., 2013; Bao et al.,
2015). All species available in RepBase-20170127 were used to identify transposable elements.
203
Overall *C. virginica* gonad methylation patterns were characterized using information from all
samples. Individual CpG dinucleotides with at least 5x coverage in each sample were classified

as methylated (≥ 50% methylation), sparsely methylated (10-50% methylation), or unmethylated

(< 10% methylation). The locations of all methylated CpGs were characterized in relation to

exons, introns, transposable elements, and putative promoter regions. We tested the null

hypothesis that there was no association between the genomic location of CpG loci and

methylation status (all CpGs versus methylated CpGs) with a chi-squared contingency test

(`chisq.test` in R Version 3.5.0).

Differential Methylation Analysis

Differential methylation analysis for individual loci was performed using `methylKit` v.1.7.9 in R

(Akalin et al., 2012) using deduplicated, sorted bam files as input. Only loci with at least 5x

coverage in each sample were considered for analysis. Methylation differences between

treatments were obtained for all loci in the CpG background using `calculateDiffMeth`, a

logistic regression built into `methylKit`. The logistic regression models the log odds ratio

based on the proportion of methylation at each locus:

220
$$log(\frac{\pi_i}{1 - \pi_i}) = \beta_0 + \beta_1 Treatment_i$$

A differentially methylated locus was defined as individual CpG dinucleotide with at least a 50%

methylation change between treatment and control groups, and a q-value < 0.01 based on

correction for false discovery rate with the SLIM method (Wang et al., 2011). Hypermethylated

DML were defined as those with significantly higher percent methylation in oysters exposed to

high pCO_2, and hypomethylated DML with significantly lower percent methylation in the high

pCO_2 treatment. A Principal Components Analysis (PCA) was used to compare oyster sample

methylation profiles between treatments for all CpG loci and for differentially methylated loci

(DML). The location of DML were characterized in relation to exons, introns, transposable

elements, and putative promoter regions using BEDtools `intersect` v2.26.0. The position of

DML within genes was characterized by scaling the position of the DML to a hypothetical gene

ranging from 0 to 100 bp. Loci that did not overlap with the aforementioned genomic features

were also identified. A chi-squared contingency test was used to test the null hypothesis of no

association between genomic location and methylation status between MBD-enriched CpGs

and DML.

Gene-specific methylation profiles were analyzed using a pipeline modified from (Liew et al.,

2018b) to determine if how whole gene methylation patterns might be influenced by ocean

acidification.For each sample, loci with 5x coverage were annotated using `intersectBED`.

Annotations included gene start and end positions, along with start and end positions for the

longest constituent mRNA. Median percent methylation was then calculated for each gene in a

sample. Any gene that was not present in all ten samples was excluded from downstream

analysis. A non-parametric Kruskal-Wallis test was used to compare median percent

methylation of a gene between pCO_2 treatments. Differentially methylated genes (DMG) were

defined as those with corrected *P*-values < 0.05.

Enrichment Analysis

Functional enrichment analyses were used to determine if any biological processes were overrepresented in genes based on individual CpG methylation levels. Enrichment was conducted with GO-MWU, a rank-based gene enrichment method initially developed for analyzing transcriptomics data (Wright et al., 2015). Instead of only using genes with DML, GO-MWU identifies GO categories that are overrepresented by genes with any hyper- or hypomethylated CpGs, allowing for more data to contribute to any trends. GO-MWU scripts and a gene ontology database were downloaded from the GO-MWU Github repository (https://github.com/z0on/GO_MWU).

A gene list, containing Genbank IDs and all associated gene ontology terms, and a table of significance measures, with Genbank IDs and signed log *P*-values for all CpGs determined by `methylKit` during the identification of DML, were used as GO-MWU analysis inputs. For the table of significance measures, *P*-values were determined by filtering methylation differences between treatments for the CpG background such that only the CpG locus with the smallest *P*-value within each gene was used in downstream enrichment. Adjusted *P*-values from `methylKit` were converted to signed log *P*-values, such that loci with positive methylation differences had positive log *P*-values, and loci with negative methylation differences had negative log *P*-values. To match the Genbank IDs to CpG loci within mRNAs and create the gene list, overlaps between the *C. virginica* mRNA track from NCBI and the CpG background used in `methylKit` were obtained using BEDtools `intersect` v2.26.0. The mRNAs were then annotated with Uniprot Accession codes using a BLASTx search (v.2.2.29; (Gish and States, 1993; UniProt Consortium, 2019)(Gish and States, 1993). The Uniprot Swiss-Prot Database (downloaded from SwissProt 2018-06-15) was used to obtain protein information and Uniprot Accession codes (Gish and States, 1993; UniProt Consortium, 2019). Genbank IDs provided by NCBI were used to match CpG background-mRNA overlaps with the annotated mRNA track. Gene ontology terms were paired to Uniprot Accession codes using the Uniprot Swiss-Prot Database (UniProt Consortium, 2019).

Once analysis inputs were created, gene ontology terms for each gene were matched with parental terms using default GO-MWU settings. Parental ontology categories with the exact same gene list were combined. Groups were further combined if they shared at least 75% of the same genes. After clustering was complete, a Mann-Whitney U test identified gene ontology categories that were significantly enriched by corresponding hyper- or hypo-methylated loci in genes using the default 10% FDR. Genes with DML were mapped to gene ontology subsets (GO Slim terms) for biological processes to further categorize gene functions.

Results

 Water Chemistry

285 Oysters experienced different water chemistry conditions between control and elevated pCO_2
286 treatments (Table 1). All oysters were subjected to acclimation pCO_2 conditions (pCO_2 = 521 ±
287 32 ppm, $\Omega_{calcite}$ = 2.82 ± 0.13) were maintained for 14 days. Oysters in control pCO_2 conditions
288 (pCO_2 = 492 ± 50 µatm; $\Omega_{calcite}$ = 3.01 ± 0.25) experienced low pCO_2 and higher $\Omega_{calcite}$ than
289 those in elevated pCO_2 conditions (pCO_2 = 2550 ± 211 µatm; $\Omega_{calcite}$ = 0.72 ± 0.06) (Table 1).
290

Experimental Stage	T (ºC)	S (PSU)	DIC	Alk_T	pH_{sw}	pCO_2 (µatm)	$\Omega_{calcite}$
Acclimation	14.6 ± 0.4	31.4 ± 0.1	1978 ± 7	2127 ± 6	7.94 ± 0.00	521 ± 32	2.82 ± 0.13
Control pCO_2 Conditions	14.5 ± 0.4	31.6 ± 0.3	1960 ± 32	2140 ± 15	7.95 ± 0.01	492 ± 50	3.01 ± 0.25
Elevated pCO_2 Conditions	14.5 ± 0.3	31.54 ± 0.32	2173 ± 37	2132 ± 42	7.29 ± 0.01	2550 ± 211	0.72 ± 0.06

Table 1. Summary of water chemistry during the 14-day acclimation period and 28-day
experimental exposure. Values indicate mean and standard error for temperature (T), salinity
(S), dissolved inorganic carbon (DIC), total alkalinity (A_T), calculated pH on seawater scale,
calculated pCO_2, and calculated calcite saturation ($\Omega_{calcite}$).

MBD-BS-Seq

DNA sequencing yielded 280 million DNA sequence reads (NCBI Sequence Read Archive:
BioProject accession number PRJNA513384). Of 276 million trimmed paired-end reads, 136
million (49.4%) were mapped to the *C. virginica* genome, providing an average of 13.6 million
reads per sample. Sequencing efforts provided data for 4,304,257 CpG loci (30.7% of
14,458,703 total CpGs in the *C. virginica* genome) with at least 5x coverage across all samples
combined. As expected, the location of CpGs with 5x coverage in the genome differed from the
distribution of all CpG motifs (Contingency test; χ_2 = 906,940, df = 4, *P*-value < 2.2e-16). Of all
loci with 5x coverage, 3,255,049 CpGs (75.6%) were found in genic regions in 33,126 out of
38,929 annotated genes in the genome.
305
The general methylation landscape was defined using all loci with a minimum 5x coverage in
each sample. The majority, 3,181,904 (73.9% of MBD-Enriched loci) loci were methylated, with
481,788 (11.2%) sparsely methylated loci and 640,565 (14.9%) unmethylated loci (Figure 1).
The *C. virginica* genome was determined to be 22% methylated. Loci methylation was
characterized in relation to exons, introns, transposable elements, and putative promoter
regions (Figure 2). Methylated CpGs were found primarily in genic regions, with 2,521,653 loci
(79.2%) in 25,496 genes. The null hypothesis — CpG methylation status was independent of

genomic location — was rejected, as the proportion of methylated CpG loci was different than
expected in exons, introns, transposable elements, putative promoters, and intergenic regions
(Contingency test; χ_2 = 872,070, df = 4, *P*-value < 2.2e-16; Figure 2). There was a larger
proportion of methylated loci found in exons compared to all CpGs in the genome (Figure 2).
Methylated loci were also found in introns (with 1,448,786 loci (47.3% of methylated loci) versus
1,013,691 CpGs (31.9%) in exons), although this was not higher than expected based on the
distribution of all CpGs. Transposable elements contained 755,222 methylated CpGs (23.7%).
Putative promoter regions overlapped with 106,111 loci (3.3%). There were 372,047 methylated
loci (11.7%) that did not overlap with either exons, introns, transposable elements, or promoter
regions.

Differential Methylation Analysis

A total of 598 CpG loci were differentially methylated between oysters exposed to control or
high pCO_2, with 51.8% hypermethylated and 48.2% hypomethylated between treatments
(Figure 3; full list available in Github repository). When considering a PCA using methylation
status of all CpG loci with 5x coverage across all samples, the first two principal components
explained 29.8% of sample variation (Figure 4A). However, the first two principal components in
a PCA with only differentially methylated loci (DML) explained 57.1% of the variation among
treatments (Figure 4B). These DML were distributed throughout the *C. virginica* genome (Figure
5). The highest number of DML were in the fifth chromosome, which also has the largest
number of genes but was not the largest chromosome (Figure 5A).

Examination of DML within genes revealed that some genes contained multiple DML (Figure
5B-F). Of the 481 genes with DML, the majority only contained one DML (Figure 5B-C). There
were 48 genes with 2 DML (Figure 5D), 16 genes with 3 DML (Figure 5E), 6 genes with 4 DML
(Figure 5F) and 1 gene with 5 DML. When multiple DML were found within a gene, they could
be methylated in either the same or opposite directions (Figure 5C-F).

Within the genome, DML were mostly present in genic regions, with 560 DML in 481 genes (368
DML in exons and 192 in introns). In addition, 42 DML were found in putative promoter regions,
57 in transposable elements, and 21 located outside of defined regions. The distribution of DML
in *C. virginica* gonad tissue was higher in exons than expected for MBD-enriched CpG loci with
minimum 5x coverage across all samples (Contingency test; χ_2 = 314.18, df = 4, *P*-value <
2.2e-16; Figure 6). Of the 598 DML, 310 were hypermethylated and 288 were hypomethylated
in the high pCO_2 treatment. The number of hyper- and hypomethylated DML was almost evenly
split within each genomic feature, with the exception of putative promoter regions that had 44
hypermethylated DML versus 23 hypomethylated DML. Within a gene, DML did not appear to
be concentrated in one particular region. The distribution of hyper- and hypomethylated DML
along a gene do not differ from each other (Figure 7). The DMG analysis did not identify any
differentially methylated genes.

The DML were found in genes responsible for various biological processes. However, no gene
ontology categories were significantly represented (Figure 8). The majority of genes with DML
were involved in protein ubiquitination processes. These genes were not consistently hyper- or

hypomethylated. Certain biomineralization genes did contain DML. The gene coding for
calmodulin-regulated spectrin-associated protein contained three hypomethylated and one
hypermethylated DML. Genes coding for EF-hand protein with calcium-binding domain,
calmodulin-binding transcription activator, and calmodulin-lysine N-methyltransferase contained
one or two hypermethylated DML.

Discussion

The present study is one of the first general descriptions of DNA methylation in *C. virginica*, and
is one of only a few that have examined epigenetic responses to ocean acidification in the
gonad tissue of a mollusk species. Five hundred ninety-eight differentially methylated loci (DML)
were identified in response to the elevated pCO_2 treatments, most of which were in exons. Not
only was DNA methylation of *C. virginica* altered in response to ocean acidification, but changes
in gonad methylation indicates potential for these methylation patterns to be inherited by
offspring.

Understanding how environmental stressors influence the epigenome is crucial when
considering potential acclimatization mechanisms in marine invertebrates. Our finding that high
pCO_2 impacts *C. virginica* DNA methylation adds to a growing body of work about ocean
acidification's impact on marine invertebrate methylomes. The coral species *P. damicornis*
demonstrated an overall increase in DNA methylation when exposed to low pH conditions (7.3 -
7.6), potentially influencing biomineralization (Putnam et al., 2016). Another coral species, *S.*
pistillata, also demonstrated an increase in genome-wide DNA methylation when exposed to
low pH conditions. Changes in the methylome also modified gene expression and altered
pathways involved in cell cycle regulation (Liew et al., 2018b). The present study on an oyster,
however, did not observe the overall genome-wide increase in methylation that was reported for
corals. Instead, we found subtle, but significant, increases or decreases in percent methylation
at several hundred individual CpGs distributed across the genome. As *C. virginica* and coral
species are adapted to different environments and ecological niches, It is possible that species-
specific differences in methylation responses contribute to the observed methylation pattern.

The *C. virginica* methylation landscape suggests a role for methylation in gene activity. Only
22% of the *C. virginica* genome is methylated, which is consistent with previous studies of
marine invertebrate genomes (Roberts and Gavery, 2012; Dimond and Roberts, 2016;
Hofmann, 2017). Methylated loci are concentrated in introns for *C. virginica*, followed by exons
and transposable elements. This location of methylated CpGs in gene bodies is consistent to
what has been reported across similar taxa (Roberts and Gavery, 2012; Eirin-Lopez and
Putnam, 2018). The concentration of methylated CpGs in gene bodies corresponds with
proposed functionality in influencing gene activity (Roberts and Gavery, 2012; Dixon et al.,
2014; Liew et al., 2018b). Our study also found methylation in transposable elements, putative
promoters and intragenic regions. In plants, transposable element methylation has been shown
to modulate the effect of transposable element insertion in genic regions (Hosaka and Kakutani,

2018). It is possible that methylation of transposable elements in *C. virginica* could also limit the
effect of transposable elements.
399
While both *C. virginica* and the closely-related Pacific oyster (*C. gigas*) have relatively sparse
transposable element and putative promoter methylation, there are differences between the
methylomes that could be indicative of species variation or genome assembly. Previously MBD-
BS-Seq of ctenidia tissue and whole genome bisulfite sequencing of sperm revealed that 15%
of the Pacific oyster genome is methylated, and that this methylation is consistent between
tissue types (Gavery and Roberts, 2013; Olson and Roberts, 2014). Our study of gonad tissue
found that 22% of the *C. virginica* genome is methylated. This difference can be attributed to the
distribution of CpG dinucleotides in both species. There are about 10 million CpG dinucleotides
in the described *C. gigas* genome, but over 14 million in the *C. virginica* genome. Additionally,
more CpG dinucleotides are located in genic regions in *C. virginica*. Gavery and Roberts (2013)
report 38% of total CpGs in *C. gigas* are found in exons and introns, with 47% of CpGs in
intergenic regions. The present study with *C. virginica* found 55% of total CpGs in exons and
introns, with 45% intergenic CpGs. However, there were more methylated loci in transposable
elements for *C. virginica*. Examination of methylation in conserved areas of both genomes may
elucidate if these differences are biologically meaningful.
415
The concentration of DML in gene bodies suggests a role for DNA methylation in gene
expression and regulation. A majority of genes with DML were involved in protein ubiquitination.
Protein ubiquitination is a post-translational protein modification that is involved in protein
synthesis and degradation (Peng et al., 2003; Komander, 2009). Previous studies in which
oysters were exposed to experimental ocean acidification conditions have demonstrated
changes in this pathway. For example, shotgun proteomic characterization of posterior gill
lamellae from adult *C. gigas* exposed to high pCO_2 revealed increased abundance of proteins
involved in ubiquitination and decreased protein degradation (Timmins-Schiffman et al., 2014).
Elevated pCO_2 levels were also found to upregulate malate dehydrogenase in adult *C. virginica*
mantle tissue (Tomanek et al., 2011). Several genes involved in protein ubiquitination, including
those for malate dehydrogenase, ubiquitin-protein ligase, RNA polymerase-associated protein,
and DNA damage-binding protein, were significantly hypermethylated in gonad tissue exposed
to elevated pCO_2. Hypermethylation of these genes may decrease transcriptional opportunities,
thus indicating a critical role in the response to ocean acidification. Given that the patterns
observed in our study were equivocal, additional research is needed to elucidate the role that
methylation plays in protein ubiquitination processes for oysters exposed to environmental
stressors.
433
Four genes involved in biomineralization contained DML, suggesting these genes can be
epigenetically regulated. Changes to calcium-binding gene expression have been previously
documented in *C. virginica* (Richards et al., 2018). Many studies examining ocean acidification-
induced carryover effects in bivalves note changes to calcification processes. For example, the
Sydney rock oyster (*Saccostrea glomerata*) larvae exhibit faster shell growth in high pCO_2
conditions when parents mature in those same conditions (Parker et al., 2012, 2015). In
contrast, larvae from other species found in the North Atlantic such as northern quahog (hard

clam; *M. mercenaria*) and bay scallops (*A. irradians*) developed slower when parents were
reproductively conditioned in low pH conditions (Griffith and Gobler, 2017). There is some
evidence to suggest that *C. virginica* larvae may be more resilient to high pCO_2 conditions than
M. mercenaria or *A. irradians (Gobler and Talmage, 2014)*. Differential methylation of
biomineralization genes in *C. virginica* reproductive tissue could be a mechanism to explain
when parental experience impacts larval calcification if in fact these DMLs are inherited.

Conclusion

Our study found that *C. virginica* demonstrates a significant epigenetic response to elevated
pCO_2 exposure, with 598 DML identified. The concentration of these DML in gene bodies
suggests that methylation may be important for transcriptional control in response to
environmental stressors. As ocean acidification induced differential methylation in *C. virginica*
gonad tissue, there is a potential for intergenerational epigenetic inheritance gene activity
control include processes such as biomineralization. As carryover effects can persist even
when stressors are long-removed (Venkataraman et al., 2019), understanding the mechanisms
involved in intergenerational acclimatization is crucial. Future work should focus on methylation
patterns in adult *C. virginica* reproductive tissue, fully-formed gametes, and larvae exposed to
various pCO_2 conditions to determine to what degree a difference in methylation influences
gene activity and how this might influence phenotypic plasticity.

Acknowledgements

461 This project was funded by National Science Foundation Biological Oceanography award
462 1635423 to KEL, JR, and SBR, and a Hall Conservation Genetics Research Award to YRV. This
463 work was facilitated through the use of advanced computational, storage, and networking
464 infrastructure provided by the Hyak supercomputer system at the University of Washington.

Data Accessibility

466 Associated information for all analyses and supplemental material can be found in the Github
467 repository: https://github.com/epigeneticstoocean/paper-gonad-meth

472 **Figures**
473
474 **Figure 1**. Frequency distribution of methylation ratios for CpG loci in *C. virginica* gonad tissue
475 DNA subjected to MBD enrichment. A total of 4,304,257 CpGs with at least 5x coverage
476 summed across all ten samples were characterized. Loci were considered methylated if they
477 were at least 50% methylated, sparsely methylated loci were 10-50% methylated, and
478 unmethylated loci were 0-10% methylated.
479
480 **Figure 2**. Proportion of CpG loci within with genomic features. All CpGs are every dinucleotide
481 in the *C. virginica* genome. Methylated CpGs refers to loci are defined as being at least 50%
482 methylated determined as part of this study.
483
Figure 3. Heatmap of DML in *C. virginica* reproductive tissue. Samples in control pCO_2
conditions are represented by grey, and samples in elevated pCO_2 conditions are represented
by a black bar. Loci with higher percent methylation are represented by darker colors. A logistic
regression identified 598 DML, defined as individual CpG dinucleotide with at least a 50%
methylation change between treatment and control groups, and a q-value < 0.01 based on
correction for false discovery rate with the SLIM method. The density of DML at each percent
methylation value is represented in the heatmap legend.
491
492 **Figure 4**. Principal Components Analysis of a) all CpG loci with 5x coverage across samples
493 and b) DML. Methylation status of individual CpG loci explained 29.2% of variation between
494 samples when considering all CpG loci. Methylation status of DML explained 57.1% of sample
495 variation.
496
Figure 5. Distribution of DML among chromosomes and genes. (a) Number of DML (bars) and
number of genes (line) in each *C. virginica* chromosome. (b) Counts of DML in genes containing
DML. Most genes that contained DML only had 1 DML. Breakdown of hyper- and
hypomethylation patterns in genes with one (c), two (d), three (e), and four (f) DML. The number
of hyper- or hypomethylated DML in a gene are indicated below the bars. (c) From left to right,
bars represent the number of genes with only hypermethylated DML and only hypomethylated
DML; (d) the number of genes with 2 hypermethylated DML, 1 hyper- and 1 hypomethylated
DML, and 2 hypomethylated DML; (e) the number of genes with 3 hypermethylated DML, 2
hyper- and 1 hypomethylated DML, 1 hyper- and 2 hypomethylated DML, and 3 hypomethylated
DML; (f) the number of genes with 4 hypermethylated DML, 3 hyper- and 1 hypomethylated
DML, 2 hyper- and 2 hypomethylated DML, 1 hyper- and 3 hypomethylated DML, and 4
hypomethylated DML. There was one gene with 5 DML (4 hypermethylated and 1
hypomethylated).
510
Figure 6. Proportion CpG loci within with exons, introns, transposable elements (TE), and
putative promoters for MBD-enriched CpGs and differentially methylated loci (DML),The
distribution of DML in *C. virginica* gonad tissue differed from distribution of MBD-enriched loci
with 5x coverage across control and treatment samples (Contingency test; χ_2 = 314.18, df = 4,
P-value < 2.2e-16).

Figure 7. Distribution of hyper- and hypomethylated DML along a hypothetical gene. The scaled

position of a DML within a gene was calculated by dividing the base pair position of the DML by

gene length. Counts of hypermethylated DML are plotted above the x-axis, and hypomethylated

DML counts are below the x-axis.

Figure 8. Biological processes represented by all genes used in enrichment background (%

Genes) and those with DML (% Genes with DML). Gene ontology categories with similar

functions are represented by the same color. Genes may be involved in multiple biological

processes. No gene ontologies were significantly enriched.

References

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., et al. (2012). methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol.* 13, R87.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Bao, W., Kojima, K. K., and Kohany, O. (2015). Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob. DNA* 6, 11.

Barton, A., Hales, B., Waldbusser, G. G., Langdon, C., and Feely, R. A. (2012). The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects. *Limnol. Oceanogr.* 57, 698–710.

Beniash, E., Ivanina, A., Lieb, N. S., Kurochkin, I., and Sokolova, I. M. (2010). Elevated level of carbon dioxide affects metabolism and shell formation in oysters *Crassostrea virginica*. *Mar. Ecol. Prog. Ser.* 419, 95–108.

Bird, A. (2002). DNA methylation patterns and epigenetic memory. *Genes Dev.* 16, 6–21.

Bossdorf, O., Richards, C. L., and Pigliucci, M. (2008). Epigenetics for ecologists. *Ecol. Lett.* 11, 106–115.

Boulais, M., Chenevert, K. J., Demey, A. T., Darrow, E. S., Robison, M. R., Roberts, J. P., et al. (2017). Oyster reproduction is compromised by acidification experienced seasonally in coastal regions. *Sci. Rep.* 7, 13276.

Byrne, M., Foo, S. A., Ross, P. M., and Putnam, H. M. (2019). Limitations of cross and multigenerational plasticity for marine invertebrates faced with global climate change. *Glob. Chang. Biol.* doi:10.1111/gcb.14882.

Deans, C., and Maggert, K. A. (2015). What Do You Mean, "Epigenetic"? *Genetics* 199, 887–896.

Dickson, A. G. (1990). Standard potential of the reaction : $AgCl_{(s)} + 12H2_{(g)} = Ag_{(s)} + Hcl_{(aq)}$, and the standard acidity constant of the ion HSO_{4^-} in synthetic sea water from 273.15 to 318.15 K. *The Journal of Chemical Thermodynamics* 22, 113–127.

Dimond, J. L., and Roberts, S. B. (2016). Germline DNA methylation in reef corals: patterns and potential roles in response to environmental change. *Mol. Ecol.* 25, 1895–1904.

Dineshram, R., Chandramouli, K., Ko, G. W. K., Zhang, H., Qian, P.-Y., Ravasi, T., et al. (2016). Quantitative analysis of oyster larval proteome provides new insights into the effects of multiple climate change stressors. *Glob. Chang. Biol.* 22, 2054–2068.

Dixon, G. B., Bay, L. K., and Matz, M. V. (2014). Bimodal signatures of germline methylation are linked with gene expression plasticity in the coral *Acropora millepora*. *BMC Genomics* 15, 1109.

Eirin-Lopez, J. M., and Putnam, H. M. (2018). Marine Environmental Epigenetics. *Ann. Rev. Mar. Sci.* doi:10.1146/annurev-marine-010318-095114.

Ekstrom, J. A., Suatoni, L., Cooley, S. R., Pendleton, L. H., Waldbusser, G. G., Cinner, J. E., et al. (2015). Vulnerability and adaptation of US shellfisheries to ocean acidification. *Nat. Clim. Chang.* 5, 207–214.

Feely, R. A., Alin, S. R., Newton, J., Sabine, C. L., Warner, M., Devol, A., et al. (2010). The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary. *Estuarine, Coastal and Shelf Science* 88, 442–449. doi:10.1016/j.ecss.2010.05.004.

Gatzmann, F., Falckenhayn, C., Gutekunst, J., Hanna, K., Raddatz, G., Carneiro, V. C., et al. (2018). The methylome of the marbled crayfish links gene body methylation to stable expression of poorly accessible genes. *Epigenetics Chromatin* 11, 57.

Gavery, M. R., and Roberts, S. B. (2013). Predominant intragenic methylation is associated with gene expression characteristics in a bivalve mollusc. *PeerJ* 1, e215.

Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J.-P., Middelburg, J. J., and Heip, C. H. R. (2007). Impact of elevated CO_2 on shellfish calcification. *Geophys. Res. Lett.* 34, L07603.

Gish, W., and States, D. J. (1993). Identification of protein coding regions by database similarity search. *Nat. Genet.* 3, 266–272.

Gobler, C. J., and Talmage, S. C. (2014). Physiological response and resilience of early life-stage Eastern oysters (*Crassostrea virginica*) to past, present and future ocean acidification. *Conserv Physiol* 2, cou004.

Gómez-Chiarri, M., Warren, W. C., Guo, X., and Proestou, D. (2015). Developing tools for the study of molluscan immunity: The sequencing of the genome of the eastern oyster, *Crassostrea virginica*. *Fish & Shellfish Immunology* 46, 2–4. doi:10.1016/j.fsi.2015.05.004.

Griffith, A. W., and Gobler, C. J. (2017). Transgenerational exposure of North Atlantic bivalves to ocean acidification renders offspring more vulnerable to low pH and additional stressors. *Sci. Rep.* 7, 11394.

Helm, M. M., and Bourne, N. (2004). *Hatchery Culture of Bivalves: A Practical Manual*. Food & Agriculture Org.

Hofmann, G. E. (2017). Ecological Epigenetics in Marine Metazoans. *Frontiers in Marine Science* 4. doi:10.3389/fmars.2017.00004.

Hosaka, A., and Kakutani, T. (2018). Transposable elements, genome evolution and transgenerational epigenetic variation. *Curr. Opin. Genet. Dev.* 49, 43–48.

IPCC, 2019: Summary for Policymakers. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate* [H.- O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama, N. Weyer (eds.)]. In press.

Johnson, K. M., and Kelly, M. W. (2019). Population epigenetic divergence exceeds genetic divergence in the Eastern oyster *Crassostrea virginica* in the Northern Gulf of Mexico. *Evol. Appl.* doi:10.1111/eva.12912.

Komander, D. (2009). The emerging complexity of protein ubiquitination. *Biochem. Soc. Trans.*

606  37, 937–953.

Krueger, F., and Andrews, S. R. (2011). Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* 27, 1571–1572.

Kurihara, H., Kato, S., and Ishimatsu, A. (2007). Effects of increased seawater pCO_2 on early development of the oyster *Crassostrea gigas*. *Aquat. Biol.* 1, 91–98.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9, 357–359.

Lee, Kitach Tae-Wook, Kim Byrne, Robert H Millero, Frank J Feely, Richard A Liu, Yong-Ming (2010). The universal ratio of boron to chlorinity for the North Pacific and North Atlantic oceans. *Geochimica et Cosmochimica Acta* 74, 1801–1811.

Lewis, E., and Wallace, D. W. (1998). R: Program developed for CO_2 system calculations ORNL/CDIAC-105. *Carbon Dioxide Information Analysis CentreOak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee.*

Liew, Y. J., Howells, E. J., Wang, X., Michell, C. T., Burt, J. A., Idaghdour, Y., et al. (2018a). Intergenerational epigenetic inheritance in reef-building corals. *bioRxiv*, 269076. doi:10.1101/269076.

Liew, Y. J., Zoccola, D., Li, Y., Tambutté, E., Venn, A. A., and Michell, C. T. (2018b). Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-building coral. *Science Advances*.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17, 10–12.

Olson, C. E., and Roberts, S. B. (2014). Genome-wide profiling of DNA methylation and gene expression in *Crassostrea gigas* male gametes. *Front. Physiol.* 5, 224.

Omoregie, E., Mwatilifange, N. S. I., and Liswaniso, G. (2019). Futuristic Ocean Acidification Levels Reduce Growth and Reproductive Viability in the Pacific Oyster (*Crassostrea gigas*). *J. Appl. Sci. Environ. Manage.* 23, 1747–1754.

Parker, L. M., O'Connor, W. A., Byrne, M., Coleman, R. A., Virtue, P., Dove, M., et al. (2017). Adult exposure to ocean acidification is maladaptive for larvae of the Sydney rock oyster *Saccostrea glomerata* in the presence of multiple stressors. *Biol. Lett.* 13, 20160798.

Parker, L. M., O'Connor, W. A., Raftos, D. A., Pörtner, H.-O., and Ross, P. M. (2015). Persistence of Positive Carryover Effects in the Oyster, *Saccostrea glomerata*, following Transgenerational Exposure to Ocean Acidification. *PLoS One* 10, e0132276.

Parker, L. M., Ross, P. M., O'Connor, W. A., Borysko, L., Raftos, D. A., and Pörtner, H.-O. (2012). Adult exposure influences offspring response to ocean acidification in oysters. *Glob. Chang. Biol.* 18, 82–92.

Parker, L. M., Ross, P. M., O'Connor, W. A., Pörtner, H. O., Scanes, E., and Wright, J. M.

(2013). Predicting the response of molluscs to the impact of ocean acidification. *Biology* 2,
651–692.

Peng, J., Schwartz, D., Elias, J. E., Thoreen, C. C., Cheng, D., Marsischky, G., et al. (2003). A
proteomics approach to understanding protein ubiquitination. *Nat. Biotechnol.* 21, 921–926.

Putnam, H. M., Davidson, J. M., and Gates, R. D. (2016). Ocean acidification influences host
DNA methylation and phenotypic plasticity in environmentally susceptible corals. *Evol.*
Appl. 9, 1165–1178.

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* 26, 841–842.

Richards, M., Xu, W., Mallozzi, A., Errera, R. M., and Supan, J. (2018). Production of Calcium-
Binding Proteins in *Crassostrea virginica* in Response to Increased Environmental CO2
Concentration. *Frontiers in Marine Science* 5, 203.

Ries, J. B. (2011). A physicochemical framework for interpreting the biological calcification
response to CO_2-induced ocean acidification. *Geochim. Cosmochim. Acta* 75, 4053–4064.

Roberts, S. B., and Gavery, M. R. (2012). Is There a Relationship between DNA Methylation
and Phenotypic Plasticity in Invertebrates? *Front. Physiol.* 2.
doi:10.3389/fphys.2011.00116.

Rondon, R., Grunau, C., Fallet, M., Charlemagne, N., Sussarellu, R., Chaparro, C., et al. (2017).
Effects of a parental exposure to diuron on Pacific oyster spat methylome. *Environ*
Epigenet 3. doi:10.1093/eep/dvx004.

Ross, P. M., Parker, L., and Byrne, M. (2016). Transgenerational responses of molluscs and
echinoderms to changing ocean conditions. *ICES Journal of Marine Science*.
doi:10.1093/icesjms/fsv254.

Roy, Rabindra N Roy, Lakshimi N Vogel, Kathleen M Porter-Moore, C Pearson, Tara Good,
Catherine E Millero, Frank J Campbell, Douglas M (1993). The dissociation constants of
carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine*
Chemistry 44, 249–267.

Smit, A. F. A., Hubley, R., and Green, P. (2013). 2013--2015. RepeatMasker Open-4.0.

Strader, M. E., Wong, J. M., Kozal, L. C., Leach, T. S., and Hofmann, G. E. (2019). Parental
environments alter DNA methylation in offspring of the purple sea urchin,
Strongylocentrotus purpuratus. J. Exp. Mar. Bio. Ecol. 517, 54–64.

Suzuki, M. M., and Bird, A. (2008). DNA methylation landscapes: provocative insights from
epigenomics. *Nat. Rev. Genet.* 9, 465–476.

Timmins-Schiffman, E., Coffey, W. D., Hua, W., Nunn, B. L., Dickinson, G. H., and Roberts, S.
B. (2014). Shotgun proteomics reveals physiological response to ocean acidification in
Crassostrea gigas. BMC Genomics 15, 951.

Tomanek, L., Zuzow, M. J., Ivanina, A. V., Beniash, E., and Sokolova, I. M. (2011). Proteomic
response to elevated PCO2 level in eastern oysters, *Crassostrea virginica*: evidence for
oxidative stress. *J. Exp. Biol.* 214, 1836–1844.

UniProt Consortium (2019). UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.*
47, D506–D515.

Van Heuven, S., Pierrot, D., Rae, J. W. B., Lewis, E., and Wallace, D. W. R. (2011). MATLAB
program developed for CO_2 system calculations. *ORNL/CDIAC-105b. Carbon Dioxide*
Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy,
Oak Ridge, Tennessee 530.

Venkataraman, Y. R., Spencer, L. H., and Roberts, S. B. (2019). Larval Response to Parental
Low pH Exposure in the Pacific Oyster *Crassostrea gigas*. *Journal of Shellfish Research*
38, 743. doi:10.2983/035.038.0325.

Waldbusser, G. G., Hales, B., Langdon, C. J., Haley, B. A., Schrader, P., Brunner, E. L., et al.
(2014). Saturation-state sensitivity of marine bivalve larvae to ocean acidification. *Nat. Clim.*
Chang. 5, 273.

Wang, H.-Q., Tuominen, L. K., and Tsai, C.-J. (2011). SLIM: a sliding linear model for estimating
the proportion of true null hypotheses in datasets with dependence structures.
Bioinformatics 27, 225–231.

Wright, R. M., Aglyamova, G. V., Meyer, E., and Matz, M. V. (2015). Gene expression
associated with white syndromes in a reef building coral, *Acropora hyacinthus*. *BMC*
Genomics 16, 371.

Zhao, L., Liu, L., Liu, B., Liang, J., Lu, Y., and Yang, F. (2019). Antioxidant responses to
seawater acidification in an invasive fouling mussel are alleviated by transgenerational
acclimation. *Aquat. Toxicol.* 217, 105331.

Zhao, L., Yang, F., Milano, S., Han, T., Walliser, E. O., and Schöne, B. R. (2018).
Transgenerational acclimation to seawater acidification in the Manila clam *Ruditapes*
philippinarum: Preferential uptake of metabolic carbon. *Sci. Total Environ.* 627, 95–103.

707
708