General DNA methylation patterns and environmentally-induced differential methylation in the eastern oyster (*Crassostrea virginica*)

Yaamini R. Venkataraman[1], Alan M. Downey-Wall[2], Justin Ries[2], Isaac Westfield[2], Samuel J. White[1], Steven B. Roberts[1], Kathleen E. Lotterhos[2]

[1]School of Aquatic & Fishery Sciences, University of Washington, 1122 NE Boat St, Seattle, WA, USA

[2]Department of Marine and Environmental Sciences, Northeastern University, 430 Nahant Road, Nahant, MA 01908

Abstract

Epigenetic modification, specifically DNA methylation, is one possible mechanism for intergenerational plasticity. Before inheritance of methylation patterns can be characterized, we need a better understanding of how environmental change modifies the parental epigenome. To examine the influence of experimental ocean acidification on eastern oyster (*Crassostrea virginica*) gonad tissue, oysters were cultured in the laboratory under control (491 ± 49 µatm) or high (2550 ± 211 µatm) pCO_2 conditions for four weeks. DNA from reproductive tissue was isolated from five oysters per treatment, then subjected to bisulfite treatment and DNA sequencing. Irrespective of treatment, DNA methylation was primarily found in gene bodies with approximately 22% of CpGs (2.7% of total cytosines) in the *C. virginica* genome predicted to be methylated. In response to elevated pCO_2, we found 598 differentially methylated loci primarily overlapping with gene bodies. A majority of differentially methylated loci were in exons (61.5%) with less intron overlap (31.9%). While there was no evidence of a significant tendency for the genes with differentially methylated loci to be associated with distinct biological processes, the concentration of these loci in gene bodies, including genes involved in protein ubiquitination and biomineralization suggests DNA methylation may be important for transcriptional control in response to ocean acidification. Changes in gonad methylation also indicate potential for these methylation patterns to be inherited by offspring. Understanding how experimental ocean acidification conditions modify the oyster epigenome, and if these modifications are inherited, allows for a better understanding of how ecosystems will respond to environmental change.

Keywords: eastern oyster, DNA methylation, epigenetics, ocean acidification

Introduction

As increased anthropogenic carbon dioxide is expected to create adverse conditions for calcifying organisms (IPCC 2019), efforts have been made to understand how ocean acidification impacts ecologically and economically important organisms like bivalves (Parker et al., 2013; Ekstrom et al., 2015). Bivalve species are sensitive to reduced aragonite saturation associated with ocean acidification, with larvae being particularly vulnerable (Barton et al., 2012; Waldbusser et al., 2014). Shell structure may be compromised in larvae, juveniles, and adults (Gazeau et al., 2007; Kurihara et al., 2007; Beniash et al., 2010; Ries, 2011). Aside from affecting calcification and shell growth, ocean acidification can impact protein synthesis, energy production, metabolism, antioxidant responses, and reproduction (Tomanek et al., 2011; Timmins-Schiffman et al., 2014; Dineshram et al., 2016; Boulais et al., 2017; Omoregie et al., 2019).

Additionally, adult exposure to ocean acidification may impact their larvae (reviewed in (Ross et al., 2016; Byrne et al., 2019). For example, adult Manila clams (*Ruditapes philippinarum*) and mussels (*Musculista senhousia*) reproductively conditioned in high pCO_2 waters yield offspring that exhibit significantly faster development or lower oxidative stress protein activity in those same conditions (Zhao et al., 2018, 2019). In contrast, northern quahog (hard clam; *Mercenaria mercenaria*) and bay scallop (*Argopecten irradians*) larvae may be more vulnerable to ocean acidification and additional stressors when parents are reproductively conditioned in high pCO_2 waters (Griffith and Gobler, 2017). Some species exhibit both positive and negative carryover effects (ex. *Saccostrea glomerata*; (Parker et al., 2012, 2017). Intergenerational effects have also been documented when adult exposure to ocean acidification does not coincide with reproductive maturity (*e.g. Crassostrea gigas*; (Venkataraman et al., 2019)). Although intergenerational carryover effects are now at the forefront of ocean acidification research in bivalve species, the mechanisms responsible for these effects are still unclear.

Epigenetics is the next frontier for understanding how environmental memory may modulate phenotypic plasticity across generations (Eirin-Lopez and Putnam, 2018). Epigenetics refers to changes in gene expression that do not arise from changes in the DNA sequence, with methylation of cytosine bases being the most studied mechanism (Bird, 2002; Deans and Maggert, 2015). Unlike highly methylated vertebrate genomes, marine invertebrate taxa have sparse methylation throughout their genomes, similar to a mosaic pattern (Suzuki and Bird, 2008). Genes that benefit from stable transcription, such as housekeeping genes, tend to be more methylated, while environmental response genes that are less methylated are prone to more spurious transcription and alternative splicing patterns, thereby possibly increasing phenotypic plasticity (Roberts and Gavery, 2012; Dimond and Roberts, 2016; Gatzmann et al., 2018). Increased levels of DNA methylation can also correlate with increased transcription. Several base pair resolution studies in *C. gigas* demonstrate a positive association between DNA methylation and gene expression that is consistent across cell types (Roberts and Gavery, 2012; Gavery and Roberts, 2013; Olson and Roberts, 2014). Since DNA methylation could provide a direct link between environmental conditions and phenotypic plasticity via influencing

79 gene activity, elucidating how invertebrate methylomes respond to abiotic factors is crucial for
80 understanding potential acclimatization mechanisms (Bossdorf et al., 2008; Hofmann, 2017).
81
While bivalve species have been used as model organisms to characterize marine invertebrate
methylomes, how ocean acidification affects bivalve DNA methylation is poorly understood.
Methylation responses to ocean acidification have been studied in multiple coral species. When
placed in low pH conditions (7.6-7.35), *Montipora capitata* did not demonstrate any differences
in calcification, metabolic profiles, or DNA methylation in comparison to clonal fragments in
ambient pH (7.9-7.65) (Putnam et al., 2016). DNA methylation increased in another coral
species, *Pocillopora damicornis*, in addition to reduced in calcification and more differences in
metabolic profiles (Putnam et al., 2016). The coral *Stylophora pistillata* also demonstrates
increased global methylation as pH decreases ($pH_{treatment}$ = 7.2, 7.4, 7.8; $pH_{control}$ = 8.0), with
methylation reducing spurious transcription (Liew et al., 2018b). Combined whole genome
bisulfite sequencing and RNA sequencing revealed differential methylation and expression of
growth and stress response pathways controlled differences in cell and polyp size between
treatments (Liew et al., 2018b). The association between DNA methylation and phenotypic
differences in these corals demonstrates that epigenetic regulation of genes is potentially
important for acclimatization and adaptation to environmental perturbation. Recent examination
of *C. virginica* methylation patterns in response to a natural salinity gradient suggests that
differential methylation may modulate environmental response in this species (Johnson and
Kelly, 2019).
100
There is evidence that suggests that methylation patterns can be inherited in marine
invertebrates. For example, purple sea urchin (*Strongylocentrotus purpuratus*) offspring have
methylomes that reflect maternal rearing conditions (Strader et al., 2019). Different parental
temperature and salinity regimes influence larval methylomes in *Platygyra daedalea* (Liew et al.,
2018a). In the Pacific oyster (*C. gigas*), parental exposure to pesticides influence DNA
methylation in spat, even though the spat were not exposed to these conditions (Rondon et al.,
2017). Methylation changes in gametes are likely the ones that could be inherited, and may play
a role in carryover effects. Before determining if DNA methylation is a viable mechanism for
altering the phenotypes of offspring or subsequent generations, the epigenome of bivalve
reproductive tissue in response to ocean acidification must be characterized.
111
The present study is the first to determine if ocean acidification induces differential methylation
in reproductive tissue in the eastern oyster (*Crassostrea virginica*). Adult *C. virginica* were
exposed to control or elevated pCO$_2$ conditions. We hypothesize that ocean acidification will
induce differential methylation in *C. virginica* gonad tissue, and that genes with differentially
methylated loci will have biological functions that could allow for acclimatization to
environmental perturbation. Understanding how experimental ocean acidification conditions
modify the oyster epigenome, and if these modifications are inherited, allows for a better
understanding of how ecosystems will respond to environmental change.

Methods

Experimental Design

Adult *C. virginica* (9.55 cm ± 0.45) were collected from an intertidal oyster reef in Plum Island
Sound, MA (42.681764, -70.813498) in mid-July 2016. The oysters were transported to the
Marine Science Center at Northeastern University (Nahant, MA), where they were cleaned and
randomly assigned to one of six flow-through tanks (50L) maintained at ambient seawater
conditions. Oysters were acclimated for 14 days under control conditions (500 µatm; 14-15ºC)
before initiating a 28-day experimental exposure. Half of the tanks remained at control pCO_2
conditions (500 µatm, $\Omega_{calcite} > 1$), while the other half were ramped up to elevated pCO_2
conditions (2500 µatm, $\Omega_{calcite} < 1$) over 24 hours. This elevated treatment is consistent with
observations in other estuarine ecosystems that oysters inhabit (Feely et al., 2010), although pH
in nature only stays as extreme for short periods of time (e.g. hours). Moreover, the extreme
treatment was also chosen to increase precision and therefore power to detect a response
(Whitlock and Schluter, 2014).
134
Treatment conditions were replicated across three tanks, with oysters distributed evenly among
tanks (1-2 oysters per tank). Each tank had an independent flow-regulator that delivered fresh,
natural seawater at approximately 150 ml min_{-1}. Carbonate chemistry was maintained
independently for each tank by mixtures of compressed CO_2 and compressed air at flow rates
proportional to the target pCO_2 conditions. Gas flow rates were maintained with *Aalborg* digital
solenoid-valve-controlled mass flow controllers (Model GFC17, precision = 0.1mL/min). Within
a treatment, tanks were replenished with fresh seawater and each tank was independently
bubbled with its own mixed gas stream, with partial recirculation and filtration with other tanks in
the treatment. As a result, the carbonate chemistry (i.e., the independent variable by which the
treatments were differentiated) of the replicate tanks were slightly different from each other,
which is evidence of their technical independence. Temperature was maintained at 15ºC using
Aqua Euro USA model MC-1/4HP chillers coupled with 50-watt electric heaters. Average salinity
was determined by the incoming natural seawater and reflected ambient ocean salinity of
Massachusetts Bay near the Marine Science Center (Latitude = 42.416100, Longitude = -
70.907737).
150
151 Oysters were fed 2.81 mL/day of a 10% Shellfish Diet 1800 twice daily following Food and
152 Agriculture Organization's best practices for oysters (Helm and Bourne, 2004). Five oysters
153 were collected from each treatment at the end of the 28 day exposure. They were immediately
154 dissected with gonadal tissue harvested and immediately flash frozen. Partial gamete
155 maturation was evident upon visual inspection.

Measurement and control of seawater carbonate chemistry

157 The carbonate chemistry of tanks was controlled by bubbling mixtures of compressed CO_2 and
158 compressed air at flow rates proportional to the target pCO_2 conditions. The control pCO_2
159 treatments were maintained by bubbling compressed ambient air only.
160
Temperature, pH, and salinity of all replicate tanks was measured three times per week for the
duration of the experiment. Temperature was measured using a glass thermometer to 0.1ºC

accuracy, pH was measured using an *Accumet* solid state pH electrode (precision = 1mV),
salinity was measured using a *YSI 3200* conductivity probe (precision = 0.1 ppt). Every two
weeks, seawater samples were collected from each replicate tank for analysis of dissolved
inorganic carbon (DIC) and total alkalinity (A_T). Samples were collected in 250 mL borosilicate
glass bottles sealed with a greased stopper, immediately poisoned with 100 µL saturated $HgCl_2$
solution, and then refrigerated. Samples were analyzed for DIC via coulometry and Alk_T via
closed-cell potentiometric Gran Titration with a VINDTA 3C (Marianda Corporation). Other
carbonate system parameters, including $\Omega_{calcite}$, pH, and *p*CO_2, were calculated from DIC, A_T,
salinity, and temperature using CO2SYS software version 2.1 (Lewis and Wallace, 1998; Van
Heuven et al., 2011), using the seawater pH scale (mol/kg-SW) with K1 and K2 values from
(Roy et al., 1993), a $KHSO_4$ value from (Dickson, 1990), and a $[B]_T$ value from (Lee et al., 2010).

MBD-BS Library Preparation
DNA was isolated from five gonad tissue samples per treatment using the E.Z.N.A. Mollusc Kit
(Omega) according to the manufacturer's instructions. Isolated DNA was quantified using a
Qubit dsDNA BR Kit (Invitrogen). DNA samples, ranging from 12.8 ng/µL to 157 ng/µL, were
placed in 1.5 mL centrifuge tubes and sonicated using a QSONICA CD0004054245 (Newtown,
CT) in 30 second interval periods over ten minutes at 4 ºC and 25% intensity. Shearing size
(350bp) was verified using a 2200 TapeStation System (Agilent Technologies). Samples were
enriched for methylated DNA with the MethylMiner kit (Invitrogen). A single-fraction elution using
400 µL of high salt buffer was used to obtain captured DNA. After ethanol precipitation, 25 µL of
buffer was used for the final elution. Library preparation and sequencing was performed by
ZymoResearch using Pico Methyl-Seq Library Prep Kit (Cat. #D5455). Libraries were then
barcoded and pooled into two lanes (eight samples in one and two in another) to generate
100bp paired-end reads on the HiSeq1500 sequencer (Illumina, Inc.).

Global Methylation Characterization
Sequences were trimmed with 10 bp removed from both the 5' and 3' ends using TrimGalore!
v.0.4.5 (Martin, 2011). Quality of sequences was assessed with FastQC v.0.11.7 (Andrews,
2010). The *C. virginica* genome (NCBI Accession GCA_002022765.4) was prepared using
Bowtie 2-2.3.4 (Linux x84_64 version; (Langmead and Salzberg, 2012)) within the
`bismark_genome_preparation` function in Bismark v.0.19.0 (Krueger and Andrews, 2011).
Trimmed sample sequences were then aligned to the genome using Bismark v.0.19.0 (Krueger
and Andrews, 2011) with non-directionality specified alignment score set using `-score_min`
`L,0,-1.2`. Alignment files (ie. bam) were deduplicated (`deduplicate_bismark`), sorted and
indexed using SAMtools v.1.9 (Li et al., 2009). Methylation calls were extracted from
deduplicated files using `bismark_methylation_extractor`.
198
Various *C. virginica* genome feature tracks were created for downstream analyses using
BEDtools v2.26.0 (Quinlan and Hall, 2010). Genes, mRNA, coding sequences, and exons were
derived directly from the *C. virginica* genome on NCBI (Gómez-Chiarri et al., 2015). The
complement of the exon track was used to identify introns, and coding sequences were
subtracted from exons to identify untranslated regions of exons (UTR). Exon locations were

removed from the complement of the gene track to define intergenic regions. Putative promoter
regions were defined as those 1kb upstream of transcription start sites. Putative transposable
elements were identified using RepeatMasker (v4.07) with RepBase-20170127 and RMBlast
2.6.0 (Smit et al., 2013; Bao et al., 2015). All species available in RepBase-20170127 were
used to identify transposable elements.
209
Overall *C. virginica* gonad methylation patterns were characterized using information from all
samples. Individual CpG dinucleotides with at least 5x coverage in each sample were classified
as methylated (≥ 50% methylation), sparsely methylated (10-50% methylation), or unmethylated
(< 10% methylation). The locations of all methylated CpGs were characterized in relation to
putative promoter regions, UTR, exons, introns, transposable elements, and intergenic regions.
We tested the null hypothesis that there was no association between the genomic location of
CpG loci and methylation status (all CpGs versus methylated CpGs) with a chi-squared
contingency test (`chisq.test` in R Version 3.5.0).
218
Methylation islands were determined to characterize overall methylation in the *C. virginica*
genome using a sliding window analysis based on (Jeong et al., 2018). Islands were defined as
areas of the genome with enriched levels of methylated CpGs (>50% methylation). To define
methylation islands, each chromosome was examined using an initial 500 bp window starting at
the first methylated CpG. If the proportion of methylated CpGs in the window was greater than
0.2, the window was extended by 50 bp; if not, the analysis proceeded to the next methylated
CpG. Windows were continually extended until the proportion of methylated CpGs in the window
fell below the 0.2 criteria. The location of methylation islands in the genome were characterized
using BEDtools `intersect` v2.26.0.

Differential Methylation Analysis
Differential methylation analysis for individual CpG dinucleotides was performed using
`methylKit` v.1.7.9 in R (Akalin et al., 2012) using deduplicated, sorted bam files as input. Only
CpGs with at least 5x coverage in each sample were considered for analysis. Methylation
differences between treatments were obtained for all loci in the CpG background using
`calculateDiffMeth`, a logistic regression built into `methylKit`. The logistic regression
models the log odds ratio based on the proportion of methylation at each locus:
235

$$log\left(\frac{\pi_i}{1 - \pi_i}\right) = \beta_0 + \beta_1 Treatment_i$$

237
A differentially methylated locus (DML) was defined as an individual CpG dinucleotide with at
least a 50% methylation change between treatment and control groups, and a q-value < 0.01
based on correction for false discovery rate with the SLIM method (Wang et al., 2011).
Hypermethylated DML were defined as those with significantly higher percent methylation in
oysters exposed to high pCO_2 conditions, and hypomethylated DML with significantly lower
percent methylation in the high pCO_2 treatment. A Principal Components Analysis (PCA) was
performed for differentially methylated loci (DML) for oyster sample methylation profiles between

treatments, then compared to a PCA for all MBD-enriched CpG loci. The location of DML were
characterized in relation to putative promoter regions, UTR, exons, introns, transposable
elements, and intergenic regions using BEDtools `intersect` v2.26.0. Loci that did not overlap
with the aforementioned genomic features were also identified. A chi-squared contingency test
was used to test the null hypothesis of no association between genomic location and
methylation status between MBD-enriched CpGs and DML. To describe the location of DML
across different gene architectures, the position of DML in the gene was scaled from 0 to 100
bp.

Enrichment Analysis
Functional enrichment analyses were used to determine if any biological processes were
overrepresented in genes based on individual CpG methylation levels. Enrichment was
conducted with GO-MWU, a rank-based gene enrichment method initially developed for
analyzing transcriptomics data (Wright et al., 2015). Instead of only using genes with DML, GO-
MWU identifies GO categories that are overrepresented by genes with any CpGs, allowing for
more data to contribute to any trends. GO-MWU scripts and a gene ontology database were
downloaded from the GO-MWU Github repository (https://github.com/z0on/GO_MWU).
261
A gene list and table of significance measures were used as GO-MWU analysis inputs. The
gene list contained Genbank IDs and all associated gene ontology terms. For the table of
significance measures, Genbank IDs were matched with the smallest *P*-value for associated
CpGs analyzed by `methylKit`. To match the Genbank IDs to CpG loci within mRNAs and
create the gene list, overlaps between the *C. virginica* mRNA track from NCBI and the CpG
background used in `methylKit` were obtained using BEDtools `intersect` v2.26.0. The
mRNAs were then annotated with Uniprot Accession codes using a BLASTx search (v.2.2.29;
(Gish and States, 1993; UniProt Consortium, 2019). The Uniprot Swiss-Prot Database
(downloaded from SwissProt 2018-06-15) was used to obtain protein information and Uniprot
Accession codes. Genbank IDs provided by NCBI were used to match CpG background-mRNA
overlaps with the annotated mRNA track. Gene ontology terms were paired to Uniprot
Accession codes using the Uniprot Swiss-Prot Database (UniProt Consortium, 2019). All GO-
MWU inputs are available in the associated Github repository (Venkataraman 2020).
275
Once analysis inputs were created, gene ontology terms for each gene were matched with
parental terms using default GO-MWU settings. Parental ontology categories with the exact
same gene list were combined. Groups were further combined if they shared at least 75% of the
same genes. After clustering was complete, a Mann-Whitney U test identified gene ontology
categories that were significantly enriched by corresponding hyper- or hypo-methylated loci in
genes using the default 10% FDR. Genes with DML were mapped to gene ontology subsets
(GO Slim terms) for biological processes to further categorize gene functions.
283

Results

Water Chemistry

All oysters were initially subjected to acclimation pCO_2 conditions (pCO_2 = 521 ± 32 ppm, $\Omega_{calcite}$
= 2.82 ± 0.13) for 14 days. Following acclimation the treatments were initiated. Oysters in
control pCO_2 conditions (pCO_2 = 492 ± 50 µatm; $\Omega_{calcite}$ = 3.01 ± 0.25) experienced low pCO_2
and higher $\Omega_{calcite}$ than those in elevated pCO_2 conditions (pCO_2 = 2550 ± 211 µatm; $\Omega_{calcite}$ =
0.72 ± 0.06) (Table 1).
291

Experimental Stage	T (ºC)	S (PSU)	DIC	A_T	pH_{sw}	pCO₂ (µatm)	$\Omega_{calcite}$
Acclimation	14.6 ± 0.4	31.4 ± 0.1	1978 ± 7	2127 ± 6	7.94 ± 0.00	521 ± 32	2.82 ± 0.13
Control pCO_2 Conditions	14.5 ± 0.4	31.6 ± 0.3	1960 ± 32	2140 ± 15	7.95 ± 0.01	492 ± 50	3.01 ± 0.25
Elevated pCO_2 Conditions	14.5 ± 0.3	31.54 ± 0.32	2173 ± 37	2132 ± 42	7.29 ± 0.01	2550 ± 211	0.72 ± 0.06

Table 1. Summary of water chemistry during the 14-day acclimation period and 28-day
experimental exposure. Values indicate mean and standard error for temperature (T), salinity
(S), dissolved inorganic carbon (DIC), total alkalinity (A_T), calculated pH on seawater scale,
calculated pCO_2, and calculated calcite saturation ($\Omega_{calcite}$).

MBD-BS-Seq

DNA sequencing yielded 280 million DNA sequence reads (NCBI Sequence Read Archive:
BioProject accession number PRJNA513384). Of 276 million trimmed paired-end reads, 136
million (49.4%) were mapped to the *C. virginica* genome, providing an average of 13.6 million
reads per sample. Sequencing efforts provided data for 4,304,257 CpG loci (30.7% of
14,458,703 total CpGs in the *C. virginica* genome) with at least 5x coverage across all samples
combined. As expected, the location of CpGs with 5x coverage in the genome differed from the
distribution of all CpG motifs (Contingency test; χ_2 = 1,306,900, df = 6, *P*-value < 2.2e-16). Of
all loci with 5x coverage, 3,255,049 CpGs (75.6%) were found in genic regions in 33,126 out of
38,929 annotated genes in the genome.
306
The general methylation landscape was defined using all loci with a minimum 5x coverage in
each sample. The majority, 3,181,904 (73.9% of MBD-Enriched loci) loci were methylated, with
481,788 (11.2%) sparsely methylated loci and 640,565 (14.9%) unmethylated loci (Figure 1A).
Median values for global percent methylation and sample methylation varied across genome
features (Figure 1B-G). Based on these parameters and data, we calculated that 22% of all
CpGs in the gonads (2.7% of total cytosines) had methylation levels greater than 50%. Loci
methylation was characterized in relation to putative promoters, UTR, exons, introns,

transposable elements, and intergenic regions (Figure 2). Methylated CpGs were found
primarily in genic regions, with 2,521,653 loci (79.2%) in 25,496 genes. We rejected the null
hypothesis that CpG methylation status was independent of genomic location, as the proportion
of methylated CpG loci was different than expected in putative promoters, UTR, exons, introns,
transposable elements, and intergenic regions (Contingency test; χ_2 = 1,311,600, df = 6, *P*-
value < 2.2e-16; Figure 2). There was a larger proportion of methylated loci found in exons
compared to all CpGs in the genome (Figure 2). Methylated loci were also found in introns (with
1,448,786 loci (47.3% of methylated loci) versus 1,013,691 CpGs (31.9%) in exons), although
this was not higher than expected based on the distribution of all CpGs. Transposable elements
contained 755,222 methylated CpGs (23.7%). Putative promoter regions overlapped with
106,111 loci (3.3%), UTR with 128,585 loci (4.0%), and intergenic regions with 660,197 loci
(20.7%). There were 372,047 methylated loci (11.7%) that did not overlap with either exons,
introns, transposable elements, or promoter regions.
327
A total of 37,063 methylation islands were identified in the *C. virginica* genome (Venkataraman
et al., 2020). Methylation islands contained between 11 and 24,777 methylated CpGs, with a
median of 30 methylated CpGs per island. Lengths of methylation islands ranged from 500 to
1,236,482 base pairs, with a median length of 1,024 base pairs. The majority of methylation
islands (36,017; 97.2%) were less than 100,000 bp in length. There were 30,773 (83.0%)
methylation islands that overlapped with genic regions.
334
335
336

337
338
339 **Figure 1**. Frequency distribution of methylation ratios for CpG loci in *C. virginica* gonad tissue
340 DNA subjected to MBD enrichment. A total of 4,304,257 CpGs with at least 5x coverage
341 summed across all ten samples were characterized. Loci were considered methylated if they
342 were at least 50% methylated, sparsely methylated loci were 10-50% methylated, and
343 unmethylated loci were 0-10% methylated.
344
345

Figure 2. Proportion of CpG loci within genomic features. All CpGs are every dinucleotide in the *C. virginica* genome. Methylated CpGs refers to a dinucleotide with a methylation level of at least 50%.

Differential Methylation Analysis

A total of 598 CpG loci were differentially methylated between oysters exposed to control or high pCO_2, with 51.8% hypermethylated and 48.2% hypomethylated between treatments (Figure 3; Venkataraman 2020). When considering a PCA using methylation status of all CpG loci with 5x coverage across all samples, the first two principal components explained 29.8% of sample variation (Figure 4A). The first two principal components in a PCA with only differentially methylated loci (DML) explained 57.1% of the variation among treatments (Figure 4B). These DML were distributed throughout the *C. virginica* genome (Figure 5). The fifth chromosome had the most DML normalized by number of CpGs in the chromosome, and had the most genes; however, this was not the largest chromosome (Figure 5A).

Examination of DML within genes revealed that some genes contained multiple DML (Figure 5B-C). Of the 481 genes with DML, the majority only contained one DML (Figure 5B). There

365 were 48 genes with 2 DML, 16 genes with 3 DML, 6 genes with 4 DML and 1 gene with 5 DML
366 (Figure 5B). When multiple DML were found within a gene, they could be methylated in either
367 the same or opposite directions (Figure 5C).
368

369
Figure 3. Heatmap of DML in *C. virginica* reproductive tissue created using a euclidean
distance matrix. Samples in control pCO$_2$ conditions are represented by grey, and samples in
elevated pCO$_2$ conditions are represented by a black bar. Loci with higher percent methylation
are represented by darker colors. A logistic regression identified 598 DML, defined as individual
CpG dinucleotide with at least a 50% methylation change between treatment and control
groups, and a q-value < 0.01 based on correction for false discovery rate with the SLIM method.
The density of DML at each percent methylation value is represented in the heatmap legend.

384
385 **Figure 4**. Principal Components Analysis of a) all CpG loci with 5x coverage across samples
386 and b) DML. Methylation status of individual CpG loci explained 29.2% of variation between
387 samples when considering all CpG loci. Methylation status of DML explained 57.1% of sample
388 variation.
389
390
391

Figure 5. Distribution of DML among chromosomes and genes. (a) Number of DML normalized by number of CpG in each chromosome (bars) and number of genes (line) in each *C. virginica* chromosome. (b) Number of genes with various numbers of DML per gene (1-5). Most genes that contained DML only had 1 DML. (c) Proportion of hypermethylated, hypomethylated DML in genes with various numbers of DML per gene (1-5). Mixed refers to a classification of a gene that has both hypermethylated and hypomethylated DML.

Within the genome, DML were mostly present in genic regions, with 560 DML in 481 genes (368 DML in exons and 192 in introns). In addition, 42 DML were found in putative promoter regions, 27 in UTR, 57 in transposable elements, and 38 in intergenic regions. There were 21 DML located outside of exons, introns, transposable elements, and putative promoters. Additionally, 537 DML were found in methylation islands. The distribution of DML in *C. virginica* gonad tissue was higher in exons than expected for MBD-enriched CpG loci with minimum 5x coverage

across all samples (Contingency test; χ_2 = 401.09, df = 6, *P*-value < 2.2e-16; Figure 6). Of the
598 DML, 310 were hypermethylated and 288 were hypomethylated in the high *p*CO$_2$ treatment.
The number of hyper- and hypomethylated DML was almost evenly split within each genomic
feature, with the exception of putative promoter regions that had 44 hypermethylated DML
versus 23 hypomethylated DML. Within a gene, DML did not appear to be concentrated in one
particular region. The distribution of hyper- and hypomethylated DML along a gene do not differ
from each other (Figure 7).

415
416   **Figure 6**. Proportion CpG loci within putative promoters, untranslated regions (UTR), exons,
417   introns, transposable elements, and intergenic regions for MBD-enriched CpGs and differentially
418   methylated loci (DML). The distribution of DML in *C. virginica* gonad tissue in response to ocean
419   acidification differed from distribution of MBD-enriched loci with 5x coverage across control and
420   treatment samples (Contingency test; χ_2 = 401.09, df = 6, *P*-value < 2.2e-16).

421
The DML were found in genes responsible for various biological processes. However, no gene
ontology categories were significantly represented (Figure 8). The majority of genes with DML
were involved in protein ubiquitination processes. These genes were not consistently hyper- or
hypomethylated. Certain biomineralization genes did contain DML. The gene coding for
calmodulin-regulated spectrin-associated protein contained three hypomethylated and one
hypermethylated DML. Genes coding for EF-hand protein with calcium-binding domain,

calmodulin-binding transcription activator, and calmodulin-lysine N-methyltransferase contained

one or two hypermethylated DML.

Figure 7. Distribution of hyper- and hypomethylated DML along a hypothetical gene. The scaled

position of a DML within a gene was calculated by dividing the base pair position of the DML by

gene length. Counts of hypermethylated DML are plotted above the x-axis, and hypomethylated

DML counts are below the x-axis.

438
439 **Figure 8**. Biological processes represented by all genes used in enrichment background (%
440 Genes) and those with DML (% Genes with DML). Gene ontology categories with similar
441 functions are represented by the same color. Genes may be involved in multiple biological
442 processes. No gene ontologies were significantly enriched.

443 **Discussion**

444
The present study is a general description of DNA methylation in *C. virginica*, and is one of the
first to examine epigenetic responses to ocean acidification in the gonad tissue of a mollusc
species. Five hundred ninety-eight differentially methylated loci (DML) were identified in
response to the elevated pCO_2 treatments, most of which were in exons. Not only was DNA
methylation of *C. virginica* altered in response to ocean acidification, but changes in gonad
methylation indicates potential for these methylation patterns to be inherited by offspring.
451
Understanding how environmental stressors influence the epigenome is crucial when
considering potential acclimatization mechanisms in marine invertebrates. Our finding that high
pCO_2 impacts *C. virginica* DNA methylation adds to a growing body of work about ocean
acidification's impact on marine invertebrate methylomes. The coral species *P. damicornis*

demonstrated an overall increase in DNA methylation when exposed to low pH conditions (7.3 -
7.6) for six weeks, potentially influencing biomineralization (Putnam et al., 2016). Another coral
species, *S. pistillata*, also demonstrated an increase in genome-wide DNA methylation when
exposed to low pH conditions for two years. Changes in the methylome also modified gene
expression and altered pathways involved in cell cycle regulation (Liew et al., 2018b). The
present study on an oyster, however, did not observe the overall genome-wide increase in
methylation that was reported for corals. Instead, we found subtle, but significant, increases or
decreases in percent methylation at several hundred individual CpGs distributed across the
genome. As *C. virginica* and coral species are adapted to different environments and ecological
niches, it is possible that species-specific differences in methylation responses contribute to the
observed methylation pattern.
467
The *C. virginica* methylation landscape suggests a role for methylation in gene activity.
Approximately 22% of CpGs in the *C. virginica* gonad genome were methylated, which is
consistent with previous studies of marine invertebrate genomes Gavery and Roberts, 2013;
Olson and Roberts, 2014; Hofmann, 2017; Dimond and Roberts, 2020). Methylated loci were
concentrated in introns for *C. virginica*, followed by exons and transposable elements. This
location of methylated CpGs in gene bodies is consistent with what has been reported across
similar taxa (Roberts and Gavery, 2012; Eirin-Lopez and Putnam, 2018). The concentration of
methylated CpGs in gene bodies corresponds with proposed functionality in influencing gene
activity (Roberts and Gavery, 2012; Dixon et al., 2014; Liew et al., 2018b). Our study also found
methylation in transposable elements, putative promoters and intragenic regions. In plants,
transposable element methylation has been shown to modulate the effect of transposable
element insertion in genic regions (Hosaka and Kakutani, 2018). It is possible that methylation
of transposable elements in *C. virginica* could also limit the effect of transposable elements. The
characterization of methylation islands in the *C. virginica* genome demonstrates the viability of
this descriptive tool for future work examining methylation in mollusc species.
483
The presence of DML suggests that exposure to experimental ocean acidification conditions
elicits an epigenetic response. Many studies have documented changes to oyster protein
synthesis, energy production, metabolism, antioxidant responses, and reproduction in response
to ocean acidification (Tomanek et al., 2011; Timmins-Schiffman et al., 2014; Dineshram et al.,
2016; Boulais et al., 2017; Omoregie et al., 2019). Examination of methylation associated with
these physiological responses could identify mechanisms that contribute to these changes. For
example, our study found a hypomethylated DML in the heat shock protein 75 kDA gene, and
gene expression responses to ocean acidification in *C. virginica* have found downregulation in a
similar molecular chaperone, heat shock protein 70kDa (Beniash et al., 2010; Ivanina et al.,
2014). Other gene expression studies in bivalves have found changes in oxidative stress
proteins such as superoxide dismutase, cytochrome c, peroxiredoxin, and NADH
dehydrogenase (Chapman et al., 2011; Clark et al., 2013; Goncalves et al., 2016, 2017).
Although we did not find any DML in these genes, combined study of DNA methylation and
transcription may reveal how changes in gene expression are regulated in response to
environmental stressors.
499

Although DML were found across various genome features, they were mostly in exons and
introns. This is consistent with a recent study of *C. virginica* gill tissue found differentially
methylated regions in response to a salinity gradient were primarily in genic regions ((Johnson
and Kelly, 2019). Interestingly, DML were not found consistently in one particular region of a
gene. Similarly, methylated positions in genic regions were evenly distributed after the coral *S.*
pistillata was exposed to low pH (Liew et al., 2018b). Examination of another coral, *P. daedalea*,
in different temperature and salinity conditions found more frequent methylation at 5' and 3'
ends of genes (Liew et al., 2018a). We also found several genes with multiple DML. These DML
were not consistently hyper- or hypomethylated in the same gene. As hyper- and
hypomethylation may result in different transcriptional outcomes, future work should examine
the role of multiple DML on alternative splicing and gene expression.
511
The concentration of DML in gene bodies suggests a role for DNA methylation in gene
expression and regulation. A majority of genes with DML were involved in protein ubiquitination.
Protein ubiquitination is a post-translational protein modification that is involved in protein
synthesis and degradation (Peng et al., 2003; Komander, 2009). Previous studies in which
oysters were exposed to experimental ocean acidification conditions have demonstrated
changes in this pathway. For example, shotgun proteomic characterization of posterior gill
lamellae from adult *C. gigas* exposed to high pCO$_2$ revealed increased abundance of proteins
involved in ubiquitination and decreased protein degradation (Timmins-Schiffman et al., 2014).
Elevated pCO$_2$ levels were also found to upregulate malate dehydrogenase in adult *C. virginica*
mantle tissue (Tomanek et al., 2011). Several genes involved in protein ubiquitination, including
those for malate dehydrogenase, ubiquitin-protein ligase, RNA polymerase-associated protein,
and DNA damage-binding protein, were significantly hypermethylated in gonad tissue exposed
to elevated pCO$_2$. Hypermethylation of these genes may decrease transcriptional opportunities,
thus indicating a critical role in the response to ocean acidification.
526
Four genes involved in biomineralization contained DML, suggesting these genes can be
epigenetically regulated. Upregulation of calcium-binding gene expression has been previously
documented in *C. virginica* (Richards et al., 2018). Since the hypermethylated DML in these
genes are typically associated with reduced transcriptional opportunities, it is unclear how
methylation changes relate to gene expression for biomineralization genes. Many studies
examining ocean acidification-induced carryover effects in bivalves note changes to calcification
processes. For example, the Sydney rock oyster (*S, glomerata*) larvae exhibit faster shell
growth in high pCO$_2$ conditions when parents mature in those same conditions (Parker et al.,
2012, 2015). In contrast, larvae from other species found in the North Atlantic such as northern
quahog (hard clam; *M. mercenaria*) and bay scallops (*A. irradians*) developed slower when
parents were reproductively conditioned in low pH conditions (Griffith and Gobler, 2017). There
is some evidence to suggest that *C. virginica* larvae may be more resilient to high pCO$_2$
conditions than *M. mercenaria* or *A. irradians (Gobler and Talmage, 2014)*. Differential
methylation of biomineralization genes in *C. virginica* reproductive tissue could be a mechanism
to explain when parental experience impacts larval calcification if in fact these DML are
inherited.
543

Although our work documents significant changes to DNA methylation in reproductive tissue
after high pCO$_2$ exposure, this finding may be confounded by secondary effects of gonad
maturation. Specimens collected were from mixed populations, and sampled tissue contained
both mature and immature gametes. Reproductive tissue likely contained both gametic and
somatic cell types. Sex-specific effects have also been documented in response to ocean
acidification in mollusc species (Parker et al., 2018; Venkataraman et al., 2019). Lack of a
reproductive phenotype precludes any interpretation of how maturation stage or sex can
influence changes DNA methylation, as previous work in *C. gigas* demonstrates these factors
as significant influences on baseline methylation patterns (Zhang et al., 2018). Nevertheless,
differential methylation in stress response and biomineralization genes suggests that our study
does record epigenetic responses to ocean acidification. Future work should pair methylation
data with reproductive phenotypes to provide additional information on sex- or stage-specific
epigenetic responses to ocean acidification.

Conclusion
558
Our study found that *C. virginica* demonstrates a significant epigenetic response to elevated
pCO$_2$ exposure, with 598 DML identified. The concentration of these DML in gene bodies
suggests that methylation may be important for transcriptional control in response to
environmental stressors. As ocean acidification induced differential methylation in *C. virginica*
gonad tissue, there is a potential for intergenerational epigenetic inheritance, which could
control the gene activity of processes such as biomineralization. As carryover effects can persist
even when stressors are long-removed ((Venkataraman et al., 2019)), understanding the
mechanisms involved in intergenerational acclimatization is crucial. Future work should focus on
methylation patterns in adult *C. virginica* fully-formed gametes and larvae exposed to various
pCO$_2$ conditions to determine to what degree a difference in methylation influences gene activity
and how this might influence phenotypic plasticity.

Acknowledgements

This project was funded by National Science Foundation Biological Oceanography award
1635423 to KEL, JR, and SBR, and a Hall Conservation Genetics Research Award to YRV. This
work was facilitated through the use of advanced computational, storage, and networking
infrastructure provided by the Hyak supercomputer system at the University of Washington. We
also thank Mackenzie Gavery and our two reviewers for their insightful feedback on the
manuscript.

Data Accessibility

Raw sequence data is available at the NCBI Sequence Read Archive under BioProject
accession number PRJNA513384, with associated metadata and information also available at
Woods Hole Open Access Server: https://hdl.handle.net/1912/25138

Associated information for all analyses and supplemental material can be found in the Github
repository which is available in an archival format (Venkataraman 2020;
https://doi.org/10.6084/m9.figshare.11923479)

References

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., et al. (2012). methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol.* 13, R87.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Bao, W., Kojima, K. K., and Kohany, O. (2015). Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob. DNA* 6, 11.

Barton, A., Hales, B., Waldbusser, G. G., Langdon, C., and Feely, R. A. (2012). The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects. *Limnol. Oceanogr.* 57, 698–710.

Beniash, E., Ivanina, A., Lieb, N. S., Kurochkin, I., and Sokolova, I. M. (2010). Elevated level of carbon dioxide affects metabolism and shell formation in oysters Crassostrea virginica. *Mar. Ecol. Prog. Ser.* 419, 95–108.

Bird, A. (2002). DNA methylation patterns and epigenetic memory. *Genes Dev.* 16, 6–21.

Bossdorf, O., Richards, C. L., and Pigliucci, M. (2008). Epigenetics for ecologists. *Ecol. Lett.* 11, 106–115.

Boulais, M., Chenevert, K. J., Demey, A. T., Darrow, E. S., Robison, M. R., Roberts, J. P., et al. (2017). Oyster reproduction is compromised by acidification experienced seasonally in coastal regions. *Sci. Rep.* 7, 13276.

Byrne, M., Foo, S. A., Ross, P. M., and Putnam, H. M. (2019). Limitations of cross and multigenerational plasticity for marine invertebrates faced with global climate change. *Glob. Chang. Biol.* doi:10.1111/gcb.14882.

Chapman, R. W., Mancia, A., Beal, M., Veloso, A., Rathburn, C., Blair, A., et al. (2011). The transcriptomic responses of the eastern oyster, *Crassostrea virginica*, to environmental conditions. *Mol. Ecol.* 20, 1431–1449.

Clark, M. S., Thorne, M. A. S., Amaral, A., Vieira, F., Batista, F. M., Reis, J., et al. (2013). Identification of molecular and physiological responses to chronic environmental challenge in an invasive species: the Pacific oyster, *Crassostrea gigas*. *Ecol. Evol.* 3, 3283–3297.

Deans, C., and Maggert, K. A. (2015). What Do You Mean, "Epigenetic"? *Genetics* 199, 887–896.

Dickson, A. G. (1990). Standard potential of the reaction : $AgCl_{(s)} + 12H2_{(g)} = Ag_{(s)} + HCl_{(aq)}$, and and the standard acidity constant of the ion $HSO4_-$ in synthetic sea water from 273.15 to 318.15 K. *The Journal of Chemical Thermodynamics* 22, 113–127.

Dimond, J. L., and Roberts, S. B. (2016). Germline DNA methylation in reef corals: patterns and potential roles in response to environmental change. *Mol. Ecol.* 25, 1895–1904.

Dimond, J. L., and Roberts, S. B. (2020). Convergence of DNA Methylation Profiles of the Reef Coral *Porites astreoides* in a Novel Environment. *Frontiers in Marine Science* 6, 792.

Dineshram, R., Chandramouli, K., Ko, G. W. K., Zhang, H., Qian, P.-Y., Ravasi, T., et al. (2016).
Quantitative analysis of oyster larval proteome provides new insights into the effects of
multiple climate change stressors. *Glob. Chang. Biol.* 22, 2054–2068.

Dixon, G. B., Bay, L. K., and Matz, M. V. (2014). Bimodal signatures of germline methylation are
linked with gene expression plasticity in the coral *Acropora millepora*. *BMC Genomics* 15,
1109.

Eirin-Lopez, J. M., and Putnam, H. M. (2018). Marine Environmental Epigenetics. *Ann. Rev.*
Mar. Sci. doi:10.1146/annurev-marine-010318-095114.

Ekstrom, J. A., Suatoni, L., Cooley, S. R., Pendleton, L. H., Waldbusser, G. G., Cinner, J. E., et
al. (2015). Vulnerability and adaptation of US shellfisheries to ocean acidification. *Nat. Clim.*
Chang. 5, 207–214.

Feely, R. A., Alin, S. R., Newton, J., Sabine, C. L., Warner, M., Devol, A., et al. (2010). The
combined effects of ocean acidification, mixing, and respiration on pH and carbonate
saturation in an urbanized estuary. *Estuarine, Coastal and Shelf Science* 88, 442–449.
doi:10.1016/j.ecss.2010.05.004.

Gatzmann, F., Falckenhayn, C., Gutekunst, J., Hanna, K., Raddatz, G., Carneiro, V. C., et al.
(2018). The methylome of the marbled crayfish links gene body methylation to stable
expression of poorly accessible genes. *Epigenetics Chromatin* 11, 57.

Gavery, M. R., and Roberts, S. B. (2013). Predominant intragenic methylation is associated with
gene expression characteristics in a bivalve mollusc. *PeerJ* 1, e215.

Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J.-P., Middelburg, J. J., and Heip, C. H. R.
(2007). Impact of elevated CO_2 on shellfish calcification. *Geophys. Res. Lett.* 34, L07603.

Gish, W., and States, D. J. (1993). Identification of protein coding regions by database similarity
search. *Nat. Genet.* 3, 266–272.

Gobler, C. J., and Talmage, S. C. (2014). Physiological response and resilience of early life-
stage Eastern oysters (*Crassostrea virginica*) to past, present and future ocean
acidification. *Conserv Physiol* 2, cou004.

Gómez-Chiarri, M., Warren, W. C., Guo, X., and Proestou, D. (2015). Developing tools for the
study of molluscan immunity: The sequencing of the genome of the eastern oyster,
Crassostrea virginica. *Fish & Shellfish Immunology* 46, 2–4. doi:10.1016/j.fsi.2015.05.004.

Goncalves, P., Anderson, K., Thompson, E. L., Melwani, A., Parker, L. M., Ross, P. M., et al.
(2016). Rapid transcriptional acclimation following transgenerational exposure of oysters to
ocean acidification. *Mol. Ecol.* 25, 4836–4849.

Goncalves, P., Thompson, E. L., and Raftos, D. A. (2017). Contrasting impacts of ocean
acidification and warming on the molecular responses of CO_2-resilient oysters. *BMC*
Genomics 18, 431.

Griffith, A. W., and Gobler, C. J. (2017). Transgenerational exposure of North Atlantic bivalves
to ocean acidification renders offspring more vulnerable to low pH and additional stressors.
Sci. Rep. 7, 11394.

Helm, M. M., and Bourne, N. (2004). *Hatchery Culture of Bivalves: A Practical Manual*. Food &
Agriculture Org.

Hofmann, G. E. (2017). Ecological Epigenetics in Marine Metazoans. *Frontiers in Marine*
Science 4. doi:10.3389/fmars.2017.00004.

Hosaka, A., and Kakutani, T. (2018). Transposable elements, genome evolution and
transgenerational epigenetic variation. *Curr. Opin. Genet. Dev.* 49, 43–48.

IPCC, 2019: Summary for Policymakers. In: *IPCC Special Report on the Ocean and*
Cryosphere   in a Changing Climate [H.- O. Pörtner, D.C. Roberts, V. Masson-
Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J.
Petzold, B. Rama, N. Weyer (eds.)]. In press.

Ivanina, A. V., Hawkins, C., and Sokolova, I. M. (2014). Immunomodulation by the interactive
effects of cadmium and hypercapnia in marine bivalves *Crassostrea virginica* and
Mercenaria mercenaria. *Fish Shellfish Immunol.* 37, 299–312.

Jeong, H., Wu, X., Smith, B., and Yi, S. V. (2018). Genomic Landscape of Methylation Islands in
Hymenopteran Insects. *Genome Biol. Evol.* 10, 2766–2776.

Johnson, K. M., and Kelly, M. W. (2019). Population epigenetic divergence exceeds genetic
divergence in the Eastern oyster *Crassostrea virginica* in the Northern Gulf of Mexico. *Evol.*
Appl. doi:10.1111/eva.12912.

Komander, D. (2009). The emerging complexity of protein ubiquitination. *Biochem. Soc. Trans.*
37, 937–953.

Krueger, F., and Andrews, S. R. (2011). Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics* 27, 1571–1572.

Kurihara, H., Kato, S., and Ishimatsu, A. (2007). Effects of increased seawater $p\mathrm{CO_2}$ on early
development of the oyster *Crassostrea gigas*. *Aquat. Biol.* 1, 91–98.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods 9, 357–359.

Lee, Kitach Tae-Wook, Kim Byrne, Robert H Millero, Frank J Feely, Richard A Liu, Yong-Ming
(2010). The universal ratio of boron to chlorinity for the North Pacific and North Atlantic
oceans. *Geochimica et Cosmochimica Acta* 74, 1801–1811.

Lewis, E., and Wallace, D. W. (1998). R: Program developed for $\mathrm{CO_2}$ system calculations
ORNL/CDIAC-105. *Carbon Dioxide Information Analysis CentreOak Ridge National*
Laboratory, US Department of Energy, Oak Ridge, Tennessee.

Liew, Y. J., Howells, E. J., Wang, X., Michell, C. T., Burt, J. A., Idaghdour, Y., et al. (2018a).
Intergenerational epigenetic inheritance in reef-building corals. *bioRxiv*, 269076.
doi:10.1101/269076.

Liew, Y. J., Zoccola, D., Li, Y., Tambutté, E., Venn, A. A., and Michell, C. T. (2018b).
Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-building
coral. *Science Advances.*

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* 17, 10–12.

Olson, C. E., and Roberts, S. B. (2014). Genome-wide profiling of DNA methylation and gene
expression in Crassostrea gigas male gametes. *Front. Physiol.* 5, 224.

Omoregie, E., Mwatilifange, N. S. I., and Liswaniso, G. (2019). Futuristic Ocean Acidification
Levels Reduce Growth and Reproductive Viability in the Pacific Oyster (*Crassostrea gigas*).
J. Appl. Sci. Environ. Manage. 23, 1747–1754.

Parker, L. M., O'Connor, W. A., Byrne, M., Coleman, R. A., Virtue, P., Dove, M., et al. (2017).
Adult exposure to ocean acidification is maladaptive for larvae of the Sydney rock oyster
Saccostrea glomerata in the presence of multiple stressors. *Biol. Lett.* 13, 20160798.

Parker, L. M., O'Connor, W. A., Byrne, M., Dove, M., Coleman, R. A., Pörtner, H.-O., et al.
(2018). Ocean acidification but not warming alters sex determination in the Sydney rock
oyster, *Saccostrea glomerata*. *Proc. R. Soc. B* 285, 20172869.

Parker, L. M., O'Connor, W. A., Raftos, D. A., Pörtner, H.-O., and Ross, P. M. (2015).
Persistence of Positive Carryover Effects in the Oyster, *Saccostrea glomerata*, following
Transgenerational Exposure to Ocean Acidification. *PLoS One* 10, e0132276.

Parker, L. M., Ross, P. M., O'Connor, W. A., Borysko, L., Raftos, D. A., and Pörtner, H.-O.
(2012). Adult exposure influences offspring response to ocean acidification in oysters. *Glob.*
Chang. Biol. 18, 82–92.

Parker, L. M., Ross, P. M., O'Connor, W. A., Pörtner, H. O., Scanes, E., and Wright, J. M.
(2013). Predicting the response of molluscs to the impact of ocean acidification. *Biology* 2,
651–692.

Peng, J., Schwartz, D., Elias, J. E., Thoreen, C. C., Cheng, D., Marsischky, G., et al. (2003). A
proteomics approach to understanding protein ubiquitination. *Nat. Biotechnol.* 21, 921–926.

Putnam, H. M., Davidson, J. M., and Gates, R. D. (2016). Ocean acidification influences host
DNA methylation and phenotypic plasticity in environmentally susceptible corals. *Evol.*
Appl. 9, 1165–1178.

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* 26, 841–842.

Richards, M., Xu, W., Mallozzi, A., Errera, R. M., and Supan, J. (2018). Production of Calcium-
Binding Proteins in *Crassostrea virginica* in Response to Increased Environmental CO_2
Concentration. *Frontiers in Marine Science* 5, 203.

Ries, J. B. (2011). A physicochemical framework for interpreting the biological calcification
response to CO_2-induced ocean acidification. *Geochim. Cosmochim. Acta* 75, 4053–4064.

Roberts, S. B., and Gavery, M. R. (2012). Is There a Relationship between DNA Methylation
and Phenotypic Plasticity in Invertebrates? *Front. Physiol.* 2.
doi:10.3389/fphys.2011.00116.

Rondon, R., Grunau, C., Fallet, M., Charlemagne, N., Sussarellu, R., Chaparro, C., et al. (2017). Effects of a parental exposure to diuron on Pacific oyster spat methylome. *Environ Epigenet* 3. doi:10.1093/eep/dvx004.

Ross, P. M., Parker, L., and Byrne, M. (2016). Transgenerational responses of molluscs and echinoderms to changing ocean conditions. *ICES Journal of Marine Science*. doi:10.1093/icesjms/fsv254.

Roy, Rabindra N Roy, Lakshimi N Vogel, Kathleen M Porter-Moore, C Pearson, Tara Good, Catherine E Millero, Frank J Campbell, Douglas M (1993). The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* 44, 249–267.

Smit, A. F. A., Hubley, R., and Green, P. (2013). 2013--2015. RepeatMasker Open-4.0.

Strader, M. E., Wong, J. M., Kozal, L. C., Leach, T. S., and Hofmann, G. E. (2019). Parental environments alter DNA methylation in offspring of the purple sea urchin, *Strongylocentrotus purpuratus*. *J. Exp. Mar. Bio. Ecol.* 517, 54–64.

Suzuki, M. M., and Bird, A. (2008). DNA methylation landscapes: provocative insights from epigenomics. *Nat. Rev. Genet.* 9, 465–476.

Timmins-Schiffman, E., Coffey, W. D., Hua, W., Nunn, B. L., Dickinson, G. H., and Roberts, S. B. (2014). Shotgun proteomics reveals physiological response to ocean acidification in *Crassostrea gigas*. *BMC Genomics* 15, 951.

Tomanek, L., Zuzow, M. J., Ivanina, A. V., Beniash, E., and Sokolova, I. M. (2011). Proteomic response to elevated PCO2 level in eastern oysters, *Crassostrea virginica*: evidence for oxidative stress. *J. Exp. Biol.* 214, 1836–1844.

UniProt Consortium (2019). UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.* 47, D506–D515.

Van Heuven, S., Pierrot, D., Rae, J. W. B., Lewis, E., and Wallace, D. W. R. (2011). MATLAB program developed for CO_2 system calculations. *ORNL/CDIAC-105b. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee* 530.

Venkataraman, Y. R. (2020). Eastern oyster (*Crassostrea virginica*) gonad DNA methylation data and analysis. Available at: https://doi.org/10.6084/m9.figshare.11923479

Venkataraman, Y. R., Spencer, L. H., and Roberts, S. B. (2019). Larval Response to Parental Low pH Exposure in the Pacific Oyster *Crassostrea gigas*. *Journal of Shellfish Research* 38, 743. doi:10.2983/035.038.0325.

Waldbusser, G. G., Hales, B., Langdon, C. J., Haley, B. A., Schrader, P., Brunner, E. L., et al. (2014). Saturation-state sensitivity of marine bivalve larvae to ocean acidification. *Nat. Clim. Chang.* 5, 273.

Wang, H.-Q., Tuominen, L. K., and Tsai, C.-J. (2011). SLIM: a sliding linear model for estimating the proportion of true null hypotheses in datasets with dependence structures. *Bioinformatics* 27, 225–231.

Wright, R. M., Aglyamova, G. V., Meyer, E., and Matz, M. V. (2015). Gene expression
associated with white syndromes in a reef building coral, *Acropora hyacinthus*. *BMC*
Genomics 16, 371.

Zhang, X., Li, Q., Kong, L., and Yu, H. (2018). DNA methylation frequency and epigenetic
variability of the Pacific oyster *Crassostrea gigas* in relation to the gametogenesis. *Fish.*
Sci. 84, 789–797.

Zhao, L., Liu, L., Liu, B., Liang, J., Lu, Y., and Yang, F. (2019). Antioxidant responses to
seawater acidification in an invasive fouling mussel are alleviated by transgenerational
acclimation. *Aquat. Toxicol.* 217, 105331.

Zhao, L., Yang, F., Milano, S., Han, T., Walliser, E. O., and Schöne, B. R. (2018).
Transgenerational acclimation to seawater acidification in the Manila clam *Ruditapes*
philippinarum: Preferential uptake of metabolic carbon. *Sci. Total Environ.* 627, 95–103.
808