**Geographic and environmental contributions to genomic divergence in mangrove forests**

Michele Fernandes da Silva[1,2], Mariana Vargas Cruz[1,2], João de Deus Vidal Júnior[1,2], Maria

Imaculada Zucchi[3], Gustavo Maruyama Mori[4], Anete Pereira de Souza[1,2]

[1]Department of Plant Biology, Institute of Biology, University of Campinas (Unicamp),

Campinas, SP, 13083-863, Brazil

[2]Center for Molecular Biology and Genetic Engineering, University of Campinas (Unicamp),

Campinas, SP, 13083-875, Brazil

[3]São Paulo Agency for Agribusiness Technology (APTA), Piracicaba, SP, 13400-790, Brazil

[4]Institute of Biosciences, São Paulo State University (Unesp), São Vicente, SP, 11330-900,

Brazil

**Corresponding author:** Anete Pereira de Souza

Center for Molecular Biology and Genetic Engineering, University of Campinas (Unicamp).

Avenue Cândido Rondom, 400, Campinas, SP, 13083-875, Brazil.

Telephone number: +55 19 3521 1132

E-mail: anete@unicamp.br

**ABSTRACT**

Assessing the relative importance of geographic and environmental factors to the spatial

distribution of genetic variation can provide relevant information about the processes that

maintain genetic variation in natural populations. With a globally wide but very restricted habitat

distribution, mangrove trees are an interesting model for studies aiming to understand the

contributions of these factors. Mangroves occur along the continent-ocean interface of tropical

and subtropical latitudes, regions considered inhospitable to many other plant types. Here, we

used landscape genomics approaches to investigate the relative contributions of geographic and

environmental variables to the genetic variation of two black mangroves, *Avicennia schaueriana*

and *A. germinans*, along the South American coast. Using single nucleotide polymorphisms

(SNPs), our results revealed an important role of ocean currents in the gene flow of *A.*

*schaueriana* and isolation by environment as the pattern that best explains the genetic

differentiation of *A. germinans*. Additionally, for both species, we found significant correlations

between genetic variation with evidence of selection and precipitation regimes, tidal variation,

solar radiation and temperature patterns. These discoveries expand our knowledge about the

evolution of mangrove trees and provide important information to predict future responses of

coastal species to the expected global changes for this century.

**KEYWORDS**

Adaptation of mangroves; isolation by distance; isolation by barrier; environmental gradient;

molecular ecology; coastal ecology.

## 1. INTRODUCTION

Environmental and geographic variations are often determinants of the distribution of allele frequencies in natural populations, facilitating or limiting gene flow across space and influencing the selection of certain genotypes [1-3]. For example, under an isolation-by-distance (IBD) model [4], geographic distance may limit the dispersal, leading to the accumulation of allele frequency divergences by genetic drift [5]. Furthermore, under the isolation-by-barrier (IBB) model, a barrier to gene flow may abruptly reduce or even disrupt connectivity between individuals of a species [6]. In addition to these models, the genetic differentiation can increase in response to environmental differences, regardless of the geographic distance. This pattern, described by the isolation-by-environment (IBE), [7] can be generated by a variety of ecological processes, such as selection against immigrants, leading to the evolution of locally adapted populations [5].

These models, IBD, IBB, and IBE, are not mutually exclusive and often co-occur in nature [8,9]. Determining the factors that control the distribution patterns of genetic variation across space can provide relevant information about the underlying processes that generate and maintain genetic variation in natural populations [7,10]. This knowledge is essential to predict future responses of current populations to environmental changes [11] and can contribute to decision-making processes aiming to minimize future biodiversity loss [12-14].

The field of research that seeks to clarify the roles of these factors in the distribution of the neutral and adaptive genetic variability of a species over space is known as landscape genomics [11,15-17]. Recently, this approach has been increasingly applied to the study of non-model organisms [18]. Nevertheless, landscape genomic studies are mostly limited to animal species, whereas studies on plants, especially tropical trees [19], remain very limited despite their

fundamental role in global biogeochemical cycles [20] and as habitat providers to most terrestrial

biodiversity [21].

As sessile organisms, trees respond directly to the environment in which they live [22].

Conversely, they often have high levels of gene flow, which tend to reduce the strength of

natural selection [23]. Thus, tree species that often occur across wide latitudinal ranges along

environmental gradients, such as mangrove forests [24], represent a promising biological model

to investigate and understand the effects of the environment on micro evolutionary processes and

populational dynamics.

Mangrove trees occur in a narrow area along the continent-ocean interface [24] of

tropical and subtropical latitudes of the world, mainly between 30ºN and 30°S [25]. These

species produce floating seeds or fruits, referred to as "propagules" [24], which can disperse over

long distances via rivers and ocean surface currents [26].Their geographic distribution is limited

by the topography of the intertidal zone [27] and frequently by the occurrence of low

temperatures [28] and precipitation patterns [29]. However, environmental variations in their

boundaries are associated with different climatic thresholds [30,31] and oceanographic

conditions [32].

*Avicennia* L. (Acanthaceae) is one of the most diverse and widely distributed mangrove

genera globally [33,34] and is very abundant on the western Atlantic coastline [35]. In this

region, *Avicennia* is mainly represented by two of the three New World *Avicennia* species,

namely, *Avicennia germinans* and *Avicennia schaueriana* [33,35]. These species present a partly

sympatric distribution on the South American coast, where they share a remarkable north-south

pattern of genetic divergence as revealed by selectively neutral microsatellites [36] and single

nucleotide polymorphisms (SNPs) [37,38]. A similar pattern of putatively neutral genetic

diversity was also observed in other coastal species, such as *Rhizophora mangle* [39,40] and the

mangrove-associated tree *Hibiscus pernambucensis* [41]. These findings likely indicate a

prominent role of dispersal of floating propagules in shaping the overlapping north-south genetic

divergence of these trees given the direction of the South Equatorial current (SEC) along the

Atlantic coast of South America [42].

In addition to neutral processes that shape the diversity of *Avicennia* species in this

region, recent studies have identified various genomic regions possibly associated with adaptive

processes relevant to the environmental context of mangroves [37,38]. These variations have

been attributed to climatic and oceanographic factors that vary widely along the latitudinal

gradient of the species distribution. Although these studies provide insights into the role of the

environment in the organization of the adaptive genetic variation in *A. schaueriana* and *A.*

*germinans*, the relative importance of neutral and nonneutral abiotic factors remains unknown.

The present study aimed to investigate the relative contributions of geographic and

environmental variables to the organization of the genetic diversity of these two dominant

species of the genus *Avicennia* that are found on the Atlantic coast of South America.

**2. METHODS**

**2.1. Biological materials and genotyping of SNP markers**

Biological materials were collected and single nucleotide polymorphism (SNP) markers

were identified within the context of previous studies by Cruz et al. [37] for *A. schaueriana* and

Cruz et al. [38] for *A. germinans*. Briefly, seventy-seven *A. schaueriana* individuals were

sampled from ten different locations and 48 *A. germinans* individuals were sampled from six

different locations along the Brazilian coast (figure 1 and table 1). Assembly, mapping and SNP

locus identification were performed using custom scripts (SNPsaurus), which created a reference

catalogue of abundant reads, retaining biallelic loci present in at least 10% of the samples. High-

quality sequences were used, allowing a maximum of 65% of missing data and one SNP per

sequence and requiring a minimum coverage of 8× and a minor allele frequency $\geq 0.05$ using

vcftools v.0.1.12b [43]. To reduce the paralogy or low-quality genotype calls, a maximum

reading coverage of 56 was used (resulting from the product with the average reading depth and

a standard deviation of 1.5 from the average). In total, 6,170 and 2,297 SNP markers were

identified for *A. schaueriana* and *A. germinans*, respectively.

**2.2. Detection of SNP loci with signatures of natural selection**

SNP loci with evidence of natural selection were identified previously by Cruz et al.

[37,38]. For *A. schaueriana*, 86 loci showed considerable deviations from neutral expectations of

interpopulation divergence. They were detected by two methods to minimize false positives:

LOSITAN [44], with a confidence interval of 0.99 and a false-discovery rate (FDR) of 0.1, and

pcadapt [45], with an FDR of 0.1. For *A. germinans*, 25 loci showed considerable deviations

from neutral expectations of interpopulation divergence. For the latter species, in addition to

LOSITAN (a confidence interval of 0.99 and an FDR of 0.05) and pcadapt (an FDR of 0.05),

SNP loci associated with ecological variables were detected using Latent Factor Mixed Models

(LFMM) implemented in the *LEA* package [46].

**2.3. Estimation of genetic distances**

To analyze the importance of geographic distance and environmental variables in spatial

genetic divergence, we evaluated which model—isolation by distance (IBD), isolation by barrier

(IBB) or isolation by environment (IBE) — best described the distribution of the genetic

diversity of each species based on genome-wide SNP markers. To that end, we estimated the

pairwise genetic differentiation (Wright's $F_{ST}$) [47] for the total set of SNP molecular markers

and the set of SNP markers with evidence of selection, using the *Hierfstat* [48] package for R v.

3.6.2 [49].

**2.4. Geographic and environmental distances and the oceanographic barrier**

Pairwise geographic distances among populations were measured using the geographic

coordinates of the sampling sites (table 1) with the global positioning system (Garmin 76CSx,

WGS-84 standard, Garmin International Inc., Olathe, KS, USA). Distances between points were

estimated based on the contour of the Brazilian coast; thus, we considered floating propagules-

mediated dispersal [50]. To determine the relative significance of the split pattern of the SEC

into the Brazil current (BC) and the North Brazil current (NBC) [42] (figure 1) for *A.*

*schaueriana* and *A. germinans* propagules, a binary matrix (0 or 1) was constructed based on the

presence (1) or absence (0) of the supposed oceanographic barrier between each pair of sampling

sites.

To evaluate the overall effect of the environment on the distribution of genetic variation,

we obtained a total of 42 environmental variables for each sampling site, with a resolution of 30

arc-seconds (~ 1 km in Ecuador). In our dataset, we included 27 climatic variables derived from

the WorldClim temperature and precipitation datasets (version 1.4 and 2.0) [51], 10

oceanographic variables derived the Marspec ocean surface salinity and temperature datasets

[52] and five variables related to tidal variations retrieved from the Environmental Climate Data

Sweden (ECDS) platform [53]. To avoid nonindependence between environmental variables, we

removed variables that showed a high correlation ($R^2 > 0.8$) (electronic supplementary material,

figure S1), as measured by the *removeCollinearity* function of the *virtualspecies* package [54] in

R [49]. We extracted the values of the environmental variables for our sample points (electronic

supplementary material, table S1 and S2) using the *raster* package [55] in R [49]. All occurrence

data have been carefully inspected to detect and correct problems associated with inconsistent

records [56]. We transformed this environmental data matrix using a principal component

analysis (PCA). We used the most representative scores from the PCA to calculate Euclidean

distances between population pairs [57]. The PCA and environmental distance measurements

were all performed in R [49].

**2.5. Association tests**

To test which model best explained our genetic data (both neutral and putatively

nonneutral), we performed Multiple Matrix Regression with Randomization (MMRR) [9]. We

employed this method to estimate the independent effect of each factor (geographic or

environmental) on a single response variable, typically genetic distance. The analysis was

performed using the *MMRR* function in R with 10,000 permutations. This function calculates the

coefficient of determination, the regression coefficients (β), and estimates significance values for

all parameters, showing to be accurate to several types of conditions [9]. The following

comparisons were performed: (I) geographic distance (km) and environmental distance, (II)

geographic distance (km) and pairwise $F_{ST}$, (III) environmental distance  and pairwise $F_{ST}$, (IV)

geographic distance (km) in conjunction with environmental distance and pairwise $F_{ST}$ and (V)

the presence/absence of the supposed oceanographic barrier and pairwise $F_{ST}$. These

comparisons were also performed separately for the set of samples located northerly from the

SEC and for the set of samples located southerly from the SEC because of previous genetic

structure evidences [36]. Finally, we tested for correlations between the genetic distances based

on the SNP marker set with selection evidence for each environmental variable separately.

**3. RESULTS**

**3.1. Genetic, geographic and environmental distances**

For the total set of SNP markers, we found $F_{ST}$ values between 0.019 and 0.189 for pairs

of *A. schaueriana* populations and between 0.047 and 0.387 for pairs of *A. germinans*

populations (figure 2). For the set of markers with evidence of selection, $F_{ST}$ ranged from 0.02 to

0.36 for *A. schaueriana* (electronic supplementary material, table S3) and from 0.03 to 0.88 for

*A. germinans* (electronic supplementary material, table S4). The pairwise geographic distances

ranged from approximately 5000 km between Bragança (SB) and Laguna (SL) to 77 km between

Cananéia (SC) and Pontal do Paraná (SP) for *A. schaueriana* (electronic supplementary material,

table S5) and from 2100 km between Ilha de Marajó in Soure (GM) and Tamandaré (GT) to 222

km between Ilha de Marajó in Soure (GM) and Bragança (GB) for *A. germinans* (electronic

supplementary material, table S6).

After removing highly correlated environmental variables ($R^2 > 0.8$), 23 variables were

retained for analyses of environmental distances (electronic supplementary material, table S7 and

S8). The first five and the three most representative axes of the PCA used to calculate the

environmental distance between the sampling sites (electronic supplementary material, table S9

and S10) retained 97% and 92% of the variation for *A. schaueriana* and *A. germinans*,

respectively. The first two PCA axis for *A. schaueriana* represented 80% of the variance and was

mainly explained by the variations in air temperature and sea surface temperature, the solar

radiation and tidal cycles (figure 3A). For *A. germinans*, the first two PCA axis represented

almost 80% of the variance and was mainly explained by the air temperature variation,

precipitation regimes, vapor pressure deficit and solar radiation (figure 3B).

**3.2. Association tests**

For *A. schaueriana*, considering all SNP loci, we observed that the geographic distances

between pairs of sampling sites played the greatest role in the describing genetic distances ($R^2 =$

0.81, $P < 0.001$; $\beta = 0.90$, $P = 0.001$; figure 4A), but environmental distance also contributed

significantly to genetic distance ($R^2 = 0.54$, $P < 0.001$; $\beta = 0.73$, $P < 0.01$; Figure 4B). However,

there was a strong significant correlation between geographic distance and environmental

difference ($R^2 = 0.82$, $P < 0.05$; $\beta = 0.88$, $P = 0.001$; Figure 4C), implying that the farther the

geographic distance, the greater the environmental difference. When tests were performed

separately for sampling locations north and south from the SEC, we found significant

correlations only between genetic and geographic distances ($R^2 = 0.81$, $P < 0.001$; $\beta = 0.89$, $P =$

0.01) for sampling sites located south of the SEC. Multiple regression combining the effects of

geographic and environmental distances on genetic distance provided a slight increase in the fit

of the model ($R^2 = 0.86$, $P < 0.05$), however, the effect of geographic ($\beta$geographic $= 0.13$, $P =$

0.001) and environmental ($\beta$environment $= -0.48$, $P = 0.01$) distances was lower than the

regressions including only one predictor variable. The variable that best explained the genetic

differentiation observed in *A. schaueriana* was the presence or absence of the oceanographic

barrier ($R^2 = 0.92$, $P < 0.01$; $\beta = 0.96$, $P < 0.01$).

For *A. germinans*, geographic ($R^2 = 0.61$, $P < 0.05$; $\beta = 0.78$, $P = 0.05$; figure 4E) and

environmental ($R^2 = 0.67$, $P < 0.05$; $\beta = 0.81$, $P = 0.01$; figure 4F) distances played significant

roles in the genetic differentiation between sampling sites in the entire SNP dataset. However,

when we removed samples from GT, which is the only sampling site south of the SEC, the

genetic divergence was not correlated with either the geographic distance ($R^2 = 0.006$, $P = 0.97$;

$\beta = 0.07$, $P = 0.88$) or the environmental distance ($R^2 = 0.03$, $P = 0.70$; $\beta = 0.17$, $P = 0.68$). The

correlation between geographic and environmental distances was significant ($R^2 = 0.74$, $P <$

$0.01$; figure 4G), but less pronounced than that observed for *A. schaueriana*, which was probably

due to the lower latitudinal amplitude between their sampling sites, and consequently, lower

environmental variation compared to the latter species. When both types of distances were

included in the model, we obtained significant values only for the environmental distance, which

showed twice the effect than the geographical distance ($R^2 = 0.69$, $P = 0.03$; $\beta_{environment} = 0.55$,

$P < 0.05$, $\beta_{geographic} = 0.28$, $P = 0.34$). The presence or absence of the oceanographic barrier did

not explain the genetic divergence among *A. germinans* sampling locations ($R^2 = 0.91$, $P = 0.15$;

$\beta = 0.95$, $P = 0.12$).

We identified a significant environmental contribution to the genetic differentiation based

on the putative nonneutral SNPs for both species. This contribution was more pronounced for *A.*

*germinans* ($R^2 = 0.62$, $P < 0.05$; $\beta = 0.78$, $P < 0.05$; figure 4H) than for *A. schaueriana* ($R^2 =$

$0.51$, $P < 0.01$; $\beta = 0.7$, $P = 0.001$; figure 4D). When we analyzed each environmental variable

separately, we found significant correlations of putative nonneutral genetic differentiation among

*A. schaueriana* sampling locations with variables that vary along the latitudinal gradient of the

species, such as the annual mean temperature ($R^2 = 0.45$, $P < 0.001$; $\beta = 0.67$, $P = 0.001$),

isothermality ($R^2 = 0.75$, $P < 0.001$; $\beta = 0.86$, $P = 0.001$), precipitation seasonality ($R^2 = 0.56$, $P$

$< 0.001$; $\beta = 0.75$, $P < 0.01$), minimum solar radiation ($R^2 = 0.49$, $P < 0.001$; $\beta = 0.70$, $P = $

$0.001$), the annual average cycle of tidal amplitude ($R^2 = 0.62$, $P < 0.001$; $\beta = 0.79$, $P < 0.05$) and

the mean annual of sea surface temperature ($R^2 = 0.46$, $P < 0.001$; $\beta = 0.68$, $P = 0.01$). For *A.*

*germinans*, the nonneutral variation was related to variations in the annual mean temperature ($R^2$

$= 0.81$, $P = 0.01$; $\beta = 0.88$, $P < 0.05$), maximum temperature of warmest month ($R^2 = 0.80$, $P < $

$0.05$; $\beta = 0.88$, $P < 0.05$), precipitation of warmest quarter ($R^2 = 0.91$, $P < 0.05$; $\beta = 0.95$, $P = $

$0.05$), maximum solar radiation ($R^2 = 0.96$, $P = 0.01$; $\beta = 0.98$, $P = 0.01$) and in the annual

average cycle of tidal amplitude ($R^2 = 0.12$, $P < 0.01$; $\beta = 0.35$, $P = 0.01$) (table 2).

**4. DISCUSSION**

The study of the geographic and environmental factors shaping neutral and adaptive

genetic variation in heterogeneous environments is one of the main approaches used to

understand the dynamics and evolutionary potential of natural populations [16,58]. In this study,

we analyzed the relative contributions of environmental and geographic distances and the

presence of an oceanographic barrier to the genetic differentiation of populations of two

dominant mangrove species in the Neotropics. We identified the relative importance of the main

environmental variables that generate adaptation in these species, providing relevant information

for decision makers when planning future efforts targeting conservation and coastal vegetation

recovery in the face of increasing challenges resulting from anthropogenic environmental and

climate changes in this century.

Pairwise $F_{ST}$ results revealed a variable degree of genetic divergence for both species,

indicating the existence of substantially structured populations, especially when samples to the

north and south of the SEC are considered. These results corroborate the genetic structure

patterns found in previous studies conducted with neutral molecular markers [36,39-41] and

indicate that regardless of the characteristics of these markers with high (microsatellite) [59] or

low (SNPs) [60] mutation rates, the evolutionary processes that led to this divergence must be

quite intense or ancient.

The geographic distances between sampling sites contributed significantly to the genetic

divergence of *A. schaueriana* ($R^2$: 0.81, $P < 0.001$; $\beta = 0.90$, $P = 0.001$) and *A. germinans* ($R^2 =$

0.61, $P < 0.05$; $\beta = 0.78$, $P = 0.05$), suggesting that spatial distance plays a fundamental role in

the genetic divergence of these species' populations. This model appears to be common in

studies of plants [8] and mangroves [61-64]. Although evidence of water-dispersed over long

distances exists for mangrove species [26,36,65,66], our results indicate that the large geographic

extension and major oceanic currents of the Brazilian coast physically limit the dispersal of

*Avicennia* species. Although these correlations were significant for the total set of sampling

locations, they were not significant when considering only the set of *A. schaueriana* and *A.*

*germinans* sampling locations north of the SEC. For the five sampling locations of *A. germinans*

in this region, which are distributed in a fairly geographically continuous habitat, the model was

not explained by the coefficient of determination values ($R^2 = 0.006$, $P = 0.97$). We hypothesized

that the genetic differentiation between 0.047 and 0.387 observed among sampling locations may

actually result from an IBB effect caused by the SEC acting as a barrier to the dispersal of

propagules. However, we did not find a significant correlation between genetic differentiation

and the presence of the SEC, probably because we were not able to sufficiently sample *A.*

*germinans* south of the SEC given that only a single location, GT, was used.

In contrast, we found statistical evidence for the action of the SEC as a barrier to gene

flow for *A. schaueriana*. For this species, the IBB is the model that best explains the genetic

differentiation between their populations. Our results corroborate Mori et al. [36], who suggested

that the neutral genetic divergence observed for *A. schaueriana* may have been shaped by marine

currents. Our results also suggest that the bifurcated flow of marine currents along the Atlantic

coast of South America may have a key role as drivers of genetic differentiation observed in

other species of mangrove or associated with this ecosystem, such as *Rhizophora mangle* [39,40]

and *Hibiscus pernambucensis* [41]. Our findings statistically showed that the SEC is an

important driver of the genetic structure of mangrove species; however, coastal and ocean

currents vary temporally in strength and directionality [50]. For example, the SEC splits into the

BC and the NBC, which have different speeds and directions [42] (Fig. 1). The NBC is faster

than the BC, favoring the spread of propagules from south to north, as observed in previous

studies [36,39]. Additionally, the flow direction of coastal currents may influence the direction of

gene flow among populations. Therefore, future investigations of the dynamics of these currents,

not as a static barrier but including different levels of resistance to gene flow, may provide more

realistic insights into their influence on the distribution of the genetic variation of species

dispersed by sea currents.

We also identified an IBE pattern in the structure of the genetic diversity of *A.*

*schaueriana* and *A. germinans*. For *A. germinans*, the IBE explained the genetic differentiation

between their populations better than the neutral processes.  Many species under IBE showed

this same pattern [8,67-73], indicating that environmental heterogeneity may be the main factor

underlying geographic distance [8,72,74]. Notably, the environmental differences of both species

were correlated with geographic distance. We suggest, therefore, future investigations that

control the effect of environmental predictors on geographic distance.

Given the significant results for IBE, we also estimated the contribution of each

environmental variable to nonneutral SNP genetic variation, allowing us to infer the degree to

which individual environmental variables contribute to these results. For *A. schaueriana*, we

found significant correlations to variations in the tidal range regimes, air and ocean surface

temperatures, water vapor pressure, solar radiation, and precipitation shape the distribution of the

genetic variation of this species (Table 2). Our results corroborate those of Cruz et al. [38] who

performed a functional annotation of loci showing deviations from neutral expectations based on

allele frequency variations and associated these loci with biological processes related to

responses to environmental variables such as drought and UV radiation. Conversely, for *A.*

*germinans*, we found that genetic variation with evidence of selection is correlated with

variations in atmospheric temperature patterns, precipitation regimes, solar radiation and tidal

amplitude cycles (Table 2). The results observed for both species corroborate those of other

studies reporting that precipitation and air and sea temperature variables were limiting factors in

the distribution of coastal species [30,32,75-78], and, together with Cruz et al. [37,38], provide

new evidence of the role of local adaptation in the distribution of the genetic diversity of these

species.

Our results are particularly relevant in light of the climate changes that have been

occurring in the last decades. With them, further investigations on the responses of these species

to future changes in predicted increases in average annual air and ocean temperatures and sea

level rise [79] can be directed. These results also have implications for conservation management

and planning decisions. With a better understanding of the role of environmental variables in the

modeling of genetic variation in *Avicennia* species on the South American coast, our results

contribute to the definition of evolutionarily significant units (ESU) for these species, thus

maintaining their evolutionary potential [80]. In addition, from the results referring to the

remarkable pattern of genetic divergence of *A. schaueriana*, resulting in part from the action of

the current of the marine surface studied here, we suggest the consideration of genetic banks

north and south from the SEC as conservation measures and maintenance of biodiversity.

**5. CONCLUSIONS**

Our results provide novel evidence that the combined action of geographic distance,

ocean currents and environmental gradient contributed to the evolution of spatial genetic

divergence within two Neotropical *Avicennia* species on the Atlantic coast of South America.

Our findings revealed the major role of ocean currents as barriers to the gene flow of *A.*

*schaueriana* and the environmental gradient as the main abiotic factor in the genetic

differentiation of *A. germinans*. For both species, we found new evidence of adaptation to

climatic and oceanographic variables that are changing rapidly in recent decades. In this context,

our results represent a starting point for understanding the role of environmental variables in

shaping genetic variation in *Avicennia* species of the South American coast, contributing to the

understanding of the evolution of genetic diversity within this genus and providing information

that can be relevant to other coastal organisms. Additionally, our findings guide further research

on the responses of these species to climate change and contribute to effective conservation plans

for mangroves and other species with similar dispersion mechanisms.

**ACKNOWLEDGEMENTS**

We thank Alessandro Alves Pereira and Prianda Rios Laborda for critically reading the

manuscript. Also, we are truly thankful for Coordination for the Improvement of Higher

Education Personnel (CAPES), São Paulo Research Foundation (FAPESP) and Brazilian

National Council for Scientific and Technological Development (CNPq) for their financial

support.

**FUNDING**

This study was financed in part by the Coordination for the Improvement of Higher Education

Personnel - CAPES Computational Biology Program (88887.177151 / 2018-00, 88882.160095 /

2013-01), the São Paulo Research Foundation - FAPESP (2013/08086-1, 2013/26793-7,

2014/22821-9, 2018 / 18431-1) and Brazilian National Council for Scientific and Technological

Development – CNPq (448286/2014-9, 153973/2018-8, 309661/2014-5).

**DATA, CODE AND MATERIALS**

*A. germinans* SNP genotypes data are available from the Dryad Digital Repository:

https://doi.org/10.5061/dryad.pk0p2nghr. *A. schaueriana* SNP genotypes data are available from

the article: doi:10.1038/s41598-019-56469-w (supplementary datasets 1 and 2). The

environmental data set and distance matrices have been uploaded as part of the electronic

supplementary material.

**COMPETING INTERESTS**

The authors declare no competing interests.

**AUTHORS' CONTRIBUTIONS**

A.P.S., G.M.M., M.F.S. and M.V.C. designed the study; M.F.S., M.V.C., G.M.M. and J.D.V.

analyzed data; A.P.S., G.M.M. and M.I.Z. contributed material/analysis tools; M.F.S. wrote the

manuscript. All authors discussed the results and contributed to the manuscript and reviewed it.

**REFERENCES**

1.  Murray KD, Janes JK, Jones A, Bothwell HM, Andrew RL, Borevitz JO. 2019 Landscape drivers of genomic diversity and divergence in woodland Eucalyptus. *Mol. Ecol.* **28**, 5232–5247. (doi:10.1111/mec.15287)

2.  Sork VL. 2016 Gene flow and natural selection shape spatial patterns of genes in tree populations: implications for evolutionary processes and applications. *Evol. Appl.* **9**, 291–310. (doi:10.1111/eva.12316)

3.  Wang IJ, Glor RE, Losos JB. 2013 Quantifying the roles of ecology and geography in spatial genetic divergence. *Ecol. Lett.* **16**, 175–182. (doi:10.1111/ele.12025)

4.  Wright S. 1943 Isolation by distance. *Genetics* **28**, 114–138.

5.  Bradburd GS, Ralph PL, Coop GM. 2013 Disentangling the effects of geographic and ecological isolation on genetic differentiation. *Evolution* **67**, 3258–3273. (doi:10.1111/evo.12193)

6.  Barton NH. 1979 Gene flow past a cline. *Heredity* **43**, 333–339. (doi:10.1038/hdy.1979.86)

7.  Wang IJ, Bradburd GS. 2014 Isolation by environment. *Mol. Ecol.* **23**, 5649–5662. (doi:10.1111/mec.12938)

405    8.      Sexton JP, Hangartner SB, Hoffmann AA. 2014 Genetic isolation by environment or

406            distance: which pattern of gene flow is most common? *Evolution* **68**, 1–15.

407            (doi:10.1111/evo.12258)

408    9.      Wang IJ. 2013 Examining the full effects of landscape heterogeneity on spatial genetic

409            variation: a multiple matrix regression approach for quantifying geographic and

410            ecological isolation. *Evolution* **67**, 3403–3411. (doi:10.1111/evo.12134)

411    10.     Lee CR, Mitchell-Olds T. 2011 Quantifying effects of environmental and geographical

412            factors on patterns of genetic differentiation. *Mol. Ecol.* **20**, 4631–4642.

413            (doi:10.1111/j.1365-294x.2011.05310.x)

414    11.     Vincent B, Dionne M, Kent MP, Lien S, Bernatchez L. 2013 Landscape genomics in

415            *Atlantic salmon* (Salmo Salar): searching for gene-environment interactions driving local

416            adaptation. *Evolution* **67**, 3469–3487. (doi:10.1111/evo.12139)

417    12.     Kovach RP, Gharrett AJ, Tallmon DA. 2012 Genetic change for earlier migration timing

418            in a pink salmon population. *Proc. R. Soc. B: Biol. Sci.* **279**, 3870–3878.

419            (doi:10.1098/rspb.2012.1158)

420    13.     Muñoz NJ, Farrell AP, Heath JW, Neff BD. 2015 Adaptive potential of a Pacific salmon

421            challenged by climate change. *Nat. Clim. Change* **5**, 163–166.

422            (doi:10.1038/nclimate2473)

423    14.     Wee AKS *et al.* 2019 The integration and application of genomic information in

424            Mangrove conservation. *Conserv. Biol.* **33**, 206–209. (doi:10.1111/cobi.13140)

425    15.     Joost S, Bonin A, Bruford MW, Després L, Conord C, Erhardt G, Taberlet P. 2007 A

426            Spatial Analysis Method (SAM) to detect candidate loci for selection: towards a

landscape genomics approach to adaptation. *Mol. Ecol.* **16**, 3955–3969.

(doi:10.1111/j.1365-294x.2007.03442.x)

16.   Schoville SD, Bonin A, François O, Lobreaux S, Melodelima C, Manel S. 2012 Adaptive

genetic variation on the landscape: methods and cases. *Annu. Rev. Ecol. Evol. Syst.* **43**,

23–43. (doi:10.1146/annurev-ecolsys-110411-160248)

17.   Lowry DB. 2010 Landscape evolutionary genomics. *Biol. Lett.* **6**, 502–504.

(doi:10.1098/rsbl.2009.0969)

18.   Storfer A, Patton A, Fraik AK. 2018 Navigating the interface between landscape genetics

and landscape genomics. *Front. Genet.* **9**. (doi:10.3389/fgene.2018.00068)

19.   Storfer A, Murphy MA, Spear SF, Holderegger R, Waits LP. 2010 Landscape genetics:

where are we now? *Mol. Ecol.* **19**, 3496–3514. (doi:10.1111/j.1365-294x.2010.04691.x)

20.   Jasechko S, Sharp ZD, Gibson JJ, Birks SJ, Yi Y, Fawcett PJ. 2013 Terrestrial water

fluxes dominated by transpiration. *Nature* **496**, 347–350. (doi:10.1038/nature11983)

21.   Mannion PD, Upchurch P, Benson RBJ, Goswami A. 2014 The latitudinal biodiversity

gradient through deep time. *Trends Ecol. Evol.* **29**, 42–50.

(doi:10.1016/j.tree.2013.09.012)

22.   Holderegger R, Buehler D, Gugerli F, Manel S. 2010 Landscape genetics of plants.

*Trends Plant Sci.* **15**, 675–683. (doi:10.1016/j.tplants.2010.09.002)

23.   Savolainen O, Pyhäjärvi T, Knürr T. 2007 Gene flow and local adaptation in trees. *Annu.*

*Rev. Ecol. Evol. Syst.* **38**, 595–619. (doi:10.1146/annurev.ecolsys.38.091206.095646)

24.   Tomlinson PB. 1986 The Botany of Mangroves. New York: Cambridge University Press

25.    Giri C, Ochieng E, Tieszen LL, Zhu Z, Singh A, Loveland T, Masek J, Duke N. 2011

449           Status and distribution of mangrove forests of the world using earth observation satellite

450           data. *Glob. Ecol. Biogeogr.* **20**, 154–159. (doi:10.1111/j.1466-8238.2010.00584.x)

451    26.    Van der Stocken T, Wee AKS, De Ryck DJR, Vanschoenwinkel B, Friess DA, Dahdouh-

452           Guebas F, Simard M, Koedam N, Webb EL. 2019 A general framework for propagule

453           dispersal in mangroves. *Biol. Rev.* **94**, 1547–1575. (doi:10.1111/brv.12514)

454    27.    Middleton BA. 2012 *Global change and the function and distribution of wetlands*.

455           Dordrecht, Netherlands: Springer.

456    28.    Morrisey DJ, Swales A, Dittmann S, Morrison MA, Lovelock CE, Beard CM. 2010 The

457           ecology and management of temperate mangroves. In: Gibson RN, Atkinson RJA,

458           Gordon JDM, editors. *Oceanography and marine biology: an annual review*. Boca

459           Raton, FL: Chapman and Hall/CRC, pp. 43–160.

460    29.    Spalding M, Blasco F, Field CD. 1997 *World mangrove Atlas*. Okinawa, Japan: The

461           International Society for Mangroves Ecosystems.

462    30.    Cavanaugh KC, Osland MJ, Bardou R, Hinojosa-Arango G, López-Vivas JM, Parker JD,

463           Rovai AS. 2018 Sensitivity of mangrove range limits to climate variability. *Glob. Ecol.*

464           *Biogeogr.* **27**, 925–935. (doi:10.1111/geb.12751)

465    31.    Osland MJ *et al.* 2017 Climatic controls on the global distribution, abundance, and

466           species richness of mangrove forests. *Ecol. Monogr.* **87**, 341–359.

467           (doi:10.1002/ecm.1248)

468    32.    Soares MLG, Estrada GCD, Fernandez V, Tognella MMP. 2012 Southern limit of the

469           Western South Atlantic Mangroves: assessment of the potential effects of global warming

from a biogeographical perspective. *Estuar. Coast. Shelf Sci.* **101**, 44–53.

(doi:10.1016/j.ecss.2012.02.018)

33.  Duke NC. 1991 A systematic revision of the mangrove genus Avicennia (Avicenniaceae)

in Australasia. *Aust. Syst. Bot.* **4**, 299. (doi:10.1071/sb9910299)

34.  Li X, Duke NC, Yang Y, Huang L, Zhu Y, Zhang Z, Zhou R, Zhong C, Huang Y, Shi S.

2016 Re-evaluation of phylogenetic relationships among species of the Mangrove Genus

Avicennia from Indo-West Pacific based on multilocus analyses. *PLoS One* **11**,

e0164453. (doi:10.1371/journal.pone.0164453)

35.  Schaeffer-Novelli Y, Cintrón-Molero G, Adaime RR, de Camargo TM, Cintron-Molero

G, de Camargo TM. 1990 Variability of Mangrove ecosystems along the Brazilian Coast.

*Estuaries* **13**, 204–218. (doi:10.2307/1351590)

36.  Mori GM, Zucchi MI, Souza AP. 2015 Multiple-geographic-scale genetic structure of

two Mangrove tree species: the roles of mating system, hybridization, limited dispersal

and extrinsic factors. *PLoS One* **10**, e0118710. (doi:10.1371/journal.pone.0118710)

37.  Cruz MV, Mori GM, Signori-Müller C, da Silva CC, Oh DH, Dassanayake M, Zucchi

MI, Oliveira RS, de Souza AP. 2019 Local adaptation of a dominant coastal tree to

freshwater availability and solar radiation suggested by genomic and ecophysiological

approaches. *Sci. Rep.* **9**, 27. (doi:10.1038/s41598-019-56469-w)

38.  Cruz MV, Mori GM, Oh DH, Dassanayake M, Zucchi MI, Oliveira RS, Souza AP. 2020

Molecular responses to freshwater limitation in the mangrove tree *Avicennia germinans*

(Acanthaceae). *Mol. Ecol.* **29**, 344–362. (doi:10.1111/mec.15330)

39.   Francisco PM, Mori GM, Alves FM, Tambarussi EV, de Souza AP. 2018 Population

genetic structure, introgression, and hybridization in the genus Rhizophora along the

Brazilian coast. *Ecol. Evol.* **8**, 3491–3504. (doi:10.1002/ece3.3900)

40.   Pil MW, Boeger MRT, Muschner VC, Pie MR, Ostrensky A, Boeger WA. 2011

Postglacial north-south expansion of populations of *Rhizophora mangle*

(Rhizophoraceae) along the Brazilian coast revealed by microsatellite analysis. *Am. J.*

*Bot.* **98**, 1031–1039. (doi:10.3732/ajb.1000392)

41.   Takayama K, Tateishi Y, Murata JIN, Kajita T. 2008 Gene flow and population

subdivision in a pantropical plant with sea-drifted seeds *Hibiscus tiliaceus* and its allied

species: evidence from microsatellite analyses. *Mol. Ecol.* **17**, 2730–2742.

(doi:10.1111/j.1365-294x.2008.03799.x)

42.   Lumpkin R, Johnson GC. 2013 Global ocean surface velocities from drifters: mean,

variance, El Niño-Southern oscillation response, and seasonal cycle. *J. Geophys. Res.:*

*Oceans* **118**, 2992–3006. (doi:10.1002/jgrc.20210)

43.   Danecek P *et al.* 2011 The variant call format and VCF tools. *Bioinformatics* **27**, 2156–

2158. (doi:10.1093/bioinformatics/btr330)

44.   Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G. 2008 LOSITAN: a workbench

to detect molecular adaptation based on a F st -outlier method. *BMC Bioinform.* **9**, 323.

(doi:10.1186/1471-2105-9-323)

45.   Luu K, Bazin E, Blum MGB. 2017 pcadapt: an R package to perform genome scans for

selection based on principal component analysis. *Mol. Ecol. Resour.* **17**, 67–77.

(doi:10.1111/1755-0998.12592)

46.   Frichot E, François O. 2015 LEA: an R package for landscape and ecological association

studies. *Methods Ecol. Evol.* **6**, 925–929. (doi:10.1111/2041-210x.12382)

47.   Wright S. 1949 The genetical structure of populations. *Ann. Eugen.* **15**, 323–354.

(doi:10.1111/j.1469-1809.1949.tb02451.x)

48.   Goudet J. 2005 Hierfstat, a package for r to compute and test hierarchical F-statistics.

*Mol. Ecol. Notes* **5**, 184–186. (doi:10.1111/j.1471-8286.2004.00828.x)

49.   R Core Team. 2019 *A language and environment for statistical computing.* Available

from: https://www.r-project.org/.

50.   Van der Stocken T, Carroll D, Menemenlis D, Simard M, Koedam N. 2019 Global-scale

dispersal and connectivity in mangroves. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 915–922.

(doi:10.1073/pnas.1812470116)

51.   Fick SE, Hijmans RJ. 2017 World Clim 2: new 1-km spatial resolution climate surfaces

for global land areas. *Int. J. Climatol.* **37**, 4302–4315. (doi:10.1002/joc.5086)

52.   Sbrocco EJ, Barber PH. 2013 MARSPEC: ocean climate layers for marine spatial

ecology. *Ecology* **94**, 979. (doi:10.1890/12-1358.1)

53.   Klein T, Langner J, Frankenberg B, Svensson J, Broman B, Bennet C, Langborg T. 2013

ECDS - a Swedish research infrastructure for the open sharing of environment and

climate data. *Data Sci. J.* **12**, 1–9. (doi:10.2481/dsj.12-045)

54.   Leroy B, Meynard CN, Bellard C, Courchamp F. 2016 Virtualspecies, an R package to

generate virtual species distributions. *Ecography* **39**, 599–607. (doi:10.1111/ecog.01388)

55.   Hijmans RJ, van Etten J. 2012 *Raster: geographic analysis and modeling with raster*

*data.* R package version 2.0-12.

56.   Chapman AD. 2005 Principles and methods of data cleaning – primary species and

speciesoccurrence data, version 1.0. Copenhagen: Report for the Global Biodiversity

Information Facility.

57.   Lira-Noriega A, Manthey JD. 2014 Relationship of genetic diversity and niche centrality:

a survey and analysis. *Evolution* **68**, 1082–1093. (doi:10.1111/evo.12343)

58.   Li Y, Zhang XX, Mao RL, Yang J, Miao CY, Li Z, Qiu YX. 2017 Ten years of landscape

genomics: challenges and opportunities. *Front. Plant Sci.* **8**, 2136.

(doi:10.3389/fpls.2017.02136)

59.   Vieira MLC, Santini L, Diniz AL, Munhoz CF. 2016 Microsatellite markers: what they

mean and why they are so useful. *Genet. Mol. Biol.* **39**, 312–328. (doi:10.1590/1678-

4685-gmb-2016-0027)

60.   Morin PA, Luikart G, Wayne RK, the SNP Workshop Group. 2004 SNPs in ecology,

evolution and conservation. *Trends Ecol. Evol.* **19**, 208–216.

(doi:10.1016/j.tree.2004.01.009)

61.   Cerón-Souza I, Rivera-Ocasio E, Medina E, Jiménez JA, McMillan WO, Bermingham E.

2010 Hybridization and introgression in New World red Mangroves,

Rhizophora(Rhizophoraceae). *Am. J. Bot.* **97**, 945–957. (doi:10.3732/ajb.0900172)

62.   Kennedy JP, Pil MW, Proffitt CE, Boeger WA, Stanford AM, Devlin DJ. 2016

Postglacial expansion pathways of red mangrove, Rhizophora mangle, in the Caribbean

Basin and Florida. *Am. J. Bot.* **103**, 260–276. (doi:10.3732/ajb.1500183)

63.   Ochoa-Zavala M, Jaramillo-Correa JP, Piñero D, Nettel-Hernanz A, Núñez-Farfán J.

2019 Contrasting colonization patterns of black mangrove (*Avicennia germinans* (L.) L.)

gene pools along the Mexican coasts. *J. Biogeogr.* **46**, 884–898. (doi:10.1111/jbi.13536)

64.    Sandoval-Castro E, Dodd RS, Riosmena-Rodríguez R, Enríquez-Paredes LM, Tovilla-

559            Hernández C, López-Vivas JM, Aguilar-May B, Muñiz-Salazar R. 2014 Post-glacial

560            expansion and population genetic divergence of Mangrove Species *Avicennia germinans*

561            (L.) stearn and *Rhizophora mangle* L. along the Mexican Coast. *PLoS One* **9**, e93358.

562            (doi:10.1371/journal.pone.0093358)

563    65.    Nettel A, Dodd RS. 2007 Drifting propagules and receding swamps: genetic footprints of

564            mangrove recolonization and dispersal along tropical coasts. *Evolution* **61**, 958–971.

565            (doi:10.1111/j.1558-5646.2007.00070.x)

566    66.    Takayama K, Tamura M, Tateishi Y, Webb EL, Kajita T. 2013 Strong genetic structure

567            over the American continents and transoceanic dispersal in the Mangrove genus

568            Rhizophora (Rhizophoraceae) revealed by broad-scale nuclear and chloroplast DNA

569            analysis. *Am. J. Bot.* **100**, 1191–1201. (doi:10.3732/ajb.1200567)

570    67.    Barker JSF, Frydenberg J, Sarup P, Loeschcke V. 2011 Altitudinal and seasonal variation

571            in microsatellite allele frequencies of *Drosophila buzzatii*. *J. Evol. Biol.* **24**, 430–439.

572            (doi:10.1111/j.1420-9101.2010.02180.x)

573    68.    Byars SG, Parsons Y, Hoffmann AA. 2009 Effect of altitude on the genetic structure of

574            an Alpine grass, Poa hiemata. *Ann. Bot.* **103**, 885–899. (doi:10.1093/aob/mcp018)

575    69.    Jiang S, Luo MX, Gao RH, Zhang W, Yang YZ, Li YJ, Liao PC. 2019 Isolation-by-

576            environment as a driver of genetic differentiation among populations of the only broad-

577            leaved evergreen shrub *Ammopiptanthus mongolicus* in Asian temperate deserts. *Sci.*

578            *Rep.* **9**, 12008. (doi:10.1038/s41598-019-48472-y)

70. Manthey JD, Moyle RG. 2015 Isolation by environment in White-breasted Nuthatches

(Sitta carolinensis) of the Madrean Archipelago sky islands: a landscape genomics

approach. *Mol. Ecol.* **24**, 3628–3638. (doi:10.1111/mec.13258)

71. Mitchell-Olds T, Willis JH, Goldstein DB. 2007 Which evolutionary processes influence

natural genetic variation for phenotypic traits? *Nat. Rev. Genet.* **8**, 845–856.

(doi:10.1038/nrg2207)

72. Shafer ABA, Wolf JBW. 2013 Widespread evidence for incipient ecological speciation: a

meta-analysis of isolation-by-ecology. *Ecol. Lett.* **16**, 940–950. (doi:10.1111/ele.12120)

73. Rodríguez-Zárate CJ, Sandoval-Castillo J, van Sebille E, Keane RG, Rocha-Olivares A,

Urteaga J, Beheregaray LB. 2018 Isolation by environment in the highly mobile olive

ridley turtle ( Lepidochelys olivacea ) in the eastern Pacific. *Proc. R. Soc. B* **285**:

20180264. (doi:10.1098/rspb.2018.0264)

74. Beheregaray LB, Cooke GM, Chao NL, Landguth EL. 2015 Ecological speciation in the

tropics: insights from comparative genetic studies in Amazonia. *Front. Genet.* **5**:477.

(doi:10.3389/fgene.2014.00477)

75. Duke NC, Kovacs JM, Griffiths AD, Preece L, Hill DJE, van Oosterzee P, Mackenzie J,

Morning HS, Burrows D. 2017 Large-scale dieback of mangroves in Australia. *Mar.*

*Freshw. Res.* **68**, 1816. (doi:10.1071/mf16322)

76. McKee K, Rogers K, Saintilan N. 2012 Response of salt marsh and Mangrove wetlands

to changes in atmospheric $CO_2$, climate, and sea level. In: Middleton BA, editor. *Global*

*change and the function and distribution of wetlands*. Dordrecht, Netherlands: Springer,

pp. 63–96.

77.    Osland MJ, Enwright NM, Day RH, Gabler CA, Stagg CL, Grace JB. 2016 Beyond just

sea-level rise: considering macroclimatic drivers within coastal wetland vulnerability

assessments to climate change. *Glob. Change Biol.* **22**, 1–11. (doi:10.1111/gcb.13084)

78.    Ximenes A, Ponsoni L, Lira C, Koedam N, Dahdouh-Guebas F. 2018 Does sea surface

temperature contribute to determining range limits and expansion of Mangroves in

Eastern South America (Brazil)? *Remote Sens.* **10**, 1787. (doi:10.3390/rs10111787)

79.    IPCC. 2014 Climate change 2014: synthesis report. In: IPCC, editor. *Climate change*

*2014 - The physical science basis*. Cambridge, MA: Cambridge University Press, pp. 1–

30.

80.    Fraser DJ, Bernatchez L. 2001 Adaptive evolutionary conservation: towards a unified

concept for defining conservation units. *Mol. Ecol.***10**, 2741–52. (doi: 10.1046/j.0962-

1083.2001.01411.x)

 **TABLES**

615   **Table 1.** Collection sites for the samples of *Avicennia schaueriana* and *A. germinans* long the

616   South American coast.

| A. schaueriana | A. germinans | Locality (City/State) | Latitude | Longitude |
|---|---|---|---|---|
| | GM | Soure, Pará | -0.723 | -48.490 |
| | GB | Bragança, Pará | -0.904 | -46.687 |
| SB | | Bragança, Pará | -0.820 | -46.615 |
| SA | GA | Alcântara, Maranhão | -2.410 | -44.406 |
| | GN | Parnaíba, Piauí | -2.778 | -41.822 |
| SR | GR | Paracuru, Ceará | -3.413 | -39.056 |
| | GT | Tamandaré, Pernambuco | -8.526 | -35.013 |
| SV | | Vera Cruz, Bahia | -12.983 | -38.684 |
| SG | | Guapimirim, Rio de Janeiro | -22.701 | -43.007 |
| SU | | Ubatuba, São Paulo | -23.489 | -45.164 |
| SC | | Cananéia, São Paulo | -25.020 | -47.918 |
| SP | | Pontal do Paraná, Paraná | -25.575 | -48.352 |
| SF | | Florianópolis, Santa Catarina | -27.576 | -48.518 |
| SL | | Laguna, Santa Catarina | -28.445 | -48.840 |

617   Sample populations of *A. schaueriana* e *A. germinans* are indicated by two capital letters corresponding

618   to Figure 1.

619 **Table 2.** Results of the MMRR between the genetic distances based on the set of nonneutral

620 molecular markers and each environmental variable for *A. schaueriana* and *A. germinans*.

| Environmental variables | A. schaueriana | | | | A. germinans | | | |
|---|---|---|---|---|---|---|---|---|
| | β | P | R² | P | β | P | R² | P |
| Annual mean temperature (ºC) | 0.67 | 0.001 | 0.45 | **0.0009** | 0.88 | 0.02 | 0.81 | **0.01** |
| Temperature annual range (ºC) | 0.60 | 0.005 | 0.36 | **0.0004** | 0.02 | 0.91 | 0.04 | 0.63 |
| Isothermality (ºC) | 0.86 | 0.001 | 0.75 | **0.0005** | 0.72 | 0.06 | 0.50 | 0.09 |
| Min temperature of coldest month (ºC) | 0.69 | 0.001 | 0.49 | **0.002** | 0.86 | 0.08 | 0.75 | 0.07 |
| Max temperature of warmest month (ºC) | 0.05 | 0.63 | 0.003 | 0.63 | 0.88 | 0.02 | 0.80 | **0.03** |
| Mean Temperature of Coldest Quarter (ºC) | 0.72 | 0.001 | 0.53 | **0.002** | 0.90 | 0.08 | 0.82 | 0.07 |
| Precipitation seasonality (mm) | 0.75 | 0.01 | 0.56 | **0.0004** | 0.65 | 0.05 | 0.42 | **0.05** |
| Precipitation of warmest quarter (mm) | 0.56 | 0.006 | 0.31 | **0.008** | 0.95 | 0.05 | 0.91 | **0.04** |
| Precipitation of coldest quarter (mm) | 0.62 | 0.01 | 0.39 | **0.01** | -0.09 | 0.88 | 0.008 | 0.88 |
| Mean solar radiation (kJ m-² day-¹) | 0.66 | 0.001 | 0.44 | **0.0005** | 0.68 | 0.05 | 0.46 | **0.04** |
| Min solar radiation (kJ m-² day-¹) | 0.70 | 0.001 | 0.49 | **0.0001** | -0.26 | 0.37 | 0.06 | 0.36 |
| Max solar radiation (kJ m-² day-¹) | 0.48 | 0.005 | 0.23 | **0.006** | 0.98 | 0.01 | 0.96 | **0.01** |
| Wind speed mean (m s-¹) | 0.19 | 0.36 | 0.04 | 0.39 | 0.27 | 0.45 | 0.07 | 0.41 |
| Max vapor pressure deficit (kPa) | 0.46 | 0.02 | 0.22 | **0.01** | 0.45 | 0.22 | 0.20 | 0.24 |
| Mean annual SSS (psu) | 0.009 | 0.96 | 0.00008 | 0.96 | -0.17 | 0.67 | 0.03 | 0.68 |
| SSS of the saltiest month (psu) | 0.12 | 0.47 | 0.01 | 0.48 | -0.31 | 0.22 | 0.09 | 0.23 |
| Annual variance in SSS (psu) | 0.19 | 0.48 | 0.03 | 0.50 | -0.22 | 0.39 | 0.05 | 0.39 |
| Mean annual SST (ºC) | 0.68 | 0.01 | 0.46 | **0.001** | -0.08 | 0.82 | 0.007 | 0.81 |
| SST of the coldest month (ºC) | 0.64 | 0.002 | 0.42 | **0.001** | 0.08 | 0.78 | 0.006 | 0.76 |
| Annual range in SST (ºC) | 0.59 | 0.003 | 0.36 | **0.006** | 0.88 | 0.13 | 0.78 | 0.15 |
| Annual average cycle amplitude (cm) | 0.79 | 0.002 | 0.62 | **0.007** | 0.35 | 0.007 | 0.12 | **0.006** |
| Annual average duration of tidal cycles (h) | -0.32 | 0.1 | 0.1 | 0.12 | -0.18 | 0.73 | 0.03 | 0.74 |
| Annual number of cycles | -0.29 | 0.22 | 0.08 | 0.23 | -0.16 | 0.84 | 0.02 | 0.82 |

621 The results are presented by the overall fit of the model (*R²*), regression coefficient (β) and significance

622 (*P*). Models with significant values are presented in bold. SST = sea surface temperature. SSS = sea

623 surface salinity

 **FIGURE LEGENDS**

**Figure 1.** Geographic distribution and characterization of *A. schaueriana* and *A. germinans*

sampling sites along the South American coast. The blue area represents the geographical

distribution of *A. schaueriana* and its sympatric region with *A. germinans* (blue and yellow).

Black circles and triangles represent the sampling sites of the plant material used for genomic

genotyping of A. *schaueriana* and A. *germinans*, respectively [37,38]. Sampling locations,

indicated by two letters, are displayed according to Table 1. Arrows represent the direction of the

main ocean currents acting on the Brazilian coast. Arrow widths illustrate the mean current speed

[42].

**Figure 2.** Heatmaps describing interpopulation genetic differentiation based on the total set SNP

molecular markers estimated by pairwise $F_{ST}$ (Wright) for (A) *A. schaueriana* (black circles) and

(B) *A. germinans* (black triangles) collection sites. Sample codes are denoted as in Table 1.

**Figure 3:** Populations location of (A) *A. schaueriana* e (B) *A. germinans* in a bidimensional

projection of the principal component analysis used to calculate the environmental distance. The

acronyms for sampling locations, indicated by two letters, are the same as those in Table 1. The

color gradient represents the contribution of each environmental variable to PC1 and PC2. V1:

Annual mean temperature (ºC). V2: Temperature annual range (ºC). V3: Isothermality (ºC).V4:

Min temperature of coldest month (ºC). V5: Max temperature of warmest month (ºC). V6: Mean

Temperature of Coldest Quarter (ºC).V7: Precipitation seasonality (mm). V8: Precipitation of

warmest quarter (mm). V9: Precipitation of coldest quarter (mm). V10: Mean solar radiation (kJ

m$^{-2}$ day$^{-1}$). V11: Min solar radiation (kJ m$^{-2}$ day$^{-1}$). V12: Max solar radiation (kJ m$^{-2}$ day$^{-1}$).

V13: Wind speed mean (m s⁻¹). V14: Max vapor pressure deficit (kPa). V15: Mean annual SSS

(psu). V16: SSS of the saltiest month (psu). V17: Annual variance in SSS (psu). V18: Mean

annual SST (ºC). V19: SST of the coldest month (ºC). V20: Annual range in SST (ºC). V21:

Annual average cycle amplitude (cm). V22: Annual average duration of tidal cycles (h). V23:

Annual number of cycles.

**Figure 4.** Plot of linear regressions showing the correlations among genetic, geographic, and

environmental distances. The pairwise comparisons were performed between (A,E) geographic

and genetic distances, (B,F) environmental and genetic distances, (C,G) geographic and

environmental distances and between (D,H) environmental and genetic distances based on the

putative nonneutral single nucleotide polymorphisms (SNPs) for *A. schaueriana* and *A.*

*germinans*, respectively. The $R^2$ and *P*-values are included.

# *A. schaueriana*

**A** $R^2 = 0.81$ $P < 0.001$ — Genetic Distance vs Geographic Distance

**B** $R^2 = 0.54$ $P < 0.001$ — Genetic Distance vs Environmental Distance

**C** $R^2 = 0.82$ $P < 0.001$ — Environmental Distance vs Geographic Distance

**D** $R^2 = 0.51$ $P < 0.01$ — Genetic Distance (nonneutral SNPs) vs Environmental Distance

# *A. germinans*

**E** $R^2 = 0.61$ $P < 0.05$ — Genetic Distance vs Geographic Distance

**F** $R^2 = 0.67$ $P < 0.05$ — Genetic Distance vs Environmental Distance

**C** $R^2 = 0.74$ $P < 0.01$ — Environmental Distance vs Geographic Distance

**H** $R^2 = 0.62$ $P < 0.05$ — Genetic Distance (nonneutral SNPs) vs Environmental Distance