1

Does seed mass drive interspecies variation in the effect of management practices on

weed demography?

Kazakou Elena[1], Fried Guillaume[2], Cheptou Pierre-Olivier[3], Gimenez Olivier[3]

[1]Montpellier Supagro CEFE, CNRS, Université de Montpellier, Université Paul Valéry

Montpellier 3, EPHE, IRD, Montpellier, France

[2] Anses, Laboratoire de la Santé des Végétaux, Unité Entomologie et Plantes Invasives,

34988, Montferrier-sur-Lez, France

[3] CNRS, CEFE, Université de Montpellier, Université Paul Valéry Montpellier 3, EPHE,

IRD, Montpellier, France

Running title : *Seed mass drive weed demography according to soil management*

Abstract

Optimizing the effect of management practices on weed population dynamics is challenging

due to the difficulties in inferring demographic parameters in seed banks and their response to

disturbance. Here, we used a long-term plant survey between 2006 and 2012 in 46 French

vineyards and quantified the effects of management practices (tillage, mowing and herbicide)

on colonization, germination and seed survival of 30 weed species in relation to their seed

mass. To do so, we used a recent statistical approach to reliably estimate demographic

parameters for plant populations with a seed bank using time series of presence–absence data,

which we extended to account for interspecies variation in the effects of management

practices on demographic parameters. Our main finding was independent of the nature of

farming practices and the demographic process. High disturbance through tillage increased

survival in the seed bank of species with high seed mass. The application of herbicides,

considered as an intermediate disturbance, increased germination, survival and colonization

probabilities of species with high seed mass. Mowing, representing habitats more competitive

for light, increased the survival of species with high seed mass. Overall, the strong

relationships between the effects of management practices and seed mass provides an

indicator for predicting the dynamics of weed communities under disturbance.

Keywords: Bayesian hidden Markov models, plant demography, seed bank, seed size, soil

management practices

Introduction

Managing weeds is one of the most challenging issues faced by farmers as they can cause

significant reductions in crop growth and yields especially in resource limited agroecosystems

(1). In vineyards, conventional weed control methods rely on different combinations of

herbicide applications, soil tillage and/or mowing applied in the vine rows and the area

between them (inter-rows) (2). However, environmental concerns have brought attention to

alternative weed management strategies, including the use of cover crop to improve soil

organic matter content and aggregation of nutrients and water and also to control weeds (3,4).

To optimize weed management, we need to not only quantify weed species presence and

abundance but also to understand species demography to determine the response of weed

population dynamics to disturbances.

   With regard to demography, buried seeds are an essential component of the

community (5) but their contribution to population dynamics is challenging to study because

the seed bank is not visible. A hidden Markov model (HMM) was recently developed to

estimate colonization, germination and seed bank survival using presence-absence

observations (6). Here, we extend this HMM with a multilevel model in a Bayesian

framework (e.g. (7)) to determine interspecies variation in the effects of management

practices on weed demography at both the plot and species level, while explicitly

considering the seed bank.

   Relatively undisturbed plant communities such as pasture generally exhibit low seed

persistence, whereas seed longevity in frequently disturbed habitats such as arable fields will

be high (8,9). Germination in most of the cases increase after soil disturbances such as tillage

and decrease after mowing (10–12). According to the disturbance intensity and frequency,

germination probability can be higher in tillage systems with a frequent disturbance than in

mowing systems. Additionnally, colonization can be more pronounced in frequently disturbed

environments, as tillage procedures increase the seed availability and may increase seedling

rates (13). From a mechanistic perspective we expect that seed mass, because of its influence

on key processes such as dispersal in space via colonization and dispersal in time via seed

bank persistence (14), will affect the influence of management practices on plant

demographic parameters. Small seeds of wind-dispersed plants should disperse at greater

distances on average than large seeds (15), whereas seedling of larger-seeded species should

better survive hazards like deep shade, physical damage and the presence of competing

vegetation (16).

To explore these hypotheses, we used a unique data set covering 46 vineyard plots in

France (Champagne, Beaujolais and Languedoc wine-growing areas) with 883 flora surveys

performed between 2006 and 2012. First, we used our novel multilevel HMM model to test

the effects of environmental variables on germination, colonization and seed persistence in

the seed bank. Second, we tested whether interspecific variation in the effects of management

practices (tillage, mowing and herbicide use) on demographic parameters could be explained

by seed mass.

Material and methods

(a) The Biovigilance dataset

Vegetation surveys were conducted in French vineyards between 2006 and 2012 in the

"Biovigilance" project (17), covering three main wine production regions – Languedoc,

Beaujolais and northern Rhône valley and Champagne – along a latitudinal gradient of pedo-

climatic conditions and management practices (for a detailed description see (18)). Languedoc

has a Mediterranean climate with a mean annual temperature of 14.1 °C, and 686 mm annual

rainfall in the surveyed plots (19) with a mean Treatment Frequency Index (TFI) for

herbicides of 0.48, i.e. the cumulative ratio of the dose applied to the recommended dose, for

all herbicide treatments applied during the growing season,. Beaujolais and northern Rhone

valley have a semi-continental climate with temperate influences, with a mean annual

temperature of 11.4 °C and 776 mm annual rainfall with a mean TFI of 1.38. Finally,

Champagne has a continental climate with oceanic influences, with a mean annual

temperature of 10.1 °C and 657 mm annual rainfall with a mean TFI of 1.24.

Forty-six vineyards plots were surveyed: 18 plots in Languedoc, 18 plots in Beaujolais

and northern Rhone valley and 10 plots in Champagne. In each of the 46 vineyards plot,

2,000m² quadrat surveys were performed along rows and inter-rows to account for different

management practices. Following (18), we focused our analyses on the 30 most abundant

species (Table 1). We extracted seed mass values from the LEDA and TRY databases (20,

21).

(b) Statistical analyses

Following (6), we built a HMM by considering three states: '1' for both seeds and standing

flora are absent in year t, '2' for seeds are present in year t but standing flora is absent in year

$t+1$ and '3' for seeds are present in year t and standing flora is present in year $t+1$, underlying

the two observations made when collecting data: (i) species not seen, (ii) species seen. The

HMM included four different parameters: the germination probability g (joint probability of

seed germination success and of plant survival to adulthood), the probability of seed bank

survival s from t to $t+1$, the probability of external colonization c (probability that at least one

seed from outside arrived on the plot and survived to the onset of the next season) and the

probability that there were seeds in the soil the year before the first observation of the existing

flora in the plot p_0. Parameters g, s and c were plot- and species-specific. First, we tested at

the plot level the effects on these demographic parameters of latitude and pedoclimatic

variables namely soil pH and soil texture with proportion of silt and clay that were shown to

be relevant in a previous study (18). Second, we assessed the effects of management practices

on the demographic parameters at the species level using species random effects on both

intercepts and slopes of these relationships, and an effect of seed mass on the slope of

management practices. Our multilevel HMM was fitted with a Bayesian approach (electronic

supplementary material). We concluded to the significance of an effect if the 95% credible

interval of the corresponding slope excluded zero. We also computed the proportion of

explained variance for multilevel models (22).

Results

Weed demography

Seed survival probability in the seed bank s varied from 0.14 (*Anisantha sterilis*) to 0.82

(*Muscari neglectum*), germination probability g varied from 0.13 (*Chenopodium album*) to

0.79 (*Diplotaxis erucoides*) and colonization probability c varied from 0 (*Mercurialis annua*)

to 0.86 (*Lamium amplexicaule*) (electronic supplementary material, Table S1). We found no

significant correlations between the different demographic parameters (p>0.05).

Effects of pedoclimatic factors on weed demography

We found no significant effect of soil parameters and latitude on colonization c and seed

survival probability s (figure 1a, figure 1c). Latitude had a significant effect on germination

probability g, with species in plots from higher latitude having a higher germination

probability (figure 1b). Silt had a positive (although non-significant) effect on germination

probability (figure 1b).

Influence of seed mass on the effect of soil management practices on weed demography

Overall, seed mass explained interspecies variation in the effects of management practices on

most of the demographic parameters (figure 2), with a positive correlation between seed mass

and the intensity of these effects. Importantly, we also found that this correlation varied

according to management practices.

The effect of the three management practices on colonization probability covaried

positively with species seed mass (figures 2a, 2b, 2c) with the strongest effet for herbicide

application ($R^2 = 0.86$). Herbicide application enhance more the rate of colonization for

species with large seed mass, like *Malva sylvestris*, *Calendula arvensis*, *Anisantha sterilis* or

Convolvulus arvensis than species with small seed mass like *Erigeron canadensis* and

Cerastium glomeratum (figure 2c).

With regard to germination, the effect of herbicide application on vineyards covaried

positively with seed mass: herbicide application increased more the germination probability

of species with large seed mass than species with small seed mass ($R^2 = 0.52$; figure 2f). The

same result was found for the effect of tillage on germination probability (except for species

Erodium cicutarium, Fumaria officinalis, Muscari neglectum or *Cerastium glomeratum*)

although the relationship was weak ($R^2 = 0.1$; figure 2d). We found no significant relationship

for the effect of mowing on germination probability (figure 2e).

Finally, the effects of the three management practices on survival probability were

better explained by seed mass than for germination and colonization. Seed survival in the seed

bank has been more increased after tillage for species with high seed mass (except from

Erodium cicutarium, Fumaria officinalis, Muscari neglectum, which also had a lower

increase in germination probability after the tillage) ($R^2 = 0.66$; figure 2h). We found the

same pattern for the effect of mowing ($R^2 = 0.66$; figure 2i) and herbicide ($R^2 = 0.96$; figure

2j) with species like *Malva sylvestris* or *Calendula arvensis* showing a higher increase in

survival in seed bank after mowing or herbicide application than *Erigeron canadensis,*

Cerastium glomeratum or *Cardamine hirsuta.*

Discussion

Effect of management practices to demographic parameters

Soil disturbance is one of the most important driver for vegetation (21, 22) and a cultivated

field can be assimilated to the first stages of secondary succession because it is regularly

disturbed. In our study, soil management practices corresponded to a gradient of disturbance

frequency and intensity. Frequent mechanical weed control is expected to reduce the

abundance of species with short-lived seeds and extend overall seed persistence (25). Mowing

directly defoliates plants and can reduce growth, decrease plant survival and reduce or prevent

seed production in two ways (26). First, mowing changes the biotic environment, such as

light, temperature and soil moisture. Second, mowing changes competitive relationships

between neighboring plants because different species die or regrow at different rates

following mowing (27). Repeated chemical control in vineyards can be considered as an

evolutionary force which results in communities secondary successions: each year the

chemical weed control restarts the succession, by the creation of gaps of bare soil but some

species escape and slowly give place to a successional process (28). The positive effects of

gaps in enhancing seedling recruitment are widely acknowledged (29).

Interestingly, our main finding was independent of the nature of farming practices and

the demographic process: when the level of disturbance increases (i.e., in plots with a higher

number of herbicide, tillage or mowing treatments), large seeded species see colonization,

germination and survival increase more quickly than small seeded species.

We found that for large seeded species, colonization, germination and survival in the

seed bank increase with the intensity of the disturbances. This means that the intensity of the

disturbances modifies the patterns of presence of these species in the seed bank and the

emerged flora. This finding can be explained by the fact that larger seeds confer an advantage

with higher seedling survival and germination probability under unfavourable environment

(29,30), at least on a relatively short term (31), and a greater success of emerging from burial

(32,33) although they disperse less due to their greater mass (34). We also found that tillage

increased the probability of the success of colonization from an external source and the

survival to the onset of the next season for species with high seed mass. One explanation is

that inversion operations, like tillage, bury weed seeds at a depth where the seeds are less

prone to predation and germination and thus persist longer (30). Additionnally, species with

large seeds are predicted to have reduced dormancy because their seedlings can draw on a

larger food reserve, and hence establish in relatively unfavorable environments. Our results

showed that herbicide application increased large seeded species germination probability and

survival in seed bank. These results are in accordance with our first hypothesis that

colonization should be more pronounced in frequently disturbed environments (15).

For very small seeded species, there is not much influence of the intensity of practices.

In fact, having small seeds is a typical characteristic of species with a ruderal strategy (37),

which are expected to be particularly adapted to disturbed environments and capable of

colonizing freshly disturbed environments (38). Our results show that for these species, the

demographic parameters remain constant regardless of the level of disturbance. Their

presence in the seed bank and in the emerged flora is therefore not modified by the intensity

of the practices.

Conclusion

Biodiversity conservation in vineyards is crucial not only for ensuring the sustainability and

stability of farming systems, but also for providing important ecological services. However, in

the last decades the vineyard has suffered an intensive management with a high

mechanisation (including frequent tilling) and/or use of herbicides, which affect species

richness and abundance (18). Here we used a novel statistical approach to determine the effect

of management practices on seed bank dynamics in weed species. The strong relationships

between these effects and seed mass provides a reliable indicator for predicting the dynamics

of weed communities.

Author's contributions

EK and OG conceived the study. GF oversaw data collection and compiled the dataset. OG

analysed the data. EK, OG, GF and P-OC interpreted results and designed model

comparisons. EK and OG framed and wrote the manuscriptwith input from GF and P-OC. All

authors approved the final versionand agree to be held accountable for the work it presents**.**

Data accessibility

The data and code used in this study are available as electronic supplementary material.

Acknowledgements

Funding

OG was funded by the French National Research Agency (grant ANR-16-CE02-0007).

References

1.   Hembree K, Lanini W. Weeds. UC IPM Pest Management Guidelines: Grape.

University of California; 2006.

2.   Steinmaus S, Elmore CL, smith RJ, Donaldson D, Weber EA, Roncoroni JA, et al.

Mulched cover crops as an alternative to conventional weed management systems in

vineyards. Weed Res. 2008 Jun 1;48(3):273–81.

3.   Guerra B, Steenwerth K. Influence of Floor Management Technique on Grapevine

Growth, Disease Pressure, and Juice and Wine Composition: A Review. Am J Enol Vitic.

2012 Jun 1;63(2):149.

4.   Sanguankeo PP, Leon RG, Malone J. Impact of Weed Management Practices on

Grapevine Growth and Yield Components. Weed Sci. 2009;57(1):103–7.

5.   Zobel M, Otsus M, Liira J, Moora M, Möls T. Is small-scale species richness limited

by seed availability or microsite availability? Ecology. 2000 Dec 1;81(12):3274–82.

6.   Pluntz M, Coz SL, Peyrard N, Pradel R, Choquet R, Cheptou P-O. A general method

for estimating seed dormancy and colonisation in annual plants from the observation of

existing flora. Ecol Lett. 2018 Sep 1;21(9):1311–8.

7.   Qian SS, Cuffney TF, Alameddine I, McMahon G, Reckhow KH. On the application

of multilevel modeling in environmental and ecological studies. Ecology. 2010;91(2):355–61.

8.   Bekker RM, Bakker JP, Grandin U, Kalamees R, Milberg P, Poschlod P, et al. Seed

size, shape and vertical distribution in the soil: indicators of seed longevity. Funct Ecol. 1998

Oct 1;12(5):834–42.

9.   Thompson K, Ceriani RM, Bakker JP, Bekker RM. Are seed dormancy and

persistence in soil related? Seed Sci Res. 2003;13(2):97–100.

10.   McConnaughay KDM, Bazzaz FA. The Relationship Between Gap Size and

Performance of Several Colonizing Annuals. Ecology. 1987 Apr 1;68(2):411–6.

11.   Reader RJ. Control of Seedling Emergence by Ground Cover and Seed Predation in

Relation to Seed Size for Some Old-Field Species. J Ecol. 1993;81(1):169–75.

12.   McIntyre S, Lavorel S, Tremont RM. Plant Life-History Attributes: Their Relationship

to Disturbance Response in Herbaceous Vegetation. J Ecol. 1995;83(1):31–44.

13.   Sheley RL, Jacobs JS, Svejcar TJ. Integrating disturbance and colonization during

rehabilitation of invasive weed-dominated grasslands. Weed Sci. 2005;53(3):307–14.

14.   Coomes DA, Grubb PJ. Colonization, tolerance, competition and seed-size variation

within functional groups. Trends Ecol Evol. 2003 Jun 1;18(6):283–91.

15.   Clark JS, Silman M, Kern R, Macklin E, HilleRisLambers J. Seed Dispersal near and

Far: Patterns across Temperate and Tropical Forests. Ecology. 1999;80(5):1475–94.

16.   Westoby M, Jurado E, Leishman M. Comparative evolutionary ecology of seed size.

Trends Ecol Evol. 1992 Nov 1;7(11):368–72.

17.   Fried G, Norton LR, Reboud X. Environmental and management factors determining

weed species composition and diversity in France. Agric Ecosyst Environ. 2008 Oct

1;128(1):68–76.

18.   Fried G, Cordeau S, Metay A, Kazakou E. Relative importance of environmental

factors and farming practices in shaping weed communities structure and composition in

French vineyards. Agric Ecosyst Environ. 2019 Apr 1;275:1–13.

19.   Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution

interpolated climate surfaces for global land areas. Int J Climatol. 2005 Dec 1;25(15):1965–

78.

20.   Kleyer M, Bekker RM, Knevel IC, Bakker JP, Thompson K, Sonnenschein M, et al.

The LEDA Traitbase: a database of life-history traits of the Northwest European flora. J Ecol.

2008 Nov 1;96(6):1266–74.

21.   Kattge J, DÍAZ S, LAVOREL S, PRENTICE IC, LEADLEY P, BÖNISCH G, et al.

TRY – a global database of plant traits. Glob Change Biol. 2011 Sep 1;17(9):2905–35.

22.   Gelman A, Rubin DB. Inference from Iterative Simulation Using Multiple Sequences.

Stat Sci. 1992;7(4):457–72.

23.   Haas H, Streibig J. Haas, H. and Streibig, J.C. (1982) . In: LeBaron H.M.and Gressel,

J. Eds., Herbicide Resistance in Plants,. In: Changing patterns of weed distribution as a result

of herbicide use and other agronomic factors. LeBaron H.M.and Gressel, J. New York, USA:

John Wiley & Sons; p. 57–9. (Herbicide Resistance in Plants).

24. Froud-Williams RJ, Chancellor RJ. A survey of weeds of oilseed rape in central

southern England. Weed Res. 1987 Jun 1;27(3):187–94.

25. Albrecht H. Seed traits in arable weed seed banks and their relationship to land-use

changes. Basic Appl Ecol. 2009 Sep;v. 10(6):516–24.

26. Ferraro DO, Oesterheld M. Effect of defoliation on grass growth. A quantitative

review. Oikos. 2002 Jul 1;98(1):125–33.

27. Vilà M, Terradas J. Neighbour effects on Erica multiflora (Ericaceae) reproductive

performance after clipping. Acta Oecologica. 1998 Mar 1;19(2):139–45.

28. Zanin G, Otto S, Riello L, Borin M. Ecological interpretation of weed flora dynamics

under different tillage systems. Agric Ecosyst Environ. 1997 Dec 15;66(3):177–88.

29. Turnbull LA, Rees M, Crawley MJ. Seed mass and the competition/colonization trade-

off: a sowing experiment. J Ecol. 1999 Oct 1;87(5):899–912.

30. Marshall DL. Effect of seed size on seedling success in three species of Sesbania

(Fabaceae). Am J Bot. 1986 Apr 1;73(4):457–64.

31. Moles A, Westoby M. Seedling survival and seed size: a synthesis of the literature. J

Ecol. 2004 Jun 1;92(3):372–83.

32. Gardarin A, Dürr C, Colbach N. Effects of seed depth and soil aggregates on the

emergence of weeds with contrasting seed traits. Weed Res. 2010 Apr 1;50(2):91–101.

33. Gardarin A, Dürr C, Mannino MR, Busset H, Colbach N. Seed mortality in the soil is

related to seed coat thickness. Seed Sci Res. 2010;20(4):243–56.

34. Fenner M. The phenology of growth and reproduction in plants. Perspect Plant Ecol

Evol Syst. 1998 Jan 1;1(1):78–91.

35. Mohler CL, Galford AE. Weed seedling emergence and seed survival: separating the

effects of seed position and soil modification by tillage. Weed Res. 1997 Jun 1;37(3):147–55.

36.   Omami, Haigh, Medd, Nicol. Changes in germinability, dormancy and viability of

Amaranthus retroflexus as affected by depth and duration of burial. Weed Res. 1999 Oct

1;39(5):345–54.

37.   Grime JP. Vegetation classification by reference to strategies. Nature. 1974 Jul

1;250(5461):26–31.

38.   Westoby M, Falster DS, Moles AT, Vesk PA, Wright IJ. Plant Ecological Strategies:

Some Leading Dimensions of Variation Between Species. Annu Rev Ecol Syst. 2002 Nov

1;33(1):125–59.

Figure 1. Effects of latitude and pedo-climatic variables (soil pH, proportions of silt and

clay) on the demographic parameters of 30 weed species in French vineyards.

Colonization (left panel), germination (middle panel) and survival (right panel) are

considered. The red circles are the posterior means and the thin grey lines are the 95%

credible intervals. The effect was considered significant when the corresponding credible

interval did not overlap 0 represented by the dashed vertical line.

Figure 2. Slopes of the response of demographic parameter to management practices for

30 weed species in French vineyards. The effect of tillage (left column), mowing (middle

column) and herbicides (right column) on colonization (upper row), germination (middle row)

and survival (bottom row) is given as a function of seed mass on the log scale (blue solid

line), as estimated by the multilevel hidden Markov model with species as a random effect,

holding latitude and pedo-climatic variables to their mean values. The red circles are the

posterior means and the thin grey lines are the 95% credible intervals. The proportion of

explained variance is also provided (bottom right corner in each panel).