1
2
3 **Propranolol promotes bone formation and limits resorption through novel mechanisms**
4 **during anabolic parathyroid hormone treatment in female C57BL/6J mice**
5
6

7 Annika Treyball, BA[1], Audrey C. Bergeron, MS[1], Daniel J. Brooks, MS[2], Audrie L. Langlais
8 BS[1,3], Hina Hashmi, BS[1], Kenichi Nagano, DDS, PhD[4], Deborah Barlow[5], Kathleen T. Nevola,
9 MS[1,6], Karen L. Houseknecht, PhD[5], Roland Baron, DDS, PhD[4], Mary L. Bouxsein, PhD[2,7],
10 Anyonya R. Guntur, PhD[1,3,8], Katherine J. Motyl, PhD[1,3,8]*
11
12
[1]Center for Molecular Medicine, Maine Medical Center Research Institute, Scarborough, ME
[2]Center for Advanced Orthopaedic Studies, Beth Israel Deaconess Medical Center, Boston, MA
[3]Graduate School of Biomedical Sciences and Engineering, University of Maine, Orono, ME
[4]Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine,
Boston, MA
[5]Department of Biomedical Sciences, University of New England, Biddeford, ME
[6]Tufts Graduate School of Biomedical Sciences, Tufts University, Boston, MA
[7]Department of Orthopedic Surgery, Harvard Medical School, Boston, MA
[8]Tufts University School of Medicine, Tufts University, Boston, MA
22
23
Current Affiliations
• Annika Treyball is presently affiliated with the Tufts University School of Medicine.
• Hina Hashmi is presently affiliated with the College of Osteopathic Medicine, University
of New England.
• Kenichi Nagano is presently affiliated with the Department of Oral Pathology and Bone
Metabolism, Institute of Biomedical Sciences, Nagasaki University.
30
Address correspondence to:
Katherine J. Motyl
81 Research Drive
Scarborough, ME 04074
(207) 396-8004
motylk@mmc.org

37
38 *Conflict of Interest Statement:*
39 The authors have declared that no conflict of interest exists.
40
41

Abstract

Although the non-selective β-blocker, propranolol, improves bone density with PTH treatment in mice, the mechanism of this effect is unclear. To address this, we used a combination of *in vitro* and *in vivo* approaches to address how propranolol influences bone remodeling in the context of PTH treatment. In female C57BL/6J mice, intermittent PTH and propranolol had complementary effects in the trabecular bone of the distal femur and L5 vertebra, with combination treatment achieving micro-architectural parameters beyond that of PTH alone. Combined treatment improved the serum bone formation marker, P1NP, but did not impact other histomorphometric parameters relating to osteoblast function at the L5. *In vitro*, propranolol amplified the acute, PTH-induced, intracellular calcium signal in osteoblast-like cells. The most striking finding, however, was suppression of PTH-induced bone resorption. Despite this, PTH-induced receptor activator of nuclear factor kappa-B ligand (RANKL) mRNA and protein levels were unaltered by propranolol, which led us to hypothesize that propranolol could act directly on osteoclasts. Using *in situ* methods, we found *Adrb2* expression in osteoclasts *in vivo*, suggesting β-blockers may directly impact osteoclasts. Taken together, this work suggests a strong anti-osteoclastic effect of non-selective β-blockers *in vivo,* indicating that combining propranolol with PTH could be beneficial to patients with extremely low bone density.

Introduction

The sympathetic nervous system (SNS) plays a critical role in the regulation of bone remodeling. The SNS suppresses bone formation and promotes receptor activator of nuclear factor kappa-B ligand (RANKL)-mediated osteoclast recruitment, resulting in low trabecular bone volume fraction in mice [1,2]. Consistent with this, tyrosine hydroxylase, one of the enzymes involved in norepinephrine production, is present in nerves within bone, marrow, and periosteum [3]. The major adrenergic receptor mediating the downstream effects of norepinephrine appears to be the β2AR in mice [1]. Deletion of β2AR, or treatment with the non-selective βAR antagonist (β-blocker) propranolol, prevents bone loss in situations of stress, antipsychotic and antidepressant treatment, and in other situations in which sympathetic signaling to bone may be elevated [1,4–8]. The majority of studies point toward a mechanism of β2AR suppressing bone formation directly in the osteoblast, and activating bone resorption indirectly, through the RANKL/OPG pathway [1,2,9]. Indeed, mice with an osteoblast-specific deletion of *Adrb2* have a high bone density phenotype as adults, with increased bone formation and reduced *Rankl* expression [10]. However, some studies have found evidence for direct effects of β2AR signaling in osteoclasts [11,12], suggesting the downstream effects of elevated SNS activity may be more complex than generally accepted.

Despite this complexity, identifying osteoporosis treatment strategies that modulate sympathetic signaling remains clinically useful. β-blockers are one such group of drugs that have been shown to reduce fracture risk and increase BMD in patients [13][14]. Furthermore, rodent studies have shown that combining β-blockers with teriparatide (intermittent truncated parathyroid hormone, PTH) may further promote bone density. Evidence from ovariectomized mice suggests that combining PTH treatment with propranolol increases bone mineral density

beyond the levels achieved by PTH alone, and histomorphometric analyses indicated that

improvement was largely due to increased bone formation and osteoblast number [15].

On a cellular level, evidence from *in vitro* and *in vivo* studies suggest that PTH efficacy is

dependent upon the presence of the β2AR. Deletion of *Adrb2,* the gene that encodes β2AR,

prevents the anabolic effect of intermittent PTH in young and aged mice, suggesting some β2AR

signaling is required [16]. One mechanism of this may be through β2AR signaling allowing G-

protein βγ (Gβγ) subunit to bind endosomal PTH1R, which sustains cAMP levels leading to

enhanced mineralization in osteoblasts [17]. cAMP signaling, however, is not the only avenue for

PTH effects to be transduced to the cell [18][19]. PTH1R also signals through phospholipase C to

increase Ca^{2+} release from intracellular stores, as well as through arrestin-mediated mechanisms

[18][19]. There is an established role for intracellular Ca^{2+} signaling in the regulation of osteoblast

differentiation through the calmodulin/CamKII pathway regulating AP-1 and CREB/ATF4

transcription factors [20]. It has been shown that silencing of *Adrb2* can increase pCREB [21], but it

remains unknown whether the combination of PTH treatment with β2AR antagonists would

impact these pathways.

To further investigate the cellular and systemic effects of co-modulation of PTH1R and

β2AR, we performed a series of *in vitro* and *in vivo* assays pharmacologically targeting these

receptors. A pharmacological approach was chosen so that any advantages of combination

treatments may eventually be translated to humans. Briefly, we determined that the combination

of PTH and propranolol *in vivo* increased bone volume fraction in the distal femur in part by

increasing markers of bone formation, which may be related to the β-blocker propranolol

enhancing PTH-induced Ca^{2+} signaling in osteoblast-like cells *in vitro*. More striking, however,

was that propranolol prevented PTH-induced bone resorption, but did not impact PTH-induced

changes in RANKL or OPG pathway members. Rather, we identified that this effect may be

through direct anti-osteoclastogenic effects of propranolol since *Adrb2* is expressed in

osteoclasts *in vivo*. In all, these findings suggest that modifications to PTH therapy that mimic

outcomes from propranolol treatment, or simply combining PTH with β-blockers, may be a

useful approach to minimizing resorption and promoting net bone accrual.

Results

Propranolol improved total BMD in the presence of PTH

To test whether propranolol would improve PTH-induced bone formation, we treated mice with

either vehicle, PTH, propranolol, or PTH + propranolol from 16-20 weeks of age. Neither body

weight, fat mass, nor fat-free mass was altered in any of the treatment groups, suggesting that

any bone changes would be independent of differences in loading (Table I). As expected, PTH

had a significant main effect on total aBMD and aBMC, as well as on femoral aBMD (Table I).

Although propranolol only had a significant main effect on total aBMD, all bone parameters

measured by DXA were highest in the PTH + propranolol group (Table I). Interestingly, co-

treatment with PTH and propranolol increased total aBMD beyond that caused by PTH or

propranolol alone.

PTH and propranolol had site-specific effects on trabecular bone microarchitecture

To investigate the impact of PTH and propranolol treatment on bone microarchitecture, we

performed μCT on L5 vertebrae and femurs. Propranolol and PTH both independently improved

trabecular BV/TV of the L5 vertebra, but the combination of PTH and propranolol improved

BV/TV and BMD above and beyond that of either treatment alone (Figure 1). This is likely due

to a combined effect of increased trabecular thickness and increased trabecular number, the latter

of which was only significantly increased by PTH when propranolol was present. Consistent

with this, PTH and propranolol together significantly reduced BS/BV ratio compared to

propranolol alone (Figure 1).

In the distal femur, we focused our examination of PTH and propranolol effects on two

distinct sites, the primary and secondary spongiosa (Figure 2). In the secondary spongiosa, PTH

increased Tb. BV/TV, BMD, and Conn.D and these parameters were all increased further by

combined treatment with propranolol (Figure 2B-D). Although some parameters (Tb.N, Tb.Th,

Tb.Sp, and BS/BV) were not altered by PTH alone, the combination reduced SMI, Tb.Sp, and

BS/BV, while significantly increasing Tb.N and Tb.Th (Figure 2E-I). The primary spongiosa

also had striking changes from PTH, including increased BV/TV, BV and BMD (Figure 2J-L),

but these were not exacerbated or diminished by combination PTH and propranolol treatment.

Tissue mineral density (TMD), a measurement of the density of the bone itself (not including any

marrow), was suppressed with PTH treatment, suggesting that mineral deposition may be

compromised (Figure 2M). Interestingly, however, propranolol significantly elevated TMD in

PTH-treated mice such that it was not different from vehicle-treated (Figure 2M). This indicates

the quality of mineralization in the primary spongiosa during combination PTH and propranolol

treatment may be improved compared to PTH alone.

Cortical bone microarchitecture

PTH significantly increased cortical area (Ct.Ar) and polar moment of inertia (pMOI) in the

femur midshaft, but did not significantly impact marrow area (Ma.Ar), total area (Tt.Ar), cortical

thickness (Ct.Th), tissue mineral density (TMD) or porosity (Figure 3 A-H). Although

propranolol treatment did not impact these parameters on its own, propranolol significantly

improved the effect of PTH on the cortical bone such that Ct.Ar, Ct.Ar/Tt.Ar, Ct.Th and TMD

were elevated in the PTH + propranolol group compared to propranolol alone (Figure 3A, D, E,

and F). The increased Ct.Ar/Tt.Ar with combination treatment is most likely due to a reduction

in Ma.Ar, because Tt.Ar was clearly unchanged (Figure 3B-D).

Propranolol promoted bone formation in PTH-treated mice.

To determine whether co-treatment with propranolol modulated the effect of PTH on bone

formation, we performed serum, histomorphometric, and mRNA analyses to evaluate bone

remodeling activity. As expected, PTH increased the serum marker of bone formation, P1NP,

and this was further increased by co-treatment with PTH and propranolol (Figure 4A).

Histomorphometric analyses in the L5 vertebrae indicated a significant effect of PTH by 2-way

ANOVA on MAR and BFR parameters (Figure 4B, C). Similarly, osteoblast parameters were

elevated by PTH, but were not significantly modulated by the combination with propranolol.

Propranolol did not have a significant main effect on these parameters (except to decrease the

N.Ob/B.Pm) and, when combined with PTH, did not increase these indices beyond the level of

PTH alone (Figure 4B, C). Propranolol did, however, reduce the amount of OS/BS in PTH-

treated mice (Figure 4C).

In general, these findings contrast the serum P1NP finding, suggesting other sites may

have elevated bone formation with the combination treatment. Therefore, we examined mRNA

markers of osteoblastogenesis and osteocyte maturation in a separate site, the tibia. Although we

did not see significant differences in *Runx2*, *Col1a1*, or *Osteocalcin* expression (Figure 4D-F),

we did observe differences in late osteoblast and osteocyte markers. First, dentin matrix acidic

phosphoprotein 1 (*Dmp1*) expression was increased with PTH treatment, but there was also a

significant interaction (p=0.01) such that PTH-induced *Dmp1* expression was higher in the

propranolol treated group, while propranolol alone had no effect in the absence of PTH (Figure

4G). Surprisingly, intermittent treatment with PTH induced *Sost* expression, in contrast to

previously reported effects with chronic elevation, and *Sost* was elevated to a greater degree in

propranolol treated mice (Figure 4H) [22]. Although serum P1NP is elevated, changes in bone

formation (histological and mRNA-based) do not convincingly explain the improved BV/TV in

mice treated with PTH and propranolol.

Recent studies have suggested PTH may suppress bone marrow adipose tissue (BMAT)

as a means or consequence of promoting bone formation. Although marrow adipocyte size was

not influenced by PTH, there was a subtle but significant interaction effect ($p < 0.05$) of PTH and

propranolol in both adipocyte volume / total volume (AV/TV) and adipocyte number parameters.

In particular, PTH tended to reduce marrow adipocyte numbers in mice not treated with

propranolol ($p = 0.08$) but this effect was blunted in propranolol-treated mice (Figure S1).

PTH-induced intracellular calcium signaling is enhanced by propranolol

Next, we sought to test the impact of propranolol and PTH directly in osteoblasts by

measuring the concentration-dependence of the intracellular Ca^{2+} response to PTH in MC3T3-E1

cells differentiated for 7 days (Figure 5). PTH induced a measurable Ca^{2+} signal at

concentrations as low as 10 nM and the peak response was observed at 1 µM. Using a log

transformation, we calculated the concentration with 50% of maximal excitation to be 66.6 nM

(Figure 5A), consistent with previously published reports (published iCa^{2+} EC_{50} = 50 nM with

dynamic range 5-500 nM)[23]. Therefore, we used a concentration of 100 nM in our studies, to be

within the dynamic range of the concentration-response curve, but also within the concentrations

used in previous publications demonstrating functional effects of PTH on osteoblasts[24]. Next,

we found that pre-treatment with propranolol significantly increased the intracellular Ca^{2+} signal

induced by PTH (Figure 5B, C). This is in contrast to treatment with propranolol alone, which

did not induce any intracellular Ca^{2+} signal (Figure 5B).

Long-term PTH treatment does not promote mineralization *in vitro*, so to get a sense

whether combined PTH and propranolol treatment had any functional consequence in

osteoblasts, we examined gene expression of relevant receptors (Figure S2) and osteoblast

markers after acute treatment with PTH, propranolol or both (Figure 5D-G). Although there were

no differences in early osteoblastogenic markers *Runx2* (Figure 5D) and *Sp7* (not shown), PTH

induced *Osteocalcin* expression (main effect of PTH p=0.016) similarly in both vehicle and

propranolol treated cells but only the group treated with PTH alone (no propranolol present)

reached pairwise statistical significance (Figure 5E). Because intracellular Ca^{2+} has been shown

to modulate AP-1 family transcription factors, we examined gene expression of *Fos*.

Interestingly, the PTH-induced increase in *Fos* was higher when propranolol was present (Figure

5F), consistent with propranolol promoting PTH-induced intracellular Ca^{2+} levels. One

established mechanism of osteoanabolism from PTH is through increased expression and

autocrine signaling of IGF-1[24]. We found a significant increase in *Igf1* expression only in the

PTH and propranolol co-treated group (Figure 5G). These findings suggest propranolol may

improve bone formation from PTH in part through increasing intracellular Ca^{2+} signaling and

promoting expression of *Igf1*.

PTH-induced osteoclast activity was blocked by propranolol

Since PTH is known to enhance resorption, we next investigated whether propranolol mitigates

resorption measurements *in vivo*. PTH increased expression of *Ctsk* and *Trap5*, as well as the

serum protein levels of the resorption marker CTx (Figure 6A-C). Interestingly, propranolol

prevented the PTH-induced increases in *Trap5* expression and serum CTx levels, suggesting

propranolol prevents PTH-induced resorption (Figure 6B, C). Consistent with the literature,

histomorphometry indicated that PTH increased the total number of osteoclasts (N.Oc/T.Ar,

Figure 6D). Propranolol alone did not have any impact on the number of osteoclasts or the other

osteoclast parameters (Figure 6D). When combined, however, the number of osteoclasts on the

bone surface and within the total area measured (N.Oc/B.Pm and N.Oc/T.Ar, respectively) and

the osteoclast surface fraction (Oc.S/BS) was lower in mice treated with PTH and propranolol

compared to those treated with PTH alone (Figure 6D), interaction $p \leq 0.02$). Notably, the PTH-

induced increase in N.Oc/T.Ar was completely blocked by co-treatment with propranolol

(*p*<0.01, Figure 6D). All three lines of evidence related to osteoclasts (serum, mRNA, and

histomorphometry) point toward a strong effect of propranolol to reduce PTH-induced bone

resorption.

Propranolol can centrally suppress sympathetic activity, therefore we measured

norepinephrine (NE) levels in serum as well. While neither PTH nor propranolol independently

modulated NE, the combination treatment resulted in significantly lower circulating NE

compared to PTH alone (Figure 6E), suggesting a systemic lowering of sympathetic activity by

propranolol might be responsible for the suppressed resorption. Because the current

understanding of the mechanism of NE signaling-induced osteoclastogenesis is through the

modulation of *receptor activator of nuclear factor kappa-B ligand* (*Rankl*) expression in the

osteoblast, we measured circulating RANKL and found that, to our surprise, propranolol had no

effect on PTH-induced RANKL levels (Figure 6F). Furthermore, although PTH induced *Rankl*

gene expression in cortical bone as expected, propranolol did not have any blunting effect on

Rankl (Figure 6G), but did lower *Opg* (Figure 6H), suggesting other mechanisms were

responsible for the reduced osteoclast numbers and CTx levels in PTH and propranolol treated

mice.

Previous *in vitro* work has suggested βARs may play a direct role in osteoclasts [11,12], but

this has not been confirmed *in vivo*. To test whether β2AR may be directly responsible for the

anti-osteoclastic effects of propranolol, we first tested marrow and serum concentrations of

propranolol achieved after dosing mice in the same manner. Propranolol concentrations achieved

in the serum of mice dosed in drinking water were 14.6 nM, while bone marrow concentrations

were 107.7 nM (Figure 7A). Within each mouse, approximately ten times more propranolol was

present in marrow compared to serum, suggesting osteoclasts are likely to have significant

propranolol exposure. We next examined whether *Adrb2* expression was present in osteoclasts *in*

vivo using *in situ* hybridization. First, we used probes specific to *Acp5*, the gene encoding

TRAP5, to identify multinucleated osteoclasts (Figure 7B). Next, we co-labeled with probes

specific to *Adrb2* and found levels of *Adrb2* expression in osteoclasts comparable to that of other

cells in the marrow (Figure 7C). Collectively, these findings suggest that lowered NE levels

combined with direct suppression of osteoclasts by propranolol may explain the reduced

resorption and improved micro-architectural parameters in mice co-treated with PTH and

propranolol.

Discussion

Here we show that propranolol and intermittent PTH co-treatment improves bone microarchitecture beyond the levels achieved by intermittent PTH treatment alone, which is consistent with previous co-treatment studies [15]. We add to this, however, the novel findings that propranolol enhances tissue mineral density, which may be through promoting Ca^{2+} signaling in response to PTH in osteoblasts. We are also the first to show that propranolol prevents PTH-induced resorption and that this may be through direct inhibition of osteoclast activity, since *Adrb2* is expressed in osteoclasts *in vivo*. This finding in particular suggests more detailed studies examining the roles of βARs in osteoclasts, as well as the impact of βAR-targeting drugs, are critically needed. In the absence of β-blockers, intermittent PTH treatment promotes both bone formation *and* resorption, which could be of concern to individuals with severely lowered bone density and/or high levels of resorption. Intracortical remodeling, which may lead to fracture, has been associated with serum PTH levels in humans, as well as with daily intermittent PTH treatment in rabbits [25,26]. Nonetheless, PTH is very effective at building new bone, so bone loss that occurs from heightened resorption needs to be weighed against its anabolic activity. Our findings suggest that combining PTH with propranolol, or potentially other β-blockers, may be an attractive option to prevent unwanted bone loss in patients with very low BMD.

The mechanism of propranolol action on bone to prevent resorption may be in part through direct effects of propranolol on osteoclasts. Neither PTH-induced serum RANKL nor *Rankl* gene expression was suppressed by propranolol, despite clear improvement in trabecular microarchitecture of the distal femur and the suppressed markers of resorption (serum CTx, *Acp5* expression, and N.Oc/T.Ar, Figure 6). The majority of publications point toward osteoblast

expression of RANKL as being the major mediator in the effects of the sympathetic nervous

system on osteoclasts [2,10]. However, our findings suggest that this is not the same mechanism

through which propranolol blocks resorption. In studies examining the osteoblast-mediated

mechanism of SNS-induced bone loss, some experiments suggested no effect of *Adrb2* in

osteoclasts. For example, Elefteriou et al. elegantly showed that the non-specific βAR agonist

isoproterenol only induces osteoclast differentiation in osteoblast and bone marrow macrophage

(BMM) co-cultures when wildtype, but not *Adrb2$^{-/-}$*, osteoblasts are present [2]. Similarly, the

effect of isoproterenol to induce osteoclast differentiation is not dependent upon the genotype of

the BMMs [2]. These data, in combination with findings from models with osteoblast-specific

deletions of *Adrb2*, have led to the generally accepted tenet that osteoblasts are the sole

mediators of the SNS effect on bone, including the mediators of osteoclast recruitment [10].

There is a relative scarcity of literature examining the effects of β-adrenergic signaling

directly in osteoclasts. However, Rodrigues, et al. showed that propranolol prevented

osteoclastogenesis of RAW 264.7 cells through suppression of NFATc1 protein expression, as

well as later markers of differentiation: *Acp5, Ctsk,* and *Mmp9* [12]. This suggests a role for βAR

signaling in controlling NFATc1, which is under the control of calcium-calmodulin and

calcineurin signaling. Kondo, et al. demonstrated that isoproterenol enhanced osteoclastogenesis

in bone marrow macrophages and RAW 264.7 cells by increasing reactive oxygen species [11].

Our findings appear more aligned with these reports because we see expression of *Adrb2* in

osteoclasts *in vivo* and propranolol-mediated suppression of resorption downstream of

osteoblast-induced RANKL. Although these findings are suggestive of a role for β2AR in

osteoclasts, additional studies are needed to establish this mechanism *in vivo* and *in vitro*, and to

test whether human osteoclasts would have similar properties.

Despite the paucity of literature relating to the effects of the SNS in osteoclasts directly,

there is extensive evidence that *Adrb2* is expressed in other cells of the myeloid lineage and that

the SNS plays a role in their function [27]. Indeed, the SNS is important for the innate immune

response required for tissue repair but can be pathologically activated during certain conditions,

such as heart failure. β-blockers are a common therapy for heart failure and work in part by

reducing inflammatory cytokine levels [27]. Systemic suppression of inflammation may also be an

interesting, yet unexplored, avenue to investigate the mechanism of efficacy of β-blockers in

improving BMD, especially with regards to resorption.

In contrast to our findings, global knockout of β2AR prevents any anabolic action of

PTH by preventing the PTH-induced increase in mineral apposition and osteoclast number [16].

Although these findings are paradoxical in some sense, they suggest that a basal level of β2-

adrenergic signaling in bone may be required for PTH action. It is also clear that pharmaceutical

inhibition of β2AR with propranolol is mechanistically distinct from the disruption of signaling

through genetic deletion. Propranolol can act as an inverse agonist, meaning it has effects

inhibitory toward cAMP signaling even when no agonist is present, reducing spontaneous β2AR

receptor activity[28]. Furthermore, propranolol also binds to the lesser expressed β1AR and is

known to have non-specific effects, including acting as a weak antagonist to some serotonin

receptors, which could, in turn, impact bone remodeling[29,30]. We cannot exclude the possibility

that propranolol is having non-specific effects in our studies, however, we did measure

propranolol exposure in serum and marrow and found serum levels comparable to those achieved

in patients (Figure 7A), suggesting clinical relevance[31]. The published binding affinity of

propranolol at the β2AR is 0.5 nM, significantly less than the 107.7 nM achieved in bone

marrow (Figure 7A), thus we are clearly inhibiting β2AR with propranolol *in vivo*[32].

Interestingly, the binding affinity of NE is 300 nM, while we were measuring levels in the 1-3

nM range in serum (Figure 6E). Collectively, these outcomes suggest that our observations are a

result of direct effects of propranolol on β2AR on osteoblasts and osteoclasts, either through

competitive inhibition of NE binding, or through inverse agonist properties (of which, IC_{50}

values are in the range of the binding affinity)[28].

Moriya et al. showed that *Adrb2* knockdown enhances PTH-induced phosphorylation of

CREB, suggesting a suppressive function of β2AR on PTH action [21]. This is consistent with our

findings that PTH effects in osteoblasts are enhanced with pharmacological β2AR inhibition.

Some intracellular scaffolding proteins, such as members of the Na^+/H^+ exchange regulatory

cofactor (NHERF) family, selectively promote coupling of the PTH1R to the $G\alpha_q$-PLC-β

signaling pathway [19]. NHERF also couples β2AR to the cytoskeleton, and agonist binding

causes uncoupling from the cytoskeleton [33]. It is unknown, however, whether propranolol might

uncouple β2AR from NHERF, G proteins and/or the other signaling components, which could

lead to the enhanced PTH-induced Ca^{2+} flux (Figure 5). Nonetheless, we observed enhanced *Igf1*

expression after co-treatment in osteoblasts, which has previously been shown to work in an

autocrine manner to stimulate aerobic glycolysis in osteoblasts in response to PTH-treatment at 6

hours [24]. In our hands, the elevation of *Igf1* after 1 hour of propranolol treatment suggests an

acceleration of this timetable, which may lead to the improved PTH effect on bone formation and

tissue mineral density that we observed *in vivo*. Further studies would be required to confirm this

mechanism.

TMD in the primary spongiosa was significantly higher with PTH and propranolol co-

treatment compared to PTH alone (Figure 2M). A similar trend was seen with cortical TMD

(Figure 3F) as well as total areal BMD assessed by DXA (Table I). Serum P1NP also indicated

global increases in bone formation with the combined treatment (Figure 4A), but this was not

observed in dynamic histomorphometric measurements of mineral apposition or bone formation

in the trabecular bone of the L5 vertebra (Figure 4C). However, gene expression in cortical bone

of the tibia suggested more mature osteoblasts/osteocytes (Figure 4G, H), indicating that

propranolol may have site-specific effects. Further studies investigating the mineral properties of

bone exposed to both PTH and propranolol would be necessary to determine if this combination

of treatments may improve strength in patients and animal models.

Clinical literature suggests that β-blockers prevent bone loss and reduce fracture risk in

humans. Yang et al. reported that β-blocker use was associated with a 17% decrease in risk of

fractures [13]. Toulis et al. analyzed 16 studies involving over 1.5 million subjects and found that

the risk of fracture was significantly reduced (by 15%) in subjects receiving β-blockers as

compared to control subjects, independent of sex, fracture site, or dose [14]. In contrast to what is

the apparent mechanism in rodent models, β1-selective blockers were found to be more effective

at reducing fractures than non-selective β-blockers (i.e. propranolol). More recently, when

evaluating adrenergic receptor mRNA expression in human bone biopsies, Khosla et al. found

that ADRB1 and ADRB2, but not ADRB3, were expressed in human bone [34]. Additionally,

patients treated with β1-selective blockers had better bone microarchitecture by quantitative

computed tomography (qCT) than non-users, had reduced CTx levels compared to placebo-

treated patients, and had increased BMD at the ultradistal radius, but these results were not seen

in propranolol (non-selective β-blockers) treated patients. Overall, this data suggests that β1-

selective blockers may be more protective against decreased BMD and increased risk of fracture

than non-selective β-blockers. Whether the perceived difference between rodents and humans is

due to β1AR being understudied in rodent models or due to an actual species-specific difference

in the relative importance of β1AR vs β2AR in bone remains unclear.

Our findings reinforce the work of others suggesting that β-blockers, which are routinely

used clinically, may be an effective treatment or co-treatment for osteoporosis. Importantly, here

we showed that propranolol prevented resorption without modulating PTH-induced RANKL,

suggesting that the current understanding of the mechanism of β-blocker and sympathetic

nervous system action on bone is incomplete. We have clear evidence of *Adrb2* expression in

osteoclasts *in vivo,* and future work should further investigate the extent to which osteoclasts

specifically mediate the impact of the SNS on bone. Investigating these mechanisms will be

necessary for the accurate interpretation of clinical and preclinical studies examining how the

sympathetic nervous system modulates bone homeostasis and contributes to bone pathologies.

Methods

Study Approval

The Institutional Animal Care and Use Committee of the Maine Medical Center Research

Institute approved all mouse protocols.

Mice and drug treatment

Sixteen-week-old C57BL/6J (Jackson Laboratories, Bar Harbor, ME) female, intact mice

underwent baseline body weight measurements and were randomly assigned to one of four

treatment groups. Mice were housed 3-4 per cage and provided standard laboratory chow *ad*

libitum. Female mice were chosen because of their higher propensity to have an anabolic

response to intermittent PTH. Mice were treated with either vehicle, PTH, propranolol or both

PTH and propranolol. PTH (H-1660, Bachem) was aliquoted and stored in glass vials, under

argon gas, at -70°C as a 10^{-4} M stock in 4 mM HCl supplemented with 0.1% bovine serum

albumin. PTH was thawed and diluted in 0.9% saline immediately prior to injection. Vehicle and

PTH (80 µg/kg) were administered by subcutaneous injection five days per week for four weeks.

Propranolol hydrochloride oral solution (West-Ward Pharmaceuticals) was dissolved in drinking

water at a concentration of 0.5 mg/mL and was delivered daily for four weeks to animals in the

propranolol and PTH + propranolol groups as described [4]. At 20 weeks-of-age, mice were

sacrificed one hour after treatment and tissues were either fixed in 10% buffered formalin or

snap-frozen in liquid nitrogen and stored at -70°C. A similar study was conducted with just 5

days of treatment to obtain additional samples for some mRNA and serum analyses and are

identified in the figure legends.

For the detection of propranolol levels in the plasma and bone marrow, sixteen-week-old

B6 female mice were administered propranolol dissolved in drinking water as described above.

Six mice were separated into three individual cages (two mice/cage). Mice were treated for four

days, and sacrificed in the morning, one hour after exchanging treated water for untreated water.

Serum was isolated from whole blood and bone marrow was collected from the tibia via

centrifugation. Samples were kept frozen until analysis was performed; propranolol levels were

detected using LC/MS-MS (see methods below).

Dual-energy X-ray absorptiometry (DXA)

Mice underwent DXA measurements at baseline and at 20 weeks of age for fat-free mass, fat

mass, bone mineral density, and bone mineral content using the PIXImus dual-energy X-ray

densitometer (GE-Lunar, Madison, WI). The instrument was calibrated daily with a mouse

phantom provided by the manufacturer. Mice were anesthetized with isoflurane and placed

ventral side down with each limb and tail positioned away from the body. Full body scans were

obtained, and X-ray energy absorptiometry data were gathered and processed with manufacturer-

supplied software (Lunar PIXImus 2, version 2.1). The head was specifically excluded from all

analyses due to concentrated mineral in skull and teeth.

Micro-computed tomography (µCT)

A high-resolution desktop micro-tomographic imaging system (µCT40, Scanco Medical AG,

Bruttisellen, Switzerland) was used to assess the microarchitecture of the fifth lumbar (L5)

vertebral body and the femur in accordance with published guidelines [35]. All analyses were

performed using the Scanco Evaluation software.

In the L5 vertebra, scans were acquired using a resolution of 10 μm^3, 70 kVp peak x-ray tube

potential, 114 mA x-ray tube current, 200 ms integration time, and were subjected to Gaussian

filtration and segmentation. The endocortical region of the vertebral body was manually selected

beginning 100 µm inferior to the cranial growth plate and extending to 100 µm superior to the

caudal growth plate. Trabecular bone was segmented from soft-tissue using a threshold of 460

$mgHA/cm^3$. Measurements included trabecular bone volume fraction (Tb.BV/TV, %), trabecular

thickness (Tb.Th, mm), trabecular number (Tb.N, mm^{-1}), trabecular separation (Tb.Sp, mm),

connectivity density (Conn.D, mm^{-3}), and trabecular bone mineral density (Tb.BMD,

$mgHA/cm^3$).

To address microarchitecture in the primary and secondary spongiosa of the distal femur and

the cortical bone of the femur midshaft, scans were acquired with the same parameters used for

the L5 vertebrae. The primary spongiosa region of interest started at the peak of the distal growth

plate and extended proximally for 500 µm (50 transverse slices) and included the whole cross-

section of the bone. The secondary spongiosa region of interest started immediately superior to

the primary spongiosa region and extended 1500 µm (150 transverse slices) proximally and

included the endocortical region of the bone. In both regions, bone was segmented from soft-

tissue using a mineral density threshold of 460 mgHA/cm^3. Trabecular bone in the endocortical

area of the secondary spongiosa region was analyzed for bone volume fraction (Tb.BV/TV, %),

trabecular thickness (Tb.Th, mm), trabecular number (Tb.N, mm^{-1}), trabecular separation (Tb.Sp,

mm), connectivity density (Conn.D, mm^{-3}), and trabecular bone mineral density (Tb.BMD,

mgHA/cm^3). The bone in the primary spongiosa was analyzed for bone volume, total volume,

and bone mineral density.

Cortical bone architecture was analyzed in a 500 µm long region at the femoral mid-

diaphysis (55% of the total femur length inferior to the top of the femoral head). Cortical bone

was segmented from soft tissue using a threshold of 700 mgHA/cm^3. Measurements included

cortical bone area (Ct.Ar, mm^2), total cortical cross-sectional area (Tt.Ar, mm^2), cortical bone

area fraction (Ct.Ar/Tt.Ar, %), cortical thickness (Ct.Th, mm) and cortical porosity (%).

Histology and Histomorphometry

Tibiae were fixed in 10% neutral-buffered formalin and transferred to 70% ethanol after 24

hours. Samples were decalcified, paraffin-embedded, sectioned, and stained with hematoxylin

and eosin. Adipocyte size and number in the marrow in the region of the secondary spongiosa

were analyzed with BIOQUANT OSTEO software (BIOQUANT Image Analysis Corporation,

Nashville, TN). Total adipocyte area was divided by total area (marrow plus trabecular bone) in

the secondary spongiosa and multiplied by 100 to calculate adipocyte volume / total volume

(AV/TV). Adipocyte number was normalized to the total area measured.

Vertebrae were fixed in 10% neutral-buffered formalin and transferred to 70% ethanol

after 24 hours. The fixed lumbar vertebrae (L2-L5) were dehydrated with acetone and embedded

in methylmethacrylate. Undecalcified 4-µm-thick sections were obtained by microtome and

stained with Von Kossa method for showing the mineralized bone. The consecutive second

section was left unstained for the analysis of fluorescence labeling and the third section was

stained with 2% Toluidine Blue (pH 3.7) for the analysis of osteoblasts, osteoid, and osteoclasts.

The bone histomorphometric analysis was performed in the lumbar vertebra under 200X

magnification in a 1.35 mm high x 1.3 mm wide region where was located 400 μm away from

the upper and lower growth plate using OsteoMeasure analyzing software (Osteometrics Inc.,

Decatur, GA), in accordance with published guidelines [36]. The structural parameters (BV/TV,

Tb.Th, Tb.N, and Tb.Sp) were obtained by taking an average from two different analysis of

consecutive sections. The structural, dynamic and cellular parameters were calculated and

expressed according to the standardized nomenclature [36].

Enzyme Immunoassays

Serum concentrations of amino-terminal propeptide of type 1 procollagen (P1NP), cross-linked

C-telopeptide (CTx) and TNF-related activation-induced cytokine (RANKL) were measured

with the Rat/Mouse P1NP enzyme immunoassay (EIA), RatLaps EIA (both from

Immunodiagnostic Systems, Scottsdale, AZ), and the Quantikine ELISA Mouse

TRANCE/RANKL/TNSFSF11 Kit (R&D Systems, Minneapolis, MN). The intraassay variations

were 6.3%, 6.9% and 4.3%, and the interassay variations were 8.5%, 12% and 6.9% respectively.

All measurements were performed in duplicate.

Osteoblast assays

MC3T3-E1 (clone 4) cells were purchased from ATCC and maintained at a low passage number

in α-MEM containing 1% penicillin streptomycin (P/S) and 10% fetal bovine serum (FBS). For

calcium assays, cells were plated overnight in 96-well black wall plates at a density of 5×10^4

cells/well. Cells were then differentiated for 7 days with ascorbic acid and β-glycerophosphate as

previously described [37]. PTH was aliquoted and stored as above, and thawed and diluted

immediately prior to use. Propranolol hydrochloride (P0884, Sigma-Aldrich) was dissolved in 1x

HBSS buffer containing 20 mM HEPES for each experiment. Calcium assays were performed

with the FLIPR Calcium 6 Assay Kit (Molecular Devices) according to the manufacturer's

protocol. In brief, on day 7 of differentiation, cells were incubated with calcium dye for two

hours at 37°C and 5% CO_2. Then, fluorescence was measured on a FlexStation 3 Multi-Mode

Microplate Reader using SoftMax Pro 7 software (Molecular Devices) in kinetic mode in

response to additions of a range of concentrations of PTH. Change in relative fluorescence units

(maximum – minimum) is indicative of intracellular calcium (iCa^{2+}) levels. Concentration-

dependent change in max-min value was plotted on a logarithmic scale and fit with a 4-

parameter logistic curve to calculate the EC_{50}. In some experiments, calcium dye incubation was

concurrent with pretreatment with propranolol. Additional 24-well plates were treated

simultaneously with PTH and propranolol for 1 hour, without the Calcium 6 reagents, and were

saved for mRNA analyses.

Propranolol extraction and measurements

Marrow was prepared for extraction by homogenization at a 1:1 ratio with 50 mM potassium

phosphate buffer pH 7.4. A volume of 20 µL of serum or marrow homogenate was combined

with 100 µL of HPLC grade acetonitrile and vortex mixed for two minutes. Subsequent to

centrifugation at 14000 rpm for five minutes, the supernatant was transferred to a 96-well plate

for liquid chromatograph tandem mass spectrometry (LC/MS-MS) analysis. A calibration curve

was formed in mouse plasma from 1.00-200 nM by serial dilution and extracted via the same

methodology. An Agilent 1200 system consisting of a binary pump, column compartment and

autosampler was used for solvent delivery and sample introduction. Chromatographic separation

was performed on an Agilent Zorbax SB C18 2.1 x 50 mm 2.7 µm column via a gradient using

0.1% formic acid in water (A) and 0.1% formic acid in acetonitrile (B). Gradient elution was

95% A ramping to 5% A from 0.0-3.0 minutes, with re-equilibration at initial conditions from

3.0 to 4.0 minutes. The flow rate was 0.75 mL/min, and column temperature was 30°C.

Detection of propranolol was obtained using an Agilent 6460 triple quadrupole mass

spectrometer, monitoring the transition $260.0 \rightarrow 116.0$ with a fragmentor of 86 V and a collision

energy of 21 V. The retention time of propranolol was 1.89 minutes.

RNA isolation and real-time PCR (qPCR)

Total RNA was isolated from whole tibia and cortical shell using the standard TRIZOL (Sigma,

St. Louis, MO) method. Total RNA from cells was isolated using an RNeasy Mini Kit (Qiagen).

cDNA was synthesized using the High Capacity cDNA Reverse Transcriptase Kit (Applied

Biosystems, Foster City, CA) according to the manufacturer's instructions. mRNA expression

analysis was performed using an iQ SYBR Green Supermix or Taqman Gene Expression Assays

with an iQ5 thermal cycler and detection system (Bio-Rad, Hercules, CA). Hypoxanthine

guanine phosphoribosyl transferase (*Hprt*) was used as an internal standard control gene[38].

Primer sequences are listed in the supplemental information (Table S1).

Norepinephrine extraction and measurements

A volume of 20 µL of serum was combined with 100 µL of HPLC grade acetonitrile and vortex

mixed for two minutes. Subsequent to centrifugation at 14000 rpm for five minutes, the

supernatant was transferred to a 96-well plate for liquid chromatograph tandem mass

spectrometry (LC/MS-MS) analysis. A calibration curve was formed in mouse plasma from

0.305-2500 nM by serial dilution and extracted via the same methodology. An Agilent 1200

system consisting of a binary pump, column compartment and autosampler was used for solvent

delivery and sample introduction. Chromatographic separation was performed on a Phenomenex

Hydro RP 2.0 x 150 mm 4 µm column via a gradient using 0.1% formic acid in water (A) and

0.1% formic acid in acetonitrile (B). Gradient elution was 98% A from 0-1 minute, ramping to

50% A from 1.1 to 3.0 minutes, holding at 50% until 5.5 minutes, with re-equilibration at initial

conditions from 5.6 to 7.5 minutes. The flow rate was 0.30 mL/min, and column temperature was

30°C. Detection of norepinephrine was obtained using an Agilent 6460 triple quadrupole mass

spectrometer, monitoring the transition 152.0 → 107.0 with a fragmentor of 94 V and a collision

energy of 18 V. The retention time of norepinephrine was 1.37 minutes.

Detection of Adrb2 mRNA transcript in femur

Femurs were isolated from 1-week-old or 6-week-old B6 mice and fixed in 10% neutral buffered

formalin for 24 hours, then decalcified in 0.5M EDTA pH 7.8 solution by rocking at 4□ C for 1-

2 weeks. Samples were dehydrated in 70% EtOH, embedded in paraffin, and sectioned in 5 μM

sections using standard methods. RNA *in situ* hybridization was performed using the RNAscope

FFPE Multiplex Fluorescent Assay V2 (ACD Bio) per manufacturer's instructions. Briefly,

sections were deparaffinized followed by pre-treatment using hydrogen peroxide and target

retrieval was achieved by steaming for 15 minutes. Tissue sections were treated with protease,

and the RNAscope assay was performed as outlined by manufacturer's instructions using probes

for *Acp5* and *Adrb2*. Images were acquired using a Leica TCS SP8 laser scanning confocal

microscope with a 20x air objective or 63x oil objective.

Sample size estimation and statistics

Sample size for mouse experiments was estimated based on previous experiments in our

laboratory using trabecular BV/TV as a primary outcome. Secondary outcomes of interest

included serum and histomorphometric indices of bone remodeling. All statistics were performed

with Prism 7 statistical software (GraphPad Software, Inc., La Jolla, CA). Normally distributed

data were analyzed for statistically significant differences using either a two-sided Student's *t*-

test or a two-way ANOVA followed by Holm-Sidak multiple comparison *post hoc* test, where

appropriate. Statistical significance was set at $p < 0.05$. All data are expressed as the mean ±

standard error of the mean (SEM). Outliers were handled as follows: first, any

scientific/biological reason for the outlier was investigated (i.e. sick mouse, degraded RNA, bad

histology section); second, if no explanation for the outlier was found we performed an outlier

test to determine if the datum was > 3 standard deviations (SD) from the mean of the other

samples in that group. If the datum was > 3 SD from the mean, it was excluded from further

analysis. Outlier handling did not significantly change the interpretation of the results.

Author Contributions

AT was responsible for experimental design, data acquisition, data analysis, interpretation, and drafting of the manuscript. ACB was responsible for experimental design, data acquisition, data analysis, interpretation, drafting and critical review of the manuscript. DJB was responsible for data acquisition, data analysis, interpretation, drafting of and critical revision of the manuscript. ALL was responsible for experimental design, data acquisition, data analysis, interpretation and drafting and critical review of the manuscript. HH was responsible for data acquisition, data analysis, interpretation and drafting of the manuscript. KTN was responsible for data analysis, drafting and critical review of the manuscript. KN was responsible for data acquisition, data analysis, interpretation, drafting of and critical revision of the manuscript. DB was responsible for experimental design, data acquisition, interpretation, drafting and critical revision of the manuscript. KLH was responsible for experimental design, data interpretation, drafting and critical revision of the manuscript. RB was responsible for data interpretation and critical revision of the manuscript. MLB was responsible for data interpretation and critical revision of the manuscript. ARG was responsible for experimental design, data interpretation and critical revision of the manuscript. KJM was responsible for experimental design, data acquisition, data analysis, interpretation, drafting and critical revision of the manuscript. All authors had final approval of the manuscript.

Acknowledgements

The authors thank Adriana Lelis Carvalho and Ryan Neilson for critical evaluation of the work.

This work was supported by the National Institute of Arthritis And Musculoskeletal And Skin

Diseases (NIAMS) and the National Institute of General Medical Sciences (NIGMS) of the

National Institutes of Health (NIH) under award numbers K01AR067858 and P20GM121301 to

KJM. This work utilized services of the Maine Medical Center Research Institute (MMCRI)

Molecular Phenotyping Core, which is supported by NIH/NIGMS P30GM106391, the

Physiology Core, which is supported by NIH/NIGMS P30GM106391 and P20GM121301, the

Histopathology and Histomorphometry Core, which is supported by NIH/NIGMS

P30GM106391, P20GM121301, and P30GM103392, and the Mouse Transgenic and In Vivo

Imaging Core which is supported by NIH/NIGMS P30GM103392. All cores also received

support from the Norther New England Clinical and Translational Research Network

NIH/NIGMS U54GM115516. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

 Figure Legends

Figure 1. Propranolol improved the positive effects of PTH on trabecular bone in the L5

vertebrae. Mice were treated with vehicle (white), 80 µg/kg PTH (light blue), 0.5 mg/ml

propranolol (gray), or PTH and propranolol (dark blue) for 4 weeks, from 16-20 weeks of age.

(A) Representative µCT images from L5 vertebrae. (B-G) Trabecular bone volume fraction (Tb.

BV/TV), bone mineral density (BMD), number (Tb.N), thickness (Tb.Th), separation (Tb.Sp),

and bone surface/bone volume (BS/BV). Bars represent mean ± standard error. *$p<0.05$,

**$p<0.01$ by Holm-Sidak *post hoc* test after a significant two-way ANOVA.

Figure 2. Trabecular bone in the femur secondary spongiosa is enhanced by combination PTH

and propranolol treatment. Mice were treated with vehicle (white), 80 µg/kg PTH (light blue),

0.5 mg/ml propranolol (gray), or PTH and propranolol (dark blue) for 4 weeks, from 16-20

weeks of age. (A) Representative 2D (top) and 3D (bottom) images of trabecular micro

architecture in the primary and secondary spongiosa of the distal femur. (B-I) Trabecular

microarchitectural parameters from the secondary spongiosa. (J-K) Volumetric and densitometric

measurements from the primary spongiosa. Bars represent mean ± standard error. *$p<0.05$,

**$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by Holm-Sidak *post hoc* test after a significant two-way

ANOVA.

Figure 3. Propranolol improved cortical thickness and density in PTH-treated mice. Mice were

treated with vehicle (white), 80 µg/kg PTH (light blue), 0.5 mg/ml propranolol (gray), or PTH

and propranolol (dark blue) for 4 weeks, from 16-20 weeks of age. (A-H) Cortical bone

microarchitectural parameters, cortical area (Ct.Ar), marrow area (Ma.Ar), total area (Tt.Ar)

cortical area fraction (Ct.Ar/Tt.Ar), cortical thickness (Ct.Th), cortical tissue mineral density

(Ct.TMD), cortical porosity, and polar moment of inertia (pMOI) were analyzed at the midshaft

of the femur. Bars represent mean ± standard error. *$p<0.05$, **$p<0.01$ by Holm-Sidak *post hoc*

test after a significant two-way ANOVA.

Figure 4. Propranolol improved bone formation markers in PTH-treated mice. Mice were treated

with vehicle (white), 80 μg/kg PTH (light blue), 0.5 mg/ml propranolol (gray), or PTH and

propranolol (dark blue) for 4 weeks, from 16-20 weeks of age. (A) Serum P1NP was measured

by ELISA. (B-C) Static and dynamic histomorphometry representative images and quantification

of architectural, bone formation and osteoblast parameters measured in L5 vertebrae. (D-H)

Gene expression of markers of bone formation and osteoblast maturity were analyzed in the tibia

and normalized by the non-modulated housekeeping gene, *Hprt*. Bars represent mean ± standard

error. *$p<0.05$, **$p<0.01$ by Holm-Sidak *post hoc* test after a significant two-way ANOVA.

Figure 5. β-blocker propranolol potentiated osteoblast intracellular Ca^{2+} and *Igf1* expression.

MC3T3-E1 cells were differentiated to day 7 in osteogenic media. (A) Fluorescence ratio

indicative of intracellular Ca^{2+} concentrations was expressed as a percentage of the maximum

level. (B) Representative traces of 100 nM PTH or vehicle-induced fluorescence over time with

pretreatment of vehicle, or 0.1, 1 or 10 μM propranolol. Note: vehicle + propranolol pretreatment

did not induce a significant signal, and traces are only slightly visible behind vehicle-only trace.

(C) Quantification of the maximum-minimum PTH-induced Ca^{2+} fluorescence after pretreatment

with vehicle or propranolol. One representative of three experiments (each performed in

triplicate) is shown. *$p<0.05$ compared to vehicle-treated (black), ^$p<0.05$ compared to 100 nM

PTH-treated (pink) by Tukey's *post-hoc* test after significant one-way ANOVA. (D-K) Gene

expression of *Runx2, Ocn, Fos,* and *Igf1* in MC3T3-E1 cells at day 7 after 1 hour treatment with

vehicle (white), 100 nM PTH (pink), 1 µM propranolol (blue) or 100 nM PTH + 1 µM

propranolol (purple). Genes are normalized to non-modulated housekeeping gene, *Hprt*. Each

data point represents the mean from an independent experiment, performed in triplicate.

Expression levels in vehicle-treated cells from each experiment were set to 1. Approximate Ct

values of vehicle-treated groups are shown above each graph. Bars represent mean of the three

independent experiments ± SEM. *$p<0.05$, **$p<0.01$ by Holm-Sidak *post hoc* test after a

significant two-way ANOVA.

Figure 6. Propranolol prevented PTH-induced resorption but not RANKL induction. Mice were

treated for 4 weeks (A-F) with vehicle (white), 80 µg/kg PTH (light blue), 0.5 mg/ml propranolol

(gray), or PTH and propranolol (dark blue). (A-B) Gene expression of *Ctsk* and *Trap5* was

analyzed in the whole tibia and normalized by the non-modulated housekeeping gene, *Hprt*. (C)

Serum CTx was measured by ELISA. (D) Histomorphometric analyses of osteoclast and

resorption parameters were measured in L5 vertebrae. (E) Serum norepinephrine (NE) was

measured by LC/MS/MS. (F) Serum Rankl was measured by ELISA. (G-H) Mice were treated

with PTH and/or propranolol for 5 days and cortical shell was collected 1 hour after the last PTH

or vehicle dose. Gene expression levels of *Rankl* and *Opg* were normalized to *Hprt*. Bars

represent mean ± standard error. *$p<0.05$, **$p<0.01$, ***$p<0.001$ by Holm-Sidak *post hoc* test

after a significant two-way ANOVA.

Figure 7. Propranolol is present in marrow, where *Adrb2* is expressed in osteoclasts *in vivo*. (A)

Left: Serum and marrow concentrations of propranolol after four days of treatment with 0.5

mg/mL in the drinking water, measured by LC/MS/MS. *p*-value calculated with a paired

Student's t-test. *Right:* Marrow to serum ratio was calculated for each individual mouse. Bar

represent mean ± SEM. N=6 mice. (B) 1-week-old mouse FFPE femurs were sectioned at 5 µm

and subjected to RNA hybridization with a probe specific to *Acp5,* red. Nuclei, blue. GP, growth

plate. TB, trabecular bone. BM, bone marrow. White arrowheads indicate multinucleated

osteoclasts. Imaged at 20x, shown is a maximum projection of a 3 µm Z-stack, with a step size of

1 µm. (C) Femurs from 5-6 week old C57BL/6J mice were decalcified and paraffin embedded.

Sections were hybridized with probes specific to *Acp5* (red) and *Adrb2* (green) to identify

osteoclasts expressing *Adrb2*. Nuclei, blue. Shown are representative of results from N=3

separate mice. TB, trabecular bone. BM, bone marrow. White arrowheads indicate

multinucleated osteoclasts with green *Adrb2* puncta. Single Z-plane, imaged at 63x.

Figure S1. Effects of PTH and propranolol on proximal tibia bone marrow adiposity. Mice were

treated for 4 weeks with vehicle (white), 80 µg/kg PTH (light blue), 0.5 mg/ml propranolol

(gray), or PTH and propranolol (dark blue). (A) Tibias were decalcified and stained with H & E.

Scale bar = 500 µm. (B-D) Adipocyte ghosts were measured using BIOQUANT OSTEO

software. (E) *Fabp4* gene expression was measured in whole tibia and normalized to non-

modulated housekeeping gene *Hprt.* Bars represent mean ± standard error. Indicated *p*-values

were determined by Holm-Sidak *post hoc* test after a significant interaction effect was found

using two-way ANOVA.

Figure S2. Relevant receptor expression levels in osteoblast-like cells. MC3T3-E1 cells were

differentiated to day 7 in osteogenic media. Gene expression of *Adrb1, Adrb2, Pth1r* and *Igf1r* in

MC3T3-E1 cells at day 7 after 1 hour treatment with vehicle (white), 100 nM PTH (pink), 1 µM

propranolol (blue) or 100 nM PTH + 1 µM propranolol (purple). Genes are normalized to non-

modulated housekeeping gene, *Hprt*. Each data point represents the mean from an independent

experiment, performed in triplicate. Expression levels in vehicle-treated cells from each

experiment were set to 1. Approximate Ct values of vehicle-treated groups are shown above each

graph. Bars represent mean of the three independent experiments ± SEM. *$p<0.05$, **$p<0.01$ by

Holm-Sidak *post hoc* test after a significant two-way ANOVA.

References

References

1.   Takeda S, Elefteriou F, Levasseur R, Liu X, Zhao L, Parker KL, Armstrong D, Ducy P,

Karsenty G. Leptin regulates bone formation via the sympathetic nervous system. Cell

[Internet]. 2002;111(3):305–17. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/12419242

2.   Elefteriou F, Ahn JD, Takeda S, Starbuck M, Yang X, Liu X, Kondo H, Richards WG,

Bannon TW, Noda M, Clement K, Vaisse C, Karsenty G. Leptin regulation of bone

resorption by the sympathetic nervous system and CART. Nature [Internet].

2005;434(7032):514–20. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15724149

3.   Mach DB, Rogers SD, Sabino MC, Luger NM, Schwei MJ, Pomonis JD, Keyser CP,

Clohisy DR, Adams DJ, O'Leary P, Mantyh PW. Origins of skeletal pain: sensory and

sympathetic innervation of the mouse femur. Neuroscience [Internet]. 2002;113(1):155–

66. Available from: http://www.ncbi.nlm.nih.gov/pubmed/12123694

4.   Motyl KJ, DeMambro VE, Barlow D, Olshan D, Nagano K, Baron R, Rosen CJ,

Houseknecht KL. Propranolol attenuates risperidone-induced trabecular bone loss in

female mice. Endocrinology. 2015;156(7).

5.   Motyl KJ, Bishop KA, Demambro VE, Bornstein SA, Le P, Kawai M, Lotinun S,

Horowitz MC, Baron R, Bouxsein ML, Rosen CJ. Altered thermogenesis and impaired

bone remodeling in Misty mice. J. Bone Miner. Res. 2013;28(9).

6.   Yirmiya R, Goshen I, Bajayo A, Kreisel T, Feldman S, Tam J, Trembovler V, Csernus V,

Shohami E, Bab I. Depression induces bone loss through stimulation of the sympathetic

nervous system. Proc Natl Acad Sci U S A [Internet]. 2006;103(45):16876–81. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/17075068

7. Ortuno MJ, Robinson ST, Subramanyam P, Paone R, Huang YY, Guo XE, Colecraft HM,

Mann JJ, Ducy P. Serotonin-reuptake inhibitors act centrally to cause bone loss in mice by

counteracting a local anti-resorptive effect. Nat Med [Internet]. 2016;22(10):1170–9.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/27595322

8. Yadav VK, Oury F, Suda N, Liu ZW, Gao XB, Confavreux C, Klemenhagen KC, Tanaka

KF, Gingrich JA, Guo XE, Tecott LH, Mann JJ, Hen R, Horvath TL, Karsenty G. A

serotonin-dependent mechanism explains the leptin regulation of bone mass, appetite, and

energy expenditure. Cell [Internet]. 2009;138(5):976–89. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/19737523

9. Ducy P, Amling M, Takeda S, Priemel M, Schilling AF, Beil FT, Shen J, Vinson C,

Rueger JM, Karsenty G. Leptin inhibits bone formation through a hypothalamic relay: a

central control of bone mass. Cell [Internet]. 2000;100(2):197–207. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/10660043

10. Kajimura D, Hinoi E, Ferron M, Kode A, Riley KJ, Zhou B, Guo XE, Karsenty G.

Genetic determination of the cellular basis of the sympathetic regulation of bone mass

accrual. J. Exp. Med. 2011;

11. Kondo H, Takeuchi S, Togari A. beta-Adrenergic signaling stimulates osteoclastogenesis

via reactive oxygen species. Am J Physiol Endocrinol Metab [Internet].

2013;304(5):E507-15. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23169789

12. Rodrigues WF, Madeira MFM, da Silva TA, Clemente-Napimoga JT, Miguel CB, Dias-

da-Silva VJ, Barbosa-Neto O, Lopes AH, Napimoga MH. Low dose of propranolol down-

modulates bone resorption by inhibiting inflammation and osteoclast differentiation. Br. J.

Pharmacol. England; 2012 Apr;165(7):2140–51.

13.   Yang S, Nguyen ND, Eisman JA, Nguyen T V. Association between beta-blockers and

771   fracture risk: A Bayesian meta-analysis. Bone. 2012.

772   14.   Toulis KA, Hemming K, Stergianos S, Nirantharakumar K, Bilezikian JP. β-adrenergic

773   receptor antagonists and fracture risk: A meta-analysis of selectivity, gender, and site-

774   specific effects. Osteoporos. Int. 2014;

775   15.   Pierroz DD, Bouxsein ML, Rizzoli R, Ferrari SL. Combined treatment with a beta-blocker

776   and intermittent PTH improves bone mass and microarchitecture in ovariectomized mice.

777   Bone [Internet]. 2006;39(2):260–7. Available from:

778   http://linkinghub.elsevier.com/retrieve/pii/S8756328206002006%5Cnpapers3://publicatio

779   n/doi/10.1016/j.bone.2006.01.145

780   16.   Hanyu R, Wehbi VL, Hayata T, Moriya S, Feinstein TN, Ezura Y, Nagao M, Saita Y,

781   Hemmi H, Notomi T, Nakamoto T, Schipani E, Takeda S, Kaneko K, Kurosawa H,

782   Karsenty G, Kronenberg HM, Vilardaga JP, Noda M. Anabolic action of parathyroid

783   hormone regulated by the beta2-adrenergic receptor. Proc Natl Acad Sci U S A [Internet].

784   2012;109(19):7433–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22538810

785   17.   Jean-Alphonse FG, Wehbi VL, Chen J, Noda M, Taboas JM, Xiao K, Vilardaga JP. β 2 -

786   adrenergic receptor control of endosomal PTH receptor signaling via Gβ 3. Nat. Chem.

787   Biol. 2017;

788   18.   Gesty-Palmer D, Luttrell LM. "Biasing" the parathyroid hormone receptor: A novel

789   anabolic approach to increasing bone mass? Br. J. Pharmacol. 2011.

790   19.   Cheloha RW, Gellman SH, Vilardaga JP, Gardella TJ. PTH receptor-1 signalling -

791   Mechanistic insights and therapeutic prospects. Nat. Rev. Endocrinol. 2015.

20.   Zayzafoon M, Fulzele K, McDonald JM. Calmodulin and calmodulin-dependent kinase

IIα regulate osteoblast differentiation by controlling c-fos expression. J. Biol. Chem. 2005;

21.   Moriya S, Hayata T, Notomi T, Aryal S, Nakamaoto T, Izu Y, Kawasaki M, Yamada T,

Shirakawa J, Kaneko K, Ezura Y, Noda M. PTH regulates β2-adrenergic receptor

expression in osteoblast-like MC3T3-E1 cells. J. Cell. Biochem. 2015;

22.   Bellido T, Ali AA, Gubrij I, Plotkin LI, Fu Q, O'Brien CA, Manolagas SC, Jilka RL.

Chronic elevation of parathyroid hormone in mice reduces expression of sclerostin by

osteocytes: A novel mechanism for hormonal control of osteoblastogenesis.

Endocrinology. 2005;

23.   Reid IR, Civitelli R, Halstead LR, Avioli L V., Hruska KA. Parathyroid hormone acutely

elevates intracellular calcium in osteoblastlike cells. Am. J. Physiol. - Endocrinol. Metab.

1987;

24.   Esen E, Lee SY, Wice BM, Long F. PTH Promotes Bone Anabolism by Stimulating

Aerobic Glycolysis via IGF Signaling. J. Bone Miner. Res. 2015;

25.   Zebaze R, Takao-Kawabata R, Peng Y, Zadeh AG, Hirano K, Yamane H, Takakura A,

Isogai Y, Ishizuya T, Seeman E. Increased cortical porosity is associated with daily, not

weekly, administration of equivalent doses of teriparatide. Bone. 2017;

26.   Osima M, Borgen TT, Lukic M, Grimnes G, Joakimsen RM, Eriksen EF, Bjørnerem.

Serum parathyroid hormone is associated with increased cortical porosity of the inner

transitional zone at the proximal femur in postmenopausal women: the Tromsø Study.

Osteoporos. Int. 2018;

27.   De Angelis E, Pecoraro M, Rusciano MR, Ciccarelli M, Popolo A. Cross-talk between

neurohormonal pathways and the immune system in heart failure: A review of the

literature. Int. J. Mol. Sci. 2019.

28.   Baker JG. Agonist and Inverse Agonist Actions of -Blockers at the Human 2-

Adrenoceptor Provide Evidence for Agonist-Directed Signaling. Mol. Pharmacol. 2003;

29.   Sozzani S, Agwu DE, McCall CE, O'Flaherty JT, Schmitt JD, Kent JD, McPhail LC.

Propranolol, a phosphatidate phosphohydrolase inhibitor, also inhibits protein kinase C. J.

Biol. Chem. 1992;

30.   Tinajero JC, Fabbri A, Dufau ML. Serotonergic inhibition of rat leydig cell function by

propranolol. Endocrinology. 1993;

31.   Rigby J, Scott A, Hawksworth G, Petrie J. A comparison of the pharmacokinetics of

atenolol, metoprolol, oxprenolol and propranolol in elderly hypertensive and young

healthy subjects. Br. J. Clin. Pharmacol. 1985;

32.   Meunier H, Labrie F. Specificity of the β2-adrenergic receptor stimulating cyclic AMP

accumulation in the intermediate lobe of rat pituitary gland. Eur. J. Pharmacol. 1982;

33.   Wheeler D, Sneddon WB, Wang B, Friedman PA, Romero G. NHERF-1 and the

cytoskeleton regulate the traffic and membrane dynamics of G protein-coupled receptors.

J. Biol. Chem. 2007;

34.   Khosla S, Monroe DG, Farr JN. Sympathetic b 1 -adrenergic signaling contributes to

regulation of human bone metabolism Graphical abstract Find the latest version□: J. Clin.

Invest. 2018;

35.   Bouxsein M, Boyd S. Guidelines for assessment of bone microstructure in rodents using

micro–computed tomography. J. bone … [Internet]. 2010;25(7):1468–86. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/20533309%5Cnhttp://onlinelibrary.wiley.com/doi/1

0.1002/jbmr.141/full

36.   Dempster DW, Compston JE, Drezner MK, Glorieux FH, Kanis JA, Malluche H, Meunier

PJ, Ott SM, Recker RR, Parfitt AM. Standardized Nomenclature, Symbols, and Units for

Bone Histomorphometry: A 2012 Update of the Report of the ASBMR Histomorphometry

Nomenclature Committee. J Bone Min. Res [Internet]. 2013;28(1):2–17. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3672237/

37.   Motyl KJ, Beauchemin M, Barlow D, Le PT, Nagano K, Treyball A, Contractor A, Baron

R, Rosen CJ, Houseknecht KL. A novel role for dopamine signaling in the pathogenesis of

bone loss from the atypical antipsychotic drug risperidone in female mice. Bone.

2017;103:168–76.

38.   Vengellur A, LaPres JJ. The role of hypoxia inducible factor 1α in cobalt chloride induced

cell death in mouse embryonic fibroblasts. Toxicol. Sci. 2004;

Tables

Table I. Body composition and areal bone parameters from mice treated with PTH and/or propranolol.

	Vehicle (N=10)	PTH (N=11)	Propranolol (N=9)	PTH + Propranolol (N=11)	2-way ANOVA p-values		
					PTH	Propranolol	Interaction
Body Mass (g)	21.8 ± 0.5	22.5 ± 0.4	21.7 ± 0.3	22.6 ± 0.4	ns	ns	ns
Lean Mass (g)	17.4 ± 0.4	17.9 ± 0.3	18.0 ± 0.3	17.9 ± 0.3	ns	ns	ns
Fat Mass (g)	2.6 ± 0.1	2.5 ± 0.2	2.6 ± 0.2	2.7 ± 0.2	ns	ns	ns
Total aBMD (g/cm^2)	0.0527 ± 0.0003	0.0544 ± 0.0004**	0.0535 ± 0.0004	0.0559 ± 0.0003^##	<0.01	<0.01	ns
Femur aBMD (g/cm^2)	0.0620 ± 0.0006	0.0655 ± 0.0008	0.0628 ± 0.0016	0.0678 ± 0.0008##	<0.01	ns	ns
Total aBMC (g)	0.438 ± 0.006	0.463 ± 0.006	0.444 ± 0.007	0.475 ± 0.009#	<0.01	ns	ns

*$p<0.05$, **$p<0.01$ compared to vehicle group.
^$p<0.05$, ^^$p<0.01$ compared to PTH group.
#$p<0.05$, ##$p<0.01$ compared to propranolol group.

Figure 1

Figure 2

Figure 4

A

B

		Propranolol	
Vehicle	PTH	Vehicle	PTH

C

	Vehicle (N=7)	PTH (N=7)	Propranolol (N=7)	PTH + Propranolol (N=9)	2-way ANOVA p-values		
					PTH	Propranolol	Interaction
BV/TV (%)	9.4 ± 0.8	14.0 ± 0.9**	10.3 ± 0.9	15.6 ± 0.4##	<0.01	*ns*	*ns*
Tb.Th (µm)	37.5 ± 1.7	43.5 ± 2.1*	39.0 ± 1.6	45.0 ± 1.5#	<0.01	*ns*	*ns*
Tb.N (mm⁻¹)	2.5 ± 0.1	3.2 ± 0.1**	2.6 ± 0.2	3.5 ± 0.1##	<0.01	*ns*	*ns*
Tb.Sp (µm)	376 ± 20	273 ± 14**	349 ± 24	246 ± 8##	<0.01	*ns*	*ns*
MAR (µm/day)†	1.19 ± 0.07	1.65 ± 0.07**	1.12 ± 0.13	1.65 ± 0.07##	<0.01	*ns*	*ns*
MS/BS (%)†	38.3 ± 2.1	42.3 ± 2.2	39.8 ± 3.3	37.3 ± 4.1	*ns*	*ns*	*ns*
BFR/BV (%/day)†	2.58 ± 0.21	3.61 ± 0.36	2.21 ± 0.26	3.08 ± 0.50	0.01	*ns*	*ns*
BFR/TV (%/day)†	0.23 ± 0.02	0.45 ± 0.05*	0.25 ± 0.05	0.46 ± 0.09#	<0.01	*ns*	*ns*
BFR/BS (µm³/µm²/day)†	0.45 ± 0.04	0.70 ± 0.06*	0.45 ± 0.06	0.62 ± 0.08	<0.01	*ns*	*ns*
N.Ob/B.Pm (mm⁻¹)	14.7 ± 1.0	22.5 ± 1.5**	12.4 ± 0.6	19.4 ± 1.2##	<0.01	0.03	*ns*
N.Ob/T.Ar (mm⁻²)	71.1 ± 8.5	147.7 ± 6.5**	64.4 ± 3.7	127.0 ± 7.9##	<0.01	*ns*	*ns*
Ob.S/B.Pm (%)	17.4 ± 1.2	26.8 ± 1.8**	14.8 ± 0.8	24.2 ± 1.4##	<0.01	*ns*	*ns*
OS/BS (%)	14.1 ± 1.2	18.8 ± 2.0	13.3 ± 1.3	12.5 ± 1.2^^	*ns*	0.02	*ns*
O.Th (µm)	2.69 ± 0.13	3.14 ± 0.14	2.71 ± 0.27	2.76 ± 0.12	*ns*	*ns*	*ns*

*$p<0.05$ compared to vehicle, **$p<0.01$ compared to vehicle
^$p<0.05$ compared to PTH, ^^$p<0.01$ compared to PTH
#$p<0.05$ compared to PRO, ##$p<0.01$ compared to PRO
†N=4-5 for dynamic measurements.

D   E   F   G   H

Figure 6

D

	Vehicle (N=7)	PTH (N=7)	Propranolol (N=7)	PTH + Propranolol (N=9)	2-way ANOVA p-values		
					PTH	Propranolol	Interaction
N.Oc/B.Pm (mm^{-1})	5.9 ± 0.3	6.2 ± 0.2	5.3 ± 0.4	4.2 ± 0.2$^{#,^{\wedge\wedge}}$	ns	<0.01	0.02
N.Oc/T.Ar (mm^{-2})	28.2 ± 2.4	41.3 ± 2.4**	27.4 ± 1.9	27.7 ± 1.5$^{\wedge\wedge}$	<0.01	<0.01	<0.01
Oc.S/B.Pm (%)	14.8 ± 0.8	15.9 ± 0.5	13.1 ± 0.9	10.6 ± 0.5$^{\wedge\wedge}$	ns	<0.01	0.02
ES/BS (%)	5.9 ± 0.5	5.8 ± 0.4	5.0 ± 0.7	4.1 ± 0.3$^{\wedge}$	ns	0.01	ns
*$p<0.05$ compared to vehicle, **$p<0.01$ compared to vehicle							
^$p<0.05$ compared to PTH, ^^$p<0.01$ compared to PTH							
#$p<0.05$ compared to PRO, ##$p<0.01$ compared to PRO							

Figure 7